1

incR: a free new R package to analyse incubation behaviour

Pablo Capilla-Lasheras

ORCiD 0000-0001-6091-7089

p.capilla@exeter.ac.uk / pacapilla@gmail.com

Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of

Exeter, Penryn, UK

Keywords: incubation, R package, software, nest temperatures

Abstract

Incubation represents a life stage of pivotal importance for the optimal development of avian

embryos. For most birds, incubation poses a trade-off between investing in self-maintenance

or offspring care. Furthermore, incubation is affected by environmental temperatures and,

therefore, will be likely impacted by climate change. Despite its relevance and readily

available temperature logging methods, avian incubation research is hindered by recognised

limitations in available software. In this paper, a new quantitative approach to analyse

incubation behaviour is presented. This new approach is embedded in a free R package,

`incR`. The core algorithm in `incR` is validated and it is shown that the method extracts

accurate metrics of incubation behaviour (e.g. number and duration of incubation bouts). The

paper also presents a suggested workflow along with detailed R code to ease the practical

implementation of the new method.

Introduction

Incubation represents a crucial life stage for egg-laying vertebrates, of which birds are a paramount example. Fine control of incubation is essential and has deep ecological and evolutionary implications, notably for developing offspring but also for their parents (Conway and Martin 2000). For embryos, the thermal environment the incubating individual provides is essential for successful development. Suboptimal incubation temperatures can lead to delayed embryonic growth (Nord and Nilsson 2011), physiological changes, for example, in immunity (Ardia et al. 2010) and long-term survival consequences (Berntsen and Bech 2016). However, incubating individuals need to divide their time budget between incubation and self-maintenance (e.g. foraging) and, therefore, they allocate time to each activity according to prevalent ecological conditions (e.g. depending on ambient temperatures (Coe et al. 2015) and food availability (Londoño et al. 2008)). Despite a long standing scientific interest in incubation, we are still elucidating subtle ecological causes and consequences of this behaviour (Smith et al. 2015, Bulla et al. 2016) which may have important practical implications, for example, in the context of global climate change (Griffith et al. 2016).

The study of avian incubation is nowadays fuelled by recent technological advances (for an account of different technologies, see Smith et al. 2015). In particular, the use of iButtons® (Maxim Integrated) and probed Tinytags (Gemini Data Loggers) allows researchers to measure incubation temperature as frequently as every second for long periods of time with little logistics involved. These technologies have the potential to expand the range of species and scientific questions that researchers can address. However, the amount of data collected is usually much larger than it was traditionally (for example, via behavioural observations) and several analytical hurdles must be overcome.

Before answering questions about the biology behind an incubation pattern, the observer first needs to summarise the data, effectively reducing their dimension to a few variables that can be correlated with a set of predictors of interest. For example, number of incubation bouts and their duration are popular metrics in avian studies (Cooper and Voss 2013). The first software for the analysis of incubation temperatures was released more than 10 years ago: Rhythm (Cooper and Mills 2005). The benefits of Rhythm were instantaneous as it allowed the automated scoring of incubation (Cooper and Voss 2013, Coe et al. 2015). This software made fast and objective an otherwise time-consuming and observer-biased activity. However, in a time when incubation data collection is easier than ever before, Rhythm lacks much of the flexibility that handling big data sets require. Rhythm also has limited analytical and graphical capabilities, which are a desire when thousands of temperature records may be available. Lastly, the use of Rhythm is made difficult by its paid license under Raven. However, apart from Rhythm, no other specialised software is currently available to analyse incubation temperature data.

To overcome these difficulties, I have developed a free R package, `incR`. This package provides a suite of R functions that i) prepare and format a raw temperature time-series (via the `incRprep` and `incRenv` functions), ii) apply an automated algorithm to score incubation (`incRscan`) and iii) calculate several biologically relevant metrics of incubation (e.g. number of incubation bouts) (Figure 1). Users can apply the whole pipeline or use any of the components of `incR` separately. `incR` takes advance of the flexibility in data handling and graphical capabilities offered by R. I first explain the `incR` workflow and the automated algorithm to score incubation. Then, I use video-recordings of incubating blue (*Cyanistes caeruleus*) and great tit (*Parus major*) females along with incubation temperature data to validate the automated algorithm. I further show how `incR` can accurately calculate several metrics of incubation behaviour. Finally, I discuss the general application of this new method

100   and its potential limitations. A stable version of the package is available on CRAN (v 1.0.2)

101   and a development version can be found on GitHub (v 1.0.2.9000.

102   https://github.com/PabloCapilla/incR).

`incR` workflow

The method implemented in `incR` exploits variation in incubation and ambient temperature

to calculate the presence or absence of the incubating individual in the nest. Ambient

temperature data are ideally collected near the nesting site but can also be obtained from web-

based sources. Code and advice to replicate the analyses presented here can be found in

Appendix 1 and 2, the package documentation (https://cran.r-

project.org/web/packages/incR/incR.pdf) and in the package vignette (accessible in R via:

`browseVignettes("incR")`).

Data preparation: `incRprep` and `incRenv`

Several temperature files from the same nest will be typically combined by the user before

the analysis starts. This combined file should consist of two columns: date-time and

temperature values. The first step in the pipeline is performed by `incRprep`, which simply

prepares the dataset for other pipeline components. Then, `incRenv` can be used to assign

environmental temperature to every incubation temperature observation. As explained below,

environmental temperatures will be used by `incRscan` to score incubation and, therefore,

`incRenv` eases this intermediate step (Figure 1).

Automated incubation scoring: `incRscan`

The algorithm implemented by `incRscan` exploits the dynamics in nest temperature that

arise from the behaviour of the incubating individual and the differences between nest (i.e.

temperature in the nest cup, ideally obtained by placing a temperature sensor inside an egg or

in between several eggs) and environmental temperatures (see Table 1 for definitions of

terms).

Four possible situations broadly exist regarding the changes in nest temperature after the

incubating individual enters (on-bout) and leaves (off-bout) the nest (see Table 1 for

definitions of commonly used terms throughout the paper). These four cases are classified as

follows: 1) incubation off-bout when nest temperature is high (close to incubation

temperature); 2) incubation on-bout when nest temperature is high (close to incubation

temperature); 3) incubation off-bout when nest temperature is low (close to environmental

temperature); 4) incubation on-bout when nest temperature is low (close to environmental

temperature). Cases 3 and 4 are especially sensitive to the assumption that environmental

temperature is lower than incubation temperature (see Results and Discussion). The change in

nest temperature that is expected after an incubation on- / off-bout differs across the four

scenarios.

Assuming that environmental temperatures are lower than incubation ones, in case 1 when

the incubating individual leaves the nest, there exists a marked difference between nest and

environmental temperatures. Thus, a sharp drop in temperature is expected to follow. At this

point, any increase in nest temperature would mean that the bird has returned to the nest (on-

bout, scenario 2). In the absence of the incubating individual, nest temperatures decrease until

they match environmental temperatures. If an off-bout occurs when nest temperature is close

to the environmental one (scenario 3, e.g. because the incubating individual has carried out a

short on-bout and then leaves the nest when temperature has not markedly increased), the

decrease in nest temperature after an incubation off-bout would be small. However, in this

situation, any incubation on-bout would be reflected in a large increase in nest temperature

(scenario 4).

These four situations represent simplified extremes in a spectrum of possible scenarios but

they illustrate the general principle. To explain the analytical approach in more practical

terms, I here describe the analysis of one day of incubation (day 1), using the terminology

employed in the R package (Table 1).

For every time point in the incubation time series, whether scenarios 1 and 2 or 3 and 4 (see

above) are applicable is determined by calculating the difference between nest and

environmental temperatures. The user has some control over this decision via the

`temp.diff` argument in incRscan, allowing for two possibilities: i) nest temperature is

higher than environmental temperature by more than `temp.diff` Celsius degrees; and, ii)

nest temperature is within `temp.diff` Celsius degrees of the environmental one.

Therefore, the argument `temp.diff` helps determine whether scenarios 1/2 or 3/4 are

applicable for a specific time point.

Comparing the change in temperature between consecutive temperature recordings against

temperature thresholds, the algorithm presented here determines whether the incubating

individual is in the nest or an off-bout has occurred. Rather than having a fixed threshold for

the entire analysis, a flexible threshold value is applied. Furthermore, no threshold choice is

required from the user but it is calculated depending on the biology of the study species for

each day of analysis. The user should decide on a time window in which the incubating

individual is assumed to be in the nest. This time window is controlled by the arguments

`lower.time` and `upper.time`, representing the start and end of such time window (for

instance, for diurnal bird species this period can be set at night, when the incubating

individual rests in the nest). Within this time window, the maximum decrease in temperature

between pairs of consecutive points is set as a threshold for incubation off-bouts (hereafter,

`maxDrop`) for scenario 1. Assuming that nest temperatures are above environmental values,

`maxDrop` is thought to effectively represent the maximum drop in temperature associated

with periods when the incubating individual is in the nest. The threshold for incubation off-

bout in situation 3 must be lower than in scenario 1 (i.e. when nest temperature is close to

environmental temperature); thus, the argument `sensitivity` (taking values from 0 to 1)

allows for such reduction, setting the off-bout threshold in scenario 3 as `maxDrop`

x `sensitivity`. Similarly, `maxIncrease` is defined as the maximum increase in

temperature between pairs of consecutive points and is set as a threshold for incubation on-

bouts in scenario 4. Any increase in nest temperature in scenario 2 would mean an incubation

on-bout (see above). Note that `maxDrop` and `maxIncrease` do not need to be chosen by

the   user   but   are   reported   by   `incR_scan`   (in   a   returned   object   named

`incRscan_threshold`)

Once these thresholds are set, temperature differences between successive pairs of data points

throughout the day and between `upper.time` and `lower.time` are calculated. These

values   are   sequentially   compared   with   the   value   of   `maxDrop`   and   `maxIncrease`,

following a set of conditions:

For scenario 1 and 2,

187              $T_t - T_{t-1} <$ `maxDrop` (A); $T_t - T_{t-1} > 0$ (B).

188   For scenario 3 and 4,

189              $T_t - T_{t-1} <$ `maxDrop` x `sensitivity` (**C**); $T_t - T_{t-1} >$ `maxIncrease` (**D**).

$T_t - T_{t-1}$ being the *i*th and *i-1*th temperature recording from *i*=2 to *i*=*I* (*I* being equal to the

total number of daily data points evaluated). Off-bout periods are, then, defined between T*i*s

satisfying A or C and the closest subsequent situation in which T*j* , when *i<j*, satisfies B or D.

On-bout periods start after an off-bout finishes and last until A or C is fulfilled again.

To make this calculation conservative and robust against highly variable `lower.time` -

`upper.time` windows, whenever |*maxDrop*| > `maxNightVariation` (in Celsius

degrees) for a particular studied day, the value of `maxDrop` and `maxIncrease` of the

previous day of incubation is instead used. The result of this algorithm is a temporal sequence

of 0's and 1's representing on-bouts (1's) and off-bouts (0's). Based on this sequence and

using other functions within the `incR` package, biological information on inferred

incubation behaviour can be calculated.

Additional functions to extract biological metrics of incubation

Regardless of whether or not `incRscan` has been used to score incubation, the `incR`

package offers a suite of functions that can be applied to any binary time-series representing

incubation. The current stable version (1.0.2) allows the user to calculate onset and offset of

daily activity (`incRact`), percentage of daily time spent in the nest (`incRatt`), number

and average duration of on/off-bouts per day as well as individual off-bout duration and

timing (`incRbouts`) and nest temperature mean and variance for a customised time

window (`incRt`). The implementation of these functions is straightforward as they only

require a variable with binary data for on and off-bouts. These data are provided by

`incRscan` under the column name `incR_score`.

Validation of `incR` using temperature and video-recording data

To show the potential of this approach to yield meaningful metrics of incubation, I assessed

the performance of `incR`'s core functions. First, I carried out a sensitivity analysis in which I

evaluated the accuracy of `incRscan` over simulated values for its main arguments. I then

chose the optimal values for these arguments and showed that the combination of `incRscan`

and other `incR` functions can yield accurate measures of incubation behaviour. I applied the

whole pipeline to incubation temperatures collected using iButton® devices. For the same

incubation events, I used video footage of these nests to visually score incubation and then

compared these results to the automatic algorithm implemented in `incRscan`.

Field protocol and incubation data collection

Incubation temperatures were recorded during 2015 and 2016 using iButton® devices in two

blue tit and six great tit clutches respectively. Blue tit data came from an urban and suburban

population in Glasgow city (n = 2 clutches; 55° 52.18'N 4° 17.22'W and 55º 9'N 4º 31'W)

(Pollock et al. 2017), whereas great tit incubation data were recorded in an oak forest at the

Scottish Centre for Ecology and the Natural Environment (n = 2 clutches; SCENE, 56°

7.73'N 4° 36.79'W) (Pollock et al. 2017) and in a mixed forest (dominated by oak, birch and

pine trees) near the Netherlands Institute for Ecology (NIOO) (n = 4; ~52º 7' N 6º 59' E)

(Spoelstra et al. 2015). iButtons® were wrapped in a piece of black cloth and placed in the

nest cup, above the lining materials and among the eggs. Nest temperatures were recorded by

iButtons® every 2 or 3 minutes. Video cameras inside the nest-boxes were used to monitor

individual females and visually score incubation (see Pollock et al. 2017 for a general

explanation about video camera deployment). In total, 14 days of incubation were completely

or partially monitored using both iButtons® and recording cameras. Environmental

temperatures for the same period in Scotland were recorded either using iButtons® placed

outside nest-boxes. For the Dutch clutches, environmental temperatures from a weather

station approximately 18 Km away from the nest-box population were used. iButtons® were

downloaded in the field using portable devices and a single file per nest was compiled before

further analysis.

Data analysis

Using video footage, I determined whether or not the incubating female had been present in

the nest at every iButton® temperature time point. After preparing incubation temperature

data using `incRprep` and `incRenv`, I applied `incRscan` to score incubation and

compared its results to the footage-based scoring. I tested the performance of `incRscan` to

changing values of its three key arguments, i) `maxNightVariation` (testing values from

0.5 to 10 every 0.5), ii) `sensitivity` (from 0 to 1 every 0.1) and iii) `temp.diff` (from

0.5 to 10 every 0.5) (see Table 1 for definitions). When testing one argument, the others were

kept to default values of 1.5, 0.15 and 3 for `maxNightVariation`, `sensitivity` and

`temp.diff` respectively. `lower.time` and `upper.time` were always fixed to 10 p.m.

and 3 a.m (night time). For every test, I calculated the percentage of correctly scored

incubation time points. After selecting the best-performing combination of argument values

(i.e. highest percentage of agreement between `incRscan` and video footage), I compared

daily incubation attendance (i.e. percentage of time spent in the nest), number of daily off-

bouts and mean daily off-bout duration between `incRscan`-based and video footage-based

incubation scores. I present Pearson's correlations coefficients between the two metrics.

`incR` functions, statistical tests and graphical illustrations (apart from the left-hand side of

Figure 1) were produced in R version 3.4.1. Detailed technical guidelines to apply the `incR`

workflow and the validation shown in this manuscript can be found in Appendix 1 and 2 as

well as in the package´s vignette (accessible in R via: `browseVignettes("incR")`).

Results and Discussion

Within nest-boxes, changing values of `maxNightVariation` did not affect the

performance of `incRscan`. Similar results were found for `sensitivity` and

`temp.diff`, with only one nest-box being markedly affected by changes in these arguments

(Figure S1A-C). It is important to note that when `maxNightVariation` is set to a very

low value (effectively not allowing for much temperature variation in the `lower.time`-

`upper.time` time window) `incRscan` fails to yield any result as no temperature

threshold would be available. This result can be seen in Figure S1A: when evaluating

`maxNightVariation` equal to 0.5℃, data from only two out of eight nest-boxes were

extracted by `incRscan`.

Consistent variation in `incRscan` best-performing argument values was found among nest-

boxes (Figure 2), suggesting that differences in, for example, iButtons® deployment may be

affecting the accuracy of the `incRscan` algorithm. This potential effect has been

qualitatively suggested before (Smith et al. 2015) and highlights the importance of collecting

high quality data in the field. However, the percentage of agreement was always high (> 80%,

Figure 2). Highly consistent results were found within nest-boxes with marked among-box

variation with only one exception (nest-box G178_GT, Figure 2) in which setting

`maxNightVariation` to 4°C improved the percentage of agreement compared to that of

the default value (3 °C). The general pattern that arises across the eight nest-boxes is that

values above 1.5°C for `maxNightVariation` give the highest accuracy (90.27%. Figure

S1D). Similarly, values below 0.3 for `sensitivity` (90.27%) and a `temp.diff` value

of 4°C (91.16%) were found to be the most accurate argument choices (Figure S1E-F).

Given these results, I chose optimal values of 1.5°C, 0.25 and 4°C for

`maxNightVariation`, `sensitivity` and `temp.diff` respectively, yielding a

percentage of agreement across nest-boxes of 91.16% (maximum = 98.56%; minimum =

80.42). With these argument values, attendance calculated based on video footage and

inferred by `incRscan` showed a Pearson's correlation coefficient of 0.992 (t = 24.81, p <

0.0001, 95% confidence interval = 0.971-0.998. Figure 3A). Likewise, the algorithm in

`incRscan` was able to provide accurate off-bout information (Figure 3B & 3C). `incR`-

estimated off-bout number and mean daily off-bout duration were highly correlated with real

off-bout number and duration as extracted from video footage (for off-bout number: r =

0.972, t_{10} = 13.04, p < 0.0001, 95% confidence interval = 0.900-0.992. For daily mean off-

bout duration: r = 0.996, t_{10} = 34.69, p < 0.0001, 95% confidence interval = 0.985-0.999).

These results show that the method presented here can yield accurate metrics of incubation

behaviour. Based on the validation of Rhythm presented in Bueno-Enciso et al. (2017), `incR`

performs better than that software and yields higher correlations between video and iButton®

data (Bueno-Enciso et al. 2017) – but note the possible influence of different environmental

temperatures across studies. Implemented in a much more flexible workspace, data validation

and visualisation of incubation temperature data using `incR` is straightforward. For number

of off-bouts, the discrepancies between `incR` and video footage seem to arise from `incR`

slightly over-estimating the number of off-bouts (Figure 3B). This effect was mainly caused

by data from two nest-boxes (G178_GT and GT173_GT) which were collected in the same

year and location. The extent to which this result represents a consistent bias of the method or

an artefact of data collection (for instance, the iButton® being placed in a suboptimal

position) is unknown. However, the magnitude of this discrepancy was small (six off-bouts of

maximum differences between estimates for whole days; estimated regression slope $\pm$ SE =

0.926 ± 0.071) and the error is unlikely to be differentially affected across groups of nests

(e.g. experimental *versus* control in an experimental setup). Apart from the ones presented

here, other metrics of incubations can be calculated using `incR` (Figure 1 and see package

documentation), for which high reliability is expected given the results of this validation.

Benefits of `incR`

The benefits of `incR` are two-fold. Firstly, it represents a quantitative improvement over

other methods. The results of the validation suggest that `incR` may perform better than other

approaches (see validation of Rhythm in Bueno-Enciso et al. 2017). No assumptions about

minimum off-bout time or off-bout temperature reductions are needed. `incRscan` uses

changes between consecutive temperature points, making the detection of short off-bouts

possible. The inclusion of environmental temperatures informs the analysis, allowing for off-

bout detection when nest and environmental temperatures are similar. In Appendix 1 and 2, I

offer detailed instructions to reproduce the analysis presented here. More generally, using a

script-based approach will improve repeatability and will ease collaboration.

Secondly, `incR` embraces the philosophy of the R project: it is completely free and is in

constant improvement. Further developments in the method to score incubation could be

embedded in or used jointly with `incR` to extract metrics of incubation.

Limitations

The capability of `incR`, or very likely of any other analytical tool to study incubation

temperatures, to yield accurate results will certainly correlate with data quality. Correct

logger deployment and data validation are, therefore, crucial. Two key assumptions underlie

the use of the algorithm implemented by `incRscan`. First, the incubating individual is

assumed to rest in the nest in the `lower.time` - `upper.time` time window. This

assumption holds for most species in temperate and tropical zones, for which activity outside

the nest is paused during night time (a reversed pattern is expected in nocturnal species).

However, careful consideration of this assumption will be needed when the species of interest

do not have a rhythmic incubation pattern or rhythms differ from 24 h (Bulla et al. 2016).

Secondly, the accuracy of `incRscan` will also depend on the difference between incubation

and environmental temperatures. Small differences between these temperatures will lead to

subtle changes in temperature after the incubating individual enters and leaves the nest,

affecting the detectability of such events. Further tests need to be carried out to evaluate the

accuracy of `incR` in these situations. For instance, comparing the performance of

`incRscan` for data collected on the same species at different latitudes (and thus with likely

large or small differences between environmental and nest temperatures) might provide

valuable information on the general applicability of this methodology.

Conclusions

We have developed a method that accurately extracts behavioural and temperature

information from series of incubation temperature recordings. This method can potentially be

used to study incubation in a broad range of species and ecological contexts and, therefore,

assist the wide community of researchers studying incubation both in the wild and in the

laboratory. For different species and environments, validation will be needed but we also

provide detailed practical advice to carry out such validation. In order to ease its application,

two appendices show in detail how researchers can easily adapt and calibrate this method to

their study species.

Acknowledgements

Barbara Helm and Davide Dominoni offered the inspiration to write this manuscript: without

their support and guidance none of this work would have been possible. Robyn Womack and

Natalie van Dis very kindly let me use their great tit data from Glasgow and The Netherlands.

They also tested the package in multiple occasions and provided thorough technical advice.

Field data collection was significantly eased by the help of Davide Dominoni, Chris Pollock,

Jessica Clark and Paul Jerem, along with many other volunteers. Many people at the Institute

of Biodiversity, Animal Health and Comparative Medicine of the University of Glasgow

provided very valuable advice. In particular, I thank Jason Matthiopoulos. Barbara Helm,

Davide Dominoni, Natalie van Dis and Natalia Bravo-Santano read an earlier draft of this

manuscript and their feedback greatly improved the text.

Funding

368　The initial stages of this project were funded by a postgraduate scholarship from Iberdrola

369　Foundation. PC-L currently thanks the British Biotechnology and Biological Science

370　Research Council (BBSRC) for a SWBio postgraduate scholarship.

Conflicts of interest

PC-L declares no conflict of interest.

References

Ardia, D. R., Pérez, J. H. and Clotfelter, E. D. 2010. Experimental cooling during incubation leads to reduced innate immunity and body condition in nestling tree swallows. - Proc. R. Soc. B 277: 1881–1888.

Berntsen, H. H. and Bech, C. 2016. Incubation temperature influences survival in a small passerine bird. - J. Avian Biol. 47: 141–145.

Bueno-Enciso, J., Barrientos, R. and Sanz, J. J. 2017. Incubation Behaviour of Blue *Cyanistes caeruleus* and Great Tits *Parus major* in a Mediterranean Habitat. - Acta Ornithol. 52: 21–34.

Bulla, M., Valcu, M., Dokter, A. M., Dondua, A. G., Kosztolányi, A., Rutten, A., Helm, B., Sandercock, B. K., Casler, B., Ens, B. J., Spiegel, C. S., Hassell, C. J., Küpper, C., Minton, C., Burgas, D., Lank, D. B., David, C., Loktionov, E. Y., Nol, E., Kwon, E., Smith, F., Gates, H. R., Vitnerová, H., Prüter, H., James, A., Clair, J. J. H. S., Lamarre, J., Rausch, J., Reneerkens, J., Conklin, J. R., Burger, J., Bêty, J., Coleman, J. T., Figuerola, J., Hooijmeijer, J. C. E. W., Alves, J. A., Smith, J. A. M., Koivula, K., Gosbell, K., Exo, K., Niles, L., Koloski, L., Mckinnon, L., Praus, L., Giroux, M., Sládeček, M., Boldenow, M. L., Goldstein, M. I., Šálek, M., Senner, N., Rönkä, N., Lecomte, N., Gilg, O., Vincze, O., Johnson, O. W., Smith, P. A., Woodard, P. F., Pavel, S., Battley, P. F., Bentzen, R., Lanctot, R. B., Porter, R., Saalfeld, S. T., Freeman, S., Brown, S. C., Yezerinac, S., Székely, T., Montalvo, T., Piersma, T., Loverti, V., Pakanen, V., Tijsen, W. and Kempenaers, B. 2016. Unexpected diversity in socially

synchronized rhythms of shorebirds. - Nature 540: 109–113.

Coe, B. H., Beck, M. L., Chin, S. Y., Jachowski, C. M. B. and Hopkins, W. A. 2015. Local

variation in weather conditions influences incubation behavior and temperature in a

passerine bird. - J. Avian Biol. 46: 1–10.

Conway, C. J. and Martin, T. E. 2000. Evolution of passerine incubation behavior: influence

of food, temperature, and nest predation. - Evolution 54: 670–685.

Cooper, C. and Mills, H. 2005. New software for quantifying incubation behavior from time-

series recordings. - J. F. Ornithol. 76: 352–356.

Cooper, C. B. and Voss, M. A. 2013. Avian Incubation Patterns Reflect Temporal Changes in

Developing Clutches. - PLoS One 8: e65521.

Griffith, S. C., Mainwaring, M. C., Sorato, E. and Beckmann, C. 2016. High atmospheric

temperatures and " ambient incubation " drive embryonic development and lead to

earlier hatching in a passerine bird. - R. Soc. Open Sci. 3: 150371.

Londoño, G. A., Levey, D. J. and Robinson, S. K. 2008. Effects of temperature and food on

incubation behaviour of the northern mockingbird, Mimus polyglottos. - Anim. Behav.

76: 669–677.

Nord, A. and Nilsson, J.-Å. 2011. Incubation temperature affects growth and energy

metabolism in blue tit nestlings. - Am. Nat. 178: 639–51.

Pollock, C., Capilla-Lasheras, P., McGill, R. A. R., Helm, B. and Dominoni, D. M. 2017.

Integrated behavioural and stable isotope data reveal altered diet linked to low breeding

success in urban-dwelling blue tits (Cyanistes caeruleus). - Sci. Rep. 7: 5014.

Smith, J. A., Cooper, C. B. and Reynolds, S. J. 2015. Advances in techniques to study

incubation. - In: Deeming, D. C. and Reynolds, S. J. (eds), Nests, Eggs, and Incubation:

New ideas about avian reproduction. Oxford University Press, pp. 312.

Spoelstra, K., van Grunsven, R. H. A., Donners, M., Gienapp, P., Huigens, M. E., Slaterus,

R., Berendse, F., Visser, M. E. and Veenendaal, E. 2015. Experimental illumination of

natural habitat--an experimental set-up to assess the direct and indirect ecological

consequences of artificial light of different spectral composition. - Philos. Trans. R. Soc.

B 370: 20140129.

Figure Captions

Figure 1. `incR` workflow and visualisation of corresponding analysis of nest temperature data at each step of the workflow. `incRprep` prepares raw data time series (1) for the pipeline and `incRenv` adds environmental temperatures (2). `incRscan` classifies data points in absence or presence of the incubating individual in the nest (3). From a sequence of 0´s and 1´s calculated by `incRscan`, `incRbouts`, `incRatt`, `incRact` and `incRt` extract information about on/off-bouts, nest attendance, start and end of activity and averaged nest temperatures for customised time windows. The green line in 2 and 3 represents environmental temperature, whereas the colour of the dots in 3 illustrates on- (light red) and off-bouts (purple).

Figure 2. Percentage of agreement between `incRscan` and video-footage across eight different nest-boxes. Colour codes represent individual nest-boxes and each point within nest-box illustrates the highest percentage of agreement for each of the three simulations, testing `maxNightVariation`, `sensitivity` and `temp.diff`. Highly consistent results are found within nest-boxes with marked among-box variation – with one exception (G178_GT) in which setting `maxNightVariation` to 4°C improved the percentage of agreement compared to that of the default value (3 °C). Points are slightly offset in the x axis to aid visualisation of overlaying points.

Figure 3. Correlations between video-footage and `incR` estimates of incubation attendance (percentage of daily time spent in the nest (A), number of daily off-bouts (B) and daily mean off-bout duration in minutes (C). Colour codes represent individual nest-boxes and each point illustrates one day of incubation. Dashed black line was drawn following an intercept of 0 and a slope of 1.

478   **Tables**

479   **Table 1.** Glossary of terms used in this manuscript.

Term	Type	Description
Incubation temperature		Maximum and stable temperature reached in the nest. Normally obtainable after a long presence of the incubating individual in the nest.
Nest temperature		Temperature inside the nest cup at any given time point. By this term I refer to a variable value that depends on whether, and for how long, the incubating individual is in or out the nest.
Environmental temperature		Air temperature outside nest.
`incR_scan`	R function	Calculates presence or absence of the incubating individual in the nest based on nest and ambient temperature variation
`temp.diff`	`incR_scan` argument	Difference allowed between nest and environmental temperatures
`lower.time`	`incR_scan` argument	Start time of a time window when the incubating individual is assumed to be in the nest
`upper.time`	`incR_scan` argument	End time of a time window when the incubating individual is assumed to be in the nest
`sensitivity`	`incR_scan` argument	Reduction in off-bout threshold when nest temperatures are close to environmental ones.
`maxNightVariation`	`incR_scan` argument	Maximum variation allowed in the `lower.time - upper.time` window. It controls for big drops in temperature within this temporal window (i.e. night-time incubation off-bouts).
`maxDrop`	Internal calculation in `incR_scan`	Maximum drop in temperature between two consecutive time points within the `lower.time – upper.time` window.
`maxIncrease`	Internal calculation in `incR_scan`	Maximum increase in temperature between two consecutive time points within the `lower.time – upper.time` window.