Long title: A revised understanding of *Tribolium* morphogenesis further

reconciles short and long germ development

Short title: A major revision to our understanding of short germ embryogenesis

Author: Matthew A. Benton

Author affiliation: Department of Zoology, University of Cologne, Cologne, NRW, Germany

Corresponding author: Matthew A. Benton, matthewabenton@gmail.com

Abstract

In *Drosophila melanogaster*, the germband forms directly on the egg surface and solely

consists of embryonic tissue (termed long germ development). In contrast, most insect

embryos undergo a complicated set of tissue rearrangements to generate a condensed, bi-

layered germband (termed short/intermediate germ development). The ventral side of the

germband is embryonic, while the dorsal side is thought to be an extraembryonic tissue

called the amnion. While this tissue organisation has been accepted for decades, and has

been widely reported in insects, its accuracy has not been directly tested in any species.

Using live cell tracking and differential cell labelling in the short germ beetle *Tribolium*

castaneum, I show that most of the cells previously thought to be amnion actually give rise

to large parts of the embryo. This process occurs via the dorsal-to-ventral flow of cells and

contributes to germband extension. In addition, I show that true 'amnion' cells in *Tribolium*

originate from a small region of the blastoderm. Together, my findings show that

development in the short germ embryos of *Tribolium* and the long germ embryos of

Drosophila is more similar than previously proposed. Dorsal-to-ventral cell flow also occurs

in *Drosophila* during germband extension, and I argue that the flow is driven by a conserved

set of underlying morphogenetic events in both species. Furthermore, the revised *Tribolium*

fatemap that I present is far more similar to that of *Drosophila* than the classic *Tribolium*

fatemap. Lastly, my findings show that there is no qualitative difference between the

structures of the blastoderm and the short/intermediate germ germband. As such, the same

tissue patterning mechanisms could function continuously throughout the cellularised

blastoderm and germband stages, and easily shift between them over evolutionary time.

Introduction

Insects are the most speciose phylum of animals and display remarkable diversity in adult

morphology (1). Insect embryo development is also very diverse, particularly in the stages

leading to the formation of the elongated, segmented embryo (called the germband) (2).

The molecular and morphogenetic basis of this process is best understood in the fly

Drosophila melanogaster. In this species, a predominantly hierarchical chain of patterning

events specifies nearly all segments more-or-less simultaneously at the syncytial blastoderm

stage (3). Cellularisation takes place near the end of this process, after which point

morphogenetic events such as germband extension (GBE) occur (see Fig 1 for schematic

summary). The *Drosophila* mode of development is termed long germ development and is

fairly representative of most true flies (4). In contrast, the vast majority of insects undergo

short or intermediate germ development, meaning that only a handful of segments are

specified at the blastoderm stage and the remaining segments are specified sequentially as

the germband elongates (5).

Short germ development has been best studied in the beetle *Tribolium castaneum,*

and recent research has shown that development in this species is more similar to

Drosophila than previously thought. For example, while previous molecular studies have

suggested that *Tribolium* segment specification occurs in a qualitatively different way to

Drosophila (6,7), recent detailed analyses have shown that a relatively conserved patterning

network may underlie segment specification in both species (8–10).

The morphogenetic events occurring during GBE are also more conserved than

previously thought. In *Drosophila*, GBE is predominantly driven by the mediolateral

intercalation of ectodermal cells (i.e. convergent extension), although cell deformation along

the anterior-posterior (AP) axis and cell divisions are also involved (11–16). In contrast to

this, *Tribolium* germband elongation was previously thought to be driven by the so-called

'growth zone' at the posterior of the germband (17). Now, however, it is clear that *Tribolium*

germband elongation is also predominantly driven by mediolateral cell intercalation (see Fig

1 for schematic summary of *Tribolium* development) (7,18,19). Furthermore, in both

Tribolium and *Drosophila*, this intercalation requires the striped expression of a specific

group of Toll genes (so-called Long Toll/Loto class genes) (20,21).

It is highly likely that germband elongation mediated by cell intercalation is

homologous in these two species, and probably in other arthropods, as well (21). As such, I

will hereafter refer to *Tribolium* 'germband elongation' as 'germband extension'/GBE,

unifying the *Drosophila/Tribolium* terminology. In addition, as there is no evidence for a

qualitatively different 'growth zone' in *Tribolium* (i.e. a specialised zone of volumetric

growth), I will refer to the posterior unsegmented region as the 'segment addition zone'

(8,23,24).

Despite the similarities described above, substantial differences in the embryonic

fatemaps of these species exist (Fig 1). In *Drosophila*, almost the entire blastoderm is fated

as embryonic tissue, and only a small dorsal region is fated as extraembryonic tissue (termed

the amnioserosa) (22). In contrast, in *Tribolium*, roughly the anterior third of the blastoderm

gives rise to an extraembryonic tissue called the serosa (25). Of the remaining blastoderm, a

large dorsal region is thought to give rise to a second extraembryonic tissue called the

amnion, with only the remaining ventral tissue giving rise to the embryo itself (26–28). Like

the amnioserosa, the serosa and the amnion are proposed to support the embryo during

development, but are thought to degenerate prior to hatching and not contribute to any

larval or adult structures (29,30).

Drosophila and *Tribolium* also exhibit dramatic differences in the morphogenetic

events occurring during early development (Fig 1). When GBE occurs in *Drosophila*, the

germband stays at the surface of the egg and the amnioserosa largely remains in place. In

Tribolium, on the other hand, germband extension begins with a complicated process called

embryo formation (see Fig 1; for a detailed description see (18,31)). During this process, the

presumptive embryonic ectoderm condenses towards the ventral side of the egg via both

mediolateral cell intercalation and a cuboidal-to-columnar cell shape transition.

Simultaneously, epithelial folding and tissue involution occurs, causing the presumptive

amnion to fold over the embryo to form the multi-layered germband. During these

movements, the serosa cells undergo a cuboidal-to-squamous transition and spread over the

entire egg surface.

The differences in fatemap and tissue folding described above show that both

fatemap shifts and reductions in early morphogenetic events have contributed to the

evolution of the long germ mode of development. Understanding how these differences

evolved is integral to understanding the short-to-long germ transition, but in order to study

how this occurred, we first need to understand how these tissues develop in each species.

The form and function of the *Tribolium* serosa has been analysed in several studies

(25,32,33), and comparative research in several fly species has revealed how the

loss/reduction of the serosa in flies may have occurred (reviewed in (29)). The amnion, on

the other hand, has proven harder to analyse, and the precise embryo/amnion boundary at

the blastoderm stage is unknown. However, a clear boundary between embryo and amnion

has been proposed to exist at the germband stage (Fig 1) (26). Cells in the ventral half of the

germband are thought to give rise to all embryonic structures, while cells in the dorsal half of

the germband are thought to form the amnion (28,34,35). This germband structure has been

described in many insects over the past century and is proposed to represent the core

conserved structure of short/intermediate germ embryos (reviewed in (2,36,37)). However,

the proposed boundary between cells fated to become embryo and those fated to become

amnion has not been directly tested.

I here set out to investigate the development of the presumptive amnion in *Tribolium*

using a combination of fluorescent live imaging and fate mapping techniques. To my great

surprise, I find that the majority of the cells previously described as 'amnion' actually form

large parts of the embryo proper. Using fate-mapping experiments, I show that true

'amnion' cells originate from a very small domain of the blastoderm, just as the *Drosophila*

amnioserosa cells do. I also show that the movement of cells from the 'amnion' side of the

germband to the 'embryo' side of the germband occurs via the large scale flow of the

ectodermal epithelium. Lastly, I describe the underlying causes of this flow, and show how

this tissue movement is likely homologous to the dorsal-to-ventral tissue flow that occurs

during *Drosophila* GBE.

This discovery forces a major shift in our view of development in *Tribolium* (and of

short/intermediate germ insects in general) and demonstrates that there is no qualitative

difference in germband tissue structure between *Tribolium* and *Drosophila*. Furthermore,

this discovery has significant consequences for how we must study short/intermediate germ

development going forward, as what was previously thought to be amnion (but is actually a

large region of the embryo) is routinely discarded or ignored during sample preparation.

Results

133 **Live cell tracking reveals movement of 'amnion' cells into the embryo**

134 To examine the development of the *Tribolium* presumptive amnion in detail, I carried out

135 high resolution live imaging of embryos transiently labelled (18) with a fluorescent histone

136 marker (H2B-venus) to label nuclei. My goal was to track presumptive amnion cells from the

137 blastoderm stage onwards. However, it was not possible to accurately track the majority of

138 cells throughout embryo condensation and GBE, due to the extensive morphogenetic

139 rearrangements that take place during this process. Instead, I focused on the stage

140 immediately following condensation, when the multi-layered germband has formed, and the

141 embryonic region where the presumptive amnion is closest to the surface of the egg.

142 Specifically, I tracked over 200 presumptive amnion cells in the central region of the embryo

143 from the closure of the serosa window until after the formation of the thoracic segments

144 (over 11 hours of development; Fig 2 Supplemental Movie 2). As previously described (18),

145 the germband and yolk exhibit pulsatile movements during this period, as well as rotating

146 within the serosa (Supplemental Movie 1).

147 The presumptive amnion initially consisted of many tightly packed cells that became

148 increasingly spread out during GBE (Supplemental Movie 2, Fig 2(A-C)). However, rather than

149 remaining restricted to the 'amnion territory', many of the tracked cells moved around the

150 edge of the germband in to the 'embryo territory'. Differential labelling of tracked cells

151 clearly showed that these cells that moved around the germband edge became part of the

152 embryo proper (Supplemental Movie 2 and Fig 2(A-C)). The cells that joined the 'embryo

153 territory' became tightly packed, continued to divide, and formed embryonic structures

154 (Supplemental Movie 3 and Fig 2(D-F)). In contrast, cells that remained in the 'amnion

155 territory' became squamous and stopped dividing. The nuclei of these latter cells became

156 enlarged (Supplemental Movie 3 and Fig 2(D-F)), suggesting that they underwent

endoreplication to become polyploid, as seen in the *Tribolium* serosa and in the *Drosophila*

amnioserosa (22,27). In addition, several germband nuclei underwent apoptosis

(Supplemental Movie 3) as has been described in fixed embryos (38). These results show

that many of the cells previously thought to be extraembryonic amnion give rise to

embryonic structures. As such, it is not accurate for the entire epithelium to be called

'amnion' and I will instead refer to it as the 'dorsal epithelium' as this region is made up of

dorsal/dorsolateral ectoderm.

Differential cell labelling confirms widespread dorsal-to-ventral cell movement

My next question was whether the movement of cells from the dorsal epithelium to the

ventral epithelium occurs throughout the AP axis or is just limited to the thoracic region. The

extensive movements of the germband made it difficult to track individual cells accurately at

the anterior and posterior poles. To overcome this problem, I combined differential cell

labelling with long term fluorescent live imaging to follow small groups of nuclei throughout

development. Specifically, I microinjected mRNA encoding a nuclear-localised

photoconvertable fluorescent protein (NLS-tdEos) into pre-blastoderm embryos to uniformly

label all nuclei, then photoconverted a small patch of nuclei at different positions along the

AP axis at the final uniform blastoderm stage. I then performed long term confocal live

imaging of both the unconverted and photoconverted forms of the fluorescent protein

throughout the period of GBE (or longer). Unlike that of *Drosophila*, the *Tribolium* egg shell

does not show any dorsoventral (DV) polarity, and I was therefore unable to specifically

target particular locations along the DV axis. Instead, I opted for a brute-force approach and

performed the photoconversion experiment for 50-150 embryos at each AP position (75%

egg length [EL] from the posterior pole, 50% EL, 25% EL, and close to the posterior pole), and

then used the subsequent live imaging to determine the DV position of the photoconverted

cells. Using a new live imaging set up (see Materials and Methods), I obtained the same

range of hatching rates as I typically obtain for other microinjection experiments

(approximately 80%, REF), even after continuous confocal live imaging for almost the

entirety of *Tribolium* embryonic development (3.5 days; Supplemental Movie 4). Both

unconverted and photoconverted protein persisted throughout germband extension and

retraction, although fluorescent signal faded over time. I have included various examples

from this data set in Supplemental Figures 1-3. In addition, I have made the raw confocal

data for a large number of timelapses available online (>300 embryos, >700 GB of data,

[upload in progress at time of submission]) for the benefit of the community. This data will

likely prove valuable for a wide range of research projects.

During my live imaging, ectodermal cell clones became elongated along the AP axis

over time, as previously reported in a *Tribolium* study that used a non-live imaging cell clone

method (19). However, Nakamoto and colleagues stated that "labelled ectodermal cells …

rarely mix with unlabelled cells" even as clones became greatly elongated. In contrast, I

frequently observed non-converted nuclei in the midst of labelled nuclei (Fig 3(A-F),

Supplemental Figures 1-3), confirming that extensive mediolateral cell intercalation occurs

throughout GBE. This difference in clone behaviour between the two approaches may be

due to differences in the size of the initial group of photoconverted cells, or in the resolution

of the subsequent imaging.

When I examined clones initially located in the dorsal epithelium, I found that

movement of cells from the dorsal epithelium to the ventral epithelium occurred throughout

the posterior of the embryo during GBE (Fig 3(A-F), Supplemental Movie 5). I also observed

the same movements at the anterior of the germband (Fig 3(G-J)), although I have focused

my analysis on the middle and posterior parts of the embryo. Together with the cell tracking

data, these results show that most of what was previously thought to be 'amnion' is in fact

embryonic tissue, and that cells move from the dorsal epithelium to the ventral epithelium

throughout the germband (summarised in Fig 3(K,L))

Support for these findings also comes from comparing the expression pattern of the

transcription factor Caudal in both *Tribolium* and *Drosophila*. In both species, this gene is

initially expressed in a broad posterior domain that then retracts along the AP axis (39,40). In

Drosophila, expression persists for longer in dorsal tissue than ventral tissue (9). Similar

dorsoventral dynamics, although not previously remarked upon, are also evident in

Tribolium; *Tc-cad* expression does indeed persist longer in the dorsal epithelium than in the

ventral epithelium (Supplemental Fig 4).

***Tribolium serpent* may mark true 'amnion'**

As described earlier, cells that remained in the dorsal epithelium became squamous, and this

cell shape change occurred progressively along the AP axis (Fig 4(A)). This change in cell

shape may be a sign of maturation of true 'amnion'. While characterising cell fate markers, I

found that the *Tribolium* ortholog of the GATA factor *serpent* (*Tc-srp*) exhibited spatial and

temporal expression dynamics that were very similar to those of the potential 'amnion' (Fig

4(B), Supplemental Fig 5). At the end of GBE, all but the posterior-most cells of the dorsal

epithelium were squamous (data not shown) and *Tc-srp* seemed to be expressed in dorsal

epithelium cells along nearly the full length of the germband (Supplemental Fig 5(I_2)).

However, this latter finding was difficult to confirm as most of the dorsal epithelium is lost

during embryo fixation at this embryonic stage (presumably due to the fragility of the

tissue). I also found *Tc-srp* to be expressed in several other domains, including in the

presumptive endoderm (Supplemental Fig 5). In *Drosophila*, *serpent* is also expressed in

extraembryonic tissue (the amnioserosa) (41–44), and, therefore, *Tc-srp* may mark 'true'

extraembryonic amnion. However, future work is required to confirm whether this putative

amnion degenerates prior to hatching (as is required to be defined as extraembryonic). For

simplicity, I will refer to this tissue as 'amnion' for the remainder of this text.

A revised *Tribolium* fatemap

To determine which blastoderm cells give rise to the amnion, I analysed 85 embryos in which

the dorsal and dorsolateral blastoderm cells were labelled by NLStdEos photoconversion as

described above. I found that amnion cells arose from a very small domain of dorsal-most

cells (that tapers from its anterior to posterior extent) and from a narrow strip of cells

between the presumptive embryo and presumptive serosa (summarised in Fig 5 and

Supplemental Fig 6). At 50% EL, only approximately the 6 dorsal-most cells (approximately

8% of the circumference of the blastoderm) gave rise to all amnion cells stretching from one

side of the thorax to the other (Fig 4(C), Supplemental Fig 1). Nearer to the posterior of the

blastoderm (25% EL), even fewer cells gave rise to amnion (approximately 3 of the most

dorsal cells; approximately 6% of the circumference; Fig 4(D), Supplemental Fig 2). The

posterior limit of the amnion was difficult to define, as although some cells from

approximately 5-10% EL appeared to become amnion (Fig 4(E)), these cells condensed

posteriorly towards the hindgut during germband retraction, and might have contributed to

the hindgut tissue (Supplemental Fig 3, Supplemental movie 6). I was unable to

unambiguously determine the fate of these cells. At the anterior of the embryo, I found that

a narrow strip of 1-2 cells between the presumptive embryo and presumptive serosa also

gave rise to amnion (Fig 3(G-J)). Substantial additional work is required to define a complete

blastoderm fatemap for *Tribolium*, however, my findings show that the 'amnion' domain is

drastically smaller than previously proposed.

Discussion

In this article, I have shown that the majority of cells that were previously thought to be

extraembryonic amnion actually give rise to embryonic tissue. The movement of cells from

the dorsal side of the germband to the ventral side was visible in live cell tracking and

differential cell labelling experiments. My results also show that the true amnion region

appears to differentiate progressively along the AP axis during GBE, as shown by differences

in cell behaviour and the expression of the gene *Tc-srp*. Lastly, the cells that give rise to this

presumptive amnion predominantly originate from a small domain on the dorsal side of the

blastoderm.

A revised understanding of the short germ embryo

The revision to the blastoderm fatemap is essentially a quantitative shift in our

understanding of where cell fate boundaries lie along the DV axis. In the revised *Tribolium*

blastoderm fatemap (Fig 5, Supplemental Fig 6), the proportion of the blastoderm that gives

rise to the presumptive amnion is much smaller than previously thought. The presumptive

amnion domain is, therefore, remarkably similar in size to the amnioserosa domain of the

Drosophila blastoderm fatemap (22). However, it is important to recognize that fatemaps

such as those presented here show a static picture of a dynamic process. There is no

evidence that the presumptive amnion is specified at the blastoderm stage in *Tribolium*.

Instead, the progressive changes in cell shape and *Tc-srp* expression in the dorsal epithelium

of the germband suggest that the amnion is specified progressively along the AP axis during

GBE. Progressive specification of DV cell fates during GBE fits with previous hypotheses

(34,45), and analysis of how this process occurs represents an exciting avenue of future

research (possible mechanisms are discussed more later).

In contrast to the fatemap revision, the observation that cells move from the dorsal

half of the germband to the ventral half of the germband represents a qualitative shift in our

understanding of development in short/intermediate germ insects. In the classic model of

short/intermediate germ development, the germband was thought of as a more-or-less flat

sheet of ectodermal cells (with mesoderm underneath) covered dorsally by the

extraembryonic amnion. Because of this, the entire dorsal epithelium is routinely removed

during embryo preparation, or not included in descriptions of gene expression patterns and

embryonic phenotypes. Based on the new data presented here, it is obvious that we have

been discarding or ignoring large parts of the embryo. Furthermore, the movement of cells

from the dorsal epithelium into the ventral epithelium must be contributing to GBE, and is,

therefore, a key aspect of the extension and overall development of the germband that has

thus far been missed.

The revised model of the germband does present some technical challenges for

future work on short/intermediate germ embryo. The flattened geometry of the germband

makes it difficult to image both the dorsal and ventral epithelium using bright-field

microscopy approaches. However, this problem can be overcome either by using

fluorescence based techniques and confocal microscopy or by mechanical sectioning of the

germband. Both approaches have been shown to work well in *Tribolium* (for example see

(7,46) and results in this manuscript). In the rest of this article, I discuss why the revised

fatemap and cell flow accord well with what we know about *Tribolium* development, and

outline the implications of this discovery on our understanding of the evolution of insect

development.

The cellular and molecular causes of tissue flow unify the blastoderm and the germband

The revised model of the *Tribolium* germband reconciles the blastoderm and germband

stages. The ectoderm of the germband is a continuous epithelium, which means that the

movement of cells from the dorsal epithelium to the ventral epithelium occurs as a tissue

level 'flow'. Such dorsal-to-ventral tissue flow also occurs during embryo condensation in

Tribolium (18), and I propose that the flow is caused by largely the same morphogenetic

processes at both stages. The evidence for this hypothesis is summarised here, but for an

extended discussion see Appendix section 1.

Three morphogenetic processes contribute to dorsal-to-ventral cell flow in *Tribolium*,

and at least two of the three occur at both the blastoderm and germband stages. First,

mediolateral cell intercalation occurs at both stages and causes tissue-wide convergence

(along the DV axis) and extension (along the AP axis). This process requires two *Toll* genes

that are expressed in rings around the entire blastoderm and germband epithelium (21).

Second, tissue specific cell shape changes occur at both stages such that ventral/lateral cells

become columnar and dorsal/dorsolateral cells become thinner (during condensation

(Supplemental Fig 7)) or squamous (during GBE). The tissue level effect of these changes is

contraction of the ventral/lateral tissue and spreading of the dorsal tissue. The flattening of

dorsal/dorsolateral cells is likely regulated by BMP signalling, as not only does BMP activity

correlate with the cell shape changes (see Appendix section 1), but functional disruption of

BMP signalling components leads to uniform cell shape changes along the DV axis (28,47). A

third major morphogenetic event is gastrulation of the mesoderm. This occurs along the

ventral midline, and as gastrulation occurs, the ectoderm moves ventrally to seal the gap left

in the epithelium (46). At the stage when a complete germband has formed, gastrulation is

complete along most of the embryo. However, current data suggests mesoderm gastrulation

may be ongoing in the segment addition zone (46). If true, the ongoing invagination would

contribute to tissue flow in this region.

It is important to note that while each of the events described here is involved in the

dorsal-to-ventral tissue flow, no single event is absolutely required for it. In the absence of

cell intercalation, embryo condensation and thinning of dorsal/dorsolateral ectoderm still

takes place yielding abnormally wide and short germbands (21). In the absence of tissue

specific cell shape changes, condensation occurs in a more radially symmetrical manner

yielding a tube-shaped germband that undergoes segment specification and convergent

extension (28,47). Finally, both condensation and GBE are only mildly affected in the

absence of mesoderm specification (48). This functional independence comes from each of

the three process being specified by different pathways (intercalation via segment

specification, dorsal thinning via dorsal tissue specification, and gastrulation via ventral

tissue specification). There may be further, as yet undiscovered morphogenetic events that

also contribute to the dorsal-to-ventral tissue flow.

Reconciling long and short germ development

I propose that the dorsal-to-ventral tissue flow occurring during embryo condensation and

GBE in *Tribolium* is homologous to the dorsal-to-ventral tissue flow that occurs during

gastrulation and GBE in *Drosophila* (Fig 1). This conclusion is based on the flow being driven

by a conserved set of morphogenetic events.

As described above, tissue flow in *Tribolium* is caused by (1) mediolateral cell

intercalation, (2) tissue specific cell shape changes along the DV axis, and (3) gastrulation at

the ventral side of the embryo. Equivalent processes are all observed in *Drosophila* as well.

In *Drosophila*, *Toll*-mediated mediolateral cell intercalation causes the tissue wide

convergence (along the DV axis) and extension (along the AP axis) of the ectoderm during

GBE (20). As in *Tribolium*, the periodic expression of the *Toll* genes is driven by the pair-rule

genes. Conservation at the level of tissue identity, morphogenetic process, and molecular

control strongly suggest *Toll*-mediated cell intercalation to be homologous.

Cell shape changes are harder to compare between these species, as unlike most

insects, cellularisation in *Drosophila* leads to the direct formation of columnar cells (22,49).

However, tissue specific cell shape changes along the DV axis do occur and are dependent on

BMP signalling ((50,51); for a detailed description see Appendix section 2). While the

intracellular effectors of these cell shape changes are unknown, the dorsal patterning

function of BMP signalling is homologous in *Drosophila* and *Tribolium*, and many dorsal cell

specification genes are conserved between these two species.

Last, *Drosophila* mesoderm gastrulation also occurs along the ventral midline and it

causes lateral/dorsolateral ectoderm to move ventrally (50). As above, the intracellular

effectors of *Tribolium* mesoderm gastrulation are unknown, but the upstream patterning

events and the tissue specification genes are highly conserved (34,48). Furthermore,

gastrulation of mesoderm at the ventral region is widely conserved within the insects, and is

undoubtedly a homologous process in each species (52).

The conservation of the tissue-level flow and the underlying processes shows that

development in *Drosophila* and *Tribolium*, which are themselves models for long and short

germ development respectively, are far more similar than previously thought. This similarity

is further emphasised when the *Tribolium* germband is represented as a cylinder rather than

its normal *in vivo* flattened shape (Fig 6). This finding helps explain how the long germ mode

of development has evolved independently many times, as it takes fewer evolutionary

'steps' to go from one form to another than previously thought.

While I have focused on *Tribolium* and *Drosophila* here, evidence exists that my new

findings in *Tribolium* also apply to other short/intermediate germ insects. For example, in

the intermediate germ bug *Oncopeltus fasciatus*, the dorsal epithelium of the germband

initially consists of a thick epithelium that then progressively becomes squamous late during

GBE ((53) and data not shown). These tissue specific cell shape changes are likely the same

as those occurring during *Tribolium* GBE. Furthermore, *Oncopeltus* pair-rule genes, Loto *Toll*

genes and even segment polarity genes are expressed in rings around the entire germband

prior to thinning of the dorsal epithelium (shown but not described in (21,54,55)). The

expression of these genes in the dorsal epithelium provides additional evidence that much of

the *Oncopeltus* dorsal epithelium is made up of embryonic tissue. Future analyses of the

molecular and morphogenetic drivers of GBE need to analyse the entire germband, rather

than focusing on the ventral half.

The *Drosophila* posterior gut fold may represent an evolutionary remnant of short germ

development

Despite the deep conservation in morphogenetic events described above, a very obvious

difference exists between the germbands of *Drosophila* and *Tribolium*. While the *Drosophila*

germband develops mostly on the egg surface with the apical side of the epithelium facing

outwards, the epithelium of the *Tribolium* germband folds onto itself such that the apical

side faces inwards (Fig 1). The last common ancestor of *Tribolium* and *Drosophila* almost

certainly exhibited short/intermediate germ development (56), which would include the

formation of a bi-layered germband like that of *Tribolium*. As such, in the lineage leading to

Drosophila, the epithelial folding that previously generated the bi-layered germband must

have become heavily reduced during evolution.

I propose that remnants of the folding seen in *Tribolium* still exists in *Drosophila*

today, in what is now termed the "posterior gut fold". The epithelial folding that causes the

formation of the bi-layered germband of *Tribolium* initiates with a deep infolding of tissue at

the posterior of the embryo (classically termed the posterior amniotic fold) (18,27).

Epithelial infolding also occurs at the posterior of the *Drosophila* blastoderm during posterior

gut formation (57), and the similarities between these folds have been pointed out before

(27). However, the evolutionary relationship of these two structures has been unclear due to

the belief that the dorsal half of the fold in *Tribolium* gave rise to the amnion. Now, with the

revised fatemap, it is clear that these folds form from the same region of the embryo.

Therefore, I propose that the epithelial fold that forms during posterior gut formation in

Drosophila is homologous to the posterior epithelial fold that forms during embryo

condensation in *Tribolium*. This hypothesis is supported by two pieces of molecular data.

First, a recent report found that a gene essential for hindgut development (the *Tribolium*

ortholog of *senseless*) is already expressed at the posterior pole of the *Tribolium* embryo at

the blastoderm stage (58). Second, the *Tribolium* posterior fold is controlled at the

morphogenetic level by the Fog signalling pathway, just like it is in *Drosophila* (Frey et al.

manuscript in preparation).

Due to this revised evolutionary relationship and the new *Tribolium* fatemap, I

propose that the posterior fold in *Tribolium* be renamed from 'posterior amniotic fold' to

'posterior fold'.

New insights in tissue patterning

While I have predominantly focused on morphogenetic events in this manuscript, the

revised model of short germ embryo development also has impacts for how we understand

tissue patterning to occur in *Tribolium*. Here, I present one example of this, but for

descriptions of pair-rule gene expression, posterior terminal gene expression, and of how DV

patterning may be occurring in the germband, see Appendix section 3.

Segment polarity genes are expressed in a single stripe in each segment, extending

from near the ventral midline of the embryo to the lateral edge of the germband (59–61).

Under the classic model, these domains appear to be static. However, the revised model

shows that cells at the lateral edges of the germband move ventrally as cells from the dorsal

epithelium move around the edges of the germband. This means that for each stripe of a

segment polarity gene to 'stay' at the edge of the germband, cells at the correct AP location

must activate segment polarity gene expression when they move from the dorsal epithelium

to the ventral. This can be thought of like Lewis Carroll's Red Queen's race, where "it takes

all the running you can do, to keep in the same place" (62). These dynamics mean that each

expression domain spreads from ventral-to-dorsal across the ectoderm, but that this occurs

simultaneously with the dorsal-to-ventral flow of tissue. A similar ventral-to-dorsal activation

of segment polarity gene expression can be seen in the *Drosophila* ectoderm (Supplemental

Fig 8), indicating that the DV position of expression is controlled by a conserved upstream

mechanism.

While I have only described a handful of examples in this manuscript, other genes are

likely to have important expression domains in both the ventral and dorsal epithelia of the

germband. Accurately characterising these domains is necessary for understanding how

these genes function in *Tribolium* and for elucidating how their functions may have evolved.

Conclusions

The deep conservation of morphogenetic events that I am proposing here may seem

unlikely, but I would argue that such conservation should have been expected from the

beginning. The vast majority of morphogenetic events are controlled by upstream patterning

networks. As such, morphogenesis can be viewed as a physical readout of such networks.

Decades of research has shown that many of the key patterning genes that function during

germband formation and extension in *Drosophila* have conserved roles in *Tribolium* and

other insects. As such, without evidence to the contrary, conservation of the cellular events

driven by these patterning genes makes perfect sense.

Materials and methods

Tribolium animal husbandry, egg collection, and RNA *in situ* hybridisation was performed as

previously described (21). The *Tc-srp* ortholog was previously described (63) and was cloned

into pGEM-t (Promega Reference A1360) with primers TCCCGCTGCTTTGATCTAGT and

TGCGATGACTGTGACGTGTA. The *Tc-cad* ortholog was as previously used (18).

The *H2B-ven* fusion was created by fusing the *D. melanogaster* histone *H2B* coding

sequence (without the stop codon) from the published H2B-RFP (18) to the *venus*

fluorescent protein (64) and cloning into the pSP64 Poly(A) (Promega Reference P1241)

expression vector. The *NLS-tdEos* fusion was kindly provided by Matthias Pechmann.

Additional details and both plasmids are available upon request to M. Pechmann or myself.

Capped mRNA synthesis was performed as previously described (18). *H2B-ven* capped mRNA

was injected at 1 µg/µL, *NLS-tdEos* capped mRNA was injected at 2-3 µg/µL.

469   Embryo microinjection was performed as previously described (18), with the

following changes. Up to 100 dechorionated embryos were mounted on a rectangular

coverslip (24 mm by 50 mm) that rested on a microscope slide. Water was allowed to dry off

the embryos before they were covered in Voltalef 10S halocarbon oil and injected as usual.

The coverslip (still resting on the slide) was then placed in a petri-dish (92 mm) containing a

base layer of 1% agarose (dissolved in water) and placed at 30-32°C until the embryos were

at the appropriate stage for imaging. The coverslip was then removed from the slide,

inverted (so that embryos were face down), and quickly but gently placed on a lumox dish

(50 mm; Sarstedt Reference 94.6077.410) that was sitting upside down. The corners of the

coverslip rested on the raised plastic lip of the dish such that the membrane and embryos

were close to each other but not touching. To ensure lateral stability of the coverslip during

the timelapse recording, approximately 5-10 µL of heptane glue (made by soaking parcel

tape in heptane) was placed at each corner. Additional Voltalef 10S halocarbon oil was then

added to fill any remaining space between the coverslip and the oxygen permeable

membrane. This contraption was then stuck to a microscope slide (using double sided tape)

for imaging on an upright microscope. This last step may be unnecessary depending on the

microscope stage and orientation.

486   Live imaging was performed on an upright Zeiss SP8 confocal microscope equipped

with Hybrid detectors in the Biocentre Imaging facility (University of Cologne). Image stacks

of 15-50 focal planes with z-steps ranging from 2-10 µm were taken with a 10x/0.3NA dry

objective or a 20×/0.7NA multi-immersion objective at intervals of 5-45 minutes. The

temperature of the sample during imaging could not be carefully regulated, but was typically

between 25-28 degrees. While this lack of temperature control is not ideal, it does not affect

the findings presented in this manuscript.

Photoconversion of NLS-tdEos protein was performed by constantly scanning the

region of interest for 20-30 seconds with the 405 wavelength laser at low power (5%). These

settings were manually determined on the above microscope, and need to be determined

independently on different systems. Photoconversions were performed during the final

uniform blastoderm stage, as photoconversion prior to this resulted in substantial diffusion

of the photoconverted protein during nuclei division. The positions of the different regions

of the embryo (75% EL etc.) were determined by measuring the length of each embryo in the

LASX software and selecting the appropriate region. Photoconversions were performed on

all embryos on the coverslip before setting up the timelapse, which led to a 0.5-2 hour delay

between performing the photoconversion and beginning the timelapse. As such, the

positions of the photoconverted region at the first time point in the timelapses in this

manuscript do not reflect the original region of photoconversion.

Imaging of fixed material was performed on an upright Zeiss SP8 confocal, an upright

Zeiss SP5 confocal microscope and an inverted Zeiss SP5 confocal microscope. The

Drosophila gooseberry expression patterns were kindly provided by Erik Clark and acquired

as in (65). Images and timelapses were analysed using FIJI (66) and Photoshop CS5. Manual

cell tracking was performed on confocal hyperstacks with MTrackJ (67). The figures were

arranged and the schematics created using Inkscape.

Acknowledgements

I would like to thank E. Clark for many stimulating discussions, extensive feedback on the

manuscript, and for providing the *Drosophila* images. I thank M. Pechmann for providing the

NLS-tdEos construct and S. Roth for supporting me during this project. In addition, I thank M.

Akam, M. Pechmann and S. Roth for comments on the manuscript.

References

1. Grimaldi D, Engel M. Evolution of the Insects. Cambridge University Press; 2005.

2. Anderson DT. Embryology and Phylogeny in Annelids and Arthropod. Pergamon Press;

1973.

3. Jaeger J. The gap gene network. Cell Mol Life Sci [Internet]. 2011 Jan [cited 2013 Mar

8];68(2):243–74. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3016493&tool=pmcentr

ez&rendertype=abstract

4. Davis GK, Patel NH. SHORT, LONG, AND BEYOND: Molecular and Embryological

Approaches to. Annu Rev Entomol. 2002;(47):669–99.

5. Peel AD, Chipman AD, Akam M. Arthropod segmentation: beyond the Drosophila

paradigm. Nat Rev Genet [Internet]. 2005 Dec [cited 2010 Jul 6];6(12):905–16.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/16341071

6. Choe CP, Miller SC, Brown SJ. A pair-rule gene circuit defines segments sequentially in

the short-germ insect Tribolium castaneum. Proc Natl Acad Sci U S A [Internet]. 2006

Apr 25;103(17):6560–4. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1564201&tool=pmcentr

ez&rendertype=abstract

7.   Sarrazin AF, Peel AD, Averof M. A segmentation clock with two-segment periodicity in

insects. Science [Internet]. 2012 Apr 20 [cited 2012 Oct 5];336(6079):338–41.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/22403177

8.   Clark E. Dynamic patterning by the Drosophila pair-rule network reconciles long-germ

and short-germ segmentation. PLOS Biol [Internet]. 2017 Sep 27;15(9):e2002439.

Available from: https://doi.org/10.1371/journal.pbio.2002439

9.   Clark E, Peel AD. Evidence for the temporal regulation of insect segmentation by a

conserved set of developmental transcription factors. bioRxiv [Internet]. 2017 Jan 1;

Available from: http://biorxiv.org/content/early/2017/06/02/145151.abstract

10. Zhu X, Rudolf H, Healey L, François P, Brown SJ, Klingler M, et al. Speed regulation of

genetic cascades allows for evolvability in the body plan specification of insects. Proc

Natl Acad Sci [Internet]. 2017 Oct 10;114(41):E8646–55. Available from:

http://www.pnas.org/content/114/41/E8646.abstract

11. Irvine KD, Wieschaus E. Cell intercalation during Drosophila germband extension and

its regulation by pair-rule segmentation genes. Development [Internet]. 1994

Apr;120(4):827–41. Available from: http://www.ncbi.nlm.nih.gov/pubmed/7600960

12. Bertet C, Sulak L, Lecuit T. Myosin-dependent junction remodelling controls planar cell

intercalation and axis elongation. Nature [Internet]. 2004 Jun 10;429(6992):667–71.

Available from: http://dx.doi.org/10.1038/nature02590

13. Collinet C, Rauzi M, Lenne P, Lecuit T. Local and tissue-scale forces drive oriented

junction growth during tissue extension. Nat Cell Biol [Internet]. 2015 Sep

21;17(10):1247–58. Available from:

http://www.nature.com/doifinder/10.1038/ncb3226

14. Butler LC, Blanchard GB, Kabla AJ, Lawrence NJ, Welchman DP, Mahadevan L, et al.

Cell shape changes indicate a role for extrinsic tensile forces in Drosophila germ-band

extension. Nat Cell Biol. 2009 Jul;11(7):859–64.

15.   Firmino AAP, Fonseca FC de A, de Macedo LLP, Coelho RR, Antonino de Souza Jr JD,

Togawa RC, et al. Transcriptome Analysis in Cotton Boll Weevil (Anthonomus grandis)

and RNA Interference in Insect Pests. PLoS One [Internet]. 2013;8(12):e85079.

Available from: http://dx.plos.org/10.1371/journal.pone.0085079

16.   da Silva SM, Vincent J-P. Oriented cell divisions in the extending germband of

Drosophila Development [Internet]. 2007 Sep 1;134(17):3049

LP-3054. Available from: http://dev.biologists.org/content/134/17/3049.abstract

17.   Liu PZ, Kaufman TC. Short and long germ segmentation: unanswered questions in the

evolution of a developmental mode. Evol Dev [Internet]. 2005;7(6):629–46. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/16336416

18.   Benton MA, Akam M, Pavlopoulos A. Cell and tissue dynamics during Tribolium

embryogenesis revealed by versatile fluorescence labeling approaches. Development

[Internet]. 2013 Jul 16 [cited 2013 Jul 16];140(15):3210–20. Available from:

http://dev.biologists.org/cgi/doi/10.1242/dev.096271

19.   Nakamoto A, Hester SD, Constantinou SJ, Blaine WG, Tewksbury AB, Matei MT, et al.

Changing cell behaviours during beetle embryogenesis correlates with slowing of

segmentation. Nat Commun [Internet]. 2015;6:6635. Available from:

http://www.nature.com.libproxy.helsinki.fi/ncomms/2015/150410/ncomms7635/full

/ncomms7635.html

20.   Paré AC, Vichas A, Fincher CT, Mirman Z, Farrell DL, Mainieri A, et al. A positional Toll

receptor code directs convergent extension in Drosophila. Nature [Internet]. 2014 Nov

2;515(7528):523–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25363762

21.   Benton MAA, Pechmann M, Frey N, Stappert D, Conrads KHKH, Chen Y-TY-T, et al. Toll

Genes Have an Ancestral Role in Axis Elongation. Curr Biol [Internet].

2016;26(12):1609–15. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0960982216304092

22.   Campos-Ortega J a, Hartenstein V. The embryonic development of Drosophila

melanogaster. 2nd ed. Springer-Verlag Berlin Heidelberg; 1997.

23.   Schönauer A, Paese CLB, Hilbrant M, Leite DJ, Schwager EE, Feitosa NM, et al. The Wnt

and Delta-Notch signalling pathways interact to direct pair-rule gene expression via

caudal during segment addition in the spider Parasteatoda tepidariorum.

Development [Internet]. 2016;2455–63. Available from:

http://dev.biologists.org/content/early/2016/06/09/dev.131656.abstract

24.   Janssen R. Gene expression suggests double-segmental and single-segmental

patterning mechanisms during posterior segment addition in the beetle Tribolium

castaneum. Int J Dev Biol. 2014;58(5).

25.   van der Zee M, Berns N, Roth S. Distinct Functions of the Tribolium zerknullt Genes in

Serosa Specification and Dorsal Closure. Curr Biol. 2005;15:624–36.

26.   Falciani F, Hausdorf B, Schröder R, Akam M, Tautz D, Denell R, et al. Class 3 Hox genes

in insects and the origin of zen. Proc Natl Acad Sci USA. 1996;93:8479–84.

27.   Handel K, Grünfelder CG, Roth S, Sander K. Tribolium embryogenesis: a SEM study of

cell shapes and movements from blastoderm to serosal closure. Dev Genes. 2000;

28.   Nunes da Fonseca R, von Levetzow C, Kalscheuer P, Basal A, van der Zee M, Roth S.

Self-Regulatory Circuits in Dorsoventral Axis Formation of the Short-Germ Beetle

Tribolium castaneum. Dev Cell [Internet]. 2008;14(4):605–15. Available from:

http://www.sciencedirect.com/science/article/pii/S1534580708001056

29.   Schmidt-Ott U, Kwan CW. Morphogenetic functions of extraembryonic membranes in

insects. Curr Opin Insect Sci [Internet]. 2016;13:86–92. Available from:

http://dx.doi.org/10.1016/j.cois.2016.01.009

30.   Panfilio K a. Extraembryonic development in insects and the acrobatics of

blastokinesis. Dev Biol [Internet]. 2008 Jan [cited 2011 Mar 10];313(2):471–91.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/18082679

31.   Benton MAA, Pavlopoulos A. Tribolium embryo morphogenesis: May the force be with

you. Bioarchitecture. 2014;4(1):16–21.

32.   Jacobs CGC, Spaink HP, van der Zee M. The extraembryonic serosa is a frontier

epithelium providing the insect egg with a full-range innate immune response. Elife

[Internet]. 2014;3:1–21. Available from:

http://elifesciences.org/lookup/doi/10.7554/eLife.04111

33.   Jacobs CGC, Van Der Zee M. Immune competence in insect eggs depends on the

extraembryonic serosa. Dev Comp Immunol. 2013;41(2).

34.   Lynch JA, Roth S. The evolution of dorsal-ventral patterning mechanisms in insects.

Genes Dev [Internet]. 2011 Jan 15 [cited 2011 Jan 18];25(2):107–18. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3022256&tool=pmcentr

ez&rendertype=abstract

35.   Horn T, Panfilio KA. Novel functions for Dorsocross in epithelial morphogenesis in the

beetle Tribolium castaneum. Development [Internet]. 2016;143(16):3002–11.

Available from: http://dev.biologists.org/lookup/doi/10.1242/dev.133280

36.   Anderson DT. The Development of Holometabolous Insects. In: Counce SJ, Waddinton

CH, editors. Developmental systems Insects, Vol 1. New York: Academic Press; 1972.

p. 165–242.

37.   Anderson DT. The Development of Hemimetabolous Insects. In: Counce SJ,

635   Waddington CH, editors. Developmental systems Insects, Vol 1. New York: Academic

636   Press; 1972. p. 95–163.

637   38.   Aranda M, Marques-Souza H, Bayer T, Tautz D. The role of the segmentation gene

638   hairy in Tribolium. Dev Genes Evol [Internet]. 2008 Sep [cited 2013 Mar

639   11];218(9):465–77. Available from:

640   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2522291&tool=pmcentr

641   ez&rendertype=abstract

642   39.   Schulz C, Schröder R, Hausdorf B, Wolff C, Tautz D. A caudal homologue in the short

643   germ band beetle Tribolium shows similarities to both, the Drosophila and the

644   vertebrate caudal expression patterns. Dev Genes Evol [Internet]. 1998 [cited 2012

645   Jan 20];208(5):283–9. Available from:

646   http://www.springerlink.com/index/u6epmj98hxfr9jmd.pdf

647   40.   Mlodzik M, Gehring WJ. Expression of the caudal gene in the germ line of Drosophila:

648   Formation of an RNA and protein gradient during early embryogenesis. Cell [Internet].

649   1987;48(3):465–78. Available from:

650   http://www.sciencedirect.com/science/article/pii/0092867487901978

651   41.   Tomancak P, Berman BP, Beaton A, Weiszmann R, Kwan E, Hartenstein V, et al. Global

652   analysis of patterns of gene expression during Drosophila embryogenesis. Genome

653   Biol. 2007;8(7):R145.

654   42.   Hammonds AS, Bristow CA, Fisher WW, Weiszmann R, Wu S, Hartenstein V, et al.

655   Spatial expression of transcription factors in Drosophila embryonic organ

656   development. Genome Biol. 2013 Dec;14(12):R140.

43.   Tomancak P, Beaton A, Weiszmann R, Kwan E, Shu S, Lewis SE, et al. Systematic

determination of patterns of gene expression during Drosophila embryogenesis.

Genome Biol. 2002;3(12):RESEARCH0088.

44. Sam S, Leise W, Hoshizaki DK. The serpent gene is necessary for progression through

the early stages of fat-body development. Mech Dev [Internet]. 1996 Dec;60(2):197–

205. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9025072

45. Sachs L, Chen YT, Drechsler A, Lynch JA, Panfilio KA, Lässig M, et al. Dynamic BMP

signaling polarized by Toll patterns the dorsoventral axis in a hemimetabolous insect.

Elife. 2015;4(MAY).

46. Handel K, Basal A, Fan X, Roth S. Tribolium castaneum twist: gastrulation and

mesoderm formation in a short-germ beetle. Dev Genes Evol. 2005;215:13–31.

47. van der Zee M, Stockhammer O, von Levetzow C, Nunes da Fonseca R, Roth S.

Sog/Chordin is required for ventral-to-dorsal Dpp/BMP transport and head formation

in a short germ insect. Proc Natl Acad Sci U S A [Internet]. 2006 Oct;103(44):16307–

12. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1637578&tool=pmcentr

ez&rendertype=abstract

48. Stappert D, Frey N, von Levetzow C, Roth S. Genome-wide identification of Tribolium

dorsoventral patterning genes. Development [Internet]. 2016 Jul 1;143(13):2443–54.

Available from: http://dev.biologists.org/lookup/doi/10.1242/dev.130641

49. van der Zee M, Benton MAA, Vazquez-Faci T, Lamers GEMEM, Jacobs CGCGC,

Rabouille C. Innexin7a forms junctions that stabilize the basal membrane during

cellularization of the blastoderm in Tribolium castaneum. Development [Internet].

2015;142(12):2173–83. Available from:

http://dev.biologists.org/cgi/doi/10.1242/dev.097113

50.   Rauzi M, Krzic U, Saunders TE, Krajnc M, Ziherl P, Hufnagel L, et al. Embryo-scale tissue

683   mechanics during Drosophila gastrulation movements. Nat Commun [Internet]. 2015

684   Oct 26;6:8677. Available from: http://dx.doi.org/10.1038/ncomms9677

685   51.   Leptin M, Grunewald B. Cell shape changes during gastrulation in Drosophila.

686   Development [Internet]. 1990 Sep 1;110(1):73 LP-84. Available from:

687   http://dev.biologists.org/content/110/1/73.abstract

688   52.   Roth S. Gastrulation in Other Insects. In: Stern CD, editor. Gastrulation From Cells to

689   Embryo. New York: Cold Spring Harbor Laboratory Press; 2004. p. 105–21.

690   53.   Ewen-Campen B, Jones TEM, Extavour CG. Evidence against a germ plasm in the

691   milkweed bug Oncopeltus fasciatus, a hemimetabolous insect. Biol Open [Internet].

692   2013;2(6):556–68. Available from:

693   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3683158&tool=pmcentr

694   ez&rendertype=abstract

695   54.   Erezyilmaz DF, Kelstrup HC, Riddiford LM. The nuclear receptor E75A has a novel pair-

696   rule-like function in patterning the milkweed bug, Oncopeltus fasciatus. Dev Biol

697   [Internet]. 2009 Oct 1 [cited 2013 May 1];334(1):300–10. Available from:

698   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2749522&tool=pmcentr

699   ez&rendertype=abstract

700   55.   Liu PZ, Kaufman TC. even-skipped is not a pair-rule gene but has segmental and gap-

701   like functions in Oncopeltus fasciatus, an intermediate germband insect. Development

702   [Internet]. 2005 May [cited 2013 Mar 7];132(9):2081–92. Available from:

703   http://www.ncbi.nlm.nih.gov/pubmed/15788450

704   56.   Peel AD. The evolution of developmental gene networks: lessons from comparative

705   studies on holometabolous insects. Philos Trans R Soc Lond B Biol Sci [Internet]. 2008

Apr 27 [cited 2012 Oct 9];363(1496):1539–47. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2615820&tool=pmcentr

ez&rendertype=abstract

57. Turner FR, Mahowald AP. Scanning Electron Microscopy of Drosophila Embryogenesis

II. Gastrulation and Segmentation. 1977;416:403–16.

58. Oberhofer G, Grossmann D, Siemanowski JL, Beissbarth T, Bucher G. Wnt/beta-

catenin signaling integrates patterning and metabolism of the insect growth zone.

Development [Internet]. 2014 Dec;141(24):4740–50. Available from:

http://dev.biologists.org/cgi/doi/10.1242/dev.112797

59. Nagy LM, Carroll S. Conservation of wingless patterning functions in the short-germ

embryos of Tribolium castaneum. Nature [Internet]. 1994;367(6462):460–3. Available

from: http://biomednet.com/db/medline/94150623

60. Brown SJ, Patel NH, Denell RE. Embryonic expression of the single Tribolium engrailed

homolog. Dev Genet [Internet]. 1994;15(1):7–18. Available from:

http://dx.doi.org/10.1002/dvg.1020150103

61. Savard J, Marques-Souza H, Aranda M, Tautz D. A segmentation gene in tribolium

produces a polycistronic mRNA that codes for multiple conserved peptides. Cell

[Internet]. 2006 Aug 11 [cited 2013 May 1];126(3):559–69. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/16901788

62. Carroll L. Alice's Adventures in Wonderland. New York: Macmillan; 1865.

63. Gillis WQ, Bowerman B a, Schneider SQ. The evolution of protostome GATA factors:

molecular phylogenetics, synteny, and intron/exon structure reveal orthologous

relationships. BMC Evol Biol [Internet]. 2008 Jan [cited 2012 Mar 4];8:112. Available

from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2383905&tool=pmcentr

ez&rendertype=abstract

64.   Nagai T, Ibata K, Park ES, Kubota M, Mikoshiba K, Miyawaki A. A variant of yellow

fluorescent protein with fast and efficient maturation for cell-biological applications.

Nat Biotechnol [Internet]. 2002 Jan 1;20:87. Available from:

http://dx.doi.org/10.1038/nbt0102-87

65.   Clark E, Akam M. Odd-paired controls frequency doubling in Drosophila segmentation

by altering the pair-rule gene regulatory network. Elife. 2016 Aug;5.

66.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods [Internet]. 2012 Jul

[cited 2013 Mar 4];9(7):676–82. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/22743772

67.   Meijering E, Dzyubachyk O, Smal I. Methods for cell and particle tracking. [Internet].

1st ed. Vol. 504, Methods in enzymology. United States: Elsevier Inc.; 2012 [cited 2013

Mar 4]. 183-200 p. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22264535

Fig 1. Schematics of development in *Drosophila* and *Tribolium*. The two left columns show

schematics of *Drosophila* embryos from the uniform blastoderm stage to the extended

germband stage. The right three columns show schematics of *Tribolium* embryos at

comparable stages. The schematics in the right-most column show dissected, flatmounted

embryos. Red arrows display cell/tissue movement. The question marks highlight two

regions where the tissue boundaries are unknown/undescribed. Several features have been

omitted, including the yolk, mesoderm gastrulation, anterior gut formation and appendage

formation. The *Drosophila* fatemap is based on data from (22) and the references therein.

Refer to text for additional details.

Fig 2. Live cell tracking reveals contribution of 'amnion' cells to embryonic tissue. (A-F)

Time series from fluorescent live imaging of a *Tribolium* embryo expressing H2B-venus. The

serosa nuclei located above the germband have been manually removed from these frames,

but left in the surrounding territory (arrowhead in (A+A')). (A'-C') show optical transverse

sections of the respective frame at the position shown by the dashed line (the surface of the

egg is to the left). In (A-C), all nuclei that lie in a region of the 'amnion territory' in (A) have

been tracked and differentially labelled depending on whether they become part of the

embryo (magenta; labels disappear when nuclei join the germband), become located at the

edge of the germband (yellow) or remain in the 'amnion territory' (cyan). In (D-F), a line of

nuclei that lie in the 'amnion territory' in (D) have been tracked and differentially labelled

depending on whether they become part of the embryo (coloured points; daughter cells are

labelled in same colour as parent) or remain in the 'amnion territory' (cyan; no division takes

place). Note that in panel (D), the orange spot is mostly hidden below the yellow spot

because the nuclei in that region are partially overlapping when viewed as projections. The

first frame of the timelapse was defined as timepoint 0. In (A-F), embryos are oriented based

on the AP/DV polarity of the egg with anterior to the left and dorsal to the top. (A-C) are

maximum intensity projections of one egg hemisphere. (D-F) are average intensity

projections of 46 microns to specifically show the germband. Scale bars are 50 μm.

Fig 3. Differential cell labelling reveals widespread movement of cells from the dorsal

epithelium to the ventral epithelium. (A-J) Time series from fluorescent live imaging of two

Tribolium embryos expressing NLS-tdEos showing unconverted protein (magenta) and

photoconverted protein (cyan). In (A-F') a patch of nuclei at the posterior-dorsal region of

the blastoderm were photoconverted. Panels (A-F) show the posterior region of the

germband during late GBE and panels (A'-F') show optical transverse sections made at the

position of the dashed line at each timepoint (roughly following the same nuclei). Serosa

nuclei are marked by white crosses in the transverse sections. In (G-J), a patch of nuclei at

the anterior-lateral region of the blastoderm were photoconverted. Panels (G-J) show the

anterior of the germband during condensation and GBE. In both embryos, all converted

nuclei are initially located in the dorsal epithelium, but most move into the ventral

epithelium during GBE. (K-L) Schematics showing the classic and revised models of the

Tribolium germband (presumptive amnion is shown in purple, presumptive embryo is shown

in grey, red arrows show the newly discovered tissue flow). The first frame of the timelapses

was defined as timepoint 0. In (A-J), embryos are oriented based on the AP/DV polarity of

the egg with anterior to the left and dorsal to the top. In (A'-F'), the surface of the egg is

oriented to the left. In (K-L), schematics show flatmounted germbands with the focus on the

dorsal epithelium, the anterior to the left and the orthogonal sections are oriented with the

dorsal half of the germand to the left. (A-J) are maximum intensity projections of one egg

hemisphere. Scale bars are 50 μm.

Fig 4. Development of the putative amnion. (A-A'') *Tribolium* embryo transiently expressing

the membrane marker GAP43-YFP. (A) shows an overview of the whole egg, (A') shows the

dorsal epithelium of the same embryo at the position of the white box, (A'') is an optical

sagittal section at the position of the dashed line in (A') showing the apical-basal height of

cells of the dorsal epithelium. (B) *Tc-srp* (red) expression in a flatmounted *Tribolium*

germband also showing nuclei (DAPI, blue). The strong *Tc-srp* signal in nuclei may suggest

nuclear or peri-nuclear localisation of the transcript, or it may be due to the cell body being

flattened. The strong patch of expression at the anterior medial region is from cells beneath

the ventral ectoderm (see Supplemental Fig 1 for a full expression series). (C-E) Extended

germband stage *Tribolium* embryos transiently expressing NLS-tdEos showing unconverted

protein (magenta) and photoconverted protein (cyan). In each embryo, the clone of

converted cells spans the entire amnion. (C_{1-2}) show both sides of the same embryo in which

a 6 nuclei wide patch of dorsal-most cells located at 50% EL were photoconverted at the

blastoderm stage. (E) shows an embryo in which a 3 nuclei wide patch of dorsal-most cells

located at 25% EL were photoconverted at the blastoderm stage. (F) shows an embryo in

which a 3 nuclei wide by 6 nuclei long patch of dorsal-most cells located at roughly 2-10% EL

were photoconverted at the blastoderm stage. In (A) and (C-E), embryos are oriented based

on the AP/DV polarity of the egg with anterior to the left and dorsal to the top. In (A''), the

surface of the egg is oriented to the bottom. In (B), the anterior of the germband is to the

left. (A) is an average intensity projection of one egg hemisphere. (A') is an average intensity

projection of 6 μm to specifically show the dorsal epithelium. (B) is a maximum intensity

projection of the whole germband. (C-E) are maximum intensity projections of one egg

hemisphere. Abbreviations are: ventral ectoderm (VE) and dorsal ectoderm (DE). Scale bars

are 50 μm.

Fig 5. Schematics showing the classic and revised *Tribolium* fatemaps and germband

models. Schematics drawn as in Fig 1 to show the classic fatemap and germband model and

the revised fatemap and model based on the results of this manuscript. Note that the

posterior amnion/embryo boundary is unclear. The schematics of the flatmounted

germbands are drawn with the focus on the dorsal epithelium. See text for additional details,

and Supplemental Fig 6 for an extended figure.

Fig 6. Representation of the *Tribolium* germband as a cylinder. (A_1) the germband of

Tribolium (and other insects) is usually drawn as it is in *in vivo*, that is, as a flattened shape

with the ventral cell fates to the top and the dorsal cell fates to the bottom. However, the

embryo can be unflattened (A_2) without changing the planar dimensions of the tissue itself,

and rotated (A_3) such that the dorsal-ventral axis is oriented in the widely accepted standard

of dorsal to the top and ventral to the bottom. When this is done, and third dimension is

incorporated to show the AP axis (A_4), the similarities in tissue boundaries and tissue

rearrangements in *Tribolium* and *Drosophila* (B) become obvious. Note that while both (A_4)

and (B) are cylindrical, the epithelium of the *Tribolium* germband inverts during embryo

condensation so the apical surface of the embryo faces inwards. Both (A) and (B) are

sections from roughly the middle of the respective embryo during GBE. Possible flattening of

dorsal tissue in *Drosophila* has been omitted due to a lack of detailed data at this stage.

Supplementary Figure 1. Results of photoconversions at 50% egg length. NLS-tdEos labelled

extended germband stage *Tribolium* embryos in which a patch of blastoderm nuclei were

photoconverted at 50% egg length (from the posterior pole) at different DV positions. The

approximate DV position of the patch and the approximate DV width of the clone (in terms

of nuclei number) are shown. The dorsal labelled embryo is shown from both sides to

demonstrate the photoconverted nuclei cover the full DV extent of the amnion (arrows).

Unconverted protein is shown in magenta, converted protein is shown in cyan. Images are

maximum intensity projections of one egg hemisphere. All eggs are oriented with the

anterior to the left and ventral to the bottom. Scale bars are 100 μm.

Supplementary Figure 2. Results of photoconversions at 25% egg length. NLS-tdEos labelled

extended germband stage *Tribolium* embryos in which a patch of blastoderm nuclei were

photoconverted at 25% egg length (from the posterior pole) at different DV positions. The

approximate DV position of the patch and the approximate DV width of the clone (in terms

of nuclei number) are shown. Unconverted protein is shown in magenta, converted protein

is shown in cyan. Images are maximum intensity projections of one egg hemisphere. All eggs

are oriented with the anterior to the left and ventral to the bottom. Scale bars are 100 μm.

Supplementary Figure 3. Results of photoconversions near the posterior pole. NLS-tdEos

labelled extended germband stage *Tribolium* embryos in which a patch of blastoderm nuclei

were photoconverted near the posterior pole at different DV positions. The approximate DV

position of the patch and the approximate DV width of the clone (in terms of nuclei number)

are shown. The second dorsally labelled embryo is shown at high magnification at two

timepoints and with a transverse section (at the position of the dashed green line) to show

the movement of tissue from the dorsal epithelium into the hindgut. Unconverted protein is

shown in magenta, converted protein is shown in cyan. Images are maximum intensity

projections of one egg hemisphere except for the bottom three embryos, which are shown

as maximum intensity projects through the germband in order to better show the labelled

nuclei. All eggs are oriented with the anterior to the left and ventral to the bottom except

for the second timepoint of the second dorsal view, which is shown with the posterior of the

germband to the left. Scale bars are 100 μm.

Supplementary Figure 4. *Tc-cad* is expressed further to the anterior in the dorsal

epithelium than in the ventral epithelium. (A-A'') show maximum intensity projections of

the posterior region of a flatmounted *Tribolium* germband shortly after serosa window

closure stained to show nuclei (DAPI, blue) and the mRNA expression pattern of *Tc-cad* (red).

The panels beside or below each main panel show optical sections through the germband at

the position of the respective coloured line in the main panel. (B-B'') show maximum

intensity projections of approximately one half of the germband from the lateral side (i.e.

projections of optical sagittal sections from the midline to the lateral edge of the germband).

Scale bar is 100 μm.

Supplementary Figure 5. RNA expression of the *Tribolium* ortholog of the GATA factor

serpent. (A-F) whole mount and (G-J) flatmount *Tribolium* embryos from the pre-blastoderm

to the retracting germband stage stained for *Tc-srp* mRNA (red) and nuclei (DAPI, blue). (G_1)

and (G_2) show the same embryo imaged from both sides. (H_1) and (H_2) show projections

from the dorsal epithelium (H_1) and the ventral epithelium (H_2) of the same embryo. *Tc-srp*

mRNA is maternally provided (A), and expression is ubiquitous until the late blastoderm

stage (B-C) when expression clears from the blastoderm but persists in the yolk nuclei

(scattered spots in (D-E)). During embryo condensation, *de novo* expression arises in a patch

of blastoderm cells at the anterior medial region (arrowhead in F). This patch of *Tc-srp*

expressing cells invaginates as part of the ventral furrow and becomes located beneath the

ectoderm (arrowhead in G_1-H_3). This expression domain is likely homologous to the anterior

ventral expression domain in *Drosophila* that marks the prohemocytes. During serosa

window closure, expression appears in a ring of dorsal epithelium cells (G_1). After serosa

window closure, expression persists in the dorsal epithelium (H_1) and (H_3). Unlike *Drosophila*,

there is no expression domain at the posterior of the blastoderm (E) or the early germband

(G_2). After germband elongation, a *de novo* expression domain appears at the posterior most

point of the embryo (I_1). Given the location of this domain at the base of the forming

hindgut, this is likely the posterior endoderm primordium. Expression can also be seen in a

patch of amnion that has remained attached to the germband (arrow in I_2), but most of the

rest of the amnion has been lost. Several other regions of expression can be seen, including

in the presumptive fat body (the segmental domains running down the body), in

presumptive hemocyte clusters (the two side-by-side domains in the anterior), and in an

anterior domain that may mark the anterior endoderm primordium. Expression also persists

in the yolk nuclei (data not shown). All embryos are shown with anterior to the left. (E-E'') is

oriented with the ventral to the bottom, (F-F'') is oriented as a ventral view. (H_3-H_3'') is an

optical sagittal section at approximately the midline of the embryo in (H_2). (A-F) are

maximum intensity projections of one egg hemisphere. (G_1-G_2''), (I1-I1'') and (J-J'') are

maximum intensity projections of flatmounted germbands. (I_2-I_2'') is a maximum intensity

projection of part of the germband to better show the amnion/dorsal epithelium. Scale bars

are 50 µm.

Supplementary Figure 6. Extended schematics showing the classic and revised *Tribolium*

fatemaps and germband models. Refer to Fig 5 and the text for details.

Supplementary Figure 7. Tissue specific cell shape changes during *Tribolium* condensation.

Stills from timelapses of two *Tribolium* embryos transiently expressing GAP43YFP to label

membranes. The second panel of each timepoint shows optical transverse sections at the

position of the dashed line in the related panel. Ventral and lateral ectoderm becomes

columnar, while dorsal ectoderm becomes flattened. The non-columnar cells at the bottom

of the left hand embryo are likely the presumptive mesoderm. The first frame of the

timelapses was defined as timepoint 0. Both embryos are oriented with the anterior to the

left and ventral to the bottom. Abbreviations are: Dorsal (Dor), Lateral (Lat), Ventral (Ven)

and Ectoderm (Ect). Scale bars are 100 μm.

Supplementary Figure 8. DV dynamics of the segment polarity gene *gooseberry* in

***Drosophila*.** Whole mount *Drosophila* embryos from the blastoderm stage (late stage 5) until

the extending germband stage (late stage 7/early stage 8). The red arrows in the 2nd and 3rd

panels highlight the dorsal-to-ventral tissue flow and the GBE. The vertical arrow is at

parasegment boundary 8/9, and is the same size in both panels. All embryos are

ventrolateral views with anterior to the left and dorsal to the top. Scale bar is 100 μm.

Images were kindly provided by Erik Clark.

Supplementary Movie 1. Confocal timelapse of a *Tribolium* embryo transiently expressing

H2B-ven to mark nuclei. A maximum intensity projection of one egg hemisphere is shown.

Anterior is to the left, the ventral side of the egg is to the bottom.

Supplementary Movie 2. Same timelapse as Supplementary Movie 1, but with nuclei of the

dorsal epithelium tracked until they join the ventral epithelium. Nuclei that join the ventral

epithelium are labelled magenta, nuclei that become located at the edge of the germband

are labelled yellow, nuclei that remain in the dorsal epithelium are labelled cyan. Anterior is

to the left, the ventral side of the egg is to the bottom. See Fig 2(A-C) for more details.

Supplementary Movie 3. Same timelapse as Supplementary Movie 1, but with a line of

nuclei of the dorsal epithelium tracked. Nuclei that remain in the dorsal epithelium are

labelled cyan. Anterior is to the left, the ventral side of the egg is to the bottom. See Fig 2(D-

F) for more details.

Supplementary Movie 4. Confocal timelapse of a *Tribolium* embryo transiently expressing

NLS-tdEos (magenta) with a line of blastoderm cells photoconverted (cyan). The brightness

increases approximately halfway through the movie due to a manual increase in laser power

at this point. A maximum intensity projection of one egg hemisphere is shown. Anterior is to

the left, the ventral side of the egg is to the bottom.

Supplementary Movie 5. Confocal timelapse of a *Tribolium* embryo transiently expressing

NLS-tdEos (magenta) with a patch of blastoderm cells photoconverted (cyan). (A) shows a

maximum intensity projection of one egg hemisphere is shown. (B) shows an optical

transverse section. (C) shows an average intensity projection of 10 optical transverse

sections. Anterior is to the left, the ventral side of the egg is to the bottom. See Fig 3(A-F) for

more details.

Supplementary Movie 6. Confocal timelapse of a *Tribolium* embryo transiently expressing

NLS-tdEos (magenta) with a patch of blastoderm cells photoconverted (cyan). Towards the

end of the timelapse, the cyan nuclei in the dorsal epithelium appear to condense

posteriorly to the hindgut. A maximum intensity projection of one egg hemisphere is shown.

Anterior is to the left, the ventral side of the egg is to the bottom.

Drosophila

Tribolium

Amnioserosa

Posterior gut

Cephalic furrow

Serosa

Amnion

Surface view

Sagittal section

Surface view

Sagittal section

Flatmount

Cell Intercalation

Planar cell stretching

Cell shape change

Cell division

K Classic model

L Revised model

VE   DE

Classic model

Revised model

Serosa
Amnion

Blastoderm dorsal view

Blastoderm dorsal view

?

Surface view

Flatmount

Surface view

Flatmount

A_1 Ventral / Dorsal / *Tribolium*

A_2 Ventral / Dorsal — *Unflatten*

A_3 Dorsal / Ventral — *Rotate*

A_4 =

B *Drosophila*