1   ***Leishmania naiffi* and *Leishmania guyanensis* reference genomes highlight genome**
2   **structure and gene evolution in the *Viannia* subgenus**
3

4   **Authors:** Simone Coughlan (1), Ali Shirley Taylor (2), Eoghan Feane (2), Mandy Sanders (3),
5   Gabriele Schonian (4), James A. Cotton (3), Tim Downing (1,2*).
6
Affiliations:
(1) School of Mathematics, Applied Mathematics and Statistics, National University of Ireland,
Galway, Ireland.
(2) School of Biotechnology, Dublin City University, Dublin, Ireland.
(3) Wellcome Trust Sanger Institute, Hinxton, UK.
(4) Charité University Medicine, Berlin, Germany.
* Corresponding author
14
Email addresses:
Simone Coughlan: coughls@gmail.com
Ali Taylor: ali.taylor7@mail.dcu.ie
Eoghan Feane: eoghan.feane2@mail.dcu.ie
Mandy Sanders: mjs@sanger.ac.uk
Gabriele Schonian: gabriele.schoenian@t-online.de
James A. Cotton: jc17@sanger.ac.uk
Tim Downing: tim.downing@dcu.ie

23
24 **Abstract**
25

The unicellular protozoan parasite *Leishmania* causes the neglected tropical disease
leishmaniasis, affecting 12 million people in 98 countries. In South America where the *Viannia*
subgenus predominates, so far only *L. (Viannia) braziliensis* and *L. (V.) panamensis* have been
sequenced, assembled and annotated as reference genomes. Addressing this deficit in
molecular information can inform species typing, epidemiological monitoring and clinical
treatment. Here, *L. (V.) naiffi* and *L. (V.) guyanensis* genomic DNA was sequenced to assemble
these two genomes as draft references from short sequence reads. The methods used were
tested using short sequence reads for *L. braziliensis* M2904 against its published reference as a
comparison. This assembly and annotation pipeline identified 70 additional genes not
annotated on the original M2904 reference. Phylogenetic and evolutionary comparisons of *L.
guyanensis* and *L. naiffi* with ten other *Viannia* genomes revealed four traits common to all
Viannia: aneuploidy, 22 orthologous groups of genes absent in other *Leishmania* subgenera,
elevated TATE transposon copies, and a high NADH-dependent fumarate reductase gene copy
number. Within the *Viannia*, there were limited structural changes in genome architecture
specific to individual species: a 45 Kb amplification on chromosome 34 was present in all bar
L. lainsoni, *L. naiffi* had a higher copy number of the virulence factor leishmanolysin, and
laboratory isolate *L. shawi* M8408 had a possible minichromosome derived from the 3' end of
chromosome 34. This combination of genome assembly, phylogenetics and comparative
analysis across an extended panel of diverse *Viannia* has uncovered new insights into the origin
and evolution of this subgenus and can help improve diagnostics for leishmaniasis surveillance.

46
47 Keywords: *Leishmania*; leishmaniasis; genome; assembly; aneuploidy.
48

Introduction

Most cutaneous (CL) and mucocutaneous leishmaniasis (MCL) cases in the Americas are the result of infection by *Leishmania* parasites belonging to the *Viannia* subgenus. The complexity of the molecular, epidemiological and ecological challenges associated with *Leishmania* in South America remains opaque due to our limited understanding of the biology of *Viannia* parasites. Nine *Viannia* (sub)species have been described so far: *L. (V.) braziliensis, L. (V.) peruviana, L. (V.) guyanensis, L. (V.) panamensis, L. (V.) shawi, L. (V.) lainsoni, L. (V.) naiffi, L. (V.) lindenbergi* and *L. (V.) utingensis*. CL and MCL are endemic in 18 out 20 countries in the Americas [1] and are mainly associated with *L. braziliensis, L. guyanensis,* and *L. panamensis,* whose frequency varies geographically. Other species are less frequently associated with human disease, and some are restricted to certain areas [2].

Human CL is partially driven by transmission from sylvatic and peridomestic mammalian reservoirs [3], via sand flies of the genus *Lutzomyia* (*sensu* Young and Duncan, 1994) in the Americas, distinct from *Phlebotomus* sand flies in the Old World [4]. Although CL has spread to domestic and peridomestic niches due to migration, new settlements and deforestation [5-7], there is still a high incidence of some *Leishmania* in sylvatic environments, such that human infection is accidentally acquired due to sand fly bites when handling livestock [8]. *L. naiffi* and *L. guyanensis* are among the *Viannia* species that show variable responses to treatment, and diversity in the types of clinical manifestations presented, and are adapting to environmental niche and transmission changes driven by humans.

L. naiffi was formally described from a parasite isolated in 1989 from its primary reservoir, the nine-banded armadillo (*Dasypus novemcinctus*), in Pará state of northern Brazil [9-11]. *L. naiffi* was initially placed in the *Viannia* subgenus based on its molecular and immunological characteristics [9]. Many phlebotomine species are likely to participate in the transmission of *L. naiffi* in Amazonia [12], including *Lu. (Psathyromyia) ayrozai* and *Lu. (Psychodopygus) paraensis* in Brazil [13], *Lu. (Psathyromyia) squamiventris* and *Lu. tortura* in Ecuador [14], and *Lu. trapidoi* and *Lu. gomezi* in Panama [30]. *L. naiffi* has been isolated from humans and armadillos [9-10], and detected in *Thrichomys pachyurus* rodents found in the same habitat as *D. novemcinctus* in Brazil [16]. The nine-banded armadillo is hunted, handled and consumed in the Americas and is regarded as a pest [11,17-18]. People in the same vector range as these armadillos could be exposed to infective sand flies: three *L. naiffi* CL cases followed contact with armadillos in Suriname [19]. *L. naiffi* causes localised CL in humans with small discrete lesions on the hands, arms or legs [10,20-21], which has been observed in Brazil, French Guiana, Ecuador, Peru and Suriname [19,22]. CL due to *L. naiffi* usually responds to treatment [10,22] and can be self-limiting [23], though poor response to antimonial or pentamidine therapy was reported in two patients in Manaus, Brazil [20].

L. guyanensis was first described in 1954 [24] and its primary hosts are the forest dwelling two-toed sloth (*Choloepus didactylus*) and the lesser anteater *Tamandua tetradactyl* [25]. Potential secondary reservoirs of *L. guyanensis* are *Didelphis marsupialis* (the common opossum) [26,27], rodents from the genus *Proechimys* [25], *Marmosops incanus* (the grey slender opposum) [28] in Brazil, and *D. novemcinctus* [29]. *Lu. umbratilis, Lu. anduzei* and *Lu. whitmani* are prevalent in forests [30] and act as vectors of *L. guyanensis* [31-33]. *L. guyanensis* has been found in French Guiana, Bolivia, Brazil, Colombia, Guyana, Venezuela, Ecuador, Peru, Argentina and Suriname [34-39].

More precise genetic screening of *Viannia* isolates is necessary to trace hybridisation between species. Infection of humans, dogs and *Lu. ovallesi* with *L. guyanensis/L. braziliensis* hybrids was reported in Venezuela [40-41]. A *L. shawi/L. guyanensis* hybrid causing CL was detected in Amazonian Brazil [42], and *L. naiffi* has produced viable progeny with *L. lainsoni* [43] and

L. braziliensis (Elisa Cupolillo, unpublished data). There is extensive evidence of interbreeding
among *L. braziliensis* complex isolates, including more virulent *L. braziliensis/L. peruviana*
hybrids with higher survival rates within hosts *in vitro* [44].

Leishmania genomes are characterised by several key features. Genes are organised as
polycistronic transcription units that have a high degree of synteny across *Leishmania* species
[45]. These polycistronic transcription units are co-transcribed by RNA polymerase II as
polycistronic pre-mRNAs that are 5'-transpliced and 3'-polyadenylated [46,47]. This means
translation and stability of these mature mRNAs determines gene expression rather than
transcription rates. In addition, *Leishmania* display extensive aneuploidy, frequently possess
extrachromosomal amplifications driven by homologous recombination at repetitive sequences,
and have variable gene copy numbers [48]. The *Leishmania* subgenus genomes of *L. infantum*,
L. donovani, and *L. major* have 36 chromosomes [49], whereas *Viannia* genomes have 35
chromosomes due to a fusion of chromosomes 20 and 34 [45,50]. In contrast to the species of
the *Leishmania* subgenus, *Viannia* parasites possess genes encoding functioning RNA
interference (RNAi) machinery that may mediate infective viruses and transposable elements
[51].

Fully annotated genomes have been described in detail for only two *Viannia* species: *L.
panamensis* [51] and *L. braziliensis* [45,48], limiting our comprehension of their evolutionary
origin, genetic diversity and functional adaptations. Consequently, we present reference
genomes for *L. guyanensis* LgCL085 and *L. naiffi* LnCL223 to address these critical gaps.
These new annotated reference genomes were compared to other *Viannia* species genomes to
examine structural variation, sequence divergence, gene synteny and chromosome copy
number changes. We contrasted the genomic configuration of *L. guyanensis* LgCL085 and *L.
naiffi* LnCL223 with the *L. braziliensis* MHOM/BR/1975/M2903 assembly, two unannotated
L. peruviana chromosome-level scaffold assemblies [52], the *L. panamensis*
MHOM/PA/1994/PSC-1 reference and the *L. braziliensis* MHOM/BR/1975/M2904 reference.
Furthermore, we assessed aneuploidy in five unassembled *Viannia* datasets originally isolated
from humans, armadillos and primates, which are commonly used in studies on *Viannia*
parasites [53–56]: *L. shawi* reference isolate MCEB/BR/1984/M8408 also known as
IOC_L1545, *L. guyanensis* MHOM/BR/1975/M4147 (iz34), *L. naiffi* MDAS/BR/1979/M5533
(IOC_L1365), *L. lainsoni* MHOM/BR/1981/M6426 (IOC_L1023), *L. panamensis*
MHOM/PA/1974/WR120 [53] (IOC stands for Instituto Oswaldo Cruz).

Results

Genome assembly from short-reads
The genomes of *L. (Viannia) guyanensis* LgCL085 and *L. (V.) naiffi* LnCL223 were assembled
from short reads, along with an assembly of *L. braziliensis* M2904 generated in the same way
as a positive control [48] (Table 1). This facilitated comparison with the published M2904
genome, which was assembled by capillary sequencing of a plasmid clone library together with
extensive finishing work and with fosmid end sequencing [45], so that the ability of short reads
to correctly and comprehensively resolve *Leishmania* genome architecture could be quantified.

Firstly, the *L. guyanensis* LgCL085, *L. naiffi* LnCL223 and the *L. braziliensis* M2904 control
reads were filtered to remove putative contaminant sequences identified by aberrant GC
content, trimmed at the 3' ends to remove low quality bases, and PCR primer sequences were
removed (see Methods for details) resulting in 26,067,692 properly paired reads for *L.
guyanensis*, 13,979,628 for *L. naiffi,* 34,592,618 for the *L. (V.) braziliensis* control (Table S1).
These filtered reads for *L. guyanensis*, *L. naiffi* and *L. braziliensis* were *de novo* assembled into
contigs using Velvet [57] with k-mers of 61 for *L. guyanensis,* 43 for *L. naiffi* and 43 for the *L.
braziliensis* control optimised for each library.

The initial contigs were scaffolded using read pair information with SSPACE [58] to yield 2,800 *L. guyanensis* scaffolds with an N50 of 95.4 Kb, 6,530 *L. naiffi* scaffolds with an N50 of 24.3 Kb, and 3,782 *L. braziliensis* scaffolds with an N50 of 20.6 Kb (Table 2). The corrected scaffolds for *L. guyanensis, L. naiffi* and the *L. braziliensis* control were contiguated (aligned, ordered and oriented) using the extensively finished *L. braziliensis* M2904 reference with ABACAS [59]. The output was split into 35 pseudo-chromosomes and REAPR [60] broke scaffolds at possible misassemblies to assess contiguation accuracy. The pseudo-chromosome lengths of each sample approximated the length of each corresponding *L. braziliensis* M2904 reference chromosome with the exceptions of shorter *L. guyanensis* chromosomes 2, 4, 12 and 21, and a longer *L. naiffi* chromosome 1 (Figure S1). Post-assembly alignment of all bin contigs using BLASTn identified 44 *L. guyanensis* sequences spanning 4,566,791 bp as putative contaminants that were removed: half had high similarity to bacterium *Niastella koreensis* (Table S2).

When the reads for each were mapped to its own assembled genome, the median read coverage was 56 for *L. guyanensis*, 36 for *L. naiffi* and 75 for the *L. braziliensis* control. The latter was on par with the 74-fold median coverage observed when M2904 short reads were mapped to the *L. braziliensis* reference [45,48] (Table S3). The differing coverage levels correlated with the numbers of gaps in the final genome assembly of *L. guyanensis* (1,557, Table 2) and *L. naiffi* (3,853).

MLSA of *L. guyanensis* LgCL085 and *L. naiffi* LnCL223 with the *Viannia* subgenus
As a first step in investigating the genetic origins of these isolates, we examined their species identity using MLSA (multi-locus sequencing analysis). Four housekeeping gene sequences published for 95 *Viannia* isolates including *L. braziliensis, L. lainsoni, L. lindenbergi, L. utingensis, L. guyanensis, L. shawi* and *L. naiffi* [56] were compared with orthologs of each gene extracted from assemblies of *L. naiffi* LnCL223, *L. guyanensis* LgCL085, the *L. braziliensis* reference, *L. panamensis* PSC-1 and *L. peruviana* PAB-4377. Among the 95 were four samples with reads available [53]: *L. shawi* MCEB/BR/1984/M8408 (IOC_L1545), *L. guyanensis* MHOM/BR/1975/M4147 (iz34), *L. naiffi* MDAS/BR/1979/M5533 (IOC_L1365) and *L. lainsoni* MHOM/BR/1981/M6426 (IOC_L1023). The genes were aligned using Clustal Omega v1.1 [61] to create a network for the 102 isolates with SplitsTree v4.13.1 [62]. This replicated the expected highly reticulated structure [56] where *L. braziliensis* M2904 and *L. peruviana* PAB-4377 were in the *L. braziliensis* cluster (Figure 1).

Previous work suggests that the *L. guyanensis* species complex includes *L. panamensis* and *L. shawi* because they show little genetic differentiation from one another [56,63-65]. The MLSA here showed that the new *L. guyanensis* LgCL085 reference clustered phylogenetically in the *L. guyanensis* species complex, had no sequence differences compared to *L. panamensis* PSC-1, and seven differences versus *L. shawi* M8408 across the 2,902 sites aligned (Figure 1). *L. guyanensis* LgCL085 grouped with isolates classified as zymodeme Z26 by multilocus enzyme electrophoresis (MLEE) associated with *L. shawi* [54]. This was supported by the number and the alleles of genome-wide SNPs called using reads mapped to the *L. braziliensis* M2904 reference for *L. guyanensis* (355,267 SNPs), *L. guyanensis* M4147 (326,491), *L. panamensis* WR120 (294,459) and *L. shawi* M8408 (296,095) (Table S4).

The *L. naiffi* LnCL223 was closest to *L. naiffi* ISQU/BR/1994/IM3936, with two differences. It clustered with MLEE zymodeme Z49 based on the correspondence between the MLSA network and previously typed zymodemes, though *L. naiffi* is associated with more zymodemes than other *Viannia*. The number and the alleles of genome-wide SNPs called using reads mapped to the *L. braziliensis* reference were similar for *L. naiffi* (548,256) and M5533 (633,560) (Table S4) and consistent with the MLSA genetic distances.

There was no evidence of recent gene flow between these three species at any genome-wide 10
Kb segment and *L. naiffi* LnCL223 had fewer SNPs compared to *L. braziliensis* M2904 than *L.*
guyanensis LgCL085 (Figure S2). Linking the MLSA network topology with previous work
[56,63-65], four genetically distinct species complexes are represented by the genome-
sequenced *Viannia* at present: (i) *braziliensis* including *L. peruviana*, (ii) *guyanensis* including
L. panamensis and *L. shawi*, (iii) *naiffi*, and (iv) *lainsoni* (Table S4), and the less explored (v)
lindenbergi and (vi) *utingensis* complexes (Figure 1).

Ancestral diploidy and constitutive aneuploidy in *Viannia*
The normalised chromosomal coverage of the *L. guyanensis* LgCL085 and *L. naiffi* LnCL223
reads mapped to *L. braziliensis* M2904 showed aneuploidy on a background of a diploid
nuclear genome (Figure 2). The coverage levels of reads for *L. peruviana* LEM1537, *L.*
peruviana PAB-4377, *L. panamensis* PSC-1 and the triploid *L. braziliensis* control mapped to
the M2904 reference, confirmed previous work (Figure S3), including the *L. braziliensis*
control (Figure S4), and demonstrated that assemblies from short read data were sufficient to
estimate chromosome copy number differences. Repeating this for *L. shawi* M8408, *L. naiffi*
M5533, *L. guyanensis* M4147, *L. panamensis* WR120 and *L. lainsoni* M6426 showed that all
these *Viannia* were predominantly disomic and thus diploidy was the likely ancestral state of
this subgenus (Figure 2).

The somy patterns were supported by the results of mapping the reads of each sample to their
own assembled genome or to the M2904 reference to produce the read depth allele frequency
(RDAF) distributions from heterozygous SNPs. The majority of *L. braziliensis* M2904 control
chromosomes had peaks with modes at ~33% and ~67% indicating trisomy, rather than a single
peak at ~50% consistent with disomy (Figure S5). The RDAF distributions from reads mapped
to its own assembly for *L. guyanensis* LgCL085 and *L. naiffi* LnCL223 had a mode of ~50%
(Figure S6), including peaks indicating trisomy for LgCL085 chromosomes 13, 26 and 35
(Figure S7).

8,262 *L. naiffi* and 8,376 *L. guyanensis* genes annotated
A total of 8,262 genes were annotated on *L. naiffi* LnCL223: of these 8,104 were protein
coding genes, 78 were tRNAs, 15 rRNA genes, four snoRNA genes, two snRNA genes, and 59
pseudogenes. 310 genes were on unassigned contigs (Table S3). 8,376 genes were annotated on
L. guyanensis LgCL085: of these 8,230 were protein coding genes, 75 tRNAs, 14 rRNA genes,
four snoRNA genes, two snRNA genes and 51 pseudogenes. 619 genes were on unassigned
contigs

There were 8,161 genes (8,001 protein coding) transferred to the control *L. braziliensis*
genome, along with 76 tRNAs, two snRNA genes, four snoRNA genes, 13 rRNA genes and 65
pseudogenes (Table 2). 7,719 of the protein coding genes (96.5%) clustered into 7,244 OGs,
whereas 8,137 of the 8,375 (97.2%) protein coding genes on the *L. braziliensis* reference
grouped into 7,383 OGs. This indicated that 97% of protein coding genes in OGs were
recovered, and only 2.8% (235) across 201 OGs were absent in the M2904 control, mainly
hypothetical or encoded ribosomal proteins (Table S5). In the same way, we found 70 protein
coding genes (Table S6) in 62 OGs on the M2904 control absent in the published *L.*
braziliensis annotation.

Few genes were present in *L. braziliensis* but absent in *L. guyanensis* LgCL085 and *L. naiffi*
LnCL223. Coverage depth was used to predict each gene's haploid copy number, such that
genes with haploid copy numbers at least twice the assembled copy number indicated partially
assembled genes in the reference assembly. Thus, we investigated all OGs with haploid copy
numbers at least twice the assembled copy number to quantity completeness of the assembly.

Only 145 genes in 92 OGs on *L. guyanensis* LgCL085 (Table S7), 142 genes in 90 OGs on *L.*
naiffi LnCL223 (Table S8) and 102 genes in 71 OGs (Table S9) on the *L. braziliensis* control
met this criterion, indicating few unassembled genes in each assembly. One hypothetical gene
(LnCL223_272760) in *L. naiffi* LnCL223 with no retrievable information had a haploid copy
number of 15 (OG5_173495), whereas all other genomes examined here had zero to two
copies.
269

A 245 Kb rearrangement akin to a minichromosome in *L. shawi* M8408
We discovered a putative minichromosome or amplification at the 3' end of *L. shawi* M8408
chromosome 34 based on elevated coverage across a pair of inverted repeats spanning 245 Kb
(Figure 3). This locus spanned at least bases 1,840,001 to 1,936,232 (the end) of *L. braziliensis*
M2904 chromosome 34 (Figure S8, Table S10). It was orthologous to a known 100 Kb
amplification on *L. panamensis* PSC-1 chromosome 34 that was predicted to produce a
minichromosome when amplified, and contained the frequently amplified LD1 (*Leishmania*
DNA 1) region [66]. In contrast to the *L. panamensis* PSC-1 minichromosome, the *L. shawi*
M8408 amplification was ~30 Kb longer and closer in length to the *L. braziliensis* M2903 245
Kb minichromosome [67].
280

A 45 Kb locus was amplified in most *Viannia* genomes
A 45 Kb amplification on chromosome 34 spanning a gene encoding a structural maintenance
of chromosome (SMC) family protein and ten hypothetical genes had between two and four
copies in all samples except *L. lainsoni* M6426 (Figure 3, Table S10). Using the *L. guyanensis*
gene annotation, putative functions were assigned to five of the ten hypothetical genes. This
duplication spanned chromosomal location 1.32-1.35 Mb in the *L. braziliensis* M2904
reference and had two additional hypothetical genes in *L. naiffi* LnCL223 (LnCL223_343280
and LnCL223_343290, Figure 4).
289

Genes exclusive to *Viannia* genomes
7,961 (96.7%) of the 8,230 genes annotated for *L. guyanensis* LgCL085 were assigned to 7,381
OGs, 7,893 (97.4%) of the 8,104 *L. naiffi* LnCL223 genes to 7,324 OGs, and 7,692 (99.3%) of
the *L. panamensis* PSC-1 7,748 to 7,245 OGs. A total of 6,835 of these OGs were shared with
nine species from the *Leishmania, Sauroleishmania* and *Viannia* subgenera: *L. (L.) major, L.*
(L.) mexicana, L. (L.) donovani (infantum), L. (V.) guyanensis, L. (V.) naiffi, L. (V.)
braziliensis, L. (V.) panamensis, L. (S.) adleri, L. (S.) tarentolae (Table S11).
297

We identified 22 OGs exclusive to *Viannia* (Table S12): three OGs contained the RNAi
pathway genes (DCL1, DCL2, RIF4). Another OG was the telomere-associated mobile
elements (TATE) DNA transposons (OG5_132061), a dynamic feature of *Viannia* genomes
[51] (Supplementary Results). Four OGs encoded a diacylglycerol kinase-like protein
(OG5_133291), a nucleoside transporter (OG5_134097), a beta tubulin / amastin
(OG5_183241), and a /zinc transporter (OG5_214682). The remaining 14 OGs contained
hypothetical genes.
305

A NADH-dependent fumarate reductase gene (OG5_128620) was amplified in the *Viannia*
examined here: *L. guyanensis* LgCL085 had 14 copies, *L. naiffi* LnCL223 had 16, *L.*
panamensis PSC-1 had 16, *L. peruviana* PAB4377 had 23, *L. peruviana* LEM1537 had 14, and
braziliensis M2904 had 12. This contrasted with the *Leishmania* and *Sauroleishmania*
subgenera for which three to four copies had been reported for *L. infantum, L. mexicana, L.*
major, L. adleri and *L. tarentolae* [68,69]. This gene has been implicated in enabling parasites
to resist oxidative stress and potentially aiding persistence, drug resistance and metastasis
[70,71].
314

Few species-specific genes in *L. guyanensis* LgCL085 and *L. naiffi* LnCL223

Four genes from four OGs unique to *L. naiffi* LnCL223 were identified compared to other
Leishmania (Table S13). Of these four, hypothetical genes LnCL223_312570 and
LnCL223_292920 had orthologs in *T. brucei* and *T. vivax,* respectively. The LnCL223_341350
protein product had 44-45% sequence identity with a *Leptomonas* transferase family protein,
and LnCL223_352070 was a methylenetetrahydrofolate reductase (OG5_128744), but had no
orthologs in the other eight *Leishmania* or five *Trypanosoma* species investigated here. *L.*
guyanensis LgCL085 had 31 unique genes in 30 OGs, 25 of which were on unplaced contigs.
Four of the six chromosomal genes were also in *Trypanosoma* genomes, encoding two
hypothetical proteins (a tuzin and a poly ADP-ribose glycohydrolase). 28 of the 31 had
orthologs in eukaryotes, of which three had orthologs in the free-living freshwater ciliate
protozoan *Tetrahymena thermophile* (Table S14) [72].

L. guyanensis LgCL085 and L. naiffi LnCL223 had over 300 gene arrays
Gene arrays are genes in the same OG with more than two haploid gene copies: they can be *cis*
or *trans*. There were 327 gene arrays on *L. naiffi* LnCL223 (Table S15), 334 on *L. guyanensis*
LgCL085 (Table S16) and 255 on the control *L. braziliensis* M2904 (Table S17) - half the
arrays on each genome had two copies of each gene. 22 of the *L. guyanensis*, 18 of the *L. naiffi*
LnCL223 and 15 of the control *L. braziliensis* gene arrays contained 10+ haploid gene copies
(Table 3). The *L. panamensis* PSC-1 genome had ~400 tandem arrays, of which 71% had more
than two copies. The *L. braziliensis* M2904 genome had 615 arrays corresponding to 763 OGs
in OrthoMCL v5. Thus, the control genome underestimated the number of gene arrays due to
either gene absence or incomplete assembly, indicating that the number of arrays on *L. naiffi*
LnCL223 and *L. guyanensis* LgCL085 was underestimated.

The most expanded array on *L. guyanensis* LgCL085 contained TATE DNA transposons
(OG5_132061) with 50 haploid gene copies (Table 3) compared with 11 on *L. naiffi* LnCL223,
21 on the *L. braziliensis* control and 16 on *L. panamensis* PSC-1. The *L. braziliensis* M2904
assembly had 40 TATE DNA transposons, but only two were annotated on the control here,
illustrating that more accurate estimates of copy number may be possible.

L. naiffi LnCL223 had the highest haploid gene copy number of the M8 family metalloprotease
leishmanolysin (GP63) array (OG5_126749) with 56 haploid gene copies, compared to 33 in *L.*
guyanensis LgCL085, 28 in *L. panamensis* PSC-1 and 31 in *L. braziliensis* M2904. This was
the sole protease-related OG amplified in all three species (Table S23). This family was not
expanded in *L. peruviana* LEM1537 or PAB4377. This was consistent with previous work on
L. guyanensis leishmanolysin [73] indicating it is a highly expressed virulence factor in
promastigotes [74] affecting the survival during the initial stages of infection [74-77].
Sauroleishmania genomes also had high array copy numbers: 37 for *L. adleri* [69] and 84 for
L. tarentolae (Table S12). *Leishmania* subgenus genomes had lower copy numbers, with 13 for
L. mexicana, 15 for *L. infantum* and five for *L. major* (OG4_10176 for *L. braziliensis* M2904,
L. mexicana, *L. infantum* and *L. major*).

A tuzin gene array (OG5_173452) had higher haploid copy numbers on *L. guyanensis*
LgCL085 (19) and *L. panamensis* PSC-1 (22) compared with the two copies in *L. naiffi*, *L.*
mexicana, *L. infantum*, *L. major*, *L. braziliensis*, *L. adleri* and *L. tarentolae*. Tuzins are
conserved transmembrane proteins in *Trypanosoma* and *Leishmania* associated with surface
glycoprotein expression [78]. They are often contiguous with δ-amastin genes, whose products
are abundant cell surface transmembrane glycoproteins potentially involved in the infection or
survival within macrophages. They are absent in *Crithidia* and *Leptomonas* species, who lack a
vertebrate host stage [78]. Tuzins may play a role in pathogenesis [79], which may be related to
leishmaniasis caused by *L. guyanensis*.

<u>Discussion</u>

L. (Viannia) guyanensis **and** *L. (V.) naiffi* **draft reference genomes**

We assembled high-quality reference genomes for two isolates, *L. (Viannia) guyanensis* LgCL085 and *L. (V.) naiffi* LnCL223, from short read sequence libraries to illuminate genomic diversity in the *Viannia* subgenus and extend previous work [52]. This process combined the *de novo* assembly with a reference-guided approach using the published genome of *L. braziliensis* M2904 to assemble the *L. guyanensis* LgCL085 and *L. naiffi* LnCL223 into 35 chromosomes each (Table 2). An essential feature of this process was to identify and remove contamination in the *L. guyanensis* and *L. braziliensis* M2904 libraries and to trim low-quality bases in *L. naiffi* LnCL223 to ensure that the reads used were informative and free of exogenous impurities. A second screen for contamination in unassigned contigs also removed several *L. guyanensis* LgCL085 contigs, which improved subsequent annotation and gene copy number estimates.

Genomes assembled from short reads capture aneuploidy and nearly all genes

Our strategy was tested by applying the same protocol to the *L. braziliensis* M2904 short read library, which acted as a positive control and quantified the precision of the final output. This facilitated the detection of structural variation or annotation problems, chiefly underestimated copy numbers at certain genes and the incorrect assembly of some loci that were fixed manually. The resulting genomes were largely complete: for comparison, the control *L. braziliensis* M2904 genome had only four homozygous SNPs, 97.2% of the protein coding genes of the reference (231 were missing) and 70 additional genes missed in the reference sequence. These findings highlight scope to resolve *Leishmania* chromosomal architecture more accurately, particularly at repetitive regions and gene arrays, using longer sequencing reads and hybrid assembly approaches.

We showed that the majority of *Viannia* were diploid and had 35 chromosomes. Aneuploidy was evident for *L. guyanensis* LgCL085, *L. guyanensis* M4147, *L. naiffi* LnCL223, *L. naiffi* M5533, *L. lainsoni* M6426, *L. panamensis* WR120 and *L. shawi* M8408 as anticipated [80]. This was verified using read depth allele frequency distributions of reads mapped to *L. braziliensis* M2904 and to their own assemblies.

The *L. guyanensis* LgCL085 genome had more protein coding genes (8,230) than *L. naiffi* LnCL223 (8,104). These numbers were similar to those for *L. panamensis* PSC-1 (7,748) [51] and *L. braziliensis* M2904 (8,357) [48]. The vast majority of protein coding gene models were computationally transferred [81] from the *L. braziliensis* M2904 reference with perfect matching, and were verified and improved manually. Both the *L. guyanensis* and *L. naiffi* reference genomes contained unassigned bin contigs, and chromosomal regions homologous to multiple chromosomal loci or containing partially collapsed gene arrays. 90 (*L. naiffi*) and 92 (*L. guyanensis*) collapsed gene arrays were identified where haploid gene copy numbers were at least twice the assembled copy number when the reads were mapped to the assembled genomes.

A better resolution of the *Viannia* species complexes

This study illustrated that high-throughput sequencing approaches, alignment methods and annotation tools can improve the accuracy of *Leishmania* gene copy number estimates, gene organistion, and genome structure resolution. This yielded insights into features differentiating the isolates examined here, including a 45 Kb duplication on chromosome 34 of most *Viannia*, variable gene repertoires across *Viannia* species, and a potential minichromosome derived from the 3' end of *L. shawi* M8408 chromosome 34. Further work is required to investigate *L. utingensis* and *L. lindenbergi* and other potential distinct lineages [82].

Both single-gene and large-scale copy number variations (CNVs) were tolerated by all

Leishmania genomes. *Leishmania* genomes have extensive conservation of gene content with few species-specific genes [45,48]: here, only 31 *L. guyanensis* LgCL085 and four *L. naiffi* LnCL223 species-specific genes were found. These four genes unique to *L. naiffi* LnCL223, its leishmanolysin hyper-amplification, the 31 genes only in *L. guyanensis* LgCL085 and its tuzin arrays all represent potential targets for improving species-specific typing and better disease surveillance. This is important because infections by the *Viannia* are spread by many hosts and all sources of infections need to be addressed. Immunological screening of anti-*Leishmania* antibodies could be enhanced by genetic testing to identify infections from non-endemic or rarer sources like *L. naiffi*, which has longer parasite survival rates in macrophages *in vitro* [83].

MLSA of 100 *Viannia* isolates across four genes and genome-wide diversity inferred from mapped reads indicated that *L. guyanensis* LgCL085 was closest to *L. panamensis* PSC-1 within the *L. guyanensis* species complex, but was assigned the *L. guyanensis* classification because *L. guyanensis, L. panamensis* and *L. shawi* were a monophyletic species complex as shown by MLSA [56], MLMT [64], *hsp70* [65], internal transcribed spacer (ITS) [84,85], MLEE [86] and RAPD data [87]. Further typing of a more extensive *L. guyanensis, L. panamensis* and *L. shawi* isolate set might clarify if these are distinct species or a single genetic group.

More precise genetic screening of *Viannia* isolates is necessary to trace hybridisation between species. Infection of humans, dogs and *Lu. ovallesi* with *L. guyanensis/L. braziliensis* hybrids was reported in Venezuela [40-41]. A *L. shawi/L. guyanensis* hybrid causing CL was detected in Amazonian Brazil [42], and *L. naiffi* has produced viable progeny with *L. lainsoni* [43] and *L. braziliensis* (Elisa Cupolillo, unpublished data). There is extensive evidence of interbreeding among *L. braziliensis* complex isolates, including more virulent *L. braziliensis/L. peruviana* hybrids with higher survival rates within hosts *in vitro* [44].

Conclusion

This study highlighted the utility of genome sequencing for the identification, characterisation and comparison of *Leishmania* species. We demonstrated that short reads were sufficient for assembly of most *Leishmania* genomes so that SNP, chromosome copy number, structural and somy changes can be investigated comprehensively. The *L. (Viannia) guyanensis* and *L. (V.) naiffi* genomes represent a further advance in refining the taxonomical complexity of the *Viannia* by illustrating their genomic characteristics and the extent to which these are shared across *Viannia* species, which will assist examining the extent to which they can hybridise. This improved understanding of *Leishmania* genomes should be used to explore the complex epidemiology of CL and MCL pathologies in the Americas and the roles of non-human reservoirs and sand flies in these processes. Future work could tackle transmission, drug resistance and pathogenesis in the *Viannia* by applying long-read high-throughput sequencing to examine broader sets of isolates, their genetic diversity, contributions to microbiome variation, and control of transcriptional dosage at gene amplifications.

Methods

L. guyanensis and L. naiffi whole genome sequencing

Extracted DNA for *L. guyanensis* LgCL085 and *L. naiffi* LnCL223 was received from Charité University Medicine (Berlin) at the Wellcome Trust Sanger Institute on 6th Feb 2012. Paired-end 100 bp read Illumina HiSeq 2000 libraries were prepared for both during which *L. guyanensis* required 12 cycles of PCR. The DNA was sequenced (run 7841_5#12) on the 15th (*L. guyanensis*, run 7841_5#12) and 23rd (*L. naiffi*, run 7909_7#9) March 2012. The library preparation, sequencing and read quality verification was conducted as outlined previously [69]. The resulting *L. guyanensis* library contained 15,272,969 reads with a median insert size

of 327.0 (NCBI accession ERX180458) and the *L. naiffi* one had 8,131,246 reads with a median insert size of 335.4 (ERX180449).

Viannia comparative genome, annotation and proteome files
The *L. braziliensis* reference genome (MHOM/BR/1975/M2904) was a positive control whose short reads were examined using the same methods. It was originally sequenced using an Illumina Genome Analyzer II [48] yielding 26,007,384 76 bp paired-end reads with a median insert size of 244.1 bp (ERX005631). Protein sequences were retrieved from the EMBL files using Artemis [88]. Two *L. panamensis* genomes, two *L. peruviana* genome assemblies and five 100 bp paired-end Illumina HiSeq 2000 read libraries of other *Viannia* isolates [53] were used for comparison (Table 1). We included the genomes of *L. panamensis* MHOM/PA/1994/PSC-1, *L. peruviana* PAB-4377 and LEM1537 (MHOM/PE/1984/LC39), and the 100 bp Illumina HiSeq 2000 paired-end reads for each *L. peruviana* PAB-4377 (16,117,316 reads) and *L. peruviana* LEM1537 (9,378,317 reads).

Library quality control, contaminant removal and screening
Figure S9 presents an overview of the bioinformatic steps used in this paper. Quality control of the *L. guyanensis* LgCL085, *L. naiffi* LnCL223, *L. braziliensis* M2904, the five *Viannia* libraries from [53], two *L. peruviana* libraries and *L. panamensis* PSC-1 read library was carried out using FastQC (www.bioinformatics.babraham.ac.uk/projects/fastqc/). No corrections were required for the other libraries. An abnormal distribution of GC content per read observed as an extra GC content peak outside the normal peak for the *L. braziliensis* M2904 and *L. guyanensis* reads indicated sequence contamination that was removed (Figure S10). Two Illumina PCR primers in the *L. braziliensis* M2904 reads were removed (Table S1). Further evaluation using GC content filtering and the non-redundant nucleotide database with BLASTn [89] to remove contaminant sequences (Figure S10) with subsequent correction of read pairing arrangements reduced the initial 52,014,768 reads to 34,592,618 properly paired reads for assembly.

The M2904 reads used to assemble a control genome were used for read mapping, error correction and SNP calling and so the contamination did not affect the published reference. However, it did reduce the number of reads mapped as shown in [48] where only 84% of the *L. braziliensis* M2904 short reads mapped to the *L. braziliensis* assembly, compared with 92% of reads for *L. infantum* reads mapped to its own assembly, 93% of *L. major* reads mapped to its own assembly, and 97% of *L. mexicana* reads mapped to its own assembly.

The 8,131,246 100 bp paired-end *L. naiffi* LnCL223 reads and 15,272,969 100 bp paired-end *L. guyanensis* LgCL085 reads were filtered (Table S1) in the same manner using BLASTn and the smoothness of the GC content distribution to remove putative contaminants. Low quality bases were trimmed at the 3' end of *L. naiffi* LnCL223 reads to remove bases with a phred base quality < 30 using Trimmomatic [90] (Table S1, Figure S11). This resulted in 13,033,846 paired-end *L. guyanensis* LgCL085 sequences and 6,989,814 paired-end *L. naiffi* LnCL223 sequences – 85% and 86% of the initial reads, respectively (Table S1).

Genome evaluation, assembly and optimisation
Processed reads were assembled into contigs using Velvet v1.2.09 and assemblies for all odd numbered k-mer lengths from 21 to 75 were evaluated. The expected k-mer coverage was determined for each assembly using the mode of a k-mer coverage histogram from the velvet-estimate-exp_cov.pl script in Velvet to maximise resolution of repetitive and unique regions [57]. This suggested optimal k-mers of 61 for *L. guyanensis* LgCL085 and 43 for both *L. naiffi* LnCL223 and *L. braziliensis*, which produced assemblies with the highest N50 lengths. Each assembly was assembled with this expected coverage, and contigs were removed if their average k-mer coverage was less than half the expected coverage levels. An expected coverage

of 16 and a coverage cutoff of 8 was applied to *L. naiffi* reads, an expected coverage of 19 and
coverage cutoff of 8.5 to *L. guyanensis* LgCL085, and an expected coverage of 28 and
coverage cutoff of 14 to *L. braziliensis*.

The assembly with the highest N50 for each was scaffolded using SSPACE [58]. In the initial
assemblies, 76% of gaps in scaffolds (3,592/4,754) were closed in for *L. guyanensis* LgCL085,
63% (4,096/6,530) for *L. naiffi* LnCL223, and 67% (4,834/8,786) for *L. braziliensis* using
Gapfiller [58]. Erroneous bases were corrected by mapping reads to the references with iCORN
[91] (Figure S12). Misassemblies detected and broken using REAPR [60] were aligned to the
L. braziliensis M2904 reference (excluding the bin chromosome 00). Scaffolds were evaluated
and broken at putative misassemblies detected from the fragment coverage distribution (FCD)
error and regions with low coverage when the reads were mapped to both broken and unbroken
options. Additionally, the *L. braziliensis* broken and unbroken scaffolds were used to verify
that removing misassemblies prior to (but not after) the contiguation of scaffolds resulted in
more accurate assembled chromosomes. Mis-assembled regions without a gap were replaced
with N bases. REAPR corrected 444 errors in *L. naiffi* LnCL223, of which 59 were caused by
low fragment coverage, 206 in *L. guyanensis* LgCL085 (eight due to low fragment coverage),
and 232 in the *L. braziliensis* control (57 caused by low fragment coverage). Each assembly
step improved the corrected N50 and percentage of error free bases (EFB%) assessed using
REAPR (Table S18), with the sole exception of *L. braziliensis* control at the error-correction
stage, likely due to its higher heterozygosity. The EFB% was the fraction of the total bases
whose reads had no mismatches, matches the expected insert length, had a small FCD error and
at least five read pairs oriented in the expected direction.

Gaps > 100 bp were reduced to 100 bp. 200 bp at the edge of each unplaced scaffold was
aligned with the 200 bp flanking all pseudo-chromosome gaps using BLASTn to verify that no
further gaps could be closed using unplaced scaffolds. Unplaced bin scaffolds < 1 Kb were
discarded, and the resulting assemblies were visualised and compared to *L. braziliensis* using
the Artemis Comparison Tool (ACT) [92]. *L. guyanensis* LgCL085 bin sequences with
BLASTn E-values < 1e-05 and percentage identities > 40% to non-*Leishmania* species in non-
redundant nucleotide database were removed as possible contaminants. The final scaffolds
were contiguated using the *L. braziliensis* reference with ABACAS [59], unincorporated
segments were labelled as unassigned "bin" contigs, and kDNA contigs were annotated as well
(Supplementary Methods).

Phylogenomic MLSA characterisation
A MLSA (multi-locus sequence analysis) approach was adopted to verify the *Leishmania*
species identity using for four housekeeping genes: glucose-6-phosphate dehydrogenase
(G6PD), 6-phosphogluconate dehydrogenase (6PGD), mannose phosphate isomerase (MPI)
and isocitrate dehydrogenase (ICD). Orthologs from other genomes and assemblies were
obtained using BLASTn alignment with thresholds of E-value < 0.05 and percentage identity >
70%. *L. peruviana* LEM-1537 genome had gaps at the MPI and 6PGD genes and was
excluded. The four housekeeping genes spanning 2,902 sites were concatenated in the order
G6PD, 6PGD, MPI and ICD, and aligned using Clustal Omega v1.1 to create a Neighbour-Net
network of uncorrected p-distances using SplitsTree v4.13.1.

Genome annotation and manual curation
Annotation of the *L. guyanensis* LgCL085, *L. naiffi* LnCL223 and *L. braziliensis* control
genomes was completed using Companion [80] using *L. braziliensis* M2904 as the reference as
outlined previously [69], including manual checking and correction of gene models. A control
run with the *L. braziliensis* M2904 reference genome using itself as a reference was performed.
In *L. naiffi* LnCL223, 13 genes and one pseudogene were removed because they overlapped
existing superior gene models that had improved sequence identity with *L. braziliensis* M2904

orthologs. 46 of the protein coding genes were also manually added. 34 of the protein coding
genes on *L. guyanensis* LgCL085 were manually added and one protein coding gene was
removed. 269 gene models on *L. naiffi* LnCL223 and 198 on *L. guyanensis* with multiple joins
mainly caused by the presence of short gaps were corrected by extending the gene model
across the gap where the gap length was known (< 100 bp). If the gap length was unknown (>
100 bp), the gene was extended to the nearest start or stop codon.
587
Measuring ploidy, chromosome copy numbers and CNVs
By mapping the reads with SMALT v5.7 (www.sanger.ac.uk/resources/software/smalt/) to *L.*
braziliensis M2904, the coverage at each site was determined to quantify the chromosome copy
numbers and RDAF distributions at heterozygous SNPs as per previous work [69]. The RDAF
distribution was based on the coverage level of each allele at heterozygous SNPs and this
feature differed across chromosomes for each isolate (Supplementary Results). The median
coverage per chromosome was obtained, and the median of the 35 values combined with the
RDAF distribution mode approximating 50% indicated that all isolates examined here were
mostly diploid (except the triploid *L. braziliensis* M2904). These were visualised with R
packages ggplot2 and gridExtra.
598
After PCR duplicate removal, the mapped reads were used to detect CNVs across genes or
within non-overlapping 10 Kb blocks for all chromosomes and bin contigs using the median
depth values normalised by the median of the chromosome (or bin contig). Loci with a copy
number ≥ 2 were analysed for *L. naiffi* LnCL223, *L. guyanensis* LgCL085 and the *L.*
braziliensis control using their reads mapped to their own assembly. This was also repeated for
reads mapped to the *L. braziliensis* M2904 reference for *L. guyanensis* M4147, *L. naiffi*
M5533, *L. shawi* M8408, *L. lainsoni* M6426, *L. panamensis* WR120, *L. panamensis* PSC-1, *L.*
peruviana LEM1537 and *L. peruviana* PAB-4377. *L. panamensis* PSC-1 reads were mapped to
its own reference genome to verify that we could find previously identified amplified loci, and
we mapped *L. panamensis* WR120 to it so that CNVs shared by both *L. panamensis* could be
obtained. The BAM files of *L. naiffi* LnCL223*, L. guyanensis* LgCL085 and *L. braziliensis*
M2904 reads mapped to its own assembly were visualised in Artemis to confirm and refine the
boundaries of amplified loci.
612
Identification of orthologous groups and gene arrays
Protein coding genes from *L. guyanensis* LgCL085, *L. naiffi* LnCL223 and the *L. braziliensis*
M2904 control genome were produced from the EMBL files for each genome and these were
submitted to the ORTHOMCLdb v5 webserver [93] to identify orthologous groups (OGs).
11,825 OGs with associated gene IDs in at least one of four *Leishmania* species (*L. major*
strain Friedlin, *L. infantum*, *L. braziliensis* and *L. mexicana)* or five *Trypanosoma* species (*T.*
vivax, T. brucei, T. brucei gambiense, *T. cruzi* strain CL Brener and *T. congolense*) were
retrieved from the OrthoMCL database and compared with OGs for each genome. The copy
number of each OG was estimated by summing the haploid copy number of each gene in the
OG. Gene arrays in each genome were identified by finding all OGs with haploid copy number
≥ 2. Large arrays (≥ 10 gene copies) were examined and arrays with unassembled gene copies
were identified by finding those with haploid gene copy number at least twice the assembled
gene number.
626
SNP screening and detection
The filtered reads with Smalt as described mapped above were used for calling SNPs using
Samtools Pileup v0.1.11 and Mpileup v0.1.18 and quality-filtered with Vcftools v0.1.12b and
Bcftools v0.1.17-dev as previously [69] such that SNPs called by both Pileup and Mpileup
post-screening were considered valid. These SNPs all had: base quality >25; mapping quality
>30; SNP quality >30; a non-reference RDAF >0.1; forward-reverse read coverage ratios >0.1
and <0.9; five or more reads; 2+ forward reads, and 2+ reverse reads. Low quality and

repetitive regions of the assemblies were identified and variants in these regions were masked as outlined elsewhere [69]. SNPs were classed as homozygous for an alternative allele to the reference if their RDAF $\geq$ 0.85 and heterozygous if it was > 0.1 and < 0.85.

The high level of nucleotide accuracy of the assembled genomes was indicated by the low rate of homozygous SNPs when the reads mapped to its own assembly (50 for *L. naiffi* LnCL223, 12 for *L. guyanensis* LgCL085, 68 for the *L. braziliensis* reference, and four for the *L. braziliensis* control). Likewise, the numbers and alleles of heterozygous SNPs for the *L. braziliensis* control (25,474) matched that for the reference (25,975), suggesting that the 705 (*L. naiffi* LnCL223) and 14,739 (*L. guyanensis* LgCL085) heterozygous SNPs were accurate. The difference in homozygous and heterozygous SNP rates for *L. braziliensis* here versus the original 2011 study [48] were likely due to differing methodology. The genetic divergence of *L. naiffi* LnCL223 and *L. guyanensis* LgCL085 compared to *L. braziliensis* was quantified using the density of heterozygous and homozygous SNPs per 10 Kb non-overlapping window on each chromosome, visualised using Bedtools.

Data Accessbility
The BioProject ID is PRJEB20208 for *L. guyanensis* LgCL085 and PRJEB20209 for *L. naiffi* LnCL223. The DNA reads are available at the NCBI Short Read Archive (SRA) and and European Nucleotide Archive at ERX180458 for *L. guyanensis* LgCL085 and ERX180449 for *L. naiffi* LnCL223 (these are associated with BioProject PRJEB2600). The consensus genome sequence FASTA files are on Figshare at 10.6084/m9.figshare.5693290 for *L. guyanensis* LgCL085 and 10.6084/m9.figshare.5693272 for *L. naiffi* LnCL223. The chromosome and bin contig annotation EMBL files are at 10.6084/m9.figshare.5693284 for *L. guyanensis* LgCL085 and 10.6084/m9.figshare.5693278 for *L. naiffi* LnCL223. The Supplementary Tables are on Figshare at 10.6084/m9.figshare.5697064. For ease of reader access, the above genome sequence and annotation files, Supplementary Tables and Supplementary Data are also available on the Dryad Digital Repository at: https://doi.org/10.5061/dryad.4bm23.

Acknowledgements
The authors thank: Matthew Berriman and members of the WTSI DNA pipelines team for generating the two sequence libraries; Elisa Cupolillo (Instituto Oswaldo Cruz, Brazil) for discussions and comments on the manuscript; Katrin Kuhls (Technical University of Applied Sciences Wildau), Cathal Seoighe (NUI Galway), Hideo Imamura and Jean-Claude Dujardin (both Institute of Tropical Medicine Antwerp) for help; Anne Stone and Kelly Harkins (both Arizona State University) for releasing valuable sequence read data; and the DJEI/DES/ SFI/HEA Irish Centre for High-End Computing (ICHEC) for computational facilities.

Funding
The authors acknowledge financial support from the NUI Galway Ph.D. Fellowship scheme (S.C.) and the Wellcome Trust core funding of the Wellcome Trust Sanger Institute (WTSI, grant 098051) (J.A.C. and M.S.).

Contributions
S.C. completed the genome assembly, comparative genomics, phylogenetic analysis, mutation investigation, helped design the study and wrote the main manuscript text. S.C., A.S.T. and E.F. completed the genome annotation. M.S. completed genome sequencing. G.S. helped design the study and wrote the main manuscript text. J.A.C. helped design the study and wrote the main manuscript text. T.D. co-ordinated and designed the study and wrote the main manuscript text. All authors gave approval for publication.

Competing interests
The authors have no competing interests.

Figure 1: Middle: A neighbor-Net network of the uncorrected p-distances from concatenated
2,902-base sequences from four housekeeping genes for 102 *Viannia* samples. The genes
were glucose-6-phosphate dehydrogenase (G6PD), 6-phosphogluconate dehydrogenase
(6PGD), mannose phosphate isomerase (MPI) and isocitrate dehydrogenase (ICD). *L. naiffi*
LnCL223 (cyan) is "New_L_naiffi_Reference" and is related to M5533 (IOC_L1365). *L.*
guyanensis LgCL085 (blue) is "New_L_guyanensis_Reference" and is related to the *L. shawi*
M8408 (IOC_L1545) assembly and the *L. panamensis* PSC-1 genome, but less so to *L.*
guyanensis M4147 (iz34). The *L. braziliensis* M2904 reference and control are
"M2904_Reference" and "M2904_Control", proximal to *L. peruviana* PAB-4377. *L. lainsoni*
M6426 (IOC_L1023) (green), *L. utingensis* (orange) and *L. lindenbergi* (pink) are shown.
The isolate names and detail for each species complex is shown by insets in red (*L.*
braziliensis), dark blue (*L. guyanensis*) and light blue (*L. naiffi*). For detailed viewing, the
nexus file can be downloaded at https://figshare.com/s/eecf1c6b42ac4deb6acc and high-
resolution PDF at https://doi.org/10.6084/m9.figshare.5687329**.**

702
703
Figure 2: Normalised chromosome copy numbers of *L. naiffi* LnCL223 reads mapped to its
assembly, *L. guyanensis* LgCL085 reads mapped to its assembly, and *L. guyanensis* M4147,
L. lainsoni M6426, *L. naiffi* M5533, *L. panamensis* WR120 and *L. shawi* M8408 reads
mapped to *L. braziliensis* M2904. Dashed lines indicate disomic, trisomic and tetrasomic
states. Results for *L. panamensis* PSC-1 and *L. peruviana* PAB-4377 were previously
published and are in Figure S3.

Figure 3: Read depth coverage (blue, y-axis) in 10 Kb blocks for reads mapped to *L.*
braziliensis M2904 chromosome 34 (x-axis) for nine *Viannia* isolates. The black horizontal
line is the median chromosome 34 coverage. *L. panamensis* PSC-1 (top left) and *L. shawi*
M8408 (middle left) showed a 3' jump in coverage (green) consistent with an amplification
of inverted repeats that could form a linear minichromosome. In addition, this pair shared a
45 Kb amplification (pink) also found in the *L. braziliensis* M2904 control (bottom left), *L.*
naiffi M5533 (top centre), *L. panamensis* WR120 (middle centre), *L. peruviana* LEM1537
(bottom centre), *L. peruviana* PAB-4377 (top right) and *L. guyanensis* M4147 (middle right).
This was absent in *L. lainsoni* M6426 (bottom right).

Figure 4: Median coverage (blue) in
10 Kb blocks for *L. guyanensis*
LgCL085 reads mapped to its own
assembled chromosome 34 (top) and
L. naiffi LnCL223 reads mapped to its
own assembled chromosome 34
(bottom). The black horizontal line is
the median chromosome 34 read
coverage. There was a 45 Kb
amplification to three copies (pink) in
L. guyanensis LgCL085 (at
chromosome 34 bases 1,195,232-
1,239,355, 44,123 bases in length).
Similarly, there was a 45 Kb four-fold
amplification (pink) in *L. naiffi*
LnCL223 (at chromosome 34 bases
1,206,328-1,251,119, 44,791 bases in
length). The latter encompassed two
additional hypothetical genes relative
to *L. guyanensis* LgCL085. Neither
had evidence of a 3' minichromosome.

Tables

Species	Source	Data Type	Name or WHO Number	SRA*	Number and length of reads	Reference
L. guyanensis	SRA	Reads	LgCL085	ERX180458	15,272,969 (100 bp paired-end)	This study
L. naiffi	SRA	Reads	LnCL223	ERX180449	8,131,246 (100 bp paired-end)	This study
L. braziliensis	Sanger FTP site	Genome & Reads	MHOM/BR/1975/M2904	ERX005631 (LbrM2904 v3)	26,007,384 (76 bp paired-end)	Rogers et al 2011 [48]
L. guyanensis	SRA	Reads	MHOM/BR/1975/M4147	SRX767379	6,225,035 (100 bp paired-end)	Harkins et al 2016 [53]
L. lainsoni	SRA	Reads	MHOM/BR/1981/M6426	SRX764333	4,630,952 (100 bp paired-end)	Harkins et al 2016 [53]
L. naiffi	SRA	Reads	MDAS/BR/1979/M5533	SRX764332	9,646,461 (100 bp paired-end)	Harkins et al 2016 [53]
L. panamensis	Genbank & SRA	Genome & Reads	MHOM/PA/1994/PSC-1	SRX681913; (CP009370: CP009404)	5,875,837 (100 bp paired-end)	Llanes et al 2015 [51]
L. panamensis	SRA	Reads	MHOM/PA/1974/WR120	SRX767384	4,536,341 (100 bp paired-end)	Harkins et al 2016 [53]
L. shawi	SRA	Reads	MCEB/BR/1984/M8408	SRX764331	5,110,479 (100 bp paired-end)	Harkins et al 2016 [53]
L. peruviana	Genbank & SRA	Genome & Reads	PAB-4377	ERX556165 (Bioproject ID: PRJEB7263)	16,117,316 (100 bp paired end)	Valdivia et al 2015 [52]
L. peruviana	Genbank & SRA	Genome & Reads	LEM1537 (MHOM/PE/1984/LC39)	ERX556164 (Bioproject ID: PRJEB7263)	9,378,317 (100 bp paired end)	Valdivia et al 2015 [52]

Table 1: Data used in this study. The World Health Organisation (WHO) numbers are
structured such that M is mammal, R is reptile, HOM is *Homo*, CAN is canine, DAS is
Dasypus (an armadillo), CEB is *Cebus* (a primate), ARV is *Arvicanthis* (a rodent), TAR is
Tarentolae and LAT is *Latastia* (a long-tailed lizard). The top two rows indicate the isolates
for *L. guyanensis* and *L. naiffi* genomes published here. * SRA stands for SRA or TriTrypDB
accession ID.

	L. braziliensis M2904	control	*L. guyanensis* LgCL085	*L. naiffi* LnCL223
Initial number of contigs		13,601	10,308	14,682
Initial contig N50 (Kb)		5.1	9.6	5.7
Number of scaffolds		3,782	2,800	6,530
Scaffold N50 (Kb)		20.6	95.4	24.3
Number of gaps	919	3,352	1,557	3,853
Median read coverage	75	74	56	36
N content (%)	0.29	0.99	0.45	1.07
Chromosomes total length (bp)	31,238,104	28,985,156	28,274,008	29,179,723
Bin sequence total length (bp)	850,747	1,024,497	2,740,314	1,161,372
Total genome length (bp)	32,088,851	30,009,653	31,014,322	30,341,095
Protein coding genes	8,357	8,001	8,230	8,104
Genes on chromosomes	8,432	7,873	7,757	7,952
Genes on bin contigs	188	288	619	310
Total number of genes	8,620	8,161	8,376	8,262

757 **Table 2:** Summary of *L. braziliensis* reference M2904, *L. braziliensis* control, *L. guyanensis*
758 LgCL085, and *L. naiffi* LnCL223 genome assembly contigs, scaffolds, gaps, read coverage,
759 assembled chromosomal and contig sequence, and levels of gene annotation.
760

OG		Genes in OG			OG haploid copy number		
ID	Description	B	G	N	B	G	N
L. guyanensis LgCL085, *L. naiffi* LnCL223 and *L. braziliensis* M2904 control							
OG5_132061	TATE DNA Transposon	2	14	3	21	50	11
OG5_126605	alpha tubulin	1	2	1	17	36	13
OG5_130729	amastin-like surface protein	8	24	20	24	26	33
OG5_126631	elongation factor 1-alpha	1	1	1	12	18	20
OG5_126703	polyubiquitin	2	2	1	21	16	43
OG5_126558	dynein heavy chain, cytosolic	14	13	14	13	13	14
OG5_129265	pteridin transporter; folate/biopterin transporter	8	10	9	14	13	13
OG5_143904	amastin-like surface protein	5	4	6	48	12	14
OG5_126623*	lipophosphoglycan biosynthetic protein / glucose regulated protein 94; heat shock protein 90 / 83-1	2	2	2	11	10	12
L. guyanensis LgCL085 and *L. naiffi* LnCL223							
OG5_126749	GP63 leishmanolysin	4	8	9	5	33	56
OG5_126617	receptor-type adenylate cyclase	3	5	5	5	14	13
OG5_126611	beta tubulin	1	1	2	0	14	27
OG5_128620	NADH-dependent fumarate reductase	4	3	4	2	14	16
OG5_126585	kinesin K39; hypothetical protein	7	10	10	7	12	11
OG5_126573	histone H4	1	7	4	3	11	10
L. naiffi LnCL223							
OG5_173495	hypothetical protein	0	1	1	0	2	15
OG5_126568	ABC1 transporter	9	10	12	10	9	14
OG5_127342	peptidase m20/m25/m40 family protein	2	2	2	6	3	10
L. guyanensis LgCL085							
OG5_173452	tuzin	2	3	1	1	19	1
OG5_145872	ATG8/AUT7/APG8/PAZ2	1	2	1	8	19	1
OG5_143922	ATP dependent DEAD-box helicase	1	2	0	6	17	0
OG5_148241	conserved hypothetical protein	1	1	1	0	14	1
OG5_137181	ATG8/AUT7/APG8/PAZ2	1	1	0	0	13	0
L. braziliensis M2904 control							
OG5_126588*	heat-shock protein hsp70; glucose-regulated protein 78	4	3	3	14	7	8
OG5_138994	tuzin	5	3	2	13	4	2
OG5_129839	phosphoglycan beta 1,3 galactosyltransferase	2	4	1	12	5	2
OG5_127067	thimet oligopeptidase; metallo-peptidase, Clan MA(E), Family M3	3	3	3	12	9	7
L. guyanensis LgCL085 and *L. braziliensis* control							
OG5_169610	surface antigen-like protein	1	1	0	16	11	0
OG5_127518	SLACS like gene retrotransposon element	2	3	1	18	10	1

Table 3: Arrays with ten or more gene copies predicted by read depth for each species. OG stands for orthologous group. Genes in OG shows the number of genes associated with that GO functional category. OG haploid copy number indicates the numbers of haploid gene copies found in each genome: B stands for the *L. braziliensis* M2904 control, G for *L. guyanensis* LgCL085, and N for *L. naiffi* LnCL223. *For OG5_126623 and OG5_126588, the elevated copy number were due to amplificated heat shock protein (*hsp*) genes rather than the glucose regulated protein (*grp*) loci, a potential limitation of OG analyses.

References

1. Pan American Health Organization. 2017 Leishmaniases: Epidemiological Report in the Americas. Available at http://www2.paho.org/hq/index.php?option=com_docman&task=doc_view&Itemid=270&gid=39646&lang=en

2. World Health Organization. 2010 Control of the leishmaniases. World Health Organ Tech Rep Ser. 949, xii-xiii, 1-186, back cover.

3. Gramiccia M, Gradoni L. 2005 The current status of zoonotic leishmaniases and approaches to disease control. Int J Parasitol. 35, 1169-80.

4. Killick-Kendrick R. 1999 The biology and control of phlebotomine sand flies. Clin Dermatol. 17, 279-89.

5. Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L. 2013 Phlebotomine sandflies and the spreading of leishmaniases and other diseases of public health concern. Med Vet Entomol. 27, 123-47. doi: 10.1111/j.1365-2915.2012.01034.x.

6. Walsh JF, Molyneux DH, Birley MH. 1993 Deforestation: effects on vector-borne disease. Parasitology. 106 Suppl, S55-75.

7. Davies CR, Reithinger R, Campbell-lendrum D, Feliciangeli D, Borges R, Rodriguez N. 2000 The epidemiology and control of leishmaniasis in Andean countries. Cad. Saude Pública. 16, 925–50. doi: http://dx.doi.org/10.1590/S0102-311X2000000400013

8. Rotureau B. 2006 Are New World leishmaniases becoming anthroponoses? Med Hypotheses. 67, 1235-41.

9. Lainson R, Shaw JJ. 1989 Leishmania (Viannia) naiffi sp. n., a parasite of the armadillo, Dasypus novemcinctus (L.) in Amazonian Brazil. Ann. Parasitol. Hum. Comp. 64, 3–9

10. Naiff RD, Freitas RA, Naiff MF, Arias JR, Barrett TV, Momen H, Grimaldi Júnior G. 1991 Epidemiological and nosological aspects of Leishmania naiffi Lainson & Shaw, 1989. Mem. Inst. Oswaldo Cruz 86, 317–21.

11. Roque AL, Jansen AM. 2014 Wild and synanthropic reservoirs of Leishmania species in the Americas. Int. J. Parasitol. Parasites Wildl. 3, 251–62. doi: 10.1016/j.ijppaw.2014.08.004.

12. de Souza AAA, da Rocha Barata I, das Graças Soares Silva M, Lima JAN, Jennings YLL, Ishikawa EAY, Prévot G, Ginouves M, Silveira FT, Shaw J, Dos Santos TV. 2017 Natural Leishmania (Viannia) infections of phlebotomines (Diptera: Psychodidae) indicate classical and alternative transmission cycles of American cutaneous leishmaniasis in the Guiana Shield, Brazil. Parasite. 24, 13. doi: 10.1051/parasite/2017016.

13. Arias JR, Miles MA, Naiff RD, Povoa MM, de Freitas RA, Biancardi CB, Castellon EG. 1985 Flagellate infections of Brazilian sand flies (Diptera: Psychodidae): isolation in vitro and biochemical identification of Endotrypanum and Leishmania. Am. J. Trop. Med. Hyg. 34, 1098–108.

14. Kato H, Gomez EA, Yamamoto Y, Calvopiña M, Guevara AG, Marco JD, Barroso PA, Iwata H, Hashiguchi Y. 2008 Natural infection of Lutzomyia tortura with Leishmania (Viannia) naiffi in an Amazonian area of Ecuador. Am. J. Trop. Med. Hyg. 79, 438–40.

15. Azpurua J, De La Cruz D, Valderama A,Windsor D. 2010 Lutzomyia Sand Fly Diversity and Rates of Infection by Wolbachia and an Exotic Leishmania Species on Barro Colorado Island, Panama ed J G Valenzuela PLoS Negl. Trop. Dis. 4, e627.

16. Cássia-Pires R, Boité MC, D'Andrea PS, Herrera HM, Cupolillo E, Jansen AM, Roque AL. 2014 Distinct Leishmania species infecting wild caviomorph rodents (Rodentia: Hystricognathi) from Brazil. PLoS Negl Trop Dis. 8(12), e3389. doi:

10.1371/journal.pntd.0003389.

17. Abba AM, Superina M. 2010 The 2009/2010 armadillo red list assessment BioOne
11, 135–84.

18. Ober HK, Degroote LW, McDonough CM, Mizell RF, Mankin RW. 2011
Identification of an attractant for the nine-banded armadillo, Dasypus novemcinctus
Wildl. Soc. Bull. 35, 421–9.

19. van Thiel PP, Gool TV, Kager PA, Bart A. 2010 First cases of cutaneous
leishmaniasis caused by Leishmania (Viannia) naiffi infection in Surinam. Am. J.
Trop. Med. Hyg. 82, 588–90.

20. Fagundes-Silva GA, Sierra Romero GA, Cupolillo E, Yamashita EP, Gomes-Silva A,
De Oliveira Guerra JA, Da-Cruz AM. 2015 Leishmania (Viannia) naiffi: Rare enough
to be neglected? Mem. Inst. Oswaldo Cruz 110, 797–800.

21. Lainson R, Shaw JJ, Silveira FT, Braga RR, Ishikawa EA. 1990 Cutaneous
leishmaniasis of man due to Leishmania (Viannia) naiffi Lainson and Shaw, 1989.
Ann. Parasitol. Hum. Comp. 65, 282–4.

22. Pratlong F, Deniau M, Darie H, Eichenlaub S, Pröll S, Garrabe E, Le Guyadec T,
Dedet JP. 2002 Human cutaneous leishmaniasis caused by Leishmania naiffi is wide-
spread in South America. Ann. Trop. Med. Parasitol. 96, 781–5.

23. Van der Snoek EM, Lammers AM, Kortbeek LM, Roelfsema JH, Bart A, Jaspers CA.
2009 Spontaneous cure of American cutaneous leishmaniasis due to Leishmania naiffi
in two Dutch infantry soldiers. Clin Exp Dermatol. 34, e889-91. doi: 10.1111/j.1365-
2230.2009.03658.x.

24. Floch H. 1954 Leishmania tropica guyanensis n.sp. agent de la leishmaniose
tegumentarie de Guyanes et de l'Amerique Centralele Arch Inst Pasteur La Guyane
Française du Teritoire L'Inni 15, 328.

25. Lainson R, Shaw JJ, Povoa M. 1981 The importance of edentates (sloths and
anteaters) as primary reservoirs of Leishmania braziliensis guyanensis, causative
agent of "pianbois" in north Brazil. Trans. R. Soc. Trop. Med. Hyg. 75, 611–2.

26. Arias JR, Naif RD, Miles MA, de Souza AA. 1981 The opossum, Didelphis
marsupialis (Marsupialia: Didelphidae), as a reservoir host of Leishmania braziliensis
guyanensis in the Amazon Basin of Brazil. Trans R Soc Trop Med Hyg. 75, 537-41.

27. Dedet JP, Gay F, Chatenay G. 1989 Isolation of Leishmania species from wild
mammals in French Guiana. Trans R Soc Trop Med Hyg. 83, 613-5.

28. Quaresma PF, Rêgo FD, Botelho HA, da Silva SR, Moura Júnior AJ, Teixeira Neto
RG, Madeira FM, Carvalho MB, Paglia AP, Melo MN, Gontijo CM. 2011 Wild,
synanthropic and domestic hosts of Leishmania in an endemic area of cutaneous
leishmaniasis in Minas Gerais State, Brazil. Trans R Soc Trop Med Hyg. 105, 579-85.

29. Lainson R, Shaw JJ, Ward RD, Ready PD, Naiff RD. 1979 Leishmaniasis in brazil:
XIII. Isolation of leishmania from armadillos (dasypus novemcinctus), and
observations on the epidemiology of cutaneous leishmaniasis in north para' state
Trans. R. Soc. Trop. Med. Hyg. 73, 239–42.

30. Quinnell RJ, Courtenay O. 2009 Transmission, reservoir hosts and control of zoonotic
visceral leishmaniasis. Parasitology 136, 1915–34.

31. Ready PD, Lainson R, Shaw JJ, Ward RD. 1986 The ecology of lutzomyia umbratilis
Ward & Fraiha (Diptera: Psychodidae), the major vector to man of Leishmania
braziliensis guyanensis in north-eastern Amazonian brazil Bull. Entomol. Res. 76, 21-
40. doi: https://doi.org/10.1017/S0007485300015248

32. Balbino VQ, Marcondes CB, Alexander B, Luna LK, Lucena MM, Mendes AC,
Andrade PP. 2001 First Report of Lutzomyia (Nyssomyia) umbratilis Ward & Frahia,
1977 outside of Amazonian Region, in Recife, State of Pernambuco, Brazil (Diptera:
Psychodidae: Phlebotominae). Mem. Inst. Oswaldo Cruz 96, 315–7

33. Young DG, Duncan MA. 1994 Guide to the identification and geographic distribution

of Lutzomyia sand flies in Mexico, the West Indies, Central and South America (Diptera: Psychodidae). Amer Entomol Inst, Gainesville.

34. Rodríguez-Barraquer I, Góngora R, Prager M, Pacheco R, Montero LM, Navas A, Ferro C, Miranda MC, Saravia NG. 2008 Etiologic agent of an epidemic of cutaneous leishmaniasis in Tolima, Colombia Am. J. Trop. Med. Hyg. 78, 276–82.

35. Fouque F, Gaborit P, Issaly J, Carinci R, Gantier JC, Ravel C, Dedet JP. 2007 Phlebotomine sand flies (Diptera: Psychodidae) associated with changing patterns in the transmission of the human cutaneous leishmaniasis in French Guiana. Mem. Inst. Oswaldo Cruz 102, 35–40.

36. Lainson R, Shaw JJ, Ready PD, Miles MA, Póvoa M. 1981 Leishmaniasis in Brazil: XVI. Isolation and identification of Leishmania species from sandflies, wild mammals and man in north Para State, with particular reference to L. braziliensis guyanensis causative agent of "pian-bois". Trans. R. Soc. Trop. Med. Hyg. 75, 530–6.

37. Van der Meide WF, Jensema AJ, Akrum RA, Sabajo LO, Lai A Fat RF, Lambregts L, Schallig HD, van der Paardt M, Faber WR. 2008 Epidemiology of cutaneous leishmaniasis in Suriname: a study performed in 2006. Am J Trop Med Hyg. 79, 192-7.

38. Garcia A L, Tellez T, Parrado R, Rojas E, Bermudez H, Dujardin JC. 2007 Epidemiological monitoring of American tegumentary leishmaniasis: molecular characterization of a peridomestic transmission cycle in the Amazonian lowlands of Bolivia Trans. R. Soc. Trop. Med. Hyg. 101, 1208–13.

39. Rotureau B, Ravel C, Nacher M, Couppié P, Curtet I, Dedet JP, Carme B. 2006 Molecular epidemiology of Leishmania (Viannia) guyanensis in French Guiana. J Clin Microbiol. 44, 468-73.

40. Delgado O, Cupolillo E, Bonfante-Garrido R, Silva S, Belfort E, Grimaldi Jr G, Momen H. 1997 Cutaneous Leishmaniasis in Venezuela Caused by Infection with a New Hybrid between Leishmania (Viannia) braziliensis and L. (V.) guyanensis Mem. Inst. Oswaldo Cruz. 92, 581–2.

41. Bonfante-Garrido R, Meléndez E, Barroeta S, de Alejos MA, Momen H, Cupolillo E, McMahon-Pratt D, Grimaldi G. 1992 Cutaneous leishmaniasis in western Venezuela caused by infection with Leishmania venezuelensis and L. braziliensis variants. Trans. R. Soc. Trop. Med. Hyg. 86, 141–8.

42. Jennings Y L, de Souza AAA, Ishikawa E A, Shaw J, Lainson R, Silveira F. 2014 Phenotypic characterization of Leishmania species causing cutaneous leishmaniasis in the lower Amazon region, western Pará state, Brazil, reveals a putative hybrid parasite, Leishmania (Viannia) guyanensis × Leishmania (Viannia) shawi shawi. Parasite 21, 39.

43. Tojal da Silva AC, Cupolillo E, Volpini AC, Almeida R, Romero GA. 2006 Species diversity causing human cutaneous leishmaniasis in Rio Branco, state of Acre, Brazil. Trop Med Int Health. 11, 1388-98.

44. Cortes S, Vaz Y, Neves R, Maia C, Cardoso L, Campino L. 2012 Risk factors for canine leishmaniasis in an endemic Mediterranean region. Vet Parasitol. 189, 189-96. doi: 10.1016/j.vetpar.2012.04.028.

45. Peacock CS, et al. 2007 Comparative genomic analysis of three Leishmania species that cause diverse human disease. Nat. Genet. 39, 839–47.

46. Martínez-Calvillo S, Yan S, Nguyen D, Fox M, Stuart K, Myler PJ. 2003 Transcription of Leishmania major Friedlin chromosome 1 initiates in both directions within a single region. Mol. Cell 11, 1291–1299.

47. Clayton C, Shapira M. 2007 Post-transcriptional regulation of gene expression in trypanosomes and leishmanias Mol. Biochem. Parasitol. 156, 93–101.

48. Rogers MB, et al. 2011 Chromosome and gene copy number variation allow major structural change between species and strains of Leishmania. Genome Res. 2, 2129–

926   42.

927 49. Wincker P, Ravel C, Blaineau C, Pages M, Jauffret Y, Dedet JP, Bastien P. 1996 The
928   Leishmania genome comprises 36 chromosomes conserved across widely divergent
929   human pathogenic species. Nucleic Acids Res. 24, 1688–94.

930 50. Britto C, Ravel C, Bastien P, Blaineau C, Pagès M, Dedet JP, Wincker P. 1998
931   Conserved linkage groups associated with large-scale chromosomal rearrangements
932   between Old World and New World Leishmania genomes. Gene. 222, 107–17.

933 51. Llanes A, Restrepo CM, Del Vecchio G, Anguizola FJ, Lleonart R. 2015 The genome
934   of Leishmania panamensis: insights into genomics of the L. (Viannia) subgenus. Sci.
935   Rep. 5, 8550.

936 52. Valdivia HO, et al. 2015 Comparative genomic analysis of Leishmania (Viannia)
937   peruviana and Leishmania (Viannia) braziliensis. BMC Genomics 16, 715.

938 53. Harkins KM, Schwartz RS, Cartwright RA, Stone AC. 2016 Phylogenomic
939   reconstruction supports supercontinent origins for Leishmania. Infect. Genet. Evol.
940   38, 101–9.

941 54. Oddone R, Schweynoch C, Schönian G, De Sousa CDS, Cupolillo E, Espinosa D,
942   Arevalo J, Noyes H, Mauricio I, Kuhls K. 2009 Development of a multilocus
943   microsatellite typing approach for discriminating strains of Leishmania (Viannia)
944   species J. Clin. Microbiol. 47, 2818–25.

945 55. Lye Lye LF, Owens K, Shi H, Murta SM, Vieira AC, Turco SJ, Tschudi C, Ullu E,
946   Beverley SM. 2010 Retention and loss of RNA interference pathways in
947   trypanosomatid protozoans. PLoS Pathog. 6, e1001161. doi:
948   10.1371/journal.ppat.1001161.

949 56. Boité MC, Mauricio IL, Miles MA, Cupolillo E. 2012 New insights on taxonomy,
950   phylogeny and population genetics of Leishmania (Viannia) parasites based on
951   multilocus sequence analysis. PLoS Negl. Trop. Dis. 6, e1888.

952 57. Zerbino DR, Birney E. 2008 Velvet: algorithms for de novo short read assembly using
953   de Bruijn graphs. Genome Res. 18, 821–9.

954 58. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011 Scaffolding pre-
955   assembled contigs using SSPACE. Bioinformatics 27, 578–9.

956 59. Assefa S, Keane TM, Otto TD, Newbold C, Berriman M. 2009 ABACAS: algorithm-
957   based automatic contiguation of assembled sequences. Bioinformatics 25, 1968–9.

958 60. Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD. 2013 REAPR: a
959   universal tool for genome assembly evaluation. Genome Biol. 14, R47.

960 61. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
961   Remmert M, Söding J, Thompson JD, Higgins DG. 2011 Fast, scalable generation of
962   high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst.
963   Biol. 7, 539.

964 62. Huson DH, Bryant D. 2006 Application of phylogenetic networks in evolutionary
965   studies. Mol. Biol. Evol. 23, 254–67.

966 63. Schönian G, Mauricio I, Cupolillo E. 2010 Is it time to revise the nomenclature of
967   Leishmania? vol 26, 466-469.

968 64. Kuhls K, Cupolillo E, Silva SO, Schweynoch C, Boité MC, Mello MN, Mauricio I,
969   Miles M, Wirth T, Schönian G. Population structure and evidence for both clonality
970   and recombination among Brazilian strains of the subgenus Leishmania (Viannia).
971   PLoS Negl Trop Dis. 2013 7(10): e2490. doi: 10.1371/journal.pntd.0002490.

972 65. Fraga J, Montalvo AM, De Doncker S, Dujardin JC, Van der Auwera G. 2010
973   Phylogeny of Leishmania species based on the heat-shock protein 70 gene. Infect
974   Genet Evol 10, 238–245.

975 66. Segovia M, Ortiz G. 1997 LDI amplifications in Leishmania Parasitol. Today 13,
976   342–8.

977 67. Fu G, Melville S, Brewster S, Warner J, Barker DC. 1998 Analysis of the genomic

organisation of a small chromosome of Leishmania braziliensis M2903 reveals two genes encoding GTP-binding proteins, one of which belongs to a new G-protein family and is an antigen. Gene 210, 325–33.

68. Raymond F, Boisvert S, Roy G, Ritt JF, Légaré D, Isnard A, Stanke M, Olivier M, Tremblay MJ, Papadopoulou B, Ouellette M, Corbeil J. 2012 Genome sequencing of the lizard parasite Leishmania tarentolae reveals loss of genes associated to the intracellular stage of human pathogenic species. Nucleic Acids Res. 40, 1131–47.

69. Coughlan S, Mulhair P, Sanders M, Schonian G, Cotton JA, Downing T .2017 The genome of Leishmania adleri from a mammalian host highlights chromosome fission in Sauroleishmania Sci. Rep. 7, 43747.

70. Hartley MA, Drexler S, Ronet C, Beverley SM, Fasel N. 2014 The immunological, environmental, and phylogenetic perpetrators of metastatic leishmaniasis Trends Parasitol. 30, 412–22.

71. Acestor N, Masina S, Ives A, Walker J, Saravia NG, Fasel N. 2006 Resistance to oxidative stress is associated with metastasis in mucocutaneous leishmaniasis. J. Infect. Dis. 194, 1160–7.

72. Eisen JA, et al. 2006 Macronuclear genome sequence of the ciliate Tetrahymena thermophila, a model eukaryote PLoS Biol. 4, e286.

73. Steinkraus HB, Greer JM, Stephenson DC, Langer PJ. 1993 Sequence heterogeneity and polymorphic gene arrangements of the Leishmania guyanensis gp63 genes Mol. Biochem. Parasitol. 62, 173–85.

74. Joshi PB, Sacks DL, Modi G, McMaster WR. 1998 Targeted gene deletion of Leishmania major genes encoding developmental stage-specific leishmanolysin (GP63) Mol. Microbiol. 27, 519–30.

75. Joshi PB, Kelly BL, Kamhawi S, Sacks DL, McMaster WR. 2002 Targeted gene deletion in Leishmania major identifies leishmanolysin (GP63) as a virulence factor. Mol. Biochem. Parasitol. 120, 33–40.

76. Olivier M, Atayde VD, Isnard A, Hassani K, Shio MT. 2012 Leishmania virulence factors: focus on the metalloprotease GP63 Microbes Infect. 14, 1377–89.

77. Brittingham A, Morrison CJ, McMaster WR, McGwire BS, Chang KP, Mosser DM. 1995 Role of the Leishmania surface protease gp63 in complement fixation, cell adhesion, and resistance to complement-mediated lysis. J. Immunol. 155, 3102–11.

78. Jackson AP. 2010 The evolution of amastin surface glycoproteins in trypanosomatid parasites. Mol. Biol. Evol. 27, 33–45.

79. Lakshmi BS, Wang R, Madhubala R. 2014 Leishmania genome analysis and high-throughput immunological screening identifies tuzin as a novel vaccine candidate against visceral leishmaniasis. Vaccine 32, 3816–22.

80. Mannaert A, Downing T, Imamura H, Dujardin JC. 2012 Adaptive mechanisms in pathogens: universal aneuploidy in Leishmania. Trends Parasitol. 28, 370–6.

81. Steinbiss S, Silva-Franco F, Brunk B, Foth B, Hertz-Fowler C, Berriman M, Otto TD. 2016 Companion: a web server for annotation and analysis of parasite genomes Nucleic Acids Res. 44, W29–34.

82. Akhoundi M, Downing T, Votýpka J, Kuhls K, Lukeš J, Cannet A, Ravel C, Marty P, Delaunay P, Kasbari M, Granouillac B, Gradoni L, Sereno D. 2017 Leishmania infections: Molecular targets and diagnosis. Mol Aspects Med. 57, 1-29. doi: 10.1016/j.mam.2016.11.012.

83. Matta NE, Cysne-Finkelstein L, Machado GM, Da-Cruz AM, Leon L. 2010 Differences in the antigenic profile and infectivity of murine macrophages of Leishmania (Viannia) parasites. J Parasitol. 96, 509-15. doi: 10.1645/GE-2241.1.

84. Cupolillo E, Grimaldi Júnior G, Momen H, Beverley SM. 1995 Intergenic region typing (IRT): a rapid molecular approach to the characterization and evolution of Leishmania. Mol. Biochem. Parasitol. 73, 145–55.

1030   85.   Berzunza-Cruz M, Cabrera N, Crippa-Rossi M, Sosa Cabrera T, Pérez-Montfort R,
1031   Becker I. 2002 Polymorphism analysis of the internal transcribed spacer and small
1032   subunit of ribosomal RNA genes of Leishmania mexicana. Parasitol. Res. 88, 918–25.
1033   86.   Cupolillo E, Grimaldi G, Momen H. 1994 A general classification of new world
1034   Leishmania using numerical zymotaxonomy Am. J. Trop. Med. Hyg. 50, 296–311.
1035   87.   Bañuls A L, Jonquieres R, Guerrini F, Le Pont F, Barrera C, Espinel I, Guderian R,
1036   Echeverria R, Tibayrenc M. 1999 Genetic analysis of leishmania parasites in Ecuador:
1037   are Leishmania (Viannia) panamensis and Leishmania (V.) Guyanensis distinct taxa?
1038   Am. J. Trop. Med. Hyg. 61, 838–45.
1039   88.   Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012 Artemis: an
1040   integrated platform for visualization and analysis of high-throughput sequence-based
1041   experimental data. Bioinformatics. 28, 464-9. doi: 10.1093/bioinformatics/btr703.
1042   89.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
1043   2009 BLAST+: architecture and applications. BMC Bioinformatics 10, 421.
1044   90.   Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: a flexible trimmer for Illumina
1045   sequence data. Bioinformatics 30, 2114-20. doi: 10.1093/bioinformatics/btu170.
1046   91.   Otto TD, Sanders M, Berriman M, Newbold C. 2010 Iterative Correction of
1047   Reference Nucleotides (iCORN) using second generation sequencing technology.
1048   Bioinformatics. 26, 1704-7. doi: 10.1093/bioinformatics/btq269.
1049   92.   Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J.
1050   2005 ACT: the Artemis Comparison Tool. Bioinformatics 21, 3422–3.
1051   93.   Li L, Stoeckert CJ Jr, Roos DS. 2003 OrthoMCL: identification of ortholog groups for
1052   eukaryotic genomes. Genome Res. 13, 2178-89.
1053