Title:

Medaka population genome structure and demographic history described via genotyping-by-

sequencing

Authors and Affiliations:

Takafumi Katsumura[1,2*], Shoji Oda[3], Mitani Hiroshi[4], Hiroki Oota[1*].

[1] Department of Anatomy, Kitasato University School of Medicine

1-15-1 Kitasato, Minami-Ku Sagamihara, Kanagawa 252-0374, Japan

[2] Graduate School of Natural Science and Technology, Okayama University, 1-1-1

Tsushimanaka, Kita-Ku, Okayama 700-8530, Japan

[3] Department of Integrated Biosciences, Graduate School of Frontier Sciences, University of

Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8562, Japan

*Authors for Correspondence:

Takafumi Katsumura, Graduate School of Natural Science and Technology, Okayama

University, Okayama, Japan, +81 86 252 7860, tk@okayama-u.ac.jp

Hiroki Oota, Department of Anatomy, Kitasato University School of Medicine, Kanagawa,

Japan, +81 42 778 9022, hiroki_oota@med.kitasato-u.ac.jp

Abstract

Medaka is a model organism in medicine, genetics, developmental biology and population

genetics.   Lab stocks composed of more than 100 local wild populations are available for

research in these fields.   Thus, medaka represents a potentially excellent bioresource for

screening disease-risk- and adaptation-related genes in genome-wide association studies.

Although the genetic population structure should be known before performing such an

analysis, a comprehensive study on the genome-wide diversity of wild medaka populations

has not been performed.   Here, we performed genotyping-by-sequencing (GBS) for 81 and

12 medakas captured from a bioresource and the wild, respectively.   Based on the GBS data,

we evaluated the genetic population structure and estimated the demographic parameters

using an approximate Bayesian computation (ABC) framework.   The autosomal data

confirmed that there were substantial differences between local populations and supported our

previously proposed hypothesis on medaka dispersal based on mitochondrial genome

(mtDNA) data.   A new finding was that a local group that was thought to be a hybrid

between the northern and the southern Japanese groups was actually a sister group of the

northern Japanese group.   Thus, this paper presents the first population-genomic study of

medaka and reveals its population structure and history based on autosomal diversity.

Key words: local population, freshwater fish, demography, RAD-seq, bioresource

Introduction

Medaka (*Oryzias latipes*) is a small fresh-water fish native to East Asia that has attracted attention as a vertebrate model for population genetics (Oota and Mitani 2011; Spivakov *et al.* 2014). Wild medaka populations have been maintained in certain universities and research institutes as a bioresource (hereafter, wild lab stocks) with funding from the Japanese government since 1985 (Shima *et al.* 1985). These populations consist of more than 100 local-wild populations that have various phenotypic traits (Watanabe-Asaka *et al.* 2014; Igarashi *et al.* 2017) and abundant genetic diversity (Kasahara *et al.* 2007). Geographical features affect the population structure of organisms. Seas and mountains restrict the movement of terrestrial animals and freshwater fish, and the climate also changes the breeding timing and foraging environment. Particularly, noticeable climate differences are observed within island groups, such as the Japanese archipelago, where a large latitudinal difference exists between the southern and northern ends. Therefore, animals are exposed to various selective pressures according to the geographic environment. The local populations have differentiated into local groups by genetic drift, resulting in genetically divergent groups. Medakas stocked as a bioresource are thought to have retained the genetically adapted traits they acquired from various environments of the Japanese archipelago.

Our ultimate goal in exploiting medaka characteristics is to establish an experimental system for testing the functional differences between alleles detected by, for instance, genome-wide association studies. Once genetic polymorphisms related to phenotypic traits in medaka are detected, *in vivo* experiments, such as genome editing experiments (Ansai and Kinoshita 2014), can be conducted to understand the functions of the genetic variants (Shimmura *et al.* 2018). Revealing the functional difference between alleles in wild populations allows us to infer the role of genetic polymorphisms in human homologous genes (Igarashi *et al.* 2017; Shimmura *et al.* 2018).

Most previous analyses of medaka genetic diversity and population structure have

been conducted using mitochondrial DNA (mtDNA).   Medaka is divided into four

mitochondrial groups: the northern Japanese (N.JPN), southern Japanese (S.JPN), eastern

Korean (E.KOR) and western Korean/Chinese groups (W.KOR) (Sakaizumi *et al.* 1980;

1983; Sakaizumi 1986; Katsumura *et al.* 2009).   These groups do not have sympatric

habitats.   Although previous allozyme-based studies show that an ambiguous group exists in

the geographic boundary region known as Tajima-Tango (see also fig. 1, supplementary fig

1), between N.JPN and S.JPN, which is thought to be a hybridization of the two groups

(Sakaizumi 1984; Takehana *et al.* 2016), the process has not been fully verified.

Based on mtDNA cytochrome *b* gene sequences, N.JPN and S.JPN are composed of 3

and 11 subgroups, respectively (Takehana *et al.* 2003).   Each subgroup is composed of local

populations, and the between-population genetic diversities are greater than the within-

population genetic diversities, indicating substantial genetic differentiation between local

populations (Katsumura *et al.* 2009).   Their habitat environments are also largely different.

For instance, there is a large amount of snowfall in the habitats of N.JPN, where the breeding

season is short.   The habitat of S.JPN is wider than that of N.JPN, and its climate

environment is also diverse; e.g., the differences of the annual average temperature and

rainfall between the south end (Nago in Okinawa) and north end (Ichinoseki in Iwate)

inhabited by S.JPN were 11.6 °C and 988 mm in 2017, respectively

(http://www.jma.go.jp/jma/indexe.html).   The phylogenetic data from these studies also

suggested that N.JPN and S.JPN have been spreading in the Japanese archipelago at different

times.   Particularly, the origin of S.JPN, which has the largest habitat, has been suggested to

be the northern part of Kyushu Island based on mtDNA ("Out of Northern Kyushu"

hypothesis) (Katsumura *et al.* 2012).   However, mtDNA is insufficient to describe the

population structure and history because of its single locus.   Genetic diversity based on

autosomes is essential to understand the medaka population structure, but the comprehensive

data of autosomes are still limited.

An inference of the population structure and history induced by autosomal information

is more robust than that induced by single loci.   To unravel the population structure based on

autosomes, we comprehensively performed a population-genetic analysis based on autosomal

single-nucleotide polymorphisms (SNPs) using all 81 local populations maintained as wild

lab stocks at the University of Tokyo.   We examined an individual sampled from each

population stock and 12 wild individuals captured from the northern part of Kyushu Island,

where (Katsumura *et al.* 2012) the medakas currently distributed along the entire Pacific side

of the Japanese archipelago originated, to estimate each population's time of expansion from

Northern Kyushu.   To obtain SNP data, we conducted genotyping-by-sequencing (GBS)

(Elshire *et al.* 2011; Narum *et al.* 2013) using a high-throughput sequencer.   This method,

which allowed us to cost-effectively genotype tens of thousands of SNPs (Andrews *et al.*

2016), resulted in an accurate enhancement of the population-genetic estimations because the

variance in the demographic parameter estimates decreased when we used many SNPs.

Eventually, we obtained more than ten thousand SNPs from eighty-one wild lab stocks and

twelve wild-captured medakas.   Here, we re-evaluated the medaka genetic diversity and

population structures based on these SNPs and reconstructed the population history by

assessing three demographic events.   These data redefine the medaka local groups and

provide a basis of the population history for discussing the role of phenotype-associated

alleles in the context of adaptation.

Materials and Methods

Samples

We sampled 81 male medakas from 81 wild lab stocks (14 from N.JPN, 56 from S.JPN, 5

from Tajima-Tango in the geographic boundary region between N.JPN and S.JPN, 3 from

E.KOR, and 3 from W.KOR; note that these groupings come from previous mtDNA

sequences and allozyme patterns) at the University of Tokyo in 2014.   The lab stocks

originated from geographically distinct populations in East Asia and have been maintained

since 1985 (Shima *et al.* 1985) as closed colonies in the Graduate School of Frontier Sciences,

the University of Tokyo, Kashiwa City (fig. 1).   These lab stocks maintain the genetic

diversity originating from their habitat (Katsumura *et al.* 2014; Igarashi *et al.* 2017) and show

less diversity within populations than between populations (Katsumura *et al.* 2009).

Because of these characteristics, we considered that one individual sampled from the lab stock

would be adequate to represent its originating population.   In addition, we used two wild-

captured medaka populations from the Saga Prefecture in the northern part of Kyushu, Japan.

One population was the Ogi (SO) population from Southern Saga, and the other population

was the Umeshiro (US) population from Northern Saga, both of which were captured in

September 2010 (see Katsumura *et al*. 2012).   Six of 48 medakas from each wild-captured

population were selected randomly and analyzed via GBS.

DNA extraction and genotyping-by-sequencing

One-third of the medaka body was dissolved in a 600 µl lysis buffer containing 1.24% SDS,

0.124 M EDTA and 0.062 mg/ml proteinase K (final concentrations).   The total genomic

DNA was extracted and purified using phenol-chloroform and isopropanol precipitation.

After a 70% EtOH wash, an isolated DNA pellet was resuspended in 100 µl TE buffer and

then treated with RNase A (final conc. 1 mg/ml) for 1 hour at room temperature. Then, the

DNA was purified again using phenol-chloroform and isopropanol precipitation. For ninety-

three samples, the GBS process was outsourced to Macrogen Japan in Kyoto. The

procedures for constructing libraries and performing Illumina HiSeq 2000 single-end

sequencing were the same as those described by Poland *et al.* (Poland *et al.* 2012) except for

the use of the restriction enzyme ApeKI instead of EcoRI–MspI. The sequence lengths were

51 bp and included each individual in-line barcode (4–9 bp) for the individual sample. The

data have been submitted to the DDBJ Sequence Read Archive (DRA) database under project

accession ID: DRA006353.

Quality filtering and SNP extraction

Our single-end reads were filtered using *FASTQ Quality Filter* in *FASTX-Toolkit version*

0.0.13 (http://hannonlab.cshl.edu/fastx-toolkit) using the following options: -Q 33 -v -z -

q 30 -p 90. The draft genome of the medaka sequenced by the PacBio sequencer (Medaka-

Hd-rR-pacbio_version2.0.fasta; http://utgenome.org/medaka_v2/#!Assembly.md) was used to

align the reads using *BWA backtrack 0.7.12-r1039* (Li and Durbin 2009) using the "-n 0.06"

option. After the mapping process, the multi-mapped reads were removed using *Samtools*

v1.2 (Li *et al.* 2009) and the "-Sq 20" option. Following this pipeline, sequencing of the

ApeKI-digested GBS libraries generated an average of 3.06 million reads per individual

before any quality filtering. The read numbers ranged from 1.89 to 4.10 million reads per

individual. After quality filtering, 2.76 million (90.2%) sequences per individual on average

were retained, and 0.30 million (9.8%) sequences were eliminated. The retained sequences

presented a mean quality score of 38.7 and a GC content of 47.8%. An average of 1.84

million of the retained sequences (66.7%) aligned to the medaka autosomal genome, and 0.92

million sequences (33.3%) were not aligned and discarded because they mapped to non-

autosomal loci (mitochondrial genome and unanchored contigs) and multiple loci.

The *Stacks* pipeline (version 1.35) and the Stacks workflow

(https://github.com/enormandeau/stacks_workflow) were used to generate SNPs and

sequences for each individual separately (Catchen *et al.* 2011; 2013).   The selected *Stacks*

parameters were as follows: minimum stack depth (-m), 3; and number of mismatches when

building the catalog (-n), 1.   In the 'rx' step of *Stacks*, we used the bounded SNP model set

to 0.1, the ε upper bound set to 0.1 and the log likelihood set to 10.   Following this pipeline,

five datasets were constructed using the *population* program in *Stacks*.

The first was the "PopStat" dataset, which was generated using the "-p 5 -r 0.66"

options and mitochondrial grouping (N.JPN, Tajima-Tango, S.JPN, E.KOR, W.KOR; see also

fig. 1) to calculate the population-genetic statistics (table 1) using the loci shared across all

groups and sequenced in one or more populations in each group.

The second was the "Global" dataset, which was generated using the "-p 58 -r 1.00"

option and without the mitochondrial grouping, to examine the phylogenetic relationships

between the geographic populations and the population structure within the species.

The third and the fourth were the "HZ-1" and "HZ-2" datasets ("HZ" is the

abbreviation of "Hybrid zone"), respectively, including the 15 boundary populations (N.JPN:

Kaga, Maiduru, Miyadu, Obama, and Sabae; Tajima-Tango: Amino, Hamasaka, Kinosaki,

Kumihama, and Toyooka; and Honshu: Ayabe, Iwami, Kasumi, Matsue, and Tottori) with or

without the Kyushu populations (Kyushu: Fukue, Hiwaki, Izumi, Kadusa, and Kikai). These

datasets were generated using the "-p 3 -r 0.70" (without Kyushu) and the "-p 4 -r 0.70" (with

Kyushu) option, respectively, to assess the genetic population structure and the history of the

boundary population for the Tajima-Tango group.

The fifth was the "Local" dataset, including two Kyushu deme samples: one was

Umeshiro (US), which was sampled in the northern part of the Saga prefecture, and the other

was Ogi (SO), which was sampled in the southern part of the Saga prefecture. These samples

were used to estimate the time of Honshu's population divergence from Kyushu to infer the

timing of the "Out of Northern Kyushu" event (Katsumura *et al.* 2012).

Note that we define the terms "deme samples" and "non-deme samples" as

"samples from the local-wild population" and "samples from the wild lab stocks",

respectively (see details in Katsumura *et al.* 2009).

Against those datasets (except "PopStat" and "HZ-1"), the SNPs with strong linkage

disequilibrium were randomly removed, one from each pair of SNPs with $r^2 > 0.2$, using

Plink1.9 (Chang *et al.* 2015) and the "--indep-pairwise 12.5 5 0.2 --autosome-num 24" option.

These values were set based on a medaka population genomics study (Spivakov *et al.* 2014).

Finally, the "Global", "HZ-2" and "Local" datasets contained 8,361 SNPs out of 13,177 SNPs,

1,014 SNPs out of 2,661 SNPs and 698 SNPs out of 2,246 SNPs, respectively.

Genetic clustering analysis

To obtain a genetic overview of the relationship of medaka geographic populations, we

performed a principal component analysis (PCA) of the "Global" dataset as implemented in

the *SNPRelate* program (Zheng *et al.* 2012) in R version 3.2.2.   Additionally, to examine the

genetic relationships between medakas in the Japanese archipelago, we used the subdataset

without the E/W.KOR and Chinese populations, which included 7,126 SNPs.   We also

performed a model-based genetic clustering analysis using *ADMIXTURE v1.23* (Alexander *et*

al. 2009) to estimate the proportions of ancestral medaka populations.   We ran 50 replicates

with random seeds for the number of clusters (K) from 1 to 9 and calculated the mean of the

lowest fivefold cross-validation errors for each K (Jeong *et al.* 2016).   Values of K = 4, 5,

and 6 showed the first-, second- and third-lowest fivefold cross-validation errors, respectively

(supplementary fig. 2).   The results of the genetic clustering analyses were visualized by a

ggplot2 package in R (Wickham 2011).

Reconstructing the phylogenetic tree using the maximum likelihood method

We considered an individual as representative of a population and generated the individual

sequences using the *population* program with the "-p 58 -r 1.00 --phylip_all" option in the

Stacks pipeline.   The dataset included 4,638 partitions and 217,986 bp of nucleotide

sequences to compensate for the loci that could not be sequenced by invariable sites across all

samples and "N" at the variable site.   The dataset including 4,638 partitions and 217,986 bp

nucleotide sequences was analyzed via the *IQ-TREE* program (Nguyen *et al.* 2015) to

reconstruct a maximum likelihood tree with model selection for each partition (Chernomor *et*

al. 2016) and 1,000 SH-aLRT/ultrafast-bootstrappings (Guindon *et al.* 2010; Minh *et al.*

2013).   Then, we used the *FigTree* program (http://tree.bio.ed.ac.uk/software/figtree/) to

visualize the phylogenetic tree.

Inference of demographic parameters of the populations in and around Tajima-Tango

and the time of "Out of Northern Kyushu" based on the ABC framework

Using the HZ-2 dataset, which added five Kyushu populations to 15 boundary populations of

the HZ-1 data set, the population history, including the possibility that the Tajima-Tango

group occurred by the admixture of N.JPN and S.JPN, was inferred from 1,014 SNPs using

DIYABC ver2.1.0 (Cornuet *et al.* 2014).   We tested four evolutionary scenarios in which

N.JPN and S.JPN diverged from an ancestral population at time t3: (I) Tajima-Tango

originated in N.JPN: Honshu and Kyushu diverged at time t2, and then Tajima-Tango

diverged from N.JPN at time t1 (Takehana *et al.* 2016); (II) Admixture of N.JPN and Honshu:

Honshu and Kyushu diverged at time t2, and then Tajima-Tango occurred at time t1 by an

admixture with rate r between N.JPN and Honshu (Sakaizumi 1984); (III) N.JPN originated in

Tajima-Tango: Honshu and Kyushu diverged at time t2 and then N.JPN diverged from

Tajima-Tango at time t1; and (IV) Honshu diverged from Kyushu and then Tajima-Tango

occurred by an admixture with Honshu of rate r.   For simplicity, all populations were

assumed to have constant effective sizes in each lineage (i.e., no bottleneck and expansion).

The same prior parameters were defined for four scenarios based on previous studies, and the

prior distribution of each parameter is presented in table 2.   In addition, we set the

conditional constraint as follows: t3 > t2, t3 > t1 and t2 ≥ t1.   In total, four million

simulations were run, which provided approximately one million simulations for each

scenario.

The summary statistics for all SNP loci included the proportion of loci with null

gene diversity, mean gene diversity across polymorphic loci, proportion of loci with null Nei's

distance between the two samples, variance across loci of non-null Nei's distances between

two samples, proportion of loci with null admixture estimates, mean across loci of non-null

admixture estimates, variance across loci of non-null admixture estimates and mean across all

locus admixture estimates.   The 1% simulated datasets closest to the observed dataset were

used to estimate the posterior parameter distributions through a weighted local linear

regression procedure (Beaumont *et al.* 2002).   Scenarios were compared by estimating their

posterior probabilities using the direct estimation and logistic regression methods

implemented in *DIYABC* (Cornuet *et al.* 2014).   We also estimated the time for "Out of

Northern Kyushu" using the Local dataset that included 698 SNPs from *DIYABC*.   We

inferred the following time-based simple evolutionary scenario: N.Saga diverged from S.Saga

at time t because this divergence event was consistent with the "Out of Northern Kyushu"

event (Katsumura *et al.* 2012).   In this case, all populations were assumed to have

unchangeable effective sizes in each lineage.

The prior distribution of each parameter is presented in table 2, and the 1%

simulated datasets closest to the observed dataset were used to estimate the posterior

parameter distributions.   We evaluated the accuracy of the demographic parameter

estimation by calculating accuracy indicators (*Bias and mean square error* option) in *DIYABC*

and accepted the parameters that were non-scaled and scaled by the mean effective population

size for the HZ-2 and Local datasets, respectively.   Using the infinite-sites model in the

Wright-Fisher populations of a constant size, a rough value of N_e was estimated using the

relationship $\pi = 4N_e\mu$ (Tajima 1983), where π is genome-wide nucleotide diversity and μ is

the mutation rate per site per generation (Osada 2015).   We calculated N_e using the above

formula with the following values: $\mu = 10^{-9}$/site/generation; generation = 1 year, which

resulted in a mean effective population size from all presented demes in northern Kyushu of

1,275,000.

Reconstruction of the phylogenetic tree for the partial mitochondrial DNA sequence

To examine the mitochondrial introgression in the tested samples, we generated and analyzed

the mitochondrial DNA (mtDNA) partial sequences from the GBS-read data.   We mapped

the reads to the complete mtDNA sequence of "Clade C" (Matsuda *et al.* 1997; Takehana *et*

al. 2003), which diverged genetically from S.JPN and N.JPN, to eliminate the mapping bias

that occurs when genetically distant sequences are used as a reference.   The genomic DNA

of "Clade C" was extracted in Katsumura *et al*. 2009.   The complete mtDNA sequence was

determined as follows.   Using PCR, five DNA fragments of mitochondrial DNA were

amplified: fragment 1—4,556 bp; fragment 2—4,527 bp; fragment 3-1—4,589 bp; fragment

3-2—4,504 bp; and fragment 4—4,546 bp (supplementary fig. 3).   Supplementary table 1

describes the primers, which were designed based on the inbred strain Hd-rR complete

mtDNA sequence (accession number: AP008938).   Approximately 20 ng of the genomic

DNA was used as a template for the PCR assay in a 50 μl solution containing dNTP at 0.2

mM, 0.2 μM of each of primer, 0.75 U of EX Taq polymerase HS (TaKaRa Shuzo Co.), and

the reaction buffer attached to the polymerase.   The reactions were conducted in a TaKaRa

PCR Thermal Cycler Dice (TaKaRa Shuzo Co.) using the following protocol: an initial

denaturing step at 95°C for 2 min, 40 cycles of denaturation at 95°C for 30 sec, annealing at

60°C for 30 sec, extension at 72°C for 300 sec, and a final extension step at 72°C for 5 min.

The PCR products were diluted 20-fold and used as templates in the sequencing reaction

(following the commercial protocol) with thirty-three primers (supplementary table 1,

supplementary fig. 3) and then analyzed in an ABI 3500xL Genetic Analyzer (Life

Technologies).   The complete mtDNA sequence of "Clade C" was reconstructed using

SeqMan Pro 10.1.2.20 (DNASTAR) and deposited into the international DNA database

DDBJ/EMBL/GenBank (accession number: LC335803).

The "Clade C" complete mtDNA sequence was used to align the reads using *BWA*

backtrack 0.7.12-r1039 (Li and Durbin 2009) using the "-n 5" option.   After the mapping

process, to remove the multi-mapped reads, we used *Samtools v1.2* (Li *et al.* 2009) using the

"-Sq 20" option.   Then, we used *Stacks* with the "-m 3" option for minimum stack depth

(Catchen *et al.* 2011; 2013) and then obtained the 322 bp nucleotide sequences.   The loci

where the sequence was missing were filled in with "N".   In addition, the nucleotide position

showing the multi-allelic state was replaced by "N".   Phylogenetic trees were constructed

using the neighbor-joining (NJ) method (Saitou and Nei 1987) with the program MEGA5

(Tamura *et al.* 2011).   The evolutionary distances were calculated using the Jukes-Cantor

method (Jukes and Cantor 1969).   The analysis also involved 13 nucleotide sequences from

the DNA database (See supplementary fig. 4).   All ambiguous positions were removed for

each sequence pair.   The reliability of the tree was evaluated using 1000 bootstrap replicates

(Felsenstein 1985).

Results

Autosomal genetic diversity of five groups of *Oryzias latipes*

To assess the genetic diversity based on autosomal SNPs within known mitochondrial and

hybridization groups—the northern Japanese (N.JPN), southern Japanese (S.JPN), eastern

Korean (E.KOR) western Korean/Chinese (W.KOR) and Tajima-Tango groups—their

population-genetic summary statistics were calculated using the "PopStat" dataset that

included the alignment regions across all groups and contained approximately 45 kb of

sequences and 2,453 SNPs (table 1).   All groups showed major allele frequencies and

heterozygosities of 0.940–0.983 and 0.023–0.054, respectively, indicating many rare alleles

and homozygous sites in our wild lab-stocks.   W.KOR showed the highest nucleotide

diversity (0.0053 ± 0.0003) in all lab-stocks originated from East Asia.   S.JPN showed the

highest nucleotide diversity (0.0036 ± 0.0001) and N.JPN the lowest in the Japanese

archipelago (table 1), which is consistent with the diversity based on mtDNA (Katsumura *et*

al. 2009).

Genetic clustering based on autosomal SNPs inferring medaka population structure

To reveal the medaka population structure based on the autosomal SNPs, we performed

genetic clustering analyses using the "Global" dataset.   First, we investigated the genetic

relationship by a principle component analysis (PCA) using 8,361 SNPs on 24 autosomes.

The PCA showed that the wild lab-stocks were divided into two clusters, N.JPN/Tajima-

Tango and S.JPN and others from E/W.KOR, which is similar to the grouping based on the

mtDNA data (Takehana *et al.* 2003; Katsumura *et al.* 2009)(fig. 2a, supplementary fig. 5).

E/W.KOR individuals were plotted dispersedly compared with other clusters, and Yongcheon

(E.KOR) was close to N.JPN/Tajima-Tango.   As PC3 was divided into E.KOR and W.KOR,

the autosomes in these mtDNA groups were differentiated into each other.

Although Tajima-Tango was considered to be a hybridization group between N.JPN

and S.JPN (Sakaizumi 1984), all individuals from Tajima-Tango were plotted with those from

N.JPN as a cluster (fig. 2a).   Next, for a fine-scale mapping of Japanese medaka, we

performed the PCA based on 7,126 SNPs, excluding E/W.KOR individuals (fig. 2b).   The

PCA showed that the Kyushu populations (Kudamatsu, Ogi, Izumi, Hiwaki, Kikai, Nago,

Gushikami, Hisayama, Umeshiro, Ashibe, Arita, Kusu and Nobeoka) in S.JPN were dispersed

along the PC2 axis (supplementary fig. 6), while that Tajima-Tango overlapped with N.JPN

again, even though we used up to PC5 (supplementary fig. 5).   This result indicated that

Tajima-Tango was not considerably differentiated from N.JPN on autosomes, suggesting that

S.JPN was differentiated within the groups and the genetic diversity was especially high

among the Kyushu populations (fig. 2b).

To examine the phylogenetic relationship among 83 populations, we constructed a

maximum likelihood (ML) tree based on 4,638 short fragment sequences (comparative

nucleotide sequence length: 217,986 bp) using *IQ-TREE*.   The maximum likelihood tree

showed relatively high bootstrap values on each branch and almost the same topology as

previous trees based on mtDNA (Takehana *et al.* 2003; Katsumura *et al.* 2009), except for

Yongcheon (fig. 3 and supplementary fig. 7; see also Discussion).   Medakas from the Korean

peninsula were genetically close to each other in both groups, whereas those from Shanghai,

which have been classified into W.KOR based on mtDNA studies, diverged from the other

individuals from W.KOR.   The medakas in the Japanese archipelago were divided into two

major clusters similar to mtDNA.   One was the N.JPN/Tajima-Tango, and the other was

S.JPN.   The N.JPN/Tajima-Tango cluster was further divided into two submajor clusters

with a 100% bootstrapping value in the ML tree based on concatenated sequences obtained

from GBS, although N.JPN and Tajima-Tango overlapped in PCA, which reduced SNP

information, i.e., N.JPN and Tajima-Tango diverged on the whole autosomal sequenced

regions.   S.JPN was also further divided into two sub-major clusters: Kyushu-only and

Kyushu & others, which included 14 Kyushu populations, represented in fig. 3 by an open

and a closed red circle, respectively.   While the Kyushu-only cluster diverged into the

northern and the southern Kyushu, the Kyushu & others cluster diverged to the Pacific side of

eastern and the northern Honshu (main-island Japan), which included several clusters

supported by high bootstrapping value (SH-aLRT $\geq$ 80% and UFboot $\geq$ 95%, supplementary

fig7).   Finally, we characterized the S.JPN ancestry in the context of East Asian genetic

diversity of medaka by performing an *ADMIXTURE* analysis, which is a model-based

unsupervised genetic clustering method.   With the optimal number of ancestral components

(K = 4), S.JPN medakas were assigned to two distinct ancestries (fig. 3 and supplementary

fig. 2).   In suboptimal runs with more ancestral components (K = 5, 6), only S.JPN in

Honshu (main-island Japan) was assigned to the other ancestral components found in the

northern Kyushu populations.   This analysis indicated that genetic diversities in the northern

Kyushu populations were the highest among S.JPN.   Thus, the genetic clustering analyses

based on genome-wide SNP data strongly suggested that S.JPN spread from northern Kyushu

to Honshu.

Boundary population genomes similar to that of the northern Japanese group

To explore hybridization signatures on autosomes, we examined allele-sharing between

boundary populations and surrounding populations using the "HZ-1" dataset (see Materials

and Methods).   We classified the fixed alleles between the groups into three states: shared by

N.JPN and Tajima-Tango, shared by S.JPN and Tajima-Tango, and shared by N.JPN and

S.JPN.   Then, we summarized each state's total numbers, as shown in table in fig. 4.   We

found 1,380 out of 4,661 SNPs that were common alleles between two of the three groups

(fig. 4).   Regarding those alleles, the majority (81.4%) were common alleles between

Tajima-Tango and N.JPN, which was a much higher frequency than that between Tajima-

Tango and S.JPN (11.2%).   The rest of them were common alleles between N.JPN and

S.JPN (7.5%), i.e., specific alleles in Tajima-Tango.   These proportions are near those of a

previous study based on 96 genomic regions (4 loci for each chromosome) (Takehana *et al.*

2016).   Although the regions with neighboring common alleles between Tajima-Tango and

S.JPN were observed for certain chromosomes, alleles on the Tajima-Tango genome were

mostly shared with N.JPN, suggesting that Tajima-Tango is a subgroup of N.JPN with a short

divergence time.

To investigate the mitochondrial introgression in boundary populations, we

reconstructed the phylogenetic tree based on partial mtDNA sequences generated by mapping

short reads from the GBS of the complete mitochondrial genome (supplementary fig. 4).

The mtDNA sequences from Toyooka and Kinosaki in Tajima-Tango were clustered and

closely related to those from Kaga, which was a root population in N.JPN.   Considering that

Tajima-Tango was a subgroup of N.JPN, which we inferred from the ML tree based on

autosomal sequences, these phylogenetic positions on the mtDNA tree would have also

reflected an evolutionary history in which N.JPN diverged from the Tajima-Tango group.

We confirmed that Amino, Kumihama and Hamasaka in Tajima-Tango had the mitochondrial

genomes of phylogenetically distant populations classified into S.JPN (supplementary fig. 4).

Additionally, the mitochondrial genome of Ayabe in S.JPN was classified into N.JPN but

diverged from Tajima-Tango.   These results suggest that mitochondrial genome

introgressions occurred reciprocally, meaning that they occurred not only from S.JPN to

Tajima-Tango but also from N.JPN to S.JPN.

Inferring medaka demographic parameters

To infer the demographic parameters, effective population size (N_e), time (T), and proportion

of the admix (r), we analyzed the "HZ-2" dataset based on the coalescent theory using an

approximate Bayesian computation (ABC) framework.   We performed the model selection

to identify the best explanation scenario for the observed data from the four scenarios; (I)

Tajima-Tango originated in N.JPN, (II) Admixture of N.JPN and Honshu, (III) N.JPN

originated in Tajima-Tango, and (IV) Admixture of Tajima-Tango and Honshu (fig. 5a).

DIYABC has two different approaches (directional and logistic) for model selection.   Based

on each criterion, scenario III (Honshu diverged from Kyushu and then N.JPN diverged from

the Tajima-Tango group without admixture with Honshu) and scenario IV (Honshu diverged

from Kyushu and then Tajima-Tango occurred by an admixture with Honshu at r rate) were

supported by the directional (posterior probability: 0.4720, 95% CI: 0.0344–0.9096) and

logistic (posterior probability: 0.4477, 95% CI: 0.4313–0.4641) approaches, respectively

(supplementary fig. 8).   Our data could not distinguish between scenarios III and IV (see the

Discussion section).

421   The posterior parameter estimates of scenarios III and IV shown in table 2 and figs.

5c and 5d were not scaled because various measures of accuracy (RRMISE, RMeanAD, and

RRMSE in the *DIYABC* output) indicated that non-scaled parameters fit the observed data

better than scaled parameters   Additionally, although we inferred the time to the most

common ancestor at three divergence events (fig. 5), these times might be overestimated

because the dataset for this estimation consisted of non-deme samples, i.e., polymorphisms

within populations were underestimated, and the random-mating assumption could not be

satisfied.   Therefore, we estimated the timing of the "Out of Northern Kyushu" event using

the "Local" dataset composed of the two deme samples (S.Saga and N.Saga), which were split

at first on the S.JPN lineage (fig. 3).   Based on a simple hypothesis, constant population size

and no migration, we obtained the estimated time (510,000 years ago, 95% CI: 337,875–

679,575) for the ancestral divergence of the two deme samples (table 2 and fig. 5e), which

was calculated from a scaled parameter by the mean effective population size because various

accuracy measures indicated that scaled parameters fit the observed data better than non-

scaled parameters.   Thus, our population-genetic estimate using ABC suggests that medakas

diverged and dispersed throughout Pacific-side Japan approximately 510 kyr ago.

Discussion

Redefined subgroups in Japanese archipelago based on the population structure

inferred from genome-wide single-nucleotide polymorphisms

Oryzias latipes can be divided into five groups by mtDNA sequences and allozymic

electrophoresis patterns (Sakaizumi 1984; Takehana *et al.* 2003). In this study, based on

autosomal SNPs, the genetic clustering analysis showed that "K = 4" was the most supportive

because it presented the lowest fivefold cross-validation error, indicating that N.JPN and

S.JPN were divided into three ancestral clusters. When the K values increased, only S.JPN

divided into more subgroups, which suggests that the S.JPN group was composed of more

divergent groups than the other groups. Considering together with the results of the ML tree

analysis, it is possible to redefine subgroups composed of each major group for our wild lab-

stocks originated from the Japanese archipelago as follows. First, Tajima-Tango, which had

been considered a hybridization group, should be included under the N.JPN group because it

shows almost the same ancestral component as N.JPN. Then, the group called N.JPN should

be assigned to a subgroup, for which we propose the name "Sea of Japan side of Northeastern

Honshu (SJ.NEH)." These two subgroups, Tajima-Tango and SJ.NEH, compose the N.JPN

group (supplementary fig. 7). Second, S.JPN can be divided into several subgroups, San-in,

San-yo/Shikoku/Kinki and the Pacific Ocean side of Northeastern Honshu (PO.NEH),

because the Kyushu & others cluster was subdivided into three sub-clusters composed of

geographically neighboring populations. Adding the Kyushu subgroup, S.JPN is composed

of four subgroups. Thus, medaka in Japanese archipelago could also be composed of the six

distinct subgroups based on autosomal genetic diversity (supplementary fig. 7).

S.JPN can be divided into finer subgroups based on the mitochondrial genome

(Takehana *et al.* 2003), likely because the effective population size of mtDNA is one quarter

that of a nuclear gene. This causes the intermingled branch pattern of the mtDNA tree,

which is not associated with geographic distance.   Although this pattern has been observed

our wild lab-stocks in a previous study based on mtDNA (Katsumura *et al.* 2009), it has been

interpreted as artificial migrations accompanied by recent human activities (Takehana *et al.*

2003).   If the branching pattern was formed by human activities, various ancestral

components should appear independent of geographic relatedness in the result of the

ADMIXTURE analysis based on autosomal SNPs.   However, no signal of recent migration

was found in the *ADMIXTURE* result.   Rather, the autosomal SNP data support another

hypothetical scenario in which the mtDNA tree topology reflects ancestral polymorphisms

only and their local fixation is caused by a small effective population size (Katsumura *et al.*

2009).   This result indicates that arguing the geographical origins of medaka based only on

mtDNA may lead to false conclusions.

From our data, it may be difficult to accurately evaluate the genetic diversities in

E/W.KOR groups because the number of populations examined in each group was small.

Therefore, though care must be taken in the interpretation, W.KOR showed the highest

nucleotide diversity and E.KOR the second highest among the four major mitochondrial

groups (table 1).   The W.KOR group included the Chinese medaka in Shanghai, which could

have elevated its value, while the latter group did not include any geographically distant

populations.   The clustering analysis showed that Shanghai from W.KOR and Yongcheon

from E.KOR had an ancestry component from N.JPN (fig.3).   This suggests two

possibilities: one is that the N.JPN ancestor was derived from the E/W.KOR ancestor, and the

other is that contamination occurred through the maintained wild lab stocks.   To investigate

these possibilities and the origin of *O. latipes*, a population-based genome wide analysis must

be conducted to increase the population numbers of E/W.KOR and include the sister species

O. curvinotus and *O. luzonensis*.

Reconstruction of the medaka population history in the Japanese archipelago

Our genome-wide analysis shows that medakas in N.JPN and S.JPN are deeply divergent and

dispersed over the Japanese archipelago from different locations at different times.   In

particular, our ABC analysis indicates that SJ.NEH originated in and diverged from Tajima-

Tango.   This inference about the history of N.JPN after divergence from S.JPN is not

consistent with previous inferences from the allozyme, mtDNA and limited autosomal SNP

analyses, which have suggested that Tajima-Tango is a hybridization group between PO.NEH

and S.JPN, or Tajima-Tango is a sub-group derived from PO.NEH.   Our GBS data show that

(i) the nucleotide diversity in Tajima-Tango is higher than that in SJ.NEH (table 1), (ii) S.JPN

is genetically more closely related to Tajima-Tango than to SJ.NEH based on the allele-

sharing rates (fig. 4), and (iii) the Tajima-Tango branch is the root in the N.JPN clade of the

phylogenetic tree based on partial mtDNA sequences (supplementary fig. 4), which is also

shown by the whole mitochondrial genome analysis (Hirayama *et al.* 2010).   These data

suggest that Tajima-Tango forms an outgroup to all present-day SJ.NEH and it spread along

the Sea of Japan side.   Furthermore, the ABC framework's estimation supports our scenario,

although the analysis does not statistically distinguish between scenario III and scenario IV.

Even if the admixture occurred, the inferred ratio of the admixture is too low (fig. 5d).

Our findings strongly support the "Out of northern Kyushu" model of S.JPN

proposed in Katsumura et al. (2012) and have revealed the dispersal route of SJ.NEH/Tajima-

Tango in N.JPN.   The genetic clustering analyses and phylogenetic tree based on the GBS

data elucidate medaka history better than previous mitochondrial DNA analyses.   In

particular, the *ADMIXTURE* analysis shows that the two ancestry components (yellow and

blue in fig. 3) were observed in northern Kyushu, suggesting that S.JPN dispersed in three

different directions after dispersing out of Northern Kyushu.   Geographically, Shikoku, San-

yo, and Kinki are separated by sea, but medakas in the three local lands share the same

ancestral component (blue in fig.6; see also supplementary fig. 1 for geographic information).

Medaka is a freshwater fish, but survival and reproduction are possible even in seawater

(Inoue and Takei 2002).   Because most of the rivers in the Japanese archipelago are steep

and short, the rivers tend to flood after heavy rain.   Although medakas are highly likely to

drift to the sea on each occasion, medakas can have survived even in the sea and may have

returned to the river because of their saltwater tolerance.   Thus, the possibility of moving

from river to river through the sea cannot be ignored.   The results of this study suggest a

history in which medakas migrated throughout the Japanese archipelago through the sea.

The different branching patterns of autosomal and mtDNA trees suggest that the

mtDNA introgression occurred not only from S.JPN to Tajima-Tango but also from N.JPN to

S.JPN.   The Tajima-Tango region is surrounded by mountains (supplementary fig. 1);

however, it contains the lowest watershed (sea level 95.45 m) in the Japanese archipelago.

The medaka has possibly moved in both directions across the watershed.   From the above,

the most plausible scenario is as follows (fig. 6).   Ancestral S.JPN and N.JPN diverged first

and independently reached their current habitats in the Japanese archipelago.   After S.JPN,

which is an ancestor of San-in, San-yo/Shikoku/Kinki and PO.NEH, dispersed from northern

Kyushu approximately 510 kyr ago, SJ.NEH diverged from Tajima-Tango in N.JPN.   Then,

their descendant populations spread rapidly to northern Honshu on the Sea of Japan side.

Meanwhile, S.JPN dispersed in as many as three different directions and then spread rapidly

northeastward from the western part of Fossa Magna.   In the process of dispersing across the

main island of Japan, certain S.JPN populations infiltrated the Tajima-Tango region from the

west and the south, and the resulting mtDNA introgression occurred independently.

Conclusion

Our genome-wide SNP analysis reconstructed the detailed population structure and reliable

history of medaka that evolved in the Japanese archipelago.   Since the distribution of the

subgroups was highly consistent with the geographical features, several adaptive traits could

have evolved in each subgroup.   Furthermore, the boundary populations were not caused by

a hybridization event but instead were the origin of the populations dispersed to a

northeastern part of the Japanese archipelago on the Sea of Japan side.   A better

understanding of the population structure and history of medaka will support association

studies for phenotypes and genotypes related to environmental adaptation.

References

Alexander, D. H., J. Novembre, and K. Lange, 2009 Fast model-based estimation of ancestry
in unrelated individuals. Genome Research 19: 1655–1664.

Andrews, K. R., J. M. Good, M. R. Miller, G. Luikart, and P. A. Hohenlohe, 2016 Harnessing
the power of RADseq for ecological and evolutionary genomics. Nature Review Genetics
17: 81–92.

Ansai, S., and M. Kinoshita, 2014 Targeted mutagenesis using CRISPR/Cas system in
medaka. Biology Open 3: 362–371.

Beaumont, M. A., W. Zhang, and D. J. Balding, 2002 Approximate Bayesian computation in
population genetics. Genetics 162: 2025–2035.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko, 2013 Stacks: an
analysis tool set for population genomics. Mol Ecol 22: 3124–3140.

Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait, 2011 Stacks:
building and genotyping Loci de novo from short-read sequences. G3 (Bethesda) 1: 171–
182.

Chang, C. C., C. C. Chow, L. C. Tellier, S. Vattikuti, S. M. Purcell *et al.*, 2015 Second-
generation PLINK: rising to the challenge of larger and richer datasets. Gigascience 4: 7.

Chernomor, O., A. von Haeseler, and B. Q. Minh, 2016 Terrace Aware Data Structure for
Phylogenomic Inference from Supermatrices. Systematic Biology 65: 997–1008.

Cornuet, J.-M., P. Pudlo, J. Veyssier, A. Dehne-Garcia, M. Gautier *et al.*, 2014 DIYABC
v2.0: a software to make approximate Bayesian computation inferences about population
history using single nucleotide polymorphism, DNA sequence and microsatellite data.
Bioinformatics 30: 1187–1189.

Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto *et al.*, 2011 A robust, simple
genotyping-by-sequencing (GBS) approach for high diversity species. PLoS ONE 6:
e19379.

Felsenstein, J., 1985 CONFIDENCE LIMITS ON PHYLOGENIES: AN APPROACH
USING THE BOOTSTRAP. Evolution 39: 783–791.

Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk *et al.*, 2010 New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. Systematic Biology 59: 307–321.

Hirayama, M., T. Mukai, M. Miya, Y. Murata, Y. SEKIYA *et al.*, 2010 Intraspecific variation
in the mitochondrial genome among local populations of Medaka Oryzias latipes. Gene
457: 13–24.

Igarashi, K., J. Kobayashi, T. Katsumura, Y. Urushihara, K. Hida *et al.*, 2017 An Approach to
Elucidate NBS1 Function in DNA Repair Using Frequent Nonsynonymous
Polymorphism in Wild Medaka (Oryzias latipes) Populations (S. D. Fugmann, Ed.). PLoS
ONE 12: e0170006–19.

Inoue, K., and Y. Takei, 2002 Diverse adaptability in oryzias species to high environmental
salinity. Zool. Sci. 19: 727–734.

Jeong, C., S. Nakagome, and A. Di Rienzo, 2016 Deep History of East Asian Populations
Revealed Through Genetic Analysis of the Ainu. Genetics 202: 261–272.

Jukes, T. H., and C. R. Cantor, 1969 Evolution of protein molecules. Mammalian protein
metabolism 3: 132.

Kasahara, M., K. Naruse, S. Sasaki, Y. Nakatani, W. Qu *et al.*, 2007 The medaka draft
genome and insights into vertebrate genome evolution. Nature 447: 714–719.

Katsumura, T., S. Oda, S. Mano, N. Suguro, K. Watanabe *et al.*, 2009 Genetic differentiation
among local populations of medaka fish (Oryzias latipes) evaluated through grid- and
deme-based sampling. Gene 443: 170–177.

Katsumura, T., S. Oda, S. Nakagome, T. Hanihara, H. Kataoka *et al.*, 2014 Natural allelic
variations of xenobiotic-metabolizing enzymes affect sexual dimorphism in Oryzias
latipes. Proceedings of the Royal Society B: Biological Sciences 281:.

Katsumura, T., S. Oda, K. Tsukamoto, T. Yamashita, M. Aso *et al.*, 2012 A population
genetic study on the relationship between medaka fish and the spread of wet-rice
cultivation across the Japanese archipelago. Anthropol. Sci.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25: 1754–1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

Matsuda, M., H. Yonekawa, S. Hamaguchi, and M. Sakaizumi, 1997 Geographic variation
and diversity in the mitochondrial DNA of the medaka, Oryzias latipes, as determined by
restriction endonuclease analysis. Zool. Sci. 14: 517–526.

Minh, B. Q., M. A. T. Nguyen, and A. von Haeseler, 2013 Ultrafast approximation for
phylogenetic bootstrap. Molecular Biology and Evolution 30: 1188–1195.

Narum, S. R., C. A. Buerkle, J. W. Davey, M. R. Miller, and P. A. Hohenlohe, 2013
Genotyping-by-sequencing in ecological and conservation genomics. Mol Ecol 22: 2841–
2847.

Nguyen, L.-T., H. A. Schmidt, A. von Haeseler, and B. Q. Minh, 2015 IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular
Biology and Evolution 32: 268–274.

Oota, H., and H. Mitani, 2011 Human Population Genetics Meets Medaka, pp. 339–350 in
Medaka: A Model for Organogenesis, Human Disease, and Evolution, edited by K.
Naruse, M. Tanaka, and H. Takeda. Medaka: A Model for Organogenesis, Human
Disease, and Evolution, Springer Japan, Tokyo.

Osada, N., 2015 Genetic diversity in humans and non-human primates and its evolutionary
consequences. Genes Genet. Syst. 90: 133–145.

Poland, J. A., P. J. Brown, M. E. Sorrells, and J.-L. Jannink, 2012 Development of High-
Density Genetic Maps for Barley and Wheat Using a Novel Two-Enzyme Genotyping-
by-Sequencing Approach (T. Yin, Ed.). PLoS ONE 7: e32253–8.

Saitou, N., and M. Nei, 1987 The neighbor-joining method: a new method for reconstructing
phylogenetic trees. Molecular Biology and Evolution 4: 406–425.

Sakaizumi, M., 1986 Genetic divergence in wild populations of Medaka, Oryzias latipes
(Pisces: Oryziatidae) from Japan and China. Genetica.

Sakaizumi, M., 1984 Rigid Isolation between the Northern Population and the Southern
Population of the Medaka, Oryzias latipes (Genetics). Zool. Sci. 1: 795–800.

Sakaizumi, M., N. EGAMI, and K. MORIWAKI, 1980 Allozymic Variation in Wild
Populations of the Fish, Oryzias latipes. Proc. Jpn. Acad., Ser. B 56: 448–451.

Sakaizumi, M., K. Moriwaki, and N. Egami, 1983 Allozymic variation and regional
differentiation in wild populations of the fish Oryzias latipes. Copeia 1983: 311.

Shima, A., A. Shimada, M. Sakaizumi, and N. Egami, 1985 First listing of wild stocks of the
medaka Oryzias latipes currently kept by the Zoological Institute, Faculty of Science,
University of Tokyo. Journal of the Faculty of Science, Imperial University of Tokyo
Sect IV, Zoology 16: 27–35.

Shimmura, T., T. Nakayama, A. Shinomiya, S. Fukamachi, M. Yasugi et al., 2018 Dynamic
plasticity in phototransduction regulates seasonal changes in color perception. Nature
Communications 1–7.

Spivakov, M., T. O. Auer, R. Peravali, I. Dunham, D. Dolle et al., 2014 Genomic and
phenotypic characterization of a wild medaka population: towards the establishment of an
isogenic population genetic resource in fish. 4: 433–445.

Tajima, F., 1983 Evolutionary relationship of DNA sequences in finite populations. Genetics
105: 437–460.

Takehana, Y., N. Nagai, M. Matsuda, K. Tsuchiya, and M. Sakaizumi, 2003 Geographic
variation and diversity of the cytochrome b gene in Japanese wild populations of medaka,
Oryzias latipes. Zool. Sci. 20: 1279–1291.

Takehana, Y., M. Sakai, T. Narita, T. Sato, K. Naruse et al., 2016 Origin of Boundary
Populations in Medaka (Oryzias latipes Species Complex). Zool. Sci. 33: 125–131.

Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei et al., 2011 MEGA5: Molecular
Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and
Maximum Parsimony Methods. Molecular Biology and Evolution 28: 2731–2739.

Watanabe-Asaka, T., Y. SEKIYA, H. Wada, T. Yasuda, I. Okubo et al., 2014 Regular
heartbeat rhythm at the heartbeat initiation stage is essential for normal cardiogenesis at
low temperature. BMC Dev. Biol. 14:.

Wickham, H., 2011 ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 3:
180–185.

659 Zheng, X., D. Levine, J. Shen, S. M. Gogarten, C. Laurie *et al.*, 2012 A high-performance
660   computing toolset for relatedness and principal component analysis of SNP data.
661   Bioinformatics 28: 3326–3328.

Acknowledgments

We thank Mrs. Shizuko Chiba, Mrs. Sumiko Tomizuka, Dr. Atsuko Shimada and Prof.

Emeritus Akihiro Shima (the University of Tokyo) for maintaining the medaka stocks from

wild populations.　This study was supported in part by grants-in-aid for Young Scientists (B),

no. 16K21352 to T.K., and by a grant-in-aid for Scientific Research (A), nos. 26251050,

25251046 and 17H01453 to H.O.　T.K. was also supported by a grant-in-aid from JSPS

Research Fellowships (16J07227).

Figure legends

FIG. 1: Map of the original locations of the wild lab stocks and wild-captured medakas.

In the upper central map, the enlarged red frame shows the boundary region between S.JPN

and N.JPN.　Each color represents the mtDNA and allozyme-based groups shown in the left

upper inset box.　The numbers on the map are consistent with the population IDs, with the

names on the right bottom inset box.

FIG. 2: Results of principle component analysis (PCA) using SNPs in East Asia (a) and

the Japanese archipelago (b).　Each plot shows PC1 versus PC2.　The population names

of each point are described in fig. S6.

FIG. 3: Phylogenetic tree using the maximum likelihood method and an ancestry

barplot with ADMIXTURE analysis.　Red closed and open circles represent the northern

and southern Kyushu populations, respectively.　"Taj-Tan" in the tree is the abbreviation for

Tajima-Tango.

FIG. 4: Shared allele distribution among boundary populations.　The gray rectangle

represents the total length of each medaka chromosome.　The table in the figure shows the

number of alleles shared between the groups observed on each chromosome.　Mark

represents the state whose group shares that allele.　*1 and *2 are the total number from

chromosomes 1 to 12 and from chromosomes 13 to 24, respectively.

FIG. 5: Scenarios for estimation of the demographic parameters.　Possible scenarios for

the population history of the Tajima-Tango group (a) and the "Out of northern Kyushu (NK)"

event (b) are shown.　The TMRCAs estimated using ABC are in scenario3 (c), scenario4 (d)

and Out of NK (e).

FIG. 6: Map representing the ancestry proportions from ADMIXTURE analysis at K =

6.   Solid and dashed lines represent the spreading patterns of S.JPN and N.JPN inferred by

GBS data, respectively.

Figure 1

Tajima-Tango region

Legend:
- N.JPN
- S.JPN
- Tajima-Tango
- E.KOR
- W.KOR

Korean peninsula

Eurasia continent

Japanese archipelago

1_Kamikita	22_Kumihama	43_Kosakai	64_Tanabe
2_Hirosaki	23_Toyooka	44_Nagashima	65_Oku
3_Odate	24_Kinosaki	45_Shingu	66_Okayama
4_Yokote	25_Hamasaka	46_Kochi	67_Kayo
5_Tsuruoka	26_Kesennuma	47_Misho	68_Tokushima
6_Yamagata	27_Ichinoseki	48_Nobeoka	69_Takamatsu
7_Nigata	28_Iwaki	49_Kusu	70_Tessei
8_Ryotsu	29_Uridura	50_Okaya	71_Miyoshi
9_Kosugi	30_Mibu	51_Ashibe	72_Matsuyama
10_Kaga	31_Sakura	52_Fukue	73_Kudamatsu
11_Sabae	32_Kobayashi_Bokujo	53_Hisayama	74_Yamaguchi
12_Obama	33_Tokyo	54_Arita	75_Sanyo
13_Maiduru	34_Toyota	55_Kadusa	76_Kasumi
14_Miyadu	35_Omati	56_Izumi	77_Iwami
15_Shanghai	36_Suwa	57_Hiwaki	78_Tottori
16_Maegok	37_Kobuchisawa	58_Kikai	79_Matsue
17_Bugang	38_Ida	59_Nago	80_Tsuma
18_Yongchon	39_Odawara	60_Gushikami	81_Hagi
19_Sacheon	40_Mishima	61_Ayabe	82_Ogi (SO-1)
20_Sinpyeong	41_Shizuoka	62_Kobe	83_Umeshiro (US-2)
21_Amino	42_Iwata	63_Kawai	

Figure 2

Figure 3

Figure 4

	Mark	Chr1	Chr2	Chr3	Chr4	Chr5	Chr6	Chr7	Chr8	Chr9	Chr10	Chr11	Chr12	Total*1
NJPN–Tajima-Tango	■	62	11	52	54	70	53	65	32	75	48	44	61	627
SJPN–Tajima-Tango	●	7	1	6	7	6	6	8	5	1	8	7	4	66
Tajima-Tango only	▲	6	1	3	6	8	3	7	5	8	7	2	4	60

	Mark	Chr13	Chr14	Chr15	Chr16	Chr17	Chr18	Chr19	Chr20	Chr21	Chr22	Chr23	Chr24	Total*2
NJPN–Tajima-Tango	■	39	47	60	50	51	28	21	44	42	52	26	36	496
SJPN–Tajima-Tango	●	10	9	6	9	13	3	5	5	9	8	5	6	88
Tajima-Tango only	▲	1	3	6	4	5	2	3	3	7	7	1	1	43

Figure 5

Figure 6

Origin of N.JPN

Origin of S.JPN

Latitude

Longitude

Table 1 Summary genetic statistics for five populations using "PopStat" dataset. These statistics include the mean number of individuals genotyped at each locus (*1), the number of variable sites in five populations (Variants), the number of polymorphic sites in each population (Polymorphic) and the number of variable sites unique to each population (Private). The number in parenthesis is Standard Error.

Group	Number of Individual[*1]	Total length of GBS loci (bp)	Variant	Polymorphic	Private	Major Allele Frequency	Observed Heterozygosity	Nucleotide Diversity
N.JPN (SJ.NEH)	11.5 (+/-0.03)	45968	2453	256	136	0.983 (+/-0.0014)	0.023 (+/-0.0022)	0.0014 (+/-0.0001)
Tajima -Tango	4.4 (+/-0.01)	44011	2453	199	71	0.980 (+/-0.0016)	0.024 (+/-0.0022)	0.0017 (+/-0.0001)
S.JPN	41.6 (+/-0.08)	45968	2453	1448	1231	0.958 (+/-0.0017)	0.029 (+/-0.0016)	0.0036 (+/-0.0001)
W.KOR	2.4 (+/-0.01)	44990	2453	467	342	0.940 (+/-0.0027)	0.054 (+/-0.0032)	0.0053 (+/-0.0003)
E.KOR	2.4 (+/-0.01)	44011	2453	346	203	0.953 (+/-0.0025)	0.041 (+/-0.0030)	0.0042 (+/-0.0002)

Table 2 Demographic parameters estimated by DIYABC under three scenarios. Performance of each estimation was evaluated by RRMISE (the square Root of the Relative Mean Integrated Square Error), RMeanAD (the Relative Mean Absolute Deviation) and RRMSE (the square Root of teh Relative Mean Square Error), which are output from DIYABC option "Compute bias and mean square error." The numbers with or without parentheses in the culunms "Performances for estimating posterior distributions of parameters" are those of the statistics computed from the prior or posterior distribution of parameters, respectively.

Scenario	Parameters	Prior distribution					Posterior parameter estimates			Performances for estimating posterior distributions of parameters		
		Type	min.	max.	mean	S.D.	Mean	95% credible interval		RRMISE	RMeanAD	RRMSE
II	$N_{\text{N.JPN}}$	Normal	10,000	3,000,000	450,000	500,000	825,000	492,000	1,240,000	0.415 (-0.586)	0.302 (-0.485)	0.302 (-0.288)
	$N_{\text{Tajima-Tango}}$	Normal	10,000	3,000,000	600,000	500,000	613,000	369,000	901,000	0.35 (-0.808)	0.256 (-0.61)	0.251 (-0.358)
	N_{Kyushu}	Normal	10,000	3,000,000	1,225,000	500,000	1,210,000	711,000	1,830,000	0.364 (-0.524)	0.264 (-0.396)	0.267 (-0.282)
	N_{honshu}	Normal	10,000	3,000,000	1,050,000	500,000	1,490,000	974,000	2,020,000	0.291 (-0.442)	0.219 (-0.368)	0.213 (-0.289)
	N_{Ancestor}	Uniform	10,000	30,000,000	-	-	432,000	59,700	922,000	5.899 (-133.368)	1.846 (-59.034)	5.119 (-115.475)
	$T_{\text{N.JPN–Hybrid}}$	Uniform	10	4,000,000	-	-	566,000	311,000	908,000	0.409 (-2.417)	0.300 (-1.725)	0.266 (-1.589)
	$T_{\text{kyushu–Honshu}}$	Uniform	10	4,000,000	-	-	1,050,000	442,000	2,290,000	0.575 (-2.398)	0.374 (-1.875)	0.284 (-2.114)
	$T_{\text{N.JPN–S.JPN}}$	Normal	1,000,000	30,000,000	4,000,000	5,000,000	4,560,000	2,820,000	6,810,000	0.329 (-1.079)	0.245 (-0.776)	0.229 (-0.743)
III	$N_{\text{N.JPN}}$	Normal	10,000	3,000,000	450,000	500,000	812,000	390,000	1,330,000	0.447 (-0.635)	0.332 (-0.506)	0.303 (-0.326)
	$N_{\text{Tajima-Tango}}$	Normal	10,000	3,000,000	600,000	500,000	480,000	266,000	739,000	0.375 (-1.208)	0.284 (-0.88)	0.261 (-0.736)
	N_{Kyushu}	Normal	10,000	3,000,000	1,225,000	500,000	1,290,000	758,000	1,910,000	0.356 (-0.483)	0.265 (-0.373)	0.265 (-0.252)
	N_{honshu}	Normal	10,000	3,000,000	1,050,000	500,000	1,450,000	913,000	2,020,000	0.316 (-0.446)	0.235 (-0.369)	0.235 (-0.279)
	N_{Ancestor}	Uniform	10,000	30,000,000	-	-	390,000	57,200	849,000	6.947 (-131.19)	2.285 (-62.259)	5.997 (-113.55)
	T_{ADMIX}	Uniform	10	4,000,000	-	-	512,000	242,000	842,000	0.397 (-4.374)	0.308 (-2.163)	0.249 (-3.309)
	$T_{\text{kyushu–Honshu}}$	Uniform	10	4,000,000	-	-	1,100,000	446,000	2,420,000	0.576 (-2.256)	0.381 (-1.76)	0.296 (-1.98)
	$T_{\text{N.JPN–S.JPN}}$	Normal	1,000,000	30,000,000	4,000,000	5,000,000	5,160,000	3,080,000	7,780,000	0.345 (-0.916)	0.262 (-0.65)	0.239 (-0.588)
	r	Uniform	0.001	0.999	-	-	0.944	0.897	0.987	0.027 (-0.559)	0.02 (-0.472)	0.014 (-0.469)
Out of NK	$N_{\text{S.Saga}}$	Normal	10,000	3,000,000	1,300,000	500,000	1,326,000	960,075	1,670,250	0.178 (-0.31)	0.138 (-0.241)	0.120 (-0.095)
	$N_{\text{N.Saga}}$	Normal	10,000	3,000,000	1,250,000	500,000	1,224,000	892,500	1,593,750	0.181 (-0.322)	0.142 (-0.251)	0.119 (-0.096)
	$T_{\text{S.Saga–N.Saga}}$	Uniform	1,000	4,000,000	-	-	510,000	337,875	679,575	0.224 (-5.918)	0.175 (-4.488)	0.143 (-4.429)