Arabidopsis leaf hydraulic conductance is regulated by xylem-sap pH, controlled, in turn, by a P-type H+-ATPase of vascular bundle sheath cells

One-sentence summary:

Bundle-sheath cells can control the leaf hydraulic conductance by proton-pump-regulated xylem sap pH

Yael Grunwald*[1], Noa Wigoda*[1,3], Nir Sade[1,2], Adi Yaaran[1], Tanmayee Torne[1], Sanbon Chaka Gosa[1], Nava Moran[1] and Menachem Moshelion[1].

[1] The R.H. Smith Institute of Plant Sciences and Genetics in Agriculture, The R.H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, 76100 Israel

[2] School of Plant Sciences and Food Security, The George S. Wise Faculty of Life Sciences at Tel-Aviv University, Tel Aviv 6997801, Israel.

[3] *Present address*: The Department of Life Sciences Core Facilities, Weizmann Institute of Science, Rehovot 76100 Israel

* Both authors contributed equally to this work.

Abstract

●The leaf vascular bundle sheath cells (BSCs) that tightly envelop the leaf veins, are a selective and dynamic barrier to xylem-sap water and solutes radially entering the mesophyll cells. Under normal conditions, xylem-sap pH of <6 is presumably important for driving and regulating the transmembranal solute transport. Having discovered recently a differentially high expression of a BSCs proton pump, AHA2, we now test the hypothesis that it regulates this pH and leaf radial water fluxes.

●We monitored the xylem-sap pH in the veins of detached leaves of WT Arabidopsis, AHA mutants, and *aha2* mutants complemented with *AHA2* gene solely in BSCs. We tested an AHA inhibitor and stimulator, and different pH buffers. We monitored their impact on the xylem-sap pH and the whole leaf hydraulic conductance (K_{leaf}), and the effect of pH on the water osmotic permeability (P_f) of isolated BSCs protoplasts.

●Our results demonstrated that AHA2 is necessary for xylem-sap acidification, and in turn, for elevating K_{leaf}. Conversely, knocking out AHA2 alkalinized the xylem-sap. Also, elevating xylem sap pH to 7.5 reduced K_{leaf} and elevating external pH to 7.5 decreased the BSCs P_f.

●All these demonstrate a causative link between AHA2 activity in BSCs and leaf radial water conductance.

Key words

AHA, AHA2, ABA, *Arabidopsis thaliana* (Thale cress), Barrier, Bundle Sheath, H^+-ATPase, Leaf hydraulic conductivity (K_{leaf}), Proton pump, Water valve, Xylem-sap pH.

Introduction

The majority (95%) of nutrients and water which enter the plant through the roots move upward in a bulk flow ("transpiration stream") via xylem vessels all the way into the leaf veins (Taiz and Zeiger, 2014), then radially, through the bundle sheath cells (BSCs) – into the leaf photosynthetic tissue, the mesophyll. The parenchymatous BSCs, tightly enwraping the entire vasculature (Kinsman and Pyke, 1998), constitute a selective barrier between the vein and the mesophyll. For example, they impeded the transport of sodium (Shapira et al., 2009) and of boron (Shapira et al., 2013) in a banana leaf, and boron – also in Thelungiella (Lamdan et al., 2012) and Arabidopsis (Shatil-Cohen and Moshelion, 2012). Importantly, the BSCs act also as a barrier to the passage of water (Shatil-Cohen et al., 2011; Pantin et al., 2013). However, it is still not clear (Geilfus, 2017) whether and how the apoplastic (xylem) milieu modulates the "barrier behavior" of the bundle sheath, and in particular, whether this behavior depends on the xylem sap pH.

Some direct determinations of leaf xylem pH placed it at 6.3-6.7 in cotton leaves (Hartung et al., 1988), or at 5.3 in sunflower leaves (Jia and Davies, 2007) but most frequently, under normal growth conditions the leaf apoplast pH is around 5.5-6 (reviewed by Grignon and Sentenac, 1991; Geilfus, 2017). The leaf xylem sap pH can change in response to changes in external conditions experienced by the root. For example, the tomato leaf xylem sap alkalinized upon nitrate supply to roots (Jia and Davies, 2007). Barley leaf sub-stomatal apoplast also alkalinized upon roots exposure to salts (KCl, NaCl, $NaNO_3$ (Felle et al., 2005)). Drying soil in tomato (Jia and Davies, 2007) and in *Vicia faba* (Karuppanapandian et al., 2017) or cold stress at the root level in barley (Felle et al., 2005) also caused alkalinization (measured, respectively, in the tomato xylem sap, in the *V. faba* xylem sap and leaf apoplast, and in the barley sub-stomatal apoplast). Finally, alkalinization of the barley leaf sub-stomatal apoplast was brought about also by applying ABA (considered not only a drought-stress hormone, but also shown to mediate cold stress (Huang et al., 2017)) both to barley roots and to detached leaves (Felle et al., 2005).

In turn, changes in the leaf xylem sap pH regulate physiological processes. Feeding high-pH solutions to detached leaves reduced stomatal conductance and transpiration both in *Commelina communis* and in tomato (Wilkinson and Davies, 1997; Wilkinson et al., 1998; Jia and Davies, 2007). As pH defines the dissociation state of weak acids, among them the major phytohormones abscisic acid (ABA) and indol acetic acid (IAA, auxin), changes in pH would affect both their distribution between the apoplast and the cellular compartments of the plant, and hence, their biological activities. For example, ABA accumulated in the alkalinizing xylem sap of a gradually pressure-dehydrated detached cotton leaf, which could be prevented by pretreatment with an H^+-ATPase-activating fungal toxin, fusicoccin (Hartung et al., 1988). Leaf xylem alkalinization in *Vicia faba* as a result of gradually drying soil was followed by elevated ABA levels in the xylem (even if with a few days delay (Karuppanapandian et al., 2017).

Already widely-accepted is the crucial role of apoplastic protons in the proton-motive force governing transmembrane transport (Serrano, 1988; Haruta and Sussman, 2012; Taiz and Zeiger, 2014), and, therefore, by extrapolation, the role of xylem sap pH – in *driving* the transport between the xylem and the surrounding BSCs (Shapira et al., 2009). Xylem pH may have also *regulatory* effects on the transport proteins in the BSCs membrane, including aquaporins. Surprisingly, however, even though aquaporins largely determine the osmotic membrane water permeability coefficient, P_f (see also Discussion), and while P_f, particularly the P_f of the BSCs, may well be important in determining the hydraulic conductance (K_{leaf}) of the entire leaf (Shatil-Cohen et al., 2011; Sade et al., 2014; suggested also by modeling, wherein K_{OX}, the lumped Outside-Xylem hydraulic conductance, is equivalent to our definition of K_{leaf} (Scoffoni et al., 2018)), the effect of *extra*cellular pH (or, rather, the lack thereof) has been mentioned only in passing in relation to the P_f of plasma membrane vesicles of *Beta vulgaris* storage root (Alleva et al., 2006) and was ignored in relation to tobacco aquaporins (NtPIP2;1 and NtPIP1) expressed in yeast (Fischer and Kaldenhoff, 2008). Thus, the effect of apoplastic pH on P_f in other plant cells, and especially the effect of xylem sap pH on the BSCs P_f, is unknown. Furthermore, in spite of the importance of apoplast pH regulation to the plant basic life processes (Haruta

and Sussman, 2012), molecular evidence linking the leaf xylem sap regulation to a specific 98
H^+-ATPase and to the ensuing physiological changes in the leaf is missing. 99

H^+-ATPases constitute a family of proton pumps driven by hydrolysis of ATP and are 100
found in the plasma membrane of plants and fungi (Axelsen and Palmgren, 2001). Among 101
the 11 H^+-ATPases isoforms reported in Arabidopsis, the AHA1 and AHA2 are by far the 102
most abundantly expressed members of this family throughout plant life and tissues 103
(Haruta et al., 2010). Both AHA2 (Wang et al., 2014) and AHA1 (Yamauchi et al., 2016) 104
take part in stomatal opening. In addition, a GUS reporter gene under the *AHA2* promoter, 105
revealed abundant expression in Arabidopsis, especially in the vascular tissue of roots and 106
leaves (Fuglsang et al., 2007). Our transcriptome analysis of protoplasts isolated from 107
Arabidopsis BSCs and mesophyll cells (MCs) showed that the BSCs express the *AHA2* 108
gene abundantly and at a threefold higher level than the MCs, while the *AHA1* gene 109
expression, though also abundant, was not different between these two cell types (the other 110
9 AHA isoforms were expressed at much lower levels in both cell types, Wigoda et al., 111
2017, resembling the finding in the whole plant, Haruta et al., 2010). 112

Here we show how the BSCs act as a dynamic barrier for water flow between the xylem 113
and mesophyll; we demonstrate a causative link between the activity of the BSCs AHA2 114
and the hydraulic conductance of the leaf and reveal that its underlying mechanism is the 115
pH-controlled osmotic water permeability of the BSCs membranes. 116

Materials and Methods 117

Plant material 118

Plant types. We used WT (wild type) *Arabidopsis thaliana* plants ecotype Colombia, Col- 119
0 (aka Col) and T-DNA insertion *AHA* mutants (Col): *AHA1* mutants: *aha1-6 (*SALK_ 120
016325), *aha1-7 (*SALK_ 065288) and *aha1-8* (SALK_ 118350), and *AHA2* mutants; 121
aha2-4 (SALK_ 082786) and *aha2-5* (SALK_022010) obtained from the Arabidopsis 122
Biological Resource Center (Ohio State University). The plants' genotypes were 123
confirmed by PCR with allele-specific primers (supplemental Table S1). The single-gene 124
mutants were confirmed further for the near-absence of *AHA2 (or AHA1)* RNA using real- 125

time PCR (RT-PCR). For single-cell P_f experiments, we used *SCR:GFP* (Col) Arabidopsis plants expressing GFP specifically in the BSCs ER generated in our lab as in Attia et al. (2018).

Plant Growth Conditions. The soil grown plants (as in (Shatil-Cohen et al., 2011)) were kept in a growth chamber under short-day conditions (10-h light) and a controlled temperature of 20–22°C, with 70% humidity. Light intensity at the plant height was 150-200 μmol m^{-2} sec^{-1} (an optimal level based on (Wu et al., 2009), achieved using either fluorescent lights (Osram cool white, T5 FH 35W/840 HE, Germany) or LED light (EnerLED 24V-5630, 24W/m, 3000K (50%) + 6000K (50%)). The plants were irrigated twice a week.

Determination of xylem sap pH in detached leaves by fluorescence imaging

To monitor the pH within the leaf minor veins, we used the fluorescence of the membrane-impermeant, ratiometric (dual-excitation) dye, fluorescein isothiocyanate conjugated to 10 kD dextran (FITC-D) perfused via petioles (at a concentration of 100 μM) into detached Arabidopsis leaves (Hoffmann and Kosegarten, 1995; Mühling et al., 1995; Lanz et al., 1997; Pitann et al., 2009b; Pitann et al., 2009a; Suppl. Figs. S1-S2; see Supplemental Materials and methods detailing our specific approach to leaf perfusion and sample preparation for fluorescence imaging).

Fluorescence microscopy was *p*erformed using an inverted microscope (Olympus-IX8 integrated within the Cell-R system, http://www.olympus-global.com), via an Olympus UPlanSApo 10*X*/0.40 (∞/0.17/FN26.5) objective. Pairs of images were recorded using a 12-bit CCD camera, Orca-AG (Hamamatsu, http:// www.hamamatsu.com), at two excitation wavelengths (488 and 450 nm, applied approx. 200 ms apart (Hoffmann and Kosegarten, 1995; Mühling et al., 1995)) and a single emission wavelength (520 nm), and saved in a linear 16 bit tiff format for further processing (Suppl. Materials and methods).

 Image analysis. The images were processed with ImageJ (ver. 1.49V; http://rsbweb.nih.gov/ij/). A pixel by pixel ratio of the two background-corrected images

in each pair was calculated using the ImageJ 'Ratio-plus' plugin (Paulo J. Magalhães, 2004; Suppl. Materials and Methods).

pH calibration curve. An *in-vivo* pH calibration curve was constructed using the dye dissolved in XPS[b] buffered to predefined pH values (5, 5.5,6, 6.5, 7 and 7.5; see Solutions below) and fed into the leaves. This calibration curve (Suppl. Fig. S2d) served to convert the mean fluorescence ratio obtained from a minor vein segment to a pH value for each leaf (biological repeat). Occasionally, we verified the system stability using an *in-vitro* calibration curve derived from imaging drops of the calibration solutions placed directly on microscope slides (Suppl. Fig. S2d).

Generation of *SCR:AHA2*-complemented plants

Vector construction: *AHA2* gene was cloned into pDONR™ 221 (Invitrogene) vector and the SCR promoter into pDONRP4P1r using Gateway® compatible by BP reactions, and later cloned into the pB7M24GW (Invitrogene) two fragment binary vector by LR reaction according to the manufacturer's instructions. The binary *SCR:AHA2* vector was transformed into Agrobacterium by electroporation; transformants were selected on LB plates containing 25 µg/mL gentamycin and 50 µg/mL spectinomycin.

aha2-4 and aha2-5 mutant lines transformation with *SCR:AHA2:* was performed using the floral dip method (Clough and Bent, 1998). Transformants were selected based on their BASTA resistance, grown on plates with MS (Murashige and Skoog, Duchefa cat# M222.0050) Basal medium + 1 % sucrose and 20 µg/mL BASTA (Glufosinate Ammonium, Sigma cat # 45520). DNA insertion was verified in selected lines by PCR targeting the junction of *AHA2* and the 35S terminator with a forward primer about 1000 bp from the 3' end of *AHA2* and reverse primer on the 35S terminator (see primer list in supplemental Table S1).

Determination of AHA2 and AHA1 gene expression level in the whole leaf by qRT-PCR

RNA extraction and quantitative real-time (qRT-) PCR. Total RNA was extracted from leaves using Tri-Reagent (Molecular Research Center, cat. #: TR 118) and treated with RNase-free DNase (Thermo Scientific™, cat. #: EN0525). Complementary DNA (cDNA)

was prepared using the EZ-First Strand cDNA synthesis kit (Biological Industries cat. #: 2080050) according to the manufacturer's instructions. qRT-PCR was performed using C1000 Thermal Cycler (Bio-Rad), in the presence of EvaGreen (BIO-RAD cat.# 172-5204) and PCR primers to amplify specific regions of the genome (Haruta et al., 2010; suppl. Table S1). The results were analyzed using Bio-Rad CFX manager™ software. Dissociation curve analysis was performed at the end of each qRT-PCR reaction to validate the presence of a single reaction product and lack of primer dimerization. Expression levels of the examined genes were normalized using two normalizing genes (AT5G12240 and AT2G07734, Wigoda et al., 2017).

Physiological characterization of the leaf (gas exchange and hydraulic conductance, K_{leaf})

Sample preparation. In all K_{leaf} determinations, we followed our previous protocols (Shatil-Cohen et al., 2011; Sade et al., 2014; Sade et al., 2015), with a necessary adaptation in the present work (detailed in the Suppl. Materials and methods) due to an approx. three-fold increase in the growth chamber light intensity (to 150-200 μmol m^{-2} sec^{-1} (Wu et al., 2009)), which resulted in increased stomatal conductance and transpiration. Prior to the gas-exchange measurements, the leaves were exposed to constant ambient conditions (1.3-1.5 kPa vapor pressure deficit (VPD), 200 μmol m^{-2} s^{-1} illumination and temperature of 24 °C) for 15-20 minutes. During this period the transpiration, E, reached steady state (the values of E varied on average within 4-5%).

The measurements were conducted between 10:00 AM to 1:00 PM (1-4 hours after Lights On).

Gas-exchange assays to determine g_s (stomatal conductance) and E (transpiration rate) were performed (as in Sade et al., 2014) using a Li-Cor 6400 portable gas-exchange system (LI-COR, USA https://www.licor.com/) equipped with a standard leaf cuvette with an aperture diameter of two cm^2, appropriate for Arabidopsis leaves, using light intensity similar to the growth chamber conditions (illumination: 200 μmol m^{-2} s^{-1}; the amount of blue light set to 10% of the photosynthetically active photon flux density, temperature:

approximately 22 °C, VPD: approximately 1.3 kPa, and [CO_2] surrounding the leaf: 400   211

µmol mol^{-1}). Readings were recorded at a steady-state E, 3-5 min after the leaf was   212

clamped in the Li-Cor chamber.   213

Measuring the leaf water potential, Ψ_{leaf}. Immediately following the gas exchange   214

measurement, the leaf was transferred to a pressure chamber (ARIMAD-3000; MRC   215

Israel) equipped with a home-made silicon adaptor especially designed to fit Arabidopsis   216

petioles into the chamber's O-ring. Ψ_{leaf} was determined as described earlier (Sade et al.,   217

2014).   218

Determination of the leaf hydraulic conductance, K_{leaf} in detached leaves. K_{leaf} was   219

calculated for each individual leaf as follows (Martre et al., 2000):   220

Eq. 1: $K_{leaf} = -E / (\Psi_{leaf} - \Psi_{AXS}) \approx -E / \Psi_{leaf}$,   221

where E is the whole-leaf's transpiration, i.e., the water flux through the leaf, Ψ_{leaf} is the   222

leaf water potential and Ψ_{AXS} is the water potential of AXS (or of the XPS^{db}); as Ψ_{AXS} is   223

negligible in comparison to the value of leaf water potential, Ψ_{leaf}, alone was used.   224

Determination of the leaf hydraulic conductance, K_{leaf}, in intact leaves in a whole plant.   225

On the evening before measurements, one leaf from each tested plant was covered by a   226

small light-proof plastic bag. In the morning (1-4 hours after Lights On), another,   227

illuminated leaf of the same size, positioned closely on the same plant underwent gas   228

exchange measurements by Licor 6400 while still attached to the plant, then it was excised   229

with a scalpel and was immediately placed in a pressure chamber to determine its water   230

potential (as described above), taken to represent the water potential of transpiring   231

illuminated leaves of the whole plant ($\Psi_{Illuminated\ Leaf}$). The water potential of the dark-   232

plastic-covered leaf – taken to represent the water potential within the plant stem (Ψ_{Dark} 233

$_{\text{Leaf}} = \Psi_{\text{Stem}}$) – was determined in an identical procedure just before the illuminated leaf. 234

K_{Leaf} was then calculated from: 235

Eq. 2: $\qquad K_{\text{Leaf}} = -E / (\Psi_{\text{Illuminated leaf}} - \Psi_{\text{Dark leaf}})$ 236

Osmotic water permeability coefficient (P_f) determination 238

Protoplast isolation for P_f experiments. Protoplasts were isolated from leaves (with 239
verified GFP-fluresceing-veins) of 6- to 8-week-old plants with GFP-labeled BSCs (see 240
'plant material' above) by the rapid and low-stress method of (Shatil-Cohen et al., 2014) 241
using an isotonic solution in the extraction process (pH 6, 600 mOsmol, see Solutions 242
below). After isolation, only cells expressing GFP – i.e., only BSCs – were assayed for 243
P_f determination. 244

P_f determination. P_f was determined as described by Shatil-Cohen et al. (2014), except 245
here we used an inverted epifluorescent microscope (Nikon eclipse TS100) with a 20x/NA 246
0.40 objective (Nikon) and a CCD 12 bit camera Manta G-235B 247
(https://www.alliedvision.com), and an image-processing software AcquireControl® 248
v5.0.0 (https://www.alliedvision.com). We recorded the BSCs swelling in response to 249
hypo-osmotic challenge (of 0.37 MPa) generated by changing the bath solutions from 250
isotonic (600 mOsm) to a hypotonic one (450 mOsm, *ibid.*). The challenges were 251
performed at pH 6 and at pH 7.5. The osmolarity of these solutions (variants of the XPS[db]; 252
see Solutions) was adjusted with D-sorbitol and was verified within 1% of the target value 253
using a vapour pressure osmometer (Wescor). P_f was determined by fitting the timecourse 254
of the cell swelling using an offline curve-fitting procedure of the P_fFit program, as 255

described in Shatil-Cohen et al., (2014) and detailed in Moshelion et al., (2004). We
present here the values of the intital P_f (P_{f_i}) obtained by fitting model 5 (ibid.).

Solutions

FITC-D dye (Sigma cat. #: FD10S) was added from a 10 mM stock in water (kept
aliquoted, protected from light, at -20 ^{o}C) to all the XPS solutions (except in the leaf-
tissue-autofluorescence (background) assays) to a final conc. of 100 µM.

XPS, basal Xylem Perfusion Solution: 1 mM KCl, 0.3 mM $CaCl_2$ and 20 mM D-sorbitol
(Sigma cat# 8143). Upon preparation, the pH of this solution was 5.6 -5.8 and its
osmolarity was approx. 23 mOsm/L.

Low-K^+ XPS: the same as XPS above (used in experiments of Figs. 1, 2, 3b, Suppl. Figs.
S1, S2a-c) .

High-K^+ XPS: XPS + 10 mM KNO_3 and D-sorbitol adjustment to approx. 23 mOsm/L.
Upon preparation, when unbuffered, the pH of this solution was 5.6 -5.8 (used in
experiments of Fig. 1).

XPS^b (pH calibration solutions): XPS buffered with 20 mM MES and HCl or NMG (N-
Methyl D-glucamine) to pH 5, 5.5, or 6), or with 20 mM HEPES and NMG to pH 6.5, 7,
or 7.5. Osmolarity of these solutions was adjusted with D-sorbitol to 23 mOsm/L (used in
experiments of Suppl. Fig. 2d-e).

AXS (*for leaf physiological characterization*): 3 mM KNO_3, 1 mM $Ca(NO_3)_2$, 1 mM
$MgSO_4$, 3 mM $CaCl_2$, 0.25 mM NaH_2PO_4, 90 µM EDFS (Sigma) and a micro-nutrient mix
of 0.0025 µM $CuSO_4$, 0.0025 µM H_2MoO_4, 0.01 µM $MnSO_4$, 0.25 µM KCl, 0.125 µM
H_3BO_3, 0.01 µM $ZnSO_4$; 21 mOsm; pH of the non-buffered solution was 5.8 (used in
experiments of Fig. 4a and Suppl. Figs. S3a, S3c, S3e).

XPS^{db} (*for leaf physiological characterization*): High-K^+ XPS buffered with 5 mM MES
+ 5 mM HEPES, and adjusted to pH 4.5, 6 or 7.5 with HCl or NMG. Each solution was

adjusted with D-sorbitol to a final osmolarity of 32 mOsm/L (used in experiments of Fig. 282

4c and Suppl. Figs. S3b, S3d, S3f). 283

Solutions for P_f determination, pH 6 or pH 7.5 solutions: XPS[db] adjusted to pH 6 or pH 284

7.5 by NMG, and adjusted with D-sorbitol to either 450 mOsmol (hypotonic) or 600 285

mOsmol (isotonic). 286

Solution for protoplasts isolation: pH 6 isotonic solution as for P_f determination. 287

All chemicals unless mentioned otherwise were from Sigma Israel. 288

Statistics 289

The Student's unpaired two-tailed t-test was used for comparison of two means, which 290

were considered to be significantly different at $P < 0.05$. For comparisons of three or more 291

population means we used ANOVA, all-pairs Tuckey HSD (JMP® Pro 13), wherein 292

different letters represent statistical differences at $P<0.05$. Images which yielded extreme 293

ratio values which were more than 2.5 SD away from the mean were discarded. Curve 294

fitting statistics was evaluated with ANOVA (OriginPro 2020, from OriginLab 295

Corporation, USA). 296

Results 299

The effect of pharmacological agents – a pump stimulator and an inhibitor – on xylem 300

pH 301

To assay the activity of AHA proton pumps in the xylem-facing membranes of bundle 302

sheath cells (BSCs) lining the Arabidopsis leaf veins, we applied fusicoccin, a known 303

fungal stimulator of P-type proton pumps (Serrano, 1988). When 10 µM fusicoccin was 304

fed into the xylem, we observed a sharp decrease in xylem sap pH of roughly 1.5 pH units 305

compared with leaves in control conditions (Figs. 1a-i, 1a-ii, 1b) approximately 30 minutes 306

after addition of fusicoccin. ETOH, the fusicoccin solvent, by itself, had no impact on xylem pH (Suppl. Fig. S4).

Further, WT leaves treated with 1 mM of vanadate (a commonly used P-type H^+-pump inhibitor (reviewed by Palmgren, 2001), in a high-K^+ solution (10 mM KNO_3), resulted in xylem sap alkalization of about one pH unit within 30-40 min (Figs. 1a-iii, 1a-iv, 1c). Interestingly, neither a similar exposure to vanadate in the low-K^+ solution (XPS without added KNO_3, suppl. Fig. S5a), nor the high-K^+ solution by itself (Suppl. Fig. S5b) had any effect on the xylem sap pH.

AHA2 acidifies the xylem pH

In an attempt to resolve between the relative contributions of the two abundant H^+-ATPAses of the BSCs, AHA1 and AHA2 (Wigoda et al., 2017), to the acidification of the xylem pH, we compared the xylem sap pH of WT plants to that in T-DNA-insertion mutants of either pump. We used two independent mutant lines with a homozygous loss of function of the *AHA2* gene (*aha2-4* and *aha2-5*) and three such lines with mutated *AHA1* (*aha1-6, aha1-7, aha1-8*). The xylem sap pH of both *AHA2* mutants, *aha2-4* and *aha2-5,* was consistently higher, by 0.5-1 pH units, compared to the WT plants (Fig. 2). We did not find a significant difference between the xylem sap pH in WT vs. the three lines of AHA1 mutants (Suppl. Fig. S6).

To further test the ability of AHA2 to acidify the xylem sap pH, we complemented the *AHA2* deficient plants (the *aha2-4* mutant), with the *AHA2* gene directed specifically to the BSCs (under the BSCs-specific promotor Scarecrow, SCR; Wysocka-Diller et al., 2000). Using qRT-PCR on whole leaves of mature transgenic plants, we confirmed *AHA2* absence in the *aha2-4* mutant (Fig. 3a, as in Haruta et al., 2010), and demonstrated successful complementation in two lines (T55 and T56). The transcript level of *AHA2* in these lines was even higher than in the WT (Fig. 3a). While the xylem sap pH in the *AHA2*-mutant, *aha2-4*, was significantly more alkaline than in WT plants (repeating the results of Fig. 2a above), upon the complementation of the *aha2-4* mutant with the BSC-directed *AHA2*, the xylem sap pH in both lines became no higher than WT (Fig 3b). The level of

AHA1 transcript in the whole leaf was not affected by the genetic manipulations of AHA2, neither by *AHA2* mutation (as in Haruta et al., 2010), nor by the BSC-specific complementation of the mutant with *AHA2* (Supl. Fig. S7).

Elevating pH lowers the leaf hydraulic conductance, K_{leaf}

To examine the impact of AHA2 on the water economy of the whole leaf we determined the leaf water potential (Ψ_{leaf}) and its transpiration (E) at steady-state, and from these two we calculated the leaf hydraulic conductance (K_{leaf}, Eq. 1 in Materials and methods).). We compared the K_{leaf} in detached leaves of WT, of the *aha2-4* and *aha2-5* mutants (respectively, 22.4, 12.0 and 14.7 mmol m^{-2} s^{-1} MPa^{-1}), and of *aha2-4* complemented with the BSCs-directed *AHA2* (i.e., *AHA2* under the Scarecrow promoter; line T56, which presented the lowest xylem pH of all genetic lines), all fed with the non-buffered AXS. Notably, while all four leaf types appeared to transpire at a similar rate, E (Suppl. Fig. S3e; but see Discussion), the K_{leaf} of the mutants was appreciably lower than that of WT leaves, about 50% of WT in *aha2-4* and about 35 % in *aha2-5*, while K_{leaf} of the AHA2-complemented T56 line (23.2 mmol m^{-2} s^{-1} MPa^{-1}) was practically as high as that of the WT leaves (Fig. 4a).

In the intact leaves in whole plants, while E also did not appear to differ among the three plant lines (Suppl. Fig. S8), K_{leaf} values were similar to those of K_{leaf} in detached leaves and we observed the same pattern: K_{leaf} of intact *aha2-4* plants was by about 60% lower than in WT (respectively, 13.5 vs. 34.7 mmol m^{-2} s^{-1} MPa^{-1}), while K_{leaf} of T56 plants (24.4 mmol m^{-2} s^{-1} MPa^{-1}) was not significantly different from WT (Fig 4b).

Furthermore, and again in contrast to E, which appeared unaffected (Suppl. Fig. S3f), pH treatments did alter the K_{leaf} of detached WT leaves. Perfusion with XPS[db] buffered at pH 7.5 lowered K_{leaf} by roughly 50% compared to K_{leaf} in WT leaves perfused with XPS[db] buffered at pH 6 or WT leaves perfused with the non-buffered XPS (respectively, 11.6, 24.9 and 26.6 mmol m^{-2} s^{-1} MPa^{-1}; Fig. 4c). In contrast to K_{leaf} in the more alkaline conditions, K_{leaf} of leaves perfused with the more acidic pH 4.5 (24.0 mmol m^{-2} s^{-1} MPa^{-1}) was no different than at the "normal" pH 6 (Fig. 4c).

Alkaline pH lowers the osmotic water permeability coefficient of BSCs protoplast membrane, P_f   364
365

In order to test the hypothesis that the pH-dependent reduction in K_{leaf} is due to a reduction   366

in the water permeability of the BSCs membranes, we determined the osmotic water   367

permeability coefficient, P_f, of BSC under two pH treatments, using GFP labeled BSCs,   368

from *SCR:GFP* plants. We verified that these plants did not differ in their physiological   369

parameters (K_{leaf} and the underlying E and Ψ_{leaf}) from the unlabeled WT plants (Suppl.   370

Fig. S9). The mean initial P_f (see Materials and methods) of BSCs assayed in pH 6 was   371

~8.5 fold higher than that of BSCs assayed in pH 7.5. ($5.39 \pm 1.0 \, \mu m \, sec^{-1}$ and 0.63 ± 0.2 372

$\mu m \, sec^{-1}$, respectively; Fig. 5).   373

Leaf veins architecture   374

We tested the hypothesis that K_{Leaf} values are proportional to the area of the bundle sheath   375

enwrapping the leaf veins, and therefore to the combined length of leaf veins per unit leaf   376

area (vein density). Thus, leaves of the *aha2-4* plants would be expected to have lower   377

vein density than leaves of WT and *SCR:AHA2* (T56) plants. However, a comparison   378

(Suppl. Materials and methods) yielded no difference among these lines (Fig. 6, Suppl.   379

Fig. S10), eliminating this structural reason for differences in K_{leaf}.   380

Discussion   382

Our conclusions from this study are based on two complementing AHA manipulation   383

approaches: (1) classical pharmacology (using fusiccocin and vanadate) and (2), AHA2   384

genetics (mutants and complementation), combined with in-vivo measurements of the   385

resulting xylem sap pH, using the petiole-fed membrane-impermeable FITC-dextran. An   386

additional approach was to assay the physiological functions in the single-cell (P_f), the   387

detached leaf (K_{leaf}), and the whole plant (as reflected in the *intact* leaf K_{leaf}), combined   388

with the modulation of the xylem sap pH (by genetic AHA2 manipulations or by direct pH   389

buffering of the BSCs-facing solutions).   390

The AHA2 of the BSCs is indispensable for the leaf xylem sap acidification

Plasma membrane H^+-ATPases (AHAs) are well-known regulators of the apoplastic pH and have a key role in cell ion homeostasis. Here we demonstrate that the BSCs AHA2 exerts a dominant role in the acidification of the xylem sap in the Arabidopsis leaf.

An AHA-specific stimulator (fusicoccin) and an inhibitor (vanadate), fed directly to the leaf xylem, established that pH acidification of the leaf xylem sap involves P-type H^+-ATPases, which extrude protons – most likely, from the BSCs – into the xylem lumen (Fig. 1). Notably, to demonstrate vanadate effect (xylem sap alkalinization) the high-K^+ XPS was necessary, possibly to dissipate the xylem sap protons by enhanced H^+/K^+ (accompanied by $2H^+/NO_3^-$) co-transport into the BSCs. That proton-coupled K^+ transmembrane co-transport plays a particularly important role in the BSCs is suggested by the 10% higher expression in BSCs (relative to mesophyll cells) of two K^+ uptake permeases, AtKT2 and AtKUP11 (likely to be H^+-coupled (Wigoda et al., 2017) and references therein). This conforms with the aforementioned general notion that the pH of an apoplastic compartment reflects, among others, a balance of bi-directional proton movements (Serrano, 1988).

While pH in the *aha1* mutant lines was not significantly different from that in WT (Suppl. Fig. S6), the pH of the xylem sap in both AHA2 mutant lines, a*ha2-4* and *aha2-5* was consistently above that in WT (Fig. 2), similar to the pH elevated by vanadate inhibition (Fig. 1C). Two lines of *aha2-4* complemented with the *AHA2* gene directed specifically into the BSCs, using the specific promotor SCR, had their xylem sap pH restored to WT-like levels. These results together underlined the importance of AHA2 in determining the xylem sap pH.

Extracellular pH regulation of the membrane osmotic water permeability

Earlier work established that low *cytosolic* pH reduces the membrane osmotic water permeability coefficient, P_f, by inhibiting aquaporin gating (Gerbeau et al., 2002; Tournaire-Roux et al., 2003; Alleva et al., 2006; Fischer and Kaldenhoff, 2008; Leitão et

al., 2012; Frick et al., 2013; Yaneff et al., 2016). This phenomenon is especially relevant   418
to roots under stress conditions such as flooding or drought, where a change in the water   419
hydraulic permeability of the roots is considered to be part of the plant stress defense   420
mechanism.   421

External pH seemed not to affect the membrane hydraulic permeability of intact cells (Lp)   422
of Nitella, Chara and maize, except a slight inhibition at pH around 4.5 in Nitella and in   423
one variety of maize (Tyerman et al., 2002 and references therein). However, P_f sensitivity   424
to pH has been also found in several mammalian cell types, be it cytosolic pH (Kaptan et   425
al., 2015; Gotfryd et al., 2018), or extracellular pH (Mosca et al., 2018). Here we   426
demonstrate, for the first time in an intact plant protoplast, the BSC, a reduction in P_f by   427
external (xylem sap) alkalinization. This pH sensitivity differs from the *lack* of external   428
pH effect on the right-side-out plasma membrane vesicles of *Beta vulgaris* root (Alleva et   429
al., 2006). However, this difference is not surprising, in view of the differences between   430
root and shoot, for example, in their opposite responses to drought: acidification in the root   431
xylem sap and alkalinization in the shoot xylem sap (Karuppanapandian et al., 2017).   432

 "Macroscopic" membrane P_f changes and, moreover, whole-organ K_{leaf} changes have   433
been tied to changes in aquaporin activity. For example, Shatil-Cohen et al., (2011),   434
reduced both P_f and K_{leaf} by a common aquaporin blocker $HgCl_2$. Also, Prado et al.   435
demonstrated that three aquaporins expressed in veins, and in particular, PIP2;1,   436
contributed to darkness-enhanced hydraulic water conductance of the Arabidopsis leaves   437
rosette (Prado et al., 2013). Additionally, BSCs-directed knockdown of aquaporins (using   438
artificial microRNAs under the SCR promoter) decreased the K_{leaf} of a detached   439
Arabidopsis leaf and, separately, the BSCs P_f (Sade et al., 2014). Therefore, it is interesting   440
to compare the external pH effect we observed on the P_f of BSCs to the effects of external   441
pH on aquaporins. For example, an external acidification (from pH 7.4 to pH 5), perceived   442
via external tyrosine and histidine, *decreased* the permeability to water and glycerol of the   443
human aquaglyceroporin, AQP7, with a half-inhibition at about pH 6 (Mósca et al., 2018);   444
in contrast, the fungal aquaporin RdAQP1 activity (when expressed in Arabidopsis   445

protoplasts) was enhanced at acidic external pH and inhibited at alkaline external pH (Turgeman et al., 2016).

Like RdAQP1, at least several Arabidopsis aquaporins have histidines (pka = 6.8) and/or cysteins (pKa=8.5) in their apoplast-facing loops, and also plenty of tyrosines (pKa=10). For example (even excluding tyrosines, because of their high pKa), based on uniprot membrane topology (www.uniprot.org/), PIP2;1 has exterior-facing Cys75 and His248; PIP2;2, has exterior-facing Cys73; His246; PIP2;3: Cys73, His148,-246, etc. Future work will clarify whether the P_f reduction in BSCs and the reduction of K_{leaf} in detached Arabidopsis leaves (see also further discussion of K_{leaf} and P_f relationship below) is also mediated by aquaporins and their external histidines/cysteins.

AHA2 role in the regulation of the whole leaf water balance

The notion that AHA pumps, in general, power secondary H^+-cotransport across cell membranes facing the apoplast is decades old (Spanswick, 1981; Serrano, 1988; Sze et al., 1999; Palmgren, 2001). The plant plasma membrane H^+-ATPase has since been recognized as essential for plant growth (reviewed by Falhof et al., 2016). The practically sole "celebrated product" of its action to date is the proton motive force (PMF) – a gradient of protons concentration without- or in combination with an electrical gradient – which drives the movement of solutes across cellular membranes. In only a few cases has a *specific* physiological role been assigned to the plasma membrane H^+-ATPase – mainly in stomatal physiology and in roots (reviewed by Falhof et al., 2016).

Here we show another novel aspect of transport activity *regulated* (rather than *driven*) by AHA2 – that of water fluxes from the xylem into the leaf and across the bundle sheath layer – evident as the leaf hydraulic conductance, K_{leaf}. Our summary figure (Fig. 7) demonstrates this regulation by juxtaposing the separate measurements of of K_{leaf} (Fig. 4a) and xylem sap pH (Figs. 2, 3b) performed in independent experiments on detached leaves of WT plants, aha2 mutants and the *SCR:AHA2*-complemented aha2-4. This combination demonstrates the inverse dependence of K_{leaf} on the xylem sap pH, in particular roughly between pH 5.5 and 6.5 (Fig. 7). The inverse K_{leaf} - pH relationship is also demonstrable

when pH is imposed directly by buffers in the xylem perfusate in the range of pH 6 to 7.5   474

(Fig. 4c, Fig. 7). Yet, while the absolute values of the maximum and the minimum K_{leaf} 475

were similar in the two experiment types, there was a two-fold difference in the K_{leaf} values   476

at the pH transition range (around pH 6): K_{leaf} was high (approximately 25 mmol m^{-2} s^{-1} 477

MPa^{-1}) in the WT leaf with the xylem sap buffered at pH 6 and it was low (approximately   478

12 mmol m^{-2} s^{-1} MPa^{-1}) in the *aha2-4* mutant leaf with the non-buffered xylem sap dye-   479

reported to be at pH 6 (Fig. 7). This apparent disparity could be bridged, if, due to the   480

ongoing activity of their AHA2, the BSCs in the leaves of the WT plants faced a nearby-   481

layer of medium more acidic than the bulk of the pH 6-buffered xylem sap. We   482

represented such possible inadequacy of the perfusate buffer by a hand-drawn elipse (Fig.   483

7). The notion that the bulk pH may differ from the pH near the membrane is supported   484

by the report of Martinière et al. (2018), who showed that the pH reported by a pH-dye in   485

the bulk of the apoplast (here: the xylem sap) did not fully represent the pH existing in the   486

immediate vicinity of the membrane. Bridged by the "elipse-correction", the results of the   487

different pH manipulations (the genetic and the imposed buffering), together attest to a   488

causative inverse correlation between the xylem sap pH and the resulting K_{leaf}.   489

What is the physiological relevance of the pH-regulated K_{leaf} changes? The relatively high   490

range of K_{leaf} values in detached leaves that we report here is characteristic of well-watered   491

Arabidopsis plants, as reported also elsewhere (e.g., (Guyot et al., 2012; Scoffoni et al.,   492

2018). This is exemplified in the resemblance between the relationships of the individual   493

values of K_{leaf} vs Ψ_{leaf} (the leaf water potential) in detached and also in intact leaves (Suppl.   494

Fig. S11) and in those of Scoffoni et al. (2018, in their Fig. 1; particularly noteworthy is   495

the quantitative agreement between our findings and Scoffoni's – in detached leaves – of   496

the values of Ψ_{leaf} at which K_{leaf} declines to 50% of its measured maximum (Scoffoni's   497

"P50"), -0.17 to -0.16 MPa).   498

Since the leaf water potential (Ψ_{leaf}) reflects the relative insufficiency of water influx to   499

the leaf in replacement for the transpired water (prior to establishing a steady-state), it is   500

meaningful to examine the inverse relationship: how Ψ_{leaf} depends on K_{leaf}. Indeed, when   501

considering the paired measurements of Ψ_{leaf} and K_{leaf} of individual leaves, both detached   502

and intact, a positive correlation can be demonstrated even among the well-hydrated plants   503

(Suppl. Figs. S12a-S12c). Similarly, the rate of transpiration (E) also increases with K_{leaf} 504

(Suppl. Figs. S12d-S12f). Incidentally, the E-K_{leaf} correlation is identical to a g_S (stomata   505

conductance)-K_{leaf} correlation, since g_S=E/ VPD (Vapor Pressure Difference, the   506

transpiration drive, which is kept constant during the Li-Cor measurements (Materials and   507

methods)). These correlations strongly suggest that pH-controlled physiological effects –   508

K_{leaf} and E changes – even if relatively subtle and masked by the large variability when   509

the averages are considered, may occur in intact plants and that, in the long run, they may   510

have cumulative effects on agricultural yield.   511

The above considerations of the AHA2 role invite a speculation with regard to the role of   512

the plant stress hormone, abscisic acid (ABA) in regulating K_{leaf}. In an earlier work, Shatil-   513

Cohen et al. (Shatil-Cohen et al., 2011) localized K_{leaf} to the BSCs layer by showing that   514

K_{leaf} decreased when the BSCs were brought into direct contact with ABA fed to the   515

detached leaf, and not when ABA was smeared on the leaf surface. Our current work   516

outlines a mechanism for this phenomenon: stress-induced xylem alkalinization (already   517

reported by others (Jia and Davies, 2007; Wang et al., 2012; Korovetska et al., 2014;   518

Karuppanapandian et al., 2017), likely due to ABA inhibition of the BSCs AHA2   519

(although not excluding other AHAs), similar to the H^+-ATPase inhibition seen in guard   520

cells (Goh et al., 1996; Schroeder et al., 2001; Zhang et al., 2004), reduces the water   521

permeability of the BSCs membranes. Consequently, the water permeability of the entire   522

bundle sheath layer declines, demonstrating that under stress conditions it is an active   523

hydraulic barrier (Shatil-Cohen et al., 2011; Pantin et al., 2013). Here we show directly,   524

using individual isolated BSCs, that P_f, the measure of the rate of water passage via the   525

BSCs membranes, declines with extracellular medium alkalinization (Fig. 5). Such a pH   526

rise could mediate the effect of ABA (in a feedback loop?) on BSCs in the detached leaf.   527

Another, slower route of generating xylem sap alkalinization would be via decreasing the   528

expression and/ or activity of AHA2 in the BSCs. In accord with this   529

prediction, Genevestigator analyses (Hruz et al., 2008) of data from a few available   530
studies, revealed that drought decreased the AHA2 expression in Arabidopsis rosettes by   531
about 20-60% (Ludwików et al., 2009 (ArrayExpress Repository, Experiment E-MEXP-   532
1863); Wilkins et al., 2010 (GEO repository, Expt. GSE19700)), which is in agreement   533
with the reported alkalization of the apoplast during drought. In contrast, under drought,   534
AHA1 expression was either unchanged (in the two above-mentioned databases) or   535
upregulated by ~25% (Pandey et al., 2013 (GEO repository, Experiment GSE40061)),   536
suggesting different roles for these pumps in the leaf, at least under this type of stress.   537
Moreover, genetic manipulation of AHA2 left the quite abundant AHA1 transcript   538
(Wigoda et al., 2017) unaffected, and AHA1 mutation left the xylem sap pH unchanged,   539
supporting such differentiation. Notwithstanding the above findings, we have not ruled out   540
completely a potential contribution of AHA1 to the xylem sap pH. Both AHAs need yet   541
to be investigated in *BSCs-focused* genetic manipulations (overcoming perhaps the   542
reported lethality of the double *aha1-aha2* mutant (Haruta et al., 2010)) and in BSCs-   543
specific responses to drought.   544

The observed variations in transpiration rate of whole plants have been proposed to reflect   545
internal adjustments – not only at the guard cell level but also at the K_{leaf} level – protecting   546
the plant against challenges of changing ambient conditions (Wallach et al., 2010; Scoffoni   547
et al., 2017; Scoffoni et al., 2018). We propose here that the dynamic adjustments in water   548
flux through the transpiring leaf are due to the dynamic changes of BSCs' P_f in response   549
to changes of xylem sap pH and the consequent changes in K_{leaf}. In a transpiring   550
plant, K_{leaf} decreases as a consequence of xylem alkalinization in the shoot, which signals   551
a decline of soil moisture. The K_{leaf} decrease, leading to leaf water potential decrease and   552
consequently to stomata closure, is what prevents further water escape from the xylem and   553
decreases the water tension in the xylem, diminishing the danger of cavitation and   554
embolism (see also, (Guyot et al., 2012; Scoffoni et al., 2018).   555

How general is drought-stress-induced shoot alkalinization? While various authors   556
demonstrated shoot xylem sap pH increase in annual plants upon gradual soil drying (for   557

example, in sunflower from around 6 to 7 (Gollan et al., 1992); in *Commelina communis* 558
from around 6 to over 6.5 (Wilkinson and Davies, 1997);   in tomato, from 5 to 8   559
(Wilkinson et al., 1998); in *Vicia faba* from 5.9 to 6.8 (Karuppanapandian et al. (2017),   560
Sharp and Davies (2009) tabulated different responses of the shoot xylem pH to drought   561
in different perrenial plants. Therefore, in view of our work results, the leaf xylem sap pH   562
response to water deficit and the dependence of K_{leaf} on xylem sap pH need to be examined   563
also in other plants in order to understand the universality of the mechanism we propose.   564

The results of this study broaden our basic understanding of how a leaf controls its water   565
influx supporting the notion that xylem sap alkalinization mediates the decline in plant   566
shoot nutrient uptake due to abiotic stress. Here we focus specifically on the stress-   567
induced cessation of activity of AHA2 in the BSCs in Arabidopsis, where we found the   568
AHA2 instrumental in generating the low pH in the leaf xylem sap.   569

  The consequence of this decline of AHA2 activity and xylem sap alkalinization, is a   570
decline in the PMF. Since AHA2 – via the PMF – powers the secondary, proton-coupled   571
export and import of solutes across the BSCs membranes, not only does it regulate plant   572
nutrition, but, very likely also drives the extrusion of toxic compounds from the xylem-   573
lining cells into the xylem, hence governing also the plant toxicity tolerance as well as   574
others potential stresses. These hypotheses await future experimentation.   575

In conclusion, our finding that the Arabidopsis AHA2 plays a major role in regulating K_{leaf} 576
via xylem sap pH provides a molecular basis for understanding a novel aspect, other than   577
just PMF, of the control that the xylem sap pH can exert in the leaf. This control, in   578
combination with the effects of PMF, is likely to underlie the plant's key physiological   579
activities. These results provide a new focus for exploration and understanding the role of   580
the involvement of BSCs in determining the xylem sap composition, and, in particular,   581
their role as a transpiration-controlling valve in series with the stomata. As the rapid   582
growth rate and high yields of crop plants are positively correlated with enhanced   583
transpiration and K_{leaf}, BSCs are likely to become a key target tissue for the development   584
of a new generation of manipulations for plant adaptation to environmental challenges.   585

Acknowledgments

We thank Ms. Dvorah Weisman of the Hebrew University of Jerusalem for help with image analysis and Dr. Shifra Ben-Dor of the Weizmann Institute of Science for comments on the manuscript.

This research was supported by the Israel Science Foundations, ISF (grants No.1312/12, and 1842/13 to NM, grant No. 878/16 to MM and a grant No. 12-01-0007 from the Israeli Chief Scientist, Ministry of Agriculture & Rural Development to MM and NM).

Author Contributions

Author contributions:

Y. Grunwald and N. Wigoda planned, performed and analyzed the vein pH monitoring and RT-PCR experiments, Y. Grunwald planned, performed and analyzed K_{leaf} and P_f experiments.Y. Grunwald and N. Wigoda wrote the paper. A. Yaaran and T. Torne participated in generating the complemented plant lines, and A. Yaaran and S. Gosa participated in the leaf hydraulics determinations. N. Sade participated in the genotyping of the mutant AHA lines. M. Moshelion and N. Moran conceived the experiments, guided the students and wrote the paper.

Competing interests

All authors declare they have no competing interests.

Supplementary Materials

References and Notes   631

Alleva K, Niemietz CM, Sutka M, Maurel C, Parisi M, Tyerman SD, Amodeo   634

 G (2006) Plasma membrane of Beta vulgaris storage root shows high water   635

 channel activity regulated by cytoplasmic pH and a dual range of calcium   636

 concentrations. J Exp Bot **57**: 609–621   637

Attia Z, Dalal A, Moshelion M (2018) Vascular bundle sheath and mesophyll   638

 regulation of leaf water balance in response to chitin. bioRxiv. doi: 639

 10.1101/337709   640

Axelsen KB, Palmgren MG (2001) Inventory of the superfamily of P-type ion   641

 pumps in Arabidopsis. Plant Physiol **126**: 696–706   642

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-   643

 mediated transformation of Arabidopsis thaliana. Plant J **16**: 735–743   644

Falhof J, Pedersen JT, Fuglsang AT, Palmgren M (2016) Plasma membrane   645

 H+-ATPase regulation in the center of plant physiology. Mol Plant **9**: 323–   646

 337   647

Felle HH, Herrmann A, Hückelhoven R, Kogel KH (2005) Root-to-shoot   648

 signalling: Apoplastic alkalinization, a general stress response and defence   649

 factor in barley (Hordeum vulgare). Protoplasma **227**: 17–24   650

Fischer M, Kaldenhoff R (2008) On the pH regulation of plant aquaporins. J Biol   651

 Chem **283**: 33889–33892   652

Frick A, Järvå M, Törnroth-Horsefield S (2013) Structural basis for pH gating   653

 of plant aquaporins. FEBS Lett **587**: 989–993   654

Fuglsang AT, Guo Y, Cuin T a, Qiu Q, Song C, Kristiansen K a, Bych K,   655

 Schulz A, Shabala S, Schumaker KS, et al (2007) Arabidopsis protein   656

 kinase PKS5 inhibits the plasma membrane H+ -ATPase by preventing   657

 interaction with 14-3-3 protein. Plant Cell **19**: 1617–34   658

Geilfus CM (2017) The pH of the apoplast: dynamic factor with functional impact   659

under stress. Mol Plant **10**: 1371–1386   660

Gerbeau P, Amodeo G, Henzler T, Santoni V, Ripoche P, Maurel C (2002)   661
The water permeability of *Arabidopsis* plasma membrane is regulated by   662
divalent cations and pH. Plant J **30**: 71–81   663

Goh CH, Kinoshita T, Oku T, Shimazaki K (1996) Inhibition of blue light-   664
dependent H+ pumping by abscisic acid in Vicia guard-cell protoplasts. Plant   665
Physiol **111**: 433–440   666

Gollan T, Schurr U, Schulze E-D (1992) Stomatal response to drying soil in   667
relation to changes in the xylem sap composition of Helianthus annuus. I. The   668
concentration of cations, anions, amino acids in, and pH of, the xylem sap.   669
Plant, Cell Environ **15**: 551–559   670

Gotfryd K, Mósca AF, Missel JW, Truelsen SF, Wang K, Spulber M, Krabbe   671
S, Hélix-Nielsen C, Laforenza U, Soveral G, et al (2018) Human adipose   672
glycerol flux is regulated by a pH gate in AQP10. Nat Commun **9**: 4749   673

Grignon C, Sentenac H (1991) pH and ionic conditions in the apoplast. Annu Rev   674
Plant Physiol Plant Mol Biol **42**: 104–121   675

Guyot G, Scoffoni C, Sack L (2012) Combined impacts of irradiance and   676
dehydration on leaf hydraulic conductance: insights into vulnerability and   677
stomatal control. Plant Cell Environ **35**: 857–871   678

Hartung W, Radin JW, Hendrix DL (1988) Abscisic acid movement into the   679
apoplastic solution of water-stressed cotton leaves: role of apoplastic pH.   680
Plant Physiol **86**: 908–913   681

Haruta M, Burch HL, Nelson RB, Barrett-Wilt G, Kline KG, Mohsin SB,   682
Young JC, Otegui MS, Sussman MR (2010) Molecular characterization of   683
mutant Arabidopsis plants with reduced plasma membrane proton pump   684
activity. J Biol Chem **285**: 17918–29   685

Haruta M, Sussman MR (2012) The effect of a genetically reduced plasma   686
membrane protonmotive force on vegetative growth of Arabidopsis. Plant   687
Physiol **158**: 1158–71   688

Hoffmann B, Kosegarten H (1995) FITC-dextran for measuring apoplast pH and   689
apoplastic pH gradients between various cell types in sunflower leaves.   690
Physiol Plant **95**: 327–335   691

Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P,   692
Gruissem W, Zimmermann P (2008) Genevestigator V3: A Reference   693
Expression Database for the Meta-Analysis of Transcriptomes. Adv   694
Bioinformatics **2008**: 1–5   695

Huang X, Shi H, Hu Z, Liu A, Amombo E, Chen L, Fu J (2017) ABA Is   696
Involved in Regulation of Cold Stress Response in Bermudagrass. Front Plant   697
Sci **8**: 1613   698

Jia W, Davies WJ (2007) Modification of leaf apoplastic pH in relation to   699
stomatal sensitivity to root-sourced abscisic acid signals. Plant Physiol **143**:   700
68–77   701

Kaptan S, Assentoft M, Schneider HP, Fenton RA, Deitmer JW, MacAulay N,   702
De Groot BL (2015) H95 Is a pH-Dependent Gate in Aquaporin 4. Structure   703
23: 2309–2318   704

Karuppanapandian T, Geilfus C-M, Mühling K-H, Novák O, Gloser V (2017)   705
Early changes of the pH of the apoplast are different in leaves, stem and roots   706
of Vicia faba L. under declining water availability. Plant Sci **255**: 51–58   707

Kinsman EA, Pyke KA (1998) Bundle sheath cells and cell-specific plastid   708
development in Arabidopsis leaves. Development **125**: 1815–1822   709

Korovetska H, Novák O, Jůza O, Gloser V (2014) Signalling mechanisms   710
involved in the response of two varieties of Humulus lupulus L. to soil drying: 711
I. changes in xylem sap pH and the concentrations of abscisic acid and anions.   712
Plant Soil **380**: 375–387   713

Lamdan NL, Attia Z, Moran N, Moshelion M (2012) The Arabidopsis-related   714
halophyte Thellungiella halophila: boron tolerance via boron complexation   715
with metabolites? Plant Cell Environ **35**: 735–46   716

Lanz E, Gregor M, Slavík J, Kotyk A (1997) Use of FITC as a Fluorescent   717

Probe for Intracellular pH Measurement. J Fluoresc **7**: 317–319   718

Leitão L, Prista C, Moura TF, Loureiro-Dias MC, Soveral G (2012) Grapevine   719
  aquaporins: Gating of a tonoplast intrinsic protein (TIP2;1) by cytosolic pH.   720
  PLoS One. doi: 10.1371/journal.pone.0033219   721

Ludwików A, Kierzek D, Gallois P, Zeef L, Sadowski J (2009) Gene expression   722
  profiling of ozone-treated Arabidopsis abi1td insertional mutant: Protein   723
  phosphatase 2C ABI1 modulates biosynthesis ratio of ABA and ethylene.   724
  Planta **230**: 1003–1017   725

Martre P, Durand JL, Cochard H (2000) Changes in axial hydraulic   726
  conductivity along elongating leaf blades in relation to xylem maturation in   727
  tall fescue. New Phytol **146**: 235–247   728

Mósca A, de Almeida A, Wragg D, Martins A, Sabir F, Leoni S, Moura T,   729
  Prista C, Casini A, Soveral G (2018) Molecular Basis of Aquaporin-7   730
  Permeability Regulation by pH. Cells **7**: 207   731

Moshelion M, Moran N, Chaumont F (2004) Dynamic changes in the osmotic   732
  water permeability of protoplast plasma membrane. Plant Physiol **135**: 2301–   733
  2317   734

Mühling KH, Plieth C, Hansen U, Sattelmacher B (1995) Apoplastic pH of   735
  intact leaves of Vicia faba as influenced by light. J Exp Bot **46**: 377–382   736

Palmgren MG (2001) Plant plasma membrane H+ -ATPases : Powerhouse for   737
  nutrient uptake. Annu Rev Plant Physiol Plant Mol Biol **52**: 817–45   738

Pandey N, Ranjan A, Pant P, Tripathi RK, Ateek F, Pandey HP, Patre U V.,   739
  Sawant S V. (2013) CAMTA 1 regulates drought responses in Arabidopsis   740
  thaliana. BMC Genomics. doi: 10.1186/1471-2164-14-216   741

Pantin F, Monnet F, Jannaud D, Costa JM, Renaud J, Muller B, Simonneau   742
  T, Genty B (2013) The dual effect of abscisic acid on stomata. New Phytol   743
  197: 65–72   744

Paulo J. Magalhães (2004) Ratio Plus (ImageJ)   745
  https://imagej.nih.gov/ij/plugins/ratio-plus.html.   746

Pitann B, Kranz T, Mühling KH (2009a) The apoplastic pH and its significance   747
     in adaptation to salinity in maize (Zea mays L.): Comparison of fluorescence   748
     microscopy and pH-sensitive microelectrodes. Plant Sci **176**: 497–504   749

Pitann B, Schubert S, Mühling KH (2009b) Decline in leaf growth under salt   750
     stress is due to an inhibition of H^+-pumping activity and increase in   751
     apoplastic pH of maize leaves. J Plant Nutr Soil Sci **172**: 535–543   752

Prado K, Boursiac Y, Tournaire-Roux C, Monneuse J-M, Postaire O, Da Ines   753
     O, Schaffner AR, Hem S, Santoni V, Maurel C (2013) Regulation of   754
     Arabidopsis Leaf Hydraulics Involves Light-Dependent Phosphorylation of   755
     Aquaporins in Veins. Plant Cell **25**: 1029–1039   756

Sade N, Shatil-Cohen A, Attia Z, Maurel C, Boursiac Y, Kelly G, Granot D,   757
     Yaaran A, Lerner S, Moshelion M (2014) The Role of Plasma Membrane   758
     Aquaporins in Regulating the Bundle Sheath-Mesophyll Continuum and Leaf   759
     Hydraulics. PLANT Physiol **166**: 1609–1620   760

Sade N, Shatil-Cohen A, Moshelion M (2015) Bundle-sheath aquaporins play a   761
     role in controlling Arabidopsis leaf hydraulic conductivity. Plant Signal   762
     Behav **10**: e1017177   763

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell   764
     signal transduction. Annu Rev Plant Physiol Plant Mol Biol **52**: 627–658   765

Scoffoni C, Albuquerque C, Brodersen CR, Townes S V., John GP, Bartlett   766
     MK, Buckley TN, McElrone AJ, Sack L (2017) Outside-Xylem   767
     Vulnerability, Not Xylem Embolism, Controls Leaf Hydraulic Decline during   768
     Dehydration. Plant Physiol **173**: 1197–1210   769

Scoffoni C, Albuquerque C, Cochard H, Buckley TN, Fletcher LR, Caringella   770
     MA, Bartlett M, Brodersen CR, Jansen S, Mcelrone AJ (2018) The Causes   771
     of Leaf Hydraulic Vulnerability and Its Influence on Gas Exchange in   772
     Arabidopsis thaliana. **178**: 1584–1601   773

Serrano R (1988) Structure and function of proton translocafing ATPase in   774
     plasma membranes of plants and fungi. Biochim Biophys Acta - Rev   775

Biomembr **947**: 1–28 776

Shapira O, Israeli Y, Shani U, Schwartz A (2013) Salt stress aggravates boron 777
toxicity symptoms in banana leaves by impairing guttation. Plant, Cell 778
Environ **36**: 275–287 779

Shapira O, Khadka S, Israeli Y, Shani U, Schwartz A (2009) Functional 780
anatomy controls ion distribution in banana leaves: Significance of Na + 781
seclusion at the leaf margins. Plant, Cell Environ **32**: 476–485 782

Sharp RG, Davies WJ (2009) Variability among species in the apoplastic pH 783
signalling response to drying soils. J Exp Bot **60**: 4363–4370 784

Shatil-Cohen A, Attia Z, Moshelion M (2011) Bundle-sheath cell regulation of 785
xylem-mesophyll water transport via aquaporins under drought stress: a target 786
of xylem-borne ABA? Plant J **67**: 72–80 787

Shatil-Cohen A, Moshelion M (2012) Smart pipes. Plant Signal Behav **7**: 1088– 788
1091 789

Shatil-Cohen A, Sibony H, Draye X, Chaumont F, Moran N, Moshelion M 790
(2014) Measuring the osmotic water permeability coefficient (Pf) of spherical 791
cells: isolated plant protoplasts as an example. J Vis Exp e51652 792

Spanswick R (1981) Eelectrogenic ion pumps. Annu Rev Plant Physiol **32**: 67–89 793

Sze H, Li X, Palmgren M (1999) Energization of plant cell membranes by H+- 794
pumping ATPases. Regulation and biosynthesis. Plant Cell **11**: 677–690 795

Taiz L, Zeiger E (2014) Plant Physiology and Development. doi: 10.3119/0035- 796
4902-117.971.397 797

Tournaire-Roux C, Sutka M, Javot H, Gout E, Gerbeau P, Luu DT, Bligny R, 798
Maurel C (2003) Cytosolic pH regulates root water transport during anoxic 799
stress through gating of aquaporins. Nature **425**: 393–397 800

Turgeman T, Shatil-Cohen A, Moshelion M, Teper-Bamnolker P, Skory CD, 801
Lichter A, Eshel D (2016) The role of aquaporins in ph-dependent 802
germination of Rhizopus delemar spores. PLoS One **11**: 1–18 803

Tyerman SD, Niemietz CM, Bramley H (2002) Plant aquaporins: multifunctional 804

water and solute channels with expanding roles. Plant, Cell Environ **25**: 173– 194

Wallach R, Da-Costa N, Raviv M, Moshelion M (2010) Development of synchronized, autonomous, and self-regulated oscillations in transpiration rate of a whole tomato plant under water stress. J Exp Bot **61**: 3439–3449

Wang Y, Liu F, Jensen CR (2012) Comparative effects of deficit irrigation and alternate partial root-zone irrigation on xylem pH, ABA and ionic concentrations in tomatoes. J Exp Bot **63**: 1907–1917

Wang Y, Noguchi K, Ono N, Inoue S, Terashima I, Kinoshita T (2014) Overexpression of plasma membrane H+-ATPase in guard cells promotes light-induced stomatal opening and enhances plant growth. Proc Natl Acad Sci U S A **111**: 533–8

Wigoda N, Pasmanik-Chor M, Yang T, Yu L, Moshelion M, Moran N (2017) Differential gene expression and transport functionality in the bundle sheath versus mesophyll – a potential role in leaf mineral homeostasis. J Exp Bot **68**: 3179–3190

Wilkins O, Bräutigam K, Campbell MM (2010) Time of day shapes Arabidopsis drought transcriptomes. Plant J **63**: 715–727

Wilkinson S, Corlett JE, Oger L, Davies WJ (1998) Effects of xylem pH on transpiration from wild-type and flacca tomato leaves1. Plant Physiol **117**: 703–709

Wilkinson S, Davies WJ (1997) Xylem sap pH increase: A drought signal received at the apoplastic face of the guard cell that involves the suppression of saturable abscisic acid uptake by the epidermal symplast. Plant Physiol **113**: 559–573

Wu F, Shen S, Lee L, Lee S, Chan T, Lin C (2009) Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation method. Plant Methods **10**: 1–10

Wysocka-Diller JW, Helariutta Y, Fukaki H, Malamy JE, Benfey PN (2000) Molecular analysis of SCARECROW function reveals a radial patterning

mechanism common to root and shoot. Development **127**: 595–603   834

Yamauchi S, Takemiya A, Sakamoto T, Kurata T, Tsutsumi T, Kinoshita T,   835

    Shimazaki K (2016) Plasma membrane H+-ATPase1 (AHA1) plays a major   836

    role in Arabidopsis thaliana for stomatal opening in response to blue light.   837

    Plant Physiol pp.01581.2016   838

Yaneff A, Sigaut L, Gómez N, Aliaga Fandiño C, Alleva K, Pietrasanta LI,   839

    Amodeo G (2016) Loop B serine of a plasma membrane aquaporin type PIP2   840

    but not PIP1 plays a key role in pH sensing. Biochim Biophys Acta -   841

    Biomembr **1858**: 2778–2787   842

Zhang X, Wang H, Takemiya A, Song C, Kinoshita T, Shimazaki K (2004)   843

    Inhibition of blue light-dependent H 1 Pumping by abscisic acid through   844

    hydrogen peroxide-induced dephosphorylation of the plasma membrane. Plant   845

    Physiol **136**: 4150–4158   846

Main Figure Legends   856

FIGURE 1. Leaf xylem sap pH reduced by petiole-fed fusicoccin (an AHA   858

    stimulator), and increased by vanadate (an AHA inhibitor) in minor veins   859

    in detached leaves of WT (wild type) Arabidopsis. **a.** Representative images   860

    of treatment effects, with color-coded pH values calculated ratiometrically for   861

    all pixels which passed the selection criteria (Materials and methods), with   862

black masking all other pixels. i, low-K^+ control (i.e., XPS, xylem perfusion
solution). ii, XPS + 10 µM fusicoccin. iii, high-K^+ XPS (XPS + 10 mM
KNO_3). iv, high-K^+ XPS + 1 mM vanadate (Na_3VO_4). **b.** Mean (±SE) xylem
sap pH without and with fusicoccin in the indicated number of leaves
(biological repeats) from at least three independent experiments. Asterisks
denote significant differences from the respective control, using Student's
two-tailed unpaired t-test (*: $P < 0.05$, **: $P < 0.001$). Other details as in a.
c. Mean (±SE) xylem sap pH without and with vanadate. Other details as in b.

**FIGURE 2. Knockout of *AHA2* increases xylem sap pH in minor leaf veins of
Arabidopsis leaf.** *aha2* knockout lines and WT plants. The mean (±SE) xylem
sap pH in the indicated number of leaves from three independent experiments.
Different letters denote significantly different pH values (P< 0.05; ANOVA).

FIGURE 3. AHA2 expression lowers the xylem sap pH. Expression levels of
AHA2 and xylem sap pH in leaves of WT, *aha2-4* and of two independent lines
(T55, T56) of *aha2-4* complemented with *AHA2* under a bundle-sheath-specific
promotor (*SCR:AHA2*). **a.** Mean normalized (±SE) *AHA2* expression levels
obtained by qRT-PCR on whole leaf RNA (n = 5 biological repetitions, leaves);
b. Mean (±SE) xylem sap pH in the indicated number of leaves from three
independent experiments. Different letters indicate significantly different
means (*P*<.0.05; ANOVA).

**FIGURE 4. The leaf hydraulic conductance (K_{leaf}) depends on the activity of
AHA2 and the pH of the xylem perfusion solution (XPS).** Mean (±SE) K_{leaf}
(calculated by Eq. 1, Materials and Methods). **a.** K_{leaf} in detached Arabidopsis
leaves of WT, *AHA2* mutants (*aha2-4, aha2-5*) and *AHA2*-complemented
aha2-4 mutant (line T56 (*SCR:AHA2*)) plants perfused with non-buffered
AXS. Numbers are those of assayed leaves from three independent

experiments. Different letters indicate significantly different means ($P<.0.05$; ANOVA). **b.** K_{leaf} determined in intact leaves of whole Arabidopsis WT, *aha2-4* and T56 (*SCR:AHA2*) plants. Other details as in a. **c.** K_{leaf} measured in detached WT Arabidopsis leaves fed with XPS[db] buffered to the indicated pH or with high-K^+ non-buffered XPS. Other details as in a. Note the lower K_{leaf} at the alkaline pH.

FIGURE 5. The BSCs membrane osmotic water permeability coefficient (P_f) depends inversely on external pH. a. Time course (60 sec) of bundle sheath protoplasts from *SCR:GFP* plants swelling upon exposure to a hypotonic XPS[db] solution at pH 6 or 7.5. The arrow indicates onset of bath flush. **b.** Time course of the osmotic concentration change in the bath (Cout) during the hypotonic challenge (calculated as in Moshelion et al., 2004). **c.** Mean ($\pm$SE) initial P_f values of the indicated number of bundle sheath protoplasts at the different pHs from three independent experiments. The asterisk denotes a significant difference between the treatments using Student's two-tailed unpaired t-test ($P < 0.01$).

FIGURE 6. Leaf vein density does not depend on AHA2. Mean vein density ($\pm$SE) averaged over the indicated number of leaves (see details in Suppl. Fig. 10). Vein density was calculated as the total vein length divided by the scanned leaf area. Note the lack of significant difference between the three genetic lines.

FIGURE 7: K_{leaf} depends inversely on xylem sap pH. K_{leaf} as a function of the xylem-sap pH in detached leaves. Symbols: **o,** K_{leaf} data from Fig. 4a (here: on the ordinate) vs. pH observed in detached leaves (of the same plant lines) in which the BSCs modified the pH of the unbuffered XPS or AXS within the xylem depending on AHA2 presence (and activity); the pH data are from Figs. 2 and 3b (here: on the abscissa); ○, K_{leaf} of detached leaves of WT plants in which the xylem sap pH was imposed by

perfusion with pH-buffered solutions; the data are from Fig. 4c. Different letters 921
denote significantly different means (a, b: K_{leaf} values, c, d: pH values). Dotted line: a 922
sigmoid $K_{leaf}=A_2+(A_1-A_2)/(1+\exp((x-x_0)/dx))$ fitted to the seven data points 923
(orthogonal fit, OriginPro v. 2020 (9.7.0.188), Origin Lab Corporation, Northampton, 924
USA); the maximum and minimum K_{leaf} indicated in the figure are $A_1=24.5$ mmol m^{-2} s^{-1}, $A_2=12.0$ mmol m^{-2} s^{-1} (R^2, the "goodness of fit" = 0.986). Note the inverse 926
relationship between K_{leaf} and the xylem sap pH around the midpoint of $x_0 = 5.87$ 927
(arrowhead). The orange ellipse (hand drawn) describes a range of possible values of 928
pH in the immediate vicinity of the WT BSCs, which could occur if the intact AHA2 929
activity in their membranes counteracted to some extent the pH of the buffered 930
perfusate). 931

FIGURE 1. Leaf xylem sap pH reduced by petiole-fed fusicoccin (an AHA stimulator), and increased by vanadate (an AHA inhibitor) in minor veins in detached leaves of WT (wild type) Arabidopsis. **a.** Representative images of treatment effects, with color-coded pH values calculated ratiometrically for all pixels which passed the selection criteria (Materials and methods), with black masking all other pixels. i, low-K^+ control (i.e., XPS, xylem perfusion solution). ii, XPS + 10 μM fusicoccin. iii, high-K^+ XPS (XPS + 10 mM KNO_3). iv, high-K^+ XPS + 1 mM vanadate (Na_3VO_4). **b.** Mean (±SE) xylem sap pH without and with fusicoccin in the indicated number of leaves (biological repeats) from at least three independent experiments. Asterisks denote significant differences from the respective control, using Student's two-tailed unpaired t-test (*: $P < 0.05$, **: $P < 0.001$). Other details as in a. **c.** Mean (±SE) xylem sap pH without and with vanadate. Other details as in b.

Fig. 1. Grunwald, Wigoda et al, 2020

FIGURE 2. Knockout of *AHA2* increases xylem sap pH in minor leaf veins of Arabidopsis leaf. *aha2* knockout lines and WT plants. The mean (±SE) xylem sap pH in the indicated number of leaves from three independent experiments. Different letters denote significantly different pH values (P< 0.05; ANOVA).

Fig. 2. Grunwald, Wigoda et al, 2020

FIGURE 3. AHA2 expression lowers the xylem sap pH. Expression levels of *AHA2* and xylem sap pH in leaves of WT, *aha2-4* and of two independent lines (T55, T56) of *aha2-4* complemented with *AHA2* under a bundle-sheath-specific promotor (*SCR:AHA2*). **a.** Mean normalized (±SE) *AHA2* expression levels obtained by qRT-PCR on whole leaf RNA (n = 5 biological repetitions, leaves); **b.** Mean (±SE) xylem sap pH in the indicated number of leaves from three independent experiments. Different letters indicate significantly different means (*P*<.0.05; ANOVA).

Fig. 3. Grunwald, Wigoda et al, 2020

FIGURE 4. The leaf hydraulic conductance (K_{leaf}) depends on the activity of AHA2 and the pH of the xylem perfusion solution (XPS). Mean (±SE) K_{leaf} (calculated by Eq. 1, Materials and Methods). **a.** K_{leaf} in detached Arabidopsis leaves of WT, *AHA2* mutants (*aha2-4, aha2-5*) and *AHA2*-complemented *aha2-4* mutant (line T56 (*SCR:AHA2*)) plants perfused with non-buffered AXS. Numbers are those of assayed leaves from three independent experiments. Different letters indicate significantly different means (*P*<.0.05; ANOVA). **b.** K_{leaf} determined in intact leaves of whole Arabidopsis WT, *aha2-4* and T56 (*SCR:AHA2*) plants. Other details as in a. **c.** K_{leaf} measured in detached WT Arabidopsis leaves fed with XPS[db] buffered to the indicated pH or with high-K^+ non-buffered XPS. Other details as in a. Note the lower K_{leaf} at the alkaline pH.

Fig. 4. Grunwald, Wigoda et al, 2020

FIGURE 5: The BSCs membrane osmotic water permeability coefficient (P_f) depends inversely on external pH. a. Time course (60 sec) of bundle sheath protoplasts from *SCR:GFP* plants swelling upon exposure to a hypotonic XPS[db] solution at pH 6 or 7.5. The arrow indicates onset of bath flush. **b.** Time course of the osmotic concentration change in the bath (Cout) during the hypotonic challenge (calculated as in Moshelion et al., 2004). **c.** Mean (±SE) initial P_f values of the indicated number of bundle sheath protoplasts at the different pHs from three independent experiments. The asterisk denotes a significant difference between the treatments using Student's two-tailed unpaired t-test (P < 0.01).

Fig. 5. Grunwald, Wigoda et al, 2020

FIGURE 6: Leaf vein density does not depend on AHA2.
Mean vein density (±SE) averaged over the indicated number of leaves (see details in Suppl. Fig. 10). Vein density was calculated as the total vein length divided by the scanned leaf area. Note the lack of significant difference between the three genetic lines.

FIGURE 7: K_{leaf} depends inversely on xylem sap pH. K_{leaf} as a function of the xylem-sap pH in detached leaves. Symbols: **O**: K_{leaf} data from Fig. 4a (here: on the ordinate) vs. pH observed in detached leaves (of the same plant lines) in which the BSCs modified the pH of the unbuffered XPS or AXS within the xylem depending on AHA2 presence (and activity); the pH data are from Figs. 2 and 3b (here: on the abscissa). **O**: K_{leaf} of detached leaves of WT plants in which the xylem sap pH was imposed by perfusion with pH-buffered solutions; the data are from Fig. 4c. Different letters denote significantly different means (a, b: K_{leaf} values, c, d: pH values). Dotted line: a sigmoid $K_{leaf}=A_2+(A_1-A_2)/(1+\exp((x-x_0)/dx))$ fitted to the seven data points (orthogonal fit, OriginPro v. 2020 (**9.7.0.188**), Origin Lab Corporation, Northampton, USA); the maximum and minimum K_{leaf} indicated in the figure are A_1=24.5 mmol m^{-2} s^{-1}, A_2= 12.0 mmol m^{-2} s^{-1} (R^2, the "goodness of fit" = 0.986). Note the inverse relationship between K_{leaf} and the xylem sap pH around the midpoint of x_0 = 5.87 (arrowhead).

The orange ellipse (hand drawn) describes a range of possible values of pH in the immediate vicinity of the WT BSCs, which could occur if the intact AHA2 activity in their membranes counteracted to some extent the pH of the buffered perfusate).