Trait components of whole plant water use efficiency are defined by unique,

environmentally responsive genetic signatures in the model C$_4$ grass *Setaria*

Max J. Feldman[1], Patrick Z. Ellsworth[2], Noah Fahlgren[1], Malia A. Gehan[1], Asaph B.

Cousins[2] and Ivan Baxter[1,3,*]

*Corresponding author

E-mail: ivan.baxter@ars.usda.gov (IB)

[1]Donald Danforth Plant Science Center, St. Louis, Missouri, USA

[2]School of Biological Sciences, Washington State University, Pullman, Washington,

USA

[3]USDA-ARS, Plant Genetics Research Unit, St. Louis, Missouri, USA

ABSTRACT

Plant growth and water use are interrelated processes influenced by the

genetic control of both plant morphological and biochemical characteristics.

Improving plant water use efficiency (WUE) to sustain growth in different

environments is an important breeding objective that can improve crop yields and

enhance agricultural sustainability. However, genetic improvements of WUE using

traditional methods have proven difficult due to low throughput and environmental

heterogeneity encountered in field settings. To overcome these limitations the study

presented here utilizes a high-throughput phenotyping platform to quantify plant

size and water use of an interspecific *Setaria italica* x *Setaria viridis* recombinant

inbred line population at daily intervals in both well-watered and water-limited

conditions. Our findings indicate that measurements of plant size and water use in

this system are strongly correlated; therefore, a linear modeling approach was used

to partition this relationship into predicted values of plant size given water use and

deviations from this relationship at the genotype level. The resulting traits

describing plant size, water use and WUE were all heritable and responsive to soil

water availability, allowing for a genetic dissection of the components of plant WUE

under different watering treatments. Linkage mapping identified major loci

underlying two different pleiotropic components of WUE. This study indicates that

alleles controlling WUE derived from both wild and domesticated accessions of the

model C_4 species *Setaria* can be utilized to predictably modulate trait values given a

specified precipitation regime.

INTRODUCTION

Improving crop productivity while simultaneously reducing agricultural

water input is essential to ensure the security of our global food supply and protect

our diminishing fresh water resources. Agriculture is by far the greatest industrial

consumer of fresh water, largely because productivity losses related to drought

stress can decrease crop yields by greater than 50% (Boyer, 1982; Hamdy et al.,

2003). Addressing these challenges will require an integrated approach that

combines irrigation practices that minimize water loss and deployment of crop

plants with superior water use efficiency (Boutraa, 2010; Davies and Bennett, 2015;

Evans and Sadler, 2008; Gregory and George, 2011; Morison et al., 2008; Stanhill,

1986).

Plant water use efficiency (WUE) can be broadly defined as the ratio of

biomass produced to total water lost by the plant (Bacon, 2009; Blum, 2009;

Condon, 2004; Evans and Sadler, 2008; Monteith, 1993; Morison et al., 2008;

Tardieu, 2013). This complex trait is determined by many factors including

photosynthetic carbon assimilated per unit of water transpired (Condon et al., 2002;

Farquhar et al., 1989; Morison et al., 2008; Penman and Schofield, 1951; Seibt et al.,

2008), leaf architecture (Brodribb et al., 2007; Sack and Holbrook, 2006), stomata

characteristics (Franks and Farquhar, 2006; Lawson and Blatt, 2014), epidermal

wax content (Premachandra et al., 1994), canopy and root architecture (White and

Snow, 2012; Martre et al., 2001), stomatal dynamics (Blatt, 2000; Hetherington and

Woodward, 2003; Lawson et al., 2010; Flood et al., 2011; Lawson et al., 2012) ,

hydraulic transport (Edwards et al., 2012; Holloway-Phillips and Brodribb, 2011),

portion of carbon lost from respiration (Escalona et al., 2012; Tomás et al., 2014)

and partitioning of photo-assimilate (Carmo-Silva et al., 2009; Chaves, 1991). Given

that plant species (Stewart et al., 1995; Winter et al., 2005; Zegada-Lizarazu and

Iijima, 2005; Zhou et al., 2012) and ecotypes within species (Kenney et al., 2014;

Lopez et al., 2015; Nakhforoosh et al., 2016; Pater et al., 2017; Ryan et al., 2016; Xu

et al., 2009) exhibit variation in WUE it is likely that the characteristics which

determine this trait are under genetic control and have evolved in response to

different environmental conditions such as water availability (Assouline and Or,

2013; Brodribb et al., 2009; Huxman et al., 2004). Therefore, WUE is likely

influenced by both genetically encoded developmental programs and changes in

growth environments throughout the plant lifecycle (Fleury et al., 2010).

The technical challenges associated with measuring plant size and

transpiration in large structured genetic populations has historically limited

experimental efforts aimed at identifying the genetic components associated with

WUE. This is particularly difficult in field settings due to year-to-year climate

fluctuation and micro-environmental variation observed within agricultural fields.

The advent of controlled environment, high-throughput phenotyping instruments

(Chen et al., 2014; Fahlgren et al., 2015; Granier et al., 2006; Pereyra-Irujo et al.,

2012; Reuzeau et al., 2006; Sadok et al., 2007; Tisné et al., 2013; Walter et al., 2007)

alleviates many of these challenges through stringent control of climatic variables

and automated, high-resolution measurement of plant size and evapotranspiration

across large breeding populations.

Evidence from studies conducted on both crop and model plants indicate that

the traits associated with WUE are heritable and largely polygenic, although

identifying the causal locus associated with differential performance has proven

difficult in crop plants due to plant size and genome complexity (Adiredjo et al.,

2014; Aparna et al., 2015; Chen et al., 2012; Coupel-Ledru et al., 2016; Honsdorf et

al., 2014; Parent et al., 2015; Schoppach et al., 2016; Xu et al., 2009). Utilization of

model plants (C_3 annuals *Arabidopsis thaliana* and *Brachypodium distachyon*) that

possess tractable genetic and experimental properties has enabled scientists to

identify QTL that contribute to WUE (Des Marais et al., 2016; Easlon et al., 2014;

Lowry et al., 2013; Mojica et al., 2016; Vasseur et al., 2014), a few of which have

been mapped to causal genes (Ruggiero et al., 2017). Species in the genus *Setaria*

also possess many of these desirable qualities and can be used as experimental

models to identify genetic components associated with WUE in a C_4 plant that is

closely related evolutionarily to C_4 crops like maize, sorghum and bioenergy grasses

(Bennetzen et al., 2012; Brutnell et al., 2010; Huang et al., 2016; Li and Brutnell,

2011; Zhu et al., 2017). However, in order to study the diversity of resource

utilization tactics present in natural and mapping populations of *Setaria* (Saha et al.,

2016) or other C4 plant species, methods to quantify plant performance and WUE

in different environments must be developed.

The objectives of this study were to use a controlled environment high-

throughput phenotyping system to characterize the genetic architecture of plant

size, water use and WUE in an interspecific *Setaria* recombinant inbred population

(RIL) under two different watering regimes. Our findings indicate that plant size,

water use and WUE are polygenic traits which are influenced by both soil water

content and greater than 10 pleiotropic loci whose effect size changes differentially

throughout development. In addition, we identify and discuss several aspects of

experimental design that should be considered when performing high-throughput

phenomics experiments to study plant WUE.

MATERIALS AND METHODS

Plant material and growth conditions

The experiment here was first described in (Feldman et al., 2017), which

focused on plant height, and the details are repeated here in quotes for clarity. "An

interspecific *Setaria* F7 RIL population comprised of 189 genotypes (1138

individuals) was used for genetic mapping. The RIL population was generated

through a cross between the wild-type green foxtail *S. viridis* accession, A10, and the

domesticated *S. italica* foxtail millet accession, B100 (Bennetzen et al., 2012; Devos

et al., 1998; Wang et al., 1998). After a six-week stratification in moist long fiber

sphagnum moss (Luster Leaf Products Inc., USA) at 4°C, *Setaria* seeds were planted

in 10 cm diameter white pots pre-filled with ~470 cm³ of Metro-Mix 360 soil

(Hummert, USA) and 0.5 g of Osmocote Classic 14-14-14 fertilizer (Everris, USA).

After planting, seeds were given seven days to germinate in a Conviron growth

chamber with long day photoperiod (16 h day/8 h night; light intensity

230 µmol/m²/s) at 31°C day/21°C night before being loaded onto the Bellwether

Phenotyping System using a random block design. Plants were grown on the system

for 25 days under long day photoperiod (16 h day/8 h night; light intensity 500

µmol/m²/s) with the same temperature regime used during germination. Relative

humidity was maintained between 40 – 80 %. Gravimetric estimation of pot weight

was performed 2-3 times per day and water was added to maintain soil volumetric

water content at either 33% full-capacity (FC) (water-limited) or 100% FC (well-

watered) as determined by (Fahlgren et al., 2015). Prescribed soil water content

across both treatment blocks was achieved by 15 days after planting (DAP).

The volume of water transpired by individual plants at each pot weighing

was calculated as the difference between the measured pot weight and the weight of

the pre-filled pot at pot capacity (100% FC) or the difference between current pot

weight and the previous weight measurement if no water was added. At the

conclusion of each weighing, if the pot weight was below the set point, water was

added to the pot to return soil water content back to its target weight. This strategy

effectively maintains soil moisture content at a consistent level within both

treatment blocks. To evenly establish seedlings before the water limitation

treatment began, equal volumes of water (100% FC) were added to all pots for two

days after transfer to the system. At 10 DAP, a dry down phase was initiated (no

watering) to establish the water-limited treatment block (40% FC) while continuing

to maintain a soil water content of 100% FC within the well-watered treatment

block."

Image acquisition and derived measurements

RGB images of individual plants were acquired using a top-view and a side-

view cameras at four different angular rotations (0°, 90° 180°, 270°) every other day

at the Bellwether Phenotyping Facility (Fahlgren et al., 2015). Optical zoom was

adjusted throughout the experiment to ensure accurate quantification of traits

throughout plant development. The unprocessed images and the details of the

configuration settings can be found at the following download site:

(https://plantcv.danforthcenter.org/pages/data-sets/setaria_height.html).

Plant objects were extracted from images and analyzed using custom PlantCV

Python scripts specific to each camera (side-view or top-view), zoom level, and lifter

height (https://github.com/maxjfeldman/Feldman_Elsworth_Setaria_WUE_2017).

Scaling factors relating pixel area and distance to ground truth measurements

calculated by (Fahlgren et al., 2015) were used to translate pixels to relative area

(pixels/cm^2) and relative distance (pixels/cm).

Biomass estimation

At the conclusion of the experiment, 176 individual plants (91 plants from

the 100% FC and 85 from the 40% FC) were harvested to measure aboveground

biomass. Gravimetric measurement of fresh weight and saturated fresh weight were

taken directly upon tissue harvest after which plant tissue was placed into

polypropylene micro-perforated bags (PJP MarketPlace #361001), dried for three

days at 60 °C and subsequently weighed to determine dry weight biomass.

Multivariate linear regression was used to evaluate, select and calibrate a predictive

model for fresh and dry weight plant biomass.

Regressing plant fresh weight biomass as a function of side-view area,

perimeter length, height, object solidity and width indicated that each of these terms

is a significant predictor of fresh weight biomass after stepwise model selection

using Akaike's Information Criterion (AIC) (Bozdogan, 1987); multiple R2 = 0.89).

Unlike fresh weight biomass, side-view area, width, and height were the only

significant terms used for prediction of dry weight biomass after using the AIC

stepwise model selection correction procedure (multiple R^2 = 0.76). Models

containing all significant terms and their interaction achieved a greater model fit,

but they introduced artifacts at earlier developmental time points due to model

over-fitting (Fig. S1). Generally, models constructed to estimate fresh weight

biomass in the well-watered treatment block exhibited greater explanatory power

than those constructed to predict dry weight biomass or those in water-limited

treatment blocks (Fig. 1).

A minimal model containing only the most significant term (side-view area)

in both fresh and dry weight models produced a goodness of fit similar to more

complex models (fresh weight R^2 = 0.86; dry weight R^2 = 0.74). To avoid

propagation of error, values that incorporated plant fresh weight biomass were

calculated based on adjusted side-view pixel area and translated to fresh weight

biomass after analysis. Cumulative biomass values calculated on a genotype within

treatment basis were interpolated using loess smoothing (Chambers and Hastie,

1992). Plant size accumulation on a per day basis was calculated as the difference

between the loess fit values on a given day and the estimates from the previous day.

Water loss tabulation

The LemnaTec instrument at the Bellwether Phenotyping Facility provided

measurements of water use based upon the gravimetric weight of each pot before

watering, the volume of water applied, and the resulting weight after watering. On

days when the volume of water added to a pot was greater than zero, the daily

volume of water added was the sum of water volume added over a single calendar

day. On days when water was not added (e.g. during the dry down period), the

water volume was calculated as the minimum gravimetric weight of the pot on the

day in question subtracted from the minimum weight value from the previous day.

The cumulative volume of water used on a specific day was the sum of all water

used prior to that day. By fifteen days after planting (DAP), the dry down phase for

the water-limited treatment block was complete and pots containing plants lost

substantially more water than their empty pot counterparts in the well-watered

treatment block (Fig. 1). This observation indicates pot water loss cannot be

considered a proximity measure of total plant transpiration before day 15 in this

experiment (Fig. 1). Examination of the ratio between fresh weight biomass

accumulated relative to the amount of water used and mathematical prediction of

the amount of water used per day over this period suggests that the amount of

water used between day 15 and 17 can be used as an approximation of cumulative

water transpired by the plant throughout this experiment up to this point (Fig. S2).

This data and the observation that at day 17 the plants are still relatively small (less

than 8% of their maximum size on average) support the rationale of starting the

analysis on this day (Fig. 1). Volumes of water use (daily and cumulative) on a

genotype within treatment basis were estimated using loess smoothing.

Heritability and trait variance partitioning

We used the same approach as in (Feldman et al., 2017) and the details are

repeated here in quotes for clarity. "During this experiment, phenotypic values for

plant area and were calculated every other day, so the number of replicates per

treatment to calculate broad sense heritability was limited. To alleviate this

technical shortcoming, trait values for each individual were interpolated across

missing days using loess smoothing.

Variance components corresponding to broad sense heritability and total

variance explained was estimated using a mixed linear model using the R package

lme4 (Bates et al., 2015). Broad sense heritability was calculated using two

methods. Within an individual experiment, broad sense heritability on a line-

estimate basis was calculated using the following formula:

Equation 1:

$$H^2_{experiment} = \sigma^2_{genotype} / (\sigma^2_{genotype} + (\sigma^2_{genotype \times treatment} / n_{treatment}) + (\sigma^2_{residual} /$$

$$n_{replicates}))$$

in which $n_{treatment}$ is the harmonic mean of the number of treatment blocks in which

each line was observed and $n_{replicates}$ is the harmonic mean of number of replicates of

each genotype in the experiment. Heritability within treatment blocks was

calculated by fitting a linear model with genotype as the only explanatory factor

within each treatment block.

Equation 2:

$$H^2_{treatment\ block} = \sigma^2_{genotype} / \sigma^2_{total\ variance}$$

The proportion of variance attributed to genotype divided by total variance

within each treatment block is reported as broad sense heritability within treatment

(equation). Total variance explained was calculated by fitting a linear model

including factors, genotype, treatment, plot and genotype x treatment effects across

all phenotypic values in all treatments. The proportion of variance that is

incorporated into these factors divided by the total variance in the experiment is

reported as total variance explained for each factor."

QTL analysis

We used the same approach as in (Feldman et al., 2017) and the details are

repeated here in quotes for clarity. "QTL mapping was performed at each time point

within treatment blocks and on the numerical difference, relative difference and

trait ratio calculated between treatment blocks using functions encoded within the

R/qtl and funqtl package (Broman et al., 2003; Kwak et al., 2016). The functions

were called by a set of custom Python and R scripts

(https://github.com/maxjfeldman/foxy_qtl_pipeline). Two complimentary analysis

methods were utilized. First, a single QTL model genome scan using Haley-Knott

regression was performed to identify QTL exhibiting LOD score peaks greater than a

permutation based significance threshold (α = 0.05, n = 1000). Next, a stepwise

forward/backward selection procedure was used to identify an additive, multiple

QTL model based upon maximization of penalized LOD score. Both procedures were

performed at each time point, within treatment blocks and on the numerical

difference relative difference and trait ratio calculated between phenotypic values

measured in treatment blocks at each time point. QTL associated with difference or

ratio composite traits may identify loci associated with genotype by environment

interaction (Des Marais et al., 2013).

The function-valued approach described by (Kwak et al., 2016), was used to

identify QTL associated with the average (SLOD) and maximum (MLOD) score at

each locus throughout the experiment. Each genotypic mean trait within treatments

was estimated using a logistic function, and the QTL significance threshold was

determined based upon permutation-based likelihood of observing the empirical

SLOD or MLOD test statistic. Separate, independent linkage mapping analysis
performed at each time point identified a larger number of QTL locations relative to
similar function valued analysis based on the SLOD and MLOD statistics calculated
at each individual marker throughout the experimental time course.
After refinement of QTL position estimates, the significance of fit for the full multiple
QTL model was assessed using type III analysis of variance (ANOVA). The
contribution of individual loci was assessed using drop-one-term, type III ANOVA.
The absolute and relative allelic effect sizes were determined by comparing the fit of
the full model to a sub-model with one of the terms removed. All putative protein
coding genes (*Setaria viridis* genome version 1.1) found within a 1.5-logarithm of
the odds (LOD) confidence interval were reported for each QTL."
289
290   **RESULTS**
291
Measuring plant size and water use throughout the plant lifecycle
Recurrent measurement of plant size and water use was performed on
individuals of a *Setaria* recombinant inbred population (Devos et al., 1998; Devos et
al., 1998) grown at two soil water content levels at the Bellwether Phenotyping
Facility (Fahlgren et al., 2015). Images of each individual plant were captured every
other day from 7 to 33 days after sowing and plant objects were isolated and
quantified using PlantCV (Fahlgren et al., 2015; Feldman et al., 2017). Weight
estimates of fresh and dry-weight aboveground biomass were calculated using a
simple linear model featuring side-view area as the only predictor (Fig. 1, Fig. S3).
Daily plant water use was inferred through gravimetric measurement of pot
weight performed two to three times each day by the LemnaTec instrument. The
amount of water used by individual plants was calculated as the difference between
the measured weight of the pot and the weight of a pre-filled pot at a fixed point that
is proportional to its water holding capacity (100% FC) or the difference between
current weight and the previous weight measurement if no water was added. At the
conclusion of each weighing event, if pot weight was below the set point, water was
added to the pot to return it to the target weight value. This strategy effectively

Figure 1. Plant size and water use can be accurately inferred throughout a majority of the plant life cycle.
A) Significant correlations between plant fresh weight and pixel area were observed in both the well-watered and water-limited treatment blocks. B) Plants exhibited a sigmoidal growth curve, characterized by an average maximal rate of growth between 23– 26 days after planting. Green lines reflect absolute average size, whereas purple lines report on growth rate. Dark and lighter shaded lines report the well-watered and water-limited treatment blocks respectively. C) Daily water loss can be accurately measured at 17 days after planting. Dark blue and orange lines correspond to average daily water lost from pots, whereas the lines with lighter shades of similar colors report the average water loss of empty pots. The dashed black line denotes the day at which dry down within the water-limited treatment block is complete whereas the dashed red line demarks when water use can be accurately measured. D) By 17 days after planting, plants have attained less than 8% of their total biomass.

maintains soil moisture potential at a consistent level within both treatment blocks.

To evenly establish seedlings before the water limitation treatment, equal volumes

of water (100% FC) were added to all pots for two days after transfer onto the

system. At 10 days after sewing, a dry down phase was initiated (no watering) to

establish uniformity within the water-limited treatment block (40% FC) while

continuing to maintain a soil water content of 100% FC within the well-watered

treatment block.

Examination of water loss from empty pots relative to those containing

plants suggested that early in the experiment a majority of water loss was

exclusively due to evaporation from the soil surface and did not informatively

report on plant transpiration (Fig. 1) (Ge et al., 2016). Beginning the analysis at day

17 enabled us to minimize the artifacts of evaporation that dominated early in the

experiment while still capturing growth attributes over a large proportion (~92%)

of the plant growth within the experiment (Fig. 1). Another potential confounding

issue was the use of a fixed set point for the pot weight, which neglected the

increasing weight of the plant when calculating the amount of water needed to

return the pot weight to the set point during watering jobs. This decreased the

volume of water present within each pot after watering by approximately 12.5%

(well-watered) and 17.5% (water-limited) on average by the end of the experiment

(Fig. S4).

Loess smoothing was used to interpolate the values of traits on a genotype

level within each treatment block across all experimental time points (Chambers

and Hastie, 1992). Rate statistics were calculated from these loess smoothed

estimates as the difference of the trait between days. Plots illustrating the mean and

variance of each trait can be observed in FIG. S5.

Plant size and water use are correlated

Over the course of this experiment cumulative plant size and water use were

highly correlated. Correlation was tightest between 21 and 27 DAP in the well-

watered treatment block (> 0.94) and quite strong between 20 and 27 DAP in the

water-limited treatment block (> 0.87, Fig. 2). In both treatment blocks, correlations

Figure 2. Plant size and water use are tightly correlated.
Pearson Correlation Coefficient both within (blue is well-watered, orange is water-limited) and between (red is across both) treatment blocks indicates strong correlation between these two characteristics, although the correlation between the rate of plant growth and daily water use decreases as plants approach maximum size. A) Correlation between cumulative plant size and water use. B) The relationship between plant size and water use at 20, 25 and 30 days after planting. C) Correlation between the rate of plant growth and daily water use. B) The relationship between plant growth rate and daily water use at 20, 25 and 30 days after planting.

between these characters were weakest at the beginning and end of the experiment

but never dropped below 0.67. The correlation of the rate statistics associated with

these traits appeared qualitatively different. Correlation between plant growth rate

and the rate of water use was initially strong (> 0.79) but rapidly decreased at about

26 DAP as the rate of growth slowed (ultimately approaching zero) by the end of the

experiment (Fig. 2) while transpiration remained high.

We implement two numerical approaches to characterize the genetic

architecture of the relationship between these traits. The first method, which is

hereafter referred to as the water use efficiency ratio (WUE$_{ratio}$), calculated the ratio

of biomass relative to the volume of water lost from the pot. This calculation was

performed on a cumulative or daily rate basis.

Equation 3:

WUE$_{ratio}$ (pixel/mL) = plant size (pixel) / plant water lost (mL)

Values of cumulative WUE$_{ratio}$ calculated during this experiment were

comparable to other experiments where plant size and water use was measured

manually at lower throughput (25-29 grams fresh weight / Liter of water, 7-9 grams

dry weight / Liter of water). On average, the cumulative and daily rate WUE$_{ratio}$ was

greater in the water-limited treatment block than in well-watered conditions. In

principle, the WUE$_{ratio}$ should attenuate the relationship between biomass and water

use, but significant correlation was still observed between these two variables,

particularly within the rate statistic over the last week of the experiment (Fig. S6,

Fig. S7).

The high correlation between plant size and water use suggests that they

were not independent traits in this experimental setup. Therefore, as a second

approach, ordinary least squares linear regression was used to model the

relationship between plant biomass and water use. For each day of the experiment,

within treatment blocks a WUE$_{model}$ was used to predict plant size

(dependent/response variable) based upon water loss (independent/explanatory

variable) (Fig. 3). The residual of this model fit was evenly distributed around zero

Figure 3. Modeling the relationship between plant size and water use results in two traits. This approach results in predicted value of water use given size (WUE$_{fit}$) colored in dark blue and deviations from this relationship (WUE$_{residual}$) plotted in red. Plot illustrates this relationship at 30 days after planting.

across the entire distribution of the predicted values suggesting minimal bias of this

approach (Fig. S8).

Equation 4:

WUE_{fit} (pixel/mL) = plant size (pixel) ~ water lost (mL) + $WUE_{residual}$ (pixel/mL)

This approach resulted in two traits: The first was the predicted model fit

(WUE_{fit}) that described the sum of squares relationship between biomass and water

use. The residual of this model ($WUE_{residual}$) can be thought of as genotype-specific

deviation from this relationship combined with measurement error. As expected,

the correlation between the fit values derived from the WUE_{model} was highly

correlated with plant size (Fig. S9). A slight correlation between cumulative plant

biomass and the residual of the WUE_{model} was observed particularly later in the

experiment demonstrating that biomass had components that were not accounted

for by the linear model fit (Fig. S10). Varying the dependence structure/assignment

or fitting of the model using major axis regression framework (Legendre, 2014) had

little effect on downstream analysis.

Each trait (biomass, water loss, WUE_{ratio}, WUE_{fit} and $WUE_{residual}$) exhibited

high average heritability over all experimental time points within and across

treatment blocks (0.28 – 0.77) (Fig. S11). Heritability tended to achieve its

maximum value in the middle of the experiment with decreased heritability

observed at the beginning and the end of the study. Proportionally, the treatment

effect of water limitation explained the largest percentage of variance within

biomass, water loss and the WUE_{fit} although genotype and genotype x treatment

interaction also explain a substantial margin of the variance (Fig. S12). Heritability

of the rate traits was generally similar but on average 5% lower than the heritability

of the cumulative traits. In all cases, the average heritability of each trait was greater

within the well-watered treatment block relative to the value calculated in water-

limited treatment block.

The genetic architecture of plant size and water use traits

For each day of the experiment, a best fit multiple QTL model was selected

for each trait (plant size, water use, WUE$_{ratio}$, WUE$_{fit}$ and WUE$_{residual}$) and the daily

rate of change of the trait within each treatment block based upon penalized LOD

score using a standard stepwise forward/backward selection procedure (Broman et

al., 2003). This approach identified 86 (cumulative Fig. 4; Table S1) and 106 (rate

Fig. S13; Table S1) unique SNPs associated with at least one of the five traits. Many

of these uniquely identified SNP positions group into clusters of tightly linked loci

that are likely representative of a single QTL location. These local clusters of SNPs

(10 cM radius) were then condensed into the most significant marker within each

cluster to simplify comparisons of genetic architecture between traits (Fig. S13; Fig.

S14). Collapsing these SNP positions yielded 23 unique QTL locations associated

with cumulative trait values (Fig. 5) and 27 unique rate QTL locations (Table S2).

Of the 23 unique QTL identified, plant biomass contributes the largest

proportion of QTL to this set (18) followed by WUE$_{ratio}$ (12), WUE$_{fit}$ (11), WUE$_{residual}$

(10) and water lost (8) (Fig. 5, Fig. S16). Despite the fact that only one QTL location

(2@96) was common across all traits and environments, the genetic architecture

that contributes to each of these characteristics was clearly related. The strong

correlation of plant size and water loss with the predicted value of plant size given

water loss (WUE$_{fit}$) are clearly reflected within the genetic architecture associated

with these traits. Plant size, water loss and WUE$_{fit}$ all shared 8 QTL (2@96, 3@48,

5@109, 6@65, 7@34, 7@51, 7@99 and 9@34) either within the well-watered or

water-limited treatment block (Fig. 5, Fig. S16). Plant size, WUE$_{ratio}$ and deviations

from the relationship between plant size and water use (WUE$_{residual}$) shared five QTL

unique to this subset (2@11, 2@113, 5@79, 5@92, and 9@127) which enable

divergence from the fundamental relationship between plant size and water loss

(Fig. 5, Fig. S16). Several QTL were identified as being uniquely associated with

plant size (3@21, 5@119, 6@80, 9@138), WUE$_{residual}$ (2@82 3@77, 6@47) and

WUE$_{fit}$ (5@39) whereas no QTL were identified as being uniquely associated with

water loss or WUE$_{ratio}$ (Fig. 5, Fig. S16).

The genetic architecture of all five traits appears to be influenced by water

availability. All traits other than water loss exhibited QTL unique to each treatment

Figure 4. Eighty-six unique QTL locations were detected across all traits in this experiment.
Each box corresponds to an individual chromosome, where the values along the x-axis are chromosome position and values along the y-axis denote the proportion of genetic variance explained by the QTL. Each triangle represents a single QTL detected, where the color indicates the trait each QTL is associated with (green = plant size, blue = water use, orange = WUE_{ratio}, black = WUE_{fit}, red = $WUE_{residual}$). The darkness of color shading is indicative of treatment block where darker represents well-watered and lighter corresponds to the water-limited block respectively. The direction of the arrow indicates the directional effect of the B100 parental allele.

Figure 5. The genetic components that contribute to subsets of traits largely overlap.
The QTL locations identified are plotted on the x-axis and the traits are plotted on the y-axis. Colored matrix entries denote at least one significant association within this experiment. A) The genetic architecture of cumulative traits. B) The genetic loci associated with trait rate of change. C) Genetic components associated with genotype x environment traits.

block (Fig. 5, Fig. S17). Biomass, water lost, and WUE_{fit} all shared four QTL in

common across environments (2@96, 5@109, 7@99 and 9@34) where as WUE_{ratio}

and the $WUE_{residual}$ shared a single QTL (2@11) between blocks not found associated

with the other traits. Two QTL (3@48, 7@34) were found specifically within the

well-watered treatment block for all traits other than $WUE_{residual}$ whereas QTL

specific to water-limited environment identified common QTL associated with

biomass and WUE_{fit} (4@52) or WUE_{ratio} and $WUE_{residual}$ (9@87, 9@127).

The identity of QTL associated with the daily rate values suggest that the

genetic architectures were largely cognate with the QTL associated with the traits

themselves, both in identity and response to treatment. In total, 28 QTL comprised

the union of all unique QTL associated with both the trait value and the daily rate of

change calculated from the trait value. Of these QTL, 22 were common between both

the trait value and rate statistic associated with the trait, whereas five are only

found associated with the rate (1@54, 2@58, 3@4, 3@61, 8@35) and only one QTL

was uniquely associated with the cumulative trait values alone (6@47) (Fig. S18).

Genotype x environment interactions

To assess the genetic architecture of genotype x environment interactions,

mapping was performed on numerical difference, relative difference and trait ratio

between the phenotypic values observed within each treatment block. In total, 148

unique SNP locations were identified as being significantly associated with at least

one of the difference trait formulations across all standard and derived plant size

and water use traits (Table S3). Substantial overlap between these categories of

genotype x interaction traits indicates that each formulation detects similar genetic

signals (Fig. S19) although the large number SNPs found uniquely associated with

the trait ratio may indicate that some of these associations may be spurious. As such,

these QTL (trait ratio genotype x environment QTL) were removed from further

analysis. The numerical difference and relative difference traits exhibited

association with 43 and 40 unique SNP positions, which were representative of 20

and 18 QTL respectively (Table S4, Fig. S20-22).

A majority of the QTL (10/15) identified as being associated with the trait

difference between treatment blocks were also found associated with the

cumulative trait in both treatment blocks (Fig. 5). The exceptions to this were QTL

located on 3@21, 3@48, 5@39, 7@34 and 9@127 that were identified as being

significantly associated with the difference between treatment blocks but only

identified in either well-watered (3@21, 3@48, 5@39, 7@34) or water-limited

conditions (9@127). Interestingly, the QTL located on 3@48, 7@34 and 9@127

were associated with more than one trait in a single treatment block which may

indicate that these QTL impart pleiotropic phenotypic effects that were dependent

upon soil water content (Fig. 5).

The temporal genetic architecture of plant growth and water usage

In order to account for the time dependence of the traits for the five plant

traits, we used a function-valued approach based upon average log-odds score

throughout across the experiment (SLOD) for each trait (Kwak et al., 2016). This

analysis parallels the individual time point analysis, although the reduction of

complexity (fewer, higher confidence QTL) provides an opportunity for

simplification and better understanding of the major loci that influence plant WUE.

SLOD based function-valued QTL models indicate that several major QTL

(2@96, 5@109, 7@99, and 9@36) influenced both plant size and water use related

traits, although the magnitude of statistical significance attributed to each loci

varied by trait and throughout plant development (Fig. 6, Fig S23). Using the SLOD

approach, we were able to partition combinations of QTL unique to related traits

(Fig. 6). For several QTL (those around 2@96 and 5@109) the positional location at

which maximal LOD score was observed changed noticeably in a trait and

environment dependent manner either due to multiple closely linked loci or noise in

our measurements. Because the confidence intervals of the QTL generally overlap,

our reporting in this section will hereafter refer to these loci by their approximate

chromosomal location.

Both plant biomass and cumulative water use exhibited almost a complete

overlap of QTL within the well-watered treatment block, whereas plant size given

Figure 6. Significant associations identified using single marker scan functional QTL mapping.
Chromosomal position is plotted on the x-axis whereas LOD score of trait association across the genome is plotted on the y-axis. Treatment block is indicated by color intensity (darker is well-watered and lighter is water-limited). Significance thresholds (based on 1000 permutations) are plotted as dashed yellow (water-limited) and red (well-watered) lines respectively.

water use (WUE$_{fit}$) and deviation of plant size from this fundamental relationship

(WUE$_{residual}$) each exhibit a unique genetic signature (Fig 6). As observed when trait

values at individual time points were treated as independent traits, a single QTL on

2@96 is the only genetic component that was shared across all five traits. The linear

modeling approach successfully partitions out QTL associated with WUE$_{fit}$ (2@96,

7@99, 9@36) from the genetic components that contribute to deviations from the

plant size ~ water use relationship (WUE$_{residual}$; 2@96, 5@109). The QTL associated

with the WUE$_{ratio}$ (2@96, 3@52, 5@109) also likely reflects deviations from the

relationship between biomass given water loss associated with the WUE$_{residual}$.

Overall, the identity of QTL associated with each trait was largely identical between

the two treatment blocks (Fig. 6, Fig. S23) as were the QTL associated with the

values of rate statistics derived from these measurements (Fig. S24, stepwise

method; Fig. S25, scanone method).

A temporal model of the genetic architecture that influences plant water use efficiency

510   Our QTL results suggest at least two components of water use efficiency with

distinct genetic architectures. In order to compare the genetic architecture across all

traits, treatments and time points in a common framework, we analyzed how each

trait was influenced by a common set of loci. Fourteen QTL were selected based

upon their association with multiple traits, robust linkage with a single trait and/or

having differential contribution to traits across treatment blocks (Table S5) and the

proportional contribution of each locus to the additive genetic variance was

calculated using drop-one-term, type III, ANOVA performed for all experimental

traits, time points and treatment. Agglomerative hierarchical clustering of the

signed proportion of additive genetic variance explained by each locus was

performed to identify modules of traits and loci that define plant phenotypes.

Examination of scree plots of the within group sum of squares suggested that the

variance within traits could be attributed to approximately six groupings although a

majority of this variance could be captured within the largest 2-3 partitions (Fig.

S26). These partitions represented the major relationships between trait classes.

The WUE$_{ratio}$ and WUE$_{residual}$ were generally grouped separately from a larger cluster

of traits that included cumulative plant size, water use and WUE$_{fit}$ (Fig. 7). The

genetic architecture of plant water use and WUE$_{fit}$ were more related to each other

than they were to plant size, which formed the third group. The influence of water

availability on these traits was apparent from the grouping of clusters whereas the

effects of time were clear but distributed within the treatment blocks. The genetic

architecture of the WUE$_{ratio}$ in the well-watered treatment block at early time points

was more similar to the architecture of plant area than itself later in development

whereas plant area in the water-limited treatment block exhibited a genetic

architecture similar to the WUE$_{ratio}$ late at the end of the experiment.

535   Examination of the signed, proportional allelic effects within the greater fixed

QTL model indicated that QTL on 2@96, 5@109, 7@99 and 9@34 contribute

medium-to-large effects on a majority of the traits examined in both treatment

blocks (Fig. 8). The B100 allele associated with QTL on 2@96 and 9@34 both

contributed to increased plant size, water loss, WUE$_{fit}$ and WUE$_{ratio}$. The QTL on

2@96 exhibited its greatest influence in the well-watered treatment block whereas

the contribution of 9@34 was greater on average in the water-limited treatment

block. Both QTL exhibited similar temporal patterns, showing an earlier effect on

plant size and WUE$_{ratio}$ but a consistent effect across water loss. Contribution of the

B100 allele on 7@99 and 5@109 decrease plant size, water use and the WUE$_{fit}$

traits; the effect of which was greater in well-watered conditions. The magnitude of

effects contributed by QTL on 7@99 on plant size decreased through time whereas

the effects on water loss and WUE$_{fit}$ peaked after 20 days and decreases slightly

thereafter. The 5@109 locus behaves similarly with little temporal variation in

plant water use and WUE$_{fit}$. A majority of the other QTL contributed minor effects

that became more prominent in one of the two treatment blocks or at a particular

developmental time points. Inheriting the B100 allele at QTL on 2@113, 3@48,

4@52, 6@65 and 9@127 increased the values while the B100 allele at the remaining

loci (2@11, 5@79, 5@95, 7@34 and 7@53) decreased the value of the traits (Fig.

S27).

555   A majority of the QTL exhibit unidirectional effects across both the well-

556   watered and water-limited treatment blocks although the direction of the effect was

Figure 7. Agglomerative hierarchical clustering defines the relationship between plant size, water use and derived water use efficiency traits.
The additive effect size of fourteen common QTLs was calculated across all traits, treatments and developmental time points through hierarchical clustering using Ward's method. Color bars on the bottom indicate trait (green = plant size, blue = water use, orange = WUE_{ratio}, black = WUE_{fit}, red = $WUE_{residual}$), treatment block (blue = well-watered, orange = water limited), and days after planting (grey scale values where white represents the trait on day 17 and black indicates the trait on day 33).

Figure 8. Additive relative effect size of the four major pleiotropic QTL plotted throughout the course of the experiment.
A model containing fourteen QTL was fit across traits, treatment blocks and days. The developmental time point (days after planting) is indicated by the x-axis whereas the proportional additive genetic effect size of the B100 allele is plotted along the y-axis. Columns are representative of traits (green = plant size, blue = water use, orange = WUE$_{ratio}$, black = WUE$_{fit}$, red = WUE$_{residual}$) while rows correspond to individual QTL. Shading within the colors denotes treatment block (darker = well-watered, lighter = water-limited).

largely dependent on the trait (Fig. S28). The exceptions to this trend represent

short periods of experimental time at which the relative effect size is near zero

within one or both treatment blocks (Fig. 8, Fig. S27).

The proportional contribution of parental alleles towards increased trait

values varied between traits, within treatment blocks and throughout plant

development. For example, B100 alleles contributed to increased trait values for all

traits other than WUE_{ratio} in the water-limited environment and the $WUE_{residual}$

across both treatment blocks (Fig. 9). Alternatively, the contributions of the A10

alleles proportionally increased the $WUE_{residual}$ value early and then again late in

plant development relative to those inherited from the B100 parent. The influence

of A10 alleles on the WUE_{ratio} was also greater their B100 counterpart under water-

limited conditions early in plant development.

DISCUSSION

The objectives of this study were to utilize technological advances in high-

throughput phenotyping (Chen et al., 2014; Fahlgren et al., 2015; Granier et al.,

2006; Pereyra-Irujo et al., 2012; Reuzeau et al., 2006; Sadok et al., 2007; Tisné et al.,

2013; Walter et al., 2007) to characterize the genetic architecture of water use

efficiency and how this architecture responds to water-limitation in an experimental

C_4 grass model system. Although considerable efforts have been made to

characterize these processes in *Arabidopsis thaliana*, C_3 grass crops and other

species (Ruggiero et al., 2017) this represents the first study performed on an

annual C_4 grass RIL population. These efforts enabled us to identify genetic loci that

contribute to differential biomass accumulation given water use in a well-watered

and water-limited environment. Our findings suggest that the major genetic

components associated with plant size, water use and water use efficiency exhibit

pleiotropic behavior and that the magnitude of their allelic effects is dependent

upon environment and developmental stage. We used two complementary

approaches to define traits, and our analysis confirmed that the genetic architecture

was similar with both approaches. We show that the loci controlling biomass

accumulation can be roughly divided into two groups: those that control the amount

Figure 9. The proportional contribution of parental alleles to increased trait values depend upon trait, environmental water content and plant developmental stage.
Alleles derived from the B100 parent contribute a greater proportional of additive genetic variance to plant size, water use and TE model fit in both well-watered and water-limited conditions than their A10 allelic counterparts. Both the WUE ratio and TE model residual traits exhibit dynamic behavior where A10 alleles contribute either greater or close to equal proportions of additive genetic variance early and late in plant development. A) The contribution of parental alleles in the water-limited treatment block. B) The contribution of parental alleles in the water-limited treatment block.

of water used to create biomass (WUE$_{fit}$) and those that control how efficiently that

water is used (WUE$_{residual}$). The results from this study indicate that alleles from

both domesticated foxtail millet and a species representative of its wild progenitor

contribute to maximal vegetative biomass yield or water use efficiency grown in

environments with different watering regimes. In addition, we highlight aspects of

our experimental design and analysis that could be improved in future studies.

The genetic architecture of plant size, water use, water use efficiency and the

relationship between these traits

Within the A10 x B100 *Setaria* RIL population, plant size, water use and the

relationship between these two variables are unique polygenic traits whose values

are all likely influenced by greater than 10 loci. Four QTL located on 2@96, 5@106,

7@99 and 9@36 exhibit strong pleiotropic influence across this suite of traits, the

relative magnitude of each is dependent upon growth environment and

developmental time point. Despite strong correlation between plant size and water

use we successfully identified genetic architectures distinct to each trait. This was

achieved by modeling plant size as a function of water use and examining the

resulting values of the model fit (plant size given water use) and deviations from

this relationship (residual of plant size given water use). This linear modeling

approach has been used much less frequently in the literature (Lopez et al., 2015;

Nakhforoosh et al., 2016) than the more commonly used WUE$_{ratio}$ (Adiredjo et al.,

2014; Honsdorf et al., 2014; Aparna et al., 2015; Fahlgren et al., 2015; Lopez et al.,

2015). While the genetic architectures associated with the WUE$_{ratio}$ and WUE$_{residual}$

in this population are closely related (Fig. 7), WUE$_{residual}$ exhibits substantial

heritability and is less correlated with plant size than the WUE$_{ratio}$ (Fig. S6 Fig. S10),

making it a more desirable metric.

By examining the model based components of WUE with function valued

single marker scan QTL analysis that accounts for multiple hypothesis testing across

time points (Kwak et al., 2016), we were able to partition the four major pleiotropic

QTL into the genetic components on 2@96, 7@99 and 9@36, which control plant

size given water use (WUE$_{fit}$) and those on 2@96 and 5@109 that contribute to

deviations from this relationship (WUE$_{residual}$). This result suggests that QTL

associated with WUE$_{fit}$ (7@99 and 9@36) potentially control the development of

transpiring plant biomass whereas the QTL associated with the WUE$_{residual}$ and

WUE$_{ratio}$ (2@96 and 5@109) influence production of non-transpiring tissues or

biological processes not directly related to transpiration. This conclusion is in

accordance with the results of other studies performed on this population which

demonstrate that these loci are largely pleiotropic (Mauro-Herrera and Doust,

2016), although the loci on 2@96 and 5@100 substantially influence plant height

(Feldman et al., 2017) and stem biomass, whereas those on 7 and 9 are not

associated with the accumulation of stem material (Banan et al., 2017).

Our study also identified many smaller effect QTL which influence biomass,

water use and WUE traits. The B100 parental allele contributes substantial positive

(3@48, 4@52, 6@65, 9@127) and negative (7@34, 7@53) effects on all traits,

whereas QTL on 2@11, 2@113, 5@79 and 5@95 contribute either to plant

size/WUE$_{ratio}$/WUE$_{residual}$ ratio to a greater degree than on plant size/water

loss/WUE$_{fit}$.

Roughly two thirds of the QTL associated with trait plasticity as a response to

water availability (difference or relative difference between treatment blocks) were

also identified as being associated with the cumulative traits within both treatment

blocks. This observation indicates that in many cases, soil water content influences

the temporal dynamics of the allelic effects by differential progression through

developmental processes that share similar genetic components (Feldman et al.,

2017). This study identifies several QTL (3@48, 7@34 and 9@127) associated with

genotype by environment traits which also exhibit significant influence on multiple

plant traits within a single treatment block. This provides relatively strong evidence

that these QTL have pleiotropic influence on size and water use related traits in an

environment specific manner. In contrast, QTL identified only by mapping on the

difference or relative difference of the traits between each environment are largely

specific to individual traits.

Evidence from this study supports an evolutionary genetic model where the

majority of QTL associated with the measured traits exhibit conditional neutrality

across both soil water potentials examined. Although all traits other than plant size

sometimes exhibit opposite directional effects across treatment blocks, the evidence

supporting a model of antagonistic pleiotropy is weak. When identified, QTL

exhibiting opposite directional effects within individual treatment blocks were

limited to short periods of experimental time and are characterized by negligible

relative effects during these time points. The contributions of alleles from both

parental lines contribute to increased WUE irrespective of soil water potential,

suggesting that neither parent was optimized for WUE. For example, alleles from the

A10 parent contribute a greater proportion of additive genetic variance to increased

WUE during early development in both well-watered and water-limited

environments, (particularly given the $WUE_{residual}$ derivation of WUE) whereas the

alleles derived from the B100 parent have greater affect on a majority of the

measured traits throughout the time course. The contribution of alleles of both

parents to water use efficiency is expected given earlier study performed on the

same platform where parental lines showed similar WUE under water-limited

conditions (Fahlgren et al., 2015).

Considerations when measuring plant size, water use and WUE

668   As observed in many other studies (Chen et al., 2012; Fahlgren et al., 2015;

Ge et al., 2016; Golzarian et al., 2011; Honsdorf et al., 2014; Lopez et al., 2015;

Parent et al., 2015), relative plant side-view pixel area provided a robust and

accurate proximity measurement of plant biomass. Although incorporation of

additional plant architectural features can improve estimates of this relationship

(Parent et al., 2015), our results indicate that caution should be taken as to not over

fit models on ground truth data collected exclusively at the end of the experiment as

was performed in this study (Fig. S1).

676   Automated or manual gravimetric measurement of pot weight has proven to

be a reliable estimator of plant transpiration but only if the evaporative loss of

moisture from soil can be accounted for. Results presented in this study indicate

that inclusion of empty pots (or pots that contain plastic plants (Parent et al., 2015)

or fabric wicks (Halperin et al., 2017)) is an appropriate empirical method to

estimate the experimental time point at which transpiration contributes

meaningfully to total pot evapotranspiration (Coupel-Ledru et al., 2016; Lopez et al.,

2015; Pereyra-Irujo et al., 2012). Estimation of evapotranspiration after this critical

time point has been effectively used by several other groups to identify and

eliminate confounding data points collected early during similar experiments

(Vasseur et al., 2014; Coupel-Ledru et al., 2016; Ge et al., 2016). Our findings

indicate that subtraction of empty pot weight (as performed by (Pereyra-Irujo et al.,

2012; Parent et al., 2015; Coupel-Ledru et al., 2016)) may overcorrect for

evaporation at early experimental time points even after the point at which plant

transpiration contributes substantially to total pot water loss. Although not applied

during this experiment, utilization of plastic covering to shield pots from

evaporative moisture loss in combination with the approaches discussed above may

improve the ability to unambiguously quantify plant transpiration (Aparna et al.,

2015; Coupel-Ledru et al., 2016; Ellsworth et al., 2017; Granier et al., 2006; Halperin

et al., 2017; Vasseur et al., 2014). In this study, the contribution of plant biomass to

overall pot weight was not accounted for during the estimation of plant water use.

Although the contribution of plant biomass to pot weight in most experiments

performed using *Arabidopsis thaliana* is negligible (Tisné et al., 2010), plant biomass

within this *Setaria* RIL population accounted for 12-18% of total average pot water

content by the end of the experiment (Fig. S4). Our inability to account for this

growth has the undesirable effect of systematically decreasing the soil water

content of larger genotypes, although in practice this small change in soil water

potential likely has minimal impact on transpiration dynamics of the plants.

Strong correlation between plant size and water use was observed in spite of

the fact that these traits can potentially be controlled by different physiological

mechanisms. A similar trend has also been described in experiments designed to

study water use efficiency in *Arabidopsis thaliana*, apple and wheat (Lopez et al.,

2015; Nakhforoosh et al., 2016; Schoppach et al., 2016; Parent et al., 2015; Vasseur

et al., 2014). The magnitude of this correlation is likely inflated in this study due to

the large differences in size between parental lines and segregants within the A10 x

B100 RIL population. Future studies aimed at investigating the genetic basis of

water use efficiency can attenuate this correlation by selecting parental lines of

similar size and flowering times that differ in their rates of transpiration within

environments of interest.

CONCLUSIONS

This study leverages recent advances in high-throughput phenotyping and

quantitative genetics to identify the genetic loci associated with plant size, water use

and water use efficiency in an interspecific RIL population of the model C_4 grass

Setaria. Our findings indicate that these traits are highly heritable and largely

polygenic, although the effects of four major pleiotropic QTL account for a

substantial proportion of the variance observed within each trait. Contribution of

parental alleles from both the domesticated and wild progenitor lines contribute to

maximization of these characteristics. Overall, the underlying genetic architecture of

each of these processes is distinct and substantially influenced by soil water content

as well as plant developmental stage. In addition, several aspects of our

experimental design which could be improved to obtain a better understanding of

the genetic components that underlie plant size, water use and water use efficiency

in future high-throughput phenotyping studies.

REFERENCE

Adiredjo AL, Navaud O, Muños S, Langlade NB, Lamaze T, Grieu P (2014)
Genetic Control of Water Use Efficiency and Leaf Carbon Isotope
Discrimination in Sunflower (Helianthus annuus L.) Subjected to Two
Drought Scenarios. PLoS ONE **9**: e101218

Aparna K, Nepolean T, Srivastsava RK, Kholová J, Rajaram V, Kumar S, Rekha
B, Senthilvel S, Hash CT, Vadez V (2015) Quantitative trait loci associated
with constitutive traits control water use in pearl millet [*Pennisetum*
glaucum (L.) R. Br.]. Plant Biol **17**: 1073–1084

Assouline S, Or D (2013) Plant Water Use Efficiency over Geological Time –
Evolution of Leaf Stomata Configurations Affecting Plant Gas Exchange. PLoS
ONE **8**: e67757

Bacon M (2009) Water Use Efficiency in Plant Biology. John Wiley & Sons, New
York, NY

Banan D, Paul R, Feldman MJ, Holmes M, Schlake H, Baxter I, Leakey ADB
(2017) High fidelity detection of crop biomass QTL from low-cost imaging in
the field. doi: 10.1101/150144

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects
Models Using lme4. J Stat Softw. doi: 10.18637/jss.v067.i01

Bennetzen JL, Schmutz J, Wang H, Percifield R, Hawkins J, Pontaroli AC, Estep
M, Feng L, Vaughn JN, Grimwood J, et al (2012) Reference genome
sequence of the model plant Setaria. Nat Biotechnol **30**: 555–561

Blatt MR (2000) Cellular Signaling and Volume Control in Stomatal Movements in
Plants. Annu Rev Cell Dev Biol **16**: 221–241

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is
the target of crop yield improvement under drought stress. Field Crops Res
112: 119–123

Boutraa T (2010) Improvement of Water Use Efficiency in Irrigated Agriculture: A
Review. J Agron **9**: 1–8

Boyer JS (1982) Plant Productivity and Environment. Science **218**: 443–448

Bozdogan H (1987) Model selection and Akaike's Information Criterion (AIC): The
general theory and its analytical extensions. Psychometrika **52**: 345–370

Brodribb TJ, Feild TS, Jordan GJ (2007) Leaf Maximum Photosynthetic Rate and
Venation Are Linked by Hydraulics. PLANT Physiol **144**: 1890–1898

Brodribb TJ, McAdam SAM, Jordan GJ, Feild TS (2009) Evolution of stomatal
responsiveness to CO2 and optimization of water-use efficiency among land
plants. New Phytol **183**: 839–847

Broman KW, Wu H, Sen S, Churchill GA (2003) R/qtl: QTL mapping in
experimental crosses. Bioinformatics **19**: 889–890

Brutnell TP, Wang L, Swartwood K, Goldschmidt A, Jackson D, Zhu X-G, Kellogg
E, Van Eck J (2010) Setaria viridis: A Model for C4 Photosynthesis. Plant Cell
22: 2537–2544

Carmo-Silva AE, Francisco A, Powers SJ, Keys AJ, Ascensao L, Parry MAJ,
Arrabaca MC (2009) Grasses of different C4 subtypes reveal leaf traits
related to drought tolerance in their natural habitats: Changes in structure,
water potential, and amino acid content. Am J Bot **96**: 1222–1235

Chambers JM, Hastie T, eds (1992) Statistical models in S. Wadsworth &
Brooks/Cole Advanced Books & Software, Pacific Grove, Calif

Chaves MM (1991) Effects of Water Deficits on Carbon Assimilation. J Exp Bot **42**:
1–16

Chen D, Neumann K, Friedel S, Kilian B, Chen M, Altmann T, Klukas C (2014)
Dissecting the Phenotypic Components of Crop Plant Growth and Drought
Responses Based on High-Throughput Image Analysis. Plant Cell Online **26**:
4636–4655

Chen J, Chang SX, Anyia AO (2012) Quantitative trait loci for water-use efficiency
in barley (Hordeum vulgare L.) measured by carbon isotope discrimination
under rain-fed conditions on the Canadian Prairies. Theor Appl Genet **125**:
71–90

Condon AG (2004) Breeding for high water-use efficiency. J Exp Bot **55**: 2447–2460

Condon AG, Richards RA, Rebetzke GJ, Farquhar GD (2002) Improving Intrinsic
Water-Use Efficiency and Crop Yield. Crop Sci **42**: 122–131

Coupel-Ledru A, Lebon E, Christophe A, Gallo A, Gago P, Pantin F, Doligez A,
Simonneau T (2016) Reduced nighttime transpiration is a relevant breeding
target for high water-use efficiency in grapevine. Proc Natl Acad Sci **113**:
8963–8968

Davies WJ, Bennett MJ (2015) Achieving more crop per drop. Nat Plants **1**: 15118

Des Marais DL, Hernandez KM, Juenger TE (2013) Genotype-by-Environment
Interaction and Plasticity: Exploring Genomic Responses of Plants to the
Abiotic Environment. Annu Rev Ecol Evol Syst **44**: 5–29

Des Marais DL, Razzaque S, Hernandez KM, Garvin DF, Juenger TE (2016)
Quantitative trait loci associated with natural diversity in water-use
efficiency and response to soil drying in Brachypodium distachyon. Plant Sci
251: 2–11

Devos KM, Wang ZM, Beales J, Sasaki T, Gale MD (1998) Comparative genetic
maps of foxtail millet (Setaria italica) and rice (Oryza sativa). Theor Appl
Genet **96**: 63–68

Easlon HM, Nemali KS, Richards JH, Hanson DT, Juenger TE, McKay JK (2014)
The physiological basis for genetic variation in water use efficiency and
carbon isotope composition in Arabidopsis thaliana. Photosynth Res **119**:
119–129

Edwards CE, Ewers BE, McClung CR, Lou P, Weinig C (2012) Quantitative
Variation in Water-Use Efficiency across Water Regimes and Its Relationship
with Circadian, Vegetative, Reproductive, and Leaf Gas-Exchange Traits. Mol
Plant **5**: 653–668

Ellsworth PZ, Ellsworth PV, Cousins AB (2017) Relationship of leaf oxygen and
carbon isotopic composition with transpiration efficiency in the C4 grasses
Setaria viridis and Setaria italica. J Exp Bot **68**: 3513–3528

Escalona JM, TomàS M, Martorell S, Medrano H, Ribas-Carbo M, Flexas J (2012)
Carbon balance in grapevines under different soil water supply: importance
of whole plant respiration: Carbon balance in grapevine. Aust J Grape Wine
Res **18**: 308–318

Evans RG, Sadler EJ (2008) Methods and technologies to improve efficiency of
water use: INCREASING WATER USE EFFICIENCIES. Water Resour Res. doi:
10.1029/2007WR006200

Fahlgren N, Feldman M, Gehan MA, Wilson MS, Shyu C, Bryant DW, Hill ST,
McEntee CJ, Warnasooriya SN, Kumar I, et al (2015) A Versatile
Phenotyping System and Analytics Platform Reveals Diverse Temporal
Responses to Water Availability in Setaria. Mol Plant **8**: 1520–1535

Farquhar GD, Hubick KT, Condon AG, Richards RA (1989) Carbon Isotope
Fractionation and Plant Water-Use Efficiency. *In* PW Rundel, JR Ehleringer,
KA Nagy, eds, Stable Isot. Ecol. Res. Springer New York, New York, NY, pp 21–
40

Feldman MJ, Paul RE, Banan D, Barrett JF, Sebastian J, Yee M-C, Jiang H, Lipka
AE, Brutnell TP, Dinneny JR, et al (2017) Time dependent genetic analysis
links field and controlled environment phenotypes in the model C4 grass
Setaria. PLOS Genet **13**: e1006841

Fleury D, Jefferies S, Kuchel H, Langridge P (2010) Genetic and genomic tools to
improve drought tolerance in wheat. J Exp Bot **61**: 3211–3222

Flood PJ, Harbinson J, Aarts MGM (2011) Natural genetic variation in plant
photosynthesis. Trends Plant Sci **16**: 327–335

Franks PJ, Farquhar GD (2006) The Mechanical Diversity of Stomata and Its
Significance in Gas-Exchange Control. PLANT Physiol **143**: 78–87

Ge Y, Bai G, Stoerger V, Schnable JC (2016) Temporal dynamics of maize plant
growth, water use, and leaf water content using automated high throughput
RGB and hyperspectral imaging. Comput Electron Agric **127**: 625–632

Golzarian MR, Frick RA, Rajendran K, Berger B, Roy S, Tester M, Lun DS (2011)
Accurate inference of shoot biomass from high-throughput images of cereal
plants. Plant Methods **7**: 2

Granier C, Aguirrezabal L, Chenu K, Cookson SJ, Dauzat M, Hamard P, Thioux J-
J, Rolland G, Bouchier-Combaud S, Lebaudy A, et al (2006) PHENOPSIS, an

automated platform for reproducible phenotyping of plant responses to soil
water deficit in *Arabidopsis thaliana* permitted the identification of an
accession with low sensitivity to soil water deficit. New Phytol **169**: 623–635

Gregory PJ, George TS (2011) Feeding nine billion: the challenge to sustainable
crop production. J Exp Bot **62**: 5233–5239

Halperin O, Gebremedhin A, Wallach R, Moshelion M (2017) High-throughput
physiological phenotyping and screening system for the characterization of
plant-environment interactions. Plant J **89**: 839–850

Hamdy A, Ragab R, Scarascia-Mugnozza E (2003) Coping with water scarcity:
water saving and increasing water productivity. Irrig Drain **52**: 3–20

Hetherington AM, Woodward FI (2003) The role of stomata in sensing and driving
environmental change. Nature **424**: 901–908

Holloway-Phillips M-M, Brodribb TJ (2011) Contrasting hydraulic regulation in
closely related forage grasses: implications for plant water use. Funct Plant
Biol **38**: 594

Honsdorf N, March TJ, Berger B, Tester M, Pillen K (2014) High-Throughput
Phenotyping to Detect Drought Tolerance QTL in Wild Barley Introgression
Lines. PLoS ONE **9**: e97047

Huang P, Shyu C, Coelho CP, Cao Y, Brutnell TP (2016) Setaria viridis as a Model
System to Advance Millet Genetics and Genomics. Front Plant Sci. doi:
10.3389/fpls.2016.01781

Huxman TE, Smith MD, Fay PA, Knapp AK, Shaw MR, Loik ME, Smith SD, Tissue
DT, Zak JC, Weltzin JF, et al (2004) Convergence across biomes to a
common rain-use efficiency. Nature **429**: 651–654

Kenney AM, McKay JK, Richards JH, Juenger TE (2014) Direct and indirect
selection on flowering time, water-use efficiency (WUE, δ13C), and WUE
plasticity to drought in Arabidopsis thaliana. Ecol Evol **4**: 4505–4521

Kwak I-Y, Moore CR, Spalding EP, Broman KW (2016) Mapping Quantitative Trait
Loci Underlying Function-Valued Traits Using Functional Principal
Component Analysis and Multi-Trait Mapping. G3amp58
GenesGenomesGenetics **6**: 79–86

Lawson T, Blatt MR (2014) Stomatal Size, Speed, and Responsiveness Impact on
Photosynthesis and Water Use Efficiency. PLANT Physiol **164**: 1556–1570

Lawson T, von Caemmerer S, Baroli I (2010) Photosynthesis and Stomatal
Behaviour. *In* UE Lüttge, W Beyschlag, B Büdel, D Francis, eds, Prog. Bot. 72.
Springer Berlin Heidelberg, Berlin, Heidelberg, pp 265–304

Lawson T, Kramer DM, Raines CA (2012) Improving yield by exploiting
mechanisms underlying natural variation of photosynthesis. Curr Opin
Biotechnol **23**: 215–220

Legendre P (2014) lmodel2: Model II Regression. R package version 1.7-2.

Li P, Brutnell TP (2011) Setaria viridis and Setaria italica, model genetic systems
for the Panicoid grasses. J Exp Bot **62**: 3031–3037

Lopez G, Pallas B, Martinez S, Lauri P-É, Regnard J-L, Durel C-É, Costes E (2015)
Genetic Variation of Morphological Traits and Transpiration in an Apple Core
Collection under Well-Watered Conditions: Towards the Identification of
Morphotypes with High Water Use Efficiency. PLOS ONE **10**: e0145540

Lowry DB, Logan TL, Santuari L, Hardtke CS, Richards JH, DeRose-Wilson LJ,
McKay JK, Sen S, Juenger TE (2013) Expression Quantitative Trait Locus
Mapping across Water Availability Environments Reveals Contrasting
Associations with Genomic Features in Arabidopsis. Plant Cell **25**: 3266–
3279

Martre P, Cochard H, Durand J-L (2001) Hydraulic architecture and water flow in
growing grass tillers (Festuca arundinacea Schreb.). Plant Cell Environ **24**:
65–76

Mauro-Herrera M, Doust AN (2016) Development and Genetic Control of Plant
Architecture and Biomass in the Panicoid Grass, Setaria. PLOS ONE **11**:
e0151346

Mojica JP, Mullen J, Lovell JT, Monroe JG, Paul JR, Oakley CG, McKay JK (2016)
Genetics of water use physiology in locally adapted Arabidopsis thaliana.
Plant Sci **251**: 12–22

Monteith JL (1993) The exchange of water and carbon by crops in a mediterranean
climate. Irrig Sci. doi: 10.1007/BF00208401

Morison JI., Baker N., Mullineaux P., Davies W. (2008) Improving water use in
crop production. Philos Trans R Soc B Biol Sci **363**: 639–658

Nakhforoosh A, Bodewein T, Fiorani F, Bodner G (2016) Identification of Water
Use Strategies at Early Growth Stages in Durum Wheat from Shoot
Phenotyping and Physiological Measurements. Front Plant Sci. doi:
10.3389/fpls.2016.01155

Parent B, Shahinnia F, Maphosa L, Berger B, Rabie H, Chalmers K, Kovalchuk A,
Langridge P, Fleury D (2015) Combining field performance with controlled
environment plant imaging to identify the genetic control of growth and
transpiration underlying yield response to water-deficit stress in wheat. J
Exp Bot **66**: 5481–5492

Pater D, Mullen JL, McKay JK, Schroeder JI (2017) Screening for Natural Variation
in Water Use Efficiency Traits in a Diversity Set of Brassica napus L. Identifies
Candidate Variants in Photosynthetic Assimilation. Plant Cell Physiol **58**:
1700–1709

Penman H, Schofield R (1951) Some physical aspects of assimilation and
transpiration. Carbon Dioxide Fixat. Photosynth. 5:

Pereyra-Irujo GA, Gasco ED, Peirone LS, Aguirrezábal LAN (2012) GlyPh: a low-
cost platform for phenotyping plant growth and water use. Funct Plant Biol
39: 905

Premachandra GS, Hahn DT, Axtell JD, Joly RJ (1994) Epicuticular wax load and
water-use efficiency in bloomless and sparse-bloom mutants of Sorghum
bicolor L. Environ Exp Bot **34**: 293–301

Reuzeau C, Frankard V, Hatzfeld Y, Sanz A, Van Camp W, Lejeune P, De Wilde C,
Lievens K, de Wolf J, Vranken E, et al (2006) Traitmill™: a functional
genomics platform for the phenotypic analysis of cereals. Plant Genet Resour
Charact Util **4**: 20–24

Ruggiero A, Punzo P, Landi S, Costa A, Van Oosten M, Grillo S (2017) Improving
Plant Water Use Efficiency through Molecular Genetics. Horticulturae **3**: 31

Ryan AC, Dodd IC, Rothwell SA, Jones R, Tardieu F, Draye X, Davies WJ (2016)
Gravimetric phenotyping of whole plant transpiration responses to
atmospheric vapour pressure deficit identifies genotypic variation in water
use efficiency. Plant Sci **251**: 101–109

Sack L, Holbrook NM (2006) LEAF HYDRAULICS. Annu Rev Plant Biol **57**: 361–381

Sadok W, Naudin P, Boussuge B, Muller B, Welcker C, Tardieu F (2007) Leaf
growth rate per unit thermal time follows QTL-dependent daily patterns in
hundreds of maize lines under naturally fluctuating conditions. Plant Cell
Environ **30**: 135–146

Saha P, Sade N, Arzani A, Rubio Wilhelmi M del M, Coe KM, Li B, Blumwald E
(2016) Effects of abiotic stress on physiological plasticity and water use of
Setaria viridis (L.). Plant Sci **251**: 128–138

Schoppach R, Taylor JD, Majerus E, Claverie E, Baumann U, Suchecki R, Fleury
D, Sadok W (2016) High resolution mapping of traits related to whole-plant
transpiration under increasing evaporative demand in wheat. J Exp Bot **67**:
2847–2860

Seibt U, Rajabi A, Griffiths H, Berry JA (2008) Carbon isotopes and water use
efficiency: sense and sensitivity. Oecologia **155**: 441–454

Stanhill G (1986) Water Use Efficiency. Adv. Agron. Elsevier, pp 53–85

Stewart G, Turnbull M, Schmidt S, Erskine P (1995) 13C Natural Abundance in
Plant Communities Along a Rainfall Gradient: a Biological Integrator of Water
Availability. Aust J Plant Physiol **22**: 51

Tardieu F (2013) Plant response to environmental conditions: assessing potential
production, water demand, and negative effects of water deficit. Front
Physiol. doi: 10.3389/fphys.2013.00017

Tisné S, Schmalenbach I, Reymond M, Dauzat M, Pervent M, Vile D, Granier C
(2010) Keep on growing under drought: genetic and developmental bases of
the response of rosette area using a recombinant inbred line population: Leaf
development and drought stress. Plant Cell Environ **33**: 1875–1887

Tisné S, Serrand Y, Bach L, Gilbault E, Ben Ameur R, Balasse H, Voisin R,
Bouchez D, Durand-Tardif M, Guerche P, et al (2013) Phenoscope: an
automated large-scale phenotyping platform offering high spatial
homogeneity. Plant J **74**: 534–544

Tomás M, Medrano H, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J
(2014) Variability of water use efficiency in grapevines. Environ Exp Bot
103: 148–157

Vasseur F, Bontpart T, Dauzat M, Granier C, Vile D (2014) Multivariate genetic
analysis of plant responses to water deficit and high temperature revealed
contrasting adaptive strategies. J Exp Bot **65**: 6457–6469

Walter A, Scharr H, Gilmer F, Zierer R, Nagel KA, Ernst M, Wiese A, Virnich O,
Christ MM, Uhlig B, et al (2007) Dynamics of seedling growth acclimation
towards altered light conditions can be quantified via GROWSCREEN: a setup
and procedure designed for rapid optical phenotyping of different plant
species. New Phytol **174**: 447–455

Wang ZM, Devos KM, Liu CJ, Wang RQ, Gale MD (1998) Construction of RFLP-
based maps of foxtail millet, Setaria italica (L.) P. Beauv. Theor Appl Genet
96: 31–36

White TA, Snow VO (2012) A modelling analysis to identify plant traits for
990   enhanced water-use efficiency of pasture. Crop Pasture Sci **63**: 63

991   **Winter K, Aranda J, Holtum JAM** (2005) Carbon isotope composition and water-
992   use efficiency in plants with crassulacean acid metabolism. Funct Plant Biol
993   **32**: 381

994   **Xu Y, This D, Pausch RC, Vonhof WM, Coburn JR, Comstock JP, McCouch SR**
995   (2009) Leaf-level water use efficiency determined by carbon isotope
996   discrimination in rice seedlings: genetic variation associated with population
997   structure and QTL mapping. Theor Appl Genet **118**: 1065–1081

998   **Zegada-Lizarazu W, Iijima M** (2005) Deep Root Water Uptake Ability and Water
999   Use Efficiency of Pearl Millet in Comparison to Other Millet Species. Plant
Prod Sci **8**: 454–460

Zhou Y, Lambrides CJ, Kearns R, Ye C, Fukai S (2012) Water use, water use
efficiency and drought resistance among warm-season turfgrasses in shallow
soil profiles. Funct Plant Biol **39**: 116

Zhu C, Yang J, Shyu C (2017) Setaria Comes of Age: Meeting Report on the Second
International Setaria Genetics Conference. Front Plant Sci. doi:
10.3389/fpls.2017.01562

1007