Partial ablation of the orexin field induces a subnarcoleptic phenotype in a conditional mouse model of orexin neurodegeneration

Sarah Wurts Black[1,*], Jessica D. Sun[1], Pamela Santiago[1], Alex Laihsu[1], Nikki Kimura[1], Akihiro Yamanaka[2], Ross Bersot[1], and Paul S. Humphries[1]

1. Reset Therapeutics, 260 Littlefield, Suite 200, South San Francisco, CA 94080, USA

2. Research Institute of Environmental Medicine, Nagoya University, Nagoya, 464-8601, Japan

*Communicating author:

Sarah Wurts Black, Ph.D.

Reset Therapeutics

260 Littlefield Ave, Suite 200

South San Francisco, CA 94080, USA

Email: sallyb@resettherapeutics.com

Abbreviated Title: DTA partial ablation

Number of Pages: 32, Figures: 6, Tables: 1, Multimedia: 0, 3D models: 0, Extended data: 1

Word count: Abstract: 250, Statement of Significance: 120, Introduction:650, Discussion: 1498

Conflict of Interest: Dr. Black, Dr. Sun, Mr. Laihsu, Ms. Kimura, Ms. Santiago, Mr. Bersot, and Dr. Humphries have been employed by Reset Therapeutics. Dr. Yamanaka has served as a paid consultant to Reset Therapeutics.

Acknowledgements: This work was funded by Reset Therapeutics and supported in part by the Michael J. Fox Foundation (grant number 11432 to Dr. Humphries). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Michael J. Fox Foundation. We thank Dr. Thomas Scammell (Harvard University) and Dr. Takeshi Sakurai (University of Tsukuba) for the donation of the orexin/ataxin-3 mice to Jackson Labs, the staff at Neuroscience Associates for the histology, Tate York (NeuroScience Associates) for the image analysis, Dr. Kevin Donohue and Dr. Bruce O'Hara (Signal Solutions and University of Kentucky) for assistance with PiezoSleep, and Tod Steinfeld and Dr. Bruce Clapham (Reset Therapeutics) for helpful discussions.

Abstract

Narcolepsy type 1 (Na-1) and 2 (Na-2) are characterized by an inability to sustain wakefulness and are likely caused by degeneration of orexin neurons. Near complete orexin neurodegeneration depletes orexin-A from the cerebrospinal fluid and produces Na-1. The pathophysiology of Na-2 is less understood, but has been hypothesized to be due to less extensive loss of orexin neurotransmission. The *orexin-tTA; TetO diphtheria toxin A* mouse allows conditional control over the extent and timing of orexin neurodegeneration. To validate partial ablation of the orexin field as a model of Na-2, orexin cell counts and sleep/wake phenotypes (determined by piezoelectric monitoring) were correlated within individual mice after different protocols of diet-controlled neurodegeneration. Sex differences were observed in the rate of orexin cell loss. Partial ablations that began during the first 8 days of study were 14% larger than partial ablations induced during the last 8 days of study, six weeks later and prior to sacrifice of all mice, suggesting orexin cell death continued despite the resumption of conditions intended to keep orexin neurons intact. Sleep/wake of mice with 71.0% orexin cell loss, initiated at the beginning of study, resembled orexin-intact controls more than that of mice with near complete neurodegeneration. Conversely, mice with 56.6% orexin cell loss, created at the end of study, had sleep/wake phenotypes that were similar to those of mice with near complete orexin cell loss. Collectively, these results suggest that compensatory wake-promotion develops in mice that have some critical portion of their orexinergic system remaining after partial ablation.

Statement of significance

The pathophysiology of narcolepsy type 2 is poorly understood but has been hypothesized to be due, at least in part, to degeneration of a smaller proportion of the orexin neuronal field than

occurs in narcolepsy type 1. To validate a transgenic mouse model of narcolepsy type 2, we correlated the sleep/wake phenotypes of individual, male and female adult mice that received diet-induced conditional ablations of orexin neurons with their orexin cell counts. Using a translatable measure of narcolepsy sleepiness severity, we demonstrated that compensatory wake-promoting responses developed in mice concurrent with progressive orexin neurodegeneration. These results provide important details necessary for preclinical drug discovery for therapeutic areas characterized by orexin insufficiency, such as narcolepsy, Parkinson's disease, and other neurodegenerative disorders.

Introduction

The sleep disorder narcolepsy features excessive daytime sleepiness (EDS), rapid-eye-movement (REM) sleep abnormalities, disturbed nighttime sleep, and is caused by diminished orexin (also known as hypocretin) neurotransmission (Black et al., 2017a). Insufficient orexin-A levels in the cerebrospinal fluid (CSF) are thought to be due to autoimmune-related destruction of the sole population of neurons that produce orexin (Peyron et al., 2000; Thannickal et al., 2000; Bonvalet et al., 2017). Patients with narcolepsy type 1 (Na-1) have low CSF orexin-A levels, cataplexy (a sudden, reversible loss of muscle tone triggered by emotions), or both. Narcolepsy type 2 (Na-2) is based on exclusion of Na-1 criteria, and its etiology is less understood than the pathology underlying narcolepsy with cataplexy. However, a limited postmortem study suggests Na-2 can be caused by 33% orexin cell loss vs. the ~90% loss observed in patients with Na-1 (Thannickal et al., 2009). Both types of narcolepsy are diagnosed, in part, by EDS that is determined by the inability of patients to sustain long periods of wakefulness during the Maintenance of Wakefulness Test (MWT) and to fall asleep with short latency on the Multiple Sleep Latency Test (MSLT).

Two transgenic mouse models have been created that recapitulate the orexin neurodegeneration and arousal state boundary dysregulation observed in Na-1. The *orexin/ataxin-3* (Atax) mouse exhibits constitutive, postnatal loss of 80-99% of orexin neurons (Hara et al., 2001; Liu et al., 2011; Black et al., 2013; Kantor et al., 2013) and express a range of narcolepsy symptom severity (Black et al., 2017a). The *orexin-tTA; TetO diphtheria toxin A* (DTA) mouse uses the double transgenic Tet Off (TetO) system to allow conditional ablation of orexin neurons (Tabuchi et al., 2014). Administration of doxycycline (DOX) to these mice blocks the tetracycline transactivator from binding to the TetO binding site and disables the expression of the DTA neurotoxin. Thus, orexin neurodegeneration can be induced by removal of doxycycline (DOX(-)). Withdrawal of dietary DOX (100 μg/g chow) for 1.5, 3.5, 7, 14 and 28 days followed by resumption of DOX has been shown to induce 36%, 64%, 86%, 95%, and 97% orexin cell loss, respectively (Tabuchi et al., 2014). Longitudinal study by the same research group on a different set of DTA mice on continual DOX(-) has shown fragmentation of wakefulness during the dark period at 1 week DOX(-) that progressively worsened over the next 3 weeks. During this time, cataplexy began at 2 weeks DOX(-)and increased in frequency over the next 5 weeks or more. Thus, conditional control of orexin neurodegeneration in DTA mice has been proposed to model either Na-1 or Na-2, depending on the extent of the ablation. However, to validate partial ablation of the orexin field as a model of Na-2, the precision of the DOX(-) timing to reliably induce a certain percentage of orexin cell loss that corresponds to a translatable phenotype of Na-2, all within the same animal, must be determined.

Recently, a translational test of the ability to sustain long bouts of wakefulness has been developed for mice using piezoelectric monitoring of sleep/wake (Black et al., 2017b). The PiezoSleep narcolepsy screen measures the ratio of time spent in long (≥ 32 min) wake bouts to time spent in short (< 8 min) sleep bouts to yield a "wake maintenance score." Sleep/wake is

sampled during the first half of the dark period, when sustained wakefulness predominates in wild type mice, to allow narcolepsy symptom severity to be assessed.

Here, orexin cell counts and sleep/wake phenotypes were correlated within individual DTA mice after partial ablation of the orexin neuronal field and sex differences were determined. The effectiveness of the Tet Off system to maintain intact orexin neurons after partial ablations induced by DOX(-) was evaluated. The stability of the sleep/wake phenotypes over time that correspond to different orexin neurodegeneration protocols were also assessed. These results further define the parameters that are essential to the validate the DTA model beyond Na-1.

Materials and Methods

All experimental procedures were approved by the Institutional Animal Care and Use Committee at Reset Therapeutics and were conducted in accordance with the principles set forth in the *Guide for Care and Use of Laboratory Animals.*

Animals. Male and female orexin-tTA mice were bred with B6.Cg-Tg(tetO-DTA)1Gfi/J mice (both strains from Nagoya University, Nagoya, JP) to produce *orexin-tTA; TetO diphtheria toxin A* (DTA) mice that were hemizygous for both transgenes (Tabuchi et al., 2014). Male and female C57BL/6NTac were used as wild type (WT) controls in Study 1. All DTA and WT mice were bred at Taconic Biosciences, Hudson, NY. All breeders and DTA mice were given doxycycline (DOX) chow (200 μg/g chow, TD.98186, Envigo, Madison, WI), to deliver a daily dose of approximately 0.6 mg DOX, until they were switched to standard chow at the start of experimental orexin neurodegeneration. Male and female hemizygous transgenic B6.Cg-Tg(HCRT-MJD)1Stak/J mice (Atax) and wild type (WT) littermates (JAX stock #023418, Jackson Labs, Bar Harbor, ME) were used in Study 2 as comparator groups that represent a standard model of narcolepsy (Hara et al., 2001) and controls. In Study 2, animals were

individually housed at 22-25 °C, 30-70% relative humidity, and LD12:12 with *ad libitum* access to food, water and nesting material.

Study Protocols. In Study 1, the effects of shipping conditions on DOX consumption and orexin neurodegeneration were assessed. Sixty DTA mice and 8 WT mice were air shipped in single-sex, group-housed compartments (N = 4-5/compartment) from Taconic in Hudson, NY to South San Francisco, CA. The WT mice (4 males and 4 females) were shipped with standard chow. Of the DTA mice, 15 males and 15 females were shipped with DOX chow; the remaining 15 males and 15 females were switched to standard chow at the time of boxing for shipment. DTA mice were weighed immediately prior to packing and upon arrival. The percentage of weight gained or lost during transit was used as a proxy measure of food consumption. The 8 WT mice, 8 DTA mice shipped with DOX chow, and 8 DTA mice that had been switched to standard chow were sacrificed for immunohistochemistry 48 h after they were packed for shipment. Half of the DTA mice in each DOX group used for histology were female. All mice used for histology remained in their original shipping compartments until the time of sacrifice. The remainder of the DTA mice were retained for other experiments, including Study 2.

In study 2 (Figure 1), the relationship between the size and timing of partial ablation of the orexin neuronal field and sleep/wake phenotype was determined. Twenty-five DTA mice were randomly assigned into groups based on DOX diet schedule; males and females were represented across groups and individually housed upon arrival for the duration of study. Mice were either: (1) maintained on DOX chow for 7 weeks (DOX(+)7wk) to keep orexin neurons intact (N = 4), (2) switched to standard chow for 8 days and then resumed DOX(+) for 6 wk (DOX(-)8d (RD)6wk) to induce partial orexin neurodegeneration during the first study week (N = 6), (3) switched to standard chow 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d) to induce partial orexin neurodegeneration during the last study week (N = 7), or (4) maintained on standard diet without doxycycline for 7 weeks (DOX(-)7wk) to induce full orexin

neurodegeneration (N = 8). Sleep/wake was assessed by piezoelectric monitoring during Study

Weeks 4 and 7. Mice were euthanized for immunohistochemistry at the end of the Week 7

recording. Previously collected piezoelectric recordings from Atax mice (N = 31) and their WT

littermates (N = 17), aged 15.4 ± 0.3 weeks and 29.4 ± 0.3 g, served as a reference as the

standard model comparator groups, as described (Black et al., 2017b).

Immunohistochemistry. In Study 1, brains from DTA mice (8 males and 8 females) and

wild type mice (4 males and 4 females), all aged 8.3 ± 0 weeks (22.0 ± 0.7 g) were used. In

Study 2, brains from DTA mice (14 males and 11 females) all aged 15.1 ± 0.1 weeks (26.1 ± 0.9

g) were used. Mice were sacrificed with 0.1 mL Beuthanasia-D (Henry Schein, Dublin, OH) and

transcardially perfused with 0.4 % paraformaldehyde. Brains were harvested, post fixed

overnight, and shipped in phosphate buffered saline to NeuroScience Associates (Knoxville,

TN), where all histology and cell counts on tissue from DTA and WT mice were performed. The

brains were embedded together per study in a single gelatin matrix block. The multi-brain blocks

were freeze-sectioned at 35 μm with brains in the coronal plane from approximately -1.23 to -

1.91 relative to Bregma. Every 4th section was retained for the detection of orexin-A positive

neurons. After blocking in normal donkey serum (5% in 0.3% triton in PBS), sections were

incubated overnight at room temperature with goat anti-donkey orexin A antibody (SC-8070,

Santa Cruz Biotechnology, CA) at 1:2000 dilution. After washing with phosphate-buffered saline,

sections were incubated for 2 hours at room temperature with biotinylated donkey anti-goat

(Jackson Immunoresearch, West Grove, PA) at 1:500 dilution in blocking solution. After further

washes, sections were developed with the avidin-biotin-peroxidase system using

diaminobenzidine as the chromagen (Vector Laboratories, Burlingame, CA). Sections were

mounted onto slides, and anatomically matched sections between brains that represented the

anterior, mid, and posterior regions of the orexin field were identified for cell counting. Counts

Figure 1. Study 2 protocol. *Orexin/tTA; TetO diphtheria toxin A* (DTA) colony mates were randomly assigned into study groups based on targeted orexin neuronal field ablation size and timing; males and females were represented across groups. DTA mice were either (1) maintained on doxycycline for 7 weeks (DOX(+)7wk) to keep orexin neurons intact (N = 4), (2) subjected to removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk) to induce partial orexin neurodegeneration during the first study week (N = 6), (3) subjected to removal of doxycycline 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d) to induce partial orexin neurodegeneration during the last study week (N = 7), or (4) maintained without doxycycline for 7 weeks (DOX(-)7wk) to induce full orexin neurodegeneration (N = 8). Sleep/wake was measured by piezoelectric monitoring at study weeks 4 and 7. For each sleep/wake recording, mice were moved from home cage to PiezoSleep cage for 20-24 h acclimation followed by experimental sampling for 30-48 h. At the end of the recording on study week 7, mice were sacrificed for histology.

were performed bilaterally on these sections using Fiji Image J automated for thresholding and particle analysis of positively stained objects. Comparisons of manual counts vs. automated counts on a control brain indicated 96-99% agreement. Immunohistochemistry for orexin A on free-floating brain sections and manual cell counting methods for Atax mice (N = 13) and WT littermates (N = 14), aged 16.9 ± 0.1 and 28.5 ± 0.8 g, have been previously described (Black et al., 2017b).

Sleep/wake recordings. In Study 2, the sleep phenotype of DTA mice with intact, partial, and full ablations of the orexin neuronal field were determined by piezoelectric recording of gross body movement and breath rate (PiezoSleep version 2.11, Signal Solutions, Lexington, KY). The PiezoSleep apparatus consisted of individual polycarbonate cages, each of which contained a polyvinylidene difluoride piezo sensor beneath a thin plastic shield. The sensor covered the entire area of the cage floor and was connected to a single channel tube amplifier (RoMo version 1.2, Signal Solutions, Lexington, KY). Each amplifier fed into a digital acquisition device connected to a computer for data collection and sleep/wake determination as previously described (Flores et al., 2007; Donohue et al., 2008; Mang et al., 2014). Briefly, signals were amplified and filtered between 0.5 and 10 Hz and analog-to-digital converted at 120 Hz sampling rate. The frequency, amplitude, and peak spectral energy of the signal were evaluated in 2 second increments over a tapered, 8 second window to automatically calculate a sleep/wake decision statistic by simple linear discriminant analysis. Sleep was classified if signals exhibited a periodicity with a consistent, low relative amplitude at a frequency in the typical breathing range (1-4 Hz). Wakefulness was associated with high amplitude signals with variable frequency and broad spectral energy. Signal features related to these properties were extracted and combined to obtain a decision statistic for each interval. Over time, the decision statistics per increment accumulated in a bimodal distribution, and the saddle point of the distribution was determined using an adaptive threshold for the classification of sleep and wake.

Bouts of contiguous intervals of sleep or wake were defined to terminate when an interval, equal to the minimum bout length of 30 s, included less than 50% of the targeted state. This method of automated, unsupervised classification of sleep/wake by piezoelectric monitoring in mice has an accuracy of 90% vs. manual EEG scoring (Mang et al., 2014) and has been used to characterize the sleep/wake phenotype of numerous transgenic mouse lines (Sethi et al., 2015b; Sethi et al., 2015a).

Piezoelectric experiments began by loading mice into individual PiezoSleep cages that contained standard rodent bedding with food and nesting material from each mouse's home cage. Mice were permitted at least 24 h of acclimation before 42-48 h of experimental data were sampled beginning at ZT6 (14:00). Prior to experimental recording and between recordings on Study Weeks 4 and 7, mice were housed individually in standard cages with nesting material in a different room.

Experimental design, analysis and statistics. Study 1 employed a between-subjects design to test the interaction of genotype/DOX chow status (WT/ standard chow (DOX(-)), DTA mice/DOX(+), or DTA mice/DOX(-) 2d) and sex (male or female) during animal shipment on body weight and orexin cell counts.

In Study 2, sleep percentages across 24 h and histograms of the percent time spent asleep or awake in bouts from 0.5 to ≥ 64 min were calculated using SleepStats version 2.11 (Signal Solutions, Lexington, KY). Based on the circadian distribution of mean sleep and wake bout durations per h, and on the increased propensity for cataplexy during the early dark period (Black et al., 2017a), sleep and wake data from ZT12-18 were used to differentiate mice based on their ability to maintain long bouts of wakefulness at that time of day. The PiezoSleep narcolepsy screen was comprised of scatterplots of time spent in short sleep bout durations (SBD, < 8 min) vs. long wake bout durations (WBD, ≥ 32 min) during ZT12-18 and was used to evaluate the severity of narcolepsy symptoms in individual mice. The ratio of time spent in long

WBD:short SBD represented a "wake-maintenance score" per individual mouse; the lower the ratio, the greater the severity of murine narcolepsy symptoms (Black et al., 2017b). Study 2 used a between-subjects design to investigate the interaction of DOX chow status (DOX(+)7wk, DOX(-)8d (RD)6wk, DOX(+)6wk DOX(-)8d, or DOX(-)7wk) and sex on orexin cell counts and wake-maintenance scores. A between-within design was used to evaluate the interaction of DOX chow status and repeated measures of circadian time, bout durations bins, and study week recordings.

Statistics were performed using GraphPad Prism version 7.03 (La Jolla, CA). Unless otherwise noted, all data are presented as the mean ± S.E.M. Body weight changes and orexin cell counts were compared between groups using two-way analysis of variance (ANOVA) on factors "genotype/DOX chow status" and "sex." For cell count analyses with both sexes combined, one-way ANOVA was used. Sleep time across the day was compared between DTA mice on different DOX chow schedules using two-way mixed-model ANOVA on factors "DOX chow status" (between subjects) and "6-h bin" (within subjects). One-way ANOVA was used to compare the 24 h total percentages of sleep between DOX groups. Histograms representing the distribution of time spent in various sleep and wake bout durations were evaluated using two-way mixed-model ANOVA on factors "DOX chow status" (between subjects) and "bout duration bin" (within subjects). For the PiezoSleep narcolepsy screen, the relationship between time spent in long WBD and short SBD, and the ratio of these variables (long WBD:short SBD, i.e., wake-maintenance score) vs. orexin cell counts, was determined using Pearson correlation. Comparison of the long WBD:short SBD ratio between DOX groups was made using one-way ANOVA; sex differences were determined using two-way ANOVA. Pearson correlations were used to evaluate the stability of sleep measures (daily sleep amount, time in long WBD, time in short SBD, long WBD:short SBD) between Study Weeks 4 and 7. For each analysis, when ANOVA indicated significance ($\alpha = 0.05$), contrasts between relevant factor levels

(genotype/DOX chow status, sex, time bin, or bout duration bin, as appropriate) were detected

post hoc with Bonferroni's multiple comparisons test.

Results

 Histology. In study 1, to determine how 48 h animal shipping conditions might affect

DOX chow consumption and therefore orexin neurodegeneration, body weight changes during

transit and orexin cell counts were compared between groups. DTA mice that had been

switched to standard chow (DTA DOX(-)2d group) lost 6.7 ± 0.6% (males) and 2.4 ± 1.0%

(females) of their body weight during transit, while DTA mice kept on DOX chow (DOX(+) group)

lost 3.4 ± 0.8% (males) or gained 0.78 ± 1.0% (females) of their body weight. Although males

lost a greater percentage of their body weight during transit than females (main effect for "sex,"

$F(1, 56) = 25$, $P < 0.0001$) and all DTA mice in the DOX(-)2d group lost more weight than those

maintained on DOX(+) (main effect for "DOX status," $F(1, 56) = 14.5$, $P = 0.0004$), there was no

interaction of these factors ($F(1, 56) = 0.009$, *n.s.*). There was a trend for DTA mice in the DOX(-

)2d group to lose 9-13% of their orexin neurons (42-63 cells lost out of 464-485 cells) compared

to WT and DTA DOX(+) mice, respectively ($F(2, 18) = 3.45$, $P = 0.05$), and females had fewer

orexin neurons than males (Table 1).

Table 1. Orexin cell counts in wild type (WT) and *orexin/tTA; TetO diphtheria toxin A* (DTA) mice in Study 1

	N/group	WT DOX(-)	DTA DOX(+)	DTA DOX(-) 2d	ANOVA	*P*
Males	4	100 ± 8.4	111.3 ± 6.5	106.5 ± 3.5	$F(1, 18) = 9.46$	0.007
Females	4	**100 ± 5.1**	**98.5 ± 2.0**	**77.3 ± 5.6**		
Both sexes	8	100 ± 4.6	104.9 ± 4.0	91.9 ± 6.3	$F(2, 21) = 1.69$	n.s.

Data are the mean ± S.E.M. expressed as the % of WT mean per sex
DOX(+) = continuous doxycycline chow
DOX(-) = standard chow
Bold = main effect for sex; no interaction or main effects for chow condition

In Study 2, orexin cell loss was quantified in DTA mice after DOX(-) protocols that were designed to produce partial and full ablation of the orexin field as compared to DTA mice maintained on DOX(+). Differences in the size of partial ablations created during the first week vs. the last week of the study were also assessed. Representative images of the posterior hypothalamus (Figure 2A) and cell counts (Figure 2B) indicate the DOX(-) protocols induced orexin cell death as expected ($F(3, 17) = 5.54$, $P = 0.008$). DTA mice maintained on normal chow for the duration of study (DOX(-)7wk) lost 97.4% of their orexin neurons compared to DTA mice maintained on DOX(+) for 7 weeks. Partial ablation of the orexin neuronal field that was initiated during Study Week 1 (DOX(-)8d (RD)6wk group) created more cell loss (71.0%) than the same duration of DOX(-) that occurred during Study Week 7 (DOX(+)6wk DOX(-)8d group, 56.6% cell loss). Female DTA mice had fewer orexin-positive cells than males in each partial ablation group (22-46% difference), but the extent of orexin cell loss after DOX(-)7wk was similar between sexes.

Figure 2. Loss of orexin neurons in *orexin/tTA; TetO diphtheria toxin A* (DTA) mice in Study 2. Representative images of Orexin-A+ cells in the posterior lateral hypothalamus **(A)** of male and female DTA colony mates maintained on doxycycline for 7 weeks (DOX(+)7wk), with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk), with removal of doxycycline 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d), or after 7 weeks without doxycycline (DOX(-)7wk). Images taken at 10x magnification (bar = 250 μm), approximately -1.5 mm from Bregma; Fornix (f), internal capsule (ic), supraoptic decussation (sox). Orexin-A+ cell counts (mean ± S.E.M.) for each group **(B)**, expressed as the percent of values from the DOX(+)7wk group. Analysis of variance with Bonferroni's multiple comparisons: * $P < 0.05$ vs. DOX(+)7wk and DOX(-) 7wk, # $P < 0.05$ vs. DOX(-)8d (RD)6wk, + $P < 0.05$ vs. female.

Sleep/wake phenotypes. The relationship between partial orexin neurodegeneration and the quantity and architecture of sleep/wake was evaluated in DTA mice after ablation was permitted by DOX(-)8d (RD)6wk (71.0% orexin cell loss) or by DOX(+)6wk DOX(-)8d (56.6% orexin cell loss) vs. DTA mice maintained as orexin-intact on DOX(+) or with 97.4% orexin cell loss from DOX(-) for 7 weeks. The percentage of time spent asleep over 24 h or within 6-h bins across the day did not depend on the extent of orexin ablation (Figure 3), $F(3, 21) = 0.57$, *n.s.* and $F(9, 60) = 0.95$, *n.s.*, respectively. Similarly, the proportion of time spent in sleep and wake bouts ranging from 0.5 to ≥ 64 min was not particularly distinguishable between DTA mice with partial ablations of the orexin neuronal field and those that were orexin intact (Figure 4). However, there were a few minor differences in time spent in wake bouts 16-64 min long during the light period (Figure 4D, $F(21, 147) = 3.77$, $P < 0.0001$) and time spent in sleep bouts 8-32 min long during the dark period (Figure 4E, $F(21, 147) = 2.59$, $P < 0.0005$) for DTA mice maintained on DOX(-) for 8 days or 7 weeks prior to sacrifice vs. orexin-intact controls.

Figure 3. Orexin neurodegeneration does not affect the daily amount nor distribution of sleep time in *orexin/tTA; TetO diphtheria toxin A* (DTA) mice. Percent of time spent asleep (mean ± S.E.M.) per 24 h **(A)** or across the day in 6-h time bins **(B)**. Male and female DTA colony mates were maintained on doxycycline for 7 weeks (DOX(+)7wk) **(white)**, with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk) **(light gray)**, with removal of doxycycline 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d) **(dark gray)**, or after 7 weeks without doxycycline (DOX(-)7wk) **(black)**.

The severity of murine narcoleptic symptoms in DTA mice with full orexin cell loss (DOX(-)7wk) vs. those with partial ablation of the orexin field (DOX(-)8d (RD)6wk and DOX(+)6wk DOX(-)7wk) or those with intact orexin neurons (DOX(+)7wk) were compared using the PiezoSleep narcolepsy screen (Figure 5). As expected, the correlation between time spent in short SBD to long WBD was significant, and mice with full orexin cell loss tended to cluster in a manner reflective of their low ratio of time in long WBD:short SBD (Figure 5A). Plots of the wake-maintenance score (ratio of time spent in long WBD:short SBD) vs. orexin cell counts further distinguished DTA mice on the basis of their narcoleptic phenotype (Figure 5B). The DTA mice with 97.4% orexin cell loss resembled Atax mice (96.6 % cell loss) and exhibited a lower wake-maintenance score than DTA mice maintained on DOX(+) with intact orexin neurons (Figure 5C), $F(3, 21) = 5.54$, $P < 0.006$. Mice with partial ablation of the orexin field clustered intermediary to DOX(+)7wk and DOX(-)7wk mice. The wake-maintenance score was 3.8x greater in mice that received the partial ablation during Study Week 1 (DOX(-)8d (RD)6wk) than in mice with near complete orexin cell loss (DOX(-)7wk) and was similar to the ratio observed in orexin-intact mice. There were no sex differences in the severity of narcoleptic symptoms in DTA littermates, $F(3, 17) = 0.43$, *n.s.,* (Figure 5-1).

The stability of the sleep/wake phenotype over time in DTA mice, in which ablation of the orexin neuronal field began in Study Week 1, and in orexin-intact controls, was evaluated by correlative comparison of data collected during Study Weeks 4 and 7. As expected, the amount of sleep per 24 h strongly correlated between study weeks (Figure 6A), $r = 0.68$, $P = 0.002$.

Figure 4. Sleep/wake architecture in *orexin/tTA; TetO diphtheria toxin A* (DTA) mice.
Percent of time during the light **(A-D)** or dark period **(E-H)** spent asleep **(A, B, E, F)** or awake
(C, D, G, H) in different bout durations from 0.5 to ≥ 64 min. Male and female DTA colony
mates were maintained on doxycycline for 7 weeks (DOX(+)7wk) **(white circles, gray dashed
line)**, with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d
(RD)6wk) **(white triangle, gray line)**, with removal of doxycycline 8 days prior to sacrifice
(DOX(+)6wk DOX(-)8d) **(black square, black line)**, or after 7 weeks without doxycycline
(DOX(-)7wk) **(black circle, black dashed line)**. Analysis of variance with Bonferroni's multiple
comparisons: *$P < 0.05$ DOX(+)6wk DOX(-)8d vs. DOX(+)7wk; +$P < 0.05$ DOX(-)7wk vs.
DOX(+)7wk.

Measures of sleep/wake fragmentation during ZT12-18 from the narcolepsy phenotyping
screen showed more variation between Study Weeks 4 and 7 than the 24-h total sleep times.
The percentage of time spent in WBD ≥ 32 min (Figure 6B) and in SBD < 8 min (Figure 6C) also
strongly correlated between Study Weeks 4 and 7, $r = 0.63$, $P = 0.005$ and $r = 0.59$, $P = 0.01$,
respectively. However, the ratio of time spent in long WBD:short SBD only reached trend level
in the comparison between study weeks (Figure 6D), $r = 0.43$, $P = 0.08$.

Discussion

For the first time, sleep/wake phenotypes and orexin cell counts have been determined
in the same individual DTA mice after partial ablation of the orexin neuronal field and
sleep/wake measurement by piezoelectric monitoring. First, the effect of animal shipment
conditions, which may disrupt the normal consumption of DOX chow, on the integrity of the Tet
Off system was assessed. The DTA mice in transit for 48 h and maintained on DOX chow (200
µg/g chow) did not lose orexin neurons despite weight loss, and presumably reduced DOX chow
consumption. Because DOX(-)2d during transit only induced 9-13% orexin cell loss, in Study 2,
DTA mice were subjected to DOX(-) 8d to produce more substantial neurodegeneration. Partial

Figure 5. Partial ablation of the orexin field induces a sub narcoleptic phenotype as revealed in the PiezoSleep narcolepsy screen. Percent of time spent in short sleep bouts (< 8 min) vs. long wake bouts (≥ 32 min) from ZT12-18 per individual *orexin/tTA; TetO diphtheria toxin A* (DTA) mouse **(A)**. Ratio of time in long wake bouts to short sleep bouts from ZT12-18 vs. orexin cell count per individual (expressed as percent of orexin-intact control mean per transgenic line) **(B)** and mean (± S.E.M.) per doxycycline/orexin ablation group **(C)**. Male **(black bars)** and female **(white bars)** DTA colony mates were maintained on doxycycline for 7 weeks (DOX(+)7wk) **(white circles)**, with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk) **(gray triangles)**, with removal of doxycycline 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d) **(gray squares)**, or after 7 weeks without doxycycline (DOX(-)7wk) **(black circles)**. Male *orexin/ataxin-3* (Atax) mice (**white diamond**) and wild type colony mates (**gray diamond**) are plotted as group means (paired sleep and cell count data from individuals not available) for reference. Pearson correlations indicated for data from DTA mice. Two-way analysis of variance with Bonferroni's multiple comparisons, main effect for "DOX chow status": * P < 0.05 vs. DOX(-)7wk.

ablations of the orexin field that were created during the first week of study (DOX(-)8d (RD)6wk group) were 14% larger than the partial ablations induced by the same duration of DOX(-) that occurred in the last week of study (DOX(+)6wk DOX(-)8d group), suggesting orexin cell death continued despite the resumption of DOX chow for 6 weeks. Despite these larger partial ablations (71.0% orexin cell loss), the wake-maintenance scores of mice in the DOX(-)8d (RD)6wk group resembled those of orexin-intact mice more than those of mice with near complete (≥ 97%) orexin neurodegeneration. Conversely, mice with smaller partial ablations (56.6% orexin cell loss) in the DOX(+)6wk DOX(-)8d group had lower wake-maintenance scores that resembled those of mice with near complete orexin cell loss.

Collectively, these results suggest that compensatory wakefulness, perhaps through the histaminergic system (John et al., 2013; Valko et al., 2013), develops over time in mice that have some critical portion of their orexinergic system remaining after partial ablation. These

Figure 6. Stability of sleep phenotype in *orexin/tTA; TetO diphtheria toxin A* (DTA) mice over time. Correlations of data sampled on Study Week 4 vs. 7 for the percent of time spent: asleep per 24 h **(A)**, in long wake bout durations (WBD, ≥ 32 min) **(B)**, or short sleep bout durations (SBD, < 8 min) **(C)**, or for the ratio of time spent in long WBD:short SBD from ZT12-18 **(D)**. Male and female DTA colony mates were maintained on doxycycline for 7 weeks (DOX(+)7wk) **(white circles)**, with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk) **(gray triangles)**, or after 7 weeks without doxycycline (DOX(-)7wk) **(black circles)**. The group in which DOX(-)8d occurred during study week 7 were not included in the analysis because of the change in DOX status between sleep sampling time points.

compensatory changes may occur mostly within the first three weeks, because sleep/wake phenotypes remained stable within individual mice between Study Weeks 4 and 7.

An important factor relevant to the interpretation of the neurodegeneration results in the current study vs. previous work concerns the different concentrations of DOX chow employed. In Study 1, DOX(-)2d resulted in ~5x less orexin cell loss than anticipated based on the rate of neurodegeneration previously reported (Tabuchi et al., 2014). Doxycycline in the current study was provided at twice the concentration as was used previously for an estimated daily dose of 24 mg/kg. An approximately 10-fold lower daily dose of DOX, as supplied by DOX (10 µg/mL) in drinking water, has been recommended as the optimal concentration for manipulation of gene expression within 1-2 days using the TetO system due to the very slow kinetics of washout and reactivation of the transgene (Hong et al., 2007). Although the pharmacokinetics of DOX used in the current study is unknown, the dose used was ~16-fold lower than the dose achieved with DOX (2 mg/mL), which has been shown to cause residual suppression of gene expression after 8 weeks washout (Chen et al., 1998). In the current study, the smaller than expected ablation sizes and the continued neurodegeneration that was observed for weeks past the resumption of DOX may be related to the slow clearance of the DOX (200 µg/g) diet.

Sex differences were observed in the rate of orexin cell loss, but not the associated

sleep/wake phenotype, in DTA mice. Female mice in Study 1 had fewer orexin cell counts

regardless of DOX chow status, consistent with a previous report indicating male C57BL/6 mice

had 20% more orexin neurons than females throughout adulthood (Brownell and Conti, 2010).

In study 2, females with partial ablations lost a greater proportion of orexin neurons than males.

The sex differences were 22% after DOX(+)6wk DOX(-)8d and 46% after DOX(-)8d (RD)6wk,

but disappeared by DOX(-)7wk when nearly all orexin neurons had degenerated. It is unclear

whether these sex differences after partial ablations are due to variation between males and

females in doxycycline pharmacokinetics or in the efficiency of the TetO system or some other

factor. The similar wake-maintenance scores between males and females in each orexin

ablation group, despite the large differences in orexin cell loss, suggests that female DTA mice

may functionally compensate for orexin neurodegeneration more than males. In mice that lack

orexin peptides, females, unlike males, weigh more than wild type controls, but exhibit similar

metabolic markers as their male counterparts (Ramanathan and Siegel, 2014). To the best of

our knowledge, sex differences have not been reported in human CSF orexin-A levels (Nishino

et al., 2000; Nishino et al., 2009; Nakamura et al., 2011; Andlauer et al., 2012; Luca et al., 2013;

Boddum et al., 2016; Lopez et al., 2017). However, women with narcolepsy have a shorter

sleep latency, younger age of EDS onset, and yet a longer diagnostic delay compared to men

(Luca et al., 2013). A recent population-based epidemiological study has identified a higher

frequency of narcolepsy in females (Lee and Radin, 2016). Further work is needed to determine

the mechanisms that underlie the sex differences in orexin system physiology and narcolepsy.

Orexin insufficiency to varying degrees, as measured through either CSF orexin-A levels

or postmortem histology, has been associated with a variety of neurological conditions in

addition to narcolepsy and expands the clinical relevance of the DTA mouse. Orexin cell loss of

27% in Huntington's disease (Petersen et al., 2005), 23-62% in Parkinson's disease (Fronczek

et al., 2007; Thannickal et al., 2007), and 69% in multiple system atrophy (Benarroch et al., 2007) has been demonstrated despite normal levels of CSF orexin-A in these patient populations (Overeem et al., 2002; Meier et al., 2005; Yasui et al., 2006; Abdo et al., 2008). This discrepancy suggests that the residual orexin neurons compensate for reduced orexin connectivity with increased peptide release. Compensation has been observed in rats with hypothalamic orexin-B-saporin lesions; 73% orexin cell loss resulted in only 50% loss of CSF orexin-A (Gerashchenko et al., 2003). If this relationship can be extrapolated to DTA mice, then it can be speculated in the current study that mice with partial ablations may have had CSF orexin-A levels that were 40-50% less than those in orexin-intact controls. DTA mice with ≥ 97% orexin neurodegeneration may have had ≥ 66% loss of CSF orexin-A. This level of remaining CSF orexin-A (~1/3 of orexin-intact controls) is very close to the cutoff value (30% of controls or 110 pg/mL) that is most predictive of narcolepsy with cataplexy in the human diagnostic test (Mignot, 2017). Future research to test these relationships between orexin cell loss and CSF orexin-A levels, and thereby increase the translational value of the partially ablated DTA mouse, will require the development of a highly sensitive assay to detect orexin-A levels in small volume (10 µL) CSF samples from individual mice, as current assays require pooling samples to obtain 5x that volume (Liu et al., 2011).

Narcolepsy type 2 is a heterogeneous disease entity with uncertain pathophysiology that makes clinical diagnosis and animal modeling challenging. Diagnosis of Na-2 requires: EDS, a positive score on the MSLT (sleep latency ≤ 8 min, and ≥ 2 sleep onset REM periods), an absence of cataplexy, and differentiation from idiopathic hypersomnia by signs of REM sleep boundary dysregulation (Baumann et al., 2014). The majority of patients with Na-2 have normal CSF orexin-A levels (Mignot, 2017), but 8% of patients with narcolepsy without cataplexy were found to have intermediary CSF orexin-A levels that were above the Na-1 cutoff value and below the normal level of ≥ 200 pg/mL (Andlauer et al., 2012). Of these patients with

intermediary orexin-A levels, 18% developed cataplexy years after their original narcolepsy-without-cataplexy diagnosis (Andlauer et al., 2012). These data, together with the intermediary number of orexin cells found postmortem from one patient with narcolepsy without cataplexy (Thannickal et al., 2009), support the hypothesis that at least some cases of Na-2 lie on a continuum between Na-1 and idiopathic hypersomnia. The current study represents an attempt to model Na-2 based on this hypothesis that partial disruption of orexinergic signaling results in pathological sleepiness associated with less severe narcolepsy. However, the current study is limited by the lack of cataplexy and REM sleep measurements. Cataplexy assessment would be necessary to fully differentiate whether the phenotypes observed in DTA mice with partial ablations of the orexin field translate as Na-1 or Na-2. REM sleep timing may further distinguish subtle differences between DTA mice with different sizes of partial ablations.

In conclusion, a subnarcoleptic phenotype was modeled in DTA mice to the extent that a range of wake-maintenance scores were found to be associated with different sizes and durations of orexin cell loss. The relationship between partial orexin neurodegeneration and sleep/wake phenotype depends on the dynamics of progressive orexin cell loss and the development of a compensatory wake response. Further work is recommended to titrate the DOX(-) RD protocol for the induction of stable 80-90% orexin neurodegeneration. This amount of orexin cell loss is predicted to distinguish partially ablated DTA mice from orexin-intact controls without permitting cataplexy. The establishment of a mouse model based on partial ablation of orexin neurons will be critical for efficacy tests of orexin modulator therapies for narcolepsy and other diseases characterized by orexin insufficiency.

References

Abdo WF, Bloem BR, Kremer HP, Lammers GJ, Verbeek MM, Overeem S (2008) CSF hypocretin-1 levels are normal in multiple-system atrophy. Parkinsonism Relat Disord 14:342-344.

Andlauer O, Moore Ht, Hong SC, Dauvilliers Y, Kanbayashi T, Nishino S, Han F, Silber MH, Rico T, Einen M, Kornum BR, Jennum P, Knudsen S, Nevsimalova S, Poli F, Plazzi G, Mignot E (2012) Predictors of hypocretin (orexin) deficiency in narcolepsy without cataplexy. Sleep 35:1247-1255f.

Baumann CR, Mignot E, Lammers GJ, Overeem S, Arnulf I, Rye D, Dauvilliers Y, Honda M, Owens JA, Plazzi G, Scammell TE (2014) Challenges in diagnosing narcolepsy without cataplexy: a consensus statement. Sleep 37:1035-1042.

Benarroch EE, Schmeichel AM, Sandroni P, Low PA, Parisi JE (2007) Involvement of hypocretin neurons in multiple system atrophy. Acta Neuropathol 113:75-80.

Black SW, Yamanaka A, Kilduff TS (2017a) Challenges in the development of therapeutics for narcolepsy. Progress in neurobiology 152:89-113.

Black SW, Morairty SR, Fisher SP, Chen TM, Warrier DR, Kilduff TS (2013) Almorexant promotes sleep and exacerbates cataplexy in a murine model of narcolepsy. Sleep 36:325-336.

Black SW, Sun JD, Laihsu A, Kimura N, Santiago P, Donohue KD, O'Hara BF, Bersot R, Humphries PS (2017b) Rapid, noninvasive, and unsupervised detection of a sleep/wake using piezoelectric monitoring for pharmacological studies in narcoleptic mice. Submitted, https://doi.org/10.1101/226522.

Boddum K, Hansen MH, Jennum PJ, Kornum BR (2016) Cerebrospinal Fluid Hypocretin-1 (Orexin-A) Level Fluctuates with Season and Correlates with Day Length. PloS one 11:e0151288.

Bonvalet M, Ollila HM, Ambati A, Mignot E (2017) Autoimmunity in narcolepsy. Current opinion in pulmonary medicine 23:522-529.

Brownell SE, Conti B (2010) Age- and gender-specific changes of hypocretin immunopositive neurons in C57Bl/6 mice. Neuroscience letters 472:29-32.

Chen J, Kelz MB, Zeng G, Sakai N, Steffen C, Shockett PE, Picciotto MR, Duman RS, Nestler EJ (1998) Transgenic animals with inducible, targeted gene expression in brain. Molecular pharmacology 54:495-503.

Donohue KD, Medonza DC, Crane ER, O'Hara BF (2008) Assessment of a non-invasive high-throughput classifier for behaviours associated with sleep and wake in mice. Biomed Eng Online 7:14.

Flores AE, Flores JE, Deshpande H, Picazo JA, Xie XS, Franken P, Heller HC, Grahn DA, O'Hara BF (2007) Pattern recognition of sleep in rodents using piezoelectric signals generated by gross body movements. IEEE Trans Biomed Eng 54:225-233.

Fronczek R, Overeem S, Lee SY, Hegeman IM, van Pelt J, van Duinen SG, Lammers GJ, Swaab DF (2007) Hypocretin (orexin) loss in Parkinson's disease. Brain : a journal of neurology 130:1577-1585.

Gerashchenko D, Murillo-Rodriguez E, Lin L, Xu M, Hallett L, Nishino S, Mignot E, Shiromani PJ (2003) Relationship between CSF hypocretin levels and hypocretin neuronal loss. Experimental neurology 184:1010-1016.

Hara J, Beuckmann CT, Nambu T, Willie JT, Chemelli RM, Sinton CM, Sugiyama F, Yagami K, Goto K, Yanagisawa M, Sakurai T (2001) Genetic ablation of orexin neurons in mice results in narcolepsy, hypophagia, and obesity. Neuron 30:345-354.

Hong HK, Chong JL, Song W, Song EJ, Jyawook AA, Schook AC, Ko CH, Takahashi JS (2007) Inducible and reversible Clock gene expression in brain using the tTA system for the study of circadian behavior. PLoS Genet 3:e33.

John J, Thannickal TC, McGregor R, Ramanathan L, Ohtsu H, Nishino S, Sakai N, Yamanaka A, Stone C, Cornford M, Siegel JM (2013) Greatly increased numbers of histamine cells in human narcolepsy with cataplexy. Annals of neurology 74:786-793.

Kantor S, Mochizuki T, Lops SN, Ko B, Clain E, Clark E, Yamamoto M, Scammell TE (2013) Orexin gene therapy restores the timing and maintenance of wakefulness in narcoleptic mice. Sleep 36:1129-1138.

Lee RU, Radin JM (2016) A population-based epidemiologic study of adult-onset narcolepsy incidence and associated risk factors, 2004-2013. Journal of the neurological sciences 370:29-34.

Liu M, Blanco-Centurion C, Konadhode R, Begum S, Pelluru D, Gerashchenko D, Sakurai T, Yanagisawa M, van den Pol AN, Shiromani PJ (2011) Orexin gene transfer into zona incerta neurons suppresses muscle paralysis in narcoleptic mice. The Journal of neuroscience : the official journal of the Society for Neuroscience 31:6028-6040.

Lopez R, Barateau L, Evangelista E, Chenini S, Robert P, Jaussent I, Dauvilliers Y (2017) Temporal Changes in the Cerebrospinal Fluid Level of Hypocretin-1 and Histamine in Narcolepsy. Sleep 40.

Luca G et al. (2013) Clinical, polysomnographic and genome-wide association analyses of narcolepsy with cataplexy: a European Narcolepsy Network study. Journal of sleep research 22:482-495.

Mang GM, Nicod J, Emmenegger Y, Donohue KD, O'Hara BF, Franken P (2014) Evaluation of a piezoelectric system as an alternative to electroencephalogram/ electromyogram recordings in mouse sleep studies. Sleep 37:1383-1392.

Meier A, Mollenhauer B, Cohrs S, Rodenbeck A, Jordan W, Meller J, Otto M (2005) Normal hypocretin-1 (orexin-A) levels in the cerebrospinal fluid of patients with Huntington's disease. Brain research 1063:201-203.

Mignot E (2017) Narcolepsy: Genetics, immunology, and pathophysiology. In: Principles and

    Practice of Sleep Medicine, 6th Edition (Kryger MH, Roth T, eds), pp 855-872.

    Philadelphia, Pennsylvania: Elsevier.

Nakamura M, Kanbayashi T, Sugiura T, Inoue Y (2011) Relationship between clinical

    characteristics of narcolepsy and CSF orexin-A levels. Journal of sleep research 20:45-

    49.

Nishino S, Ripley B, Overeem S, Lammers GJ, Mignot E (2000) Hypocretin (orexin) deficiency

    in human narcolepsy. Lancet 355:39-40.

Nishino S, Sakurai E, Nevsimalova S, Yoshida Y, Watanabe T, Yanai K, Mignot E (2009)

    Decreased CSF histamine in narcolepsy with and without low CSF hypocretin-1 in

    comparison to healthy controls. Sleep 32:175-180.

Overeem S, van Hilten JJ, Ripley B, Mignot E, Nishino S, Lammers GJ (2002) Normal

    hypocretin-1 levels in Parkinson's disease patients with excessive daytime sleepiness.

    Neurology 58:498-499.

Petersen A, Gil J, Maat-Schieman ML, Bjorkqvist M, Tanila H, Araujo IM, Smith R, Popovic N,

    Wierup N, Norlen P, Li JY, Roos RA, Sundler F, Mulder H, Brundin P (2005) Orexin loss

    in Huntington's disease. Human molecular genetics 14:39-47.

Peyron C et al. (2000) A mutation in a case of early onset narcolepsy and a generalized

    absence of hypocretin peptides in human narcoleptic brains. Nature medicine 6:991-997.

Ramanathan L, Siegel JM (2014) Gender differences between hypocretin/orexin knockout and

    wild type mice: age, body weight, body composition, metabolic markers, leptin and

    insulin resistance. Journal of neurochemistry 131:615-624.

Sethi M, Joshi SS, Webb RL, Beckett TL, Donohue KD, Murphy MP, O'Hara BF, Duncan MJ

    (2015a) Increased fragmentation of sleep-wake cycles in the 5XFAD mouse model of

    Alzheimer's disease. Neuroscience 290:80-89.

Sethi M, Joshi SS, Striz M, Cole N, Ryan J, Lhamon ME, Agarwal A, R. S, J. S, Denegre JM, Braun RE, Fardo DW, Donohue KD, Chesler EJ, Svenson KL, O'Hara BF (2015b) Analysis of sleep traits in knockout mice from the large-scale KOMP2 population using a non-invasive, high-throughput piezoelectric system. BMC Bioinformatics 16:P15.

Tabuchi S, Tsunematsu T, Black SW, Tominaga M, Maruyama M, Takagi K, Minokoshi Y, Sakurai T, Kilduff TS, Yamanaka A (2014) Conditional ablation of orexin/hypocretin neurons: a new mouse model for the study of narcolepsy and orexin system function. The Journal of neuroscience : the official journal of the Society for Neuroscience 34:6495-6509.

Thannickal T, Moore R, Y., Nienhuis R, Ramanathan L, Gulyani S, Aldrich M, Cornford M, Siegel JM (2000) Reduced number of hypocretin neurons in human narcolepsy. Neuron 27:469-474.

Thannickal TC, Lai YY, Siegel JM (2007) Hypocretin (orexin) cell loss in Parkinson's disease. Brain : a journal of neurology 130:1586-1595.

Thannickal TC, Nienhuis R, Siegel JM (2009) Localized loss of hypocretin (orexin) cells in narcolepsy without cataplexy. Sleep 32:993-998.

Valko PO, Gavrilov YV, Yamamoto M, Reddy H, Haybaeck J, Mignot E, Baumann CR, Scammell TE (2013) Increase of histaminergic tuberomammillary neurons in narcolepsy. Annals of neurology 74:794-804.

Yasui K, Inoue Y, Kanbayashi T, Nomura T, Kusumi M, Nakashima K (2006) CSF orexin levels of Parkinson's disease, dementia with Lewy bodies, progressive supranuclear palsy and corticobasal degeneration. Journal of the neurological sciences 250:120-123.

Extended Data

Figure 5-1. Narcoleptic phenotypes in male vs. female *orexin/tTA; TetO diphtheria toxin A* (DTA) mice after partial, full, or no ablation of the orexin neuronal field. Ratio of time spent in long wake bout durations (WBD) to short sleep bout durations (SBD) from ZT12-18 vs. orexin cell count per individual (expressed as percent of orexin-intact control mean per sex). Male **(triangles)** and female **(circles)** DTA colony mates were maintained on doxycycline for 7 weeks (DOX(+)7wk) **(white symbols)**, with removal of doxycycline for 8 days and resumption of DOX(+) for 6 wk (DOX(-)8d (RD)6wk) **(light gray symbols)**, with removal of doxycycline 8 days prior to sacrifice (DOX(+)6wk DOX(-)8d) **(dark gray symbols)**, or after 7 weeks without doxycycline (DOX(-)7wk) **(black symbols)**.