Title:

Germ layer specific regulation of cell polarity and adhesion: insight into the evolution of

mesoderm.

Authors:

Miguel Salinas-Saavedra[1*], Amber Q. Rock[1], and Mark Q Martindale[1†]

Affiliation:

[1] The Whitney Laboratory for Marine Bioscience, and the Department of Biology, University of

Florida, 9505 N, Ocean Shore Blvd, St. Augustine, FL 32080-8610, USA

*Corresponding author

Email: mssaavedra@whitney.ufl.edu (MS-S)

†Lead contact:

Email: mqmartin@whitney.ufl.edu (MQM)

Summary

In triploblastic animals, Par proteins regulate cell-polarity and adherens junctions of both

ectodermal and endodermal epithelia. But, in embryos of the diploblastic cnidarian *Nematostella*

vectensis, Par proteins are eliminated altogether in the bifunctional endomesodermal epithelia.

Using immunohistochemistry, CRISPR/Cas9, and overexpression of specific mRNAs, we describe

the functional association between Par proteins, ß-catenin, and snail genes in *N.*

vectensis embryos. We demonstrate that the aPKC/Par complex regulates the localization of ß-

catenin in the ectoderm by directing its role in cell adhesion, and that endomesodermal epithelia

are organized by a different cell adhesion system. We also show that *snail* genes, which are

mesodermal markers in bilaterians, are sufficient to downregulate Par proteins and translocate ß-

catenin from the junctions to the cytoplasm in *N. vectensis*. These data provide insight into the

evolution of epithelial structure and the evolution of mesoderm in metazoan embryos.

Introduction

Bilaterian animals comprise more than the 95% of the extant animals on earth and exhibit

enormous body plan diversity (Martindale and Lee, 2013). One of the most important

morphological features in bilaterian evolution is the emergence of the mesoderm, an embryological

tissue that gives rise important cell types such as muscle, blood, cartilage, bone, and kidneys. The

emergence of mesoderm clearly contributed to the explosion of biological diversity throughout

evolution (Martindale, 2005; Martindale and Lee, 2013). Cnidarians (e.g., sea anemones, corals,

hydroids, and "jellyfish") are the sister group to bilaterians, and despite their surprisingly complex

genomes (Putnam et al., 2007), do not possess a distinct mesoderm. Instead, the gastrodermal

lining to their gut cavity consists of a bifunctional endomesoderm with molecular characteristics of

both bilaterian endodermal and myoepithelial mesodermal cells (Jahnel et al., 2014; Martindale

and Lee, 2013; Martindale et al., 2004; Technau and Scholz, 2003). For example, *brachyury* and

snail, among other genes, contribute to the specification of the endomesodermal fates in both

bilaterian and cnidarian embryos (Magie et al., 2007; Martindale et al., 2004; Servetnick et al.,

2017; Technau and Scholz, 2003; Yasuoka et al., 2016). Yet in bilaterians, mesodermal cells

segregate from a endomesodermal precursor (Davidson et al., 2002; Maduro and Rothman, 2002;

Martindale et al., 2004; Rodaway and Patient, 2001; Solnica-Krezel and Sepich, 2012) to form both

endoderm and a third germ layer not present in the embryos of diploblastic cnidarians. How

mesodermal cells originally segregated from the endomesodermal epithelium during animal

evolution is still unclear (Martindale, 2005; Martindale and Lee, 2013; Technau and Scholz, 2003).

During the last decade, several studies have described molecular and cellular characteristics

related to the segregation of mesoderm during bilaterian development (Darras et al., 2011; Keller

et al., 2003; Schäfer et al., 2014; Solnica-Krezel and Sepich, 2012). Here we investigate the

cellular basis of morphogenesis during embryogenesis of the "diploblastic" sea anemone,

Nematostella vectensis.

In most bilaterian embryos described to date, after a series of synchronous and stereotyped

cleavage patterns, maternal determinants redistribute and induce the localization of nuclear ß-

catenin to blastomeres of the vegetal pole (Martindale and Lee, 2013). Hence, gastrulation and the

specification of endomesodermal fates is restricted to the vegetal pole. In these species, *brachyury*

is expressed at the border of the blastopore and *snail* is expressed in the prospective mesodermal

tissues (Technau and Scholz, 2003). The formation of mesoderm involves a variety of cellular

processes including the downregulation of E-cadherin, loss of apicobasal cell polarity, and in some

cases, the induction of epithelial-to-mesenchymal transition (EMT) (Acloque et al., 2009; Lim and

Thiery, 2012; Schäfer et al., 2014; Solnica-Krezel and Sepich, 2012).

Embryos of the cnidarian starlet sea anemone *N. vectensis* develop under non-synchronous and

random cleavage pattern with non-comparable cleavage stages from one embryo to another

(Fritzenwanker et al., 2007; Salinas-Saavedra et al., 2015). During blastula formation, embryonic

cells of *N. vectensis* begin to organize to form an embryo composed of a single hollow epithelial

layer. Epithelial cells of the animal pole, characterized by the nuclear localization of *Nv*ß-catenin

prior to gastrulation (Lee et al., 2007; Wikramanayake et al., 2003), invaginate by apical

constriction to form the endomesodermal epithelium (Magie et al., 2007; Tamulonis et al., 2011).

The expression of *Nvbrachyury* around the presumptive border of the blastopore and *Nvsnail*

genes in the presumptive endomesodermal gastrodermis of *N. vectensis* embryos (Röttinger et al.,

2012; Scholz and Technau, 2003) occurs even before the morphological process of gastrulation

begins.

Interestingly, the components of the Par system (*Nv*aPKC, *Nv*Par-6, *Nv*Par-3, *Nv*Par-1, and

*Nv*Lgl), which show a highly polarized "bilaterian-like" sub cellular distribution throughout all

epithelial cells at the blastula stage in *N. vectensis*, are degraded or down-regulated from the

endomesoderm during the gastrulation process (Figure 1A)(Salinas-Saavedra et al., 2015). We

have previously suggested that the expression of bilaterian "mesodermal genes" (e.g. *Nvsnail)*

might induce the loss of apicobasal cell-polarity indicated by the absence of the components of the

Par system in the endomesoderm of *N. vectensis* embryos (Salinas-Saavedra et al., 2015).

Compelling studies in *N. vectensis* and bilaterians have given information that supports this

hypothesis. For example, recent studies have shown that *snail* is necessary and sufficient to

downregulate Par3 in *Drosophila* mesoderm, inducing the disassembly of junctional complexes in

these tissues (Weng and Wieschaus, 2016, 2017). In addition, we have shown that *Nvbrachyury*

regulates epithelial apicobasal polarity of *N. vectensis* embryos, suggesting conserved

mechanisms with bilaterian systems (Servetnick et al., 2017). Together, this evidence suggests a

plausible cellular and molecular mechanism for the segregation of a distinct cell layer in bilaterian

evolution from an ancestral endomesodermal tissue. Thus, in this study we describe the functional

association between the components of the Par system, apical junctions, epithelial integrity, and

the nuclearization of *Nv*ß-catenin in a cnidarian embryo. In addition, we demonstrate that the

endomesoderm of *N. vectensis* embryos is organized by different junctional complexes that

confers different functional properties than the overlying ectoderm. And finally, we investigate the

putative interactions between the components of the Par system, Wnt canonical pathway, and *snail*

gene expression, giving insights on the evolution of the mesoderm and EMT.

Results and Discussion

Ectodermal and endomesodermal epithelium are joined by different cell-cell adhesion

complexes.

Components of the Par system are not present in the cells of endomesodermal epithelium of *N.*

vectensis during gastrulation, even though the very same cells express these proteins during the

blastula stage (Salinas-Saavedra et al., 2015) (Figure 1). This absence is consistent with the

absence of apical Adherens Junctions (AJs) in the endomesoderm of *N. vectensis* (Figure S1) and

other cnidarians (Chapman et al., 2010; Ganot et al., 2015; Magie et al., 2007). AJs are composed

of transmembrane cadherin molecules that recruit alpha and ß-catenin proteins that link the Par

system to the actin cytoskeleton (Magie and Martindale, 2008; Ohno et al., 2015; Ragkousi et al.,

2017). At polyp stages, neither ß-catenin (an AJ-associated protein) (Figure 1B and 1C) nor the

Par proteins (Figure S1C) are detectable in endomesodermal cells of either the gastrodermis or the

pharynx. When *N. vectensis* embryos are stained with antibodies to ß-catenin (Figure 1) or if *Nv*ß-

catenin::GFP mRNA is expressed in uncleaved zygotes (Figure S1B), clear localization of ß-

catenin can be seen in the cortex of ectodermally derived epithelial cells (Figure 1B, 1C, 1D, 1G,

and 1I), but not in endomesodermal cells (Figure 1B and 1C). In pharyngeal cells that are located

between the epidermis and gastrodermis, *Nv*ß-catenin (Figure 1B and 1D), *Nv*Par-6 (Figure 1E),

and *Nv*Par-1 (Figure 1F) expression begins to disappear, and is localized only in the most apical

regions, indicating that AJs are being disassembled/degraded during the gastrulation process

(Figure 1G and 1H). During planula stages, ß-catenin and the components of the Par system

display scattered patterns in the cytoplasm of a small subset of endomesodermal cells (Figure 1I

and 1J). Even though we do not know the origin and identity of these cells, this expression

temporally coincides with the transient activation of Wnt signaling emanating from the oral pole

(Kusserow et al., 2005; Marlow et al., 2013) at those developmental stages. In bilaterians (Acloque

et al., 2009; Lim and Thiery, 2012) and *N. vectensis* (Kusserow et al., 2005; Marlow et al., 2013),

the later activation of Wnt signaling is also associated with neurogenesis, and may cause the

observed changes in protein localization.

Regardless of this scattered expression, it is clear that cells that undergo gastrulation in *N.*

vectensis lose their polarized ectodermal cell-cell adhesion complex and components of the Par

system, including ß-catenin, are downregulated from endomesodermal tissues (Figure 1). In

bilaterians, the proper formation of an epithelial paracellular barrier (essential for tissue

homeostasis) depends on the establishment of adhesive complexes between adjacent cells

(Higashi et al., 2016), which are regulated by the aPKC/Par complex (Ohno et al., 2015). To test if

this absence of protein expression is correlated to differential cell-cell adhesion in the

endomesodermal epithelium of *N. vectensis*, we assessed potential functional differences in their

role in regulating paracellular movements between ectodermal and endomesodermal epithelia by

using a fluorescent tracer dye penetration assay (Figure 2A)(Higashi et al., 2016). For the

purposes of these experiments, in order to avoid unwanted results related to tissue specification

and cell movements, we used newly hatched juvenile polyps where the gastrodermis

(endomesodermally derived) is fully differentiated.

N. vectensis polyps were exposed to media containing 10,000 MW fluorescent dextran (Molecular

Probes, Inc.). When juvenile polyps are incubated in dextran for 5-10 minutes (Figure 2B),

fluorescent dextran solution moves into the gastric cavity and then spreads into the mesoglea

through the gastrodermal epithelium (Figure 2C), but does not enter the mesoglea through the

outer ectodermally-derived epidermis (Figure 2C and 2D). These results suggest that cell-cell

adhesion is differentially regulated between the epidermis and gastrodermis. Similar results were

obtained in *N. vectensis* polyps when we overexpressed *Nv*Par-3::mVenus by injection of mRNA

into uncleaved eggs which is normally expressed in ectodermal but not endodermal epithelial

tissue (Figure 2D and 2E). However, in polyps expressing a dominant negative version of *Nv*Par-

3::mVenus (dn*Nv*Par-3; microinjected into uncleaved eggs) dye penetrated between epithelial cells

in both the gastrodermis and the outer epidermis (Figure 2D, 2F, and 2G), demonstrating an

ancestral role of the aPKC/Par complex in the maintenance of cell-cell adhesion and the

maintenance of the paracellular boundary of epithelial cells during animal development. These

data suggest that *Nv*Par-3 regulates the formation and maintenance of AJs during *N. vectensis*

embryogenesis.

The *Nv*aPKC/Par complex regulates the formation and maintenance of cell-cell junctions.

Our results suggest that the absence of Par proteins in the endomesoderm is associated with

changes in cell-cell adhesion complexes. Pharmacological treatment of *N. vectensis* embryos with

the aPKC activity inhibitor blocks cytokinesis but not mitosis in cleaving embryos (Figure 3A). In

addition, a dominant negative version of *Nv*Par-1 (dn*Nv*Par-1), that lacks its kinase domain,

localizes only to the cortex of cell-cell contacts (Figure 3B). dn*Nv*Par-1 can be phosphorylated by

aPKC, but cannot phosphorylate the aPKC/Par complex. Thus, dn*Nv*Par-1 is able to localize to the

cell cortex where aPKC is inactive (Figure S2A). These results together suggest that, similar to

bilaterian animals, the formation of cell-cell contacts is regulated by the activity of the aPKC/Par

complex in *N. vectensis* embryos (Figure 3C).

We further tested these observations by using genome editing by CRISPR/Cas9 targeting *Nvpar-6*

and *Nvpar-3* genes (Figure 3D). We did not observe any effects on the embryo until 36 hpf at 16ºC

(late blastula stage), indicating the activity of maternal loaded proteins up until that stage. When

*Nv*Par-6 and *Nv*Par-3 are knocked-out, the ectodermal epithelium loses its integrity, presenting

changes in thickness (Figure S2B and S3A), and interestingly, the endomesodermal epithelium

(which does not express these proteins) becomes a mesenchymal-like endomesoderm (Figure

3D). In *Nvpar-6* and *Nvpar-3* mutant embryos, we also observed the disruption of microtubules and

actin cytoskeleton (Figure S3B), and AJs (visualized with the ß-catenin antibody in Figure 3I) that

confirms our previous observations. Since we did not observe significant changes in the

expression of germ layer markers when these genes were disrupted (e.g. *Nvbra*, *Nvsnail*,

NvSix3/6, and *Nvfz10*; Figure S3E), we believe that only cell adhesion, but not cell specification,

was affected by these experiments. Similar results were obtained when we overexpressed the

mRNA encoding for a dominant negative version *Nv*Par-6 (dn*Nv*Par-6) and *Nv*Par-3 (dn*Nv*Par-3)

into the eggs of *N. vectensis* (Figure S2 and S4). However, dominant negative effects on the

injected embryos were observed at earlier stages (10-12 hpf) than the CRISPR/Cas9 mutants

(zygotic expression) because the mutant proteins must compete with the wild type proteins

(maternally loaded). Hence, in these experiments, lethality (90%) was higher and the phenotypes

were more striking than the phenotypes generated by CRISPR/Cas9 experiments.

Furthermore, the changes observed in the ectodermal and endodermal epithelium after disrupting

*Nv*Par-6 and *Nv*Par-3 (Figure 3) suggests some sort of trans-epithelial regulation of cell-cell

adhesion (most likely involving AJs) because these Par genes are not expressed in the

endomesoderm. The polarizing activity of the aPKC/Par complex in the ectoderm is thus necessary

to maintain the integrity of both ecto- and endodermal epithelia during cellular movements

associated with gastrulation.

To assess whether or not the observed phenotypes on cell-cell adhesion are related to non-

autonomous cell regulation (trans-epithelial interactions), we repeated the above experiments

randomly injecting single blastomeres at 3-4 hpf (8-16 cell-stage). In these experiments, only the

cell-lineage of the injected blastomere would be affected and would exhibit defective cell-cell

adhesion in an otherwise undisturbed wild-type background. If endomesodermal cells derived from

an injected blastomere display fibroblast/mesenchymal cell morphology, it would indicate that the

organization of the endomesodermal epithelium is not dependent on the ectoderm but, rather, an

intrinsic activity of the aPKC/Par complex (Figure 3E). Our results show that only ectodermal- but

not endomesodermal- lineages are affected by these mutations (Figure 3F and S3D). Presumptive

ectodermal cells derived from an injected blastomere fail to maintain AJs and the resulting clone of

epithelial cells loses its structural integrity. In contrast, presumptive endomesodermal cells derived

from an injected blastomere develop into a normal endomesodermal epithelium (Figure 3F). Our

results demonstrate that the proper cell-cell adhesion of the ectodermal layer somehow regulates

trans-epithelially the integrity of the endomesodermal layer. This regulation may maintain the

tension between cells during invagination at gastrula stages, or, in conjunction with the

extracellular matrix (ECM), it may influence signaling patterns necessary to organize epithelial

layers during *N. vectensis* embryogenesis.

NvaPKC/Par complex regulates ß-catenin localization.

In bilaterians, AJs that are forming recruit members of the aPKC/Par complex and the direct

interaction between Par-3 and aPKC/Par-6 is required for the maintenance and maturation of AJs

(Ohno et al., 2015; Ragkousi et al., 2017). AJs are characterized by the binding between cadherin

and ß-catenin: cadherin sequesters ß-catenin from the cytoplasm to the cortex, making it

unavailable for nuclear signaling and endomesoderm specification (Kumburegama et al., 2011;

Wikramanayake et al., 2003). Therefore, using ß-catenin as a marker for AJs, we separately co-

injected *Nv*Par-3::mVenus or dn*Nv*Par-3::mVenus, with *Nv*ß-catenin::RFP into uncleaved zygotes.

We observed cortical co-localization of *Nv*Par-3 and *Nv*ß-catenin at the cell boundaries in the

ectodermal epithelium of embryos co-injected with *Nv*Par-3::mVenus and *Nv*ß-catenin::RFP

(Figure 4A). However, in embryos co-injected with *Nv*ß-catenin::RFP and dn*Nv*Par-3::mVenus, we

observed an alteration of the sub-cellular expression of *Nv*ß-catenin::RFP in all cells as indicated

by the translocation of ß-catenin from the AJs into the nuclei (Figure 4A).

Interestingly, the association between the canonical Wnt signaling pathway and the Par system

has been poorly studied. Two studies, one in *Drosophila* (Sun et al., 2001) and the another in

Xenopus (Ossipova et al., 2005) embryos, have shown by immunoblotting that the kinase Par-1 is

associated with Dishevelled protein and might act as a positive regulator of Wnt signaling. Here, for

the first time, we show *in vivo* embryonic evidence suggesting that *Nv*Par-3 (whose cortical

localization is normally inhibited by Par-1) recruits *Nv*ß-catenin protein and stabilizes its localization

at the apico-lateral cortex of ectodermal cells through the formation of AJs. Furthermore, the

putative disassembly of the aPKC/Par complex induced by dn*Nv*Par-3 overexpression, induces the

nuclearization of *Nv*ß-catenin protein (Figure 4A) due to its failure to be incorporated into cell

adhesion complexes. Strikingly, we also observed the detachments of individual cells from the

ectodermal epithelium of dn*Nv*Par-3 treated-embryos (Figure 4C) and single injected-blastomere

CRISPR/Cas9 *Nv*Par-3 knock-out (Figure 4D). This suggests that these treatments induce EMT-

like processes, not observed under control conditions (Figure 4C).

GSK-3ß regulates the stabilization of cell-cell junctions of epithelial cells in *N. vectensis*

embryos.

Our results and previous studies suggest that the localization of ß-catenin is regulated by its

cytoplasmic availability. Inhibiting the expression of ß-catenin affects both the stabilization of AJs

and the nuclearization of this protein (Leclère et al., 2016). Hence, increasing the concentration of

cytosolic ß-catenin may affect the localization of the aPKC/Par complex. To assess this

hypothesis, we treated *N. vectensis* embryos with 5µm 1-azakenpaullone (AZ), an inhibitor of

GSK-3ß and a canonical Wnt agonist. AZ has been used to upregulate the Wnt/ß-catenin pathway

in many systems, including *N. vectensis* (Leclère et al., 2016; Röttinger et al., 2012).

Approximately 60% of the treated embryos showed an increase in the number of endomesodermal

cells, in agreement with previous studies where ß-catenin could be detected in the nucleus

(Leclère et al., 2016; Röttinger et al., 2012). However, we noticed that around 40% of the embryos

showed two phenotypes not reported by previous studies: 27% (27/100) of the embryos gastrulate

but do not elongate (Figure 5a'), and in the other 13% (13/100) the endomesoderm does not

invaginate at all (Figure 5b'). Both phenotypes displayed severe changes in morphology, loss of

the aboral apical tuft, and an enhanced stabilization of apical microtubules in the endomesodermal

cells.

Surprisingly, we also observed an expansion of the expression domain of Par-6 (Figure 5A) and a

stabilization of AJs (labeled with ß-catenin in Figure 5B) in endomesodermal cells of treated

embryos, which was never observed in control embryos. Par-1 was also detected

immunohistochemically in a few endomesodermal cells, however its ectodermal expression was

clearly not affected by this treatment (Figure 5C). Early cleavage stages (0-4 hpf) were not affected

by this treatment (Figure S2A), thus, we believe that GSK-3ß must maintain apicobasal polarity

once an epithelium is formed. GSK-3ß may regulate the apicobasal cell polarity of epithelial cells

by a direct interaction with the aPKC/Par complex or regulating the cytoskeleton and the

maintenance of AJs. GSK-3ß has been proposed to have different roles during gastrulation and

mesoderm specification by regulating ß-catenin (Röttinger et al., 2012). In other cases, differential

regulation of these protein can inhibit the formation of endomesoderm by regulating cell junctions

(Przybyla et al., 2016). Changes in ECM stiffness can lead to an increment of integrin-based

junctions that up-regulates GSK-3ß and SFKs. SFKs induce the release of ß-catenin from AJs to

the cytoplasm where is degraded by GSK-3ß (Przybyla et al., 2016). Our results are

complimentary to these hypotheses but further research is required.

AJs are down-regulated in mesoderm and neural crest of bilaterian animals.

The segregation of different germ layers during embryogenesis of many bilaterian animals is

carried out by similar cellular mechanisms. EMT is a shared mechanism utilized by mesoderm,

neural crest, and tumorigenesis to generate migratory cells in bilaterian animals (triploblastic

animals). During EMT, the nuclearization of ß-catenin induces the expression of 'endomesodermal'

genes like *brachyury* and *snail* (Acloque et al., 2009). The expression of these genes

downregulates epithelial cadherins, disrupts apicobasal polarity (mediated by the aPKC/Par

complex), disassembles AJs, and induces changes in cytoskeleton organization (Acloque et al.,

2009; Lim and Thiery, 2012). A rearrangement of the actin-myosin cytoskeleton induces apical

constriction of cells (generating a bottle-like shape), which detach from the epithelial sheet, break

down the basal membrane, and invade a specific tissue as mesenchymal cells (Acloque et al.,

2009; Lim and Thiery, 2012).

Interestingly, mesoderm formation, tumorigenesis, and EMT have never been described as natural

processes during *N. vectensis* (a diploblastic animal) embryogenesis. During *N. vectensis*

gastrulation (Magie et al., 2007; Tamulonis et al., 2011), cells around the edge of the blastopore at

the animal pole (which expresses *Nvbrachyury*) acquire a bottle-like shape by apical constriction

(similar to EMT in bilaterians), leading to epithelial buckling and the invagination of presumptive

endomesoderm (which expresses *Nvsnail*). However, throughout this process the endomesoderm

remains as a monolayer of epithelial cells and individual mesenchymal cells never detach and

invade the blastocoel.

Nvsnail genes induce the translocation of _Nv_ß-catenin from AJs to the cytoplasm.

We have shown that by disrupting the aPKC/Par complex (apicobasal cell-polarity) in *N. vectensis*

(Figure 3 and 4), we are able to convert cells from the endomesodermal epithelium into

mesenchymal-like cells, translocate *Nv*ß-catenin (Figure 4A), and emulate EMT-like processes

(apical constriction and individual cell-detachments) in the ectodermal epithelium of *N. vectensis*

treated-embryos (Figure 4C and 4D). These results demonstrate that the cnidarian *N. vectensis*

possesses mechanisms necessary to segregate individual germ layers (e.g. mesoderm and neural

crest) described in bilaterians via the downregulation of Par proteins and AJs from the

endomesodermal epithelium. We recently showed that *Nvbrachyury* regulates apicobasal polarity

of epithelial cells in *N. vectensis* embryos (Servetnick et al., 2017). In bilaterians, it has been

shown that EMT is controlled by the genes *brachyury* and *snail* (Acloque et al., 2009; Lim and

Thiery, 2012). We therefore examined the role of *Nvsnail* genes on the localization of ß-catenin,

components of the Par system, and the stabilization of AJs. Our hypothesis is that expression of *N.*

vectensis snail genes will destabilize AJs, and induce the nuclearization of ß-catenin in ectodermal

epithelial cells.

N. vectensis has two *snail* genes, *Nvsnail-A* and *Nvsnail-B*, which are both expressed in the

endomesodermal plate prior to and throughout the gastrulation process, and which define the

boundary between gastrodermis and ectodermal pharynx (Amiel et al., 2017; Magie et al., 2007;

Röttinger et al., 2012). The overexpression of both *Nvsnail-A and Nvsnail-B* together induce the

ectopic translocation of *Nv*ß-catenin to the nuclei of ectodermal cells when the mRNA of the three

proteins are co-injected into uncleaved eggs (Figure 6A). This treatment also delocalizes *Nv*Par-3

from the cell cortex (Figure 6B). Although all of the cells expressed the injected mRNAs, not every

cell was affected by this treatment, suggesting that this patched pattern is a result of ectodermal vs

endomesodermal differences in response to *Nvsnail* over-expression (or the co-expression with

*Nv*ß-catenin and *Nv*Par-3). Supporting this idea is the fact that the effects of the *Nvsnail* gene on

AJs were not observed earlier than 36 hpf. When gastrulation is complete, nuclear ß-catenin is no

longer observable in either ecto or endomesodermal cells, and only its cortical localization (AJs)

remains in the ectodermal epithelium.

To segregate potential spatial differences in the effects of *Nvsnail* genes on cell adhesion/epithelial

polarity, we randomly injected single blastomeres at the 8-16 cell-stage with mRNA from both

Nvsnail-A and Nvsnail-B together. In this way, we can use the fluorescent dextran to detect the

clones where the over-expression of the co-injected mRNAs occurred in a "wild-type" background

(Figure 6D). Similar to the *Nvpar-3* knock-out (Figure 3F), the expression of *Nvsnail* genes together

is sufficient to induce the degradation of Par proteins and AJs (ß-catenin) from the ectoderm, and

disrupts its epithelial integrity (Figure 6D). Concordantly with our *in vivo* observations, not all

regions of the embryo are affected by this treatment (Figure 6E). However, nuclear ß-catenin was

not observed under these treatments. Thus, nuclear ß-catenin observed in Figure 6A is a

consequence of the high cytosolic availability generated by its overexpression and release from

AJs.

These results suggest that the role of *Nvsnail* genes on AJs and apicobasal cell polarity is spatially

and temporally constrained to the site of gastrulation in *N. vectensis* embryos, and that these

genes in may be required for gastrulation movements. Therefore, we predict that ß-catenin (AJs)

and Par proteins will be retained in the cells of the *N. vectensis* endomesodermal plate if *Nvsnail*

genes are disrupted together.

Nvsnail genes downregulate apicobasal cell polarity and AJs in the endomesodermal

epithelium of *N. vectensis* embryos.

Snail genes temporally down-regulate E-cadherin during mesoderm segregation and EMT in

bilaterian animals (Lim and Thiery, 2012). Subsequently, the apicobasal polarity and AJs of the

cells expressing *snail* are disrupted, leading to the detachment of individual (or a group of) cells

from the epithelium (Lim and Thiery, 2012). As we have shown here, as well as in previous studies

(Magie et al., 2007; Magie and Martindale, 2008), the cells comprising the endomesodermal plate

lose their cell-cell adhesion during gastrulation in *N. vectensis* embryos. It may be possible that

temporal regulation of the endomesodermal patterning might act upon the AJs as it has been

proposed in other bilaterian systems. Our data suggest that once gastrulation is complete and the

pharynx forms, components of the Par system and ß-catenin (AJs) are degraded from both the

cortex and cytoplasm of endomesodermal cells (Figure 1 and S1). Hence, it could be possible that

Nvbrachyury induces the disruption of apicobasal polarity (Servetnick et al., 2017), remnant AJs

maintain the endomesodermal-plate cells together, and *Nvsnail* genes degrades and prevents the

reassembly of AJs in the endomesoderm of *N. vectensis.*

To address these issues, we used CRISPR/Cas9 knock-out of *Nvsnail-A* and *Nvsnail-B* genes

together to inhibit zygotic function of these genes and investigate their role on the temporal

regulation of AJs and cell polarity. In CRISPR/Cas9 mutants, the endomesodermal plate forms but

gastrulation does not proceed (Figure 6F). Furthermore, AJs (labeled with ß-catenin) and apical

Par proteins (labeled with anti*Nv*Par-6 and anti*Nv*aPKC) are retained at the apical cortex of the

cells of the endomesodermal plate (Figure 6F and S5). Surprisingly, *Nv*Par-1 and *Nv*Lgl were not

detected in those cells (Figure 6F), suggesting that the degradation of these basolateral proteins

precede or do not depend on the activity of the *Nvsnail* genes. In other words, *Nvsnail* regulates

apical cell-cell adhesion controlling the elongation of columnar cell shape ('zippering') but not the

invagination of the endomesodermal plate during gastrulation of *N. vectensis* embryos.

Interestingly, the invagination of the endomesodermal plate (controlled by the Wnt/PCP pathway)

is uncoupled from its specification (Wnt/ß-catenin pathway) in *N. vectensis* embryos

(Kumburegama et al., 2011; Wijesena, 2012). Furthermore, components of the Wnt/PCP pathway

are expressed only in the endomesoderm (Kumburegama et al., 2011; Wijesena, 2012), while

components of the Par system are expressed only in the ectoderm (Salinas-Saavedra et al., 2015).

This suggests these differences regulate the integrity of both ecto- and endomesodermal epithelia,

and the endomesodermal specification (mediated by *Nvsnail* and other endomesodermal genes,

e.g., *Nvbrachyury* and *Nvtwist*) regulates and maintains this differential cell-cell adhesion between

both germ layers, allowing gastrulation movements and differentiation.

Therefore, in this scenario, the *N. vectensis* embryo is composed of two independent

morphogenetic modules, which are integrated and organized together by the pharynx. One, the

ectoderm, whose apicobasal polarity (and thus, AJs and epithelial integrity) is regulated by

Nvbrachyury allowing ectodermal epiboly and pharynx formation (Servetnick et al., 2017). The

other one, the endomesoderm, whose differentiation and cell-movements are regulated by *Nvsnail*.

This is supported by the expression *Nvbrachyury* in *Nvsnail* knock-out embryos (Figure S6), and

Nvsnail knock-out phenotypes where ectodermal pharynx but no clear endomesoderm is formed

(Figure S5). Additional work is required to elucidate any differences in function between *Nvsnail-A*

and *Nvsnail-B* genes. Furthermore, both modules are specified by nuclear ß-catenin (Röttinger et

al., 2012), suggesting that the nuclear ß-catenin translocation from the animal pole in cnidarians to

the vegetal pole in bilaterians is mechanistically plausible and sufficient to re-specify the site of

gastrulation and germ-layers along the animal-vegetal axis during Metazoan evolution (Lee et al.,

2007; Martindale and Lee, 2013).

The dual identity and collective migration of the endomesodermal cells.

Bilaterian-EMT has been a focus of study during the last decade as the mechanism to segregate

different cell layers involved in different biological processes (e.g., formation of the mesoderm,

neural crest, and tumorigenesis). However, this process appears to depend on the fine regulation

of the level or expression of *snail* genes and their temporal activity, allowing 'intermediate' cell

states. For example, during neural crest migration, cells display 'partial-EMT' phenotypes where

cells are maintained together but the apicobasal polarity and AJs are down-regulated (Lee et al.,

2006; Nieto et al., 2016; Ribeiro and Paredes, 2014; Theveneau and Mayor, 2013). Our data

suggest that a partial-EMT may be the mechanism by which the endomesodermal epithelium

moves into the blastocoel in *N. vectensis* embryos (Figure 7). In this scenario, mesodermal

specifiers (e.g. *snail* and *brachyury*) may down-regulate AJs and induce EMT-like processes, and

endodermal characteristics may keep the necessary cell adhesion systems together in the *N.*

vectensis endomesodermal epithelium.

In bilaterian animals, there are many other pathways in addition to the cWnt that activate *snail*

transcription and induce the disruption of AJs and apicobasal cell polarity. For example, TGFß,

BMP, NANOS, FGF, and MEK/ERK/ERG take on roles during the specification of mesoderm,

neural crest, tumorigenesis, and other EMT-related processes (Barrallo-Gimeno and Nieto, 2005;

Lim and Thiery, 2012; Nieto et al., 2016). Changes in cell-polarity, cell-cell adhesion, cytoskeleton

organization, nuclear shape, and permeability, among others, characterize the cells of tissues that

are being specified by those signaling pathways (e.g., neurons, gonads, regenerative tissues,

cardiac and vascular system, and muscles). Concordantly in *N. vectensis* embryos, cells of the

pharyngeal and endomesodermal tissues present all these characteristics and express

components of all these pathways (Amiel et al., 2017; Extavour et al., 2005; Matus et al., 2006,

2007; Röttinger et al., 2012; Wijesena et al., 2017). For example, one cadherin (*Nv*CDH2 (Clarke

et al., 2016): 1g244010), and kinases that modify tubulin and histones are differentially regulated

between ecto- and endomesodermal epithelium (Wijesena et al., 2017).

In conclusion, *N. vectensis* has all the components and cellular mechanisms to form a distinct

mesoderm as seen in triploblastic bilaterian animals. However, the mechanism(s) that regulates

the segregation of a distinct mesoderm must have been absent during the evolution of cnidarians.

This mechanism must have evolved to segregate mesodermal cells from the endoderm or to

tighten cell-cell junctions and segregate the endoderm from mesodermal cells. Intriguingly,

ctenophores segregate a mesodermal cell population during embryogenesis but do not have the

genes that encode for all cell-cell adhesion complexes and specify for mesoderm in bilaterians

(Figure S7) (Ganot et al., 2015; Ryan et al., 2013). Further research and funding is needed to

understand the cellular mechanisms that evolve to segregate mesoderm and control epithelial cell

polarity at the base of the metazoan tree.

Acknowledgments

444 We thank Leslie S. Babonis for TEM data in Figure S1, A. Wikramanayake for *Nvß*-catenin

445 expression constructs, and C. Magie and E. Röttinger for *Nvsnail-A* expression constructs. We

446 thank E. Seaver, C.E. Schnitzler, and the members of Martindale's lab for helpful discussion. We

447 also thank to 'choose development' program for AQR. This research was supported by the NIH

448 GM093116, NASA 16-EXO16_2-0041, and Synthego grant from MQM.

Author Contributions

451 MS-S., and MQM. designed research and analyzed data. MS-S. performed research with help of

452 AQR and MQM. MS-S., and MQM. wrote the manuscript with help of AQR. All authors read and

453 approved the final manuscript.

Declaration of Interests

456 The authors declare no competing interests.

Bibliography

Acloque, H., Adams, M.S., Fishwick, K., Bronner-Fraser, M., and Nieto, M.A. (2009). Epithelial-

mesenchymal transitions:the importance of changing cell statein development and disease.

Journal of Clinical I*Nv*estigation *119*, 1438-1449.

Amiel, A.R., Johnston, H., Chock, T., Dahlin, P., Iglesias, M., Layden, M., Röttinger, E., and

Martindale, M.Q. (2017). A bipolar role of the transcription factor ERG for cnidarian germ layer

formation and apical domain patterning. Developmental Biology *430*, 346-361.

Barrallo-Gimeno, A., and Nieto, M.A. (2005). The Snail genes as inducers of cell movement and

survival: implications in development and cancer. Development (Cambridge, England) *132*, 3151-

3161.

Chapman, J.A., Kirkness, E.F., Simakov, O., Hampson, S.E., Mitros, T., Weinmaier, T., Rattei, T.,

Balasubramanian, P.G., Borman, J., Busam, D.*, et al.* (2010). The dynamic genome of Hydra.

Nature *464*, 592-596.

Clarke, D.N., Miller, P.W., Lowe, C.J., Weis, W.I., and Nelson, W.J. (2016). Characterization of the

Cadherin-Catenin Complex of the Sea Anemone Nematostella vectensis and Implications for the

Evolution of Metazoan Cell-Cell Adhesion. Molecular Biology and Evolution *33*, 2016-2029.

Darras, S., Gerhart, J., Terasaki, M., Kirschner, M., and Lowe, C.J. (2011). β-catenin specifies the

endomesoderm and defines the posterior organizer of the hemichordate Saccoglossus

kowalevskii. Development (Cambridge, England) *138*, 959-970.

Davidson, E.H., Rast, J.P., Oliveri, P., Ransick, A., Calestani, C., Yuh, C.-H., Minokawa, T.,

Amore, G., Hinman, V., Arenas-Mena, C.*, et al.* (2002). A Provisional Regulatory Gene Network for

Specification of Endomesoderm in the Sea Urchin Embryo. Developmental Biology *246*, 162-190.

Doench, J.G., Hartenian, E., Graham, D.B., Tothova, Z., Hegde, M., Smith, I., Sullender, M., Ebert,

B.L., Xavier, R.J., and Root, D.E. (2014). Rational design of highly active sgRNAs for CRISPR-

Cas9–mediated gene inactivation. Nature Biotechnology, 1-8.

Extavour, C.G., Pang, K., Matus, D.Q., and Martindale, M.Q. (2005). vasa and nanos expression

patterns in a sea anemone and the evolution of bilaterian germ cell specification mechanisms.

Evolution & Development *7*, 201-215.

Fritzenwanker, J.H., Genikhovich, G., Kraus, Y., and Technau, U. (2007). Early development and

axis specification in the sea anemone Nematostella vectensis. Developmental Biology *310*, 264-

279.

Ganot, P., Zoccola, D., Tambutté, E., Voolstra, C.R., Aranda, M., Allemand, D., and Tambutté, S.

(2015). Structural molecular components of septate junctions in cnidarians point to the origin of

epithelial junctions in eukaryotes. Molecular Biology and Evolution *32*, 44-62.

Hand, C., and Uhlinger, K.R. (1992). The Culture, Sexual and Asexual Reproduction, and Growth

of the Sea Anemone Nematostella vectensis. Biological Bulletin *182*, 169-176.

Higashi, T., Arnold, T.R., Stephenson, R.E., Dinshaw, K.M., and Miller, A.L. (2016). Maintenance

of the Epithelial Barrier and Remodeling of Cell-Cell Junctions during Cytokinesis. Current Biology,

1-31.

Ikmi, A., McKinney, S.A., Delventhal, K.M., and Gibson, M.C. (2014). TALEN and CRISPR/Cas9-

mediated genome editing in the early-branching metazoan Nematostella vectensis. Nature

communications *5*, 5486.

Jahnel, S.M., Walzl, M., and Technau, U. (2014). Development and epithelial organisation of

muscle cells in the sea anemone Nematostella vectensis. Frontiers in zoology *11*, 44-15.

Keller, R., Davidson, L.A., and Shook, D.R. (2003). How we are shaped: the biomechanics of

gastrulation. Differentiation; research in biological diversity *71*, 171-205.

Kumburegama, S., Wijesena, N., Xu, R., and Wikramanayake, A.H. (2011). Strabismus-mediated

primary archenteron i*nv*agination is uncoupled from Wnt/β-catenin-dependent endoderm cell fate

specification in Nematostella vectensis (Anthozoa, Cnidaria): Implications for the evolution of

gastrulation. EvoDevo *2*, 2.

Kusserow, A., Pang, K., Sturm, C., Hrouda, M., Lentfer, J., Schmidt, H.A., Technau, U., von

Haeseler, A., Hobmayer, B., Martindale, M.Q.*, et al.* (2005). Unexpected complexity of the Wnt

gene family in a sea anemone. Nature *433*, 156-160.

Layden, M.J., Röttinger, E., Wolenski, F.S., Gilmore, T.D., and Martindale, M.Q. (2013).

Microinjection of mRNA or morpholinos for reverse genetic analysis in the starlet sea anemone,

Nematostella vectensis. Nature Protocols *8*, 924-934.

Leclère, L., Bause, M., Sinigaglia, C., Steger, J., and Rentzsch, F. (2016). Development of the

aboral domain in Nematostellarequires β-cateninand the opposing activities of Six3/6and

Frizzled5/8. Development (Cambridge, England) *143*, 1766-1777.

Lee, J.M., Dedhar, S., Kalluri, R., and Thompson, E.W. (2006). The epithelial–mesenchymal

transition: new insights in signaling, development, and disease. The Journal of Cell Biology *172*,

973-981.

Lee, P.N., Kumburegama, S., Marlow, H.Q., Martindale, M.Q., and Wikramanayake, A.H. (2007).

Asymmetric developmental potential along the animal-vegetal axis in the anthozoan cnidarian,

Nematostella vectensis, is mediated by Dishevelled. Developmental Biology *310*, 169-186.

Lim, J., and Thiery, J.P. (2012). Epithelial-mesenchymal transitions: insights from development.

Development (Cambridge, England) *139*, 3471-3486.

Maduro, M.F., and Rothman, J.H. (2002). Making Worm Guts: The Gene Regulatory Network of

the Caenorhabditis elegans Endoderm. Developmental Biology *246*, 68-85.

Magie, C.R., Daly, M., and Martindale, M.Q. (2007). Gastrulation in the cnidarian Nematostella

vectensis occurs via i*nv*agination not ingression. Developmental Biology *305*, 483-497.

Magie, C.R., and Martindale, M.Q. (2008). Cell-cell adhesion in the cnidaria: insights into the

evolution of tissue morphogenesis. The Biological Bulletin *214*, 218-232.

Marlow, H., Matus, D.Q., and Martindale, M.Q. (2013). Ectopic activation of the canonical wnt

signaling pathway affects ectodermal patterning along the primary axis during larval development

in the anthozoan Nematostella vectensis. Developmental Biology *380*, 324-334.

Martindale, M. (2005). The evolution of metazoan axial properties. Nature reviews Genetics *6*, 917-

927.

Martindale, M.Q., and Lee, P.N. (2013). The Development of Form: Causes and Consequences of

Developmental Reprogramming Associated with Rapid Body Plan Evolution in the Bilaterian

Radiation. Biological Theory *8*, 253-264.

Martindale, M.Q., Pang, K., and Finnerty, J.R. (2004). I*nv*estigating the origins of triploblasty:

'mesodermal' gene expression in a diploblastic animal, the sea anemone Nematostella vectensis

(phylum, Cnidaria; class, Anthozoa). Development (Cambridge, England) *131*, 2463-2474.

Matus, D.Q., Thomsen, G.H., and Martindale, M.Q. (2006). Dorso/ventral genes are

asymmetrically expressed and i*nv*olved in germ-layer demarcation during cnidarian gastrulation.

Current Biology *16*, 499-505.

Matus, D.Q., Thomsen, G.H., and Martindale, M.Q. (2007). FGF signaling in gastrulation and

neural development in Nematostella vectensis, an anthozoan cnidarian. Development Genes and

Evolution *217*, 137-148.

Moreno-Mateos, M.A., Vejnar, C.E., Beaudoin, J.-D., Fernandez, J.P., Mis, E.K., Khokha, M.K.,

and Giraldez, A.J. (2015). CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9

targeting in vivo. Nature Methods *12*, 982-988.

Nieto, M.A., Huang, R.Y.-J., Jackson, R.A., and Thiery, J.P. (2016). EMT: 2016. Cell *166*, 21-45.

Ohno, S., Goulas, S., and Hirose, T. (2015). The PAR3-aPKC-PAR6 Complex. In Cell Polarity 1:

Biological Role and Basic Mechanisms, K. Ebnet, ed. (Cham: Springer International Publishing),

pp. 3-23.

Ossipova, O., Dhawan, S., Sokol, S., and Green, J.B.A. (2005). Distinct PAR-1 proteins function in

different branches of Wnt signaling during vertebrate development. Developmental Cell *8*, 829-841.

Przybyla, L., Lakins, J.N., and Weaver, V.M. (2016). Tissue Mechanics Orchestrate Wnt-

Dependent Human Embryonic Stem Cell Differentiation. Stem Cell, 1-27.

Putnam, N.H., Srivastava, M., Hellsten, U., Dirks, B., Chapman, J., Salamov, A., Terry, A., Shapiro,

H., Lindquist, E., Kapitonov, V.V.*, et al.* (2007). Sea anemone genome reveals ancestral

eumetazoan gene repertoire and genomic organization. Sup. Science *317*, 86-94.

Ragkousi, K., Marr, K., McKinney, S., Ellington, L., and Gibson, M.C. (2017). Cell-Cycle-Coupled

Oscillations in Apical Polarity and Intercellular Contact Maintain Order in Embryonic Epithelia.

Current Biology, 1-19.

Ribeiro, A.S., and Paredes, J. (2014). P-Cadherin Linking Breast Cancer Stem Cells and I*nv*asion:

A Promising Marker to Identify an "Intermediate/Metastable" EMT State. Frontiers in

oncology *4*, 371.

Rodaway, A., and Patient, R. (2001). Mesendoderm. an ancient germ layer? Cell *105*, 169-172.

Röttinger, E., Dahlin, P., and Martindale, M.Q. (2012). A framework for the establishment of a

cnidarian gene regulatory network for 'endomesoderm' specification: the inputs of ß-catenin/TCF

signaling. PLoS genetics *8*, e1003164.

Roure, A., Rothbächer, U., Robin, F., Kalmar, E., Ferone, G., Lamy, C., Missero, C., Mueller, F.,

and Lemaire, P. (2007). A multicassette Gateway vector set for high throughput and comparative

analyses in ciona and vertebrate embryos. PLoS One *2*, e916.

Ryan, J.F., Pang, K., Schnitzler, C.E., Nguyen, A.D., Moreland, R.T., Simmons, D.K., Koch, B.J.,

Francis, W.R., Havlak, P., Program, N.C.S.*, et al.* (2013). The Genome of the Ctenophore

Mnemiopsis leidyi and Its Implications for Cell Type Evolution. Science *342*, 1242592-1242592.

Salinas-Saavedra, M., Stephenson, T.Q., Dunn, C.W., and Martindale, M.Q. (2015). Par system

components are asymmetrically localized in ectodermal epithelia, but not during early development

in the sea anemone Nematostella vectensis. EvoDevo *6*.

Schäfer, G., Narasimha, M., Vogelsang, E., and Leptin, M. (2014). Cadherin switching during the

formation and differentiation of the Drosophila mesoderm - implications for epithelial-to-

mesenchymal transitions. Journal of Cell Science *127*, 1511-1522.

Scholz, C.B., and Technau, U. (2003). The ancestral role of Brachyury: expression of NemBra1 in

the basal cnidarian Nematostella vectensis (Anthozoa). Archiv für Entwickelungsmechanik der

Organismen *212*, 563-570.

Servetnick, M.D., Steinworth, B., Babonis, L.S., Simmons, D., Salinas-Saavedra, M., and

Martindale, M.Q. (2017). Cas9-mediated excision of Nematostella brachyury disrupts

endoderm development, pharynx formation, and oral-aboral patterning. Development.

Solnica-Krezel, L., and Sepich, D.S. (2012). Gastrulation: making and shaping germ layers. Annual

review of cell and developmental biology *28*, 687-717.

Sun, T.Q., Lu, B., Feng, J.J., Reinhard, C., Jan, Y.N., Fantl, W.J., and Williams, L.T. (2001). PAR-1

is a Dishevelled-associated kinase and a positive regulator of Wnt signalling. Nature Cell Biology *3*,

628-636.

Tamulonis, C., Postma, M., Marlow, H.Q., Magie, C.R., de Jong, J., and Kaandorp, J. (2011). A

cell-based model of Nematostella vectensis gastrulation including bottle cell formation,

i*Nv*agination and zippering. Developmental Biology *351*, 217-228.

Technau, U., and Scholz, C.B. (2003). Origin and evolution of endoderm and mesoderm. The

International Journal of Developmental Biology *47*, 531-539.

Theveneau, E., and Mayor, R. (2013). Collective cell migration of epithelial and mesenchymal

cells. Cellular and molecular life sciences : CMLS *70*, 3481-3492.

Weng, M., and Wieschaus, E. (2016). Myosin-dependent remodeling of adherens junctions

protects junctions from Snail-dependent disassembly. The Journal of Cell Biology *212*, 219-229.

Weng, M., and Wieschaus, E. (2017). Polarity protein Par3/Bazooka follows myosin-dependent

junction repositioning. Developmental Biology *422*, 125-134.

Wijesena, N., Simmons, D.K., and Martindale, M.Q. (2017). Antagonistic BMP-cWNT signaling in

the cnidarian Nematostella vectensis reveals insight into the evolution of mesoderm. Proceedings

of the National Academy of Sciences of the United States of America *114*, E5608-E5615.

Wijesena, N.M. (2012). Wnt Signaling in the Cnidarian Nematostella vectensis: Insighits into the

Evolution of Gastrulation (University of Miami), pp. 1-120.

Wikramanayake, A.H., Hong, M., Lee, P.N., Pang, K., Byrum, C.A., Bince, J.M., Xu, R., and

Martindale, M.Q. (2003). An ancient role for nuclear catenin in the evolution of axial polarity and

germ layer segregation. Nature *426*, 446-450.

Wolenski, F.S., Layden, M.J., Martindale, M.Q., Gilmore, T.D., and Finnerty, J.R. (2013).

Characterizing the spatiotemporal expression of RNAs and proteins in the starlet sea anemone,

Nematostella vectensis. Nature Protocols *8*, 900-915.

Yasuoka, Y., Shinzato, C., and Satoh, N. (2016). The Mesoderm-Forming Gene brachyury

Regulates Ectoderm-Endoderm Demarcation in the Coral Acropora digitifera. Current Biology, 1-

21.

Figure 1. Components of the Par system and ß-catenin are downregulated from the *N.

vectensis endomesoderm during gastrulation.** A-F. Confocal images of immunofluorescent

staining (IFS) of lateral views of gastrulation embryos (animal pole up). The * marks the site of

gastrulation in all cases. Samples are counterstained with Phalloidin (Phall) staining (white) to

626 show cell boundaries, DAPI to visualize cell nuclei (blue), and Tubulin antibody (Tub) staining is

627 shown as counterstain (green). All images are a single optical section from a z-stack confocal

628 series. All scale bars, 20 μm.

629 (A) Summary diagram depicting the localization of ß-catenin and Par proteins at the observed

630  stages. Pale boxes denote changes observed in the endomesoderm.

631 (B) IFS for ß-catenin (magenta) in primary polyps. High magnification images from boxed region

632  (endomesoderm, Endo) are shown on the bottom. Arrows indicate the absence of ß-catenin

633  expression in the endomesoderm. Arrowheads indicate the ß-catenin expression in the

634  ectodermal pharynx (EP). Star indicates the endomesodermal pharynx (EnP). Histone antibody

635  (Hist) staining is shown as counterstain to show the penetrability in the fixed tissue. See also

636  Figure S1.

637 (C) IFS for ß-catenin (magenta) in the ecto and endomesoderm (arrow) of primary polyps.

638 (D) IFS for ß-catenin (magenta) at 24 hpf shows localization to the apical domain where adherens

639  junctions reside in all cells of the blastula. High magnification images from boxed region

640  (prospective endomesoderm) are shown on the right.

641 (E) IFS for *Nv*Par-6 (magenta) at 24 hpf showing the same sub-cellular localization as ß-catenin

642  (A). High magnification images from boxed region in (A) (prospective endomesoderm) are

643  shown on the right. Merged image shown on upper left.

644 (F) IFS for *Nv*Par-1 at 24 hpf shows a complementary basolateral expression. High magnification

645  images from boxed region (prospective endomesoderm) are shown on the right.

646 (G) IFS for ß-catenin at 30 hpf shows the loss of expression of ß-catenin (magenta) in invaginating

647  endomesoderm (box). The arrow (D-F) marks the boundary between ectoderm and

648  invaginating endomesoderm. High magnification images from boxed region (prospective

649  endomesoderm) are shown on the right.

650 (H) IFS for *Nv*Par-6 and *Nv*Par-1(magenta) at 30 hpf show that all Par proteins are down regulated

651  at the site of gastrulation. IFS for *Nv*Par-6 shows an even earlier down regulation than ß-

652  catenin (D). High magnification images from boxed region (prospective endomesoderm) are

653  shown on the right. Merged image shown on upper left.

(I) Oral view of IFS for ß-catenin (magenta) at 96 hpf showing apical localization in overlying

ectoderm, but absence in endomesodermal tissues. The two bottom panels show high

magnifications of the endomesoderm region (image inverted). Arrowheads indicate the

localization of ß-catenin expression (black) in some scattered endomesodermal cells.

(J) Lateral view of IFS for *Nv*aPKC and *Nv*Lgl (magenta) at 96 hpf showing loss of expression in

invaginating epithelial cells. The four bottom panels show high magnifications of the

endomesoderm region (image inverted). Arrowheads indicate the localization of *Nv*aPKC and

*Nv*Lgl proteins (black) in some scattered endomesodermal cells.

Figure 2. The aPKC/Par complex maintains Adherens Junctions (AJs) of ectodermal

epithelial cells.

Arrows indicate the direction of the flow: from gastric cavity (gc) to mesoglea (white) and from

external media (ex) to ectoderm (blue). Dashed lines indicate the base of the epidermis. All images

are single optical section from the z-stack confocal series. See also Figure S2 for dn*Nv*Par-3

description. Scale bars, 20 μm.

(A) Diagram depicting the hypothesis that when the aPKC/Par complex is functional (top row), AJs

are present (blue stripes) and a paracellular epithelial barrier is formed. When aPKC/Par

complex is not functional (bottom row), AJs are disrupted, the epithelial barrier is perturbed,

and the extracellular solution moves paracellularly into the mesoglea.

(B) Penetration assay of wild type (uninjected) primary polyps at low magnification showing the

movement of 10,000MW fluorescent dextran. Top row, no dextran. Bottom row, dextran

(yellow) in the gc moves in to the mesoglea through paracellular spaces between gastrodermal

cells (arrows).

(C) High magnification images from box shown in (B). *: mesoglea (purple band) that separates the

ectoderm (ECT, dashed line) from gastrodermis (GDR). Note the dye moving between cells

from the gc media (arrows)

(D) Low magnification images comparing polyps expressing *Nv*Par-3::mVenus and a dominant

negative version of *Nv*Par-3 (dn*Nv*Par-3::mVenus) expressing-embryos. Dextran media (DM;

extracellular) is pseudo-colored yellow. Dextran (red) was co-injected with mRNAs to label the

cells and differentiate intracellular regions. mVenus channel was omitted for better

visualization. Lower concentrations of dn*Nv*Par-3 were injected to preserve endomesodermal

tissues. Note that the dextran media was found between the cells labeled in red.

(E) High magnification images from (F) boxed region in (E). Purple band depicts Mesoglea.

(F) High magnification images from (G) boxed region in (E).

(G) High magnification images from (H) boxed region in (E).

*: Paracellular spaces of both, the epidermis (blue) and gastrodermis (white).

Figure 3. Ectodermal *Nv*aPKC/Par complex polarity regulates the epithelial integrity of both

ecto- and endomesoderm.

(A) IFS for *Nv*aPKC at 4 hpf showing that the aPKC inhibitor (Sigma P1614) blocks cytokinesis but

not cell cycle.

(B) In vivo expression of dn*Nv*Par-1 shows precocious localization to zones of cell contact during

cleavage stages, well before wild-type *Nv*Par genes do. See also Figure S2A.

(C) Diagram depicting the suggested the working hypothesis.

(D) CRISPR/Cas9 knock-out for *Nv*Par-6 and *Nv*Par-3 at 48 hpf. Controls show no effect on

gastrulation. Tubulin (Tub), Phalloidin (Phall), and DAPI are used as counterstains.

CRISPR/Cas9 mutants: tubulin stained low magnification of CRISPR phenotype. High

magnification images from boxed region shows mesenchymal-like cells. Arrowheads indicate

filopodia-like structures. Number of cases observed for each gene are shown. See also Figure

S2, S3, and S4.

(E) Diagram depicts the hypothesis addressed in (I). The cell lineage derived from a single

injected-blastomere is in green.

(F) IFS for ß-catenin (ß-cat), *Nv*Par-6, *Nv*aPKC, and *Nv*Par-1 in single injected-blastomere

CRISPR/Cas9 *Nv*Par-3 knock-outs at 40 hpf. Streptavidin-Biotin TxRed Dextran (Dex) is

shown in green. Arrowheads indicate the absence of the protein and disrupted epithelium.

Arrows indicate bottle-like shape cells. * indicate the orientation of the site of gastrulation. See

also Figure S3D.

Morphology is shown by DAPI, Tub, and Phall IFS. Except for 3B and 3D, all images are single

optical sections from the z-stack confocal series. (B) and (D) are 3D reconstructions from a z-stack

confocal series. All scale bars, 20 µm.

Figure 4. *Nv*aPKC/Par complex regulates ß-catenin localization and cell attachment.

(A) In vivo co-localization of *Nv*Par-3venus co-injected with *Nv*ß-cateninRFP, and dn*Nv*Par-3venus

co-injected with *Nv*ß-cateninRFP. Arrowheads indicate junctions (AJs). Arrows indicate nuclear

ß-catenin.

(B) Diagram of the suggested interpretation for A.

(C) In vivo time series of ectodermal epithelial layers of embryos injected with *Nv*Par-3venus and

dn*Nv*Par-3venus mRNA demonstrating epithelial delamination in the absence of functional

*Nv*Par3. Lifeact::mTq2 mRNA was co-injected to visualize cell boundaries. Pink arrows indicate

the absence cell detachments. A subset of cells was artificially colored. The purple cell

detaches from the epithelium and the red arrow indicates a second cell

detachment. See also Movie S1.

(D) IFS of an embryo in which a single blastomere was injected with *Nv*Par-3 guide RNAs and

Cas9 and green dextran. Red arrowheads indicate the apical constriction and delamination of

ectodermal cells in the mutated clone of cells. Note the different layers of nuclei stained with

DAPI. Asterisks indicate the site of gastrulation.

(E) Diagram of the suggested interpretation for D.

All images are 3D reconstructions from a z-stack confocal series. All scale bars, 20 µm.

Figure 5. Apical junctions are regulated by GSK-3ß in epithelial cells during gastrulation in

***N. vectensis* embryos.**

(A) IFS for *Nv*Par-6 in AZ-treated embryos.

(B) IFS for ß-catenin in AZ-treated embryos.

(C) IFS for *Nv*Par-1 in AZ-treated embryos.

(D) Graphical summary of the observed results.

Arrowheads indicate the localization of *Nv*Par-6, ß-catenin, and *Nv*Par-1 in the endomesodermal

epithelium of AZ-treated embryos. Arrows indicated the stabilization of microtubules at the apical

cortex of endomesodermal cells. Two distinct phenotypes were observed after AZ treatment: (a')

gastrulation without elongation, (b') no invagination of the endomesoderm.

Figure 6. *Nvsnail* genes downregulate AJs and *Nv*aPKC/Par polarity allowing

endomesodermal migration.

(A) in vivo localization of *Nv*ß-cateninGFP co-injected with both *Nv*Snail-A::mCherry and *Nv*Snail-

B::mCherry mRNA together in zygotes at 40 hpf. White arrowheads indicate AJs. Patched

patterns of cytosolic and nuclear ß-catenin (Blue arrowheads) were observed.

(B) In vivo localization of *Nv*Par-3::mVenus co-injected with both *Nv*Snail-A::mCherry and *Nv*Snail-

B::mCherry mRNA together at 40 hpf. Patched patterns of AJs (White arrowheads) were

observed.

(C) Diagram depicts the suggested interpretation for A and B.

(D) IFS for ß-catenin (ß-cat), *Nv*Par-6, *Nv*aPKC, and *Nv*Par-1 in embryos at 40 hpf where *Nv*Snail-

A::mCherry and *Nv*Snail-B::mCherry mRNA were overexpressed together into a single

ectodermal blastomere lineage (followed by green Streptavidin-TxRed Dextran (Dex).

Arrowheads indicate the absence of the protein, cytosolic ß-cat, and disrupted epithelium.

Arrows indicate bottle-like shape cells. *site of gastrulation.

(E) IFS for ß-cat in embryos at 40 hpf where *Nv*Snail-A::mCherry and *Nv*Snail-B::mCherry mRNA

were overexpressed together into a single blastomere lineage and no affects were observed.

See also Figure S6.

(F) Embryo wide CRISPR/Cas9 knock-out for both *Nvsnail-A* and *Nvsnail-B* at 40 hpf showing that

AJs form in presumptive endomesodermal region similar to ectodermal cells. High

magnification images from boxed region (endomesodermal plate) are shown on the right.

Arrowheads indicate protein localization.

(G) Graphical summary of the observed results with previous published data (Servetnick et al.,

2017).

Morphology is shown by DAPI, Tub, and Phall IFS. Except from 6A and 6B, all images are single

optical sections from the z-stack confocal series. (A) and (B) are 3D reconstructions from a z-stack

confocal series. All scale bars, 20 μm.

Figure 7. The differences between epithelial structure in ectoderm and endomesoderm in _N._

vectensis embryos are due to the lack of mechanisms to segregate a distinct mesoderm.

(A) Diagram depicting key cellular and molecular mechanisms involved during gastrulation of

bilaterian and _N. vectensis_ (a cnidarian) embryos. See also Figure S7.

Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Mark Q. Martindale (mqmartin@whitney.ufl.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Culture and spawning of _Nematostella vectensis_.

Spawning, gamete preparation, fertilization and embryo culturing of _N. vectensis_

(RRID:SCR_005153) embryos was performed as previously described (Hand and Uhlinger, 1992;

Layden et al., 2013; Röttinger et al., 2012; Wolenski et al., 2013). Adult _N. vectensis_ were

cultivated at the Whitney Laboratory for Marine Bioscience of the University of Florida (USA).

Males and females were kept in separate glass bowls (250 ml) in 1/3x seawater (salinity: 12pp)

reared in dark at 16ºC. Animals were fed freshly hatched _Artemia_ 3 times a week and macerated

oyster the day before spawning. Spawning was induced by incubating the adults under an eight-

hour light cycle at 25°C the night before the experiment. Distinct groups of animals were spawned

once every 2 weeks. Oocytes and sperm were collected separately and fertilized _in vitro_ by adding

sperm to egg masses for 25 minutes. The jelly mass surrounding the fertilized eggs was removed

by incubating the eggs in 4% L-Cysteine (in 1/3x seawater; pH 7.4) for 15-17 minutes and then

washed 3 times with 1/3x seawater. De-jellied eggs were kept in glass dishes (to prevent sticking)

in filtered 1/3 seawater at 16ºC until the desired stage.

METHOD DETAILS

Immunohistochemistry

All immunohistochemistry experiments were carried out using the previous protocol for *N.*

vectensis (Salinas-Saavedra et al., 2015) with a slight modification in the glutaraldehyde

concentration to allow better antibody penetration. Embryos were fixed on a rocking platform at

room temperature in two consecutive steps. Embryos of different stages were fixed for no longer

than 3 minutes in fresh Fix-1 (100mM HEPES pH 6.9; 0.05M EGTA; 5mM MgSO4; 200mM NaCl;

1x PBS; 3.7% Formaldehyde; <u>0.2%</u> Glutaraldehyde; 0.5% Triton X-100; and pure water). Then,

Fix-1 was removed and replace with fresh Fix-2 (100mM HEPES pH 6.9; 0.05M EGTA; 5mM

MgSO4; 200mM NaCl; 1x PBS; 3.7% Formaldehyde; <u>0.05%</u> Glutaraldehyde; 0.5% Triton X-100;

and pure water). Embryos were incubated in Fix-2 for 1 hour. Fixed embryos were rinsed at least

five times in PBT (PBS buffer plus 1% BSA and 0.2% Triton X-100) for a total period of 3 hours.

PBT was replaced with 5% normal goat serum (NGS; diluted in PBT) and fixed embryos were

blocked for 1 to 2 hours at room temperature with gentle rocking. Primary antibodies were diluted

in 5% NGS to desired concentration. Blocking solution was removed and replaced with primary

antibodies diluted in NGS. All antibodies incubations were conducted over night on a rocker at 4°C.

After incubation of the primary antibodies, samples were washed at least five times with PBT for a

total period of 3 hours. Secondary antibodies were then applied (1:250 in 5% NGS) and samples

were left on a rocker overnight at 4°C. Samples were then washed with PBT and left on a rocker at

room temperature for an hour. To visualize F-actin, samples were incubated then for 1.5 hours in

Phalloidin (Invitrogen, Inc. Cat. # A12379) diluted 1:200 in PBT. Samples were then washed once

with PBT and incubated with DAPI (0.1µg/µl in PBT; Invitrogen, Inc. Cat. # D1306) for 1 hour to

allow nuclear visualization. Stained samples were rinsed again in PBS two times and dehydrated

quickly into isopropanol using the gradient 50%, 75%, 90%, and 100%, and then mounted in

Murray's mounting media (MMM; 1:2 benzyl benzoate:benzyl alcohol) for visualization. Note that

MMM may wash DAPI out of your sample. For single blastomere microinjection experiments, after

Phalloidin staining, samples were incubated with Texas Red Streptavidin (1:200 in PBT from

1mg/ml stock solution; Vector labs, Inc. Cat.# SA-5006. RRID:AB_2336754) for 1 hour to visualize

the injected dextran. We scored more than 1,000 embryos per each antibody staining and confocal

imaged more than 50 embryos at each stage.

The primary antibodies and concentrations used were: mouse anti-alpha tubulin (1:500; Sigma-

Aldrich, Inc. Cat.# T9026. RRID:AB_477593), rabbit anti-ß-catenin (1:300; Sigma-Aldrich, Inc.

Cat.# C2206. RRID:AB_476831), mouse anti-histone H1 (1:300; F152.C25.WJJ, Millipore, Inc.

RRID:AB_10845941).

Rabbit anti-*Nva*PKC, rabbit anti-*Nv*Lgl, rabbit anti-*Nv*Par-1, and rabbit anti-*Nv*Par-6 antibodies are

custom made high affinity-purified peptide antibodies that were previously raised by the same

company (Bethyl Inc.). All these four antibodies are specific to *N. vectensis* proteins (Salinas-

Saavedra et al., 2015) and were diluted 1:100.

Secondary antibodies are listed in Table S1.

Fluorescent tracer dye penetration assay

Primary polyps were incubated and mounted in 1/3 sea water with fluorescent dextran solution (0.5

mg/ml). For uninjected embryos we used Dextran, Alexa Fluor® 555 (Molecular Probes, INC. Cat.#

D34679). For injected embryos, expressing fluorescent proteins, we used Dextran, Alexa Fluor®

647 (Molecular Probes, INC. Cat.# D22914). Animals were observed within 10 minutes of

incubation. 15 animals were recorded per treatment. For better visualization of the dextran solution

inside the gastric cavity as shown in Figure 2B, we delivered additional dextran solution by

microinjecting dye through the polyp's mouth. For the rest of the experiments, we mainly focused in

the ectodermal permeability and we let the polyps to eat the solution by themselves as grown

babies.

mRNA microinjections

The coding region for each gene of interest was PCR-amplified and cloned into pSPE3-mVenus or

pSPE3-mCherry using the Gateway system (Roure et al., 2007). Eggs were injected directly after

fertilization as previously described (Layden et al., 2013; Salinas-Saavedra et al., 2015) with the

mRNA encoding one or more proteins fused in frame with reporter fluorescent protein (N-terminal

tag) using final concentrations of 450 ng/µl for each gene. Fluorescent dextran was also co-

injected to visualize the embryos. For single blastomere microinjections, we raised the embryos

until 8-16 cell stages (3-4 hpf) and co-injected the mRNA solution with Biotinylated Dextran Amine-

Texas Red (10 µg/µl; Vector labs, Inc. Cat.# SP-1140. RRID:AB_2336249). Live embryos were

kept at 16ºC and visualized after the mRNA of the FP was translated into protein (2-3 hours). To

avoid lethality, lower mRNA concentrations of the mutant proteins (250 ng/µl) were used to image

the specimens for Figures 2 and 4, and Movie S1. Live embryos were mounted in 1/3 sea water for

visualization. Images were documented at different stages from 3-96 hrs. post fertilization. We

injected and recorded more than 500 embryos for each injected protein and confocal imaged

approximately 20 specimens for each stage for detailed analysis of phenotypes *in vivo*. We

repeated each experiment at least five times obtaining similar results for each case. The

fluorescent dextran and primers for the cloned genes are listed in Table S1.

CRISPR/Cas9 knock-outs

To target our gene of interest, we used synthetic guide RNAs (sgRNA; Synthego, Inc.) and

followed the instructions obtained from the manufacturer to form the RNP complex with Cas9

(Cas9 plus sgRNAs). Target sites, off-target sites, and CFD scores were identified and sgRNA

were designed using CRISPRscan (Doench et al., 2014; Moreno-Mateos et al., 2015). We

delivered the RNP complex by microinjection as previously described (Ikmi et al., 2014; Servetnick

et al., 2017; Wijesena et al., 2017). Lyophilized Cas9 (PNA Bio., Inc. Cat.# CP01) was

reconstituted in nuclease-free water with 20% glycerol to a final concentration of 2µg/µl.

Reconstituted Cas9 was aliquoted for single use and stored at -80ºC. Embryos were injected, as

described for mRNA microinjections, with a mixture (12.5µl) containing sgRNAs (80 ng/µl of each

sgRNA), Cas9 (3 µg), and Alexa Fluor 488-dextran (0.2 µg/µl; Molecular Probes, Inc. Cat.#

D22910). Cas9 and sgRNA guides only controls were injected alongside each round of

experiments. sgRNA guides controls are only shown in figures as Cas9 had no significative effects.

3 sgRNA were used to knock out *Nvpar-3*, 3 sgRNA were used to knock out *Nvpar-6*, 6 sgRNA

were used to knock out *Nvsnail-A*, and 6 sgRNA were used to knock out *Nvsnail-B*. Single-embryo

genomic DNA was analyzed as previously described (Servetnick et al., 2017). Gene expression

was confirmed by i*n situ* hybridization. We injected and recorded more than 1000 embryos for each

treatment. We repeated each experiment at least six times obtaining similar results for each case.

sgRNAs' sequences and PCR primers flanking the targeted region are listed in Table S1.

***In situ* hybridization**

In situ hybridization was carried out following a previously published protocol for *N. vectensis*

(Wolenski et al., 2013). Animals were fixed in ice-cold 4% paraformaldehyde with 0.2%

glutaraldehyde in 1/3x seawater for 2 min, followed by 4% paraformaldehyde in PBTw for 1 hour at

4°C. Digoxigenin (DIG)-labeled probes, previously described (Röttinger et al., 2012; Salinas-

Saavedra et al., 2015), were hybridized at 63°C for 2 days and developed with the enzymatic

reaction of NBT/BCIP as substrate for the alkaline phosphatase conjugated anti-DIG antibody (

Roche, Inc. Cat.#11093274910. RRID:AB_514497). Samples were developed until gene

expression was visible as a purple precipitate.

Drug treatment

We incubated *N. vectensis* embryos in 20µM of aPKC pseudosubstrate inhibitor (Protein kinase Cζ

pseudosubstrate, myristoyl trifluoroacetate salt, Sigma, Cat.#P1614) from 0 to 4 hpf. Controls and

1-azakenpaullone (AZ; Sigma, Cat.#A3734) drug treatment of *N. vectensis* embryos was

performed as previously described (Leclère et al., 2016; Röttinger et al., 2012). Embryos were

developed in 5µm AZ from 3 to 76 hpf. Controls were incubated in 0.08% DMSO.

Imaging of *N. vectensis* embryos

Images of live and fixed embryos were taken using a confocal Zeiss LSM 710 microscope using a

Zeiss C-Apochromat 40x water immersion objective (N.A. 1.20). Pinhole settings varied between

1.2-1.4 A.U. according to the experiment. The same settings were used for each individual

experiment to compare control and experimental conditions. Results from *in situ* hybridization

studies were imaged using a Zeiss Imager.M2 with a Zeiss 425 HRc color digital camera run by

Zeiss Zen 2012 software. Z-stack images were processed using Imaris 7.6.4 (Bitplane Inc.)

software for three-dimensional reconstructions and FIJI for single slice and movies. Final figures

were assembled using Adobe Illustrator and Adobe Photoshop.

Supplemental Information

Figure S1. Related to Figure 1. Epidermal and gastrodermal cells are joined by different set of

junctional complexes.

(A) TEM micrographs of *N. vectensis* primary polyps. Epidermal cells (ectodermally derived) are joined

most likely by AJs (Red arrowheads). Gastrodermal cells (endomesodermally derived) are

interconnected by fewer and shorter contacts, most likely by septate junctions (yellow arrowheads). a

and b: two different types of epidermal cells. Note the ectodermally-derived cnidocyte in b. c and d:

two different types of gastrodermal cells. b' and d' are high magnification images from boxed region in

b and d, respectively. gc: gastric cavity. Scale bars in a, b, c, and d: 500 nm. Scale bars in b' and d': 200

nm.

(B) *In vivo* localization of *Nv*ß-catenin::GFP after 36 hpf in *N. vectensis* embryos. Arrowheads indicate the

cortical localization of *Nv*ß-catenin::GFP (AJs) in the ectoderm that was not observed in the

endomesoderm. High magnification image from boxed region is shown on the right. Scale bars: 20

μm.

(C) Immunofluorescent staining for *Nv*Par-6 and *Nv*Par-1(red) at late planula and polyp stages show that

both Par proteins are absented from the endomesoderm. Phalloidin is shown in gray. Histone and

tubulin antibody staining are shown in Figure 1B and 1C as counterstain to show the penetrability in

the fixed tissue. Scale bars: 20 μm.

Figure S2. Related to Figure 2 and 3. Disruption of the aPKC/Par complex in *N. vectensis* embryos.

(A) Diagram depicting the modifications made to *Nv*Par-1, *Nv*Par-6, and *Nv*Par-3 sequence to generate

the dominant negative version of each protein (dn*Nv*Par-1, dn*Nv*Par-6, and dn*Nv*Par-3, respectively),

which lack the putative interaction domain with *Nv*aPKC. In the right, a diagram depicts the

localization of *Nv*Par-1, *Nv*aPKC, *Nv*Par-6, and *Nv*Par-3 proteins in epithelial cells. The putative

interaction with *Nv*aPKC, restricts the localization of *Nv*Par-6 and *Nv*Par-3 strictly to the apical cortex

of the cell, and *Nv*Par-1 to the lateral cortex of the cell. See also Figure S3.

(B) Immunofluorescent staining for Tubulin and Phalloidin at gastrula stage of embryos expressing

dn*Nv*Par-6, and dn*Nv*Par-3. The overexpression of either dn*Nv*Par-6::mVenus or dn*Nv*Par-3::mVenus

induced phenotypes where the endomesodermal cells (yellow arrows) are disorganized during

gastrulation. We observed a disorganized endoderm formed by 1) cells with fibroblast-like

morphologies, 2) stellate shaped mesenchymal-like cells, or 3) a mass of round dead cells. Cell

morphotypes are outlined in red and their schematic representation is presented below them. The

penetrance of the obtained phenotypes when either dn*Nv*Par-6::mVenus (blue) or dn*Nv*Par-3::mVenus

(magenta) are overexpressed is indicated for each case.

(C) Immunofluorescent staining for Tubulin and Phalloidin at polyp stage of embryos expressing dn*Nv*Par-

6, and dn*Nv*Par-3. We observed 'Endoderm-less' polyps from phenotype 3: absence of an organized

endoderm, tentacles or complete mesenteries in injected animals that survived to this stage (2 weeks

post fertilization). A recognizable ectodermal pharynx (yellow arrowheads) was detected in some of

these polyps.

(D) The position of the sgRNAs (red) and primers (green) used for the PCR assay are shown on the diagram

depicting the genomic sequence of *Nvpar-6*. Note the absence of fragments of *Nvpar-6* (arrow)

resulting from CRISPR/Cas9 mediated mutagenesis. The presence of other bands suggests mosaicism.

Black rectangles correspond to *Nvpar-6* exon. PF: primer forward. PF: primer reverse.

(E) The position of the sgRNAs (red) and primers (green) used for the PCR assay are shown on the diagram

depicting the genomic sequence of *Nvpar-3*. Note the absence/truncation of fragments of *Nvpar-3*

(arrow) resulting from CRISPR/Cas9 mediated mutagenesis. The presence of other bands suggests

mosaicism as shown in Figure S3E. Blue rectangles correspond to *Nvpar-3* exon. PF: primer forward.

PF: primer reverse.

Figure S3. Related to Figure 3. CRISPR/Cas9 mediated mutagenesis of *Nvpar-3.* Morphology is shown

by DAPI, Tubulin, and Phalloidin staining. *: site of gastrulation is up.

(A) Different ectodermal thickness (yellow) observed in CRISPR/Cas9 mutants. Control and

affected (thick and thin) epithelia were aligned at the base of the ectoderm (red line) for better

visualization.

(B) Immunofluorescent staining for cytoskeleton and *Nv*Par-6 in CRISPR/Cas9 *Nvpar-6* knock-out

embryos (mosaic phenotype). Arrowheads indicate the absence of *Nv*Par-6. High magnification

images are shown on the right. The cytoskeleton is apically organized only where *Nv*Par-6 is

apically localized.

(C) Diagram depicts the role of aPKC/Par complex (green) on cytoskeleton.

(D) Immunofluorescent staining for ß-catenin (ß-cat) in single injected-blastomere control and

CRISPR/Cas9 *Nvpar-3* knock-outs at 40 hpf. Streptavidin-Biotin TxRed Dextran (Dex) is shown

in green. Arrowheads indicate the absence of the protein and disrupted epithelium. Arrows

indicate bottle-like shape cells. Note the displacement of nuclei (yellow arrow) and changes in

tubulin (Tub) staining.

(E) *In situ* hybridization of *Nvpar-3* knockout embryos (Cas9 and gRNAs) compared with control embryos

at 40 hpf. The disruption of the *N. vectensis* Par/aPKC complex modified the morphology but did not

modify the cell-fate specification of endomesodermal cells.

Figure S4. Related to Figure 3 and S2. *In vivo* localization of dn*Nv*Par-6 and dn*Nv*Par-3 proteins at

different embryonic stages.

Images of the whole embryo correspond to a 3D reconstruction from a z-stack series. Side panels are a

single optical section from the z-stack series. An aboral view is shown for all gastrula stages. Yellow arrows

indicate the apico-lateral cortex labeled with Lifeact::mTq2. Scale bars: 10µm.

(A) *In vivo* localization of *Nv*Par-6::mVenus and dn*Nv*Par-6::mVenus at cleavage, blastula, and gastrula

stages. *Nv*Par-6::mVenus distributes uniformly at the apical region of the cell but displays a scattered

pattern. However, dn*Nv*Par-6::mVenus displays stronger cortical localization due to its interaction

with *Nv*Cdc42 at the apical and apico-lateral cortex of the cells. This was confirmed by their co-

expression with *Nv*Cdc42::mCherry. White arrowheads indicate the scattered localization of *Nv*Par-

6::mVenus. White arrows indicate the cortical and stronger localization of dn*Nv*Par-6::mVenus.

(B) *In vivo* localization of Lifeact::mTq2 in *Nv*Par-6::mVenus and dn*Nv*Par-6::mVenus expressing embryos

shown in (A) at cleavage, blastula, and gastrula stages. The actin cytoskeleton was also affected by the

overexpression of dn*Nv*Par-6::mVenus.

(C) *In vivo* co-distribution of *Nv*Cdc42::mCherry with *Nv*Par-6::mVenus at blastula and gastrula stages.

(D) I*n vivo* co-distribution of *Nv*Cdc42::mCherry with dn*Nv*Par-6::mVenus at blastula and gastrula stages.

*Nv*Cdc42::mCherry localization is also affected when dn*Nv*Par-6::mVenus is overexpressed.

(E) Graphical summary of the observed results for *Nv*Par-6::mVenus and dn*Nv*Par-6::mVenus.

(F) *In vivo* localization of *Nv*Par-3::mVenus and dn*Nv*Par-3::mVenus at cleavage, blastula, and gastrula

stages. dn*Nv*Par-3 displays broader localization, resembling the localization of *Nv*Par-6 and indicating

its release from AJs (compare with *Nv*Par-3). White arrowheads indicate the punctate localization of

*Nv*Par-3::mVenus. White arrows indicate the broader localization of dn*Nv*Par-3::mVenus.

(G) *In vivo* localization of Lifeact::mTq2 in *Nv*Par-3::mVenus and dn*Nv*Par-3::mVenus expressing embryos

shown in (F) at cleavage, blastula, and gastrula stages. The actin cytoskeleton was also affected by the

overexpression of dn*Nv*Par-3::mVenus.

(H) *In vivo* co-distribution of *Nv*Cdc42::mCherry with *Nv*Par-3::mVenus at blastula and gastrula stages.

(I) I*n vivo* co-distribution of *Nv*Cdc42::mCherry with dn*Nv*Par-3::mVenus at blastula and gastrula stages.

*Nv*Cdc42::mCherry localization is not affected when dn*Nv*Par-3::mVenus is overexpressed.

(J) Graphical summary of the observed results for *Nv*Par-3::mVenus and dn*Nv*Par-3::mVenus.

1013

Figure S5. Related to Figure 6. *Nvsnail-A and Nvsnail-B* together regulate AJs.

*site of gastrulation.

(A) Immunofluorescent staining of embryos at 40 hpf where *Nv*Snail-A::mCherry and *Nv*Snail-

B::mCherry mRNA were overexpressed together into a single blastomere lineage (followed by

green Streptavidin-TxRed Dextran (Dex). No affects were observed in approximately 1/8 of the

injected embryos. Arrowheads indicate the disrupted epithelium. Arrows indicate bottle-like

shape cells.

(B) Embryo wide CRISPR/Cas9 knock-out for both *Nvsnail-A* and *Nvsnail-B* at 40 hpf showing

embryos that did not gastrulate and retain ß-catenin (ß-cat) and *Nv*aPKC at the apical cell-

cortex of the endomesodermal plate (arrowheads).

(C) Embryo wide CRISPR/Cas9 knock-out for both *Nvsnail-A* and *Nvsnail-B* at 40 hpf showing

embryos that developed a pharynx (arrow) but not an organized endomesoderm.

(D) Embryo wide CRISPR/Cas9 knock-out for both *Nvsnail-A* and *Nvsnail-B* at 2 weeks hpf

showing embryos that did not developed a gastrodermal epithelium.

1028

Figure S6. Related to Figure 6. CRISPR/Cas9 mediated mutagenesis of *Nvsnail-A and Nvsnail-B*.

(A) The position of the sgRNAs (red) and primers (green) used for the PCR assay are shown on the diagram

depicting the genomic sequence of *Nvsnail-A*. Note the absence of fragments of *Nvsnail-A* (arrow)

resulting from CRISPR/Cas9 mediated mutagenesis. The presence of other bands suggests mosaicism.

Black rectangles correspond to *Nvsnail-A* exon. Blue double arrow depicts UTR regions. PF: primer

forward. PF: primer reverse.

(B) The position of the sgRNAs (red) and primers (green) used for the PCR assay are shown on the diagram

depicting the genomic sequence of *Nvsnail-B*. Note the absence of fragments of *Nvsnail-B* (arrow)

resulting from CRISPR/Cas9 mediated mutagenesis. The presence of other bands suggests mosaicism

as shown in (C). Black rectangles correspond to *Nvsnail-B* exon. Blue double arrow depicts UTR

regions. PF: primer forward. PF: primer reverse.

(C) *In situ* hybridization of *Nvsnail-A+Nvsnail-B* knockout embryos (Cas9 and gRNAs) compared with

control embryos at 40 hpf. *Nvsnail-A+Nvsnail-B* knockout embryos display no expression and mosaic

expression of *Nvsnail-A* and *Nvsnail-B,* but did not modify the cell-fate specification of ectodermal

cells.

1044

Figure S7. Related to Figure 7. Suggested model for mesoderm specification in Metazoa.

(A) Differential regulation of cell adhesion by the endomesoderm GRN is mediated by changes in cell

polarity that regulate ß-catenin localization. These mechanisms emerged at the Bilateria + Cnidaria

node.

(B) Ctenophores do not possess a full complement of cell adhesion, cell-polarity, and endomesodermal

GRN components present in the most common ancestor between Cnidaria and Bilateria. However,

ctenophores possess a distinct mesoderm, suggesting the emergence of different mechanisms to

segregate mesoderm in Metazoa.

1053

1054

1055   **Movie S1. Related to Figure 4. EMT-like process occurs in dn*Nv*Par-3::mVenus expressing cells.**

1056   Lifeact::mTq2 is shown in grey. Arrows indicate the apical constriction of detaching cells. Scale bars:10µm

A

*Nv*ß-Catenin | *Nv*ß-Catenin + *Nv*Snail::mCherry

GFP | GFP

B

*Nv*Par-3 | *Nv*Par-3 + *Nv*Snail::mCherry

mVenus | mVenus

C

Ectoderm: Without Snail

When Snail is not functional, Cadherins are stabilized, and ß-catenin translocates from the cytoplasm to AJs.

Endomesoderm: With Snail

When Snail is expressed, it destabilizes Cadherins. ß-catenin translocates from the adherens junctions to the cytoplasm and nucleus.

D | Single blastomere injection *Nv*Snail::mCherry

Dex DAPI | ß-cat | Phall | 84/134

*Nv*Par-6

*Nv*aPKC

*Nv*Par-1

E

Dex DAPI | ß-cat | Phall | 17/134

F | Control/Guides only

DAPI Phall | ß-cat / Tub | 185/200

CRISPR/Cas9 *Nvsnail* genes

ß-cat | 95/200

*Nv*Par-6

*Nv*aPKC

*Nv*Par-1

*Nv*Lgl

G

Nvbrachyury → Ectoderm epiboly → Pharynx formation

Nuclear ß-cat

aPKC/Par AJs

Nvsnail-A+B → Endomesoderm → Migration & Zippering

Invagination | Migration | Zippering
24 hpf | 30 hpf | 48 hpf

A

Bilateria

Invagination — Molecular & tissue Segregation — Migration — Triploblasty: Distinct mesoderm

Animal — Vegetal

Endomesoderm

cWNT & WNT/PCP nuclear ß-catenin → **Brachyury** → **Snail**

AJs ⇄ aPKC/Par ⇄ Cytoskeleton

Segregation by **EMT** Integrin-based junctions → Mesoderm / Endoderm

time

Cnidaria
N. vectensis

cWNT nuclear ß-catenin → **Brachyury** --?--> **Snail**

? → GSK-3ß ⊣ AJs ⇄ aPKC/Par ⇄ Cytoskeleton

Partial-EMT? Integrin-based junctions? → EndoMesoderm

Ectoderm — Epithelial integrity — ECM?

Invagination — Molecular Segregation NO tissue segregation — Migration — Diploblasty: Dual endomesoderm

Animal — Vegetal

Porifera Ctenophora **?**