1 Title: S1PR3 mediates inflammatory pain and itch via distinct TRP channel-
2 dependent pathways
3
4 Abbrev. Title: S1PR3 mediates inflammatory pain and itch
5

6 Authors and Affiliations: Rose Z. Hill[1], Takeshi Morita[1#], Rachel B. Brem[2, 3], Diana M.
7 Bautista[1, 4*]
8
1. Department of Molecular and Cell Biology, University of California, Berkeley, CA
94720, USA.
2. Department of Plant and Microbial Biology, University of California, Berkeley, CA
94720, USA.
3. Buck Institute for Research on Aging, Novato, CA 94945, USA.
4. Helen Wills Neuroscience Institute, University of California, Berkeley, CA 94720,
USA.
Current Address: The Rockefeller University, New York, NY 10065, USA.
*   Correspondence should be addressed to dbautista@berkeley.edu

18
19
20
21 Number of pages: 36
22
23 Number of figures: 7
24
25 Number of tables: 0
26
27 Number of words:
28
29   Abstract: 210
30   Introduction: 641
31   Discussion: 824
32
33 Conflict of interest: The authors declare no competing financial interests.
34

35 Acknowledgements: We would like to thank all members of the D.M.B laboratory for
36 constructive feedback and criticism.

Abstract

In this work, we establish a role for the bioactive lipid sphingosine 1-phosphate (S1P) and its receptor S1PR3 in inflammatory thermal pain and itch. S1P injection elicited robust heat hypersensitivity, acute pain, and itch behaviors in mice, which were wholly dependent on S1PR3. S1P-evoked pain was entirely dependent on TRPV1 ion channels, whereas S1P-evoked itch was dependent on TRPA1. Consistent with our *in vivo* findings, we found that S1P activated subsets of both heat- and itch-sensitive dorsal root ganglion (DRG) neurons, and that activation was solely dependent on S1PR3, and not S1PR1. Our data support a model in which S1P elicits neuronal activation via two pathways in distinct subsets of somatosensory neurons. Gβγ-dependent activation of TRPA1 mediates S1PR3-evoked calcium responses in one subpopulation, whereas PLC-dependent activation of TRPV1 mediates S1PR3-evoked calcium responses in the other subpopulation. In the Complete Freund's Adjuvant (CFA) -induced inflammatory pain model, S1PR3-deficient mice failed to develop heat hypersensitivity, whereas mechanical hypersensitivity developed normally. Pharmacological blockade of S1PR3 activity or S1P production rapidly and selectively ameliorated heat hypersensitivity in wild-type mice treated with CFA. These data definitively identify S1PR3 as a mediator of thermal pain and itch, and, furthermore, demonstrate for the first time that elevated peripheral S1P is an essential driver of chronic inflammatory pain.

Significance Statement

Chronic pain is a major health problem with few effective treatments. Here, we show that the pro-inflammatory lipid, S1P, and its receptor S1PR3 are essential for thermal hypersensitivity and acute itch. Our results suggest that S1PR3 inhibitors or S1P modulators may be of interest as powerful and heretofore untested treatments for inflammatory pain and itch.

Introduction

Hypersensitivity to thermal and mechanical stimuli is a hallmark of inflammatory pain. During

inflammation, immune cells, skin cells, and neurons release a slew of small molecules, lipids,

and proteins into the extracellular milieu collectively referred to as the "inflammatory soup" (Ji,

Xu, & Gao, 2014). A handful of these factors are known to directly modulate the activity of

peripheral sensory neurons, resulting in thermal and/or mechanical hypersensitivity (Chuang et

al., 2001; Tominaga, Wada, & Masu, 2001). While several transduction channels and sensory

neuronal populations mediating acute pain and itch have been identified (Basbaum, Bautista,

Scherrer, & Julius, 2009; Meixiong & Dong, 2017; Pinho-Ribeiro, Verri, & Chiu, 2017), the

molecular factors that drive thermal and mechanical hypersensitivity under inflammatory states

remains enigmatic (Ghasemlou, Chiu, Julien, & Woolf, 2015). For example, it is known that mice

lacking TRPV1 fail to develop thermal but not mechanical sensitization in a model of chronic

inflammatory pain (Caterina et al., 2000), yet the physiologically relevant molecular pathways

upstream of TRPV1 sensitization are not fully understood. And likewise, while the immune cells

that promote mechanical hypersensitivity in chronic inflammatory pain have been identified, the

signaling factors that promote that sensitization remain unclear (Ghasemlou et al., 2015).

Sphingosine 1-phosphate (S1P) is a bioactive sphingolipid associated with asthma, neuropathic

pain, inflammatory bowel disease, and psoriasis (Checa et al., 2015; Donoviel et al., 2015; Hou

& Fang, 2015; Myśliwiec et al., 2016; Patti et al., 2008; Rivera, Proia, & Olivera, 2008; F.

Roviezzo et al., 2015). S1P signaling via S1P Receptor 1 (S1PR1) facilitates the migration,

recruitment, and activation of immune cells (Cyster & Schwab, 2012; Matloubian et al., 2004).

Intriguingly, expression of both S1PR1 and S1PR3 has been reported in the somatosensory

ganglia (Camprubi-Robles et al., 2013; Usoskin et al., 2015), and S1P modulates sensory

neuronal excitability (Li, Li, Kays, Guerrero, & Nicol, 2015; Wang et al., 2017). However, the role

of S1P signaling in somatosensation remain unclear.

In this study, we elucidate the molecular mechanisms of S1P signaling in pain and itch. While

the subject of a number of previous studies, whether and how S1P signaling directly mediates

acute and inflammatory pain remains enigmatic for several reasons (Camprubi-Robles et al.,

2013; Canlas et al., 2015; Finley et al., 2013; Janes et al., 2014; Mair et al., 2011; Weth et al.,

2015). First, and foremost, the S1P receptors in sensory ganglia involved in S1P-evoked heat

hyperalgesia (Mair et al., 2011) and spontaneous pain behaviors (Camprubi-Robles et al., 2013)

have remained controversial. While both S1PR1 and S1PR3 have been implicated in

nociception, the aforementioned studies showed incomplete loss of S1P-evoked behaviors or

partial phenotypes. Localization studies were performed with anti- S1P receptor antibodies that

stain S1PR3-deficient sensory ganglia. As such, the sensory neuronal subtype(s) that

participate in S1P signaling are unknown. Additionally, these studies were cursory and did not

probe the downstream signaling pathways activated by S1P receptor(s) in sensory neurons.

These studies also employed excessively high concentrations of S1P in behavioral experiments,

two to three orders of magnitude higher than levels reported in mammals (0.1 nM – 10 µM), and

beyond the solubility limits of this lipid in saline (micromolar). Finally, no study has examined the

role of S1P signaling in itch sensation.

Here, we aimed to perform a rigorous and systematic characterization of the behavioral effects

and underlying mechanisms of S1P signaling in acute pain and itch and chronic inflammatory

pain. We have found that elevated S1P triggers acute pain, heat hypersensitivity, and,

surprisingly, itch behaviors that are wholly dependent on S1PR3. Intriguingly, S1PR3-deficient

mice do not develop thermal hypersensitivity in the CFA model of inflammatory pain. In addition,

acute pharmacological blockade of peripheral S1PR3 signaling or S1P production reverses

CFA-evoked thermal hypersensitivity. Our findings define new roles for S1PR3 signaling in pain

and itch and show that S1P itself is an essential contributor to inflammatory pain in the

periphery.

Materials and Methods

Behavioral studies & mice

$S1P_3^{mcherry/+}$ (Sanna et al., 2015) and $S1pr3^{-/-}$ (Kono et al., 2004) mice were obtained from

Jackson Laboratory and backcrossed to C57bl6/J, $Trpv1^{-/-}$ and $Trpa1^{-/-}$ mice were described

previously (Bautista et al., 2006; Caterina et al., 2000) and $Trpv1^{-/-}/Trpa1^{-/-}$ mice were bred from

crossing $Trpv1^{-/+}$ and $Trpa1^{-/+}$ mice. Wherever possible, wild-type (for *Trpa1* and *Trpv1* behavior)

or wild-type/heterozygous (*S1pr3*) littermate controls were used in behavioral experiments. Mice

(20–30 g) were housed in 12 h light-dark cycle at 21°C. Mice were singly housed one week prior

to all behavioral experiments and were between 8-10 weeks at the time of the experiment. All

mice were acclimated in behavioral chambers on the 2 subsequent days for at least 1 hour prior

to treatment for itch/pain behavior and radiant heat. Male mice were used for all behavioral

studies, due to the effects of the female estrus cycle on itch and pain behaviors.

Itch and acute pain behavioral measurements were performed as previously described (Wilson

et al., 2011). Mice were shaved one week prior to itch behavior. Compounds injected: 200 nM, 2

µM, 10 µM S1P (Tocris, Avanti Polar Lipids), 500 µM TY 52156 (Tocris), and 50 µM SKI II

(Tocris) in PBS with either DMSO- or Methanol-PBS vehicle controls. Pruritogens were injected

using both the neck/rostral back model (20 µl), and the cheek model (20 µl) of itch, as

previously described (Wilson et al., 2011). Behavioral scoring was performed while blind to

experimental condition and mouse genotype. All scratching and wiping behavior videos were

recorded for 1 hour and scored for either the first 30 minutes (scratching) or the first five minutes

(wiping). Bout number, time spent scratching, and bout length were recorded for scratching

behavior. Wiping was recorded as number of wipes.

For radiant heat hypersensitivity behavior, S1P was injected intradermally into the plantar

surface of the hindpaw (20 µl). Radiant heat paw withdrawal latencies, before and after

application of compound or vehicle were performed as previously described (Morita et al., 2015;

Tsunozaki et al., 2013). Radiant heat assays were performed using the Hargreaves test system

(IITC Life Science). Mice were injected with compound of interest into the hind paw, and their

paw withdrawal latencies were measured 15 min pre- and 20 min-30 min post-injection. Valid

responses for radiant heat included fast paw withdrawal, licking/biting/shaking of the affected

paw, or flinching. Mice were allowed to acclimate on platform for 1 hour before injection.

For the CFA model of chronic inflammatory pain, 15 µl of CFA (Sigma-Aldrich) or PBS control

was injected into either the left or right hindpaw of 8-9 week old mice briefly anesthetized with

2% isoflurane. Experimenter was blind to genotype. Von Frey paw withdrawal latencies were

recorded as previously described (Tsunozaki et al., 2013) using the up-down method (Chaplan,

SR, Bach, FW, Pogrel, JW, Chung, JM, Yaksh, 1994; Dixon, 1980) on a mesh grate (IITC Life

Science) with calibrated filaments (Touch Test). Radiant heat withdrawal latencies were

measured at baseline, 24-, and 48-hours post-injection for both paws. Paw swelling was

measured using calipers while experimenter was blind to genotype. Photographs were taken

after mice were euthanized using CO_2 while photographer was blinded to genotype.

All behavior experiments were carried out using age-matched or littermate cohorts of male mice.

Mice were tested in 4-part behavior chambers (IITC Life Science) with opaque dividers (Tap

Plastics). Itch and acute pain behavior was filmed from below using high-definition cameras. All

experiments were performed under the policies and recommendations of the International

Association for the Study of Pain and approved by the University of California, Berkeley Animal

Care and Use Committee.

Analysis of gene expression vs. itch correlation data

A previous study examined the correlation of transcript expression of individual genes in dorsal

root ganglion (DRG) neurons with itch behavior among BXD mouse strains (Morita et al., 2015).

One of the top correlated genes from the screen was *Fam57b* (r = 0.57), a recently identified

ceramide synthase (Yamashita-Sugahara et al., 2013) and component of the S1P pathway that

is robustly expressed in somatosensory neurons (Gerhold et al., 2013). To assess whether

expression of the S1P pathway genes as a group was correlated with somatosensory behaviors

across the mouse strains of the BXD population, we first tabulated the absolute value of the

Pearson's correlation r between expression of each S1P pathway gene in turn (*Fam57b,*

Ppargc1a, Spns1, Spns2, Sphk2, S1pr3, S1pr1, Esrrb, Esrrg, Lrp2) and itch behavior (Morita et

al., 2015), and calculated the median of these correlation values, r_{true}. We then drew 10 random

genes from the set of all 16,220 genes with detected expression and computed the median

correlation as above using this null set, r_{null}. Repeating the latter 10,000 times established a null

distribution of median correlations; we took the proportion of resampled gene groups that

exhibited ($r_{true} \geq r_{null}$) as an empirical p-value reporting the significance of enriched correlation

between expression and itch in the genes of the S1P pathway.

In situ hybridization (ISH)

Fresh DRG were dissected from 8-12 week old mice, flash frozen in OCT embedding medium,

and sectioned at 14 µm onto slides. ISH was performed using Affymetrix Quantigene ViewISH

Tissue 2-plex kit according to manufacturer's instructions with Type 1 and Type 6 probes. The

following probes against mouse mRNAs were created by Affymetrix and used for ISH: *S1pr3,*

Mrgpra3, Trpv1, and *Trpa1.* Slides were mounted in Fluoromount with No. 1.5 coverglass.

Imaging of ISH experiments and all other live- and fixed-cell imaging was performed on an

Olympus IX71 microscope with a Lambda LS-xl light source (Sutter Instruments). Images were

analyzed using ImageJ software. Briefly, DAPI-positive cells were circled and their fluorescence

intensity (AFU) for all channels was plotted against cell size using IgorPro software. Co-labeling

analysis was performed using ImageJ. Intensity thresholds were set based on the negative

control staining slide. Cells were defined as "co-expressors" if their maximum intensities were

greater than the threshold for both the Type 1 and Type 6 probe channels.

Cell culture

Cell culture was carried out as previously described (Wilson et al., 2011). Briefly, neurons from

dorsal root ganglia of 2-8 week old male and female mice, or trigeminal ganglia from P0-P4

neonates were dissected and incubated for 10 min in 1.4 mg ml−1 Collagenase P (Roche) in

Hanks calcium-free balanced salt solution, followed by incubation in 0.25% standard trypsin

(vol/vol) -EDTA solution for 2 min with gentle agitation. Cells were then triturated, plated onto

Poly D-Lysine coated glass coverslips and used within 20 h. (media: MEM Eagle's with Earle's

BSS medium, supplemented with 10% horse serum (vol/vol), MEM vitamins,

penicillin/streptomycin and L-glutamine).

Electrophysiology

Current clamp experiments were carried out as previously described (Wilson et al., 2011).

Briefly, gap-free current-clamp recordings were collected at 5 kHz and filtered at 2 kHz

(Axopatch 200B, pClamp software). Electrode resistance ranged between 1.5–5 MΩ. Internal

solution contained 140 mM KCl, 5 mM EGTA and 10 mM HEPES (pH 7.4). Bath solution was

physiological Ringer's solution. The pipette potential was canceled before seal formation.

Experiments were carried out only on cells with a series resistance of less than 30 MΩ. Analysis

of electrophysiology data was performed in pClamp and IgorPro.

Calcium imaging

Ca^{2+} imaging experiments were carried out as previously described (Wilson et al., 2011). DRG

neurons from 2-8 week old male and female mice were used for all experiments. Where

indicated, TG neurons were from P0-P4 neonates. When experiments were performed on non-

wild-type mice, an age-matched wild-type control was also prepared and imaged the same day.

Cells were loaded for 60 min at room temperature with 10 µM Fura-2AM supplemented with

0.01% Pluronic F-127 (wt/vol, Life Technologies) in a physiological Ringer's solution containing

(in mM) 140 NaCl, 5 KCl, 10 HEPES, 2 CaCl2, 2 MgCl2 and 10 D-(+)-glucose, pH 7.4. All

chemicals were purchased from Sigma. Acquired images were displayed as the ratio of 340 nm/

380 nm. Cells were identified as neurons by eliciting depolarization with high potassium Ringer's

solution (75 mM) at the end of each experiment. Responding neurons were defined as those

having a > 15% increase from baseline ratio. Image analysis and statistics were performed

using automated routines in Igor Pro (WaveMetrics). Fura-2 ratios were normalized to the

baseline ratio F340/F380 = (Ratio)/(Ratio t = 0).

Flow cytometry of CFA-treated paws

CFA injections were performed as described above. Briefly, hindpaw skin and underlying fascia

of treated and PBS-injected paws were removed from freshly euthanized mice. Skin was placed

in RPMI media (Gibco) on ice before mincing with dissection scissors. Digestions were

performed for 90 minutes at 37°C on a rotating platform in 1 mL RPMI supplemented with

1:1000 DNaseI enzyme (Roche) and 1 unit LiberaseTM (Roche). Skin was then filtered through

70 µm nylon mesh (Falcon), washed in RPMI, and resuspended in PBS for Aqua Live-Dead

staining. Samples were then transferred to FACS buffer (PBS with 0.5% FBS and 2 mM EDTA),

blocked, then surface stained with the following antibodies: cKit-Biotin, FceRI-PE, CD49b-

PECy7, SiglecF-APC, SA-FITC, Ly6C-PerCP, CD11b-PB, Ly6G-780, CD45.2-AF700.

Compensation tubes (single-stain) were prepared for each fluorophore using positive and

negative control beads. A spleen from a wild-type animal was also prepared by crushing

between frosted glass slides, straining through 70 µm nylon mesh, and lysing of erythrocytes in

ACK (Gibco). A portion of spleen sample was heat-killed for 10 minutes at 65°C and stained

with Aqua viability stain and set aside. The rest of the spleen was stained normally with the

other skin samples. Samples were then run through a flow cytometer (BD Fortessa). Data were

analyzed using FlowJo (Prism) and Microsoft Excel. Neutrophils were defined as live single cells

with the following staining profile: $CD45.2^+/CD11b^+/Ly6G^+/Ly6C^+$. Inflammatory monocytes were

defined as the following: $CD45.2^+/CD11b^+/Ly6G^-/Ly6C^{high}$. Total number of immune cells was

reported since neither genotype differed significantly between total number of live cells or total

number of CD45.2+ immune cells.

Experimental design & statistical analyses

All statistical analyses were performed using IgorPro software or Prism. Values are reported as

the mean ± SEM where multiple experiments are reported, and mean ± SD where one

experiment was performed with multiple wells (for calcium imaging) or mice (for behavior). For

comparison between two groups, Student's unpaired 2-tailed t-test was used. For single-point

comparison between > 2 groups, a one-way ANOVA followed by the Sidak's, Dunnett's or

Tukey's post hoc tests (where appropriate) were employed. For the time course comparison

between 2 groups, 2-way ANOVA was used and single comparison p-values were derived using

Sidak's multiple comparisons. For comparing distributions, a type 2 Kolmogorov-Smirnov test

was used. Number of mice or samples required to attain significance was not calculated

beforehand and was based on numbers used in similar behavioral studies. Significance was

labeled as: n.s., not significant, $p \geq 0.05$; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

Results

Our group previously harnessed natural variation in somatosensory behaviors among genetically distinct mouse strains to identify candidate transducers in DRG neurons (Morita et al., 2015). Analysis of this dataset revealed the sphingosine 1-phosphate (S1P) signaling pathway (Fig. 1a) had transcript expression that was significantly correlated with increased itch behaviors (see Methods; $p = 0.0116$; Fig. 1b). We focused our study on S1PR3 because it was the most highly expressed S1P receptor in the dorsal root ganglia (DRG) (Morita et al., 2015; Usoskin et al., 2015).

S1PR3 is the sole mediator of acute S1P-evoked pain and itch

A handful of previous studies have suggested that S1P may promote acute pain and/or thermal sensitization via various S1PRs in sensory neurons, neutrophils, and/or platelets (Camprubi-Robles et al., 2013; Finley et al., 2013; Mair et al., 2011; Weth et al., 2015). Here, we sought to address the specific contribution of S1PR3 signaling to S1P-evoked somatosensory behaviors using physiologically relevant concentrations of S1P (Checa et al., 2015; Myśliwiec et al., 2016; Sanna et al., 2004).

We first used the cheek assay, which allows for simultaneous discrimination of pain-evoked wiping and itch-evoked scratching (Shimada & LaMotte, 2008). We observed that S1P injection resulted in wiping behaviors in response to S1P concentrations at or above 2 µM in the first five minutes after injection (Fig. 2a). We also found that S1P elicited robust scratching behaviors (Fig. 2b) that developed with an average latency of 4 minutes 29 seconds ± 1 minute 37 seconds and persisted for at least 30 minutes. Scratching was elicited in response to a wider range of S1P concentrations than wiping (0.2-10 µM versus 2-10 µM). Furthermore, we did not observe scratching behaviors in response to the lowest tested dose, 50 nM S1P (data not shown). For all subsequent S1P scratching behavioral assays where wiping was not quantified,

we injected 0.2 µM S1P into the rostral back and observed an average scratching time of 55.5 ±

7.6 seconds (N = 29 animals), which was significantly greater than scratching evoked by vehicle

(0.1% methanol-PBS; 11.1 ± 3.3 seconds, N = 23 animals). Our results from the cheek and

rostral back models are consistent with a role for S1P in promoting both acute pain and itch.

Consistent with a previous study (Finley et al., 2013), we observed robust heat hypersensitivity

after injection of S1P (Fig. 2c). However, the concentration at which we observed

hypersensitivity, 10 µM, is at least 20-fold lower than sensitizing or pain-inducing concentrations

previously reported in the literature (Camprubi-Robles et al., 2013; Finley et al., 2013; Mair et

al., 2011). Mechanical hypersensitivity was not observed (data not shown).

We next tested the requirement of S1PR3 to S1P-evoked itch and pain behaviors. *S1pr3* [-/-] mice

displayed significantly attenuated S1P-evoked scratching indistinguishable from vehicle injection

(10.6 ± 12.3 s versus 6.3 ± 6.2 s, respectively; Fig. 2e). *S1pr3* [-/-] mice additionally lacked the

robust heat hypersensitivity and cheek wiping in response to injection of 10 µM S1P we

observed in both *S1pr3* [+/+] and *S1pr3* [+/-] littermates (Fig 2d,f). As previously reported, we

observed no defects in baseline thermal sensitivity in *S1pr3* [-/-] mice. The complete loss of S1P-

evoked behaviors in S1PR3-deficient mice suggests that S1PR3 is the sole receptor by which

S1P signals in somatosensory neurons to drive heat hypersensitivity, pain, and itch.

S1PR3 is expressed and functional in subsets of TRPV1+ thermal nociceptors and

TRPA1+ pruriceptors

Our acute behavioral data suggest that S1P-evoked behaviors are mediated by activation of

thermal nociceptors and pruriceptors. We demonstrated by *in situ* hybridization (ISH) that 43%

of dorsal root ganglion cells express *S1pr3.* These neurons comprise two main populations:

medium-to-large diameter mechanonociceptors as well as small-diameter neurons. Here, in

order to address the role of S1PR3 in thermal pain and itch, we characterized the small-

diameter *S1pr3+* neurons by performing co-ISH of *S1pr3* with known markers of C-nociceptor

populations. We found that 51% of *S1pr3+* cells express *Trpv1*, 34.3% express *Trpa1*, and

11.6% express *Mrgpra3* (Fig. 3a). These were indeed small-diameter nociceptors and/or

pruriceptors (mean diameter of *Trpv1+/S1pr3+* cells = 23.4 ± 8.2 μm). In summary, our cellular

localization data reveal two main populations of small-diameter *S1pr3+* neurons: 1) *Trpv1-*

positive thermal nociceptors and, 2) *Trpa1*-positive cells, including *Mrgpra3+* pruriceptors.

These results are consistent with our behavioral studies showing a role for S1PR3 in acute pain

and itch.

We next employed whole cell current-clamp measurements to test whether S1P was sufficient

for action potential firing. Indeed, 1 μM S1P evoked action potential firing in capsaicin-sensitive

neurons (Fig. 3b). As an independent strategy to complement our ISH and electrophysiology

studies, we employed an *S1pr3* mCherry reporter mouse, which produces functional S1PR3-

mCherry fusion protein. Using sensory neurons this mouse in ratiometric calcium imaging, we

found that 56.1 ± 22.4 % of mCherry-expressing neurons were activated by 1 μM S1P (Fig. 3c-d

and these cells had a mean diameter of 22.4 ± 1.0 μm. Furthermore, 51.1 ± 16.8% of capsaicin-

responsive neurons were mCherry+, indicating functional overlap with TRPV1+ thermal

nociceptors. This suggests that only small-diameter thermal nociceptors and/or pruriceptors are

activated by S1P.

Further characterization via calcium imaging in wild-type sensory neurons revealed that 1 μM

S1P activates 34.2% of neurons, which fall into two main populations: 22.6% are

TRPV1+/TRPA1+ neurons, which are capsaicin- and AITC-sensitive, and 11.6% are

TRPV1+/TRPA1- neurons, which are capsaicin-sensitive and AITC-insensitive (Fig. 3e). The

complete overlap of S1P-responsive neurons with TRPV1+ neurons suggests a role for S1P in

heat hypersensitivity. Previous studies have shown that the MrgprA3+ subpopulation of

TRPA1+ neurons are required for many forms of non-histaminergic acute and chronic itch (Han

et al., 2013; Wilson et al., 2011). In keeping with our hypothesized role for S1P in itch, 80% of

chloroquine-responsive MrgprA3+ neurons responded to S1P. Our findings are consistent with

a role for S1P in activation of thermal nociceptors that express TRPV1 and pruriceptors that

express TRPA1.

We thus sought to examine the contribution of S1PR3 to S1P-evoked neuronal activation, for

which the literature is contradictory and suggestive of contributions of both S1PRs 1 and 3.

Incubating cells with the S1PR3-selective antagonist TY 52156 significantly attenuated S1P-

evoked calcium transients (Fig. 3f). In contrast, the S1PR1-selective antagonist W146 had no

effect on S1P responses (Fig. 3f). Thus, we hypothesized that S1PR3 is required for S1P-

evoked calcium signals. In support of our hypothesis, cultured DRG and TG neurons isolated

from *S1pr3*[-/-] mice exhibited no calcium responses to 1 μM S1P, and displayed indistinguishable

AITC- and capsaicin-evoked responses compared to wild-type controls (Fig. 3g). These findings

show that S1PR3, but not S1PR1, is required for neuronal activation by S1P.

S1P activates sensory neurons via distinct TRPV1- and TRPA1-dependent mechanisms.

We set out to identify the molecular pathway underlying S1P-evoked calcium transients in

cultured mouse DRG sensory neurons. S1P responses were mediated solely by calcium influx,

as chelation of extracellular calcium with 10 mM EGTA abolished all responses (Fig. 4a),

suggesting that S1P triggers the opening of plasma membrane cation channels. We also

observed similar results with application of Ruthenium Red (Fig. 4b), a blocker of a number of

calcium-permeable ion channels, including TRP channels.

Since TRPA1 and TRPV1 have been shown to mediate GPCR-dependent calcium signaling

and activation in small-diameter sensory neurons, we asked whether either of these channels

mediated S1P-evoked neuronal activation. We found that the percentage of sensory neurons

exhibiting calcium transients in response to S1P application were reduced in neurons isolated

from both *Trpv1*[-/-] (24.4 ± 5.84% vs. 34.8 ± 6.25%) and *Trpa1*[-/-] mice (8.6 ± 4.3% vs. 34.8 ±

3.3%; Fig. 4c-d).

By ratiometric calcium imaging, we observed a dose-dependent increase in the percentage of

DRG neurons activated by S1P, with an EC_{50} of 104 nM and maximal activation at 1 µM (Fig.

4e). However, neurons which were responsive to both AITC and S1P exhibited a slightly lower

EC_{50} for S1P (102 nM), and were activated by lower S1P concentrations (10 nM) than neurons

which were responsive to capsaicin and S1P (EC_{50} : 155 nM). This supports the notion that

S1P-evoked neuronal activation via TRPV1 and TRPA1 may occur in different populations of

sensory neurons, rather than through A1/V1 crosstalk in the same cells.

We also performed ratiometric calcium imaging on wild-type DRG neurons treated with the

TRPV1 antagonist capsazepine (50 µM) or the TRPA1 antagonist HC-030031 (50 µM). HC-

030031 significantly reduced S1P responses by ~78% (HC: 3.673 ± 1.568% vs. Veh: 17 ±

2.19%) whereas capsazepine significantly attenuated S1P responses by only ~50% (8.69 ±

0.865% vs. 17 ± 2.19%; Fig. 4f). Neurons isolated from mice lacking both TRPA1 and TRPV1

showed S1P responses similar to those evoked by vehicle (Fig. 4g).

We asked whether activation of sensory neurons by S1P/S1PR3 could be due to different G-

protein coupled mechanisms in distinct neuronal subpopulations. We found that inhibition of

PLC signaling using the drug U73122 blocked 42.4% of S1P-evoked calcium responses,

whereas the Gβγ blocker gallein attenuated 73.2% of S1P responses (Fig. 4h). Further

experiments showed that gallein and U73122 attenuate calcium responses to S1P in AITC-

sensitive neurons (S1P + vehicle: 22% S1P+/AITC+; S1P + Gallein: 6.2% S1P+/AITC+; S1P +

U73122: 11.2% S1P+/AITC+) with gallein causing much greater attenuation, whereas U73122

attenuates the majority of responses in capsaicin-sensitive, AITC-insensitive neurons (S1P +

vehicle: 14.5% S1P+/CAP+; S1P + U73122: 1.6% S1P+/CAP+). Our results with the G-protein

inhibitors mirror our observed proportions of TRPA1+ and TRPA1- S1P-responsive neurons,

suggesting that S1PR3 may signal primarily via PLC to activate TRPV1 and via Gβγ and/or PLC

to activate TRPA1, similar to the behavior of other excitatory GPCRs that signal via these

channels in somatosensory neurons (Imamachi et al., 2009; Morita et al., 2015; Wilson et al.,

2011). Overall, our data suggest that S1PR3 signaling opens TRPA1 or TRPV1 channels in two

distinct subpopulations of DRG neurons via canonical G-protein coupled mechanisms.

S1P elicits pain and itch via distinct molecular pathways

We hypothesized that the heat-sensor TRPV1 would be required for S1P-evoked acute pain

and heat hypersensitivity; however, we could not predict whether S1P-evoked itch would be

transduced via the TRPV1-dependent or TRPA1-dependent itch pathways. To this end, we

tested the requirement of TRPA1 and TRPV1 to S1P-evoked itch and pain behaviors. In both

the cheek wiping and radiant heat assays, S1P injection elicited robust acute pain and heat

hypersensitivity behaviors in wild type and *Trpa1*[-/-] mice, but not in *Trpv1*[-/-] mice (Fig. 5a,c).

Wild-type and *Trpv1*[-/-], but not *Trpa1*[-/-] mice, exhibited robust scratching behaviors to S1P that

were significantly greater than vehicle (Fig. 5b). Thus we concluded that S1P-evoked

spontaneous pain behaviors and heat hypersensitivity are mediated by TRPV1 and that S1P-

evoked itch is mediated by TRPA1, a known transducer of acute and chronic itchy stimuli

(Wilson et al., 2011, 2013). Overall, our results highlight a key role for S1PR3 signaling through

distinct molecular pathways in S1P-evoked acute pain, heat hypersensitivity, and itch.

S1P and S1PR3 are essential for chronic thermal hypersensitivity

The Complete Freund's Adjuvant (CFA) model of chronic inflammatory hypersensitivity is widely

used as a model of peripheral inflammation, but the molecular basis of pain hypersensitivity in

this model is incompletely understood. Injection of CFA into the hindpaw causes significant

thermal and mechanical hypersensitivity and swelling in wild type animals. In this model,

thermal hypersensitivity is dependent on TRPV1 ion channels (Caterina et al., 2000), although

the upstream mechanisms of TRPV1 sensitization in CFA are unclear (Boyce et al., 1996).

Additionally, paw edema in this model is due to extensive plasma extravasation, in which

soluble components of blood plasma are exposed to the skin. S1P, which is present at

concentrations 10- to 100-fold higher in blood plasma than in peripheral tissue, presents itself as

an attractive but unstudied component of the CFA inflammatory soup.

We therefore explored the role of S1PR3 in the CFA model of inflammatory pain. A variety of

immune cells infiltrate into the hindpaw following CFA treatment, and a recent study has shown

that myeloid cells are the principal driver of CFA-evoked mechanical but not thermal

hypersensitivity (Ghasemlou et al., 2015). Because S1P plays a key role in immune cell

migration, we asked whether $S1pr3^{-/-}$ displayed normal CFA-evoked immune cell infiltration into

the hindpaw. We observed no difference between $S1pr3^{-/-}$ and $^{+/-}$ animals among all immune cell

subtypes quantified (Fig. 6a-b), in support of previous studies showing that S1PR1 is the

primary mediator of S1P signaling in immune cells.

We next compared CFA-evoked hypersensitivity between $S1pr3^{+/-}$ and $S1pr3^{-/-}$ littermates, as no

significant differences were observed between $S1pr3^{+/+}$ and $S1pr3^{+/-}$ animals (Fig. 6). Strikingly,

$S1pr3^{-/-}$ animals failed to develop heat hypersensitivity (Fig. 6c), and displayed mild attenuation

of hindpaw swelling (Fig. 6d), similar to what has been reported for CFA-treated TRPV1-

deficient animals (Caterina et al., 2000). We therefore hypothesized that S1P levels become

elevated under inflammatory conditions, and that this promotes heat hypersensitivity via S1PR3-

mediated sensitization of TRPV1. In support of our hypothesis, acute blockade of S1P

production with SKI II (Chiba, Takeuchi, Sakai, & Misawa, 2010) rapidly reversed CFA heat

hypersensitivity (Fig. 6e). Furthermore, acute S1PR3 blockade with TY elevated heat-evoked

withdrawal latencies to pre-CFA, baseline levels (Fig. 6e). These results demonstrate that

peripheral S1P promotes CFA-evoked heat hypersensitivity via S1PR3 signaling and is a

component of the inflammatory soup that actively signals to promote pain.

Discussion

S1P is a bioactive signaling lipid best known for its essential roles in immune cell migration,

maturation, and activation via S1PR1 (Matloubian et al., 2004; Roviezzo et al., 2011; Schwab et

al., 2005). Recent studies have linked aberrant S1P signaling to a variety of diseases, including:

asthma, multiple sclerosis, cancer, psoriasis, and pain (Brinkmann et al., 2010; Checa et al.,

2015; Liang et al., 2013; Myśliwiec et al., 2016; Patti et al., 2008; Rivera et al., 2008; F.

Roviezzo et al., 2015). Yet the cellular and molecular targets of S1P in these disorders are

largely unknown. Here, we show that S1PR3 is the sole mediator of S1P-evoked pain and itch

and that S1P signaling is essential to chronic heat hypersensitivity (Fig. 7a), highlighting the

potential for a direct contribution of S1P to sensitization, pain and/or itch in these diseases.

S1P evokes pain and itch via S1PR3

The role of S1P in acute pain and pain hypersensitivity has been controversial. There are

conflicting reports of whether S1P mediates acute flinching and licking pain behaviors, and/or

thermal and mechanical hypersensitivity (Camprubi-Robles et al., 2013; Finley et al., 2013; Mair

et al., 2011; Weth et al., 2015). Likewise, these previous studies employed high-micromolar or

millimolar concentrations of S1P in mouse behavioral experiments, leaving it unclear whether

physiologically relevant S1P levels could contribute to somatosensation. Here we have shown

that physiological levels of S1P (nanomolar to low micromolar) are sufficient to evoke

somatosensory behaviors. We demonstrate that S1P triggers acute pain and heat

hypersensitivity via the noxious heat-sensing ion channel TRPV1. Additionally, S1P promotes

itch-evoked scratching that is dependent on the irritant-sensing ion channel TRPA1. While a

number of previous studies have implicated both S1PR1 and 3 in somatosensory neuron

function (Camprubi-Robles et al., 2013; Mair et al., 2011), several lines of evidence

unequivocally show that S1PR3 is the sole mediator of S1P-evoked pain and itch. First, DRG

neurons isolated from S1PR3-deficient animals do not respond to S1P. Second, application of

an S1PR3-selective antagonist, but not an S1PR1-selective antagonist, attenuates S1P

responses in cultured DRG neurons. Finally, S1PR3-deficient mice display no S1P-evoked

scratching, wiping, or heat hypersensitivity.

We also show for the first time that S1P acts as a pruritogen to evoke acute itch. Recent studies

have shown that circulating S1P levels are increased in chronic itch patients (from ~0.1 to 2 µM

in healthy versus chronic itch patients; Checa et al., 2015; Myśliwiec et al., 2016). S1PR3-

deficient mice displayed a complete loss of S1P-evoked itch behaviors. Future studies will

explore the role of S1P/S1PR3 signaling in chronic itch models to ascertain the pathological

relevance of S1P-evoked itch.

S1PR3 mediates inflammatory thermal pain

In the CFA model of inflammatory pain, both mechanical and thermal hypersensitivity occur with

distinct time courses, suggesting differential modes of regulation (Ghasemlou et al., 2015). It is

well-established that TRPV1 is required for CFA-evoked thermal hypersensitivity (Caterina et

al., 2000). We now show that S1PR3 activation can lead to opening of TRPV1 channels, and

that both S1PR3 and elevated S1P are required for CFA-evoked thermal hypersensitivity. These

data suggest that S1P may be an important contributor to inflammatory pain. Consistent with

this hypothesis, S1PR3-deficient mice display partial attenuation of paw swelling, a hallmark of

neurogenic inflammation, but still develop mechanical hypersensitivity, which has been

previously been shown to depend on infiltration of myeloid cells which display normal infiltration

in the S1PR3 knockout animals. S1PR3 signaling in neurons may generally contribute to pain

under inflammatory conditions. Plasma extravasation and edema, common symptoms of

inflammation, may increase local S1P concentrations and contribute to sensitization. Moreover,

we use pharmacology to show that active S1P signaling is required to maintain chronic heat

hypersensitivity.

Our findings suggest sensory neuronal S1PR3 may play roles in a host of disorders linked to

S1P signaling. Although S1P is produced by nearly every cell type, blood plasma contains the

highest levels of S1P (low micromolar), and S1P is secreted in large amounts by activated mast

cells (Olivera, Allende, & Proia, 2013; F. Roviezzo et al., 2015). Others have shown that

sensory neurons innervating the lung express *S1pr3* and that administration of a non-selective

S1P receptor agonist can enhance asthma-like airway hyperreactivity after acetylcholine

challenge (Trankner, Hahne, Sugino, Hoon, & Zuker, 2014). It is also well known that TRPV1+

sensory afferents are required for airway hypersensitivity (Rogerio, Andrade, & Calixto, 2011).

Furthermore, mast cells, which release S1P when activated (Kalesnikoff & Galli, 2008), play

important roles in allergic asthma as well as itch, and we propose that S1PR3 may play an

important role in mediating itch and/or airway hyperreactivity under these conditions.

Several S1PR modulators are currently in clinical trial for diseases linked to pain and/or itch,

including multiple sclerosis, IBD, and psoriasis (Brinkmann et al., 2010; Degagné & Saba, 2014;

Kunkel, Maceyka, Milstien, & Spiegel, 2013). While these drugs are intended to target the

immune system, we propose neuronal S1P/S1PR3 signaling as a potential target for the

treatment of inflammatory pain and itch.

References

Basbaum, A. I., Bautista, D. M., Scherrer, G., & Julius, D. (2009). Cellular and Molecular

Mechanisms of Pain. *Cell*, *139*(2), 267–284. http://doi.org/10.1016/j.cell.2009.09.028

Bautista, D. M., Jordt, S. E., Nikai, T., Tsuruda, P. R., Read, A. J., Poblete, J., … Julius, D.

(2006). TRPA1 Mediates the Inflammatory Actions of Environmental Irritants and

Proalgesic Agents. *Cell*, *124*(6), 1269–1282. http://doi.org/10.1016/j.cell.2006.02.023

Boyce, S., Rupniak, N. M., Carlson, E. J., Webb, J., Borkowski, J. A., Hess, J., … Hill, R. G.

(1996). Nociception and inflammatory hyperalgesia in B2 bradykinin receptor knockout

mice. *Immunopharmacology*, *33*, 333–335. http://doi.org/10.1007/s00467-008-0901-

3.Dopamine

Brinkmann, V., Billich, A., Baumruker, T., Heining, P., Schmouder, R., Francis, G., … Burtin, P.

(2010). Fingolimod (FTY720): discovery and development of an oral drug to treat multiple

sclerosis. *Nat Rev Drug Discov*, *9*(11), 883–897. http://doi.org/10.1038/nrd3248

Camprubi-Robles, M., Mair, N., Andratsch, M., Benetti, C., Beroukas, D., Rukwied, R., …

Camprubí-Robles, M. (2013). Sphingosine-1-phosphate-induced nociceptor excitation and

ongoing pain behavior in mice and humans is largely mediated by S1P3 receptor. *J*

Neurosci, *33*(6), 2582–2592. http://doi.org/10.1523/jneurosci.4479-12.2013

Canlas, J., Holt, P., Carroll, A., Rix, S., Ryan, P., Davies, L., … Haberberger, R. V. (2015).

Sphingosine kinase 2-deficiency mediated changes in spinal pain processing. *Frontiers in*

Molecular Neuroscience, *8*(August), 1–13. http://doi.org/10.3389/fnmol.2015.00029

Caterina, M. J., Leffler, A., Malmberg, A. B., Martin, W. J., Trafton, J., Petersen-Zeitz, K. R., …

Julius, D. (2000). Impaired Nociception and Pain Sensation in Mice Lacking the Capsaicin

Receptor. *Science*, *288*(5464), 306–313. http://doi.org/10.1126/science.288.5464.306

Chaplan, SR, Bach, FW, Pogrel, JW, Chung, JM, Yaksh, T. (1994). Quantitative assessment of

tactile allodynia evoked by unilateral ligation of the fifth and sixth lumbar nerves in the rat.

Journal of Neuroscience, *53*, 55–63.

Checa, A., Xu, N., Sar, D. G., Haeggström, J. Z., Ståhle, M., & Wheelock, C. E. (2015).

Circulating levels of sphingosine-1-phosphate are elevated in severe, but not mild psoriasis

and are unresponsive to anti-TNF-α treatment. *Scientific Reports*, *5*(July), 12017.

http://doi.org/10.1038/srep12017

Chiba, Y., Takeuchi, H., Sakai, H., & Misawa, M. (2010). SKI-II, an inhibitor of sphingosine

kinase, ameliorates antigen-induced bronchial smooth muscle hyperresponsiveness, but

not airway inflammation, in mice. *Journal of Pharmacological Sciences*, *114*, 304–310.

http://doi.org/10.1254/jphs.10202FP

Chuang, H. H., Prescott, E. D., Kong, H., Shields, S., Jordt, S. E., Basbaum, a I., … Julius, D.

(2001). Bradykinin and nerve growth factor release the capsaicin receptor from

PtdIns(4,5)P2-mediated inhibition. *Nature*, *411*(6840), 957–962.

http://doi.org/10.1038/35082088

Cyster, J. G., & Schwab, S. R. (2012). Sphingosine-1-Phosphate and Lymphocyte Egress from

Lymphoid Organs. *Annual Review of Immunology*, *30*(1), 69–94.

http://doi.org/10.1146/annurev-immunol-020711-075011

Degagné, E., & Saba, J. D. (2014). S1 pping fire: Sphingosine-1-phosphate signaling as an

emerging target in inflammatory bowel disease and colitis-associated cancer. *Clinical and*

Experimental Gastroenterology, *7*(1), 205–214. http://doi.org/10.2147/CEG.S43453

Dixon, W. J. (1980). The Up-and-Down Method for Small Samples. *Journal of the American*

Statistical Association, *60*(312), 967–978.

Donoviel, M. S., Hait, N. C., Ramachandran, S., Maceyka, M., Takabe, K., Milstien, S., …

Spiegel, S. (2015). Spinster 2, a sphingosine 1-phosphate transporter, plays a Critical Role

in inflammatory and Autoimmune Diseases. *The FASEB Journal : Official Publication of the*

Federation of American Societies for Experimental Biology, *29*(12), 5018–5028.

http://doi.org/10.1096/fj.15-274936

Finley, a, Chen, Z., Esposito, E., Cuzzocrea, S., Sabbadini, R., & Salvemini, D. (2013).

Sphingosine 1-phosphate mediates hyperalgesia via a neutrophil-dependent mechanism.

PLoS One, *8*(1), e55255. http://doi.org/10.1371/journal.pone.0055255

Gerhold, K. a., Pellegrino, M., Tsunozaki, M., Morita, T., Leitch, D. B., Tsuruda, P. R., …

Bautista, D. M. (2013). The Star-Nosed Mole Reveals Clues to the Molecular Basis of

Mammalian Touch. *PLoS ONE*, *8*(1). http://doi.org/10.1371/journal.pone.0055001

Ghasemlou, N., Chiu, I. M., Julien, J.-P., & Woolf, C. J. (2015). CD11b [+] Ly6G [−] myeloid cells

mediate mechanical inflammatory pain hypersensitivity. *Proceedings of the National*

Academy of Sciences, 201501372. http://doi.org/10.1073/pnas.1501372112

Han, L., Ma, C., Liu, Q., Weng, H., Cui, Y., Tang, Z., … Dong, X. (2013). A subpopulation of

nociceptors specifically linked to itch. *Nature Neuroscience*, *16*(2), 174–182.

http://doi.org/10.1038/nn.3289.A

Hou, J.-C., & Fang, X.-M. (2015). S1P-S1PRs Alliance in Neuropathic Pain Processing. *Journal*

of Anaesthesia and Perioperative Medicine, *2*, 1–9.

Imamachi, N., Park, G. H., Lee, H., Anderson, D. J., Simon, M. I., Basbaum, A. I., & Han, S.-K.

600 (2009). TRPV1-expressing primary afferents generate behavioral responses to pruritogens

601 via multiple mechanisms. *Proceedings of the National Academy of Sciences of the United*

602 *States of America*, *106*(27), 11330–11335. http://doi.org/10.1073/pnas.0905605106

603 Janes, K., Little, J. W., Li, C., Bryant, L., Chen, C., Chen, Z., … Salvemini, D. (2014). The

604 development and maintenance of paclitaxel-induced neuropathic pain require activation of

605 the sphingosine 1-phosphate receptor subtype 1. *Journal of Biological Chemistry*, *289*(30),

606 21082–21097. http://doi.org/10.1074/jbc.M114.569574

607 Ji, R.-R., Xu, Z.-Z., & Gao, Y.-J. (2014). Emerging targets in neuroinflammation-driven chronic

608 pain. *Nature Reviews. Drug Discovery*, *13*(7), 533–48. http://doi.org/10.1038/nrd4334

609 Kalesnikoff, J., & Galli, S. J. (2008). New developments in mast cell biology. *Nature*

610 *Immunology*, *9*(11), 1215–23. http://doi.org/10.1038/ni.f.216

611 Kono, M., Mi, Y., Liu, Y., Sasaki, T., Allende, M. L., Wu, Y. P., … Proia, R. L. (2004). The

612 sphingosine-1-phosphate receptors S1P1, S1P2, and S1P3 function coordinately during

613 embryonic angiogenesis. *Journal of Biological Chemistry*, *279*(28), 29367–29373.

614 http://doi.org/10.1074/jbc.M403937200

615 Kunkel, G. T., Maceyka, M., Milstien, S., & Spiegel, S. (2013). Targeting the sphingosine-1-

616 phosphate axis in cancer, inflammation and beyond. *Nature Reviews. Drug Discovery*,

617 *12*(9), 688–702. http://doi.org/10.1038/nrd4099

618 Li, C., Li, J., Kays, J., Guerrero, M., & Nicol, G. D. (2015). Sphingosine 1-phosphate enhances

619 the excitability of rat sensory neurons through activation of sphingosine 1-phosphate

620 receptors 1 and/or 3. *Journal of Neuroinflammation*, *12*(1), 1–20.

621 http://doi.org/10.1186/s12974-015-0286-8

Liang, J., Nagahashi, M., Kim, E. Y., Harikumar, K. B., Yamada, A., Huang, W. C., … Spiegel, S. (2013). Sphingosine-1-Phosphate Links Persistent STAT3 Activation, Chronic Intestinal Inflammation, and Development of Colitis-Associated Cancer. *Cancer Cell*, *23*(1), 107–120. http://doi.org/10.1016/j.ccr.2012.11.013

Mair, N., Benetti, C., Andratsch, M., Leitner, M. G., Constantin, C. E., Camprubí-Robles, M., … Haberberger, R. V. (2011). Genetic evidence for involvement of neuronally expressed s1P1 receptor in nociceptor sensitization and inflammatory pain. *PLoS ONE*, *6*(2). http://doi.org/10.1371/journal.pone.0017268

Matloubian, M., Lo, C. G., Cinamon, G., Lesneski, M. J., Xu, Y., Brinkmann, V., … Cyster, J. G. (2004). Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature*, *427*(6972), 355–360. http://doi.org/10.1038/nature02284

Meixiong, J., & Dong, X. (2017). Annual Review of Genetics Mas-Related G Protein – Coupled Receptors and the Biology of Itch Sensation. http://doi.org/10.1146/annurev-genet-120116-024723

Morita, T., McClain, S. P., Batia, L. M., Pellegrino, M., Wilson, S. R., Kienzler, M. A., … Bautista, D. M. (2015). HTR7 Mediates Serotonergic Acute and Chronic Itch. *Neuron*, *87*(1), 124–138. http://doi.org/10.1016/j.neuron.2015.05.044

Myśliwiec, H., Baran, A., Harasim-Symbor, E., Choromańska, B., Myśliwiec, P., Milewska, A. J., … Flisiak, I. (2016). Increase in circulating sphingosine-1-phosphate and decrease in ceramide levels in psoriatic patients. *Archives of Dermatological Research*. http://doi.org/10.1007/s00403-016-1709-9

Olivera, A., Allende, M. L., & Proia, R. L. (2013). Shaping the landscape: Metabolic regulation of S1P gradients. *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids*,

1831(1), 193–202. http://doi.org/10.1016/j.bbalip.2012.06.007

Patti, G., Yanes, O., Shriver, L., Courade, J.-P., Tautenhahn, R., Manchester, M., & Siuzdak, G.

(2008). Metabolomics implicates altered sphingolipids in chronic pain of neuropathic origin.

Nature Chemical Biology, *8*(3), 232–234. http://doi.org/10.1038/nchembio.767

Pinho-Ribeiro, F. A., Verri, W. A., & Chiu, I. M. (2017). Nociceptor Sensory Neuron–Immune

Interactions in Pain and Inflammation. *Trends in Immunology*, *38*(1), 5–19.

http://doi.org/10.1016/j.it.2016.10.001

Rivera, J., Proia, R. L., & Olivera, A. (2008). The alliance of sphingosine-1-phosphate and its

receptors in immunity. *Nature Reviews Immunology*, *8*(10), 753–763.

http://doi.org/10.1038/nri2400

Rogerio, A. P., Andrade, E. L., & Calixto, J. B. (2011). C-fibers, but not the transient potential

receptor vanilloid 1 (TRPV1), play a role in experimental allergic airway inflammation.

European Journal of Pharmacology, *662*(1–3), 55–62.

http://doi.org/10.1016/j.ejphar.2011.04.027

Roviezzo, F., Brancaleone, V., De Gruttola, L., Vellecco, V., Bucci, M., D'Agostino, B., … Cirino,

G. (2011). Sphingosine-1-phosphate modulates vascular permeability and cell recruitment

in acute inflammation in vivo. *The Journal of Pharmacology and Experimental*

Therapeutics, *337*(3), 830–837. http://doi.org/10.1124/jpet.111.179168

Roviezzo, F., Sorrentino, R., Bertolino, A., De Gruttola, L., Terlizzi, M., Pinto, A., … Cirino, G.

(2015). S1P-induced airway smooth muscle hyperresponsiveness and lung inflammation in

vivo: Molecular and cellular mechanisms. *British Journal of Pharmacology*, *172*(7), 1882–

1893. http://doi.org/10.1111/bph.13033

Sanna, M. G., Liao, J., Jo, E., Alfonso, C., Ahn, M. Y., Peterson, M. S., … Rosen, H. (2004).

Sphingosine 1-Phosphate (S1P) Receptor Subtypes S1P1 and S1P3, Respectively,

Regulate Lymphocyte Recirculation and Heart Rate. *Journal of Biological Chemistry*,

279(14), 13839–13848. http://doi.org/10.1074/jbc.M311743200

Sanna, Vincent, K. P., Repetto, E., Nhan, N., Brown, S. J., Abgaryan, L., … Gonzalez-Cabrera,

P. J. (2015). Bitopic S1P3 Antagonist Rescue from Complete Heart Block: Pharmacological

and Genetic Evidence for Direct S1P3 Regulation of Mouse Cardiac Conduction. *Molecular*

Pharmacology, *89*, 176–186. http://doi.org/10.1124/mol.115.100222

Schwab, S. R., Pereira, J. P., Matloubian, M., Xu, Y., Huang, Y., & Cyster, J. G. (2005).

Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients.

Science (New York, N.Y.), *309*(5741), 1735–1739. http://doi.org/10.1126/science.1113640

Shimada, S. G., & LaMotte, R. H. (2008). Behavioral differentiation between itch and pain in

mouse. *Pain*, *139*(3), 681–687. http://doi.org/10.1016/j.pain.2008.08.002

Tominaga, M., Wada, M., & Masu, M. (2001). Potentiation of capsaicin receptor activity by

metabotropic ATP receptors as a possible mechanism for ATP-evoked pain and

hyperalgesia. *Proceedings of the National Academy of Sciences of the United States of*

America, *98*(12), 6951–6956. http://doi.org/10.1073/pnas.111025298

Trankner, D., Hahne, N., Sugino, K., Hoon, M. A., & Zuker, C. (2014). Population of sensory

neurons essential for asthmatic hyperreactivity of inflamed airways. *Proceedings of the*

National Academy of Sciences, *111*(31), 11515–11520.

http://doi.org/10.1073/pnas.1411032111

Tsunozaki, M., Lennertz, R. C., Vilceanu, D., Katta, S., Stucky, C. L., & Bautista, D. M. (2013). A

"toothache tree" alkylamide inhibits Aδ mechanonociceptors to alleviate mechanical pain.

The Journal of Physiology, *591*(Pt 13), 3325–40.

http://doi.org/10.1113/jphysiol.2013.252106

Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., … Ernfors, P. (2015).

Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing

Dmitry. *Nature Neuroscience*, *18*(1), 145–153. http://doi.org/10.1038/nn.3881

Wang, J., Kollarik, M., Ru, F., Sun, H., Mcneil, B., Dong, X., … Undem, B. (2017). Distinct and

common expression of receptors for inflammatory mediators in vagal nodose versus

jugular capsaicin-sensitive / TRPV1- positive neurons detected by low input RNA

sequencing, 1–20.

Weth, D., Benetti, C., Rauch, C., Gstraunthaler, G., Schmidt, H., Geisslinger, G., … Kress, M.

(2015). Activated platelets release sphingosine 1-phosphate and induce hypersensitivity to

noxious heat stimuli in vivo. *Frontiers in Neuroscience*, *9*(April), 1–8.

http://doi.org/10.3389/fnins.2015.00140

Wilson, S. R., Gerhold, K. a, Bifolck-Fisher, A., Liu, Q., Patel, K. N., Dong, X., & Bautista, D. M.

(2011). TRPA1 is required for histamine-independent, Mas-related G protein-coupled

receptor-mediated itch. *Nature Neuroscience*, *14*(5), 595–602.

http://doi.org/10.1038/nn.2789

Wilson, S. R., Nelson, A. M., Batia, L., Morita, T., Estandian, D., Owens, D. M., … Bautista, D.

M. (2013). The Ion Channel TRPA1 Is Required for Chronic Itch. *Journal of Neuroscience*,

33(22), 9283–9294. http://doi.org/10.1523/JNEUROSCI.5318-12.2013

Yamashita-Sugahara, Y., Tokuzawa, Y., Nakachi, Y., Kanesaki-Yatsuka, Y., Matsumoto, M.,

Mizuno, Y., & Okazaki, Y. (2013). Fam57b (Family with sequence similarity 57, member B),

a novel peroxisome proliferator-activated receptor y target gene that regulates

adipogenesis through ceramide synthesis. *Journal of Biological Chemistry*, *288*(7), 4522–

4537. http://doi.org/10.1074/jbc.M112.440792

a

Gene Name
Ppargc1a
Fam57b
Esrrg
Esrrb
Lrp2
Spns1
Spns2
S1pr3
S1pr1

b

Figure 1

Figure 1. S1P pathway genes are positively correlated with chloroquine-evoked itch behaviors. a. Tabular list of the top 10 most highly expressed S1P pathway genes in DRG neurons. b. Histogram showing median absolute r-value for gene expression vs. itch plotted against frequency for 10,000 permutations of random subsets of 10 genes in the BXD DRG transcriptome. Red line indicates median r-value for S1P genes (r_{median} = 0.399, see Methods).

Figure 2. S1PR3 is the sole mediator of S1P-evoked pain and itch. a. Intradermal cheek injection of 10 μM S1P, 2 μM, 0.2 μM, and 20 μL 0.3% methanol PBS (vehicle), with quantification of number of forepaw wipes over the 5 minute post-injection interval; $p < 0.0001$ (one-way ANOVA ($F_{(3,8)} = 30.67$); N = 3 mice per condition). Dunnett's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. b. For same experiment in (a), quantification of time spent scratching over the 30-minute post-injection interval; $p = 0.0027$ (one-way ANOVA ($F_{(3,14)} = 7.798$); N = 3-5 mice per condition). Dunnett's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. c. Radiant heat normalized paw withdrawal latencies 20-30 minutes post injection of 15 μL 10 μM S1P, 0.2 μM S1P. or 0.3% methanol-PBS vehicle i.d. into the hind paw of wild-type mice; $p = 0.0129$ (one-way ANOVA ($F_{(2,6)} = 9.8$); N = 3 mice per condition). Dunnett's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. d. Number of wipes in response to intradermal injection of 20 μL 10 μM S1P or 0.3% methanol-PBS vehicle in cheek of age-matched $S1pr3^{+/-}$ and $S1pr3^{-/-}$ mice; $p < 0.0001$ (one-way ANOVA ($F_{(3,9)} = 40.4$); N = 3-4 mice per condition). Dunnett's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. e. Time spent scratching in response to intradermal injection of 20 μL 0.2 μM S1P or 0.1% methanol-PBS vehicle in rostral back of age-matched $S1pr3^{+/+}$ and $S1pr3^{-/-}$ mice; $p = 0.0089$ (one-way ANOVA ($F_{(3,9)} = 7.274$); N = 3-4 mice per treatment). Dunnett's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. f. Radiant heat normalized paw withdrawal latencies 20-30 minutes post injection of 15 μL 10 μM S1P or 0.3% methanol-PBS vehicle i.d. into the hind paw of age matched $S1pr3^{+/+}$ and $S1pr3^{-/-}$ mice; $p = 0.0004$ (one-way ANOVA ($F_{(3,32)} = 7.964$); N = 8-10 mice per condition). Tukey's multiple comparisons p-values are represented on graph for comparisons made between treated and vehicle groups. Unless otherwise indicated, all error bars represent mean ± SD.

Figure 3

Figure 3. S1PR3 is expressed and functional in subsets of TRPV1+ thermal nociceptors and TRPA1+ pruriceptors. a. Co-ISH of *S1pr3* (green) with *Trpv1*, *Trpa1*, and *Mrgpra3* (magenta) in sectioned whole dorsal root ganglia from adult wild-type mice. Third column: overlay. Scale bar = 100 μm. Images were acquired using a 10x air objective. b. Current-clamp trace of a single neuron firing action potentials in response to bath addition of 1 μM S1P and 1 μM capsaicin. N = 4 of 10 neurons responded to S1P and N = 1 of 1 S1P-responsive also responded to capsaicin. c. (Top) Fura-2 AM calcium imaging before (left) and after (right) addition of 1 μM S1P in S1P3 $^{mCherry/+}$ cultured mouse DRG neurons. Bar indicates fluorescence ratio. Right-hand image indicates mCherry fluorescence. Arrows indicate an mCherry+/S1P- cell (right) and an mCherry+/S1P- cell (left). d. Percentage of mCherry+ DRG neurons which respond to 1 μM S1P (N = 274 neurons). Error bars represent mean response ± SD. e. Venn diagram demonstrating overlap of neuronal subpopulations activated by capsaicin, AITC, chloroquine, and S1P. Relative proportion of overlap represents % overlap as calculated from ratiometric calcium imaging data (N = 4,000 neurons). f. Effects of 100 μM TY 52156 (p = 0.0016 (two-tailed unpaired Student's t-test ($t(6)$ = 5.42); N = 4 wells of 100 cells each)) or 1 μM W146 (p = 0.7921 (two-tailed unpaired Student's t-test ($t(4)$ = 0.2817; N = 3 wells of 100 cells each)) on S1P responses of cultured DRG neurons. Error bars represent mean ± SD. g. Percent responders to S1P in age-matched wild-type and *S1pr3* $^{-/-}$ cultured DRG and TG neurons; p < 0.0001 (one-way ANOVA ($F(7,95)$ = 4.764); N = 2 DRG and 2 TG preparations of 8 wells each). Sidak's multiple comparisons p-values are represented on graph for comparisons made between genotypes. Error bars represent mean ± SEM.

Figure 4. S1P activates sensory neurons via distinct TRPV1- and TRPA1-dependent mechanisms. a. Percent responders for vehicle, 1 μM S1P, AITC, and Cap in Calcium Ringer's- and 10 μM EGTA-Ringer's-treated DRG neurons; $p < 0.0001$ (one-way ANOVA ($F_{(3,11)} = 8.291$); N = 4 wells of ~100 neurons each per treatment). Error bars represent mean ± SD. b. Percent responders for vehicle, 1 μM S1P, AITC, and Cap in vehicle treated and 10 μM Ruthenium Red-treated DRG neurons; $p < 0.0001$ (one-way ANOVA ($F_{(5,17)} = 17.42$); N = 5 wells of ~100 neurons each per treatment). Error bars represent mean ± SD. c. Percent responses for 1 μM S1P in +/+ and *Trpv1* -/- neurons; $p < 0.0001$ (one-way ANOVA ($F_{(7, 179)} = 72.67$); N = 3 age-matched mice per genotype). Error bars represent mean responses ± SEM. d. Percent responses for 1 μM S1P in +/+ and *Trpa1* -/- neurons; $p < 0.0001$ (one-way ANOVA ($F_{(7,210)} = 105$); N = 3 age-matched mice per genotype). Error bars represent mean responses ± SEM. e. Dose-response curve of mean neuronal responses to varying concentrations of S1P. Concentrations used: 1, 10, 50, 100, 200, 1000, and 10,000 nanomolar. Error bars represent mean ± SD (N = 2 animals). Black dotted line indicates sigmoidal fit for all S1P responders from which EC_{50} was derived. Colored traces indicate proportion of neurons responding to S1P and AITC or S1P and capsaicin at indicated concentrations. f. Effect of 50 μM Capsazepine (TRPV1 antagonist) and 50 μM HC-030031 (TRPA1 antagonist) vs. DMSO-Ringer's vehicle on S1P-evoked calcium responses in cultured DRG neurons; $p < 0.0001$ (one-way ANOVA ($F_{(8,30)} = 84.32$); N = 3 wells of ~100 neurons each per treatment) Error bars represent mean responses ± SD. g. Percent responses for vehicle, 1 μM S1P, AITC, and Cap in *Trpa1/Trpv1* [+/+] and [-/-] DRG neurons; $p < 0.0001$ (one-way ANOVA ($F_{(7,40)} = 8.555$); N = 2 age-matched mice per genotype) Error bars represent mean ± SD. h. Percent S1P responders as a fraction of total neurons pre-incubated 15 minutes with either DMSO-Ringer's vehicle, 1 μM U73122 (PLC blocker), or 100 μM Gallein (Gβγ blocker); $p = 0.0030$ (one-way ANOVA ($F_{(3,8)} = 11.32$); N = 3 wells of ~100 neurons each per treatment). Error bars represent mean responses ± SD. Unless otherwise indicated, Sidak's multiple comparisons were made between treated and vehicle for each agonist.

Figure 5. S1P elicits pain and itch via distinct molecular pathways. a. Number of wipes in response to intradermal injection of 20 μL 10 μM S1P or 0.3% methanol-PBS vehicle in cheek of age-matched wild-type, *Trpa1* [-/-] and *Trpv1* [-/-] mice; $p < 0.0001$ (one-way ANOVA ($F_{(5,11)} = 33.98$); N = 3 age-matched mice per condition). Error bars represent mean ± SD. b. Time spent scratching in response to intradermal injection of 20 μL 0.2 μM S1P or 0.1% methanol-PBS vehicle in rostral back of age-matched wild-type, *Trpa1* [-/-] and *Trpv1* [-/-] mice; $p < 0.0001$ (one-way ANOVA ($F_{(5,91)} = 14.13$); N = 8 or more age-matched mice per condition). Error bars represent mean ± SD. c. Radiant heat normalized paw withdrawal latencies 20-30 minutes post injection of 15 μL 10 μM S1P or 0.3% methanol-PBS vehicle i.d. into the hind paw of age matched wild-type, *Trpa1* [-/-] and *Trpv1* [-/-] mice; $p < 0.0001$ (one-way ANOVA ($F_{(5,66)} = 31.23$); N = 7 or more age-matched mice per condition). Error bars represent mean ± SD. For all graphs, Sidak's multiple comparisons were made between treated and vehicle for each genotype.

Figure 6. S1P and S1PR3 are essential for chronic thermal hypersensitivity. a. Recruitment of neutrophils (Neut.) or inflammatory monocytes (IMs) to hindpaw skin 24 hours post-CFA administration in HET and KO mice, as a dot plot of CD11b+ cells plotting Ly6G fluorescence intensity vs. Ly6C intensity (AFU). Boxes are for illustrative purposes. b. Recruitment of immune cells, including neutrophils (Neut.) and inflammatory monocytes (IMs) to hindpaw skin 24 hours post-CFA administration in HET (grey) and KO (green) mice, displayed as total number of cells; N = 8 mice per genotype. Sidak's multiple comparisons were made between HET and KO for each cell type. c. Thermal latency before and after CFA treatment (indicated by dotted line); $p_{genotype}$ = 0.0053 (two-way ANOVA ($F_{(1,8)}$ = 14.36); N = 5 mice per genotype). Sidak's multiple comparison between genotypes for specific time points indicated on graph. Error bars represent mean ± SD. d. (Top) Images of HET and KO littermate mice 48 hours after CFA injection into the left paw (arrows). (Bottom) Hindpaw swelling of CFA-treated paw compared to PBS-treated paw as measured to the nearest 1/10th mm 24- and 48-hours post CFA injection; ; $p_{genotype}$ = 0.0003 (two-way ANOVA ($F_{(1,8)}$ = 9.64); N = 5 mice per genotype). Error bars represent mean ± SD. e. (Left) Thermal latency assessed before and after CFA injection (arrow) with either vehicle (VEH) or SKI II acutely administered on Day 1; $p < 0.0001$ (one-way ANOVA ($F_{(2,21)}$= 19.16); N = 5-7 mice per treatment). Dunnett's test comparisons to baseline are indicated on graph. (Right) Thermal latency assessed before ("Baseline") and 24 hours post CFA injection with either vehicle (CFA + VEH) or TY 52156 (CFA + TY) acutely administered; $p < 0.0001$ (one-way ANOVA ($F_{(2,17)}$= 50.44); N = 5 mice per treatment). Dunnett's test comparisons to baseline are indicated on graph. Error bars represent mean ± SD.

a

Figure 7

Figure 7. Model of the effects of elevated S1P on S1PR3-dependent thermal hypersensitivity and acute itch in sensory neurons. a. Elevated S1P can 1) elicit thermal hypersensitivity or acute pain via S1PR3-dependent activation of TRPV1 (top) and 2) elicit itch-evoked scratching via S1PR3-dependent activation of TRPA1.