Human Herpes Virus 6 (HHV-6) - Pathogen or Passenger? A pilot study of clinical laboratory data and next generation sequencing

Colin Sharp[a,b], Tanya Golubchik[c,d], William F. Gregory[a],

Anna L. McNaughton[e], Nicholas Gow[f], Mathyruban Selvaratnam[f],

Alina Mirea[f], Dona Foster[g], Monique Andersson[f],

Paul Klenerman[e,f], Katie Jeffery[f], Philippa C. Matthews[e,f#]

[a] The Roslin Institute, University of Edinburgh, Easter Bush, Midlothian, Edinburgh, Scotland, UK

[b] Edinburgh Genomics, Ashworth Laboratories, University of Edinburgh, Edinburgh, EH9 3FL, Scotland, UK

[c] The Wellcome Trust Centre for Human Genetics, Roosevelt Drive, Oxford, OX3 7BN, UK

[d] Big Data Institute, University of Oxford, Old Road, Oxford, OX3 7FZ, UK

[e] Nuffield Department of Medicine, Peter Medawar Building for Pathogen Research, University of Oxford, South Parks Road, Oxford, OX1 3SY, UK

[f] Department of Infectious Diseases and Microbiology, Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Headley Way, Headington, Oxford, OX3 9DU, UK

[g] Modernising Medical Microbiology Research Group, Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Headley Way, Headington, Oxford, OX3 9DU, UK

 # Corresponding author: Philippa Matthews; Telephone: 0044 1865 271973;

Email: philippa.matthews@ndm.ox.ac.uk

KEY WORDS: Human herpesvirus 6, HHV-6, screening, diagnosis,

encephalitis, haematology, epidemiology, metagenomics, next generation

sequencing, Illumina, haematology

RUNNING TITLE: HHV-6 epidemiology and diagnostics in a UK cohort

ABSTRACT

Human herpes virus 6 (HHV-6) is a ubiquitous organism that can cause a variety of clinical syndromes ranging from short-lived rash and fever through to life-threatening encephalitis. In order to determine patterns of HHV-6 infection and detection, we took two approaches. First, we scrutinized HHV-6 requests in samples submitted to our UK hospital laboratory through routine diagnostic pathways. Of 72 samples tested for HHV-6 by PCR at the request of a clinician, 24 (33%) were positive for HHV-6. The majority of these patients were under the care of the haematology team (73%), and there were borderline associations between testing positive for HHV-6 DNA and both Graft versus Host Disease (GvHD) and central nervous system (CNS) disease (p=0.05 in each case). Second, we undertook a pilot study using Illumina next generation sequencing to investigate the application of this tool to detection of HHV-6 in CSF and respiratory samples that were initially submitted to the laboratory for other diagnostic tests. We confirmed detection of HHV-6, but highlight challenges in interpretation of metagenomic data and the need for further controls to define sensitivity and specificity. Our results lead to recommendations for the improvement of existing laboratory protocols, and underpin ongoing efforts to explore next generation sequencing technology for viral diagnostics.

ABBREVIATIONS

- cDNA – complementary DNA

- CiHHV-6 – Chromosomally integrated HHV-6

- CMV – Cytomegalovirus

- CNS – central nervous system

- CSF – cerebrospinal fluid

- EPR – Electronic patient record

- GvHD – graft versus host disease

- HHV-6 - Human herpes-virus-6

- HLA – human leucocyte antigen

- HSCT - Haematopoietic stem cell transplant

- HSV – Herpes Simplex Virus

- HTLV – Human T cell lymphotrophic virus

- NGS – Next Generation Sequencing

- OUH – Oxford University Hospitals NHS Foundation Trust

- PCR – Polymerase Chain Reaction

- RSV – Respiratory Syncytial Virus

- VZV – Varicella Zoster Virus

INTRODUCTION

Human herpes-virus-6 (HHV-6) is a human beta-herpesvirus, first described in 1986 in patients with Human T cell lymphotrophic virus (HTLV) infection and/or haematological malignancy (1). Like its close relative human cytomegalovirus (CMV), it is ubiquitous (frequently infecting >95% of the population by the age of two years), has the potential for latency followed by chronic low level replication or reactivation, and may modulate immune responses to other pathogens (2-5).

In children, clinical consequences of HHV-6 range from a trivial (or asymptomatic) infection, to a more significant febrile illness with rash ('roseola'), or occasionally encephalitis (6). HHV-6 reactivation has also been described in the setting of encephalitis or meningitis caused by other pathogens, such as enteroviruses, in which context it appears to follow a generally benign course (7). At the other end of the spectrum, HHV-6 can reactivate in the context of severe sepsis (8) and is a cause of potentially life-threatening pathology in patients with haematological malignancy, usually following bone marrow transplantation (9-15).

HHV-6 transmission is predominantly via respiratory secretions or saliva, but can also be vertical. The majority of mother-to-child transmission occurs as a result of high-level germline integration, referred to as chromosomally-integrated HHV-6, 'CiHHV-6' (16, 17). This chromosomal integration, present in an estimated 1% of the population, can lead either to reactivation (17), or to persistent HHV-6 viraemia, with a suggested threshold of ≥5.5 log10

copies/ml (approx. 300,000 copies/ml) being indicative of CiHHV-6 (18). HHV-

6 has two variants, A and B, that share approximately 90% sequence

homology (19). HHV-6A accounts for more CiHHV-6 (17), while HHV-6B is

more likely to be acquired as an acute infection, is associated with childhood

rash/fever (4) and with reactivation following bone marrow transplantation

(19). Although HHV-6A is less common as a cause of CNS disease (20), it

may be more aggressive when present (19).

Given that exposure to the virus is common, HHV-6 laboratory diagnostics

raises a number of challenges: what sample type to test, in which patient

groups to focus, and how to should interpret a positive test result. There is

increasing interest in metagenomic ('next generation sequencing', NGS)

approaches to clinical microbiology diagnosis of many pathogens (21-25) but

ongoing development is required to optimize these pipelines, including

tackling the challenge of high proportion of human DNA reads, differentiating

between pathogenic organisms and commensal or environmental flora, and

determining thresholds at which the identification of an organism is likely to be

clinically significant (26) (Suppl data set 1). The latter challenge is particularly

pertinent for herpes viruses, which are ubiquitous and potentially integrated

into the host.

We here present data first from the local diagnostic microbiology laboratory, to

review the patterns of clinical testing for HHV-6, and to describe the patients

who test HHV-6 DNA positive. Second, we report the results of screening

CSF and respiratory samples, in cases where HHV-6 testing had not been

specifically requested, using PCR and NGS in a small pilot study to ascertain

the extent to which this virus can be detected in routine diagnostic samples.

Together, these aim to describe the distribution of HHV-6 in local clinical

samples, and to evaluate the contribution made by different diagnostic tools,

thereby informing ongoing development of laboratory protocols for detection

of this organism.

MATERIALS AND METHODS

Study site, cohorts and ethics

Clinical data and samples from between 2013-2016 were collected from the

microbiology department at Oxford University Hospitals (OUH) NHS

Foundation Trust, a large tertiary referral teaching centre in the UK

(http://www.ouh.nhs.uk/). This study pertains to the analysis of two separate

sample cohorts:

i.　Samples submitted to the laboratory by clinicians with a request for

HHV-6 screening (study ID numbers prefixed by **HHV**); the metadata

file for this cohort is provided on-line (Suppl data set 1).

ii.　Samples which had no request for HHV-6 testing, but had completed

routine diagnostic laboratory testing for other indications and were

used for **v**iral **s**equencing studies (study ID numbers prefixed by **VS**);

supporting datafiles for metagenomic work are provided online (Suppl

data set 1).

Approval for retrospective collection of clinical and laboratory data was

granted by the Oxford University Hospitals NHS Foundation Trust Clinical

Audit Committee (HHV cohort samples). Testing of consecutive anonymised

laboratory samples (VS cohort samples) was approved by local Research

Services and through review via the UK Integrated Research Application

System (REC Reference 14/LO/1077).

Collection of local laboratory data (HHV cohort)

We undertook an electronic search of the OUH Microbiology laboratory

system to identify all instances of an HHV-6 test (antibody or viral load) being

requested over a three year period commencing 1-Jan-2013, and recorded

age, sex, sample type and patient location. We interrogated the hospital

Electronic Patient Record (EPR) to determine the patient's underlying

diagnosis. Follow-up data for survivors were available for a median period of

25 months (range 316 – 1374 days). During the period of the study, diagnostic

tests were sent to two different National Reference Laboratories (Colindale

from 1-Jan-2013 to 1-Oct-2014, and Bristol for the remainder of the study

period). Testing in these centres was undertaken by in-house real time PCR

assays (27).

Data collection and statistical analysis

We used GraphPad Prism v.6.0f for statistical analysis with Fisher's Exact

Test applied to identify differences between binary groups, and the Mann

Whitney U test for continuous variables. Multivariate regression analysis was

undertaken   using   open   access   on-line   software

(https://docs.google.com/spreadsheets/u/0/).

Testing CSF and respiratory samples (VS cohort)

We identified 100 CSF samples and 100 respiratory samples (comprising

throat swabs (n=22), nasopharyngeal aspirates (n=42), endotracheal

aspirates (n=4), and bronchoalveolar lavage samples (n=32)) that had been

processed by the diagnostic microbiology laboratory in 2013/2014. These

represented a subgroup of patients deemed to be at 'high risk' of viral

infection based on the following criteria:

i. For CSF, the clinical request for testing included viral causes of

meningitis and encephalitis;

ii. For respiratory samples, the patient had a clinical history of

immunocompromise and/or was on the intensive care unit.

Samples had all undergone clinical laboratory testing as described in our

supplementary methods file (Suppl data set 2) and were stored at -80°C prior

to further processing. We selected consecutive samples; the only exclusion

criterion was inadequate sample volume (<200 ul). For each sample, we

documented patient age group and clinical location, and recorded the clinical

information that had been supplied with the sample and microbiology

laboratory data generated as part of the primary clinical testing pipeline.

PCR for HHV6 (VS cohort)

Additional experimental testing was undertaken at the Roslin Institute,

University of Edinburgh. Nucleic acids were extracted from 200µl of each

sample using the AllPrep DNA/RNA Mini Kit (Qiagen) and recovered in 30µl

of nuclease-free water. To allow for broad detection of HHV-6 and other

known herpesviruses, 4ul of DNA was used as template for a consensus PCR

primer as previously published (28). The results were visualised on 2%

agarose gels and amplicons from positive reactions were cut directly from the

gels for sequencing. Direct amplicon sequencing was performed using BigDye

Terminator v3.1 (Applied Biosystems) according to the manufacturer's

instructions with both second round primers. Sequencing reactions were read

by Edinburgh Genomics and assembled using SSE v1.2 (29).

Next generation sequencing using Illumina HiSeq and Bioinformatic

analysis of NGS dataset (VS cohort)

Our methods are described in supplementary data files, available on-line

(Suppl data set 2).

RESULTS

Routine clinical samples received for HHV-6 testing (HHV cohort)

During a three-year study period, our clinical laboratory received 85 samples

with a request for HHV-6 testing (Fig 1A). In total, 41 different patients were

tested; 22 M:19 F; median age 52 years (range 2-71; IQR 39-61). For full

clinical dataset see Suppl data set 1. Most patients were under the care of the

haematology team (n=30; 73%), among whom five had Graft versus Host

Disease (GvHD). Nine patients had a report of central nervous system (CNS)

disease, including encephalitis, encephalopathy, meningism and seizures.

Results of HHV-6 testing (HHV cohort)

Of 85 samples received, 72 were tested by PCR at the regional reference

laboratory, 24 were positive for HHV-6 DNA (33%). Among 41 patients tested,

15 (37%) were PCR positive at some point during the three-year time period

under review (Fig 1A). Although subtyping is not now undertaken routinely, of

seven samples that were subtyped, all were reported as HHV-6B. Among

patients who tested PCR positive, this was more frequently derived from

blood (9/15 patients) than from CSF (3/15 patients) (Table 1). Positive HHV-6

status was not statistically associated with patient age, sex, or haematological

malignancy, but there was a borderline association with both GvHD and CNS

disease (p=0.05 in both cases; Table 1; Fig. 2A and B). On multiple logistic

regression analysis, there was no relationship between HHV-6-positivity and

any patient or laboratory characteristic (Table 1).

In this cohort, 23/41 (56%) of patients died, at a median age of 56 years. Of

those who died, 11/23 (48%) had tested HHV-6-PCR positive at some point

during our study period, compared to surviving patients of whom 4/18 (22%)

had a positive HHV-6 PCR, although this difference did not reach statistical

significance (p=0.1). All those who died did so within 19 months of the HHV-6

test (range 1-552 days, median 163 days). We did not have sufficient clinical

data to determine cause of death.

Quantification of HHV-6 viral load in blood and CSF (HHV cohort)

HHV-6 DNA quantification was undertaken in 23 samples from ten different

patients (Fig 2C). The levels varied from below the threshold for accurate

quantification in serum and bone marrow, to an individual with $>1.0 \times 10^{6}$

copies/ml in blood and $2.0x10^8$ DNA copies/ml in CSF (patient ID HHV-012),

suggesting CiHHV-6. Another patient had HHV-6 DNA detected in both a

blood and CSF sample (patient ID HHV-007, $9x10^4$ DNA copies/ml in blood

vs. $3x10^5$ copies/ml in CSF). This patient, an adult with haematological

malignancy, was reported to have limbic encephalitis, and the raised HHV-6

titre in CSF compared to blood is in keeping with a localized CNS pathology

caused by the virus.

Longitudinal testing (HHV cohort)

Among a total of 72 tests, 46 were longitudinal samples from thirteen

individual patients. Among these thirteen, the median number of serial tests

was three, but one individual (ID HHV-045) had seven EDTA blood samples

and a CSF sample tested within an 8 week time period (of these, only one

blood sample was positive for HHV-6 DNA). There was no statistical

association between multiple HHV-6 tests and other factors that might predict

disease severity (age, sex, mortality, haematological malignancy, transplant,

intensive care location, documented co-infection; $p>0.1$ for all parameters in

multiple regression analysis).

Screening laboratory samples for HHV-6 by PCR (VS cohort)

We screened consecutive laboratory CSF (n=100) and respiratory (n=100)

samples for herpesvirus DNA by PCR, and subsequently confirmed HHV-6 by

sequencing (workflow shown in Fig 1B, and Suppl data set 1). We identified

HHV-6 DNA in eight (4%) of these, comprising three CSF samples and five

respiratory samples (the samples testing PCR positive are shown in Table 2).

Four of these PCR-positive cases were children age <5 years, representing

the age group in whom primary HHV-6 infection is most likely to occur.

However, in four samples (two from adults and two from children) an

alternative potential pathogen was identified by the clinical diagnostic lab

(VZV, HSV, Rhinovirus, Metapneumovirus). This illustrates the difficulty in

distinguishing between HHV-6 as a primary pathogen, as a co-infecting agent

contributing to pathology, or as an innocent bystander (potentially as a result

of reactivation).

The overall HHV-6 prevalence of 4% in this group of unselected samples is

significantly lower than the 33% rate obtained from the samples in which the

clinician had specifically requested HHV-6 testing ($p<0.0001$; Fisher's Exact

Test). This suggests that the targeting of the highest risk groups (primarily

patients with haematological malignancy) for HHV-6 testing is appropriate,

and that the high prevalence of detectable HHV-6 DNA in this group is not

merely reflective of universal reactivation of herpesviruses in an unwell

hospitalized population.

Screening laboratory samples for HHV-6 DNA by NGS (VS cohort)

We screened a subset of 20 samples (10 CSF and 10 respiratory; Fig 1B)

using a metagenomic (NGS) approach. In these 20, we identified HHV-6 in

four samples that we deemed 'positive'; one CSF and three respiratory

samples (Table 3). The representation of HHV-6 reads as a proportion of the

virome is shown for one representative respiratory sample in Fig 3, and the

number and distribution of reads in Fig 4. Sample VS183 was designated

HHV-6-B by Kraken, while the other three positive samples were identified as

HHV-6-A (Fig 4B). The coverage of the HHV-6 genome was only 1% in each

case, but multiple reads distributed across the genome, including in

conserved regions (Fig 4B), add confidence to the conclusion that HHV-6

DNA is genuinely present in these samples. The equivocal outcomes,

comprising low numbers of HHV-6 reads, were all respiratory samples, and

the sequences clustered in the repeat region of the genome, suggesting lower

specificity for HHV-6.

In sample VS067 the clinical syndrome was not explained by other diagnostic

results, and HHV-6 infection was a plausible agent of the clinical syndrome (in

keeping with meningoencephalitis). In the three other cases, other primary

pathogens had been identified (Table 3), although it is plausible that HHV-6

could have been a contributory agent.

The metagenomic sequence data have been uploaded to European

Nucleotide Archive (ENA) https://www.ebi.ac.uk/ena; HHV6 sequence

accession numbers ERS1980462 (sample ID VS067), ERS1980463 (sample

ID VS183), ERS1980464 (sample ID VS200) and ERS1980465 (sample ID

VS207).

Concordance between PCR and NGS (VS cohort)

Among the total of eight samples that were HHV-6 positive by conventional

PCR, three were tested on the NGS pipeline (sample IDs VS114, VS121 and

VS183). HHV-6 reads were detected by NGS in all three, although two had

low numbers of reads leading us to classify them as equivocal (Fig 4A,B).

Conversely, in four samples where HHV-6 was detected by NGS, only one of

these was positive by PCR.

DISCUSSION

Summary of findings

This preliminary study provides insight into the patterns of HHV-6 testing in a

large UK teaching hospital, with the aim of updating recommendations for

consistent laboratory practice and translating this into improvements in clinical

practice. We also describe the application of two different approaches

(targeted PCR, vs. a sequence-independent NGS pipeline) to report the

'background' presence of HHV-6 DNA in CSF and respiratory samples at the

point of discard from a routine clinical microbiology laboratory. Although our

NGS data only represent a small pilot study, to date, we have identified only a

single other reference that describes a metagenomic approach to the

diagnosis of HHV-6 infection (6).

Although it is recognized that HHV-6 can reactivate in the context of critical

illness (e.g. septic shock (8)), we here found a higher prevalence in clinical

samples taken from a high-risk group (HHV cohort) that in laboratory samples

representing an unwell and immunocompromised hospital population (VS

cohort), suggesting that HHV-6 is not simply reactivating across the board in

unwell hospitalized patients. Although we cannot attribute mortality to HHV-6

in the context of this study, a recent study documents poor outcomes of HHV-

6 disease in the bone marrow transplant population (15).

Implications for HHV-6 screening and treatment

The combination of a high rate of HHV-6 DNA detection and the high mortality

(HHV cohort), suggest that we should consider lowering our threshold for

testing in this context, in order to identify patients who may benefit from

treatment earlier in the course of disease. In future, enhanced prospective

surveillance may help shed more light on the clinical significance of HHV-6 in

patients with haematological malignancy. This study was not designed to

evaluate the use or outcome of antiviral therapy as this information could not

robustly be captured retrospectively for the time period in question. There are

some data to suggest good outcomes from treatment of symptomatic viraemia

(30), most commonly with ganciclovir and/or foscarnet. However, there is no

universally agreed definition of clinical disease or threshold for therapy and

this area is not well informed by clinical trials (31); the gains made by testing

early have to be carefully balanced against the possibility of identifying

patients with bystander viral reactivation in whom the toxicity and side-effects

of treatment would not be justified.

Suggested applications for current clinical diagnostic pipeline

Our recommendations following this study include:

i.   Diagnostic laboratories should only offer testing for HHV-6 PCR, not for

IgG testing. The rationale for this is that the majority of individuals are

likely to have been exposed, and therefore will be HHV-6 IgG positive.

In the setting of profound immunocompromise, where there may be a

wish to identify patients at risk of reactivation, IgG may be falsely

negative due to the reduced pool of productive B cells.

ii.   Longitudinal HHV-6 testing should generally be reserved for monitoring

response to therapy, but if a high index of suspicion exists (e.g. a

profoundly immunosuppressed patient who becomes encephalopathic

in the absence of an alternative explanation), then serial testing for

diagnostic purposes may be helpful.

Significance of HHV-6 subtype

Currently, HHV-6 subtype has no consistent bearing on the approach to

clinical management and is therefore not part of routine testing. However,

further studies, including NGS, may determine whether different clinical

syndromes are consistently associated with the two subtypes. In this study, all

viruses subtyped by the clinical laboratory were HHV-6B, which reflects the

anticipated pattern. In contrast, the NGS primarily detected HHV-6A,

potentially reflecting reactivation of integrated virus.

Sensitivity and specificity of HHV-6 detection by different methods

Assigning a sensitivity or specificity to different methods is difficult in the

absence of a single gold standard test that includes application to

metagenomic data. For samples with low HHV-6 copy numbers, PCR is

anticipated to be more sensitive than the current metagenomic approach; this

is illustrated by two samples that tested positive by PCR but were equivocal

by NGS. However, the overall proportion of samples testing positive for HHV-

6 DNA was higher by NGS than by PCR. This may at least in part be

accounted for by use of highly degenerate herpesvirus primers that were used

to amplify DNA, followed by sequencing primers that have a higher degree of

sequence homology for HHV-6B than HHV-6A (28). Although we cannot

exclude the possibility of cross-contamination by samples on the HiSeq

platform, this is more frequently encountered in the context of samples with

high viral loads, and the low copy numbers in our samples generally make this

unlikely.

Clinical significance of NGS data

In screening unselected samples by NGS, we identified HHV-6 at levels

deemed to be potentially clinically significant in 4/20 (20%) samples, although

further studies are needed, and these thresholds remain to be determined

with confidence. Importantly, positivity cannot be inferred by read numbers

alone (32), and we have also used genome coverage to help infer which

samples were truly positive for HHV-6. The four patients positive for HHV-6

by NGS all had clinical syndromes that could be compatible with HHV-6

infection, but due to a limited dataset we cannot attribute causality.

Interestingly, positive HHV-6 PCR from young children with severe respiratory

tract infections suggest a potential pathological role of the virus that is not well

described in this population to date. When HHV-6 detection is deemed

significant, this could support the introduction of antiviral therapy, and/or

reduce exposure to broad-spectrum antibiotics.

In parallel with wet lab methods, bioinformatic approaches for interpreting

metagenomic data derived from clinical samples are still evolving. Multiple

considerations feed into this interpretation, some of which we have

summarized as an algorithm in Fig 5. More robust approaches will evolve as

the volume and quality of data from clinical metagenomic testing improves.

Caveats and limitations of this study

The number of clinical requests for HHV-6 testing each year is small – even in

a tertiary referral teaching hospital. By nature, our clinical sample set (HHV

cohort) was strongly skewed towards sampling a high-risk population.

Together, these factors reduce statistical power in the identification of

significant risk factors or clinical associations of HHV-6. Our data quality is

dependent upon the completeness of electronic data, and we may have

under-represented certain clinical groups due to lack of complete information.

For our VS cohort, we screened only CSF and respiratory samples; inclusion

of blood samples remains an important avenue for future enquiry.

Further evaluation of optimum methods for NGS is required; some of the

potential caveats are presented in more detail in our supplementary methods

file (Suppl data set 2). For our own future metagenomic work, we are currently

developing positive internal controls, and for HHV-6 specifically, an

international control standard is currently being prepared by the National

Institute for Biological Standards and Control (NIBSC).

Summary

This study improves our understanding of HHV-6 detection both in a clinically

selected population and in a cross-section of laboratory samples. On the

basis of the clinical component of the study, we have improved the

consistency of our laboratory approach to these requests. Although it is

doubtless at times a benign passenger, HHV-6 is indeed significantly

associated with the clinical syndromes that arise in patients with profound

immunocompromise in the haematology setting, and may have a role in other

syndromes, including respiratory infections in children aged <5 years.

Metagenomic approaches to clinical diagnostics are accelerating, and as

additional data become available, increasing insights will be gained into the

interpretation of these results.

TABLES

Table 1: Relationship between HHV-6 status and clinical/demographic
features in 41 patients with samples submitted for HHV-6 testing to a
clinical UK microbiology lab.
465

	HHV-6 PCR positive[a] (n=15 patients)	HHV-6 PCR negative (n=26 patients)	p-value[b] (univariate analysis by Fisher's Exact test)	p-value[b] (multivariate analysis by logistic regression)
Sample type[c]: blood	9/15 (60%)	15/26 (58%)	1.0	0.7
Sample type[c]: CSF	3/15 (20%)	4/26 (15%)	0.7	0.8
Age ≥50 years	10/15 (67%)	13/26 (50%)	0.3	0.3
Male sex	8/15 (53%)	14/26 (54%)	1.0	0.2
Haematological malignancy present	13/15 (87%)	17/26 (65%)	0.2	0.9
Graft versus host disease present	4/15 (27%)	1/26 (4%)	**0.05**	0.1
CNS[d] disease present	6/15 (40%)	3/26 (12%)	**0.05**	0.1
Solid organ transplant present	0/15 (0%)	2/26 (8%)	0.5	1.0
Death[e]	11/15 (73%)	11/25 (44%)	0.1	0.2

466
467 [a] In the case of patients tested more than once, any instance of PCR positive
468 is recorded as positive; [b] Sample types were not documented for 3 patients
469 testing PCR-positive, and for 7 patients testing PCR-negative; [c] CNS=central
470 nervous system; [d] Prospective survival data were available for 40/41 patients.
471

Table 2: Summary of eight clinical samples (from a total of 200 consecutive samples selected for screening) that tested positive for HHV-6 DNA by PCR

Study ID number	Sample type	Age group (years)	Location	Clinical details	CSF cell count: total WCC (cells/µl)	CSF cell count: lymphs (cells/µl)	CSF cell count: RBC (cells/µl)	Bacterial culture results from routine clinical lab	Molecular diagnostics (PCR) from routine clinical lab
VS013	CSF	20-24	Emergency Department	Headache, ? SAH	346	342	70	No growth	VZV detected
VS014	CSF	<5	Paediatric ICU	? Meningitis	8	6	8	No growth	No virus detected
VS022	CSF	60-64	Medicine	Confusion, temporal attenuation on CT brain	28	26	28	No growth	HSV type 1 detected
VS114	NPA	<5	Paediatric ICU	? Brochiolitis	n/a	n/a	n/a	Not requested	No organism detected
VS121	NPA	<5	Paediatric ICU	LRTI	n/a	n/a	n/a	Not requested	Rhinovirus detected
VS127	BAL	65-69	Chest Unit	Pneumonia	n/a	n/a	n/a	No growth	No organism detected
VS152	BAL	55-59	Chest Unit	? TB	n/a	n/a	n/a	*Streptococcus viridans*	No organism detected
VS183	NPA	<5	Paediatrics	Viral infection	n/a	n/a	n/a	Not requested	Meta-pneumovirus detected

CSF=cerebrospinal fluid, NPA=nasopharyngeal aspirate, BAL=broncho-alveolar lavage, HSV = Herpes Simplex Virus, ICU=intensive care unit, SAH=subarachnoid haemorrhage, CT=computer tomography, LRTI=lower respiratory tract infection, TB=tuberculosis, PCR=polymerase chain reaction, VZV=varicella zoster virus. '?' in clinical details column reflects potential but unconfirmed diagnosis being raised as a query by the requesting clinician.

Table 3: Summary of HHV-6 results from twenty clinical samples screened for pathogens by a metagenomic (NGS) approach.

Study ID number	Sample type	Age (years)	Clinical details on request form	CSF Lympho-cyte count (cells/µl)	Clinical laboratory virology result	HHV-6 PCR result	HHV-6 detection by NGS approach
VS011	CSF	<5	Fever	2	No virus detected	Negative	Negative
VS012	CSF	<5	VP shunt	4	No virus detected	Negative	Negative
VS033	CSF	60-64	No data	0	No virus detected	Negative	Negative
VS036	CSF	<5	No data	0	No virus detected	Negative	Negative
VS047	CSF	5-9	No data	0	No virus detected	Negative	Negative
VS060	CSF	40-44	Previous high CSF white cell count (50).	32	No virus detected	Negative	Negative
VS067	CSF	50-54	Confusion, headache, malaise, fever	438	No virus detected	Negative	Positive (BWA HHV-6A, 3653 reads, 1% coverage)
VS071	CSF	70-74	Headache, fever, cognitive impairment	0	No virus detected	Negative	Negative
VS079	CSF	40-44	Sepsis, Confusion, Encephalopathy	2	No virus detected	Negative	Negative
VS088	CSF	20-24	Mixed sensory and motor symptoms	0	No virus detected	Negative	Negative
VS114	Resp (NPA)	75-79	Possible bronchiolitis	N/A	No organism detected	Positive	Equivocal (HHV-6A; BWA mapping within repeat region only)
VS121	Resp (NPA)	<5	Lower respiratory tract infection	N/A	Rhinovirus detected	Positive	Equivocal (HHV-6A; BWA mapping within repeat region only)
VS134	Resp (BAL)	55-59	Neutropenic; sudden respiratory deterioration	N/A	No organism detected	Negative	Equivocal (HHV-6A; BWA mapping within repeat region only)
VS147	Resp (BAL)	50-54	?TB	N/A	No organism detected	Negative	Equivocal (HHV-6A; BWA mapping within repeat region only)
VS167	Resp (NPA)	<5	Tracheostomy, NIV, increasing secretions	N/A	Rhinovirus detected	Negative	Negative
VS177	Resp (NPA)	<5	LRTI, CXR changes	N/A	Meta-pneumo-virus detected	Negative	Negative
VS183	Resp (NPA)	<5	Viral infection	N/A	Meta-pneumo-	Positive	Positive (HHV-6B; 93 reads;

						virus detected		BWA 1% coverage)
VS200	Resp (NPA)	<5	DKA, increased respiratory secretions,	N/A	RSV detected	Negative	Positive (HHV-6A; 7077 reads; BWA 1% coverage)
VS207	Resp (BAL)	<5	Respiratory arrest	N/A	Rhinovirus detected	Negative	Positive (HHV-6A; 11305 reads; BWA 1% coverage)
VS216	Resp (NPA)	<5	LRTI,? TB	N/A	No organism detected	Negative	Equivocal (HHV-6A; BWA mapping within repeat region only)

491 BAL = broncho-alveolar lavage; BWA = Burrows-Wheeler Aligner (33); DKA =
492 diabetic ketoacidosis; LRTI = lower respiratory tract infection; NIV = non
493 invasive ventilation; NGS = next generation sequencing; NPA = naso-
494 pharyngeal aspirate; N/A = not applicable; TB = tuberculosis; VP = ventriculo-
495 peritoneal. Read counts and thresholds for positivity are shown in Fig 4.
496

 FIGURE LEGENDS

Figure 1: Schematic summary of work flow and output of HHV-6 testing

among clinical samples from a UK teaching hospital. A: Flow diagram

showing number of samples submitted to a hospital diagnostic microbiology

laboratory with a clinical request for HHV-6 testing (sample ID's prefixed with

HHV). *'Wrong test' indicates request for HHV-6 antibody (rather than PCR).

B: Flow diagram showing consecutive random samples (sample ID's prefixed

with VS) screened for herpesviruses (i) by PCR with HHV-6 confirmed by

sequencing, and (ii) by a metagenomic approach. These samples had been

submitted to the clinical laboratory for other reasons, and had reached the

end of their diagnostic testing pathway.

Figure 2: Results of clinical laboratory testing for HHV-6 in blood and

CSF samples at the request of the clinical team. A: Relationship between

presence or absence of Graft vs Host Disease (GvHD) and HHV-6 status; P

value by Fisher's exact test. B: Relationship between presence or absence of

a clinical Central Nervous System (CNS) syndrome and HHV-6 status; P

value by Fisher's exact test. C: HHV-6 viral loads in blood and CSF; P value

by Mann Whitney test; individuals who died are shown in red. In all three

panels, the numbers at the top of each column show the total number of

patients represented.

Figure 3: Multi-layered pie charts generated to visualize the metavirome

from a respiratory sample taken from a child. Krona was used to generate

the metagenomic visualizations of these data (34). The sample was a

nasopharyngeal aspirate taken from patient ID VS183 (a child age <5 with a

clinical syndrome described by the requesting clinician as 'viral infection'). (A)

HHV-6 B (in pink) shown as a proportion of all viral reads; (B) HHV-6-B (in

pink) shown as a proportion of all virus contigs. The other two predominant

viruses represented in both panels are Torque Teno Mini Virus (in red), a

ubiquitous and probably non-pathogenic virus, and Enterobacteria phage Phi

X (in purple), which is an artefact of the sequencing method (spike used for

positive control in Illumina sequencing run); these illustrate a high proportion

of sequence reads generated from organisms that are not clinically significant.

Figure 4: Detection of HHV-6 DNA in unselected CSF and respiratory

samples. A: Read numbers determined by next generation sequencing

(NGS) according to sample type. The y-axis refers to the number of uniquely

mapping HHV6 reads, representing deduplicated read numbers (Q>30). The

areas shaded green, blue and yellow represent suggested thresholds for

samples to be classified as negative, equivocal or positive, respectively. White

gaps between the colours illustrate that the exact position of the boundaries

between these areas are uncertain. B: Plots generated with Burrows-Wheeler

Aligner (BWA) to illustrate coverage of HHV-6 genomes against consensus.

The x-axis of the plot represents the full length genome for HHV-6, with HHV-

6A shown in triangles, and HHV-6B shown in circles. From a total of 20

samples tested, we show data for samples from which we generated any

HHV-6 reads. The beginning position of each read is indicated. The grey bars

indicate repeat regions (low variability) as defined in the methods; for a

sample to be considered positive, we stipulated that the reads should fall

outside these regions of the genome.

Figure 5: Suggested algorithm showing process of determining the

significance of an organism identified from a clinical sample by next

generation sequencing (NGS). This represents a structure that can be

applied to bioinformatics and clinical interpretation of metagenomic data. We

recognize that the approach and thresholds are different for different

organisms, and that robust output also depends also on optimization of *in vitro*

sample preparation.

SUPPLEMENTARY RESOURCES

Suppl data set 1. Human Herpes Virus 6 (HHV-6): a study of clinical laboratory data and next generation sequencing; DOI: 10.6084/m9.figshare.5671153

This fileset includes the following:

- HHV6 cohort metadata (clinical cohort data to describe patients undergoing clinical testing for HHV-6 infection) as .xlsx and .csv files;
- VS cohort metadata (research cohort data to describe CSF and respiratory samples underdoing screening for HHV-6 infection using PCR and next generation sequencing) as .xlsx and .csv files.

Suppl data set 2. Oxford Screening of CSF and Respiratory Samples ('OSCAR'): Supporting data files for a project using Next Generation Sequencing (NGS) for identification of viruses from clinical laboratory samples; DOI 10.6084/m9.figshare.5670007

This fileset includes the following:

- OSCAR protocol as .pdf file (detailed laboratory methods);
- OSCAR table benefits and challenges (summary of benefits and challenges of NGS approach to clinical samples);
- OSCAR 20 samples metadata (subset of 20 samples from VS cohort for which next generation sequencing was undertaken) as .xlsx and .csv files.

Data note in submission to BMC Research Notes

This short note, currently in submission, comprises an overview of the metagenomic screening project presented here, with links to the methods (as presented above in suppl data set 2) and to the complete dataset in two formats as follows:

- An interactive html file (Krona (34))
- Metagenomic sequences uploaded to European Nucleotide archive (35)

A citation / DOI will be added to this manuscript on acceptance by BMC Research Notes.

ACKNOWLEDGEMENTS

PCM is funded by a Wellcome Trust Intermediate Fellowship, grant number 110110 and received salary from the NIHR during the clinical data collection phase of this work. A project grant to PCM from The British Infection Association (BIA; https://www.britishinfection.org/) covered the costs of laboratory staff and consumables for undertaking PCR and metagenomic screening of CSF and respiratory samples. We are grateful to the Centre for Genomic Research (CGR), University of Liverpool, UK for running the HiSeq platform. Thank you to Dr Peter Muir (Public Health England South West Regional Laboratory, Southmead Hospital, Bristol), who provided valuable advice and discussion during data collection. A subset of these work were presented at the UK Federation of Infection Meeting, 2017 (36). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1. **Salahuddin SZ, Ablashi DV, Markham PD, Josephs SF,**

Sturzenegger S, Kaplan M, Halligan G, Biberfeld P, Wong-Staal F,

Kramarsky B, et al. 1986. Isolation of a new virus, HBLV, in patients

with lymphoproliferative disorders. Science **234:**596-601.

2. **De Bolle L, Naesens L, De Clercq E.** 2005. Update on human

herpesvirus 6 biology, clinical features, and therapy. Clinical

microbiology reviews **18:**217-245.

3. **Adland E, Klenerman P, Goulder P, Matthews PC.** 2015. Ongoing

burden of disease and mortality from HIV/CMV coinfection in Africa in

the antiretroviral therapy era. Front Microbiol **6:**1016.

4. **Abdel Massih RC, Razonable RR.** 2009. Human herpesvirus 6

infections after liver transplantation. World journal of gastroenterology :

WJG **15:**2561-2569.

5. **Loutfy SA, Fawzy M, El-Wakil M, Moneer MM.** 2010. Presence of

human herpes virus 6 (HHV6) in pediatric lymphomas: impact on

clinical course and association with cytomegalovirus infection. Virol J

7:287.

6. **Kawada J, Okuno Y, Torii Y, Okada R, Hayano S, Ando S, Kamiya**

Y, Kojima S, Ito Y. 2016. Identification of Viruses in Cases of Pediatric

Acute Encephalitis and Encephalopathy Using Next-Generation

Sequencing. Sci Rep **6:**33452.

7. **Labska K, Roubalova K, Picha D, Maresova V.** 2015. Presence of

herpesvirus DNA in cerebrospinal fluid of patients with tick-borne

encephalitis and enteroviral meningoencephalitis. Journal of medical

virology **87:**1235-1240.

8.   **Ong DSY, Bonten MJM, Spitoni C, Verduyn Lunel FM, Frencken**

JF, Horn J, Schultz MJ, van der Poll T, Klein Klouwenberg PMC,

Cremer OL, Molecular D, Risk Stratification of Sepsis C. 2017.

Epidemiology of Multiple Herpes Viremia in Previously

Immunocompetent Patients With Septic Shock. Clinical infectious

diseases : an official publication of the Infectious Diseases Society of

America **64:**1204-1210.

9.   **Ogata M, Satou T, Kadota J, Saito N, Yoshida T, Okumura H, Ueki**

T, Nagafuji K, Kako S, Uoshima N, Tsudo M, Itamura H, Fukuda T.

2013. Human herpesvirus 6 (HHV-6) reactivation and HHV-6

encephalitis after allogeneic hematopoietic cell transplantation: a

multicenter, prospective study. Clinical infectious diseases : an official

publication of the Infectious Diseases Society of America **57:**671-681.

10.   **de Pagter PJ, Schuurman R, Keukens L, Schutten M, Cornelissen**

JJ, van Baarle D, Fries E, Sanders EA, Minnema MC, van der Holt

BR, Meijer E, Boelens JJ. 2013. Human herpes virus 6 reactivation:

important predictor for poor outcome after myeloablative, but not non-

myeloablative allo-SCT. Bone marrow transplantation **48:**1460-1464.

11.   **Schonberger S, Meisel R, Adams O, Pufal Y, Laws HJ, Enczmann**

J, Dilloo D. 2010. Prospective, comprehensive, and effective viral

monitoring in children undergoing allogeneic hematopoietic stem cell

transplantation. Biology of blood and marrow transplantation : journal of

the American Society for Blood and Marrow Transplantation **16:**1428-

1435.

12.   **Betts BC, Young JA, Ustun C, Cao Q, Weisdorf DJ.** 2011. Human

herpesvirus 6 infection after hematopoietic cell transplantation: is

routine surveillance necessary? Biology of blood and marrow

transplantation : journal of the American Society for Blood and Marrow

Transplantation **17:**1562-1568.

13.   **Toriumi N, Kobayashi R, Yoshida M, Iguchi A, Sarashina T, Okubo**

H, Suzuki D, Sano H, Ogata M, Azuma H. 2014. Risk factors for

human herpesvirus 6 reactivation and its relationship with syndrome of

inappropriate antidiuretic hormone secretion after stem cell

transplantation in pediatric patients. Journal of pediatric

hematology/oncology **36:**379-383.

14.   **Mori Y, Miyamoto T, Nagafuji K, Kamezaki K, Yamamoto A, Saito**

N, Kato K, Takenaka K, Iwasaki H, Harada N, Abe Y, Teshima T,

Akashi K. 2010. High incidence of human herpes virus 6-associated

encephalitis/myelitis following a second unrelated cord blood

transplantation. Biology of blood and marrow transplantation : journal of

the American Society for Blood and Marrow Transplantation **16:**1596-

1602.

15.   **Ogata M, Oshima K, Ikebe T, Takano K, Kanamori H, Kondo T,**

Ueda Y, Mori T, Hashimoto H, Ogawa H, Eto T, Ueki T, Miyamoto T,

Ichinohe T, Atsuta Y, Fukuda T. 2017. Clinical characteristics and

outcome of human herpesvirus-6 encephalitis after allogeneic

679   hematopoietic stem cell transplantation. Bone marrow transplantation

680   **52:**1563-1570.

681 16. **Hall CB, Caserta MT, Schnabel K, Shelley LM, Marino AS,**

682   **Carnahan JA, Yoo C, Lofthus GK, McDermott MP.** 2008.

683   Chromosomal integration of human herpesvirus 6 is the major mode of

684   congenital human herpesvirus 6 infection. Pediatrics **122:**513-520.

685 17. **Tweedy J, Spyrou MA, Pearson M, Lassner D, Kuhl U, Gompels**

686   **UA.** 2016. Complete Genome Sequence of Germline Chromosomally

687   Integrated Human Herpesvirus 6A and Analyses Integration Sites

688   Define a New Human Endogenous Virus with Potential to Reactivate

689   as an Emerging Infection. Viruses **8**.

690 18. **Pellett PE, Ablashi DV, Ambros PF, Agut H, Caserta MT,**

691   **Descamps V, Flamand L, Gautheret-Dejean A, Hall CB, Kamble RT,**

692   **Kuehl U, Lassner D, Lautenschlager I, Loomis KS, Luppi M, Lusso**

693   **P, Medveczky PG, Montoya JG, Mori Y, Ogata M, Pritchett JC,**

694   **Rogez S, Seto E, Ward KN, Yoshikawa T, Razonable RR.** 2012.

695   Chromosomally integrated human herpesvirus 6: questions and

696   answers. Rev Med Virol **22:**144-155.

697 19. **Ablashi D, Agut H, Alvarez-Lafuente R, Clark DA, Dewhurst S,**

698   **DiLuca D, Flamand L, Frenkel N, Gallo R, Gompels UA, Hollsberg**

699   **P, Jacobson S, Luppi M, Lusso P, Malnati M, Medveczky P, Mori Y,**

700   **Pellett PE, Pritchett JC, Yamanishi K, Yoshikawa T.** 2014.

701   Classification of HHV-6A and HHV-6B as distinct viruses. Arch Virol

702   **159:**863-870.

20. **Lou J, Wu Y, Cai M, Wu X, Shang S.** 2011. Subtype-specific, probe-

based, real-time PCR for detection and typing of human herpesvirus-6

encephalitis from pediatric patients under the age of 2 years. Diagn

Microbiol Infect Dis **70:**223-229.

21. **Fridholm H, Ostergaard Sorensen L, Rosenstierne MW, Nielsen H,**

Sellebjerg F, Bengard Andersen A, Fomsgaard A. 2016. Human

pegivirus detected in a patient with severe encephalitis using a

metagenomic pan-virus array. J Clin Virol **77:**5-8.

22. **Graf EH, Simmon KE, Tardif KD, Hymas W, Flygare S, Eilbeck K,**

Yandell M, Schlaberg R. 2016. Unbiased Detection of Respiratory

Viruses by Use of RNA Sequencing-Based Metagenomics: a

Systematic Comparison to a Commercial PCR Panel. J Clin Microbiol

54:1000-1007.

23. **Lecuit M, Eloit M.** 2015. The potential of whole genome NGS for

infectious disease diagnosis. Expert Rev Mol Diagn **15:**1517-1519.

24. **Lewandowska DW, Zagordi O, Zbinden A, Schuurmans MM,**

Schreiber P, Geissberger FD, Huder JB, Boni J, Benden C, Mueller

NJ, Trkola A, Huber M. 2015. Unbiased metagenomic sequencing

complements specific routine diagnostic methods and increases

chances to detect rare viral strains. Diagn Microbiol Infect Dis **83:**133-

138.

25. **Mulcahy-O'Grady H, Workentine ML.** 2016. The Challenge and

Potential of Metagenomics in the Clinic. Front Immunol **7:**29.

26. **Bukowska-Osko I, Perlejewski K, Nakamura S, Motooka D,**

Stokowy T, Kosinska J, Popiel M, Ploski R, Horban A, Lipowski D,

Caraballo Cortes K, Pawelczyk A, Demkow U, Stepien A,

Radkowski M, Laskus T. 2017. Sensitivity of Next-Generation

Sequencing Metagenomic Analysis for Detection of RNA and DNA

Viruses in Cerebrospinal Fluid: The Confounding Effect of Background

Contamination. Adv Exp Med Biol **944:**53-62.

27.   **Collot S, Petit B, Bordessoule D, Alain S, Touati M, Denis F,**

Ranger-Rogez S. 2002. Real-time PCR for quantification of human

herpesvirus 6 DNA from lymph nodes and saliva. J Clin Microbiol

40:2445-2451.

28.   **VanDevanter DR, Warrener P, Bennett L, Schultz ER, Coulter S,**

Garber RL, Rose TM. 1996. Detection and analysis of diverse

herpesviral species by consensus primer PCR. J Clin Microbiol

34:1666-1671.

29.   **Simmonds P.** 2012. SSE: a nucleotide and amino acid sequence

analysis platform. BMC Res Notes **5:**50.

30.   **Colombier MA, Amorim S, Salmona M, Thieblemont C, Legoff J,**

Lafaurie M. 2017. HHV-6 reactivation as a cause of fever in

autologous hematopoietic stem cell transplant recipients. J Infect

75:155-159.

31.   **Ljungman P, de la Camara R, Cordonnier C, Einsele H, Engelhard**

D, Reusser P, Styczynski J, Ward K, European Conference on

Infections in L. 2008. Management of CMV, HHV-6, HHV-7 and

Kaposi-sarcoma herpesvirus (HHV-8) infections in patients with

hematological malignancies and after SCT. Bone marrow

transplantation **42:**227-240.

32.   **Lu J, Breitwieser FP, Thielen P, Salzberg SL.** 2017. Bracken:

estimating species abundance in metagenomics data. PeerJ Comput.

Sci. **3:e104; DOI 10.7717/peerj-cs.104**.

33.   **Li H, Durbin R.** 2009. Fast and accurate short read alignment with

Burrows-Wheeler transform. Bioinformatics **25:**1754-1760.

34.   **Ondov BD, Bergman NH, Phillippy AM.** 2011. Interactive

metagenomic visualization in a Web browser. BMC Bioinformatics

12:385.

35.   European Nucleotide Archive; https://www.ebi.ac.uk/ena. [Online.]

36.   **McNaughton A, Golubchik T, Sharp C, Gregory WF, Selvaratnam**

M, Mirea A, Klenerman P, Jeffery K, Matthews PC. 2017. Oxford

screening of CSF and Respiratory Samples ('OSCAR'): comparison of

routine laboratory diagnostics, multiplex PCR and next generation

sequencing for identification of viruses from clinical samples.

F1000Research 2017, 6:2130 (poster) (doi:

10.7490/f1000research.1115150.1).