Title

Ancient tropical extinctions contributed to the latitudinal diversity gradient

Authors

Andrea S. Meseguer[1,2]†* and Fabien L. Condamine[2]†*

Affiliations

[1]INRA, UMR 1062 CBGP (INRA, IRD, CIRAD, Montpellier SupAgro), 755 Avenue du

Campus Agropolis, 34988 Montferrier-sur-Lez, France.

[2]CNRS, UMR 5554 Institut des Sciences de l'Evolution (Université de Montpellier), Place

Eugène Bataillon, 34095 Montpellier, France.

†These authors contributed equally.

Running title

Asymmetric gradient of tropical extinction

21 *Corresponding authors (*)*

22 Andrea   S.   Meseguer   (asanchezmeseguer@gmail.com),   Fabien   L.   Condamine

23 (fabien.condamine@gmail.com)

Abstract

Biodiversity currently peaks at the equator and decreases towards the poles. Accumulating fossil evidence reveals that this hump-shaped latitudinal diversity gradient (LDG) has not been persistent through time, but there was comparable species diversity across latitudes such that the LDG flattened during greenhouse periods. Rather than seeing the LDG as the result of limited dispersal, higher equatorial diversification and turnover in Holarctic regions, we propose the modern-steep LDG be explained by climate-driven extinctions of warm-adapted clades from the Holarctic during coldhouse periods, coupled with equator-ward dispersals of organisms that tracked their climatic preferences when tropical biomes got restricted to the equator. Elevated rates of speciation and pole-ward dispersals, conversely, could explain the formation of a flat LDG during greenhouse periods. We studied the LDG of Testudines, Crocodilia and Squamata and found the high latitudes have been a source of tropical diversity for these groups. During greenhouse periods, diversification rates were equivalent between latitudes. During the transition to cold climates, we detected important levels of extinction for ancient tropical lineages distributed at high latitudes, coupled with high levels of range contractions and equator-ward dispersals. Our results further demonstrate the inclusion of fossils in macroevolutionary studies allows detecting the signal of ancient high-latitude tropical extinctions, otherwise vanish from analyses based on present-day data. Our results challenge the current consensus, which considers the evolutionary processes behind the LDG have acted constantly over time and suggest the LDG should be regarded as a time-varying biodiversity pattern.

Keywords: climate change; fossils; biodiversity; extinction; speciation; dispersal; tropics; Holarctic; phylogeny; niche.

Introduction

The current increase of species richness occurring from the poles towards the equator, referred to as the latitudinal diversity gradient (LDG), is one of the most conspicuous patterns in ecology and evolution[1]. This pattern has been described for microbes[2], insects[3], vertebrates[4], and plants[5], and for marine, freshwater, and terrestrial ecosystems[1,6,7].

During decades, it has been considered that the current steep shape of the LDG (with higher diversity at the equator) rooted back several million years ago to the respective origin of the groups, and has been maintained through time[7]. This idea was grounded on previous studies of the fossil record suggesting that the steep LDG has persisted through the Phanerozoic, the last 540 Myrs[8,9]. However, methodological limitations on fossil sampling cast doubts on this conclusion[10,11]. Analyses controlling for sampling biases suggest that the LDG of many groups was less pronounced than the one observed today or was even flat during some periods in the past, *i.e.* there was comparable species diversity across latitudes (see ref.[12] for a review). This is the case for groups like non-avian dinosaurs[10], mammals[11,13], tetrapods[14], insects[15-17], brachiopods[18,19], bivalves[20], coral reefs[21], crocodiles[22], turtles[23,24], or plants[25,26].

The emerging pattern from fossil studies suggests the steep modern LDG was restricted to the relatively few and short coldhouse intervals of the Earth history: the Ordovician/Silurian, the Carboniferous/Permian, the end of the Jurassic, and the Neogene[12,27-29]. Most of the Phanerozoic was characterized by warm greenhouse climates accompanied by a less pronounced (more flatten) LDG[10,11,13,16] (**Fig. 1**), such that the LDG shape fluctuates following climatic transitions. The new fossil evidence set a starting point on LDG research. Most evolutionary hypotheses published to date seek to understand why diversity did not accumulate in the Holarctic at the same levels than in the equator[7,30,31], while recent fossil

Figure 1 | Changes in global temperature and extension of the tropical belt during the Cenozoic, in relation with the shape of the LDG. During the early Cenozoic, global temperatures were higher than the present and warm paratropical conditions extended over northern and southern latitudes. From the early Eocene climatic optimum (EECO; ca. 53-51 Ma), a global cooling trend was intensified on Earth, punctuated by sharp declines of temperatures, such as the dramatic drop of the terminal Eocene Event (TEE, ca. 33.9 Ma). Warm-equable regimes got then restricted to the equator. The Neogene is characterized by a continuous decrease of temperatures, punctuated with periods of relative warmth, such as the middle Miocene climatic optimum, but also by sharp declines during the late Miocene cooling events. Neogene climate cooling culminated with major ice-sheet expansions during Pleistocene glaciations. The LDG of many groups evolved following these global changes[12]; during the greenhouse period diversity was uniform across latitudes, such that the LDG flattened, while during cold periods diversity peaked at the equator (a steep LDG). The question mark denotes the focus of this study, which is to unveil the processes that mediated the transition between a flat and steep LDG. The relative temperature curve of the Cenozoic is adapted from ref.[59]. Maps represent the extension of the tropical belt and Earth tectonic changes as derived from refs.[57,58].

investigations show that, for many lineages, we rather need to explain why, how and when

diversity was lost at high latitudes to shape the current LDG.

Phylogenetic investigations proposed potential mechanisms to explain the transition

from high to low Holarctic diversity. Recent studies consider the high latitudes as a possible

source of diversity, and suggest the elevated species richness in the tropics today could be due

to differential dispersal (d) from high to low latitudes, following the contraction of the

tropical biome[32-35]. Disproportionate high-latitude extinctions – a scenario in which extinction

(μ) overcomes speciation (λ), resulting in negative net diversification rates ($r = \lambda - \mu; r < 0$)

– could also explain this transition. Often considered behind the LDG[36], diversity losses in the

Holarctic have been suggested early in the literature and attributed to Pleistocene

glaciations[37], but this hypothesis was strongly challenged given that the LDG has more

ancient origins than the Pleistocene[7,12]. More ancient extinctions were also considered[38-42].

For example, Hawkins *et al.*[41] proposed the avian LDG resulted from the differential

extirpation of older warm-adapted clades from the temperate regions newly formed in the

Neogene. Unfortunately, the difficulty to infer extinction rates from present-day data[43,44] and

the assumption that diversity in the Holarctic was always low in comparison with equatorial

diversity have made this hypothesis seldom considered (but see ref.[45]). On the contrary,

widely accepted views of equatorial regions as the source of world diversity[46,47] and the long-

term persistence of the steep LDG, permeate the current consensus explaining the LDG as the

result of higher diversification and limited dispersal out of the equator[7,30,48] (**Table 1**).

Meanwhile, elevated turnover rates (comparable speciation and extinction rates) have been

proposed to maintain the low diversity in the Holarctic through time: *e.g.* amphibians[49],

birds[31,36,50], butterflies[32], conifers[51], fishes[52], mammals[48], and squamates[34].

From a flat to hump-shaped LDG

Table 1. Predictions of the most common LDG hypotheses, including the *Asymmetric gradient of tropical extinction* (AGE) hypothesis proposed in this study. The AGE model includes various diversification and dispersal parameters for each temporal interval (greenhouse, coldhouse, transition). Abbreviations: λ = speciation, μ = extinction, d = dispersal, t = time, e = equatorial, h = Holarctic, Trop = Tropical biome, Temp = Temperate biome.

| | *Cradle of diversity* (Stebbins 1974) | *Museum of diversity* (Stebbins 1974) | *Time for speciation* (Stephens & Wiens 2003)[1] | *Tropical niche conservatism* (Wiens & Donoghue 2004) | *Out of the tropics* (Jablonsky et al. 2006) | *Into the tropics* (Rolland et al. 2015) | *Holarctic turnover* (Weir & Schluter, 2007) | *Asymmetric gradient of extinction* | | |
								Greenhouse	Transition	Coldhouse
Predictions	$r_e > r_h$ $(\lambda_e > \lambda_)$ $d_{eh} = d_{he}$	$r_e > r_h$ $(\mu_e < \mu_h)$ $d_{eh} = d_{he}$	$t_{trop} = t_{temp}$ $(r_{trop} = r_{temp})$ $(d_{trtp} = d_{tptr})$	$r_e = r_h$ $d_{eh} > d_{he}$	$r_e > r_h$ $(\lambda_e > \lambda_h)$ $(\mu_e < \mu_h)$ $d_{eh} > d_{he}$	$r_e = r_h$ $d_{eh} < d_{he}$	$r_h = 0$ $(\mu_h = \lambda_h)$ $d_{eh} = d_{he}$	$r_e = r_h$ $d_{eh} = d_{he}$	$r_e > r_h$ $(\mu_h > \lambda_h)$ $d_{eh} < d_{he}$	$r_e > r_h$ $(r_h = 0)$ $d_{eh} = d_{he}$

[1] No specific predictions on diversification differences across regions are provided in this hypothesis but across taxa adapted to particular climates, as it assumes that tropical-adapted clades originated before temperate clades, and thus had more time to diversify than temperate clades.

Across Earth history, the extension of the tropical biome has fluctuated geographically around

the equator, with periods of pole-ward expansions when warm paratropical conditions and

tropical-adapted taxa appeared at high latitudes[53-56], followed by periods of equatorial-ward

contractions[57,58] (**Fig. 1**). The last 100 Myrs have witnessed the last transition, with a global

cooling trend set on Earth since the Cretaceous–early Cenozoic period that culminated with

the Pleistocene glaciations[59].

The expansion of tropical-like conditions to the higher latitudes induced species

diversification in the new paratropical areas (**Fig. 2a**) and facilitated movements within the

broad "*paleotropical belt*", such that equatorial clades were able to disperse "*out of the*

equator" into high-latitude warm regions[28,46] – or towards the south in the case of northern

origins. A greenhouse equable climate thus would have triggered the formation of a flat LDG

(**Fig. 2a**). On the contrary, tropical biome retractions following climate cooling (**Fig. 2b**)

induced periods of declining diversity at high latitudes (where climate change was more

intensively felt), and biotic movements "*into the equator*" prevailed over northern migrations.

Higher extinction rates impacted tropical lineages at high latitudes, whereas low-latitude

tropical lineages experienced lower extinction. Climate change would have thus driven an

asymmetric extinction pattern within the tropical biome, and mediated the formation of a

steep LDG shape (**Fig. 2c**). Biodiversity dynamics may shift between these two patterns

during climatic transitions. This framework imposes a time-variable vision of the LDG in

which prevailing speciation, extinction and dispersal dynamics switch during warm and cool

intervals. Therefore we propose the hypothesis that the current hump-shaped LDG results

from a combination of ancient tropical extinctions in high latitudes and equator-ward

dispersals, both due to the global cooling of the Cenozoic. It would have led to diversity loss

in the Holarctic and southern temperate regions, and contribute to the transition from a flat

LDG to a steep LDG with lower diversity towards the south and north.

Figure 2 | Prevalent evolutionary processes behind the latitudinal diversity gradient under the AGE time-variable framework. The graphic shows the hypothetic change in evolutionary dynamics (diversification "r", speciation "λ", extinction "μ" and dispersal "d") between Holarctic and equatorial regions (denoted with "h" and "e" subscripts, respectively) and across main climatic intervals: **a**, the greenhouse late Cretaceous-early Cenozoic period. **b**, the late Eocene-Oligocene climatic transition. **c**, the Neogene coldhouse interval. For each period, inset Figs. represent the distribution of species richness across latitudes (the LDG shape) and the hypothetical change in global evolutionary dynamics under the AGE hypothesis. In the centre, the average model represents the prevalent evolutionary dynamics expected under the AGE hypothesis under a single, constant rate, evolutionary model.

We refer this hypothesis to the "Asymmetric Gradient of Extinction" (AGE). The

predictions (P) of the AGE model could be formulated as follows (**Fig. 2; Table 1**): **P1**;

similar diversification rates between Holarctic ($_H$) and equatorial ($_E$) regions during

greenhouse periods ($r_E = r_H$), **P2**; similar dispersal rates between Holarctic and equatorial

regions during greenhouse periods ($d_{HE} = d_{EH}$), **P3**; extinction increases relative to

speciation at high latitudes during climatic transitions towards cooler climates, *i.e.* declining

diversity ($r_H < 0$), while equatorial regions remain stable, and **P4**; dispersal "*into the*

equator" being more important than "*out of the equator*" during climatic transitions

($d_{HE} > d_{EH}$).

The AGE hypothesis builds up from previous ideas but here framed in the context of a

time-variable scenario. As noted earlier, the idea of tropical lineages occurring at higher

latitudes going extinct during climatic transitions has been suggested before[38,39,41,42].

Similarly, behind this hypothesis prevails the idea of niche conservatism[30,47]: when the

tropical biome retreated towards the equator, most of the tropical-adapted taxa at high

latitudes were unable to adapt and either went extinct or got their distribution restricted. The

dynamic history of the tropical belt has been considered before to explain different dispersal

opportunities for tropical lineages through time[32,35,60,61], as well as the wider extension of the

paleotropical belt to explain the large number of groups originated in the tropics, as the

product of increased area size[62]. To our knowledge, however, diversity losses mediated by

tropical biome contractions have seldom been considered a major mechanism behind the

LDG. This idea might seem quite straightforward, but its direct formulation and corroboration

has been lacking. In addition, although it is widely accepted that the processes shaping

biodiversity varied over time and space[63], this has been overlooked in the context of the LDG,

generally explained by the action of time-constant process[33,34,45,64]. Here, we studied the LDG

of Testudines, Crocodilia and Squamata using comparative methods for both the phylogeny

and fossil record, and we confronted the AGE hypotheses to classical LDG hypotheses.

Results

Phylogeny-based diversification analyses: are diversification rates higher in the equator?

According to current distribution data, species richness of turtles, squamates and crocodiles

peaks near the equator, with 84% of all extant species living in the tropics, whereas only 15%

live in temperate regions and 1% span both biomes. We categorized each species reported in

the phylogeny (Supplementary Tables 1-3) as living in the equator (tropical biome) or the

Holarctic and Southern Hemisphere (temperate biome). In turtles, we found that there are 239

tropical species, 84 temperate species and 6 widespread species (7 are marine); while in

squamates, there are 7955 tropical species, 1337 temperate species and 124 widespread. For

the species-poor crocodile clade, there are 23 tropical species and 2 temperate species.

We analyzed differences in diversification rates between Holarctic and equatorial

regions for turtles, squamates and crocodiles using the Binary State change Speciation and

Extinction model (BiSSE[65,66], see *Methods*). We did not use the Geographic State change

Speciation and Extinction model[67], which is appropriate when dealing with widespread

species, because most species in our datasets are endemic to a region and to consider a

character state in SSE models, it must represent at least 10% of the total diversity[68]. We did

not apply the BiSSE model on crocodiles because simulation studies have shown that trees

containing fewer than 300 species may lack sufficient phylogenetic signal to produce enough

statistical power[68].

First, we used the time-constant BiSSE model as is generally applied to studies of the

LDG[33,34,35,45,48]. For turtles, net diversification rates were found higher in the Holarctic than in

the equator (**Table 2**, Supplementary Fig. 1a), although differences were not significant, and

Table 2. Results of the diversification and biogeographic analyses performed in this study. Abbreviations: λ = speciation, μ = extinction, d = dispersal, $R\mu$ = range extirpations, e = equatorial, h = Holarctic.

	Turtles			Squamates			Crocodiles		
	Greenhouse	**Transition**	**Coldhouse**	**Greenhouse**	**Transition**	**Coldhouse**	**Greenhouse**	**Transition**	**Coldhouse**
PyRate	$\lambda_e > \lambda_h$ $\mu_e > \mu_h$ $r_e = r_h$	$\lambda_e > \lambda_h$ $\mu_e < \mu_h$ $r_e > r_h$	$\lambda_e > \lambda_h$ $\mu_e > \mu_h$ $r_e > r_h$	$\lambda_e > \lambda_h$ $\mu_e > \mu_h$ $r_e = r_h$	$\lambda_e = \lambda_h$ $\mu_e > \mu_h$ $r_e < r_h$	$\lambda_e > \lambda_h$ $\mu_e < \mu_h$ $r_e > r_h$	$\lambda_e > \lambda_h$ $\mu_e > \mu_h$ $r_e = r_h$	$\lambda_e > \lambda_h$ $\mu_e < \mu_h$ $r_e > r_h$	$\lambda_e > \lambda_h$ $\mu_e > \mu_h$ $r_e > r_h$
BiSSE (time-variable)	$\lambda_e = \lambda_h$ $\mu_e = \mu_h$ $d_{he} = d_{eh}$	$\lambda_e = \lambda_h$ $\mu_e = \mu_h$ $d_{he} > d_{eh}$	$\lambda_e < \lambda_h$ $\mu_e = \mu_h$ $d_{he} > d_{eh}$	$\lambda_e > \lambda_h$ $\mu_e < \mu_h$ $d_{he} < d_{eh}$	$\lambda_e > \lambda_h$ $\mu_e < \mu_h$ $d_{he} > d_{eh}$	$\lambda_e < \lambda_h$ $\mu_e = \mu_h$ $d_{he} > d_{eh}$	-	-	-
BiSSE (constant)		$\lambda_e < \lambda_h \, \mu_e < \mu_h, r_e < r_h$, $d_{he} > d_{eh}$			$\lambda_e < \lambda_h \, \mu_e < \mu_h, r_e > r_h, d_{he} > d_{eh}$			-	
DEC		$R\mu_e < R\mu_h \, (R\mu_e < R\mu_h)^*$ $d_{he} < d_{eh} \, (d_{he} < d_{eh})^*$			$R\mu_e > R\mu_h \, (R\mu_e = R\mu_h)^*$ $d_{he} < d_{eh} \, (d_{he} = d_{eh})^*$			$R\mu_e = R\mu_h \, (R\mu_e = R\mu_h)^*$ $d_{he} = d_{eh} \, (d_{he} = d_{eh})^*$	
DEC fossil		$R\mu_e < R\mu_h \, (R\mu_e < R\mu_h)^*$ $d_{he} > d_{eh} \, (d_{he} > d_{eh})^*$			$R\mu_e > R\mu_h \, (R\mu_e < R\mu_h)^*$ $d_{he} < d_{eh} \, (d_{he} > d_{eh})^*$			$R\mu_e < R\mu_h \, (R\mu_e < R\mu_h)^*$ $d_{he} > d_{eh} \, (d_{he} > d_{eh})^*$	

* Prevalent dynamics when the number of events is calculated relative to the number of taxa currently distributed in each region

dispersal '*into the equator*' was 10-fold higher than '*out of the equator*'. For squamates, the

same dispersal pattern is recovered but net diversification rates are significantly higher in the

equator (Supplementary Fig. 1b). To test the AGE hypothesis, we introduced two shift times

at 51 and 23 Ma to detect different diversification dynamics between greenhouse, transitional,

and coldhouse periods. This model indicates that net diversification of turtles was equivalent

between the Holarctic and equator, while it was lower in the Holarctic for squamates (**Table**

2, Supplementary Fig. 2) until the coldhouse period, when Holarctic diversification increased.

Dispersal was estimated symmetrical between regions (*into the equator = out of the equator*)

during greenhouses, and asymmetrical (*into the tropics > out of the tropics*) during the

climatic transition and coldhouse period. We inferred the same patterns by applying the same

model but with different combination of shift times (51 Ma and 34/23 Ma; Supplementary

Fig. 3).

Estimations of ancestral origins: did groups originate preferentially in the equator?

To further test the predictions of the AGE hypothesis, we performed biogeographic analyses

using the Dispersal-Extinction-Cladogenesis (DEC) model[69] (see *Methods*). We first analyzed

the data using an unconstrained DEC analysis in which all ranges comprised of three areas

can be an ancestral state. We inferred an equatorial distribution for most basal nodes of turtles

and squamates, from where lineages colonized the other regions (**Fig. 3a**, **Table 2**,

Supplementary Fig. 4). Crocodile ancestors were reconstructed widespread during the

Cretaceous, but an early vicariant event separated *Alligator* in the Holarctic from the other

genera in equatorial regions (Supplementary Fig. 5).

The biogeographic estimates based on extant data only conflict with the fossil record

of turtles, squamates, and crocodiles[22-24,70]. To overcome this bias, we introduced information

about the distribution of fossils in DEC, in the form of hard (HFC) and soft (SFC) geographic

(a) Unconstrained biogeographical model

(b) Fossil-informed biogeographical model

Pleurodira

Cryptodira

Chelidae
Podoc.
Trionychidae
Cheloniidae
Emydidae
Testudinidae
Geoemydidae

Range dispersal event
Range extinction event
Range inheritance scenario
Ancestral range reconstruction:
Northern temperate regions
Equatorial and tropical regions
Southern temperate regions

Tr | J | K | Pg | Ng

Figure 3 | Biogeographic reconstruction of Testudines showing the effects of the incorporation of fossil information into biogeographic inference. **a**, Biogeographic reconstruction inferred with DEC based on the distribution of extant taxa. **b**, Biogeographic reconstruction under the fossil-informed HFC (*hard fossil constraint*) model. Coloured circles at tips and nodes represent current and ancestral ranges, respectively, while squares represent range inheritance scenarios. Colours correspond with the discrete areas in the legend. Black circles indicate fossil range constraints included in the analysis, with numbers corresponding with taxa in Table Supplementary S4. The reconstruction under the *soft fossil constraint* (SFC, see text) model is presented in Supplementary Fig. 7.

fossil constraints placed at specific nodes (see *Methods*; Supplementary Tables 4-6). By

including fossil information, we recovered very different biogeographic histories for the three

groups (**Table 2**;turtles: **Fig. 3**, Supplementary Fig. 7; squamates: Supplementary Figs. 8, 9;

and crocodiles: Supplementary Figs. 10, 11). We found that turtles originated in the Northern

Hemisphere under the SFC model (they are widespread in the HFC model), from where

lineages migrated into the equator and southern regions (**Fig. 3b**, Supplementary Fig. 7). Most

dispersal is thus '*into the equator*' (Supplementary Fig. 6, Supplementary Table 7). We also

detected a higher number of geographic extinctions when fossil ranges were considered,

affecting predominantly turtle lineages in the Holarctic (53 and 11 lineages disappeared from

the region under the HFC and SFC model, respectively) and in southern temperate regions (9

in the HFC model; Supplementary Fig. 6, Supplementary Table 8). The same trend is

observed when the number of extinction/dispersal events is controlled by the number of

lineages currently distributed in each region (**Fig. 4**).

Squamates are recovered widespread at their origin in both SFC and HFC analyses

(Supplementary Figs. 8, 9). During the greenhouse period, dispersal '*into the equator*' equals

(or overcomes, in the HFC model) dispersal '*out of the equator*', and afterwards dispersals

'*out of the equator*' prevail (Supplementary Fig. 6, Supplementary Table 7). Range extinction

is estimated very high in this group under the unconstrained model with 30 lineages extirpated

from the Holarctic, 2 from southern temperate regions and 152 from the equator

(Supplementary Fig. 6, Supplementary Table 8). Under the fossil-informed models, the

number of Holarctic extinctions increases (to 109 and 66 lineages in the HFC and SFC

models, respectively), while the lineages extirpated from the equator remains relatively stable

(144 and 109 from the equator in HFC and SFC models; Supplementary Fig. 6). When the

number of extinction is controlled by the actual number of lineages distributed in each region,

the number of Holarctic extinctions and dispersal '*into the equator*' increases dramatically

Figure 4 | Estimated number of range-extinction and dispersal events through time. Analyses were performed for Testudines, Squamata and Crocodiles under the *unconstrained model* (Unc.), based on present evidence only, and the fossil-based *hard* (HFC) and *soft fossil constraint* (SFC) biogeographic models. **a**, Inferred number of range extinction events through time and across regions relative to the number of lineages currently distributed in each region. The global mean temperature curve is modified from ref.[59]. **b**, Inferred number of dispersal events from the Holarctic into the equator (IntoEq) and out of the equatorial zone (OutEq), relative to the current number of lineages distributed in the Holarctic and equatorial zones, respectively. Abbreviations: Tr, Triassic; J, Jurassic; K, Cretaceous; Pg, Paleogene; and Ng, Neogene, Trans. = climatic transition, and C. = coldhouse.

Figure 5 | Global pattern of turtles, squamates and crocodiles diversification through time based on the fossil record, and analysed with a Bayesian model. Origination (blue) and extinction (red) rates were estimated using time bins as defined by epochs of the geological timescale (on the top, main climatic periods are shown as follows: Greenhouse, Tran. = climatic transition, and C. = coldhouse). Solid lines indicate mean posterior rates, whereas the shaded areas show 95% credibility intervals. Net diversification rates (black) are the difference between origination and extinction. The vertical lines indicate the boundaries between geological periods and major mass extinction events. Tr, Triassic; J, Jurassic; K, Cretaceous; Pg, Paleogene; and Ng, Neogene.

and overpass equatorial dispersal/extinctions (**Fig. 4**). For crocodiles, including fossil ranges,

we recovered that all the early nodes are distributed in the Holarctic (Supplementary Figs. 10,

11), and we detected range extinctions; four lineages disappeared from the Holarctic, three

from southern temperate regions, and two from the equator (HFC model; Supplementary Fig.

6, Supplementary Tables 7, 8). Only two lineages disappeared from the Holarctic in the SFC

model. The same tendencies are observed when the number of events is controlled by the

current number of lineages distributed in each region (**Fig. 4**).

Fossil-based diversification analyses: evidence for ancient tropical extinctions?

The turtle fossil dataset comprised 4084 occurrences for 420 genera (65 extant and 355

extinct) (Supplementary Table 9). The squamate fossil dataset comprised 4798 occurrences

for 638 genera (120 extant and 518 extinct) (Supplementary Table 10). The crocodile fossil

dataset comprised 1596 occurrences for 121 genera (9 extant and 112 extinct) (Supplementary

Table 11). To infer the global diversification dynamics, we first analysed the fossil datasets as

a whole using a Bayesian approach that infers the temporal dynamics of origination and

extinction rates based on fossil occurrences[71] (see *Methods*). The results of the turtles show

that origination rates peaked during the Jurassic and decreased since then until the present.

Extinction was generally low and constant during the Mesozoic but increased during the

coldhouse intervals of the Jurassic and Paleogene resulting in negative net diversification

during the Paleogene (**Fig. 5**, **Table 2**, Supplementary Figs. 12, 13). For squamates,

origination rates peaked in the Jurassic and the Late Cretaceous, while extinction increased

steadily until the Late Cretaceous. In the Paleogene, net diversification approaches zero,

suggesting a high turnover (**Fig. 5**, Supplementary Figs. 14, 15). Crocodilia origination

peaked in the Early Cretaceous and decreased towards the present, and extinction was

generally low and constant. We also found diversity losses in the Paleogene that extend the

Holarctic *vs.* equatorial

Figure 6 | Global pattern of turtle, squamate and crocodile diversification between Holarctic and equatorial regions, based on the fossil record. Diversification dynamics are compared between fossils distributed in Holarctic and equatorial regions. Origination (blue) and extinction (red) rates were estimated using time bins as defined by the main climatic intervals since the Mesozoic (on the top, climatic periods are shown as follows: Greenhouse, Tran. = climatic transition, and C. = coldhouse). Solid lines indicate mean posterior rates, whereas the shaded areas show 95% credibility intervals. Net diversification rates (black) are the difference between origination and extinction. The vertical lines indicate the boundaries between climatic intervals. Tr, Triassic; J, Jurassic; K, Cretaceous; Pg, Paleogene; and Ng, Neogene

Temperate *vs.* tropical

Figure 7 | Global pattern of turtle, squamate and crocodile diversification across temperate and tropical climates, based on the fossil record. Diversification dynamics are compared between fossils inhabiting under temperate and tropical macroclimates. Origination (blue) and extinction (red) rates were estimated using time bins as defined by the main climatic intervals since the Mesozoic (on the top, climatic periods are shown as follows: Greenhouse, Trans. = climatic transition, and C. = coldhouse). Solid lines indicate mean posterior rates, whereas the shaded areas show 95% credibility intervals. Net diversification rates (black) are the difference between origination and extinction. The vertical lines indicate the boundaries between climatic intervals. Tr, Triassic; J, Jurassic; K, Cretaceous; Pg, Paleogene; and Ng, Neogene.

present, suggesting the crocodiles are declining in diversity since then (**Fig. 5**, Supplementary

Figs. 16, 17).

We performed additional analyses with different subsets of the three fossil datasets to

tease apart the signal of speciation and extinction between geographic regions (equator or

Holarctic) and ecological conditions (temperate or tropical, see *Methods*). These analyses

evidenced that Paleogene diversity losses in turtles and crocodiles are mostly attributed to

species living in the Holarctic and under tropical conditions (**Figs. 6**, **7, Table 2**). The global

diversity loss inferred for crocodiles during the Neogene is attributed to both taxa living in

Holarctic and equatorial regions, and adapted to tropical and temperate conditions, reinforcing

the hypothesis that the group as a whole is in decline. Diversification rates of temperate taxa

are estimated high in all groups during the Oligocene but decrease during the Neogene. For

the equator dataset, we found a decrease in extinction and speciation rates through time, and

more or less constant net diversification (except for squamates, which show a decrease in

diversification during the Paleogene, followed by a diversification increase in the Neogene).

Discussion

The genesis of the current LDG

Recent fossil investigations show that, for many groups, diversity levels at high latitudes were

more elevated than in the present, or even equivalent to diversity levels in the equator, during

some periods of the past[11,12]. This suggests that the LDG should be regarded as a biodiversity

pattern that has changed through time, while the previous studies have generally assumed the

processes shaping the LDG have acted constantly. Limited dispersal to the Holarctic[30], high

Holarctic turnover[34], or high equatorial diversification[31,33,34,45,48,49,64] cannot explain the

formation of a flatten LDG, nor the transition to lower diversity in the Holarctic observed in

many groups[11,12].

To account for temporal changes in the global distribution of biodiversity, we propose

the AGE hypothesis that captures components of previous studies[12,30,33,35,46] in the context of a

new time-variable framework. When confronting the AGE hypothesis with the LDG of

turtles, crocodiles and squamates, this evolutionary scenario receives little support from

analyses based on extant species only (**Table 2**). Biogeographic analyses of turtles support the

'*tropical niche conservatism*' hypothesis, with an equatorial origin and recent invasion of

high latitude regions; thereby lineages had less time to diversify[30] (**Figs. 3**, **4**, **Table 2**), in

agreement with recent investigations[72,73]. Time-constant and time-variable BiSSE analyses

support a higher Holarctic diversification and the '*into the tropics*' hypothesis for the turtle

LDG (Supplementary Figs. 1a, 2a, 3a). For squamates, biogeographic analyses support the

'*out of the tropics*' hypothesis, while BiSSE supports both a higher equatorial diversification

and the '*into the tropics*' hypothesis to explain their LDG (**Fig. 4**, Supplementary Figs. 1-4).

Crocodile biogeographic results support the diversification hypothesis, with higher origination

rates in equator and no effect of dispersal (**Fig. 4**, **Table 2**, Supplementary Fig. 5). In contrast,

results from the fossil-based analyses are overall in agreement with the AGE hypothesis

(**Table 2**).

The AGE hypothesis predicts that "*on average*" (i.e. integrated over time under a

constant-rate evolutionary model), the diversification of a group was lower in the Holarctic

(overall extinction being higher; the "average AGE hypothesis" of **Fig. 2**), as generally

accepted for the origin of the LDG[31,32,50,74]. However, the time-variable scenario proposed

here allows disentangling the relative contribution of speciation, extinction and dispersal in

particular periods for the formation of the LDG. As predicted in the AGE hypothesis (**P1**), we

found equivalent diversification rates for turtles, squamates and crocodiles between Holarctic

and equatorial regions during the greenhouse period (rates overlap in their credibility

intervals; **Fig. 6**; Supplementary Figs. 1-3, 12-17), in agreement with the existence of a flatten

LDG during this phase[12]. We found mixed evidence for equivalent dispersals between

Holarctic and equatorial regions during greenhouse periods (**P2**) for these groups, with BiSSE

showing equal dispersal rates between biomes (**Table 2**, Supplementary Figs. 1-3), but fossil

biogeographic analyses indicating higher dispersal rates '*into the tropics*' (**Fig. 4**,

Supplementary Table 7). This might not necessarily conflict with the AGE hypothesis, but it

rather reflects the idiosyncratic history of these groups with many ancestors distributed in the

Holarctic from where they colonized southern regions.

Subsequently, our analyses reveal that diversification rates decreased in all regions

during the transition to cold climates (**Fig. 5**). Nonetheless, for turtles and crocodiles,

diversification slowdowns were more dramatic in the Holarctic than in the equator, where

extinction exceeded speciation ($r < 0$) and diversity losses occurred (**P3**; **Fig. 6**).

Additionally, our results suggest that turtles and crocodiles had a widespread distribution at

their origins that subsequently contracted to the equator due to elevated levels (both in

absolute and relative terms) of range extirpations at higher latitudes and "*into the equator*"

dispersals[35] during Cenozoic cooling (**P4**; **Figs. 3**, **4**). This result is in agreement with BiSSE

results (Supplementary Figs. S2, S3), and with previous fossil investigations on turtles[23,24,70]

and crocodiles[22,39]. Hence our results suggest the combined effect of widespread Mesozoic

tropical origins, followed by diversity losses and range extirpations at high-latitudes (**Table 2**,

Figs. 6, 7), and elevated dispersal rates "*into the equator*" (**Fig. 4**) during the climatic

transition to cold could have shaped the current steep LDG of turtles and crocodiles.

Paleontological evidence attests that paratropical conditions and associated warm-

adapted taxa disappeared from high latitudes at some point between the mid-late Eocene (51-

34 Ma) to the Neogene (23 Ma)[53,57,58]. Our results with high-latitude extinctions, range-

contractions and southward dispersals peaking by the end of the Paleogene, concur with these

observations, and indicate that the Cenozoic climate cooling had a profound effect on shaping

the current LDG. We cannot discard that diversity losses at high latitudes occurred earlier (as

suggested by our fossil-based diversification analyses with time intervals defined by the main

geological periods; Supplementary Figs. 12, 14, 16), as the result of the K-Pg mass extinction,

because of low temperature excursion brought by the impact winter[75,76]. This scenario,

however, might not conflict with our interpretations because our results are consistent

showing extinction in the high latitudes to have been superior than in the low latitudes,

independently on how time intervals were defined (Supplementary Figs. 12–17). Moreover,

both hypotheses are not mutually exclusive; the number of range extinctions and southward

dispersals increases from the K-Pg to the Neogene (**Fig. 4**), suggesting an additive effect of

K-Pg and Neogene cooling on the depopulation of the Holarctic. Nonetheless, we favour the

former hypothesis because *(i)* Eusuchia and Testudines are thought to be marginally affected

by the K-Pg boundary events[39,77,78]; and *(ii)* evidence for other organisms point towards

diversity recoveries to Cretaceous levels in the high latitudes by the Eocene warming[79,80].

Ancestors of turtles and crocodiles were adapted to tropical conditions since the Late

Cretaceous[39,81]. Our results show that extinction has not been random, but preferably affected

taxa living under a tropical-like climate at high latitudes (**Figs. 6, 7**), which suggest that many

of the Holarctic warm-adapted species were unable to adapt to the new temperate regimes and

went extinct or escaped extinction contracting their ranges to the south (**Fig. 4**).

While our fossil-based results for turtles and crocodiles support the AGE hypothesis,

this scenario is only partially supported for squamates. During Cenozoic cooling, squamates

in the higher latitudes did not go extinct more than in the equator but the opposite, with

diversity losses inferred in the equator (**Fig. 6**). Diversity dynamics for species distributed in

the equator, however, are difficult to trust because the equatorial dataset is the poorest of our

study due to the small number of fossil lineages and the small number of records per lineage

(Supplementary Table 12). As a result, there are uncertainties on the estimated rates with wide

credibility intervals probably due to geographic biases in the fossil record[82]. In the Holarctic,

turnover was very elevated, indicating that species did disappear from high latitudes but were

replaced by a new squamate community. It is possible the number of Squamata species was

always imbalanced between regions. The high Holarctic turnover thus contributing to

maintain this pattern, together with the increases in diversification inferred in the equator

towards the present (**Fig. 6**), as previously hypothesized[34].

Nonetheless, high-latitude diversity losses could not be completely ruled out.

Although in absolute terms the Holarctic received more immigrant species than species

emigrated from the region (Supplementary Fig. 6), the number of species in the equator today

is 4-fold the number of lineages in other areas. When controlled for this imbalance sampling

of species in our tree, a relatively higher number of species lost their ancestral Holarctic

distribution and emigrated towards the equator than the opposite way around (**Fig. 4**). This is

in agreement with our BiSSE results and those of Pyron[34] pointing towards a higher rate of

dispersal '*into the equator*'. Altogether, the current LDG of squamates could results from the

Cenozoic climate cooling through range extinctions and southward dispersals on species

distributed in the Northern Hemisphere (**Fig. 4**), and not necessarily by promoting lineage

extinctions (**Fig. 6**). In the aim to reconcile the phylogeny and fossil record, we used a

number of fossil constrains in our biogeographic analyses of Squamata, but these constraints

were relatively low in comparison with the size of the tree (30 fossils for 4161 nodes). The

few nodes constrained with fossils increased the absolute number of Holarctic range-

extinction and dispersal events '*into the equator*' in comparison with the estimations when

fossils were not considered (from 30 to 109 and from 40 to 124, respectively; Supplementary

Tables 7, 8). Nevertheless, the inclusion of fossils did not significantly alter the number of

events estimated for equatorial taxa. This suggests that a more profound understanding of the

taxonomy of squamate fossils could help to assign more fossils in the tree and detect other

high-latitude range extinctions probably missed in this study.

Reconciling fossil and phylogenetic evidence

Our results unequivocally evidence the inclusion of fossils in macroevolutionary studies

allows detecting the signal of ancient high-latitude extinctions and range extirpations (**Figs. 4-**

6), otherwise hidden from the analyses based on present-day data. Only '*into the equator*'

dispersals were detected with BiSSE models, but not with unconstrained biogeographic

analyses[72,73] (**Figs. 3**, **4**, Supplementary Figs. 4, 5).

This conflict between extant and fossil evidence might not only pervade our study but

also the LDG literature. On the one hand, high extinction rates have seldom been inferred

among tropical lineages (but see ref.[83-85]). Hypotheses on extinction have often focused on

temperate taxa and on recent times, notably on the effects of recent Pleistocene

glaciations[33,34,36,37,48]. When extinction was reported, it was accompanied by elevated

origination rates (high turnover) in high latitude groups[31-34,36,48-51]. Importantly, to our

knowledge, diversity losses ($r < 0$) have never been inferred in phylogenetic studies of the

LDG (note that the negative diversification rates, inferred for the entire history of temperate

bird groups[45], are likely an error of the time-constant BiSSE model, because this result

conflicts with the fact that these group are still extant).

On the other hand, ancient tropical extinction in high latitudes is supported by fossil

studies documenting comparable diversity levels across latitudes during the greenhouse

Paleogene, followed by diversity losses at higher latitudes during the climatic transition. For

instance, Archibald *et al.*[15,16] sampled insect diversity at an Eocene site in Canada, and in

present-day temperate Massachusetts (USA) and tropical sites of Costa Rica. Insect diversity

was higher at the Eocene paleotropical site than the modern temperate locality, and

comparable to the modern-day tropical locality. Post-Eocene insects have thus suffered

greater levels of extinction in the Nearctic than around the equator. This pattern is consistent

with results of other studies on various groups like birds[41,86], invertebrates[42,87], mammals[11,88]

or plants[89,90].

If extinction of tropical clades in the high latitudes was paramount and trait dependent

(tropical-adapted lineages having higher risk of extinction), entire tropical clades could have

disappeared without leaving any phylogenetic footprint. This might be the case for crocodiles,

turtles and squamates, in which extinction preferably affected warm-adapted lineages at high

latitudes during a cooling climate (**Fig. 7**), and in which most evidence of an ancient Holarctic

and southern temperate distributions were erased by species extinctions and range

contractions (**Figs. 4, 6**). Although the recent decade has witnessed many efforts to reconcile

fossil and phylogenetic evidence[91,92], simulation and empirical studies have shown that under

such asymmetric-rate scenarios, phylogenetic comparative methods tend to fail to recover the

correct processes, simply because there is not enough information in the phylogeny to predict

the loss of the focal character along each terminal branch[44,69,93-95]. Consequently, if the LDG

was shaped by asymmetric extinction and dispersal, as we hypothesized here, the diversity

patterns we observe today for many groups might be a poor representation of the historical

diversification processes (**Figs. 3**, **4**), and the inclusion of fossils might be crucial to the study

of the LDG[96]. Although we acknowledge the fossil records are not complete: fossils still

represent the unique evidence of the diversity that existed in the past. In contrast to molecular

phylogenies, the incompleteness of the fossil record have a less problematic effect on the

estimation of speciation and extinction rates because removing a random set of taxa does not

affect the observed occurrences of other lineages[71]. Simulations have shown that PyRate

estimates correctly the dynamics of speciation and extinction rates under low levels of

preservation or severely incomplete taxon sampling.

Studies on the LDG are often based on state-dependent speciation and extinction

models[33,34,35,45,48,97]. Though these models are conceived to test scenarios of differential

diversification and asymmetric transitions as the one suggested here, studies on the LDG

often assumed that diversification parameters remain constant through time[33,34]. If the

evolutionary processes shaping the LDG have varied across latitudes and time, using time-

constant models has little power to test more complex scenarios behind the LDG. Testing our

scenario requires the implementation of time-variable models. When applied to the study of

diversity patterns, these models have revealed notable signatures of extinction among

ancestral tropical plant clades, showing that large temperate groups do not form monophyletic

radiations, but rather polyphyletic clusters of lineages derived from the tropics and masked by

extinction[98]. Incorporating time-shifts in our BiSSE analyses, however, has not completely

reconciled fossil and extant evidence, although the time-variable BiSSE analyses correctly

identified the main direction of dispersal in different periods. While providing the causes and

solutions to this issue goes beyond the purpose of this study, this potential artefact highlights

the importance of fossils in macroevolutionary inferences.

Diversification differences between temperate- and tropical-adapted taxa

A long-standing question in ecology and evolution, directly related to the formation of the

LDG, is whether diversification rates differ between taxa living under temperate or tropical

climates. The metabolic theory of ecology predicts that warm tropical environments prompt

higher speciation rates than colder climates, mainly due to increased rates of molecular

evolution[99], and/or to higher mutational robustness[100]. This theory received support in the

literature[101], but other studies later suggested per capita speciation rates are highest in the

temperate zone[34,36,102,103]. Answering this question requires to consider that the today's

equatorial belt merely represents a fraction of the wider extension that the tropics had in

deeper times and the Holarctic once housed lineages with tropical-like affinities[57,58,62] (**Fig.**

1). Hence, geographic-based diversification results cannot be directly extrapolated to assess

diversification differences between ecological preferences, because if high extinction rates are

estimated for taxa in the Holarctic, extinction could have affected lineages living under both

tropical and temperate conditions (depending on the time interval). In this sense, we think it

has been overlooked that the main hypotheses on the LDG have sought to answer two related

but slightly different questions, which could explain some of the discrepancies found between

ecological and phylogenetic perspectives.

Here, we found that diversification rates of turtles, crocodiles and squamates living

under temperate climatic conditions were significantly higher than rates of tropical-adapted

taxa right after the late Eocene settlement of temperate climates (**Fig. 7**, **Table 2**). The new

temperate niches could have constituted an opportunity for diversification because of

increased geographic ranges and ecological opportunities[49]. Several radiations have been

identified in other temperate groups following the apparition of the temperate biome such as

plants[51,104-106], mammals[107,108] or insects[32]. After this period, speciation dramatically

decreased for temperate lineages, maybe due to the effect of the Pleistocene glaciations, and

no difference in diversification between tropical- and temperate adapted lineages could be

currently observed (**Fig. 7**, Supplementary Figs. 1-3).

Conclusion

The foundations of a plausible explanation to understand large-scale diversity patterns in

terms of ecology and evolution are progressively assembling as we develop analytical

tools[95,109,110], accumulate data from several fields, and combine them into a single

framework[96]. We propose a framework explaining the origin of the today's LDG, where its

shape is mostly determined by changes in the global diversification and dispersal dynamics

imposed by large-scale climatic transitions. Our analyses suggest that the LDG is probably the

product of ancient tropical extinctions and Holarctic range contractions as the consequence of

the retraction of the tropical biome. We acknowledge the AGE hypothesis might not be

applicable to all organisms. As exemplified here, tropical extinction is not the most influential

cause behind the LDG of squamates. But the AGE hypothesis will be useful to explain the

LDG of groups that once were diverse at high latitudes. We acknowledge as well we consider

large geographical areas and long-time intervals that likely encompass high variability in

habitats and climates. The AGE hypothesis represents an effort to increase the complexity of

evolutionary models behind the LDG.

Methods

Time-calibrated phylogenies and fossil record. We confront the predictions of the AGE

hypothesis with the LDG of three vertebrate groups: turtles (order Testudines), crocodiles

(order Crocodilia), and the scaled reptiles (order Squamata), which are ancient clades that

experienced early Cenozoic climatic transitions[22,23,34,111]. They display contrasting patterns of

species richness: turtles and crocodiles are species-poor (330 and 25 species, respectively),

while squamates are species-rich (9400+ species). They all have a rich fossil record going

back to the Triassic for turtles and squamates (Early Cretaceous for crocodiles), which can

inform on how the latitudinal accumulation of species richness has varied through its

evolution[22,23,39]. Turtles are also remarkable because they show today a hump-shaped

diversity gradient centred around subtropical latitudes (10°S–30ºN)[111], while the

paleolatitudinal distribution of turtles was concentrated in the Holarctic, between 30–60ºN

during the Cretaceous[23,24].

Time-calibrated phylogeny for each group was obtained from published studies. For

turtles, we used the time-calibrated phylogeny of Jaffe *et al.*[112] including 233 species. We

preferred this phylogeny over other more recent and slightly better sampled trees[113] because

the divergence times estimates in Jaffe *et al.*[112] are more congruent with recent estimates

based on genomic datasets[72,114]. For squamates, we retrieved the most comprehensive dated

tree to date including 4161 species[34], and for crocodiles we obtained a complete phylogeny[115].

Fossil occurrences were downloaded from the *Paleobiology Database*

(https://paleobiodb.org/#/, last accessed October 25[th] 2017). To reduce potential biases in the

taxonomic assignation of turtle, crocodiles and squamates fossils, we compiled occurrence

data at the genus level. The fossil datasets were cleaned by checking for synonymies among

all taxa and for assignment to particular genus or family based on the literature

(Supplementary Table 9–11).

Estimation of origination and extinction rates with the phylogenies. To address whether

diversification differs between Holarctic and equatorial regions, we used the turtle and the

squamate phylogeny combined with distributional data (Supplementary Tables 1, 2) to fit

trait-dependent diversification models in BiSSE[65]. We also accounted for incomplete taxon

sampling, which is informed as a global sampling fraction of species at present having a given

trait[116].

To make our approach comparable with previous LDG studies[33-35,45,48], we first relied

on a constant-rate trait-dependent diversification model. The constant-rate BiSSE model has

six distinct parameters: two speciation rates (without range shift, or *in situ* speciation)

associated with Holarctic (hereafter "H", λ_H), and species occurring in other equatorial and

subtropical regions (hereafter "equator" or "E", λ_E), two extinction rates associated with

Holarctic (μ_H) and equator (μ_E), and two transition rates (dispersal or range shift) with one out

of Holarctic to equator (q_{H-E}), and from equator to Holarctic (q_{E-H}).

We also assessed the effect of species distribution on diversification, while allowing

for rates to change at specific points of time, which reduces the bias of using constant rates,

and allow testing the role of extinction and climate change on the generation of the LDG. We

used the time-dependent BiSSE (BiSSE.td) model, in which speciation, extinction, and

dispersal rates are allowed to vary between regions and to change after the shift times[98]. We

introduced two shift times to model different diversification dynamics between greenhouse,

transitional, and coldhouse periods; we assumed that a global warm tropical-like climate

dominated the world from the clades' origin to 51 Ma (the peak of temperature in the

Cenozoic), which then progressively cooled until 23 Ma (the transitional period), when

climate definitely shifted to a temperate-like biome in the Holarctic[59]. The shift times at 51

Ma and at 23 Ma are initial values that are re-estimated by the model during the likelihood

calculation. The climatic transition in the Cenozoic can have different temporal boundaries,

which might influence the results. We thus applied the same model but with different

combination of shift times (we tested 51/66 Ma and 34/23 Ma for the upper and lower bounds

of the climatic transition).

Analyses were performed using the R-package *diversitree* 0.9-7[66] using the functions

make.bisse to construct the likelihood functions for each model based on the data, and the

functions constrain and *find.mle* to apply different diversification scenarios. Finally, we used

a Markov Chain Monte Carlo (MCMC) approach to examine the credibility intervals of the

parameter estimates. Following the recommendations from FitzJohn[66], we used an

exponential prior $1/(2r)$ and started the chain with the parameters obtained by maximum

likelihood. We ran 10,000 steps of MCMC and applied a burn-in of 10%.

Inferring ancestral geographic distribution with and without fossils. We performed

biogeographic analyses with the parametric likelihood method DEC[69] using the fast C++

version[117]. Turtle, squamate, and crocodile species distributions were obtained from online

databases (www.iucnredlist.org and www.reptile-database.org). We chose 23.4°N and 23.4°S

as the threshold latitudes defining the tropics and categorized each species as living in the

Holarctic, in the southern temperate regions, or in the equatorial tropics and subtropical

regions[48]. We considered that all ranges comprised of three areas could be an ancestral state

(*maxareas* =3).

We set up three different DEC analyses: first, we run DEC without particular

constraints using only the distribution of extant species. Then, we performed DEC analyses

including fossil information in the form of 'fossil constraints' placed on certain nodes,

according to the range of distribution of fossil occurrences assigned to a particular taxon

during the relevant time frame. For example the crown age of Carettochelyidae (Testudines)

dates back to the Late Jurassic (150 Ma, *node 5*, **Fig. 3**; Supplementary Table 4), and we set a

constraint on this node reflecting the distribution of all the Late Jurassic fossils attributed to

this clade. Similarly, for the origin of turtles (210 Ma, *node 1*), distribution constraints

represent the range of Late Triassic fossils assigned to turtles. For the crown of Trionychidae,

which is Early Cretaceous age (123 Ma, *node 2*), the early fossils assigned to the clade were

used to constraint the geographic origin of Trionychidae (Supplementary Table 4). In total,

we implemented 23 fossil constraints for turtles (Supplementary Table 4), 30 fossil

constraints for squamates (Supplementary Table 5), and 8 for crocodiles (Supplementary

Table 6).

We included the fossil distribution following two different approaches: *(i)* a hard

(HFC), and *(ii)* a soft fossil constraints (SFC). With HFC, estimation of ancestral areas was

fixed to the fossil occurrence location. This was done using existing functions in the C++

version of Lagrange[117] (https://github.com/rhr/lagrange-cpp), with the command "*fixnode*".

By fixing nodes to the area of distribution of fossils, we assume fossil occurrences reflect the

distribution of the ancestors, *i.e.* the fossil record is complete. For SFC approach, fossil data

are incorporated into the anagenetic component of the likelihood framework. The direct

impact of a given fossil is limited to the particular branch to which it has been assigned,

although it may indirectly influence other branches. The inclusion of a fossil conditions the

estimated geographic-transition probability matrix for that branch by imposing a spatio-

temporal constraint on the simulation process. Only those simulations that realize a

geographic range that includes the area of fossil occurrence will contribute to the geographic-

range transition probability matrix of that branch; simulations that fail this constraint will be

discarded[118]. For SFC, we used the command "*fossil*" on DEC. We consider this as a "soft"

constraint because other areas different to that of the fossil could be included in the ancestral

states.

We then compared the timing and number of range extinction and dispersal events

inferred with the three different biogeographic analyses. In DEC, range-subdivision

(inheritance) scenarios (vicariance, duplication and peripatric isolation) occur at cladogenetic

events, while extinction (range contraction) and dispersal (range expansion) are modelled as

stochastic processes that occur along the branches of the tree[44,119]. Since the probability of any

extinction/dispersal event to occur is equal across all the branch length[69], to estimate the

periods at which range extinction and dispersal occurred, we divided the phylogeny in

intervals of 25 Ma and calculated the number of branches in which extinction/dispersal was

inferred that cross a particular time interval (the same branch could cross two continuous

intervals).

Estimation of origination and extinction rates with fossils. We analysed the three fossil

records using a Bayesian model to simultaneously infer the temporal dynamics of origination

and extinction, as well as the preservation rates[71]. This approach, implemented in PyRate[120],

uses fossil occurrences that can be assigned to a taxon, in this case fossil genera, and it jointly

models the preservation and diversification processes. The preservation process infers the

individual origination and extinction times of each taxon based on all fossil occurrences and

on an estimated preservation rate, and expressed as expected occurrences per taxon per Myr.

We followed the birth-death shift approach developed by Silvestro *et al.*[121], which is

focused on variation in origination and extinction at global scale and large temporal ranges.

We used a homogeneous Poisson process of preservation (-mHPP option). We also accounted

for varying preservation rates across taxa using the Gamma model with gamma-distributed

rate heterogeneity (-mG option). We used four rate categories to discretize the gamma

distribution, to accommodate the potential for more variability of preservation rates across

taxa.

Because of the number of occurrences analysed and of the vast timescale considered,

we dissected the birth–death process into time intervals, and estimated origination and

extinction rates within these intervals. In one set of analyses we defined the time intervals

using the geological epochs of the stratigraphic timescale[122] (Supplementary Figs. 12, 14, 16).

In another set of analyses, we defined the intervals with the different climatic periods that

characterized the Cenozoic (Supplementary Figs. 13, 15, 17), as discussed in the main text.

We adopted this solution as an alternative to the algorithms implemented in the original

PyRate that jointly estimate the number of rate shifts and the times at which origination and

extinction undergo a shift[71,120]. The estimation of origination and extinction rates within fixed

time intervals improved the mixing of the MCMC and allowed us to obtain an overview of the

general trends of rate variation throughout a long timescale[121]. Both the preservation and the

birth–death processes are still modelled in continuous time and are not based on boundary

crossings. Thus, the origination and extinction rates are measured as the expected number of

origination and extinction events per lineage per Myr. One potential problem in fixing *a*

priori the number of rate shifts is over-parameterization. We overcame this issue by assuming

that the rates of origination and extinction are part of two families of parameters following a

common prior distribution, with parameters estimated from the data using hyper-priors[123].

We ran PyRate for 10 million of MCMC generations on each of the 10 randomly

replicated datasets. We monitored chain mixing and effective sample sizes by examining the

log files in Tracer 1.6[124]. After excluding the first 20% of the samples as burn-in period, we

combined the posterior estimates of the origination and extinction rates across all replicates to

generate rates-through-time plots. Rates of two adjacent intervals were considered

significantly different when the mean of one lay outside of the 95% credibility interval of the

other, and conversely. We looked at the marginal posterior distributions of origination and

extinction rates through the evolutionary history of the three groups and looked at the effect

of the major environmental changes like the greenhouse world (Cretaceous), the climatic

transition (Paleogene), and the icehouse world (Neogene until the present).

In the context of the LDG and the hypothesis of tropical extinction, we performed

additional analyses with different subsets of fossils to tease apart the signal of speciation and

extinction between geographic regions (equator or Holarctic) and ecological conditions

(temperate or tropical). For example, for turtles, we split the global fossil dataset into four

subsets: one for the fossil genera occurring in the equator (429 occurrences), one for the

fossils occurring in the Holarctic (3558 occurrences), one for the fossil genera considered as

being adapted to temperate conditions (993 occurrences), and one for the fossils considered

being tropically adapted (2996 occurrences). We excluded the few fossil occurrences for the

southern regions of the South Hemisphere (about 180) for the different subset analyses, as

they are poorly represented in our dataset. Note that a given fossil can be present both in the

'Holarctic' and in the 'tropical' datasets. To encode tropical/temperate preferences, we

considered that conditions in the Holarctic were paratropical until the end of the Eocene

following[57-59] (and references therein), and assumed that taxa inhabiting the warm Holarctic

were adapted to tropical-like conditions (*i.e.* a high global temperature would indicate

probable adaptations to tropical climates). This is, of course, a simplifying assumption that

may cause bias in the analysis, but general patterns might still emerge from it[125]. Further, this

assumption could be justified by the presence of early Cenozoic vegetation assemblages in the

Holarctic that represent tropical vegetation, as evidenced by paleofloristic composition[53,56].

For turtles, this assumption is supported by a recent study that modelled their climatic niche

during the Late Cretaceous and found that Holarctic ancestors of turtles were adapted to

tropical conditions[81]. After the late Eocene, we categorized each species as living in the

temperate biome or the tropical biome following the threshold latitudes defining the tropics

(23.4°N and 23.4°S) following ref.[48]. This delineation is also overall congruent with the

Köppen climate classification. With these datasets, we reproduced the exact same PyRate

analyses than for the whole datasets (see above). In general, the fossil datasets include mostly

Holarctic fossils, with a more limited number of occurrences for the equator. Therefore we

remain cautious with the conclusions derived from the equatorial datasets.

References

1. Hillebrand, H. On the generality of the latitudinal diversity gradient. *Am. Nat.* **163**, 192-211

(2004).

2. Fuhrman, J. A. *et al.* A latitudinal diversity gradient in planktonic marine bacteria. *Proc.*

Natl Acad. Sci. USA **105**, 7774-7778 (2008).

3. Novotny, V. *et al.* Why are there so many species of herbivorous insects in tropical

rainforests? *Science* **313**, 1115-1118 (2006).

4. Jenkins, C. N., Pimm, S. L. & Joppa, L. N. Global patterns of terrestrial vertebrate diversity

and conservation. *Proc. Natl Acad. Sci. USA* **110**, E2602-E2610 (2013).

5. Kreft, H. & Jetz, W. Global patterns and determinants of vascular plant diversity. *Proc.*

Natl Acad. Sci. USA **104**, 5925-5930 (2007).

6. Willig, M. R. *et al.* Latitudinal gradients of biodiversity: pattern, process, scale, and

synthesis. *Annu. Rev. Ecol. Evol. Syst.* **34**, 273-309 (2003).

7. Mittelbach, G. G. *et al.* Evolution and the latitudinal diversity gradient: speciation,

extinction and biogeography. *Ecol. Lett.* **10**, 315-331 (2007).

8. Crame, J. A. Taxonomic diversity gradients through geological time. *Divers. Distrib.* **7**,

175–189 (2001).

9. Alroy, J. *et al.* Phanerozoic trends in the global diversity of marine invertebrates. *Science*

321, 97-100 (2008).

10. Mannion, P. D. *et al.* A temperate palaeodiversity peak in Mesozoic dinosaurs and

evidence for Late Cretaceous geographical partitioning. *Glob. Ecol. Biogeogr.* **21**, 898-

908 (2012).

11. Marcot, J. D., Fox, D. L. & Niebuhr, S. R. Late Cenozoic onset of the latitudinal diversity

gradient of North American mammals. *Proc. Natl Acad. Sci. USA* **113**, 7189-7194

(2016).

12. Mannion, P. D. *et al.* The latitudinal biodiversity gradient through deep time. *Trends Ecol.*

Evol. **29**, 42-50 (2014).

13. Rose, P. J., Fox, D. L., Marcot, J. & Badgley, C. Flat latitudinal gradient in Paleocene

mammal richness suggests decoupling of climate and biodiversity. *Geology* **39**, 163-166

(2011).

14. Brocklehurst, N., Day, M. O., Rubidge, B. S. & Fröbisch, J. Olson's Extinction and the

latitudinal biodiversity gradient of tetrapods in the Permian. *Proc. R. Soc. Biol. Sci.* **284**,

20170231 (2017).

15. Archibald, S. B. *et al.* Seasonality, the latitudinal gradient of diversity, and Eocene

insects. *Paleobiology* **36**, 374-398 (2010).

16. Archibald, S. B. *et al.* Seasonality, montane beta diversity, and Eocene insects: Testing

Janzen's dispersal hypothesis in an equable world. *Palaeogeogr. Palaeoclimatol.*

Palaeoecol. **371**, 1-8 (2013).

17. Labandeira, C. C. & Currano, E. D. The fossil record of plant-insect dynamics. *Annu. Rev.*

Earth Planet. Sci. **41**, 287-311 (2013).

18. Krug, A. Z. & Patzkowsky, M. E. Geographic variation in turnover and recovery from the

Late Ordovician mass extinction. *Paleobiology* **33**, 435–454 (2007).

19. Powell, M. G. Latitudinal diversity gradients for brachiopod genera during late Palaeozoic

time: links between climate, biogeography and evolutionary rates. *Glob. Ecol.*

Biogeogr. **16**, 519-528 (2007).

20. Crame, J. A. Evolution of taxonomic diversity gradients in the marine realm: evidence

from the composition of recent bivalve faunas. *Paleobiology* , 26, 188–214 (2000).

21. Leprieur, F. *et al.* Plate tectonics drive tropical reef biodiversity dynamics. *Nature*

Commun. **7**, 11461 (2016).

22. Mannion, P. D. *et al.* Climate constrains the evolutionary history and biodiversity of

crocodylians. *Nature Commun.* **6**, 8438 (2015).

23. Nicholson, D. B., Holroyd, P. A., Benson, R. B. & Barrett, P. M. Climate-mediated

diversification of turtles in the Cretaceous. *Nature Commun.* **6**, 7848 (2015).

24. Nicholson, D. B., Holroyd, P. A., Valdes, P. & Barrett, P. M. Latitudinal diversity

gradients in Mesozoic non-marine turtles. *R. Soc. Open Sci.* **3**, 160581 (2016).

25. Coiffard, C. & Gomez, B. Influence of latitude and climate on spread, radiation and rise to

dominance of early angiosperms during the Cretaceous in the Northern Hemisphere.

Geol. Acta **10**, 181–188 (2012).

26. Peralta-Medina, E. & Falcon-Lang, H. J. Cretaceous forest composition and productivity

inferred from a global fossil wood database. *Geology* **40**, 219–222 (2012).

27. Benson, R. B. J. & Upchurch, P. Diversity trends in the establishment of terrestrial

vertebrate eco-systems: interactions between spatial and temporal sampling biases.

Geology **41**, 43–46 (2013).

28. Jablonski, D. *et al.* Out of the tropics, but how? Fossils, bridge species, and thermal

ranges in the dynamics of the marine latitudinal diversity gradient. *Proc. Natl Acad. Sci.*

USA **110**, 10487-10494 (2013).

29. Montañez, I. P. *et al.* Climate, pCO_2 and terrestrial carbon cycle linkages during late

Palaeozoic glacial-interglacial cycles. *Nature Geosci.* **9**, 824-828 (2016).

30. Wiens, J. J. & Donoghue, M. J. Historical biogeography, ecology and species richness.

Trends Ecol. Evol. **19**, 639-644 (2004).

31. Ricklefs, R. E. Global variation in the diversification rate of passerine birds. *Ecology* **87**,

2468-2478 (2006).

32. Condamine, F. L., Sperling F. A. H., Whalberg, N., Rasplus, J.-Y. & Kergoat, G. J. What

causes latitudinal gradients in species diversity? Evolutionary processes and ecological

constraints on swallowtail biodiversity. *Ecol. Lett.* **15**, 267-277 (2012).

33. Pyron, R. A. & Wiens, J. J. Large-scale phylogenetic analyses reveal the causes of high

tropical amphibian diversity. *Proc. R. Soc. Biol. Sci.* **280**, 20131622 (2013).

34. Pyron, R. A. Temperate extinction in squamate reptiles and the roots of latitudinal

diversity gradients. *Glob. Ecol. Biogeogr.* **23**, 1126-1134 (2014).

35. Rolland, J., Condamine, F. L., Beeravolu, C. R., Jiguet, F. & Morlon, H. Dispersal is a

major driver of the latitudinal diversity gradient of Carnivora. *Glob. Ecol. Biogeogr.* **24**,

1059-1071 (2015).

36. Weir, J. T. & Schluter, D. The latitudinal gradient in recent speciation and extinction rates

of birds and mammals. *Science* **315**, 1574-1576 (2007).

37. Martin, P. S. & Klein, R. G. (Eds.) Quaternary extinctions: a prehistoric revolution.

University of Arizona Press (1989).

38. Wing, S. L. in *Biodiversity II: Understanding and Protecting our Biological Resources*

(ed. Reaka-Kudla, M. L. Wilson, D. E. & Wilson, E. O.) 163-185 (Joseph Henry Press,

Washington, D.C., 1997).

39. Markwick, P. J. Fossil crocodilians as indicators of Late Cretaceous and Cenozoic

climates: implications for using palaeontological data for global change. *Palaeogeogr.*

Palaeoclimatol. Palaeoecol. **137**, 205-271 (1998).

40. Roy, K. and Pandolfi, J.M. (2005). Responses of marine species and ecosystems to past

climate change. In: *Climate Change and Biodiversity* (1 edn.) (ed. Lovejoy, T.E and

Hannah, E.L.). Yale University Press, pp. 160-175.

41. Hawkins, B. A. *et al.* Post-Eocene climate change, niche conservatism, and the latitudinal

diversity gradient of New World birds. *J. Biogeogr.* **33**, 770-780 (2006).

42. Dunn, R.R., Agosti, D., Andersen, A.N., Arnan, X., Bruhl, C.A., Cerdá, X. et al. (2009).

Climatic drivers of hemispheric asymmetry in global patterns of ant species richness.

Ecol. Lett., **12**, 324-333.

43. Rabosky, D. L. Extinction rates should not be estimated from molecular phylogenies.

Evolution **64**, 1816-1824 (2010).

44. Sanmartín, I. & Meseguer, A. S. Extinction in phylogenetics and biogeography: From

lineages to the assembly of regional biotas. *Front. Genet.*, **7**, 35 (2016).

45. Pulido-Santacruz, P. and Weir, J.T. (2016). Extinction as a driver of avian latitudinal

diversity gradients. *Evolution*, **70**, 860-872.

46. Jablonski, D. *et al.* Out of the tropics: evolutionary dynamics of the latitudinal diversity

gradient. *Science* **314**, 102-106 (2006).

47. Donoghue, M. J. A phylogenetic perspective on the distribution of plant diversity. *Proc.*

Natl Acad. Sci. USA **105**, 11549-11555 (2008).

48. Rolland, J., Condamine, F. L., Jiguet, F. & Morlon, H. Faster speciation and reduced

extinction in the tropics contribute to the mammalian latitudinal diversity gradient.

PLoS Biol. **12**, e1001775 (2014).

49. Wiens, J. J. Global patterns of diversification and species richness in amphibians. *Am.*

Nat. **70**, S86-S106 (2007).

50. Cardillo, M., Orme, C. D. L. & Owens, I. P. F. Testing for latitudinal bias in

diversification rates: an example using New World birds. *Ecology* **86**, 2278–2287

(2005).

51. Leslie, A. B. *et al.* Hemisphere-scale differences in conifer evolutionary dynamics. *Proc.*

Natl Acad. Sci. USA **109**, 16217-16221 (2012).

52. Siqueira, A. C., Oliveira-Santos, L. G. R., Cowman, P. F. & Floeter, S. R. Evolutionary

processes underlying latitudinal differences in reef fish biodiversity. *Global Ecol.*

Biogeogr. **25**, 1466-1476 (2016).

53. Greenwood, D. R. & Wing, S. L. Eocene continental climates and latitudinal temperature

gradients. *Geology* **23**, 1044-1048 (1995).

54. Johnson, K. R. & Ellis, B. A tropical rainforest in Colorado 1.4 million years after the

Cretaceous-Tertiary boundary. *Science* **296**, 2379-2383 (2002).

55. Sluijs, A. *et al.* Subtropical Arctic Ocean temperatures during the Palaeocene/Eocene

thermal maximum. *Nature* **441**, 610-613 (2006).

56. Suan, G. *et al.* Subtropical climate conditions and mangrove growth in Arctic Siberia

during the early Eocene. *Geology* **45**, 539-542 (2017).

57. Ziegler, A. *et al.* Tracing the tropics across land and sea: Permian to present. *Lethaia* **36**,

227-254 (2003).

58. Morley, R. J. (2007). Cretaceous and Tertiary climate change and the past distribution of

megathermal rainforests. In: *Tropical Rainforest Responses to Climatic Changes* (ed.

Bush, M.B. and Flenley, J.). Praxis Publishing, Chichester, pp. 1–31.

59. Zachos, J. C., Dickens, G. R. & Zeebe, R. E. An early Cenozoic perspective on

greenhouse warming and carbon-cycle dynamics. *Nature* **451**, 279-283 (2008).

60. Couvreur, T. L., Forest, F. & Baker, W. J. Origin and global diversification patterns of

tropical rain forests: inferences from a complete genus-level phylogeny of palms. *BMC*

Biol. **9**, 44 (2011).

61. Christenhusz, M. J. & Chase, M. W. Biogeographical patterns of plants in the Neotropics–

dispersal rather than plate tectonics is most explanatory. *Bot. J. Linn. Soc.* **171**, 277-286

(2013).

62. Fine, P. V. A. & Ree, R. H. Evidence for a time-Integrated species-area effect on the

Latitudinal Gradient in tree diversity. *Am. Nat.* **186**, 796–804 (2006).

63. Rosenzweig, M. L. *Species Diversity in Space and Time* (Cambridge University Press,

1995).

64. Jansson, R., Rodríguez-Castañeda, G. & Harding, L. E. What can multiple phylogenies

say about the latitudinal diversity gradient? A new look at the tropical conservatism, out

of tropics, and diversification rate hypotheses. *Evolution* **67**, 1741-1755 (2013).

65. Maddison, W. P., Midford, P. E. & Otto, S. P. Estimating a binary character's effect on

speciation and extinction. *Syst. Biol.* **56**, 701-710 (2007).

66. FitzJohn, R. G. Diversitree: comparative phylogenetic analyses of diversification in R.

Methods Ecol. Evol. **3**, 1084-1092 (2012).

67. Goldberg, E. E., Lancaster, L. T. & Ree, R. H. Phylogenetic inference of reciprocal effects

between geographic range evolution and diversification. *Syst. Biol.* **60**, 451-465 (2011).

68. Davis, M. P., Midford, P. E. & Maddison, W. Exploring power and parameter estimation

of the BiSSE method for analyzing species diversification. *BMC Evol. Biol.* **13**, 38

(2013).

69. Ree, R. H. & Smith, S. A. Maximum likelihood inference of geographic range evolution

by dispersal, local extinction, and cladogenesis. *Syst. Biol.* **57**, 4-14 (2008).

70. Joyce, W. G., Rabi, M., Clark, J. M. & Xu, X. A toothed turtle from the Late Jurassic of

China and the global biogeographic history of turtles. *BMC Evol. Biol.* **16**, 236 (2016).

71. Silvestro, D., Schnitzler, J., Liow, L. H., Antonelli, A. & Salamin, N. Bayesian estimation

of speciation and extinction from incomplete fossil occurrence data. *Syst. Biol.* **63**, 349–

367 (2014).

72. Pereira, A. G., Sterli, J., Moreira, F. R. & Schrago, C. G. Multilocus phylogeny and

statistical biogeography clarify the evolutionary history of major lineages of turtles.

Mol. Phylogenet. Evol. **113**, 59-66 (2017).

73. Rodrigues, J. F. M., Olalla-Tárraga, M. Á., Iverson, J. B., Akre, T. S. & Diniz-Filho, J. A.

F. Time and environment explain the current richness distribution of non‐marine

turtles worldwide. *Ecography* **40**, 1402–1411 (2017).

74. Jansson, R. & Davies, T. J. Global variation in diversification rates of flowering plants:

energy vs. climate change. *Ecol. Lett.* **11**, 173-183 (2008).

75. Vellekoop, J. *et al.* Rapid short-term cooling following the Chicxulub impact at the

Cretaceous–Paleogene boundary. *Proc. Natl Acad. Sci. USA* **111**, 7537-7541 (2014).

76. Kaiho, K. *et al.* Global climate change driven by soot at the K-Pg boundary as the cause

of the mass extinction. *Sci. Rep.* **6**, 28427 (2016).

77. Lyson, T. R. Joyce, W. G. Knauss, G. E. & Pearson, D. A. *Boremys* (Testudines,

Baenidae) from the latest Cretaceous and early Paleocene of North Dakota: an 11-

million-year range extension and an additional K/T survivor. *J. Vertebrate Paleontol.*

31, 729-737 (2011).

78. Holroyd, P. A., Wilson, G. P. & Hutchison, J. H. Temporal changes within the latest

Cretaceous and early Paleogene turtle faunas of northeastern Montana. *Geol. Soc. Am.*

Spec. Pap. **503**, 299–312 (2014).

79. Longrich, N. R., Bhullar, B. A. S. & Gauthier, J. A. Mass extinction of lizards and snakes

at the Cretaceous–Paleogene boundary. *Proc. Natl Acad. Sci. USA* **109**, 21396-21401

(2012).

80. D'Hondt, S., Donaghay, P., Zachos, J. C., Luttenberg, D. & Lindinger, M. Organic carbon

fluxes and ecological recovery from the Cretaceous-Tertiary mass extinction. *Science*

282, 276-279 (1998).

81. Waterson, A. M. *et al.* Modelling the climatic niche of turtles: a deep-time perspective.

Proc. R. Soc. B Biol. Sci. **283**, 20161408 (2016).

82. Albino, A. M. & Brizuela, S. An overview of the South American fossil squamates. *Anat.*

Rec. **297**, 349-368 (2014).

83. Antonelli, A. & Sanmartín, I. Mass extinction, gradual cooling, or rapid radiation?

Reconstructing the spatiotemporal evolution of the ancient angiosperm genus

Hedyosmum (Chloranthaceae) using empirical and simulated approaches. *Syst. Biol.* **60**,

596-615 (2011).

84. Sun, Y. *et al.* Lethally hot temperatures during the Early Triassic greenhouse. *Science*

338, 366-370 (2012).

85. Crisp, M. D. & Cook, L. G. How was the Australian flora assembled over the last 65

million years? A molecular phylogenetic perspective. *Annu. Rev. Ecol. Evol. Syst.* **44**,

303-324 (2013).

86. Mayr, G. The early Eocene birds of the Messel fossil site: a 48 million-year-old bird

community adds a temporal perspective to the evolution of tropical avifaunas. *Biol. Rev.*

92, 1174-1188 (2017).

87. Wilf, P., Labandeira, C. C., Johnson, K. R. & Cúneo, N. R. Richness of plant–insect

associations in Eocene Patagonia: a legacy for South American biodiversity. *Proc. Natl*

Acad. Sci. USA **102**, 8944-8948 (2005).

88. Blois, J. L. & Hadly, E. A. Mammalian response to Cenozoic climatic change. *Annu. Rev.*

Earth Planet. Sci. **37**, 181–208 (2009).

89. Smith, R. Y., Basinger, J. F. & Greenwood, D. R. Early Eocene plant diversity and

dynamics in the Falkland flora, Okanagan Highlands, British Columbia, Canada.

Palaeobio. Palaeoenv. **92**, 309-328 (2012).

90. Xing, Y., Onstein, R. E., Carter, R. J., Stadler, T. & Linder, H. P. Fossils and a large

molecular phylogeny show that the evolution of species richness, generic diversity, and

turnover rates are disconnected. *Evolution* **68**, 2821-2832 (2014).

91. Morlon, H., Parsons, T. L. & Plotkin, J. B. Reconciling molecular phylogenies with the

fossil record. *Proc. Natl Acad. Sci. USA* **108**, 16327-16332 (2011).

92. May, M. R., Höhna, S. & Moore, B. R. A Bayesian approach for detecting the impact of

mass-extinction events on molecular phylogenies when rates of lineage diversification

may vary. *Methods Ecol. Evol.* **7**, 947-959 (2016).

93. Betancur-R, R., Ortí, G. & Pyron, R. A. Fossil-based comparative analyses reveal ancient

marine ancestry erased by extinction in ray-finned fishes. *Ecol. Lett.* **18**, 441-450

(2015).

94. Eiserhardt, W. L., Borchsenius, F., Plum, C. M., Ordonez, A. & Svenning, J.-C. Climate-

driven extinctions shape the phylogenetic structure of temperate tree floras. *Ecol. Lett.*

18, 263–272 (2015).

95. Meseguer, A. S., Lobo, J. M., Ree, R., Beerling, D. J. & Sanmartín, I. Integrating fossils,

phylogenies, and niche models into biogeography to reveal ancient evolutionary history:

the case of *Hypericum* (Hypericaceae). *Syst. Biol.* **64**, 215-232 (2015).

96. Fritz, S. A. *et al.* Diversity in time and space: wanted dead and alive. *Trends Ecol. Evol.*

28, 509-516 (2013).

97. Ng, J. & Smith, S. D. How traits shape trees: new approaches for detecting character state-

dependent lineage diversification. *J. Evol. Biol.* **27**, 2035-2045 (2014).

98. Spriggs, E. L. *et al.* Temperate radiations and dying embers of a tropical past: the

diversification of *Viburnum*. *New Phytol.* **207**, 340-354 (2015).

99. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Towards a

metabolic theory of ecology. *Ecology* **85**, 1771-1789 (2004).

100. Puurtinen, M. *et al.* Temperature-dependent mutational robustness can explain faster

molecular evolution at warm temperatures, affecting speciation rate and global patterns

of species diversity. *Ecography* **39**, 1025-1033 (2016).

101. Allen, A. P., Gillooly, J. F., Savage, V. M. & Brown, J. H. Kinetic effects of temperature

on rates of genetic divergence and speciation. *Proc. Natl Acad. Sci. USA* **103**, 9130–

9135 (2006).

102. Bromham, L. & Cardillo, M. Testing the link between the latitudinal gradient in species

richness and rates of molecular evolution. *J. Evol. Biol.* **16**, 200-207 (2003).

103. Schluter, D. Speciation, ecological opportunity, and latitude. *Am. Nat.* **187**, 1-18 (2016).

104. Spriggs, E. L., Christin, P. A. & Edwards, E. J. C_4 photosynthesis promoted species

diversification during the Miocene grassland expansion. *PLoS One* **9**, e97722 (2014).

105. Spalink, D. *et al.* Biogeography of the cosmopolitan sedges (Cyperaceae) and the area-
richness correlation in plants. *J. Biogeogr.* **43**, 1893-1904 (2016).

106. Favre, A. *et al.* Out-of-Tibet: the spatio-temporal evolution of *Gentiana* (Gentianaceae).
J. Biogeogr. **43**, 1967-1978 (2016).

107. Ge, D. *et al.* Evolutionary history of lagomorphs in response to global environmental
change. *PLoS One* **8**, e59668 (2013).

108. Pisano, J. *et al.* Out of Himalaya: the impact of past Asian environmental changes on the
evolutionary and biogeographical history of Dipodoidea (Rodentia). *J. Biogeogr.* **42**,
856-870 (2015).

109. Slater, G. J., Harmon, L. J. & Alfaro, M. E. Integrating fossils with molecular
phylogenies improves inference of trait evolution. *Evolution* **66**, 3931-3944 (2012).

110. Condamine, F. L., Rolland, J. & Morlon, H. Macroevolutionary perspectives to
environmental change. *Ecol. Lett.* **16**, 72-85 (2013).

111. Angielczyk, K. D., Burroughs, R. W. & Feldman, C. R. Do turtles follow the rules?
Latitudinal gradients in species richness, body size, and geographic range area of the
world's turtles. *J. Experimental Zool. B* **324**, 270-294 (2015).

112. Jaffe, A. L., Slater, G. J. & Alfaro, M. E. The evolution of island gigantism and body
size variation in tortoises and turtles. *Biol. Lett.* **7**, 558–561 (2011).

113. Rodrigues, J. F. M. & Diniz-Filho, J. A. F. Ecological opportunities, habitat, and past
climatic fluctuations influenced the diversification of modern turtles. *Mol. Phylogenet.*
Evol. **101**, 352-358 (2016).

114. Shaffer, H. B., McCartney-Melstad, E., Near, T. J., Mount, G. G. & Spinks, P. Q.
Phylogenomic analyses of 539 highly informative loci dates a fully resolved time tree
for the major clades of living turtles (Testudines). *Mol. Phylogenet. Evol.* **115**, 7-15
(2017).

939 115. Oaks, J. R. A time-calibrated species tree of Crocodylia reveals a recent radiation of the

940   true crocodiles. *Evolution* **65**, 3285-3297 (2011).

941 116. FitzJohn, R. G., Maddison, W. P. & Otto, S. P. Estimating trait-dependent speciation and

942   extinction rates from incompletely resolved phylogenies. *Syst. Biol.* **58**, 595-611 (2009).

943 117. Smith, S. A. Taking into account phylogenetic and divergence-time uncertainty in a

944   parametric biogeographical analysis of the Northern Hemisphere plant clade

945   Caprifolieae. *J. Biogeogr.* **36**, 2324-2337 (2009).

946 118. Moore, B. R., Smith, S. A., Ree, R. H. & Donoghue, M. J. Incorporating fossil data in

947   biogeographic inference: A likelihood approach. *Proc. 52nd Annu. Syst. Symposium*

948   Available at: http://www.phylo.org/pdf_docs/2006_MoBot_biogeo.pdf (2006).

949 119. Ree, R. H. & Sanmartín, I. Prospects and challenges for parametric models in historical

950   biogeographical inference. *J. Biogeogr.* **36**, 1211-1220 (2009).

951 120. Silvestro, D., Salamin, N. & Schnitzler, J. PyRate: a new program to estimate speciation

952   and extinction rates from incomplete fossil data. *Methods Ecol. Evol.* **5**, 1126-1131

953   (2014).

954 121. Silvestro, D., Cascales-Miñana, B., Bacon, C. D. & Antonelli, A. Revisiting the origin

955   and diversification of vascular plants through a comprehensive Bayesian analysis of the

956   fossil record. *New Phytol.* **207**, 425-436 (2015).

957 122. Gradstein, F. M., Ogg, J. G., Schmitz, M. & Ogg, G. (Eds.). *The Geologic Time Scale*

958   *2012*. Elsevier (2012).

959 123. Gelman, A. Parameterization and Bayesian modeling. *J. Am. Stat. Assoc.* **99**, 537-545

960   (2004).

961 124. Rambaut, A., Suchard, M. A., Xie, D. & Drummond, A. J. Tracer v1.6. Available at:

962   http://tree.bio.ed.ac.uk/software/tracer/ (2017).

963   125. Romdal, T. S., Araújo, M. B. & Rahbek, C. Life on a tropical planet: niche conservatism

964   and the global diversity gradient. *Global Ecol. Biogeogr.* **22**, 344-350 (2013).

Acknowledgements

We are grateful to Gary Mittlebach, Emmanuelle Jousselin and Jonathan Rolland for their comments on early versions of the manuscript. Financial support was provided by a Marie-Curie FP7–COFUND (AgreenSkills fellowship–26719) to A.S.M. and Marie Curie FP7-IOF (project 627684 BIOMME) to F.L.C. This work has benefited from an "Investissements d'Avenir" grant managed by Agence Nationale de la Recherche (CEBA, ref. ANR-10-LABX-25-01). This is contribution ISEM 2017-xxx of the Institut des Sciences de l'Evolution de Montpellier (UMR 5554 – CNRS). This is Paleobiology Database contribution no. xxx.

Author contributions

Both authors designed the study, analysed the data and wrote the manuscript.

Competing interests

The authors declare no competing financial interests.