1
2
3
The signaling lipid sphingosine 1-phosphate regulates mechanical pain

Authors and Affiliations: Rose Z. Hill[1], Benjamin U. Hoffman[2, 3], Takeshi Morita[1#], Stephanie M. Campos[4‡], Ellen A. Lumpkin[2, 4], Rachel B. Brem[5, 6], Diana M. Bautista[1, 4, 7, 8*]

1. Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.

2. Department of Physiology & Cellular Biophysics, Columbia University College of Physicians & Surgeons, New York, NY 10032

3. Medical Scientist Training Program, Columbia University, New York, NY 10032

4. Neurobiology Course, Marine Biological Laboratory, Woods Hole, MA

5. Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA.

6. Buck Institute for Research on Aging, Novato, CA 94945, USA.

7. Helen Wills Neuroscience Institute, University of California, Berkeley, CA 94720, USA.

8. Corresponding author

[#] Current Address: Rockefeller University, New York, NY 10065, USA.

[‡] Current Address: Indiana University, Bloomington, IN 47405, USA.

[*] Correspondence: dbautista@berkeley.edu

Abstract

Somatosensory neurons mediate responses to diverse mechanical stimuli, from innocuous touch to noxious pain. While recent studies have identified distinct populations of A mechanonociceptors (AMs) that are required for mechanical pain, the molecular underpinnings of mechanonociception remain unknown. Here, we show that the bioactive lipid sphingosine 1-phosphate (S1P) and S1P Receptor 3 (S1PR3) are critical regulators of acute mechanonociception. Genetic or pharmacological ablation of S1PR3, or blockade of S1P production, significantly impaired the behavioral response to noxious mechanical stimuli, with no effect on responses to innocuous touch or thermal stimuli. These effects are mediated by fast-conducting A mechanonociceptors, which displayed a significant decrease in mechanosensitivity in S1PR3 mutant mice. We show that S1PR3 signaling tunes mechanonociceptor excitability via modulation of KCNQ2/3 channels. Our findings define a new role for S1PR3 in regulating neuronal excitability and establish the importance of S1P/S1PR3 signaling in the setting of mechanical pain thresholds.

Introduction

Pain is a complex sensation. It serves to protect organisms from harmful stimuli, but can also become chronic and debilitating following tissue injury and disease. Distinct cells and molecules detect noxious thermal and mechanical stimuli. Thermal pain is detected by thermosensitive TRP channels in subsets of nociceptors[1,2], and gentle touch is detected by Piezo2 channels in low-threshold mechanoreceptors (LTMRs)[3,4]. Aδ high-threshold mechanoreceptors (HTMRs) have been shown to play a key role in responses to painful mechanical stimuli[5,6].

Recent studies have shown that there are at least two populations of HTMRs that mediate responses to noxious mechanical stimuli. The *Npy2r*[+] subpopulation of HTMRs mediates fast paw withdrawal responses to pinprick stimulation and terminates as free nerve endings in the epidermis[5]. The *Calca*[+] subpopulation of circumferential-HTMRs responds to noxious force and hair pulling, and terminates as circumferential endings wrapped around guard hair follicles[6]. Additionally, somatostatin-expressing interneurons of laminae I-III in the dorsal horn of the spinal cord receive input from nociceptors and are required for

behavioral responses to painful mechanical stimuli[7]. Despite these advances in defining the cells and

circuits of mechanical pain, little is known about the molecular signaling pathways in mechanonociceptors.

Here, we show that sphingosine 1-phosphate (S1P) is required for mechanical pain sensation. S1P is a

bioactive lipid that signals via 5 G-protein coupled S1P Receptors (S1PRs 1-5). S1P signaling, mainly via

S1PR1, plays a well-known role in immune cell migration and maturation[8–10]. Additionally, recent studies

have shown that S1PRs are expressed throughout the nervous system[11–13] and S1P signaling is associated

with a variety of neuroinflammatory disorders, including multiple sclerosis[14] and Alzheimer's disease[15]. S1P

has been implicated in spontaneous pain[13] and thermal pain hypersensitivity[12,16,17], but due to conflicting

accounts of S1P receptor expression in the CNS[11,18] and PNS[12,13,19] as well as inconsistent reports on the

effects of S1P on neuronal excitability[13,20,21] and pain behaviors[12,13,16,17], the role of S1P in somatosensation

remains controversial.

We found that mice lacking the S1P receptor S1PR3 display striking and selective deficits in behavioral

responses to noxious mechanical stimuli. Likewise, peripheral blockade of S1PR3 signaling or S1P

production impairs mechanical sensitivity. We found that S1P constitutively enhances the excitability of

A mechanonociceptors (AMs) via closure of KCNQ2/3 potassium channels to tune mechanical pain

sensitivity. The effects of S1P are completely dependent on S1PR3. While previous studies have shown

that elevated S1P triggers acute pain and injury-evoked thermal sensitization[12,13], we now show that

baseline levels of S1P are necessary and sufficient for setting normal mechanical pain thresholds. In

contrast, elevated S1P selectively triggers thermal sensitization via activation of TRPV1[+] heat nociceptors,

with no effect on mechanical hypersensitivity. Our findings uncover an essential role for constitutive S1P

signaling in mechanical pain.

Results

To identify candidate genes underlying mechanosensation, we previously performed transcriptome

analysis of the sensory ganglia innervating the ultra-sensitive tactile organ (the star) of the star-nosed

mole[22]. Immunostaining revealed the tactile organ is preferentially innervated by myelinated Aδ fibers[22],

which are primarily mechanosensitive. While our original analysis focused on ion channels enriched in the

neurons of the star organ, our dataset also revealed enrichment of several components of the S1P

pathway, including *S1pr3*. Likewise, single-cell RNA seq of mouse dorsal root ganglion (DRG) neurons

revealed *S1pr3* expression in a subset of myelinated mechanoreceptors[19] in addition to a subpopulation of

peptidergic C nociceptors.

S1P promotes excitability in small-diameter, capsaicin-sensitive nociceptors[12,13,20,21]. In addition, S1PR3

has been shown to mediate spontaneous pain triggered by elevated S1P and thermal sensitization

following sterile tissue injury[13]. However, no studies have examined the role of S1PR3 in

mechanosensation or in regulating somatosensory behaviors under normal conditions. Given the

enrichment of S1PR3 in specialized mechanosensory neurons of the star-nosed mole and mouse, we

hypothesized that S1P signaling via S1PR3 may also play a role in mechanosensation. Thus, we set out to

define the role of S1P signaling and S1PR3 in somatosensory mechanoreceptors.

S1PR3 mediates acute mechanical pain

We first examined a variety of somatosensory behaviors in mice lacking S1PR3[23] (*S1pr3$^{tm1Rlp/Mmnc}$;* referred

to herein as S1PR3 KO)*.* We initially investigated baseline responses to mechanical stimuli. S1PR3 KO

mice displayed a dramatic loss of mechanical sensitivity (**Figure 1A; see Figure 1-source data 1**), as von

Frey paw withdrawal thresholds were significantly elevated in S1PR3 KO mice relative to WT and S1PR3

HET littermates (mean thresholds: 1.737 g vs. 0.736 and 0.610 g, respectively). Moreover, S1PR3 KO

mice demonstrated decreased responses to a range of noxious tactile stimuli (2-6 g; **Figure 1B**) and to

noxious pinprick stimulation (**Figure 1C**), but normal responsiveness to innocuous tactile stimuli (0.6-1.4g;

Figure 1B). S1PR3 KO mice exhibited normal tape removal attempts[3] (**Figure 1D**), righting reflexes

(**Figure 1-figure supplement 1A**), radiant heat withdrawal latencies (**Figure 1E**), and itch-evoked

scratching (**Figure 1-figure supplement 1B**). These results demonstrate a selective role for S1PR3 in

acute mechanical pain.

As a complement to our analysis of somatosensation in S1PR3 KO animals, we employed a

pharmacological approach, using the S1PR3-selective antagonist TY 52156 (TY)[24]. Similar to the

phenotype of knockout animals, intradermal injection of 500 µM TY into the mouse hindpaw (the site of

testing) triggered a rapid and significant elevation in von Frey paw withdrawal thresholds (**Figure 1F**) and

decreased responsiveness to noxious (2-6 g), but not innocuous (0.6-1.4g), tactile stimuli (**Figure 1G**),

without affecting noxious heat sensitivity (**Figure 1-figure supplement 1C**). In contrast, blockade of

S1PR1 with the selective antagonist W146[16] had no effect on baseline mechanical or thermal thresholds

(**Figure 1F; Figure 1-figure supplement 1C**). Overall, these data show that S1PR3 signaling sets

mechanical pain sensitivity.

Endogenous S1P mediates acute mechanical pain

We next asked whether peripheral S1P was required for the S1PR3-dependent effects on

mechanosensation. We decreased S1P levels via injection of the sphingosine kinase inhibitor SKI II to

block local production of S1P[25] or elevated S1P levels via intradermal injection of S1P and measured

behaviors 30 minutes after injection. Decreasing local S1P levels with SKI II significantly reduced

mechanical sensitivity (**Figure 2A; see Figure 2-source data 1**), comparable to the hyposensitivity

phenotype observed in S1PR3 KO mice (**Figure 1A**). Again, similar to what was observed in S1PR3 KO

animals (**Figure 1E**), peripheral blockade of S1P production had no effect on baseline thermal sensitivity

(**Figure 1-figure supplement 1C**). Surprisingly, injecting exogenous S1P (10 µM; maximum solubility in

saline vehicle) had no effect on mechanical sensitivity (**Figure 2A-B**). However, as previously reported[12,13],

S1P injection triggered S1PR3-dependent thermal hypersensitivity and spontaneous pain (**Figure 2C-D**),

demonstrating that the lack of effect on mechanical hypersensitivity is not due to problems with S1P

delivery or degradation.

These data support a model whereby S1P constitutively activates S1PR3 to set normal mechanical pain

thresholds. To further test this model, we asked if the mechanical hyposensitivity elicited after endogenous

S1P depletion (via SKI II) could be rescued by local injection of exogenous S1P. Indeed, we found that

injection of exogenous S1P reversed SKI II-induced mechanical hyposensitivity in a dose-dependent

manner, and observed a maximal effect with 200 nM S1P (**Figure 2E**). Although quantification of native

S1P levels in skin is inaccurate owing to avid lyase activity[26], our data establish that baseline S1P levels

are sufficient to maximally exert their effect on S1PR3-dependent mechanical pain, such that increased of

S1P does not evoke mechanical hypersensitivity, but diminished S1P leads to mechanical hyposensitivity.

These data show that constitutive activation of S1PR3 by S1P is required for normal mechanosensitivity.

S1PR3 is expressed in A mechanonociceptors and thermal nociceptors

Our behavioral data showing distinct roles for S1PR3 in mechanonociception and thermal hypersensitivity

suggest that S1PR3 is expressed in distinct subsets of somatosensory neurons. While a previous study

suggested that all somatosensory neurons express S1PR3[13], single cell RNA seq data suggests *S1pr3* is

not expressed by all DRG neurons[19], and no studies have performed quantitative analysis of S1PR3

staining or co-staining to define subpopulations of S1PR3[+] neurons. We thus set out to characterize the

somatosensory neuron subtypes expressing *S1pr3* using *in situ* hybridization (ISH) of wild-type

somatosensory ganglia and immunohistochemistry (IHC) in an *S1pr3*[mCherry/+] reporter mouse[27].

We first used *in situ* hybridization (ISH) with a specific *S1pr3* probe to examine expression patterns of

S1pr3 (**Figure 3A-B**; see **Supplementary File 1**). In our experiments, 43% of cells from wild-type DRG

expressed *S1pr3*.   Co-ISH revealed that one population of *S1pr3*[+] neurons represents Aδ

mechanonociceptors (AMs). These cells expressed *Scn1a* (39.9% of all *S1pr3*[+]), a gene that encodes the

Nav1.1 sodium channel, which mediates mechanical pain in Aδ fibers[28]. *S1pr3*[+] cells also co-expressed

Npy2r (20.4% of all *S1pr3*[+]), a marker of a subset of mechanonociceptive A fibers[5]. *S1pr3* was expressed

in 70.6% of *Scn1a*[+] cells and 72% of *Npy2r*[+] cells, comprising a majority of both of these populations.

Interestingly, a subset of cells co-expressed *S1pr3* and the mechanically sensitive channel *Piezo2*, which is

expressed by Aβ, Aδ, and C fibers[3]. The remaining *S1pr3*[+] cells were *Trpv1*[+] and/or *Trpa1*[+] C nociceptors

(67.1% of all *S1pr3*[+]), which are reported to overlap minimally with the *Scn1a*[+] and *Npy2r*[+] populations[5,28].

We next used an *S1pr3*[mCherry/+] reporter mouse, which produces a functional S1PR3-mCherry fusion

protein[27], as an independent strategy to explore S1PR3 expression and localization. This strategy was

used because we found that anti-S1PR3 antibodies showed broad immunoreactivity in DRG from mice

lacking S1PR3, and so we instead used anti-DsRed antibodies to probe expression of the S1PR3 fusion

protein (**Figure 3-figure supplement 1E**). We found that 42.4% S1PR3$^+$ cells co-stained with anti-Peripherin, demonstrating that S1PR3 is expressed in a subset of small-diameter neurons. We also observed that 69.5% of S1PR3$^+$ cells co-stained with anti-NF200, which marks medium and large-diameter myelinated neurons. Furthermore, we observed that S1PR3$^+$ cells were primarily of small to medium diameter (11.3-35.1 µm), whereas all cells in the DRG ranged from 11.3-53.9 µm. Overall, these data support the expression of S1PR3 in subsets of small-diameter thermal nociceptors and medium-diameter mechanonociceptors (**Figure 3F**). Additionally, no significant differences were observed between WT and S1PR3 KO DRG in number of *Trpa1, Trpv1,* Peripherin-, NF200-, or IB4-positive cells (**Figure 3-figure supplement 1B-C, F, G**). The mean diameters of *Trpv1*$^+$ neurons (**Figure 3-figure supplement 1D**, left), NF200$^+$ neurons (**Figure 3-figure supplement 1G**), or all neurons (**Figure 3-figure supplement 1D**, right) in WT versus KO DRG were not significantly different, suggesting no loss of major sensory neuronal subtypes in the S1PR3 KO.

We then visualized S1PR3 expression in nerve fibers that innervate the skin using anti-DsRed antibodies in whole-mount immunohistochemistry (IHC; **Figure 3D**). The reporter animals showed no specific antibody staining in epidermal or dermal cells (**Figure 3-figure supplement 1I**), and single-cell RNA seq of a diverse array of mouse epidermal and dermal cells corroborates this lack of expression[29]. We observed overlap of S1PR3-expressing free nerve endings with NF200-positive myelinated free nerves and NF200-negative putative C-fiber endings (**Figure 3F**), but did not observe expression of S1PR3 in NF200-positive circumferential or lanceolate hair follicle receptors, or in putative Merkel afferents (**Figure 3D-E**). β-tubulin III, PGP9.5 (pan-neuronal markers), and NF200 staining in S1PR3 KO skin displayed patterns of epidermal and dermal innervation similar to WT skin, suggesting the phenotypes observed in the S1PR3 KO mice are not due to developmental loss of sensory neuronal innervation ($p_{PGP9.5}$ = 0.443 (n = 93, 38 fibers); p_{NefH} = 0.405 (n = 61, 28 fibers); p_{BTIII} = 0.353 (n = 104, 89 fibers); two-tailed t-tests based on average number of fibers per field of view). These results support expression of S1PR3 in subsets of myelinated A mechanonociceptors and unmyelinated C nociceptors that terminate as free nerve endings.

S1P activates thermal nociceptors but not putative AMs

Live imaging of cultured DRG neurons from adult reporter animals showed expression of S1PR3-mCherry fusion protein in 48.3% of neurons, mirroring our ISH and IHC results (**Figure 4A**). To examine the effects of S1P on A mechanonociceptors and C nociceptors, we performed ratiometric calcium imaging and electrophysiology on DRG cultures from reporter mice. Interestingly, only 56.1 ± 22.4 % of mCherry-expressing neurons were activated by 1 µM S1P (Representative trace in **Figure 4B**; representative images in **Figure 4C**), which our dose-response showed to be the saturating concentration for calcium influx (**Figure 4D**; EC_{50} = 155 nM). All S1P-responsive neurons were also capsaicin-sensitive (n > 2000 neurons). And while sensory neurons from S1PR3 KO animals did not respond to S1P, as expected[13], they exhibited capsaicin responses that were not significantly different from WT neurons (**Figure 3-figure supplement 1H**). The mean diameter of S1P-responsive mCherry[+] neurons was 22.4 ± 1.0 µm, whereas the mean diameter of non-responsive mCherry[+] neurons was 28.7 ± 3.2 µm (p = 0.0002, two-tailed t-test). We also performed whole cell current clamp experiments and, consistent with other studies[12,20,21], found that S1P evoked action potential firing in capsaicin-sensitive small diameter cells (**Figure 4E**). This shows that only the small-diameter, S1PR3[+] putative nociceptors are excited by S1P. We next asked whether the S1PR3[+] medium-large diameter neurons represent the mechanonociceptors observed by ISH (**Figure 3A**). To this end, we asked whether the spider toxin Hm1a, a selective activator of AM nociceptors[28], triggers calcium influx in S1PR3-expressing trigeminal neurons. Indeed, we found that 44.2 ± 15.1% of Hm1a-responsive neurons expressed mCherry (**Figure 4F**), consistent with our staining showing expression of *S1pr3* in AM nociceptors and the role of Hm1a-responsive neurons in mediating mechanical pain *in vivo*[28].

S1PR3 modulates KCNQ2/3 channels to regulate AM excitability

We next interrogated the molecular mechanism by which S1P signaling in AM nociceptors may regulate mechanical pain. We performed whole-cell current clamp on the medium-diameter *S1pr3*[mCherry/+] dissociated DRG neurons (membrane capacitance = 61.05 ± 1.92 pF), which did not display S1P-evoked calcium influx (**Figure 4B-C**). In these cells, 1 µM S1P application did not change membrane potential (**Figure 5-figure supplement 1A;** see **Figure 5-source data 1**) or elicit firing in the absence of current injection (**Figure 5-figure supplement 1A**; **Figure 5A**). However, S1P dramatically lowered the threshold to fire action

potentials (rheobase) in an S1PR3-dependent manner (**Figure 5A**, **Figure 5-figure supplement 1B**).

We then set out to determine the mechanism by which S1PR3 activity increases neuronal excitability using

whole-cell voltage clamp recording. Previous studies showed that S1P excites capsaicin-sensitive

nociceptors by increasing voltage-gated sodium currents and reducing steady-state potassium currents[20,21].

We found that S1P had no such effects on S1PR3[+] medium-diameter cells (**Figure 5-figure supplement**

1C-F). In contrast, S1P triggered a robust increase in input resistance (**Figure 5B**), consistent with the

closure of potassium channels. I-V analysis revealed that the current inhibited by S1P application was

carried by potassium (**Figure 5C**). Additionally, S1P significantly reduced slow, voltage-dependent tail

current amplitudes (**Figure 5D**; **Figure 5E** (top)) in an S1PR3-dependent manner (**Figure 5D**, center).

As tail currents in Aδ neurons are primarily mediated by KCNQ2/3 potassium channels[30,31], we postulated

that S1P may alter tail currents through modulation of these channels. Furthermore, the above properties

of the S1P-sensitive current were consistent with the reported electrophysiological properties of KCNQ2/3

channels in DRG neurons[30,32,33]. To address whether KCNQ2/3 channels mediated S1P-dependent

neuronal excitability, we applied the KCNQ2/3-selective inhibitor XE 991 and found that it completely

occluded the effects of S1P on tail current (**Figure 5E**). Similar results were observed with the related

antagonist, linopirdine (**Figure 5E**). These findings are consistent with S1P/S1PR3-dependent inhibition of

KCNQ2/3 in somatosensory neurons.

We also found that the effect of S1P on KCNQ2/3 currents was mediated by low levels of S1P, exhibiting

an IC_{50} of 48 nM with saturation at 100 nM (**Figure 5-figure supplement 1F**). While S1P cannot be

accurately measured in non-plasma tissues, this is similar to estimated levels of S1P in peripheral

tissues[10,34], and to levels which were sufficient to rescue mechanosensitivity after local S1P depletion

(**Figure 2E**). Thus, our *in vitro* IC_{50} supports our behavioral observations that baseline S1P levels are

sufficient to maximally exert their effect on mechanical pain. In summary, our electrophysiological and

behavioral observations support a model in which baseline S1P/S1PR3 signaling governs mechanical pain

thresholds through modulation of KCNQ2/3 channel activity in AM neurons (**Figure 8**).

274 **S1PR3 is required for nociceptive responses of high-threshold AM nociceptors**

275 Given the effects of S1P on putative AM neurons *in vitro* and the selective attenuation of baseline

276 mechanical pain in S1PR3 KO animals, we hypothesized that S1PR3 would play a role in AM afferent

277 function. To test this, we utilized *ex vivo* skin-nerve recordings to analyze the effects of genetic ablation of

278 S1PR3 on AM afferents, which mediate fast mechanical pain sensation. As no significant differences were

279 observed between S1PR3 WT and S1PR3 HET mice in any behavioral assays (**Figure 1**) or in AM force-

280 response relationships[28,35–38] (**Figure 6-figure supplement 1A;** see **Figure 6-source data 1**), we

281 compared AM nociceptor function between S1PR3 HET and S1PR3 KO littermates. S1PR3 KO AM

282 nociceptors displayed reduced sensitivity in their force-response relation (slope for HET versus KO: 50

283 Hz/N versus 35 Hz/N), as well as attenuated firing over the noxious, but not innocuous, range of

284 mechanical stimulation (**Figure 6A**). Strikingly, the median von Frey threshold to elicit firing in AM

285 nociceptors was significantly higher in S1PR3 KO animals (3.92 mN) compared to littermate controls (1.56

286 mN; **Figure 6B**).

287

288 Furthermore, S1PR3 KO AM nociceptors displayed a right-shifted cumulative response curve (50%

289 effective force for HET versus KO: 33.7 versus 60.0 mN; **Figure 6C**), consistent with the

290 mechanonociceptive hyposensitivity observed *in vivo*. A recent report suggested that A-nociceptors are

291 composed of two distinct neuronal populations that differ genetically, in mechanically evoked response

292 patterns, and in conduction velocity[5]. Accordingly, we found in the S1PR3 KO that a proportion of "Adapting

293 AM" nociceptors, characterized by adapting responses to static mechanical stimuli, were unresponsive to

294 controlled force stimulation (**Figure 6E-F**), and only responded to stimulation with a blunt glass probe or to

295 suprathreshold stimulation with von Frey filaments (see Methods). However, we found no differences in

296 conduction velocity between genotypes (**Figure 6D**) or the proportions of Aβ, Aδ, and C fibers between

297 wild-type and KO animals (**Figure 6-figure supplement 1B**). We conclude that S1PR3 is an essential

298 regulator of both mechanical threshold and sensitivity in a distinct population of AM nociceptors.

299

300

301

S1PR3 is required for inflammatory pain hypersensitivity

Having examined the mechanisms of S1P/S1PR3 signaling in acute mechanonociception, we next sought to evaluate S1P/S1PR3 signaling in pain hypersensitivity. For this purpose, we used an experimental model of inflammatory pain triggered by Complete Freund's Adjuvant (CFA) injection into the hindpaw, which elicits infiltration of immune cells and thermal and mechanical hypersensitivity[39]. While one previous study proposed that S1PR3 promotes injury-evoked heat and mechanical hypersensitivity, they did not measure or compare post-injury mechanical thresholds to pre-injury baselines for the knockout or control animals[13]. Here, we compared development of CFA-evoked hypersensitivity between S1PR3 HET and KO littermates, since no significant behavioral differences were observed between WT and HET animals in CFA experiments ($p_{\text{von Frey}}$ = 0.12; $p_{\text{radiant heat}}$ = 0.12; two-tailed t-tests). Strikingly, S1PR3 KO mice failed to develop thermal hypersensitivity (**Figure 7A; see Figure 7-source data 1**) relative to heterozygous littermates at both 24 and 48 hours post-CFA injection. In stark contrast, S1PR3 KO animals developed robust mechanical hypersensitivity when thresholds were normalized to account for the dramatic baseline differences between knockouts and control animals (**Figure 7B**). Our data demonstrate that S1PR3 mediates baseline mechanical sensitivity and is not required for the development of CFA-evoked mechanical hypersensitivity.

Influx of myeloid lineage (Cd11b$^+$/Ly6G$^-$) cells is required for the development of mechanical hypersensitivity in the CFA model[39]. In the immune system, S1P signaling via S1PR1 plays a key role in immune cell migration[9]. Consistent with the development of mechanical hypersensitivity, flow cytometry experiments showed robust infiltration of immune cells into hindpaw skin from both S1PR3 KO and littermate controls (**Figure 7-figure supplement 1A-B**). These data suggest that the phenotypes observed in S1PR3 KO mice cannot be attributed to compromised immune cell infiltration.

Next, we tested whether active S1P/S1PR3 signaling was required to maintain CFA-evoked thermal hypersensitivity using pharmacology. Acute blockade of S1P production with SKI II or S1PR3 with TY also reversed CFA heat hypersensitivity (**Figure 7C**), demonstrating that peripheral S1P actively signals via S1PR3 to promote CFA-evoked heat hypersensitivity. Furthermore, acute S1P/S1PR3 blockade with SKI II

or TY elevated mechanical thresholds to pre-CFA, baseline levels (**Figure 7D**) showing that S1PR3 tunes

mechanical pain under normal and inflammatory conditions. These results are consistent with the distinct

roles for AM and C nociceptors in mechanical pain. Under normal conditions, AM nociceptors set

mechanical pain thresholds[28,40]. In contrast, under inflammatory conditions, the combined activity of both

non-sensitized AM fibers and sensitized C fibers determines overall post-inflammatory mechanical

thresholds[41,40]. Consistent with this model, acute blockade of S1P production or S1PR3 activity under

normal conditions induces mechanical hyposensitivity and under inflammatory conditions returns

mechanical sensitivity to normal levels.

Discussion

We now show that S1P signaling via S1PR3 is a key pathway that tunes mechanical pain sensitivity.

Overall, our data reveal two new key findings. First, S1P/S1PR3 sets baseline mechanical pain thresholds.

Depletion of baseline, endogenous S1P induces mechanical hyposensitivity and nanomolar levels of

exogenous S1P are sufficient to restore normal mechanical pain sensitivity after depletion. Second,

elevated micromolar S1P levels, such as those produced during inflammation or disease, promote thermal,

but not mechanical hypersensitivity. The effects of S1P on acute mechanical pain and thermal

hypersensitivity are completely lost in S1PR3 knockout animals, which are otherwise normal with respect to

other somatosensory behaviors.

What is the local source of S1P in the skin that constitutively modulates mechanical pain? Even in the

mature field of S1P signaling in the vascular and immune systems, the cellular source of S1P, while an

intriguing question, remains unclear. All cells in the body, including somatosensory neurons, immune cells,

and skin cells, express sphingosine kinases 1 and 2 which are essential for S1P production[42]. Deletion of

both kinases is lethal and attempts to conditionally knockout these kinases fail to completely eliminate S1P

in tissues[43]. While RNA seq data suggests that somatosensory neurons contain all of the enzymatic

machinery required to produce and export local S1P[19,44], future work will be needed to identify the key cell

types that are important for maintaining baseline S1P levels in the skin to regulate mechanical sensitivity

and for increasing S1P under inflammatory/injury conditions to promote pain hypersensitivity.

Recent studies have identified distinct populations of AM nociceptors that are required for mechanical

pain[5,6]. Likewise, it was discovered that a subset of somatostatin-expressing spinal interneurons is required

for mechanical pain transduction[7]. Although these papers delineate the cells and circuitry of mechanical

pain, the molecular underpinnings of mechanonociception in the periphery are poorly understood. While

the identity of the transduction channel(s) in AM nociceptors remains enigmatic, understanding molecular

mechanisms that regulate excitability will no doubt provide key insights into the function and specialization

of the diverse subtypes of mechanosensitive nerve fibers. For example, although *Piezo2*-hypomorphic

animals exhibit normal mechanical pain behaviors[3], *ex vivo* skin-nerve recordings show that their AM

nociceptors display decreased force-responsiveness[3], and a recent study found that subpopulations of

sensory neurons express different splice variants of *Piezo2* that exhibit different force sensitivities[45]. These

studies suggest that mechanosensitive neurons exhibit functional specialization on multiple levels. Our

study demonstrates that S1PR3 is indispensable for normal function of AM nociceptors, including the

adapting AM population, recently discussed in Arcourt *et al.,* that innervates the epidermis and encodes

noxious touch[5]. We show that S1PR3 signaling modulates KCNQ2/3 channels to regulate excitability of

these A mechanonociceptors (**Figure 8**).

GPCR-mediated inhibition of KCNQ2/3 potassium channels is a well-known mechanism by which neuronal

excitability is regulated[46]. Other studies have shown that KCNQ channels mediate excitability of Aδ

fibers[30,31] and are required for normal mechanonociceptive responses in dorsal horn neurons receiving Aδ

input[31], and that opening KCNQ2/3 channels directly with retigabine alleviates pain *in vivo*[33,47,48]. Our

results not only complement previous work implicating KCNQ2/3 channels in pain, but also define the

upstream mechanisms that promote the regulation of KCNQ2/3 channels to tune mechanical pain

thresholds. Our data thus highlight S1PR3 as a novel and attractive target for the treatment of mechanical

pain and describe a new signaling pathway by which S1P regulates AM nociceptor excitability.

Interestingly, the neurons that innervate the ultra-sensitive tactile organ of the star-nosed mole are highly

enriched in transcripts for S1PR3 and KCNQ channels, as well as for a variety of other potassium

channels[22]. While it is difficult to directly examine the physiological basis for heightened mechanosensitivity in the star-nosed mole, S1PR3-dependent modulation of KCNQ may represent an important mechanism underlying the high tactile sensitivity of the star organ. Moreover, link between *S1pr3* and *Kcnq2/3* are echoed in single-cell RNA seq datasets from mouse DRG neurons, which show co-expression of *S1pr3* and *Kcnq2/3* in a subset of myelinated mechanoreceptors[19]. These cells are distinct from the *S1pr3/Trpv1* subset that mediates S1P-evoked acute pain and heat hypersensitivity. In addition to being transcriptionally distinct, we show that the mechanisms underlying S1P's activities in these cells are functionally distinct. Finally, *S1pr3* and *Kcnq2/3* are highly expressed in human sensory ganglia[49,50] and recordings from human stem cell-derived sensory neurons show that KCNQ2/3 channels play a key role in mediating their excitability[51]. Thus S1PR3 signaling may represent a new target for modulating mechanical pain.

Previous studies of S1P signaling in DRG neurons focused on S1P-evoked excitation of small diameter and/or capsaicin-sensitive neurons, and pain behaviors triggered by elevated S1P. While our new data affirms the effects S1P in thermal nociceptors observed by others, our manuscript highlights a novel effect of baseline levels of S1P in modulation of rheobase and KCNQ2/3 currents in mechanonociceptors. We are also the first to examine the role of S1PR3 in a variety of somatosensory behaviors under normal conditions (non-injury, no algogen injection), and to demonstrate a key role for S1PR3 in mechanonociception. We also go beyond previous studies[12,13,16,17] in showing that elevated S1P selectively promotes thermal and not mechanical hypersensitivity. Previous studies have shown that there are distinct cells and molecular pathways that trigger thermal versus mechanical hypersensitivity. For example, thermal hypersensitivity in CFA is dependent on TRPV1 ion channels[1] and independent of immune cell infiltration[39]. In contrast, infiltration of a subset of myeloid immune cells is required for mechanical hypersensitivity in an inflammatory pain model[39]. Here we show that S1P via S1PR3 signaling is a key component of the inflammatory soup that triggers thermal hypersensitivity in the CFA model. Our observation that S1PR3 KO animals display normal immune cell infiltration and develop mechanical hypersensitivity after CFA is consistent with these previous studies showing distinct mechanisms of inflammatory thermal and mechanical hypersensitivity.

Outside of the nervous system, S1P signaling via S1PR1 allows for the continuous circulation of lymphocytes between blood, lymph, and peripheral tissues[9]. Our findings that S1P plays a key role in noxious mechanosensation are in line with recent studies showing that sensory neurons co-opt classical immune pathways to drive chronic itch or pain[52,53]. What distinguishes this study from the others is that S1P signaling is critical for acute mechanical pain, even in the absence of inflammation or exogenously elevated S1P. In the immune system, disruptions in S1P levels or S1PR1 signaling result in significant immune dysfunction and disease[54-56]. Accordingly, in the somatosensory system, excessive, high levels of S1P (micromolar), such as those present in inflammation, evokes thermal pain and sensitization. Intermediate, baseline levels (nanomolar) regulate AM excitability and are required for normal mechanical pain sensation. In contrast, lowering S1P levels reduces mechanical pain sensation, while sparing innocuous touch sensation.

We propose that S1PR3 signaling may contribute to a variety of inflammatory diseases. S1P has been linked to a wide range of human inflammatory disorders[57-63]. Canonically, S1P signaling via S1PR1 is thought to promote inflammation via the immune system[58], however, we propose that S1P signaling via S1PR3 in neurons may also contribute to inflammatory disease. Indeed, fingolimod, a non-selective S1PR modulator, is prescribed as an immunosuppressant for multiple sclerosis treatment[14], but the possibility that some of its therapeutic effects may also be mediated via the nervous system has not been fully explored. Likewise, one study found that intrathecal fingolimod reduces bone cancer pain[64], and while analgesia was attributed to effects on S1PR1 in glia, some of the benefits may be due to S1PR3 signaling in DRG neurons. S1PR3 antagonism may also be useful in the treatment of inflammatory pain due to its selective dampening of acute mechanical pain and inflammatory thermal hypersensitivity, while preserving innocuous touch and normal thermal sensitivity. S1PR3 inhibitors may also be beneficial for treating other inflammatory disorders where S1PR3-expressing somatosensory neurons have been shown to contribute to neurogenic inflammation, such as asthma[65]. Our study demonstrates a crucial role for S1P signaling in the peripheral nervous system and highlights the potential of S1PR3 as a target for future pain therapies.

Materials and Methods

Behavioral studies & mice

S1pr3^{mcherry/+} and *S1pr3*^{-/-} mice were obtained from Jackson Laboratory and backcrossed to C57bl6/J. Wherever possible, wild-type/heterozygous (*S1pr3)* littermate controls were used in behavioral experiments. Mice (20–25 g) were housed in 12 h light-dark cycle at 21°C. Mice were singly housed one week prior to all behavioral experiments and were between 8-10 weeks at the time of the experiment. All mice were acclimated in behavioral chambers (IITC Life Sciences) on 2 subsequent days for 1 hour prior to all behavioral experiments.

Itch and acute pain behavioral measurements were performed as previously described[44,66,67]. Mice were shaved one week prior to itch behavior. Compounds injected: 500 μM TY 52156 (Tocris), 50 μM SKI II (Tocris), 0.2-10 μM S1P (Tocris, Avanti Polar Lipids), 50 mM chloroquine (Sigma), and 27 mM histamine (Tocris) in PBS with either 0.01-0.1% Methanol- (S1P) or 0.1-0.5% DMSO-PBS (all other compounds) vehicle controls. Pruritogens were injected using the cheek model (20 μL) of itch, as previously described[68]. Behavioral scoring was performed while blind to experimental condition and mouse genotype. All scratching and wiping behavior videos were recorded for 1 hour. Itch behavior was scored for the first 30 minutes and acute pain was scored for the first five minutes. Bout number and length were recorded.

For radiant heat and von Frey hypersensitivity behavior, drugs were injected intradermally into the plantar surface of the hindpaw (20 μL). Radiant heat assays were performed using the IITC Life Science Hargreaves test system. Mechanical threshold was measured using calibrated von Frey monofilaments (Touch Test) on a metal grate platform (IITC). Von Frey was performed as previously described[67,69] using the up-down method[70] while blinded to compound injected and genotype, or a descending force-series of 4 trials per force from 0.4 g to 6 g. Valid responses for both von Frey and radiant heat included fast paw withdrawal, licking/biting/shaking of the affected paw, or flinching. For radiant heat and von Frey, mice were allowed to acclimate on platform for 1 hour before injection. Measurements were taken 15 minutes pre-injection and 20-30 minutes post-injection for all compounds used.

The pinprick assay[7] was conducted on a von Frey testing platform (IITC). The mouse hindpaw was poked

with a 31 g syringe needle without breaking the skin to induce fast acute mechanical pain. Each paw was

stimulated 10 times with the needle, with five minutes rest in between trials, and the % withdrawal (fast

withdrawal, licking/biting/shaking of paw, squeaking, and/or flinching) was calculated from the total number

of trials.

The tape assay was conducted according to previously described methods[3]. Number of attempts to remove

a 3 cm piece of lab tape was recorded for 10 minutes after manual tape application to the rostral back.

Scorer and experimenter were blinded to genotype.

For righting reflex measurements, age-matched $S1pr3^{-/-}$ and $^{+/+}$ P6-7 neonates were used. Briefly, pups

were overturned one at a time on the home cage lid while experimenter was blinded to genotype. The time

to righting was measured to the nearest $1/10^{th}$ of a second with a stopwatch.

For the CFA model of hypersensitivity, mice were lightly anesthetized with isoflurane (2%) and injected with

15 µL CFA (Sigma) into one hindpaw using a Hamilton syringe (30 g) at 5pm. Radiant heat latencies and

von Frey 50% withdrawal thresholds were recorded one day prior to CFA, the morning of CFA (prior to

injection), and one and two days post-CFA. Von Frey measurements were acquired before radiant heat

latencies, and mice were allowed a one-hour recovery period in home cage with access to food and water

in between testing. Both ipsilateral and contralateral paw were measured. Experimenter was blind to

genotype for injections and recording.

All behavior experiments were carried out using age-matched or littermate cohorts of male mice and

conducted between 8am and 1pm. Mice were tested in 4-part behavior chambers (IITC Life Sciences) with

opaque dividers (TAP Plastics) with the exception of righting reflex measurements. Scratching and wiping

behaviors were filmed from below using high-definition cameras. All experiments were performed under the

policies and recommendations of the International Association for the Study of Pain and approved by the

University of California, Berkeley Animal Care and Use Committee.

In situ hybridization (ISH)

Fresh DRG were dissected from 8-12 week old mice, flash frozen in OCT embedding medium, and

sectioned at 14 μm onto slides. ISH was performed using Affymetrix Quantigene ViewISH Tissue 2-plex kit

according to manufacturer's instructions with Type 1 and Type 6 probes. The following probes against

mouse mRNAs were created by Affymetrix and used for ISH: *S1pr3, Scn1a, Npy2r, Piezo2, Trpv1, Trpa1.*

Immunohistochemistry (IHC) of DRG

DRG were dissected from 8-12 week old adult mice and post-fixed in 4% PFA for one hour. DRG were

cryo-protected overnight at 4°C in 30% sucrose-PBS, embedded in OCT, and then sectioned at 12 μm onto

slides. Briefly, slides were washed 3x in PBST (0.3% Triton X-100), blocked in 2.5% horse serum + 2.5%

BSA PBST, and incubated overnight at 4°C in 1:1000 primary antibody in PBST + 0.5% horse serum +

0.5% BSA. Slides were washed 3X in PBS then incubated 1-2 hours at RT in 1:1000 secondary antibody.

Slides were washed 3X in PBS and mounted in Fluoromount-G + DAPI with No. 1.5 coverglass. Primary

antibodies used: Rabbit anti-DsRed (Clontech #632496), Rabbit anti-S1PR3 (Abcam #38324; #108370),

Mouse anti-NF200 (Sigma #N5389), Chicken anti-Peripherin (Abcam #39374). Secondary antibodies used:

Goat anti-Mouse Alexa 488 (Abcam #150117), Goat anti-Chicken Alexa 488 (ThermoFisher #A11039),

Goat anti-Rabbit Alexa 594 (Invitrogen #R37117). Slides were mounted in Fluoromount with No. 1.5

coverglass. Imaging of ISH and DRG IHC experiments, and all live-cell imaging, was performed on an

Olympus IX71 microscope with a Lambda LS-xl light source (Sutter Instruments). For ISH and DRG IHC

analysis, images were analyzed using ImageJ software. Briefly, DAPI-positive cells were circled and their

fluorescence intensity (AFU) for all channels was plotted against cell size using Microsoft Excel software.

Co-labeling analysis was performed using ImageJ. Intensity thresholds were set based on the negative

control (no probe) slide. Cells were defined as co-expressing if their maximum intensities exceeded the

threshold for both channels of interest.

IHC of sectioned skin

Skin was dissected from 8 week old adult mice and post-fixed in 4% PFA for 30 minutes at RT. DRG were

cryo-protected overnight at 4°C in 30% sucrose-PBS, embedded in OCT, and then sectioned at 18 µm onto

slides. Briefly, slides were blocked in 5% normal goat serum in PBST (0.1% Triton X-100) and incubated

overnight at 4°C in 1:1000 primary antibody in blocking buffer. Slides were washed 3X in PBS then

incubated 45 minutes at RT in 1:1000 secondary antibody. Slides were washed 5X in PBS and mounted in

Fluoromount-G + DAPI with No. 1.5 coverglass. Primary antibodies used: Rabbit anti-DsRed (Clontech

#632496), Chicken anti-NefH (Abcam #4680), Chicken anti-β-tubulin III (Abcam #107216), mouse anti-

PGP9.5 (Abcam #8189). Secondary antibodies used: Goat anti-Mouse Alexa 488 (Abcam #150117), Goat

anti-Rabbit Alexa 594 (Invitrogen #R37117), Goat anti-Chicken Alexa 488 (ThermoFisher #A11039). For

co-localization analysis, only fibers for which > 50% of the length of the visible fiber contained co-localized

(white) pixels were counted. Image analysis was performed using ImageJ.

Whole mount skin IHC

Staining was performed according to Marshall and Clary *et al*[71]. Briefly, 8 week old mice were euthanized

and the back skin was shaved, depilated, and tape-stripped. The removed skin was fixed overnight in 4%

PFA, then washed in PBS (3X for 10 minutes each). Dermal fat was scraped away with a scalpel and skin

was washed in PBST (0.3% Triton X-100; 3X for two hours each) then incubated in 1:500 primary antibody

(Rabbit anti DsRed: Clontech #632496; Chicken anti-Nefh: Abcam #4680) in blocking buffer (PBST with

5% goat serum and 20% DMSO) for 5.5 days at 4°C. Skin was washed as before and incubated in 1:500

secondary antibody (Goat anti-Rabbit Alexa 594; Invitrogen #R37117; Goat anti-Chicken Alexa 488;

ThermoFisher #A11039) in blocking buffer for 3 days at 4°C. Skin was washed in PBST, serially dried in

methanol: PBS solutions, incubated overnight in 100% methanol, and finally cleared with a 1:2 solution of

benzyl alcohol: benzyl benzoate (BABB; Sigma) before mounting between No. 1.5 coverglass. Sectioned

and whole mount skin samples were imaged on a Zeiss LSM 880 confocal microscope with OPO using a

20x water objective. Image analysis was performed using ImageJ.

Cell culture

Cell culture was carried out as previously described[72]. Briefly, neurons from dorsal root ganglia (2-8 week

old adults) or trigeminal ganglia (P0) were dissected and incubated for 10 min in 1.4 mg ml−1 Collagenase

P (Roche) in Hanks calcium-free balanced salt solution, followed by incubation in 0.25% standard trypsin

(vol/vol) STV versene-EDTA solution for 2 min with gentle agitation. Cells were then triturated, plated onto

Poly D-Lysine coated glass coverslips and used within 20 h. Media: MEM Eagle's with Earle's BSS

medium, supplemented with 10% horse serum (vol/vol), MEM vitamins, penicillin/streptomycin and L-

glutamine.

Calcium imaging

Ca^{2+} imaging experiments were carried out as previously described[72]. Cells were loaded for 60 min at room

temperature with 10 µM Fura-2AM supplemented with 0.01% Pluronic F-127 (wt/vol, Life Technologies) in

a physiological Ringer's solution containing (in mM) 140 NaCl, 5 KCl, 10 HEPES, 2 CaCl2, 2 MgCl2 and 10

D-(+)-glucose, pH 7.4. All chemicals were purchased from Sigma. Acquired images were displayed as the

ratio of 340 nm/ 380 nm. Cells were identified as neurons by eliciting depolarization with high potassium

Ringer's solution (75 mM) at the end of each experiment. Responding neurons were defined as those

having a > 15% increase from baseline ratio. Image analysis and statistics were performed using

automated routines in Igor Pro (WaveMetrics). Fura-2 ratios were normalized to the baseline ratio

F340/F380 = (Ratio)/(Ratio t = 0).

In vitro electrophysiology

Electrophysiological experiments were carried out as previously described[72]. Briefly, recordings were

collected at 5 kHz and filtered at 2 kHz (Axopatch 200B, pClamp software). Electrode resistance ranged

between 1.5–5 MΩ. Internal solution contained 140 mM KCl, 2 mM MgCl2, 1 mM EGTA, 5 mM HEPES, 1

mM Na2ATP, 100 µM GTP, and 100 µM cAMP (pH 7.4). Bath solution was physiological Ringer's solution.

The pipette potential was canceled before seal formation. Cell capacitance was canceled before whole cell

voltage-clamp recordings. For mechanonociceptors experiments, only cells which were visually identified

as mCherry expressing and which had a capacitance between 40-80 pF were used. Rheobase was

calculated as the smallest current step required to elicit an action potential using current steps of 50 pA. M

currents were measured and analyzed using standard protocols for DRG neurons reported in the

literature[30,32,73]. Experiments were carried out only on cells with a series resistance of less than 30

MΩ. Analysis of electrophysiology data was performed in pClamp and IgorPro.

Ex vivo skin-nerve electrophysiology

Touch-evoked responses in the skin were recorded after dissecting the hind limb skin and saphenous

nerve from 7-10 week old mice, according to published methods[74,75]. The skin was placed epidermis-side-

up in a custom chamber and perfused with carbogen-buffered synthetic interstitial fluid (SIF) kept at 32 °C

with a temperature controller (model TC-344B, Warner Instruments). The nerve was kept in mineral oil in a

recording chamber, teased apart, and placed onto a gold recording electrode connected with a reference

electrode to a differential amplifier (model 1800, A-M Systems). The extracellular signal was digitized using

a PowerLab 8/35 board (AD Instruments) and recorded using LabChart software (AD Instruments).

For these studies, we focused on A-mechanonociceptors (AMs). To identify responses from these afferents

in mutant and control genotypes, we used a mechanical search paradigm with a fine glass probe. Afferents

were classified as AMs according to the following criteria: (1) conduction velocity (approximately, 1 to(≤12

m/s^{-1}), (2) medium-sized receptive fields, (3) sustained response to mechanical indentation[74,76,77].

Touch-sensitive afferents that did not meet these criteria were not analyzed further. Responses were

classified as Adapting AMs if the ratio of mean firing rate in the dynamic phase of stimulation (first 0.2 s) to

the static phase of stimulation (last 4.8 s) was greater than 2, and Non-Adapting AMs if the ratio was less

than or equal to 2. Non-responders (**Figure 6F**) responded to suprathreshold mechanical stimulation with

von Frey monofilaments (tip diameter <0.5 mm), but not to maximal controlled mechanical stimulation (256

mN, tip diameter 2 mm). All recordings and analyses were performed blind to genotype.

Mechanical responses were elicited with von Frey monofilaments and a force controlled custom-built

mechanical stimulator. Mechanical thresholds were defined as the lowest von Frey monofilament to reliable

elicit at least on action potential. Force controlled mechanical stimuli were delivered using a computer

controlled, closed-loop, mechanical stimulator (Model 300C-I, Aurora Scientific, 2 mm tip diameter). Low-

pass filtered, 5-second long, length control steps (square wave) simultaneously delivered with permissive

force control steps (square wave) were generated using LabChart software (AD Instruments). An arbitrarily

selected force step-and-hold protocol (8, 32, 4, 64, 128, 16, 256 mN) was delivered to all fibers. The period

between successive displacements was 60 seconds.

Conduction velocity was measured by electrically stimulating identified receptive fields. Spike sorting by

principal component analysis (PCA) and density based clustering, and data analysis was performed off-line

with custom-made software in MATLAB. Statistics were performed in Prism.

Flow cytometry of CFA-treated hind paws

CFA injections were performed as described above. Briefly, hindpaw skin and underlying fascia of treated

and PBS-injected paws were removed from freshly euthanized mice. Skin was placed in RPMI media

(Gibco) on ice before mincing with dissection scissors. Digestions were performed for 90 minutes at 37°C

on a rotating platform in 1 mL RPMI supplemented with 1:1000 DNaseI enzyme (Roche) and 1 unit

LiberaseTM (Roche). Skin was then filtered through 70 μm nylon mesh (Falcon), washed in RPMI, and

resuspended in PBS for Aqua Live-Dead staining. Samples were then transferred to FACS buffer (PBS

with 0.5% FBS and 2 mM EDTA), blocked, then surface stained with the following antibodies: cKit-Biotin,

FceRI-PE, CD49b-PECy7, SiglecF-APC, SA-FITC, Ly6C-PerCP, CD11b-PB, Ly6G-780, CD45.2-AF700

(eBiosciences). Compensation tubes (single-stain) were prepared for each fluorophore using positive and

negative control beads. A spleen from a wild-type animal was also prepared by crushing between frosted

glass slides, straining through 70 μm nylon mesh, and lysing of erythrocytes in ACK (Gibco). A portion of

spleen sample was heat-killed for 10 minutes at 65°C and stained with Aqua viability stain and set aside.

The rest of the spleen was stained normally with the other skin samples. Samples were then run through a

flow cytometer (BD Fortessa). Data were analyzed using FlowJo (Prism) and Microsoft Excel. Neutrophils

were defined as live single cells with the following staining profile: $CD45.2^+/CD11b^+/Ly6G^+/Ly6C^+$.

Inflammatory monocytes were defined as the following: $CD45.2^+/CD11b^+/Ly6G^-/Ly6C^{high}$. Total number of

immune cells was reported since neither genotype differed significantly between total number of live cells

or total number of CD45.2$^+$ immune cells.

Statistical analyses

All statistical analyses, except for skin nerve data (see above), were performed using IgorPro software or

Microsoft Excel. Values are reported as the mean ± SEM where multiple independent experiments are

pooled and reported (for whole cell electrophysiology), and mean ± SD where one experiment was

performed with multiple wells (for calcium imaging) or mice (for behavior). For comparison between two

groups, Student's unpaired 2-tailed t-test was used. A paired t-test was employed only for measurements

within the same biological replicate and after a given treatment. For single-point comparison between >2

groups, a one-way ANOVA followed by appropriate post hoc test was used, depending on comparison. For

the time course comparison between 2 groups, 2-way ANOVA was used and single comparison p-values

were derived using Tukey's HSD or appropriate statistical test, depending on comparison. Number of mice

or samples required to attain significance was not calculated beforehand, and where multiple statistical

tests were performed, a Bonferroni correction was applied. In figure legends, significance was labeled as:

n.s., not significant, $p \geq 0.05$; *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Acknowledgements

We thank Z. Rifi (UC Berkeley) for assistance with scoring itch behavior, R. P. Dalton (UC Berkeley) for

assistance with confocal microscopy, and P. Lishko and M. Miller (UC Berkeley) for advice on lipid stability

and usage. We would also like to thank R. Clary (Columbia) for whole mount skin staining protocols, B.

Jenkins (Columbia) for advice on the immunohistochemistry experiments, and D. Julius (UC San

Francisco) for the gift of Hm1a spider toxin. Confocal imaging experiments were conducted at the CRL

Molecular Imaging Center, supported by the Helen Wills Neuroscience Institute (UC Berkeley). We would

like to thank Holly Aaron and Jen-Yi Lee for their microscopy training and assistance. We are grateful to all

members of the D.M.B Laboratory (UC Berkeley) for constructive feedback and criticism. The National

Institutes of Health grants NS077224 (to D.M.B and R.B.), AR059385 (to D.M.B.), AR051219 (to E.A.L),

NS105449 and GM007367 (to B.U.H), and NS063307 (to the Neurobiology Course at the Marine Biological

Laboratory); and a Howard Hughes Medical Institute Faculty Scholars grant (to D.M.B) supported this work.

Competing interests

666 The authors declare no competing interests at this time.

References

1.   Caterina, M. J. *et al.* Impaired Nociception and Pain Sensation in Mice Lacking the Capsaicin

Receptor. *Science.* **288,** 306–313 (2000).

2.   Vriens, J. *et al.* TRPM3 Is a Nociceptor Channel Involved in the Detection of Noxious Heat. *Neuron*

70, 482–494 (2011).

3.   Ranade, S. S. *et al.* Piezo2 is the major transducer of mechanical forces for touch sensation in mice.

Nature **516,** 121–125 (2014).

4.   Woo, S.H. *et al.* Piezo2 is required for Merkel-cell mechanotransduction. *Nature* **509,** 622–626

(2014).

5.   Arcourt, A. *et al.* Touch Receptor-Derived Sensory Information Alleviates Acute Pain Signaling and

Fine-Tunes Nociceptive Reflex Coordination. *Neuron* **93,** 179–193 (2017).

6.   Ghitani, N. *et al.* Specialized Mechanosensory Nociceptors Mediating Rapid Responses to Hair Pull.

Neuron **95,** 944–954.e4 (2017).

7.   Duan, B. *et al.* Identification of Spinal Circuits Transmitting and Gating Mechanical Pain. *Cell* **159,**

1417–1432 (2014).

8.   Spiegel, S. & Milstien, S. Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat Rev Mol Cell*

Biol **4,** 397–407 (2003).

9.   Matloubian, M. *et al.* Lymphocyte egress from thymus and peripheral lymphoid organs is dependent

on S1P receptor 1. *Nature* **427,** 355–360 (2004).

10. Schwab, S. R. *et al.* Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P

gradients. *Science* **309,** 1735–1739 (2005).

11. Janes, K. *et al.* The development and maintenance of paclitaxel-induced neuropathic pain require

activation of the sphingosine 1-phosphate receptor subtype 1. *J. Biol. Chem.* **289,** 21082–21097

(2014).

12. Mair, N. *et al.* Genetic evidence for involvement of neuronally expressed S1P1 receptor in nociceptor

sensitization and inflammatory pain. *PLoS One* **6,** (2011).

13. Camprubi-Robles, M. *et al.* Sphingosine-1-phosphate-induced nociceptor excitation and ongoing

pain behavior in mice and humans is largely mediated by S1P3 receptor. *J Neurosci* **33,** 2582–2592

(2013).

14. Brinkmann, V. *et al.* Fingolimod (FTY720): discovery and development of an oral drug to treat

multiple sclerosis. *Nat Rev Drug Discov* **9,** 883–897 (2010).

15. Couttas, T. a *et al.* Loss of the neuroprotective factor Sphingosine 1-phosphate early in Alzheimer's

disease pathogenesis. *Acta Neuropathol. Commun.* **2,** 9 (2014).

16. Finley, A. *et al.* Sphingosine 1-phosphate mediates hyperalgesia via a neutrophil-dependent

mechanism. *PLoS One* **8,** e55255 (2013).

17. Weth, D. *et al.* Activated platelets release sphingosine 1-phosphate and induce hypersensitivity to

noxious heat stimuli in vivo. *Front. Neurosci.* **9,** 1–8 (2015).

18. Weth-Malsch, D. *et al.* Ablation of Sphingosine 1-Phosphate Receptor Subtype 3 Impairs

Hippocampal Neuron Excitability In vitro and Spatial Working Memory In vivo. *Front. Cell. Neurosci.*

10, 1–12 (2016).

19. Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-cell RNA

sequencing. *Nat. Neurosci.* **18,** 145–153 (2015).

20. Zhang, Y. H., Fehrenbacher, J. C., Vasko, M. R. & Nicol, G. D. Sphingosine-1-phosphate via

activation of a G-protein-coupled receptor(s) enhances the excitability of rat sensory neurons. *J.*

Neurophysiol. **96,** 1042–1052 (2006).

21.   Li, C., Li, J., Kays, J., Guerrero, M. & Nicol, G. D. Sphingosine 1-phosphate enhances the excitability

of rat sensory neurons through activation of sphingosine 1-phosphate receptors 1 and/or 3. *J.*

Neuroinflammation **12,** 1–20 (2015).

22.   Gerhold, K. *et al.* The Star-Nosed Mole Reveals Clues to the Molecular Basis of Mammalian Touch.

PLoS One **8,** (2013).

23.   Kono, M. *et al.* The sphingosine-1-phosphate receptors S1P1, S1P2, and S1P3 function coordinately

during embryonic angiogenesis. *J. Biol. Chem.* **279,** 29367–29373 (2004).

24.   Nussbaum, C. *et al.* Sphingosine-1-phosphate receptor 3 promotes leukocyte rolling by mobilizing

endothelial P-selectin. *Nat. Commun.* **6,** 6416 (2015).

25.   Chiba, Y., Takeuchi, H., Sakai, H. & Misawa, M. SKI-II, an inhibitor of sphingosine kinase,

ameliorates antigen-induced bronchial smooth muscle hyperresponsiveness, but not airway

inflammation, in mice. *J. Pharmacol. Sci.* **114,** 304–310 (2010).

26.   Shaner, R. L. *et al.* Quantitative analysis of sphingolipids for lipidomics using triple quadrupole and

quadrupole linear ion trap mass spectrometers. *J. Lipid Res.* **50,** 1692–1707 (2009).

27.   Sanna, M. *et al.* Bitopic S1P3 Antagonist Rescue from Complete Heart Block: Pharmacological and

Genetic Evidence for Direct S1P3 Regulation of Mouse Cardiac Conduction. *Mol. Pharmacol.* **89,**

176–186 (2015).

28.   Osteen, J. D. *et al.* Selective spider toxins reveal a role for the Nav1.1 channel in mechanical pain.

Nature **534,** 494–9 (2016).

29.   Joost, S. *et al.* Single-Cell Transcriptomics Reveals that Differentiation and Spatial Signatures

Shape Epidermal and Hair Follicle Heterogeneity. *Cell Syst.* **3,** 221–237.e9 (2016).

30.   Schutze, S., Orozco, I. J. & Jentsch, T. J. KCNQ Potassium Channels Modulate Sensitivity of Skin

Down hair (D-hair) Mechanoreceptors. *J. Biol. Chem.* **291,** 5566–5575 (2016).

31.   Passmore, G. M. *et al.* Functional significance of M-type potassium channels in nociceptive

cutaneous sensory endings. *Front. Mol. Neurosci.* **5,** 1–12 (2012).

32.   Crozier, R. A., Ajit, S. K., Kaftan, E. J. & Pausch, M. H. MrgD Activation Inhibits KCNQ / M-Currents

and Contributes to Enhanced Neuronal Excitability. *J Neurosci* **27,** 4492–4496 (2007).

33.   Xu, W. *et al.* Activation of Voltage-Gated KCNQ/Kv7 Channels by Anticonvulsant Retigabine

741   Attenuates Mechanical Allodynia of Inflammatory Temporomandibular Joint in Rats. *Mol. Pain* **6,**

742   1744-8069-6–49 (2010).

743 34. Ramos-Perez, W. D., Fang, V., Escalante-Alcalde, D., Cammer, M. & Schwab, S. R. A map of the

744   distribution of sphingosine 1-phosphate in the spleen. *Nat. Immunol.* **16,** 1245–52 (2015).

745 35. McIlwrath, S. L., Lawson, J. J., Anderson, C. E., Albers, K. M. & Koerber, H. R. Overexpression of

746   neurotrophin-3 enhances the mechanical response properties of slowly adapting type 1 afferents

747   and myelinated nociceptors. *Eur. J. Neurosci.* **26,** 1801–1812 (2007).

748 36. Kwan, K. Y., Glazer, J. M., Corey, D. P., Rice, F. L. & Stucky, C. L. TRPA1 Modulates

749   Mechanotransduction in Cutaneous Sensory Neurons. *J. Neurosci.* **29,** 4808–4819 (2009).

750 37. Smith, A. K., O'Hara, C. L. & Stucky, C. L. Mechanical sensitization of cutaneous sensory fibers in

751   the spared nerve injury mouse model. *Mol. Pain* **9,** 1–6 (2013).

752 38. Garrison, S. R., Dietrich, A. & Stucky, C. L. TRPC1 contributes to light-touch sensation and

753   mechanical responses in low-threshold cutaneous sensory neurons. *J. Neurophysiol.* **107,** 913–922

754   (2012).

755 39. Ghasemlou, N., Chiu, I. M., Julien, J.-P. & Woolf, C. J. CD11b [+] Ly6G [−] myeloid cells mediate

756   mechanical inflammatory pain hypersensitivity. *Proc. Natl. Acad. Sci.* 201501372 (2015).

757   doi:10.1073/pnas.1501372112

758 40. Abrahamsen, B. *et al.* The Cell and Molecular Basis of Mechanical, Cold, and Inflamatory Pain.

759   *Science.* **321,** 702 (2008).

760 41. Lennertz, R. C., Kossyreva, E. A., Smith, A. K. & Stucky, C. L. TRPA1 mediates mechanical

761   sensitization in nociceptors during inflammation. *PLoS One* **7,** 1–11 (2012).

762 42. Chalfant, C. E. & Spiegel, S. Sphingosine 1-phosphate and ceramide 1-phosphate: expanding roles

763   in cell signaling. *J. Cell Sci.* **118,** 4605–4612 (2005).

764 43. Pappu, R. *et al.* Promotion of lymphocyte egress into blood and lymph by distinct sources of

765   sphingosine-1-phosphate. *Science* **316,** 295–298 (2007).

766 44. Morita, T. *et al.* HTR7 Mediates Serotonergic Acute and Chronic Itch. *Neuron* **87,** 124–138 (2015).

767 45. Szczot, M. *et al.* Cell-Type-Specific Splicing of Piezo2 Regulates Mechanotransduction. *Cell Rep.*

768   **21,** 2760–2771 (2017).

769   46.   Passmore, G. M. *et al.* KCNQ/M currents in sensory neurons: significance for pain therapy. *J.*

770   *Neurosci.* **23,** 7227–7236 (2003).

771   47.   Hayashi, H., Iwata, M., Tsuchimori, N. & Matsumoto, T. Activation of peripheral KCNQ channels

772   attenuates inflammatory pain. *Mol. Pain* **10,** 15 (2014).

773   48.   Blackburn-Munro, G. & Jensen, B. S. The anticonvulsant retigabine attenuates nociceptive

774   behaviours in rat models of persistent and neuropathic pain. *Eur. J. Pharmacol.* **460,** 109–116

775   (2003).

776   49.   Ray, P. *et al.* Comparative transcriptome profiling of the human and mouse dorsal root ganglia: An

777   RNA-seq-based resource for pain and sensory neuroscience research. *bioRxiv* 165431 (2017).

778   doi:10.1101/165431

779   50.   Flegel, C. *et al.* RNA-Seq analysis of human trigeminal and dorsal root ganglia with a focus on

780   chemoreceptors. *PLoS One* **10,** 1–30 (2015).

781   51.   Young, G. T. *et al.* Characterizing human stem cell-derived sensory neurons at the single-cell level

782   reveals their ion channel expression and utility in pain research. *Mol. Ther.* **22,** 1530–1543 (2014).

783   52.   Oetjen, L. K. *et al.* Sensory Neurons Co-opt Classical Immune Signaling Pathways to Mediate

784   Chronic Itch. *Cell* **129,** 742–751 (2017).

785   53.   Pinho-Ribeiro, F. A., Verri, W. A. & Chiu, I. M. Nociceptor Sensory Neuron–Immune Interactions in

786   Pain and Inflammation. *Trends Immunol.* **38,** 5–19 (2017).

787   54.   Donoviel, M. S. *et al.* Spinster 2, a sphingosine 1-phosphate transporter, plays a Critical Role in

788   inflammatory and Autoimmune Diseases. *FASEB J.* **29,** 5018–5028 (2015).

789   55.   Gräler, M. H. & Goetzl, E. J. The immunosuppressant FTY720 down-regulates sphingosine 1-

790   phosphate G-protein-coupled receptors. *FASEB J.* **18,** 551–553 (2004).

791   56.   Olivera, A., Allende, M. L. & Proia, R. L. Shaping the landscape: Metabolic regulation of S1P

792   gradients. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1831,** 193–202 (2013).

793   57.   Allende, M. L. *et al.* Sphingosine-1-phosphate lyase deficiency produces a pro-inflammatory

794   response while impairing neutrophil trafficking. *J. Biol. Chem.* **286,** 7348–7358 (2011).

795   58.   Kunkel, G. T., Maceyka, M., Milstien, S. & Spiegel, S. Targeting the sphingosine-1-phosphate axis in

796   cancer, inflammation and beyond. *Nat. Rev. Drug Discov.* **12,** 688–702 (2013).

59.   Roviezzo, F. *et al.* S1P-induced airway smooth muscle hyperresponsiveness and lung inflammation

in vivo: Molecular and cellular mechanisms. *Br. J. Pharmacol.* **172,** 1882–1893 (2015).

60.   Liang, J. *et al.* Sphingosine-1-Phosphate Links Persistent STAT3 Activation, Chronic Intestinal

Inflammation, and Development of Colitis-Associated Cancer. *Cancer Cell* **23,** 107–120 (2013).

61.   Rivera, J., Proia, R. L. & Olivera, A. The alliance of sphingosine-1-phosphate and its receptors in

immunity. *Nat. Rev. Immunol.* **8,** 753–763 (2008).

62.   Myśliwiec, H. *et al.* Increase in circulating sphingosine-1-phosphate and decrease in ceramide levels

in psoriatic patients. *Arch. Dermatol. Res.* (2016). doi:10.1007/s00403-016-1709-9

63.   Checa, A. *et al.* Circulating levels of sphingosine-1-phosphate are elevated in severe, but not mild

psoriasis and are unresponsive to anti-TNF-α treatment. *Sci. Rep.* **5,** 12017 (2015).

64.   Grenald, S. A. *et al.* Targeting the S1P/S1PR1 axis mitigates cancer-induced bone pain and

neuroinflammation. *Pain* **158,** 1 (2017).

65.   Trankner, D., Hahne, N., Sugino, K., Hoon, M. A. & Zuker, C. Population of sensory neurons

essential for asthmatic hyperreactivity of inflamed airways. *Proc. Natl. Acad. Sci.* **111,** 11515–11520

(2014).

66.   Wilson, S. R. *et al.* The Ion Channel TRPA1 Is Required for Chronic Itch. *J. Neurosci.* **33,** 9283–

9294 (2013).

67.   Tsunozaki, M. *et al.* A 'toothache tree' alkylamide inhibits Aδ mechanonociceptors to alleviate

mechanical pain. *J. Physiol.* **591,** 3325–40 (2013).

68.   Shimada, S. G. & LaMotte, R. H. Behavioral differentiation between itch and pain in mouse. *Pain*

139, 681–687 (2008).

69.   Chaplan, S.R., Bach, F.W., Pogrel, J.W., Chung, J.M., Yaksh, T. Quantitative assessment of tactile

allodynia evoked by unilateral ligation of the fifth and sixth lumbar nerves in the rat. *J. Neurosci.* **53,**

55–63 (1994).

70.   Dixon, W. J. The Up-and-Down Method for Small Samples. *J. Am. Stat. Assoc.* **60,** 967–978 (1980).

71.   Marshall, K. L. *et al.* Touch Receptors Undergo Rapid Remodeling in Healthy Skin. *Cell Rep.* **17,**

1719–1727 (2016).

72.   Wilson, S. R. *et al.* TRPA1 is required for histamine-independent, Mas-related G protein-coupled

receptor-mediated itch. *Nat. Neurosci.* **14,** 595–602 (2011).

73.   Zheng, Q. *et al.* Suppression of KCNQ/M (Kv7) potassium channels in dorsal root ganglion neurons

827   contributes to the development of bone cancer pain in a rat model. *Pain* **154,** 434–448 (2013).

828   74.   Wellnitz, S. A., Lesniak, D. R., Gerling, G. J. & Lumpkin, E. A. The Regularity of Sustained Firing

829   Reveals Two Populations of Slowly Adapting Touch Receptors in Mouse Hairy Skin. *J.*

830   *Neurophysiol.* **103,** 3378–3388 (2010).

831   75.   Maksimovic, S. *et al.* Epidermal Merkel cells are mechanosensory cells that tune mammalian touch

832   receptors. *Nature* **509,** 617–621 (2014).

833   76.   Koltzenburg, M. *et al.* Receptive Properties of Mouse Sensory Neurons Innervating Hairy Skin

834   Receptive Properties of Mouse Sensory Neurons Innervating Hairy Skin. *J. Neurophysiol.* **78,** 1841–

835   1850 (1997).

836   77.   Zimmermann, K. *et al.* Phenotyping sensory nerve endings in vitro in the mouse. *Nat. Protoc.* **4,**

837   174–196 (2009).

Figure captions

Figure 1. S1PR3 mediates acute mechanical pain. A. von Frey 50% withdrawal threshold

measurements for *S1pr3*$^{+/+}$ (WT, N = 8), $^{+/-}$ (HET, N = 7) and $^{-/-}$ (KO, N = 12) mice. $p < 0.0001$ (one-way

ANOVA). Tukey-Kramer post hoc comparisons for KO and HET to WT indicated on graph. **B.** von Frey

force-response graph for WT (N = 8) versus KO (N = 12) animals; $p_{genotype} < 0.0001$ (two-way ANOVA).

Tukey HSD comparisons between genotypes are indicated for given forces. **C.** % withdrawal to pinprick

stimulation of hindpaw for HET versus KO animals; $p < 0.0001$ (unpaired t-test; N = 5-7 mice per group). **D.**

Number of attempted removal bouts in tape assay for WT (N =2), HET (N = 2), and KO (N = 5) mice; $p =$

0.172 (one-way ANOVA). **E.** Baseline radiant heat measurements for WT, HET, and KO mice. N = 8 WT, 3

HET, and 5 KO mice. $p = 0.444$ (one-way ANOVA). **F.** von Frey 50% withdrawal threshold measurements

for mice pre- and post-injection of 500 μM TY 52156 (N = 10), 10 μM W146 (N = 6), or 1% DMSO-PBS

vehicle (N = 17); $p = 0.016, 0.650$ (two-tailed paired t-test comparing vehicle- vs. drug-injected paw). **G.**

von Frey force-response graph for mice injected with either 1% DMSO-PBS or 500 μM TY 52156; $p_{treatment} <$

1e-05 (two-way ANOVA; N = 4 mice per group). Tukey HSD comparisons were made between treatment

groups and significant differences at a given force are indicated on graph. Error bars represent mean ± SD.

Figure 1-source data 1. S1PR3 mediates acute mechanical pain. *Related to Figure 1.*

Figure 1-figure supplement 1. Loss of S1PR3 selectively impairs mechanonociception. *Related to*

Figure 1. **A.** Time to righting in seconds for N = 6 P7 pups per genotype for WT and KO mice; p = 0.575

(two-tailed unpaired t-test). **B.** (Left) Time spent scratching in response to injection of 50 mM chloroquine or

PBS vehicle (VEH) in WT and KO mice; p = 0.36, 0.98, (unpaired t-tests; N = 3-4 mice per group). (Right)

Time spent scratching in response to injection of 27 mM Histamine or 0.1% DMSO-PBS in WT and KO

mice; p = 0.51, 0.06 (unpaired t-tests; N = 3-4 mice per group). **C.** Normalized paw withdrawal latencies

post-injection of SKI II, TY 52156, or 0.1% DMSO-PBS vehicle into the hind paw of wild-type animals; p =

0.65 (one-way ANOVA); N = 5 mice per group. Unless otherwise indicated, error bars represent mean ±

SD.

Figure 2. Endogenous S1P mediates acute mechanical pain. A. von Frey 50% withdrawal

measurements for mice pre- and post-injection of 50 µM SKI II (N = 8) or 10 µM S1P (N = 7); p = 0.003,

0.604 (two-tailed paired t-tests). **B.** von Frey force-response graph for animals injected with 10 µM S1P or

0.1% MeOH-PBS; $p_{genotype}$ > 0.05 (two-way ANOVA; N = 8 mice per group). No Tukey HSD comparisons

at any force between genotypes were significant. **C.** Intradermal cheek injection of 10 µM S1P, 2 µM, 0.2

µM, and 20 µL 0.3% methanol PBS (vehicle), with quantification of number of forepaw wipes over the 5

minute post-injection interval; p < 0.0001 (one-way ANOVA; N = 3 mice per condition). Dunnett's multiple

comparisons p-values are represented on graph for comparisons made between treated and vehicle

groups. **D.** Radiant heat normalized paw withdrawal latencies 20-30 minutes post injection of 15 µL 10 µM

S1P, 0.2 µM S1P. or 0.3% methanol-PBS vehicle (i.d.) into the hind paw of S1PR3 WT or KO mice; p =

0.0129 (one-way ANOVA; N = 3-10 mice per condition). Dunnett's multiple comparisons p-values are

represented on graph for comparisons made between treated and vehicle groups. **E.** von Frey 50%

withdrawal measurements for mice pre- (baseline) and post-injection of 50 µM SKI II (N = 14) and 0, 10,

75, or 200 nM S1P (one-way ANOVA, p = 0.0001; N = 4, 3, 4, 3 animals). Tukey Kramer comparisons are

indicated on graph. Error bars represent mean ± SD.

Figure 2-source data 2. Endogenous S1P mediates acute mechanical pain. *Related to Figure 2.*

Figure 3. *S1pr3* is expressed in A mechanonociceptors and C thermal nociceptors. A. (Top)

Representative co-ISH of *S1pr3* (green; left) with *Scn1a*, *Npy2r*, *Piezo2*, and *Trpv1* (magenta; center) in

sectioned DRG. Right column: overlay with co-localized regions colored white (10x air objective; scale =

100 μm). **B.** Bar chart showing the % of total cells expressing the indicated marker (grey) and the % of total

cells co-expressing both marker and *S1pr3* (green). See Table S1 for quantification. **C.** Representative IHC

images of sectioned DRG from *S1pr3$^{mCherry/+}$* animals stained with anti-DsRed (green, S1PR3) and anti-

Peripherin (left, magenta) or anti-NF200 (right, magenta). Arrows indicate co-stained cells. Images were

acquired using a 10x air objective (scale = 100 μm). **D.** Whole-mount skin IHC confocal images with anti-

DsRed antibody (S1PR3, green) and anti-NefH antibody (NF200, magenta) in an *S1pr3$^{mCherry/+}$* animal (20x

water objective; scale = 50 μm). Arrows indicate co-positive free nerves (left image). Arrowheads indicate

NF200- free nerves (left) or S1PR3- circumferential fibers (right image). **E.** Sectioned skin IHC with anti-

DsRed (S1PR3) and anti-NefH (NF200, left, top right) or anti-DsRed (S1PR3) and anti-beta-tubulin III

(BTIII, bottom right) antibody (magenta) in *S1pr3$^{mCherry/+}$* skin (20x air objective; scale = 50 μm). Arrows

indicate co-positive free nerve endings (left), S1PR3-negative lanceolate/circumferential hair follicle

endings (top right, arrow = circumferential, arrowhead = lanceolate), or S1PR3-negative putative Merkel

afferent (bottom right). **F.** (Left) Quantification of sectioned DRG IHC experiments showing % of S1PR3+

cells that co-stained with indicated markers (n > 250 cells per marker). (Right) Quantification of sectioned

skin IHC experiments showing % of fibers positive for indicated marker that co-stained with S1PR3 (anti-

DsRed; n = 10 images per marker from 2 animals).

Figure 3-figure supplement 1. S1PR3 KO animals display normal representation of somatosensory

neuronal subtypes. *Related to Figure 3.* **A.** ISH of sectioned adult DRG from WT and S1PR3 KO animals

showing specificity of *S1pr3* probes (20x air objective, scale = 50 μm). **B.** (Image) Representative ISH

Trpv1 (top) and *Trpa1* (bottom) from sectioned DRG of wild-type (left) and S1PR3 KO animals (right; 20x

air objective; scale = 50 µm). **C.** % of total cells expressing *Trpv1* and *Trpa1* in sectioned DRG of wild-type

and S1PR3 KO animals. **D.** (Left) Average diameter ± SEM of all cells in sectioned DRG from WT and

S1PR3 KO animals (p = 0.36, t-test; n = 437 and 679 cells from two animals each, respectively). (Right)

Average diameter ± SD of *Trpv1+* cells in WT and S1PR3 KO DRG (p = 0.63, t-test; n = 127 and 85 cells

respectively). **E.** Representative staining with anti-S1PR3 antibody (1:2000) in sectioned adult DRG from

WT and S1PR3 KO animals (10x air objective, scale = 100 µm). **F.** Representative IHC images of

sectioned DRG from S1PR3 KO animals stained with anti-DsRed (green) and anti-Peripherin (left,

magenta) or anti-NF200 (right, magenta). Images were acquired using a 10x air objective (scale = 100 µm).

G. (Left) Quantification of total percentage of cells stained with indicated markers in sectioned DRG from

S1pr3$^{mCherry/+}$ and S1PR3 KO animals (n > 250 cells per condition). (Right) Average diameter of anti-

NF200+ cells in *S1pr3$^{mCherry/+}$* and S1PR3 KO DRG (p = 0.15, t-test; n = 256 and 194 cells, respectively). **H.**

Percent responders to S1P and capsaicin in ratiometric calcium imaging of wild-type and *S1pr3 $^{-/-}$* cultured

DRG and TG neurons; $p < 0.0001$ (one-way ANOVA; N = 2 DRG and 2 TG preparations of 8 wells each).

Sidak's multiple comparisons p-values are represented on graph for comparisons made between

genotypes. Error bars represent mean ± SEM. **I.** (Left) No primary control showing robust staining of hair

follicles in whole mount skin in contrast to specific neuronal staining shown in **Figure 3D**. (Right) No

primary control showing staining around hair follicles and in epidermis in sectioned skin. Scale = 50 µm

(20x water objective).

Supplementary file 1. Co-ISH quantification for sectioned DRG from adult wild-type mice. *Related to*

Figure 3.

Figure 4. S1P activates thermal nociceptors but not mechanonociceptors. A. (Left) Representative

image of mCherry signal in live, cultured adult DRG neurons from one *S1pr3$^{mCherry/+}$* animal. (Right)

Quantification of % of total cells expressing *S1pr3* from DRG ISH and mCherry from dissociated DRG

cultures (N = 2 animals each experiment). **B.** Representative traces depicting F340/F380 signal from

Fura2-AM calcium imaging showing two neurons, one which responded to 1 µM S1P, 1 µM Capsaicin, and

high K+ Ringer's (red) and one which only responded to high K+ (black). **C.** (Left) Fura-2 AM calcium

imaging before (left) and after (center) addition of 1 μM S1P in $S1pr3^{mCherry/+}$ cultured mouse DRG neurons.

Bar indicates fluorescence ratio. Right-hand image indicates mCherry fluorescence. (Right) % of mCherry

neurons that are responsive to 1 μM S1P in ratiometric calcium imaging (n > 1000 cells from 16 imaging

wells from three animals). **D.** Dose-response curve of mean neuronal calcium responders to varying

concentrations of S1P. Concentrations used: 1, 10, 50, 100, 200, 1000, and 10,000 nanomolar. Error bars

represent mean ± SD (N = 2 animals). Black dotted line indicates sigmoidal fit for all S1P responders from

which EC_{50} was derived. All S1P responders were also capsaicin-responsive. **E.** Current-clamp trace of a

single wild-type neuron firing action potentials in response to bath addition of 1 μM S1P and 1 μM

capsaicin, with Ringer's wash in-between. n = 4 of 10 neurons responded to S1P and N = 1 of 1 S1P-

responsive also responded to capsaicin. Bar = 2 seconds. **F.** (Left) Fura-2 AM calcium imaging after

addition of 500 nM Hm1a in $S1pr3^{mCherry/+}$ P0 TG neurons, which were instead of adult DRG neurons

because they respond to Hm1a without prior PGE_2 sensitization. Right-hand image indicates mCherry

fluorescence. (Right) % of Hm1a-responsive P0 TG neurons that are mCherry+ (N = 1 animal, 1230 total

neurons).

Figure 5. S1PR3 modulates KCNQ2/3 channels to regulate AM excitability. All experiments were

performed in $S1pr3^{mCherry/+}$ or $^{-/-}$ DRG neurons. **A.** (Left) Example traces of a single mCherry+ neuron in

whole cell current clamp before and after S1P application. (Right) % change in rheobase after S1P

application for $S1pr3^{mCherry/+}$ (left) and KO (right) neurons ($p_{WT,KO}$ = 0.012, 0.287; two-tailed paired t-tests; N

= 7, 12 cells). **B.** % Δ in input resistance after S1P or vehicle application (p = 0.017; two-tailed paired t-

test; N = 4 cells per group). **C.** Instantaneous current-voltage relationship after subtraction of the post-S1P

current from the pre-S1P current. Current reverses at -60.125 mV; N = 6 cells. Data were fitted with a

Boltzmann equation. A pre-pulse stimulation of +100 mV for 100 ms was used. Unless indicated otherwise,

all error bars represent mean ± SEOM. **D.** % Δ in inward tail current after S1P or 1% DMSO vehicle

application for $S1pr3^{mCherry/+}$ and KO medium-diameter neurons using a step to +80 mV followed by a step

to -80 mV (p = 0.014; one-way ANOVA; N = 10, 13, 10 cells). Tukey Kramer post hoc p-values indicated

on graph. **E.** (Graphic, top) Averaged current traces of a single mCherry+ neuron in whole cell voltage

clamp recording pre- and post-S1P using indicated voltage step protocol. (graphic, bottom) Averaged

current traces of a single mCherry+ neuron in whole cell voltage clamp recording with XE991 treatment.

Holding phase (-40 mV, 150 ms) was truncated in traces. (Left graph) % Δ in outward tail current (average

+/- SD after indicated treatments (1 µM S1P, 3 µM XE 991, or both) for $S1pr3^{mCherry/+}$ medium-diameter

neurons; (p = 0.58; one-way ANOVA; n = 6, 8, 14 cells) using protocol depicted at right. (Right graph) % Δ

in inward tail current after indicated treatments (LINO = 100 µM linopirdine) for $S1pr3^{mCherry/+}$ medium-

diameter neurons; (p = 0.47; two-tailed paired t-test; N = 12 cells).

Figure 5-source data 1. S1PR3 modulates KCNQ2/3 channels to regulate AM excitability. *Related to*

Figure 5.

Figure 5-figure supplement 1. S1P selectively modulates potassium tail currents to increase DRG

neuron excitability. *Related to Figure 5.* **A.** (Left) Example trace of a single mCherry+ neuron in S1P

before and after current injection. (Right) Resting membrane potential (RMP) in millivolts before and after

addition of S1P (p = 0.23; two-tailed paired t-test; n = 6 cells). **B.** Rheobase pre- and post-S1P application

in DRG neurons; p_{WT} = 0.011; p_{KO} = 0.28 (two-tailed paired t-test). Same data are represented in Figure

5A. **C.** Sodium I-V relationship for a representative $S1pr3^{mCherry/+}$ medium-diameter neuron pre- and 5

minutes post- 1 µM S1P using voltage step from -100 to +80 mV (150 ms steps, -80 mV holding). **D.**

Steady-state I-V relationship for same neuron. **E.** (Left) % Δ in peak instantaneous sodium current after

S1P or 1% DMSO vehicle application for medium-diameter mCherry+ neurons; p = 0.39 (two-tailed paired

t-test; n = 7 cells per group). (Right) % Δ in peak steady-state current after S1P or 1% DMSO vehicle

application for medium-diameter mCherry+ neurons; p = 0.948 (two-tailed paired t-test; n = 7 cells per

group). **F.** Dose-response relationship between % Δ in tail current and S1P concentration for 1 nM, 50 nM,

100 nM, and 1 µM S1P (n = 7 cells). EC_{50} (48.8 nM), marked by thin dotted lines, was estimated from

sigmoidal fit (thick dotted line).

Figure 6. S1PR3 is required for nociceptive responses of high-threshold AM nociceptors. A. (Left)

Representative traces of AM fiber activity over time in *ex vivo* skin-saphenous nerve recording in response

to stimulation (128 mN, top) from HET (middle) and KO (bottom) mice. (Right) Mean firing rate of AM fibers

in response to force controlled stimulation (4,8,16,32,64,128,256 mN). **p=0.001, ***p=0.0002 (two-way

ANOVA, Sidak's post-hoc); lines, linear regression (HET: slope, 50 Hz/N, R^2, 0.99; KO: slope, 35 Hz/N, R^2,

0.95). **B.** von Frey threshold of AM fibers in S1PR3 HET and KO skin. ***$p < 0.0001$ (Mann-Whitney test);

lines, median; boxes, 25-75 percentile; whiskers, min-max. **C.** Conduction velocity (CV) of AM fibers in

S1PR3 HET and KO mice. p = 0.65 (two-tailed t-test); n = 40, 36 fibers; errors, mean ± SEM. **D.**

Cumulative response plot of AM fibers to force controlled stimulation (solid lines); four-parameter logistic fit

from which half-maximal force was estimated for each genotype (dotted lines). **E.** Representative traces of

Non-Adapting and Adapting AMs in response to force controlled stimulation (256 mN, top) for S1PR3 HET

and KO mice; blue regions, dynamic phase of stimulation. **F**. Proportion of fibers classified by pattern of

mechanically evoked responses to 256 mN of stimulation: Non-Responder (HET, 2/40 fibers; KO 5/36),

Non-Adapting AM (HET, 18/40; KO, 29/36), Adapting AM (HET, 20/40; KO, 2/36). Non-Responders fired

action potentials to large magnitude von Frey monofilaments (<0.5 mm tip diameter), but not to 256 mN of

controlled mechanical stimulation (2 mm tip diameter). Classification of Non-Adapting vs. Adapting AMs

detailed in methods. ***p<0.00001 (Chi-square test).

1008

1009   **Figure 6-source data 1. S1PR3 is required for nociceptive responses of high-threshold AM**

1010   **nociceptors.** *Related to Figure 6.* Table provides properties of all S1PR3 HET and KO AM fibers that were

1011   recorded.

1012

Figure 6-figure supplement 1. S1PR3 +/- AM nociceptors display normal nociceptive responses.

Related to Figure 6. **A.** Mean firing rate of AM fibers in response to force controlled stimulation

(4,8,16,32,64,128,256 mN) from **Figure 6A**, with additional data from one wild-type animal (n_{WT} = 4 fibers)

(WT: slope, 57 Hz/N, R^2, 0.98; p = 0.90, two-way ANOVA comparing HET and WT). **B.** (Left) Collected

conduction velocities from teased fibers from 6 S1PR3 KO animals (purple, Ab, centroid=13.1 m/s; yellow,

Ad, centroid=5.7 m/s; orange, C, centroid=0.4 m/s; R^2 =0.69, N = 65 fibers). (Right) Collected conduction

velocities from teased fibers from 1 S1PR3 HET and 6 WT animals (purple, Ab, centroid=14.5 m/s; yellow,

Ad, centroid=6.8 m/s; orange, C, centroid=0.3 m/s; R^2 =0.82, N = 76 fibers). Three-term Gaussian model.

X-axis plotted on a log 1.2 scale.

1022

Figure 7. S1PR3 is dispensable for development of chronic mechanical hypersensitivity. A. Thermal

latency before and after CFA treatment (indicated by dotted line); $p_{genotype}$ = 0.0053 (two-way ANOVA; N = 5

mice per genotype). Sidak's multiple comparison between genotypes for specific time points indicated on

graph. Error bars represent mean ± SD. **B.** (Left) Normalized 50% withdrawal threshold before and after

CFA treatment (indicated by dotted line); p(genotype) < 0.001 (two-way ANOVA). (Right) 50% withdrawal

thresholds for same experiment (p(genotype) = 0.1634; two-way ANOVA). **C.** (Left) Thermal latency

assessed before ("Baseline") and 24 hours post CFA injection with either vehicle (CFA + VEH) or TY 52156

(CFA + TY) acutely administered; p < 0.0001 (one-way ANOVA N = 5 mice per treatment). (Right) Thermal

latency assessed before and after CFA injection with either vehicle (VEH) or SKI II acutely administered on

Day 1; p < 0.0001 (one-way ANOVA; N = 5-7 mice per treatment). Dunnett's test comparisons to baseline

are indicated on graph. Error bars represent mean ± SD. **D.** (Left) 50% withdrawal threshold assessed

before and 24 hours post CFA injection with either vehicle (CFA + VEH) or TY 52156 (CFA + TY) acutely

administered on Day 1; p < 0.0001 (one-way ANOVA; N = 5 mice per treatment). Dunnett's test

comparisons to baseline are indicated on graph. (Right) 50% withdrawal threshold assessed before and

24 hours post CFA injection with either vehicle (CFA + VEH) or SKI II (CFA + SKI II) acutely administered;

p-values indicated on graph (two-tailed unpaired t-test; N = 5 mice per group.

1039

1040   **Figure 7-source data 1. S1PR3 is dispensable for development of chronic mechanical**

1041   **hypersensitivity.** *Related to Figure 7.*

1042

Figure 7-figure supplement 1. S1PR3 KO animals display normal CFA-evoked immune cell

recruitment. *Related to Figure 7.* **A.** Recruitment of neutrophils (Neut.) or inflammatory monocytes (IMs)

to hindpaw skin 24 hours post-CFA administration in S1PR3 HET and KO mice, as a dot plot of CD11b+

cells plotting Ly6G fluorescence intensity vs. Ly6C intensity (AFU). Boxes are for illustrative purposes. **B.**

Recruitment of immune cells, including neutrophils (Neut.) and inflammatory monocytes (IMs) to hindpaw

skin 24 hours post-CFA administration in HET and KO mice, displayed as total number of cells; N = 8 mice

per genotype. Sidak's multiple comparisons were made between HET and KO for each cell type.

1050

1051   **Figure 8. Proposed model illustrating a key role for S1PR3 in regulating mechanical pain in AM**

1052   **nociceptors.** (Top) S1P promotes activation of S1PR3, which leads to inhibition of KCNQ2/3 currents and

1053   promotes normal mechanical pain sensitivity. (Bottom) Diminished S1P or S1PR3 antagonism alleviates

1054   inhibition of KCNQ2/3, leading to mechanical pain hyposensitivity.

Figure 1

Figure 1-figure supplement 1

Figure 2

A In situ hybridization

S1pr3 Marker Merge

Scn1a

Npy2r

Piezo2

Trpv1

B In situ quantification

□ = % marker
□ = % co-loc

% of total cells

Scn1a *Npy2r* *Piezo2* *Trpv1* *Trpa1*

C DRG Immunohistochemistry

Peripherin   NF200

S1PR3   S1PR3
Peripherin   NF200

D Skin Immunohistochemistry

Superficial skin layers   Dermis

S1PR3   S1PR3
NF200   NF200

E Free nerves   Hair follicle innervation

S1PR3   S1PR3 NF200 DAPI Merge

NF200

DAPI   Merkel afferent

Merge   S1PR3 BTIII DAPI Merge

F Immunohistochemistry quantification

DRG   Skin

% co-stained   % of marker

Peripherin NF200 IB4   BT III PGP NF200

Figure 3

Figure 3–figure supplement 1

Figure 4

Figure 5

Figure 5-figure supplement 1

Figure 6

A

B

Figure 6-figure supplement 1

Figure 7

Figure 7-figure supplement 1

In AM nociceptors:

Figure 8