1

The signaling lipid sphingosine 1-phosphate regulates mechanical pain

Authors and Affiliations: Rose Z. Hill[1], Benjamin Hoffman[2,3], Takeshi Morita[1#], Stephanie M. Campos[4‡], Ellen A. Lumpkin[2, 4], Rachel B. Brem[5, 6], Diana M. Bautista[1, 4, 7, 8*]

1. Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.
2. Department of Physiology & Cellular Biophysics, Columbia University College of Physicians & Surgeons, New York, NY 10032
3. Medical Scientist Training Program, Columbia University, New York, NY 10032
4. Neurobiology Course, Marine Biological Laboratory, Woods Hole, MA
5. Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA.
6. Buck Institute for Research on Aging, Novato, CA 94945, USA.
7. Helen Wills Neuroscience Institute, University of California, Berkeley, CA 94720, USA.
8. Lead Contact

Current Address: Rockefeller University, New York, NY 10065, USA.

‡ Current Address: Indiana University, Bloomington, IN 47405, USA.

* Correspondence: dbautista@berkeley.edu

Abstract

Somatosensory neurons mediate responses to diverse mechanical stimuli, from innocuous touch to noxious pain. While recent studies have identified distinct populations of A mechanonociceptors (AMs) that are required for mechanical pain, the molecular underpinnings of mechanonociception remain unknown. Here, we show that the bioactive lipid sphingosine 1-phosphate (S1P) and S1P Receptor 3 (S1PR3) are critical regulators of acute mechanonociception. Genetic or pharmacological ablation of S1PR3, or blockade of S1P production, significantly impaired the behavioral response to noxious mechanical stimuli, with no effect on responses to innocuous touch or thermal stimuli. These effects are mediated by fast-conducting Aδ mechanonociceptors, which displayed a significant decrease in mechanosensitivity in S1PR3 mutant mice. We show that S1PR3 signaling tunes mechanonociceptor excitability via modulation of KCNQ2/3 channels. Our findings define a new role for S1PR3 in regulating neuronal excitability and establish the importance of S1P/S1PR3 signaling in the setting of mechanical pain thresholds.

Introduction

Pain is a complex sensation. It serves to protect organisms from harmful stimuli, but can also become chronic and debilitating following tissue injury and disease. Distinct cells and molecules detect noxious thermal and mechanical stimuli. Thermal pain is detected by thermosensitive TRP channels in subsets of nociceptors[1–4], and gentle touch is detected by Piezo2 channels in low-threshold mechanoreceptors (LTMRs)[5–7]. Aδ high-threshold mechanoreceptors (HTMRs) have been shown to play a key role in responses to painful mechanical stimuli[8–10].

Recent studies have shown that there are at least two populations of HTMRs that mediate responses to noxious mechanical stimuli. The *Npy2r+* subpopulation of HTMRs mediates fast

paw withdrawal responses to pinprick stimulation and terminate as free nerve endings in the

epidermis[11]. The *Calca+* subpopulation of circumferential-HTMRs respond to noxious force and

hair pulling, and terminate as circumferential endings wrapped around guard hair follicles[12].

Additionally, somatostatin-expressing interneurons of laminae I-III in the dorsal horn of the

spinal cord receive input from nociceptors and are required for behavioral responses to painful

mechanical stimuli[13]. Despite these advances in defining the cells and circuits of mechanical

pain, little is known about the molecular signaling pathways in mechanonociceptors.

Here, we show that the bioactive signaling lipid sphingosine 1-phosphate (S1P) is required for

mechanical pain. Mice lacking the S1P receptor S1PR3 display striking and selective deficits in

behavioral responses to noxious mechanical stimuli. Likewise, peripheral blockade of S1PR3

signaling or S1P production impairs mechanical sensitivity. We found that S1P constitutively

enhances the excitability of A mechanonociceptors (AMs) via closure of KCNQ potassium

channels to tune mechanical pain sensitivity. The effects of S1P are completely dependent on

S1PR3. Our findings uncover an essential role for S1P signaling in noxious mechanosensation.

Results

To identify candidate genes underlying mechanosensation, we previously performed

transcriptome analysis of the sensory ganglia innervating the ultra-sensitive tactile organ (the

star) of the star-nosed mole[14]. Immunostaining revealed the tactile organ is preferentially

innervated by myelinated Aδ fibers[14]. While our original analysis focused on ion channels

enriched in the neurons of the star organ, our dataset also revealed enrichment of several

components of the S1P pathway, including S1PR3. Likewise, single-cell RNA seq of mouse

DRG neurons revealed expression of *S1pr3* in neurofilament H-expressing myelinated

mechanoreceptors[15], which includes Aβ as well as the Aδ neuronal populations. Thus, we set

out to define the role of S1P signaling and S1PR3 in somatosensory mechanoreceptors.

S1PR3 mediates acute mechanical pain

We first examined a variety of somatosensory behaviors in mice lacking S1PR3[16]

(*S1pr3^{tm1Rlp/Mmnc}*; referred to herein as *S1pr3^{-/-}*). We initially investigated baseline responses to

mechanical stimuli. *S1pr3^{-/-}* mice displayed a dramatic loss of mechanical sensitivity (Fig. 1A),

as von Frey paw withdrawal thresholds were significantly elevated in *S1pr3^{-/-}* mice relative to

S1pr3^{+/+} and *S1pr3^{+/-}* littermates (mean thresholds: 1.737 g vs. 0.736 and 0.610 g, respectively).

Moreover, *S1pr3^{-/-}* mice demonstrated decreased responses to a range of noxious tactile stimuli

(2-6 g; Fig. 1B) and to noxious pinprick stimulation (Fig. 1C), but normal responsiveness to

innocuous tactile stimuli (0.6-1.4g; Fig. 1B). *S1pr3^{-/-}* mice exhibited normal tape removal

attempts[6] (Fig. 1D), righting reflexes (Fig. S1A), radiant heat withdrawal latencies (Fig. S1B),

and itch-evoked scratching[17] (Fig. S1C). These results demonstrate a selective role for S1PR3

in acute mechanical pain.

As a complement to our analysis of somatosensation in *S1pr3^{-/-}* animals, we employed a

pharmacological approach, using the S1PR3-selective antagonist TY 52156 (TY). Similar to the

phenotype of knockout animals, intradermal injection of 500 μM TY into the mouse hindpaw (the

site of testing) triggered a rapid and significant elevation in von Frey paw withdrawal thresholds

(Fig. 1E) and decreased responsiveness to noxious (2-6 g), but not innocuous (0.6-1.4g), tactile

stimuli (Fig. 1F). In contrast, blockade of S1PR1 with the selective antagonist W146 had no

effect on mechanical thresholds (Fig. 1E). Overall, these data show that S1PR3 signaling sets

mechanical pain sensitivity.

Endogenous S1P mediates acute mechanical pain

We next asked whether peripheral S1P was required for the S1PR3-dependent effects on

mechanosensation. We decreased S1P levels via injection of the sphingosine kinase inhibitor

SKI II to block local production of S1P[18] or elevated S1P levels via intradermal injection of S1P

and measured behaviors 30 minutes after injection. Decreasing local S1P levels significantly

reduced mechanical sensitivity (Fig. 2A), comparable to the hyposensitivity phenotype observed

in $S1pr3^{-/-}$ mice (Fig. 1A). Again, similar to what was observed in $S1pr3^{-/-}$ animals (Fig. S1B),

peripheral blockade of S1PR3 or S1P production had no effect on thermal sensitivity (Fig. S1D).

Surprisingly, injecting exogenous S1P (10 µM; maximum solubility in saline vehicle) had no

effect on mechanical sensitivity (Fig. 2A-B). However, as previously reported[19], S1P injection

triggered thermal hypersensitivity (data not shown) demonstrating that the lack of effect on

mechanical hypersensitivity is not due to problems with S1P delivery or degradation. Although

quantification of native S1P levels in skin is inaccurate owing to avid lyase activity[20], our data

establish that baseline S1P levels are sufficient to maximally exert their effect on S1PR3-

dependent mechanical pain, such that increased levels of S1P do not evoke mechanical

hypersensitivity, but diminished S1P leads to mechanical hyposensitivity. These data show that

constitutive S1PR3 signaling is required for normal mechanosensitivity.

S1PR3 is expressed in A mechanonociceptors

We next aimed to characterize the somatosensory neuron subtypes expressing $S1pr3$ as a

window onto the role of S1PR3 in acute mechanical pain. We used in $situ$ hybridization (ISH)

with a specific $S1pr3$ probe to examine expression patterns of $S1pr3$. In our experiments, 43%

of cells from wild-type dorsal root ganglia (DRG) expressed $S1pr3$ (Fig. 3A; top, Fig. 3E). We

observed no reactivity in DRG isolated from $S1pr3^{-/-}$ mice (Fig. 3A; bottom), and no significant

differences in the distribution of cell sizes between $S1pr3^{-/-}$ and $S1pr3^{+/+}$ ganglia (p = 0.18;

Kolmogorov-Smirnov type II test), suggesting no loss of major neuronal subtypes. While a

previous study using antibody staining concluded that S1PR3 protein was expressed by all DRG

neurons[21], we found that anti-S1PR3 antibodies showed broad immunoreactivity in DRG from

mice lacking S1PR3 (data not shown).

Co-ISH revealed that a population of *S1pr3+* neurons represents Aδ mechanonociceptors

(AMs). These cells were medium-diameter (25.9 ± 4.5 μm) and expressed *Scn1a* (39.9% of all

S1pr3+; Fig. 3B-C), a gene that encodes the Nav1.1 sodium channel which mediates

mechanical pain in Aδ fibers[10], as well as *Npy2r* (20.4% of all *S1pr3+*; Fig. 3B-C), a marker of a

specialized subset of mechanonociceptive A fibers[7]. *S1pr3* was expressed in 70.6% of *Scn1a+*

cells and 72% of *Npy2r+* cells, comprising a majority of both these populations. Interestingly, a

substantial subset of cells co-expressed *S1pr3* and the mechanically sensitive channel *Piezo2*

(Fig. 3C), which is expressed in Aβ, Aδ, and C fibers[6]. The remaining *S1pr3+* cells were small

diameter (23.4 ± 8.2 μm), *Trpv1+* C nociceptors (27.3% of total cells). To elucidate the role of

S1PR3 in mechanical pain, we focused our analysis on Aδ mechanonociceptors, as C-fibers do

not contribute to baseline mechanical thresholds[22].

We also used an *S1pr3*^mCherry/+ reporter mouse[23] as an independent strategy to explore *S1pr3*

expression and localization. Analysis of this line detected expression of S1PR3-mCherry fusion

protein in 48.3% of cultured DRG neurons (Fig. 3E), mirroring the results from ISH. Additionally,

we observed S1PR3 expression in nerve fibers that innervate the skin using antibodies against

mCherry in whole-mount immunohistochemistry (Fig. 3D). We observed overlap of S1PR3-

expressing free nerve endings with NefH+ myelinated A fibers, but did not observe expression

of S1PR3 in circumferential hair follicle receptors (Fig. 3D). These results suggest that S1PR3 is

not expressed in the circ-HTMR subpopulation of mechanonociceptors identified by Ghitani *et*

al. In addition to our mRNA and protein localization experiments, calcium imaging revealed that

a subset of S1PR3+ neurons were activated by the Nav1.1-selective toxin Hm1a[10] (Fig. 3E). We

conclude that a subset of *S1pr3*+ neurons represents AMs that terminate as free nerve endings.

S1PR3 is required for nociceptive responses of high-threshold AM nociceptors

Given that the behavioral deficit of *S1pr3*[-/-] animals was restricted to noxious mechanosensation,

we utilized *ex vivo* skin-nerve recordings to analyze the effects of genetic ablation of S1PR3 on

AM afferents, which mediate fast mechanical pain sensation (Fig. 4A). We hypothesized that

S1PR3 may play a role in AM afferent function. Strikingly, the median von Frey threshold to

elicit firing in AM nociceptors was significantly higher in *S1pr3*[-/-] animals (3.92 mN) compared to

littermate controls (1.56 mN; Fig. 4C). Additionally, *S1pr3*[-/-] AM nociceptors displayed reduced

sensitivity in their force-response relation (slope for [+/-] versus [-/-]: 50 Hz/N versus 35 Hz/N), as

well as attenuated firing over the noxious, but not innocuous, range of mechanical stimulation

(Fig. 4D). Furthermore, *S1pr3*[-/-] AM nociceptors displayed a right-shifted cumulative response

curve (50% effective force for [+/-] versus [-/-]: 33.7 versus 60.0 mN; Fig. 4E), consistent with the

mechanonociceptive hyposensitivity observed *in vivo*. A recent report suggested that A-

nociceptors are composed of two distinct neuronal populations that differ genetically, in

mechanically evoked response patterns, and in conduction velocity[11]. Accordingly, we found

that a proportion of AM nociceptors, characterized by adapting responses to static mechanical

stimuli (Adapting AM), were selectively lost in *S1pr3*[-/-] animals (Fig. 4F-G). However, we found

no differences in conduction velocity between genotypes (Fig. 4B). We conclude that S1PR3 is

an essential regulator of both mechanical threshold and sensitivity in a distinct population of AM

nociceptors.

S1PR3 modulates KCNQ2/3 channels to regulate AM excitability

To define the molecular pathway(s) by which S1PR3 modulates mechanonociceptor function*,*

we next examined S1P signaling in cultured DRG neurons. Unlike the environment of sensory

neurons in intact skin, the physiological Ringer's solutions used *in vitro* lack S1P. Thus for our *in*

vitro experiments we tested the effects of exogenous S1P on neuronal excitability. We

interrogated the molecular mechanism by which S1P signaling regulates mechanical pain using

current-clamp recording of medium-diameter *S1pr3*^(mCherry/+) dissociated DRG neurons

(membrane capacitance = 61.05 ± 1.92 pF), which ISH identified as putative

mechanonociceptors (Fig. 3B-C). In these cells, 1 µM S1P application did not elicit firing in the

absence of current injection (Fig. 5A, Fig. S2A). However, S1P dramatically lowered the

threshold to fire action potentials (rheobase) in an S1PR3-dependent manner (Fig. 5A, Fig.

S2B).

We then set out to determine how S1PR3 activity increases neuronal excitability using whole-

cell voltage clamp recording. While we found that S1P application had no effect on

instantaneous sodium currents or steady state potassium currents (Fig. S2C-D), S1P

significantly reduced slow, voltage-dependent tail current amplitudes (Fig. 5B; Fig. 5D (top)) in

an S1PR3-dependent manner (Fig. 5B, center).

As tail currents in Aδ neurons are primarily mediated by KCNQ2/3 potassium channels[24], we

postulated that S1P might alter tail currents through modulation of these channels. To address

whether KCNQ2/3 channels mediated S1P-dependent neuronal excitability, we examined

several other aspects of the S1P-sensitive current. S1P triggered a robust increase in input

resistance (Fig. 5E), consistent with the closure of potassium channels. Likewise, I-V analysis

revealed that the current inhibited by S1P application was carried by potassium (Fig. 5F).

Additionally, the lack of an effect of S1P on resting membrane potential (Fig. S2E) and steady

state potassium current (Fig. S2D) were consistent with the electrophysiological properties of

KCNQ2/3 channels in DRG neurons[24–27]. Finally, application of the KCNQ2/3-selective inhibitor

linopirdine completely occluded the effects of S1P on tail current (Fig. 5C, Fig. 5d (bottom)).

These findings are consistent with S1P/S1PR3-dependent inhibition of KCNQ2/3 in

somatosensory neurons.

We also found that the effect of S1P on KCNQ2/3 currents was mediated by low levels of S1P,

exhibiting an IC_{50} of 48 nM with saturation at 100 nM (Fig. S2F). While S1P cannot be

accurately measured in non-plasma tissues, this is similar to estimated levels of S1P in

peripheral tissues[28,29]. Thus, our *in vitro* IC_{50} supports our behavioral observations that baseline

S1P levels are sufficient to maximally exert their effect on mechanical pain. In summary, our

electrophysiological and behavioral observations support a model in which baseline S1P/S1PR3

signaling governs mechanical pain thresholds through modulation of KCNQ2/3 channel activity

in AM neurons (Fig. 6).

Discussion

This study identifies S1P/S1PR3 as a key pathway that tunes mechanical pain sensitivity.

Recent studies have identified distinct populations of AM nociceptors that are required for

mechanical pain[11,12]. Likewise, it was discovered that a subset of somatostatin-expressing

spinal interneurons are required for mechanical pain transduction[13]. While these papers

delineate the cells and circuitry of mechanical pain, the molecular underpinnings of

mechanonociception in these neurons were unknown. We now demonstrate that S1PR3 is

required for normal mechanosensitivity in a majority of mechanonociceptors, including the

Npy2r+ population, recently identified by Arcourt *et al.,* that innervates the epidermis and

encodes noxious touch[11].

While the transduction channels that detect noxious force remain enigmatic, we show that

S1PR3 signaling modulates KCNQ2/3 channels to regulate excitability of mechanonociceptors

(Fig. 6). GPCR-mediated inhibition of KCNQ2/3 potassium channels is a well-known mechanism

by which neuronal excitability is regulated[30]. Other studies have shown that KCNQ channels

mediate excitability of Aδ fibers[24] and that opening KCNQ2/3 channels directly with retigabine

alleviates pain *in vivo*[25,31,32]. Our results not only complement previous work implicating

KCNQ2/3 channels in pain, but also define the upstream mechanisms that promote the

regulation of KCNQ2/3 channels to tune mechanical pain thresholds. Our data thus highlight

S1PR3 as a novel and attractive target for the treatment of mechanical pain and describe a new

signaling pathway by which S1P regulates AM nociceptor excitability.

Interestingly, the neurons that innervate the ultra-sensitive tactile organ of the star-nosed mole

are highly enriched in transcripts for S1PR3 and KCNQ channels, as well as for a variety of

other potassium channels[14]. While it is difficult to directly examine the physiological basis for

heightened mechanosensitivity in the star-nosed mole, S1PR3-dependent modulation of KCNQ

may represent an important mechanism underlying the high tactile sensitivity of the star organ.

Outside of the nervous system, S1P signaling via S1PR1 allows for the continuous circulation of

lymphocytes between blood, lymph, and peripheral tissues[33]. Our findings that S1P plays a key

role in somatosensation is in line with recent studies showing that sensory neurons co-opt

classical immune pathways to drive chronic itch or pain[34,35]. What distinguishes this study from

the others is that S1P signaling is critical for acute mechanical pain, even in the absence of

inflammation. In the immune system, disruptions in S1P levels or S1PR1 signaling results in

significant immune dysfunction and disease[36–38]. Paralleling the role of S1PR1 in the immune

system, we propose that S1PR3 might constitutively regulate the excitability of a variety of

neurons and that aberrant S1P signaling may trigger nervous system dysfunction and disease.

For example, S1P has been proposed to constitutively modulate synaptic transmission and

excitability of hippocampal neurons in slice recordings via S1PR3[39–42], and S1PR3-deficient

animals display learning and memory deficits[39]. However, these previous studies, unlike ours,

did not identify a molecular mechanism by which S1P signaling alters neuronal activity.

Abnormal S1P signaling is also linked to a host of neurological disorders, including thermal

hypersensitivity[19], multiple sclerosis[43], Huntington[44], Parkinson's[45], and Alzheimer's disease[46].

In addition to expression in the somatosensory system, S1PR3 is expressed in brain regions

relevant to these disorders (Allen Brain Atlas) and may thus represent an important mechanism

by which S1P signaling regulates excitability.

The relationship between lipids and ion channel activity in the context of mechanotransduction

is well-established and thought to occur through direct channel-lipid interactions[47,48]. Here, we

highlight the importance an indirect pathway in which a lipid activates a GPCR to modulate

mechanoreceptor activity. Our findings complement very recent work demonstrating the diverse

and complex roles that GPCRs play in somatosensory signaling[49]. Our study demonstrates a

crucial role for S1P signaling in the peripheral nervous system and highlights the potential of

S1PR3 as a target for new therapies for mechanical pain.

Materials and Methods

Behavioral studies & mice

S1pr3$^{mcherry/+}$ and *S1pr3$^{-/-}$* mice were obtained from Jackson Laboratory and backcrossed to

C57bl6/J. Wherever possible, wild-type/heterozygous (*S1pr3*) littermate controls were used in

behavioral experiments. Mice (20–25 g) were housed in 12 h light-dark cycle at 21°C. Mice were

singly housed one week prior to all behavioral experiments and were between 8-10 weeks at

the time of the experiment. All mice were acclimated in behavioral chambers on 2 subsequent

days for 1 hour prior to recording for itch behavior, von Frey, and radiant heat.

Itch and acute pain behavioral measurements were performed as previously described[50,51]. Mice

were shaved one week prior to itch behavior. Compounds injected: 500 µM TY 52156 (Tocris),

50 µM SKI II (Tocris), 10 µM S1P (Tocris, Avanti Polar Lipids), 50 mM chloroquine (Sigma), and

27 mM histamine (Tocris) in PBS with either DMSO- or Methanol-PBS vehicle controls.

Pruritogens were injected using the cheek model (20 µL) of itch, as previously described[52].

Behavioral scoring was performed while blind to experimental condition and mouse genotype.

All scratching behavior videos were recorded for 1 hour and scored for the first 30 minutes. Bout

number and length were recorded.

For radiant heat and von Frey hypersensitivity behavior, drugs were injected intradermally into

the plantar surface of the hindpaw (20 µL). Radiant heat assays were performed using the IITC

Life Science Hargreaves test system. Mechanical threshold was measured using calibrated von

Frey monofilaments (Touch Test) on a metal grate platform. Von Frey was performed as

previously described[53,54] using the up-down method while blinded to compound injected and

genotype. Valid responses for both von Frey and radiant heat included fast paw withdrawal,

licking/biting/shaking of the affected paw, or flinching. For radiant heat and von Frey, mice were

allowed to acclimate on platform for 1 hour before injection.

The pinprick assay was conducted on a von Frey testing platform (IITC). The mouse hindpaw

was poked with a 31 g syringe needle without breaking the skin to induce fast acute mechanical

pain. Each paw was stimulated 10 times with the needle, and the % withdrawal (fast withdrawal,

licking/biting/shaking of paw, squeaking, and/or flinching) was calculated from the total number

of trials.

The tape assay was conducted according to previously described methods[6]. Number of

attempts to remove a 3 cm piece of lab tape was recorded for 10 minutes after manual tape

application to the rostral back. Scorer and experimenter were blinded to genotype.

For righting reflex measurements, age-matched *S1pr3*[-/-] and [+/+] P6-7 neonates were used.

Briefly, pups were overturned one at a time on the home cage lid while experimenter was

blinded to genotype. The time to righting was measured to the nearest 1/10[th] of a second with a

stopwatch.

All behavior experiments were carried out using age-matched or littermate cohorts of male mice.

Mice were tested in 4-part behavior chambers (IITC Life Sciences) with opaque dividers (TAP

Plastics) with the exception of righting reflex measurements. Itch behavior was filmed from

below using high-definition cameras. All experiments were performed under the policies and

recommendations of the International Association for the Study of Pain and approved by the

University of California, Berkeley Animal Care and Use Committee.

In situ hybridization (ISH)

Fresh DRG were dissected from 8-12 week old mice, flash frozen in OCT embedding medium,

and sectioned at 14 μm onto slides. ISH was performed using Affymetrix Quantigene ViewISH

Tissue 2-plex kit according to manufacturer's instructions with Type 1 and Type 6 probes. The

following probes against mouse mRNAs were created by Affymetrix and used for ISH: *S1pr3,*

Scn1a, and *Piezo2.* Slides were mounted in Fluoromount with No. 1.5 coverglass. Imaging of

ISH experiments and all other live- and fixed-cell imaging was performed on an Olympus IX71

microscope with a Lambda LS-xl light source (Sutter Instruments). Images were analyzed using

ImageJ software. Briefly, DAPI-positive cells were circled and their fluorescence intensity (AFU)

for all channels was plotted against cell size using Microsoft Excel software. Co-labeling

analysis was performed using ImageJ. Intensity thresholds were set based on the negative

control (no probe) slide. Cells were defined as co-expressing if their maximum intensities

exceeded the threshold for both the Type 1 and Type 6 probe channels.

Immunohistochemistry (IHC) of DRG

8-12 week old mice were deeply anesthetized and transcardially perfused with ice-cold PBS

followed by ice-cold 4% PFA. DRG were dissected and post-fixed in 4% PFA for one hour. DRG

were cryo-protected overnight at 4°C in 30% sucrose-PBS, embedded in OCT, and then

sectioned at 12 μm onto slides. Briefly, slides were washed 3x in PBST (0.3% Triton X-100),

blocked in 2.5% horse serum + 2.5% BSA PBST, and incubated overnight at 4°C in 1:1000

primary antibody in PBST + 0.5% horse serum + 0.5% BSA. Slides were washed 3X in PBS

then incubated 1-2 hours at RT in 1:1000 secondary antibody. Slides were washed 3X in PBS

and mounted in Fluoromount with No. 1.5 coverglass. Primary antibodies used: Rabbit anti-

S1PR3, Mouse anti-NF200, Chicken anti-Peripherin (Abcam). Secondary antibodies used: Goat

anti-Mouse Alexa 350, Goat anti-Chicken Alexa 547, Goat anti-Rabbit Alexa 488 (Abcam). Cells

labeled with anti-peripherin were circled to define regions of interest.

Whole mount skin IHC

Staining was performed according to Marshall and Clary *et al*[55]. Briefly, 8-12 week old mice

were euthanized and the back skin was shaved, depilated, and tape-stripped. The removed skin

was fixed overnight in 4% PFA, then washed in PBS (3X for 10 minutes each). Dermal fat was

scraped away with a scalpel and skin was washed in PBST (0.3% Triton X-100; 3X for two

hours each) then incubated in 1:500 primary antibody (Rabbit anti DsRed Polyclonal antibody;

Clontech #632496) in blocking buffer (PBST with 5% goat serum and 20% DMSO) for 5.5 days

at 4°C. Skin was washed as before and incubated in 1:500 secondary antibody (Goat anti-

Rabbit Alexa 594; Invitrogen #R37117) in blocking buffer for 3 days at 4°C. Skin was washed in

PBST, serially dried in methanol: PBS solutions, incubated overnight in 100% methanol, and

finally cleared with a 1:2 solution of benzyl alcohol: benzyl benzoate (BABB; Sigma) before

mounting.

Cell culture

Cell culture was carried out as previously described[17]. Briefly, neurons from dorsal root ganglia

(2-8 week old adults) or trigeminal ganglia (P0) were dissected and incubated for 10 min in 1.4

mg ml−1 Collagenase P (Roche) in Hanks calcium-free balanced salt solution, followed by

incubation in 0.25% standard trypsin (vol/vol) STV versene-EDTA solution for 2 min with gentle

agitation. Cells were then triturated, plated onto Poly D-Lysine coated glass coverslips and used

within 20 h. Media: MEM Eagle's with Earle's BSS medium, supplemented with 10% horse

serum (vol/vol), MEM vitamins, penicillin/streptomycin and L-glutamine.

Calcium imaging

Ca^{2+} imaging experiments were carried out as previously described[17]. Cells were loaded for 60

min at room temperature with 10 µM Fura-2AM supplemented with 0.01% Pluronic F-127

(wt/vol, Life Technologies) in a physiological Ringer's solution containing (in mM) 140 NaCl, 5

KCl, 10 HEPES, 2 CaCl2, 2 MgCl2 and 10 D-(+)-glucose, pH 7.4. All chemicals were purchased

from Sigma. Acquired images were displayed as the ratio of 340 nm/ 380 nm. Cells were

identified as neurons by eliciting depolarization with high potassium Ringer's solution (75 mM) at

the end of each experiment. Responding neurons were defined as those having a > 15%

increase from baseline ratio. Image analysis and statistics were performed using automated

routines in Igor Pro (WaveMetrics). Fura-2 ratios were normalized to the baseline ratio

F340/F380 = (Ratio)/(Ratio t = 0).

Ex vivo skin-nerve electrophysiology

Touch-evoked responses in the skin were recorded after dissecting the hind limb skin and

saphenous nerve from 7-10 week old mice, according to published methods[56,57]. The skin was

placed epidermis-side-up in a custom chamber and perfused with carbogen-buffered synthetic

interstitial fluid (SIF) kept at 32 °C with a temperature controller (model TC-344B, Warner

Instruments). The nerve was kept in mineral oil in a recording chamber, teased apart, and

placed onto a gold recording electrode connected with a reference electrode to a differential

amplifier (model 1800, A-M Systems). The extracellular signal was digitized using a PowerLab

8/35 board (AD Instruments) and recorded using LabChart software (AD Instruments).

For these studies, we focused on A-mechanonociceptors (AMs). To identify responses from

these afferents in mutant and control genotypes, we used a mechanical search paradigm with a

fine glass probe. Afferents were classified as AMs according to the following criteria: (1) Aδ

conduction velocity (approximately, 1 to (≤ 12 m/s^{-1}), (2) medium-sized receptive fields, (3)

sustained response to mechanical indentation[9,57,58].

Touch-sensitive afferents that did not meet these criteria were not analyzed further. Responses

were classified as Adapting AMs if the ratio of mean firing rate in the dynamic phase of

stimulation (first 0.2 s) to the static phase of stimulation (last 4.8 s) was greater than 2, and Non-

Adapting AMs if the ratio was less than or equal to 2. Non-responders (Fig. 4g) responded to

suprathreshold mechanical stimulation with von Frey monofilaments (tip diameter <0.5 mm), but

not to maximal controlled mechanical stimulation (256 mN, tip diameter 2 mm). All recordings

and analyses were performed blind to genotype.

Mechanical responses were elicited with von Frey monofilaments and a force controlled custom-

built mechanical stimulator. Mechanical thresholds were defined as the lowest von Frey

monofilament to reliable elicit at least on action potential. Force controlled mechanical stimuli

were delivered using a computer controlled, closed-loop, mechanical stimulator (Model 300C-I,

Aurora Scientific, 2 mm tip diameter). Low-pass filtered, 5-second long, length control steps

(square wave) simultaneously delivered with permissive force control steps (square wave) were

generated using LabChart software (AD Instruments). An arbitrarily selected force step-and-hold

protocol (8, 32, 4, 64, 128, 16, 256 mN) was delivered to all fibers. The period between

successive displacements was 60 seconds.

Conduction velocity was measured by electrically stimulating identified receptive fields. Spike

sorting by principal component analysis (PCA) and density based clustering, and data analysis

was performed off-line with custom-made software in MATLAB. Statistics were performed in

Prism.

In vitro electrophysiology

Electrophysiological experiments were carried out as previously described[17]. Briefly, recordings

were collected at 5 kHz and filtered at 2 kHz (Axopatch 200B, pClamp software). Electrode

resistance ranged between 1.5–5 MΩ. Internal solution contained 140 mM KCl, 2 mM MgCl2, 1

mM EGTA, 5 mM HEPES, 1 mM Na2ATP, 100 µM GTP, and 100 µM cAMP (pH 7.4). Bath

solution was physiological Ringer's solution. The pipette potential was canceled before seal

formation. Cell capacitance was canceled before whole cell voltage-clamp recordings.

Experiments were carried out only on cells with a series resistance of less than 30 MΩ. Analysis

of electrophysiology data was performed in pClamp and IgorPro.

Statistical analyses

All statistical analyses, except for skin nerve data (see above), were performed using IgorPro

software or Microsoft Excel. Values are reported as the mean ± SEOM where multiple

independent experiments are pooled and reported (for whole cell electrophysiology), and mean

± SD where one experiment was performed with multiple wells (for calcium imaging) or mice (for

behavior). For comparison between two groups, Student's unpaired 2-tailed t-test was used. A

paired t-test was employed only for measurements within the same biological replicate and after

a given treatment. For single-point comparison between >2 groups, a one-way ANOVA followed

by Tukey Kramer post hoc test was used. For the time course comparison between 2 groups, 2-

way ANOVA was used and single comparison p-values were derived using Tukey's HSD. For

comparing distributions, a type II Kolmogorov-Smirnov test was used. Number of mice or

samples required to attain significance was not calculated beforehand, and where multiple

statistical tests were performed, a Bonferroni correction was applied. In figure legends,

significance was labeled as: n.s., not significant, $p \geq 0.05$; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

Author Contributions

460   R.Z.H, D.M.B., and R.B. conceived experiments and wrote the manuscript. R.Z.H. performed

461   behavioral, immunostaining, whole cell electrophysiology, calcium imaging, and ISH, and

462   analyzed data and made figures. T.M. also performed ISH experiments. E. A. L. & B.H.

463   conceived and analyzed *ex vivo* recordings. B.U.H. and S. C. performed *ex vivo* recordings. All

464   authors contributed to the final version of the manuscript.

Acknowledgements

We thank Z. Rifi (UC Berkeley) for assistance with scoring itch behavior, R. P. Dalton (UC

Berkeley) for assistance with confocal microscopy, and P. Lishko and M. Miller (UC Berkeley)

for advice regarding lipid stability and usage. We would also like to thank R. Clary (Columbia)

for whole mount skin staining protocols and D. Julius (UC San Francisco) for the gift of Hm1a

spider toxin. We are grateful to all members of the D.M.B Laboratory (UC Berkeley) for

constructive feedback and criticism. The National Institutes of Health grants NS077224 (to

D.M.B and R.B.), AR059385 (to D.M.B.), AR051219 (to E.A.L), NS105449 and GM007367 (to

B.U.H), and NS063307 (to the Neurobiology Course at the Marine Biological Laboratory); and a

Howard Hughes Medical Institute Faculty Scholars grant (to D.M.B) supported this work.

Competing interests

The authors declare no competing interests at this time.

478
References

1.   Caterina, M. J., Schumacher, M. a, Tominaga, M., Rosen, T. a, Levine, J. D. & Julius, D.

The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* **389,**

816–824 (1997).

2.   Caterina, M. J, Leffler. A., Malmberg, A. B., Martin, W. J., Trafton, J., Petersen-Zeitz, K.

R., Koltzenburg, M., Basbaum, A. I., Julius, J. D. Impaired Nociception and Pain

Sensation in Mice Lacking the Capsaicin Receptor. *Science* **288,** 306–313 (2000).

3.   McKemy, D. D., Neuhausser, W. M. & Julius, D. Identification of a cold receptor reveals a

general role for TRP channels in thermosensation. *Nature* **416,** 52–58 (2002).

4.   Bautista, D. M., Siemens, J., Glazer, J. M., Tsuruda, P. R., Basbaum, A. I., Stucky, C. L.,

Jordt, S.-E. & Julius, D. The menthol receptor TRPM8 is the principal detector of

environmental cold. *Nature* **448,** 204–208 (2007).

5.   Coste, B., Mathur, J., Schmidt, M., Earley, T. J., Ranade, S., Petrus, M. J., Dubin, A. E. &

Patapoutian, A. Piezo1 and Piezo2 Are Essential Components of Distinct Mechanically

Activated Cation Channels. *Science* **330,** 55–60 (2010).

6.   Ranade, S. S., Woo, S., Dubin, A. E., Moshourab, R. a, Wetzel, C., Petrus, M., Mathur,

J., Bégay, V., Coste, B., Mainquist, J., Wilson, a J., Francisco, A. G., Reddy, K., Qiu, Z.,

Wood, N., Lewin, G. R. & Patapoutian, A. Piezo2 is the major transducer of mechanical

forces for touch sensation in mice. *Nature* **516,** 121–125 (2014).

7.   Woo, S.-H., Ranade, S., Weyer, A. D., Dubin, A. E., Baba, Y., Qiu, Z., Petrus, M.,

Miyamoto, T., Reddy, K., Lumpkin, E. a., Stucky, C. L. & Patapoutian, A. Piezo2 is

required for Merkel-cell mechanotransduction. *Nature* **509,** 622–626 (2014).

8.　Basbaum, A. I., Bautista, D. M., Scherrer, G. & Julius, D. Cellular and Molecular
Mechanisms of Pain. *Cell* **139,** 267–284 (2009).

9.　Koltzenburg, M., Stucky, C. L., Lewin, G. R., Bautista, M., Lechner, S. G., Gemes, G.,
Koopmeiners, A., Rigaud, M., Lirk, P., Sapunar, D., Latha, M., Vilceanu, D., Garrison, S.
R., Ljubkovic, M., Mueller, S. J., Cheryl, L. & Hogan, Q. H. Receptive Properties of Mouse
Sensory Neurons Innervating Hairy Skin Receptive Properties of Mouse Sensory
Neurons Innervating Hairy Skin. *J. Neurophysiol.* **78,** 1841–1850 (1997).

10.　Osteen, J. D., Herzig, V., Gilchrist, J., Emrick, J. J., Zhang, C., Wang, X., Castro, J.,
Garcia-Caraballo, S., Grundy, L., Rychkov, G. Y., Weyer, A. D., Dekan, Z., Undheim, E.
A. B., Alewood, P., Stucky, C. L., Brierley, S. M., Basbaum, A. I., Bosmans, F., King, G.
F. & Julius, D. Selective spider toxins reveal a role for the Nav1.1 channel in mechanical
pain. *Nature* **534,** 494–9 (2016).

11.　Arcourt, A., Gorham, L., Dhandapani, R., Prato, V., Taberner, F. J., Wende, H.,
Gangadharan, V., Birchmeier, C., Heppenstall, P. A. & Lechner, S. G. Touch Receptor-
Derived Sensory Information Alleviates Acute Pain Signaling and Fine-Tunes Nociceptive
Reflex Coordination. *Neuron* **93,** 179–193 (2017).

12.　Ghitani, N., Barik, A., Szczot, M., Thompson, J. H., Li, C., Le Pichon, C. E., Krashes, M.
J. & Chesler, A. T. Specialized Mechanosensory Nociceptors Mediating Rapid
Responses to Hair Pull. *Neuron* **95,** 944–954.e4 (2017).

13.　Duan, B., Cheng, L., Bourane, S., Britz, O., Padilla, C., Garcia-campmany, L., Krashes,
M., Knowlton, W., Velasquez, T., Ren, X., Ross, S. E., Lowell, B. B., Wang, Y., Goulding,
M. & Ma, Q. Identification of Spinal Circuits Transmitting and Gating Mechanical Pain.
Cell **159,** 1417–1432 (2014).

14.　Gerhold, K. a., Pellegrino, M., Tsunozaki, M., Morita, T., Leitch, D. B., Tsuruda, P. R.,
Brem, R. B., Catania, K. C. & Bautista, D. M. The Star-Nosed Mole Reveals Clues to the
Molecular Basis of Mammalian Touch. *PLoS One* **8,** (2013).

15.   Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., Hjerling-Leffler, J.,
Haeggström, J., Kharchenko, O., Kharchenko, P. V, Linnarsson, S. & Ernfors, P.
Unbiased classification of sensory neuron types by large-scale single-cell RNA
sequencing. *Nat. Neurosci.* **18,** 145–153 (2015).

16.   Kono, M., Mi, Y., Liu, Y., Sasaki, T., Allende, M. L., Wu, Y. P., Yamashita, T. & Proia, R.
L. The sphingosine-1-phosphate receptors S1P1, S1P2, and S1P3 function coordinately
during embryonic angiogenesis. *J. Biol. Chem.* **279,** 29367–29373 (2004).

17.   Wilson, S. R., Gerhold, K. a, Bifolck-Fisher, A., Liu, Q., Patel, K. N., Dong, X. & Bautista,
D. M. TRPA1 is required for histamine-independent, Mas-related G protein-coupled
receptor-mediated itch. *Nat. Neurosci.* **14,** 595–602 (2011).

18.   Chiba, Y., Takeuchi, H., Sakai, H. & Misawa, M. SKI-II, an inhibitor of sphingosine kinase,
ameliorates antigen-induced bronchial smooth muscle hyperresponsiveness, but not
airway inflammation, in mice. *J. Pharmacol. Sci.* **114,** 304–310 (2010).

19.   Mair, N., Benetti, C., Andratsch, M., Leitner, M. G., Constantin, C. E., Camprubí-Robles,
M., Quarta, S., Biasio, W., Kuner, R., Gibbins, I. L., Kress, M. & Haberberger, R. V.
Genetic evidence for involvement of neuronally expressed s1P1 receptor in nociceptor
sensitization and inflammatory pain. *PLoS One* **6,** (2011).

20.   Shaner, R. L., Allegood, J. C., Park, H., Wang, E., Kelly, S., Haynes, C. a, Sullards, M. C.
& Merrill, A. H. Quantitative analysis of sphingolipids for lipidomics using triple quadrupole
and quadrupole linear ion trap mass spectrometers. *J. Lipid Res.* **50,** 1692–1707 (2009).

21.   Camprubi-Robles, M., Mair, N., Andratsch, M., Benetti, C., Beroukas, D., Rukwied, R.,
Langeslag, M., Proia, R. L., Schmelz, M., Ferrer Montiel, A. V, Haberberger, R. V, Kress,
M. & Camprubí-Robles, M. Sphingosine-1-phosphate-induced nociceptor excitation and
ongoing pain behavior in mice and humans is largely mediated by S1P3 receptor. *J.*
Neurosci. **33,** 2582–2592 (2013).

22.   Abrahamsen, B., Zhao, J., Asante, C. O., Cendan, C. M., Marsh, S., Martinez-Barbera, J.

P., Nassar, M. A., Dickenson, A. H. & Wood, J. N. The Cell and Molecular Basis of

Mechanical, Cold, and Inflamatory Pain. *Science* **321,** 702 (2008).

23. Sanna, M. G., Vincent, K. P., Repetto, E., Nhan, N., Brown, S. J., Abgaryan, L., Riley, S.

W., Leaf, N. B., Cahalan, S. M., Kiosses, W. B., Kohno, Y., Heller Brown, J., McCulloch,

A. D., Rosen, H. & Gonzalez-Cabrera, P. J. Bitopic S1P3 Antagonist Rescue from

Complete Heart Block: Pharmacological and Genetic Evidence for Direct S1P3

Regulation of Mouse Cardiac Conduction. *Mol. Pharmacol.* **89,** 176–186 (2015).

24. Schu, S., Orozco, I. J. & Jentsch, T. J. KCNQ Potassium Channels Modulate Sensitivity

of Skin Down hair (D-hair) Mechanoreceptors. *J. Biol. Chem.* **291,** 5566–5575 (2016).

25. Zheng, Q., Fang, D., Liu, M., Cai, J., Wan, Y., Han, J. S. & Xing, G. G. Suppression of

KCNQ/M (Kv7) potassium channels in dorsal root ganglion neurons contributes to the

development of bone cancer pain in a rat model. *Pain* **154,** 434–448 (2013).

26. Crozier, R. A., Ajit, S. K., Kaftan, E. J. & Pausch, M. H. MrgD Activation Inhibits KCNQ /

M-Currents and Contributes to Enhanced Neuronal Excitability. *J. Neurosci.* **27,** 4492–

4496 (2007).

27. Heidenreich, M., Lechner, S. G., Vardanyan, V., Wetzel, C., Cremers, C. W., De

Leenheer, E. M., Aránguez, G., Moreno-Pelayo, M. Á., Jentsch, T. J. & Lewin, G. R.

KCNQ4 K(+) channels tune mechanoreceptors for normal touch sensation in mouse and

man. *Nat. Neurosci.* **15,** 138–145 (2012).

28. Bäumer, W., Rossbach, K., Mischke, R., Reines, I., Langbein-Detsch, I., Lüth, A. &

Kleuser, B. Decreased concentration and enhanced metabolism of sphingosine-1-

phosphate in lesional skin of dogs with atopic dermatitis: disturbed sphingosine-1-

phosphate homeostasis in atopic dermatitis. *J. Invest. Dermatol.* **131,** 266–268 (2011).

29. Ramos-Perez, W. D., Fang, V., Escalante-Alcalde, D., Cammer, M. & Schwab, S. R. A

map of the distribution of sphingosine 1-phosphate in the spleen. *Nat. Immunol.* **16,**

1245–52 (2015).

30.   Suh, B.-C., Inoue, T., Meyer, T. & Hille, B. Rapid Chemically Induced Changes of

580   PtdIns(4,5)P2 Gate KCNQ Ion Channels. *Science* **314,** 1454–1457 (2006).

581   31.   Blackburn-Munro, G. & Jensen, B. S. The anticonvulsant retigabine attenuates

582   nociceptive behaviours in rat models of persistent and neuropathic pain. *Eur. J.*

583   *Pharmacol.* **460,** 109–116 (2003).

584   32.   Hayashi, H., Iwata, M., Tsuchimori, N. & Matsumoto, T. Activation of peripheral KCNQ

585   channels attenuates inflammatory pain. *Mol. Pain* **10,** 15 (2014).

586   33.   Matloubian, M., Lo, C. G., Cinamon, G., Lesneski, M. J., Xu, Y., Brinkmann, V., Allende,

587   M. L., Proia, R. L. & Cyster, J. G. Lymphocyte egress from thymus and peripheral

588   lymphoid organs is dependent on S1P receptor 1. *Nature* **427,** 355–360 (2004).

589   34.   Pinho-Ribeiro, F. A., Verri, W. A. & Chiu, I. M. Nociceptor Sensory Neuron–Immune

590   Interactions in Pain and Inflammation. *Trends Immunol.* **38,** 5–19 (2017).

591   35.   Oetjen, L. K., Mack, M. R., Feng, J., Whelan, T. M., Niu, H., Guo, C. J., Chen, S., Trier, A.

592   M., Xu, A. Z., Tripathi, S. V., Luo, J., Gao, X., Yang, L., Hamilton, S. L., Wang, P. L.,

593   Brestoff, J. R., Council, M. L., Brasington, R., Schaffer, A., Brombacher, F., Hsieh, C.-S.,

594   Gereau, R. W., Miller, M. J., Chen, Z.-F., Hu, H., Davidson, S., Liu, Q. & Kim, B. S.

595   Sensory Neurons Co-opt Classical Immune Signaling Pathways to Mediate Chronic Itch.

596   *Cell* **129,** 742–751 (2017).

597   36.   Donoviel, M. S., Hait, N. C., Ramachandran, S., Maceyka, M., Takabe, K., Milstien, S.,

598   Oravecz, T. & Spiegel, S. Spinster 2, a sphingosine 1-phosphate transporter, plays a

599   Critical Role in inflammatory and Autoimmune Diseases. *FASEB J.* **29,** 5018–5028

600   (2015).

601   37.   Schwab, S. R., Pereira, J. P., Matloubian, M., Xu, Y., Huang, Y. & Cyster, J. G.

602   Lymphocyte sequestration through S1P lyase inhibition and disruption of S1P gradients.

603   *Science* **309,** 1735–1739 (2005).

38.   Allende, M. L., Bektas, M., Lee, B. G., Bonifacino, E., Kang, J., Tuymetova, G., Chen, W.,

Saba, J. D. & Proia, R. L. Sphingosine-1-phosphate lyase deficiency produces a pro-

inflammatory response while impairing neutrophil trafficking. *J. Biol. Chem.* **286,** 7348–

7358 (2011).

39.   Weth-Malsch, D., Langeslag, M., Beroukas, D., Zangrandi, L., Kastenberger, I., Quarta,

S., Malsch, P., Kalpachidou, T., Schwarzer, C., Proia, R. L., Haberberger, R. V. & Kress,

M. Ablation of Sphingosine 1-Phosphate Receptor Subtype 3 Impairs Hippocampal

Neuron Excitability In vitro and Spatial Working Memory In vivo. *Front. Cell. Neurosci.* **10,**

1–12 (2016).

40.   Mitroi, D. N., Deutschmann, A. U., Raucamp, M., Karunakaran, I., Glebov, K., Hans, M.,

Walter, J., Saba, J., Gräler, M., Ehninger, D., Sopova, E., Shupliakov, O., Swandulla, D.

& van Echten-Deckert, G. Sphingosine 1-phosphate lyase ablation disrupts presynaptic

architecture and function via an ubiquitin- proteasome mediated mechanism. *Sci. Rep.* **6,**

37064 (2016).

41.   Chan, J. P. & Sieburth, D. Localized Sphingolipid Signaling at Presynaptic Terminals Is

Regulated by Calcium Influx and Promotes Recruitment of Priming Factors. *J. Neurosci.*

32, 17909–17920 (2012).

42.   Riganti, L., Antonucci, F., Gabrielli, M., Prada, I., Giussani, P., Viani, P., Valtorta, F.,

Menna, E., Matteoli, M. & Verderio, C. Sphingosine-1-Phosphate (S1P) Impacts

Presynaptic Functions by Regulating Synapsin I Localization in the Presynaptic

Compartment. *J. Neurosci.* **36,** 4624–4634 (2016).

43.   Choi, J. W., Gardell, S. E., Herr, D. R., Rivera, R., Lee, C.-W., Noguchi, K., Teo, S. T.,

Yung, Y. C., Lu, M., Kennedy, G. & Chun, J. FTY720 (fingolimod) efficacy in an animal

model of multiple sclerosis requires astrocyte sphingosine 1-phosphate receptor 1 (S1P1)

modulation. *Proc. Natl. Acad. Sci. U. S. A.* **108,** 751–6 (2011).

44.   Di Pardo, A., Amico, E., Favellato, M., Castrataro, R., Fucile, S., Squitieri, F. & Maglione,

V. FTY720 (fingolimod) is a neuroprotective and disease-modifying agent in cellular and

mouse models of huntington disease. *Hum. Mol. Genet.* **23,** 2251–2265 (2014).

45.   Zhao, P., Yang, X., Yang, L., Li, M., Wood, K., Liu, Q. & Zhu, X. Neuroprotective effects

of fingolimod in mouse models of Parkinson's disease. *FASEB J.* **31,** 172–179 (2017).

46.   Couttas, T. a, Kain, N., Daniels, B., Lim, X. Y., Shepherd, C., Kril, J., Pickford, R., Li, H.,

Garner, B. & Don, A. S. Loss of the neuroprotective factor Sphingosine 1-phosphate early

in Alzheimer's disease pathogenesis. *Acta Neuropathol. Commun.* **2,** 9 (2014).

47.   Nomura, T., Cranfield, C. G., Deplazes, E., Owen, D. M., Macmillan, a., Battle, a. R.,

Constantine, M., Sokabe, M. & Martinac, B. Differential effects of lipids and lyso-lipids on

the mechanosensitivity of the mechanosensitive channels MscL and MscS. *Proc. Natl.*

Acad. Sci. **109,** 8770–8775 (2012).

48.   Brohawn, S. G., Su, Z. & MacKinnon, R. Mechanosensitivity is mediated directly by the

lipid membrane in TRAAK and TREK1 K+ channels. *Proc. Natl. Acad. Sci. U. S. A.* **111,**

3614–9 (2014).

49.   Scholz, N., Guan, C., Nieberler, M., Grotemeyer, A., Maiellaro, I., Gao, S., Beck, S.,

Pawlak, M., Sauer, M., Asan, E., Rothemund, S., Winkler, J., Prömel, S., Nagel, G.,

Langenhan, T. & Kittel, R. J. Mechano-dependent signaling by Latrophilin/CIRL quenches

cAMP in proprioceptive neurons. *Elife* **6,** 1–21 (2017).

50.   Morita, T., McClain, S. P., Batia, L. M., Pellegrino, M., Wilson, S. R., Kienzler, M. A.,

Lyman, K., Olsen, A. S. B., Wong, J. F., Stucky, C. L., Brem, R. B. & Bautista, D. M.

HTR7 Mediates Serotonergic Acute and Chronic Itch. *Neuron* **87,** 124–138 (2015).

51.   Tsunozaki, M., Lennertz, R. C., Vilceanu, D., Katta, S., Stucky, C. L. & Bautista, D. M. A

'toothache tree' alkylamide inhibits Aδ mechanonociceptors to alleviate mechanical pain.

J. Physiol. **591,** 3325–40 (2013).

52.   Shimada, S. G. & LaMotte, R. H. Behavioral differentiation between itch and pain in

mouse. *Pain* **139,** 681–687 (2008).

53.   Dixon, W. J. The Up-and-Down Method for Small Samples. *J. Am. Stat. Assoc.* **60,** 967–

978 (1980).

54.   Chaplan, SR, Bach, FW, Pogrel, JW, Chung, JM, Yaksh, T. Quantitative assessment of

tactile allodynia evoked by unilateral ligation of the fifth and sixth lumbar nerves in the rat.

J. Neurosci. **53,** 55–63 (1994).

55.   Marshall, K. L., Clary, R. C., Baba, Y., Orlowsky, R. L., Gerling, G. J. & Lumpkin, E. A.

Touch Receptors Undergo Rapid Remodeling in Healthy Skin. *Cell Rep.* **17,** 1719–1727

(2016).

56.   Maksimovic, S., Nakatani, M., Baba, Y., Nelson, A. M., Marshall, K. L., Wellnitz, S. A.,

Firozi, P., Woo, S.-H., Ranade, S., Patapoutian, A. & Lumpkin, E. A. Epidermal Merkel

cells are mechanosensory cells that tune mammalian touch receptors. *Nature* **509,** 617–

621 (2014).

57.   Wellnitz, S. A., Lesniak, D. R., Gerling, G. J. & Lumpkin, E. A. The Regularity of

Sustained Firing Reveals Two Populations of Slowly Adapting Touch Receptors in Mouse

Hairy Skin. *J. Neurophysiol.* **103,** 3378–3388 (2010).

58.   Zimmermann, K., Hein, A., Hager, U., Kaczmarek, J. S., Turnquist, B. P., Clapham, D. E.

& Reeh, P. W. Phenotyping sensory nerve endings in vitro in the mouse. *Nat. Protoc.* **4,**

(2009).

Figure 1. S1PR3 mediates acute mechanical pain. A. von Frey 50% withdrawal threshold measurements for *S1pr3*$^{+/+}$ (WT, N = 8), $^{+/-}$ (HET, N = 7) and $^{-/-}$ (KO, N = 12) mice. $p < 0.0001$ (one-way ANOVA). Tukey Kramer post hoc comparisons for KO and HET to WT indicated on graph. B. von Frey force-response graph for WT (N = 8) versus KO (N = 12) animals; $p_{genotype}$ < 0.0001 (two-way ANOVA). Tukey HSD comparisons between genotypes are indicated for given forces. C. % withdrawal to pinprick stimulation of hindpaw for HET versus KO animals; $p < 0.0001$ (unpaired t-test; N = 5-7 mice per group). D. Number of attempted removal bouts in tape assay for WT (N =2), HET (N = 2), and KO (N = 5) mice; $p = 0.172$ (one-way ANOVA). E. von Frey 50% withdrawal threshold measurements for mice pre- and post-injection of 500 μM TY 52156 (N = 10), 10 μM W146 (N = 6), or 1% DMSO-PBS vehicle (N = 17); p = 0.016, 0.650 (two-tailed paired t-test comparing vehicle-vs. drug-injected paw). F. von Frey force-response graph for mice injected with either 1% DMSO-PBS or 500 μM TY 52156; $p_{treatment}$ < 1e-05 (two-way ANOVA; N = 4 mice per group). Tukey HSD comparisons were made between treatment groups and significant differences at a given force are indicated on graph. Unless otherwise indicated, error bars represent mean ± SD. See Source Data Table 1 for additional details.

Figure 2. Endogenous S1P mediates acute mechanical pain. A. von Frey 50% withdrawal measurements for mice pre- and post-injection of 50 µM SKI II (N = 8) or 10 µM S1P (N = 7); p = 0.003, 0.604 (two-tailed paired t-test). B. von Frey force-response graph for animals injected with 10 µM S1P or 0.1% MeOH-PBS; $p_{genotype}$ > 0.05 (two-way ANOVA; N = 8 mice per group). No Tukey HSD comparisons at any force between genotypes were significant. Error bars represent mean ± SD. See Source Data Table 2 for additional details.

Figure 3. S1PR3 is expressed in A mechanonociceptors. A. ISH of *S1pr3* (green) in sectioned DRG from adult wild-type (top) and S1PR3 KO (bottom) mice (20x air objective; scale = 50 μm). B. Representative Co-ISH of *S1pr3* (green; left) with *Scn1a*, *Npy2r,* and *Piezo2* (magenta; center). Right column: overlay with co-localized regions colored white (10x air objective; scale = 100 μm). C. Bar chart showing the % of total cells expressing the indicated marker (grey) and the % of total cells co-expressing both marker and *S1pr3* (green) from the images in Fig. 3B. D. Whole-mount skin immunohistochemistry with anti-DsRed antibody (green) and anti-NefH antibody (magenta) in an *S1pr3*$^{mCherry/+}$ animal (20x water objective; scale = 50 μm). Arrows indicate S1PR3+ epidermal fibers (left image) or S1PR3- circumferential fibers (right image). E. Quantification of % of total cells expressing *S1pr3* from DRG ISH and mCherry from dissociated DRG cultures (N = 2 animals each experiment). F. (Left) Fura-2 AM calcium imaging before (left) and after (center) addition of 500 nM Hm1a in *S1pr3*$^{mCherry/+}$ P0 TG neurons. Right-hand image indicates mCherry fluorescence. Arrows indicate one mCherry+/Hm1a-responsive cell. (Right) % of Hm1a-responsive P0 TG neurons that are mCherry+ (N = 1 animal, 1230 total neurons).

Figure 4. S1PR3 is required for nociceptive responses of high-threshold AM nociceptors. A. Representative traces of AM fiber activity over time in *ex vivo* skin-saphenous nerve recording in response to stimulation (128 mN, top) from S1PR3[+/-] (HET) (middle) and [-/-] (KO) (bottom) mice. B. Conduction velocity (CV) of AM fibers in S1PR3 HET and KO mice. $p = 0.65$ (two-tailed t-test); N = 40, 36 fibers; errors, mean ± SEM. C. von Frey threshold of AM fibers in S1PR3 HET and KO skin. ***$p < 0.0001$ (Mann-Whitney test); lines, median; boxes, 25-75 percentile; whiskers, min-max. D. Mean firing rate of AM fibers in response to force controlled stimulation (4,8,16,32,64,128,256 mN). **$p=0.001$, ***$p=0.0002$ (two-way ANOVA, Sidak's post-hoc); lines, linear regression (HET: slope, 50 Hz/N, R^2, 0.99; KO: slope, 35 Hz/N, R^2, 0.95). E. Cumulative response plot of AM fibers to force controlled stimulation (solid lines); four-parameter logistic fit from which half-maximal force was estimated for each genotype (dotted lines). F. Representative traces of Non-Adapting and Adapting AMs in response to force controlled stimulation (256 mN, top) for S1PR3 HET and KO mice; blue regions, dynamic phase of stimulation. G. Proportion of fibers classified by pattern of mechanically evoked responses to 256 mN of stimulation: Non-Responder (HET, 2/40 fibers; KO 5/36), Non-Adapting AM (HET, 18/40; KO, 29/36), Adapting AM (HET, 20/40; KO, 2/36). Non-Responders fired action potentials to large magnitude von Frey monofilaments (<0.5 mm tip diameter), but not to 256 mN of controlled mechanical stimulation (2 mm tip diameter). Classification of Non-Adapting vs. Adapting AMs detailed in methods. ***$p<0.00001$ (Chi-square test). See Source Data Table 4 for additional details.

Figure 5. S1PR3 modulates KCNQ2/3 channels to regulate AM excitability. All experiments were performed in *S1pr3*[mCherry/+] or [-/-] DRG neurons. A. (Left) Example traces of a single mCherry+ neuron in whole cell current clamp before and after S1P application. (Right) % change in rheobase after S1P application for *S1pr3*[mCherry/+] (left) and KO (right) neurons ($p_{WT,KO}$ = 0.012, 0.287; two-tailed paired t-test; N = 7, 12 cells). B. % Δ in tail current after S1P or 1% DMSO vehicle application for *S1pr3*[mCherry/+] and KO medium-diameter neurons (p = 0.014; one-way ANOVA; N = 10, 13, 10 cells). Tukey Kramer post hoc *p*-values indicated on graph. C. % Δ in tail current after indicated treatments (LINO = 100 μM linopirdine) for *S1pr3*[mCherry/+] medium-diameter neurons; (p = 0.47; two-tailed paired t-test; N = 12 cells). D. (Top) Averaged tail current traces of a single mCherry+ neuron in whole cell voltage clamp recording pre- and post-S1P. (Bottom) Averaged tail current traces of a single mCherry+ neuron in whole cell voltage clamp recording with indicated treatment. E. % Δ in input resistance after S1P or vehicle application (p = 0.017; two-tailed paired t-test; N = 4 cells per group). F. Instantaneous current-voltage relationship after subtraction of the post-S1P current from the pre-S1P current. Current reverses at -60.125 mV; N = 6 cells. Data were fitted with a Boltzmann equation. A pre-pulse stimulation of +100 mV for 100 ms was used. Unless indicated otherwise, all error bars represent mean ± SEOM. See Source Data Table 5 for additional details.

In AM nociceptors:

Figure 6. Proposed model illustrating a key role for S1PR3 in regulating mechanical pain in AM nociceptors. S1P promotes activation of S1PR3, which leads to inhibition of KCNQ2/3 currents and promotes normal mechanical pain sensitivity. Diminished S1P or S1PR3 antagonism alleviates inhibition of KCNQ2/3, leading to mechanical pain hyposensitivity.

Figure S1. Loss of S1PR3 or S1P selectively impairs mechanonociception. *Related to Figure 2.* A. Time to righting in seconds for N = 6 P7 pups per genotype for WT and KO mice; $p = 0.575$ (two-tailed unpaired t-test). B. Baseline radiant heat measurements for WT, HET, and KO mice. N = 8 WT, 3 HET, and 5 KO mice. 3 measurements from both paws were averaged for each mouse. $p = 0.444$ (one-way ANOVA). C. (Left) Time spent scratching in response to injection of 50 mM chloroquine or PBS vehicle (VEH) in KO mice; $p = 0.042$ (unpaired t-test; N = 3,4 mice). (Right) Time spent scratching in response to injection of 27 mM Histamine or 0.1% DMSO-PBS in KO mice; $p = 0.019$ (unpaired t-test; N = 3,4 mice). D. Normalized paw withdrawal latencies post-injection of SKI II, TY 52156, or 0.1% DMSO-PBS vehicle into the hind paw of wild-type animals; $p = 0.65$ (one-way ANOVA); N = 5 mice per group. Unless otherwise indicated, error bars represent mean ± SD.

Figure S2. S1P selectively modulates potassium tail currents to increase DRG neuron excitability. *Related to Figure 5.* A. Example trace of a single mCherry+ neuron in S1P before and after current injection. B. Rheobase pre- and post-S1P application in DRG neurons; p_{WT} = 0.011; p_{KO} = 0.28 (two-tailed paired t-test). Same data are represented in Figure 5A. C. % Δ in instantaneous sodium current after S1P or 1% DMSO vehicle application for medium-diameter mCherry+ neurons; p = 0.39 (two-tailed paired t-test; N = 7 cells per group). D. % Δ in steady-state current after S1P or 1% DMSO vehicle application for medium-diameter mCherry+ neurons; p = 0.948 (two-tailed paired t-test; N = 7 cells per group). E. Resting membrane potential (RMP) in millivolts before and after addition of S1P (p = 0.23; two-tailed paired t-test; N = 6 cells). F. Dose-response relationship between % Δ in tail current and S1P concentration for 1 nM, 50 nM, 100 nM, and 1 μM S1P (N = 7 cells). EC_{50} (48.8 nM), marked by thin dotted lines, was estimated from sigmoidal fit (thick dotted line).