1   Defining the seafloor microbiome of the Gulf of Mexico and its response to oil

2   perturbation

Will A. Overholt[1], Patrick Schwing[2], Kala M. Raz[1], David Hastings[3], David J. Hollander[2], Joel E. Kostka[1,4*]

[1]School of Biological Sciences, Georgia Institute of Technology, Atlanta GA; [2]College of Marine Science,

University of South Florida, St. Petersburg, FL; [3]Department of Marine Science, Eckerd College, St.

Petersburg, FL; [4]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA

Keywords: biogeography, deep ocean sediment, microbial community, Deepwater Horizon, response

and recovery, perturbations

Running title: Deepsea benthic microbial biogeography

*Corresponding author: Joel E. Kostka

Georgia Institute of Technology
310 Ferst Drive
Atlanta, Georgia 30332
+1 (404) 385-3325
joel.kostka@biology.gatech.edu
20
21

22   Conflict of interest: The authors declare that they have no conflict of interest.

Abstract

The microbial ecology of oligotrophic deep ocean sediments is understudied relative to their shallow counterparts, and this lack of understanding hampers our ability to predict responses to current and future perturbations. The Gulf of Mexico has experienced two of the largest accidental marine oil spills, the 1979 Ixtoc-1 blowout and the 2010 Deepwater Horizon (DWH) discharge. Here, microbial communities were characterized for 29 sites across multiple years in >700 samples. The composition of the seafloor microbiome was broadly consistent across the region and was well approximated by the overlying water depth and depth within the sediment column, while geographic distance played a limited role. Biogeographical distributions were employed to generate predictive models for over 4000 OTU that leverage easy-to-obtain geospatial variables which are linked to measured sedimentary oxygen profiles. Depth stratification and putative niche diversification are evidenced by the distribution of taxa that mediate the microbial nitrogen cycle. Further, these results demonstrate that sediments impacted by the DWH spill had returned to near baseline conditions after two years. The distributions of benthic microorganisms in the Gulf can be constrained, and moreover deviations from these predictions may pinpoint impacted sites and aid in future response efforts or long-term stability studies.

Introduction

Deep ocean sediments are the largest ecosystems on Earth and cover approximately 65 % of the Earth's surface (1). Solute transport rates in these sediments are dominated by diffusion, while larger particles are transported through burial or via bioturbation (2). The biogeochemistry regimes of deep ocean sediments are governed by the quality and quantity of deposited organic matter, most of which originated in the photic zone via photosynthesis or via transport from terrestrial sources (3,4). Furthermore, most organic matter is decomposed in the water column during transport to the seafloor (5). Therefore, oligotrophic sediments typically have very low organic carbon contents and < 20 % of sedimentary organic matter is recognizable as one of the main biomolecular classes (5,6).

In contrast to the complexity and patchiness of the sedimentary carbon content, the oligotrophic deep ocean is characterized by consistent physico-chemical gradients (temperature, pressure, salinity, pH) that are often noted as the primary variables in controlling bacterial diversities (7,8). In part, this temporal and spatial stability is thought to contribute to the higher diversity and lower variation in community structure observed in deep marine sediments compared to their coastal and pelagic counterparts (3,9,10). The main environmental variables that are considered to drive microbial community composition in the deepsea are the organic carbon content, water depth, geographic distance, and overlying ecosystem productivity (9–11).

While deep marine sediments are thought to be relatively stable, they are not impervious to ecological disturbances (12–14). One glaring example of such a disturbance is the Deepwater Horizon oil spill (DWH), which represents the largest accidental marine oil spill in history, releasing an estimated 4.1 – 4.9 million barrels of crude oil and a third as much natural gas at approximately 1500 m water depth (15,16). Although the amounts and chemical composition of DWH oil remaining in the environment are uncertain, the largest reservoir is thought to be primarily in deep ocean sediments with a considerable, but unknown, amount deposited near coastal sediments (17). Approximately 12% of the 2 million

barrels entrained in the deep ocean was transported to a 3200 km^2 region of the seafloor (18,19). The

fate of this oil will largely be determined by microbial metabolisms in these environments and by burial.

A substantial challenge in determining the impacts of this oil being deposited into deep ocean

sediments is the paucity of studies on benthic microbial communities performed prior to the DWH

discharge. Due to interest in petroleum exploration, previous studies in the Gulf focused almost entirely

on deep subsurface sediments, or on sediments impacted by natural hydrocarbon seeps (20–22). Due to

the lack of knowledge on surficial microbial communities, the majority of studies on deep ocean

sediments associated with the DWH event used samples that were collected outside of impacted areas

as controls (23–26). While the datasets on benthic microbes are limited to relatively few samples,

results indicate that microbial communities responded quickly to oil perturbation and shifts in

community composition as well as total metabolic potential were observed (23–25).

The Ixtoc-I well blowout that began in 1979 in the Gulf shares many similarities with DWH. The

event released approximately 3.5 million barrels of crude oil, and represents the second largest

accidental marine oil spill (27). Like DWH, Ixtoc-I crude oil was saturated with gaseous hydrocarbons and

had a similar chemical composition. Approximately 25% of the total oil released is thought to have

remained in seafloor sediments (27). Unfortunately, the long-term fate of Ixtoc-I oil is not well

understood (28), and due to technical limitations at the time, little is known about the *in situ* microbial

community response (29).

Here, we present the largest benthic marine microbial community study conducted to date, and

by far the most extensive in the Gulf of Mexico. We focus on two Gulf regions that were both impacted

by massive marine oil spills. Over 700 samples were analyzed representing 29 distinct sites, across four

years, including archaeal and bacterial communities. Our objectives were to (1) characterize un-

impacted sedimentary microbial communities to establish a baseline, (2) map the biogeographical

patterns in microbial community structure across the Gulf of Mexico, (3) using this map, generate a Gulf

of Mexico biogeography model of microbial community structure that can predict the abundance of

dominant microbial populations, and (4) determine if impacted regions had returned to baseline

conditions. We determine the environmental parameters structuring microbial communities, including

the stratification of microbial communities according to sediment depth and across oxygen gradients. In

addition, we address the temporal stability of microbial communities across the region.

Material and Methods

Sample Collection

Samples were collected during five research cruises from 2012 to 2015 across the Northern and Southern Gulf of Mexico (see Table S1, Figure 1). Sediments were sampled using a MC-800 multicorer capable of collecting up to eight, 10-cm diameter by 70 cm long cores per deployment. Cores were sectioned at either sub-centimeter (2 mm sections) or 1 cm resolution using a calibrated threaded rod attached to a core clamp stand with a tight fitting plunger (30). Sectioned sediments were frozen immediately in a portable -80 °C freezer onboard the vessel and transported back to the laboratory for DNA extraction (Table S1).

In the northern Gulf of Mexico (GoM), samples were collected across approximately 100,000 square kilometers, primarily focusing on the Desoto Canyon feature and near the Deepwater Horizon well-head (Table S1, Figure 1). Sampling ranged from 56 m (SL1040) near the Mississippi River Delta to 2293 m (DSH09) water depth southeast of the DWH wellhead, spanning three years (2012 – 2014) and four sampling time points (Table S1). In the southern Gulf of Mexico, sampling sites spanned approximately 40,000 square kilometers ranging from 16-1470 m water depth, primarily focusing on the Ixtoc-I well blowout site and along the western transect in the direction surface oil traveled (Figure 1) (31,32). Cores from the southern GoM were all collected from one cruise in 2015. For each region, sediment core profiles extended to at least 10 cm depth below the seafloor, with a subset including sections up to 20 cm of the sediment column. Our experimental design incorporated multiple levels of replication to address variation due to sampling effort, as well as spatial and temporal heterogeneity. These include: (1) technical replicates with duplicate DNA extractions performed and separate libraries generated on the same depth section, (2) multicore cast replicates where triplicate cores from the same cast were analyzed separately representing variation across approximately $1m^2$ of the seafloor, (3) replicates at the site level where triplicate casts were deployed at the same site and cores from each

cast were analyzed separately, and (4) temporal replicates where the same site was sampled up to 4

times in 3 years (Figure S1).

Oxygen Profiles

Oxygen concentrations were profiled in the sediments using a Presens Microx4 needle-

microoptode (Presens, Regensburg, Germany) along with a CHPT micromanipulator (Composite High

Pressure Technologies, Lewes, Delaware, USA). Immediately after the cores were retrieved from the

multicorer, a core was mounted on the extruder for stabilization and oxygen was profiled at mm-scales

until the sensor measured 0 umol/L oxygen concentration. Between sites, the optode was calibrated

following the manufacturer's protocol. The depth of oxygen penetration was imported into Ocean Data

View v4.6.3 (33) and predictive surfaces were generated using DIVA gridding with automatic scale

lengths and with a signal to noise ratio of 50.

DNA Extraction, SSU rRNA Gene Sequencing and Processing

Frozen sediment was sub-sectioned using a flame sterilized drillbit and total genomic DNA

extractions performed on approximately 0.25 g using a MoBio Powersoil DNA extraction kit following

the manufacturer's protocol. The V4 region of the SSU rRNA was amplified using the Earth Microbiome

Project 515F/806R primer set and sequenced using Illumina MiSeq 2x250 v2 chemistry (34,35). Steps are

briefly outlined here and are described in detail in the Supplemental Information. Following QA/QC,

operational taxonomic units (OTUs) were picked using Swarm v2.1.8 (36,37). Taxonomic affiliations for

each OTU were determined using the SILVA v123 database (38). OTU counts were normalized using the

metagenomeSeq package in R (39). Beta diversity was determined using Bray Curtis dissimilarity and

visualized with nonmetric multidimensional scaling (NMDS) plots using the R package 'vegan' (40,41).

Patterns in Bray Curtis dissimilarities were tested using multiple linear regressions in R (41). The most

significant regression used the formula (dissimilarity ~ sediment depth difference * water depth

difference * geographic distance * Sampling Year) which was simplified to (dissimilarity ~ sediment

depth difference + water depth difference + geographic distance + sampling year + sed:water) by

including only variables contributing >1% variance. Linear regression models were compared using

Analysis of Variance (ANOVA), and variable contribution was determined with ANOVA (Table S2, Table

S3). All plots generated use the ggplot2 package in R (41,42). All raw sequences have been uploaded to

NCBI under Bioproject PRJNA414249 and all associated data and scripts can be accessed at

https://doi.org/10.7266/N70G3HN3.

Constructing a Gulf-Wide Model for Microbial Community Composition

A model for microbial community composition across the Gulf of Mexico was constructed using

random forest regression implemented in the R package 'randomForest' at the OTU level (43). Random

Forest regression is an ensemble machine learning method where multiple individual decision trees are

constructed and ultimately averaged to minimize over-fitting and improve prediction accuracy.

Individual decision trees are generated from a subset of the response variable (OTU abundance) and

each fork or node in the tree is guided by predictor variables (randomized subset) to maximize

differences in the branches. Random Forest is robust to outliers, does not assume specific distributions

(nonparametric), and tests/trains the model by subsampling the response variable at each step and

testing the decision node with the remaining data (11).

Water depth, sediment depth, latitude, and longitude were chosen as predictor variables and

were used as proxies for sediment carbon input, oxygen concentrations, geological regime shifts from E-

W and N-S as well as riverine inputs. Latitude, longitude, and water depths were extracted from the

NGDC Coastal Relief model (44). The Random Forest regression model for each response variable was

constructed with 500 trees, a minimum of 5 terminal nodes per tree, and randomly subsampling 1/3 of

the response variable (OTU normalized counts) for each node of each tree. Only response variables of

which >50% of the variance could be explained by the regression forest were retained. Details

concerning the evaluation of the models are included in the Supplemental Information.

Results and Discussion

Environmental parameters structuring benthic microbial distributions across the Gulf of Mexico

This study incorporated sediment samples across approximately 100,000 and 40,000 square kilometers in the northern and southern Gulf of Mexico, respectively. Sampling sites ranged from 16 m to 2300 m water depth, focusing on regions adjacent to the Deepwater Horizon (DWH) and Ixtoc I blowouts. The variation in community structure across a range of spatial scales was tested at multiple replication levels (Figure S1). Technical and cast level (1 m^2 bottom area) replicates showed similar Bray-Curtis dissimilarity values (mean = 0.23, sd= ± 0.05; 0.25 ± 0.05, respectively), while site level replicates (10-100 m^2) were less similar (0.34 ± 0.09). When comparing the same site over time (across 3 years), lower similarity (0.54 ± 0.12) was observed, although this value may be inflated due to a small change in DNA extraction technique between 2013 and 2014 sampling years. All sites sampled at greater than 500 m water depth for the entire study show a mean dissimilarity of 0.61 ± 0.15 in comparison to a mean dissimilarity of 0.51 ± 0.13 for sites sampled at less than 100 m.

Overall, the distribution of microbial populations across these geographically distant regions of the Gulf were remarkably consistent and were structured primarily based on sediment depth along with overlying water depth (Figure 2 A, B). The sampling design focused primarily on shallow sites (<100 m) and deep sites (>1000 m), and thus the down core sediment progressions seen in Figure 2 A are structured into two main clusters primarily along the Y-axis. However, a continuous gradient with increasing water depth is expected, and indeed, is observed at the sites >1000 m water depth. Performing a multiple linear regression analysis and partitioning the variance resulted in water depth explaining 22.1% of the total variation (F = 1.29e5, p < 2e-16, Table S2).

The effect of sediment depth is primarily observed along the x-axis with surficial sediment samples clustering in the upper right of Figure 2 proceeding down core towards the bottom left, consistent with other studies (Figure 2 B) (6). Sediment depth explained 14.2% of the total microbial

community variation (F = 8.3e4, p < 2e-16), and together with water depth, could explain 36% of the

observed variance (74% of the total explained variance, Table S2). In contrast, geographic distance was

of limited effectiveness in distinguishing microbial community compositions, explaining just 6% of the

variance (F = 3.6e4, p < 2e-16), or 4% if using a discrete variable for basin region (Figure 2 C). Similarly,

the sampling time point explained only 5% of the variance (F = 2.9e4, p < 2e-16) and differences were

not visible in Figure 2 D. In total, water depth, sediment depth, geographic distance, sampling time

point, and their interactions explained 52% of the observed variation, and a simplified model could

explain 49% of the observed variance (Table S2, see Supplementary Information). Patterns in alpha

diversity (Shannon, estimated OTUs, rarefaction curves) revealed lower metrics in samples collected at

less than 100 m water depth compared to those collected at greater than 400 m, while changes in

sediment depth were minor (Figure S2). The remarkable consistency in the Gulf seafloor microbiome

may be attributed to the stability of deepsea environments, which are fueled by organic matter and

nutrients supplied by the surface ocean (8). Unlike previous research, our dataset reveals that the

downcore distribution of deepsea microbial communities below the surface across a large area is also

likely determined by the same environmental forcings.

Previous research has mainly focused only on the biogeographic patterns of microbial

communities in surface sediments, both within ocean basins and globally, and this is discussed in detail

in the Supplemental Information (8,9,45–47). In general, water depth appears to have more explanatory

power within a region while geographic distance and carbon bioavailability tends to have more power in

global studies (1,9,10). The results generated in this study greatly contribute to the growing body of

knowledge surrounding the environmental parameters structuring the seafloor microbiome. In contrast

to previous work, this is the first study to examine large scale biogeographic patterns within a sediment

core-profile from the shallow to the deep ocean. The larger number of samples and high depth of

sequencing employed in this study along with focusing on a relatively small ocean region provided

higher spatial resolution than previous work, which likely contributes to the strong biogeographic

patterns in microbial community composition observed. Future global studies are encouraged to explore

constraining the variation in microbial communities using a basin-based nested approach that maintains

water and sediment depths as explanatory variables that facilitate the construction of predictive models

for microbial communities.

Sedimentary oxygen concentrations linked to intra-basin microbial biogeography in the Gulf of
Mexico

Sedimentary oxygen profiles were determined for 42 sites across the northern and southern

regions of the Gulf of Mexico using a micro-optode and micromanipulator at millimeter resolution.

Oxygen profiles were obtained at most sites where cores were sectioned for microbial community

composition analysis, as well as over a substantially larger area (Figure 1, Table S1). Typical core profiles

are found in Figure 4 (B, D), while the interpolated maximum oxygen penetration depths across the

study area are shown in Figure 3 A. At shallow, neritic sites, oxygen was depleted within the first

millimeter, while oxygen penetration depths extended to 16 cm at the deepest sites (~ 3700 m). As

anticipated, oxygen penetration increased nearly linearly with increasing water depths (Figure 3 B) (48).

This likely results from a greater amount of phytodetritus reaching the seafloor in shallower

environments as well as increases in primary productivity closer to the coast (49,50). On the shelf,

elevated chlorophyll α and β concentrations and depleted major nutrients are indicative of higher

primary production (51). The oxic-anoxic interface was closer to the sediment surface in the northern

Gulf in comparison to the southern Gulf at similar water depths, likely due to a lower sedimentary

carbon deposition regulated by riverine inputs (Figure 3 C). The northern Gulf of Mexico is greatly

influenced by the Mississippi River, the fifth largest river by volume in the world and the primary source

(~90%) of both sediment and freshwater to the entire Gulf (52–54). This region also has one of the

largest and best studied anthropogenically induced "dead zones" in the world, linked to the high

nutrient loadings, large phytoplankton blooms, and massive deposition events to the northern Gulf of

Mexico (55,56). Furthermore, in late 2010 following the Deepwater Horizon oil spill, a massive offshore

(>300m – 1500m sites) sedimentation event was observed whereby oily marine snow was deposited at

4-5 fold higher rates than typically observed (57,58). Relative to the northern Gulf of Mexico, the

southern Gulf of Mexico generally receives lower riverine inputs (~10% of the northern Gulf input) (54).

However, our southern Gulf samples were collected in close proximity to the second largest riverine

system in the Gulf of Mexico, the Grijalva-Usumacinta, as well as the smaller Coatzacoalcos and

Papaloapan systems (31,59).

In agreement with contrasting oxygen penetration depths (Figure 3), significant differences in

microbial community composition (p < 2e-16, var. explained = 4%) were observed between the northern

and southern Gulf of Mexico, although these differences were small relative to the impacts of overlying

water column depth and sediment depth (Figure 2). When comparing the microbial community

composition of sediments from similar water depths in each region, community composition in the

northern Gulf of Mexico more closely resembled shallower sites in the southern Gulf of Mexico (Figure

4, Supplemental Information). Following a simple adjustment for water depth, the sedimentary

microbial communities in these geographically distinct regions with such different quantities of riverine

inputs were remarkably similar.

The core seafloor microbiome across the Gulf of Mexico

In corroboration of previous work conducted in deep ocean oligotrophic sediments across the

globe, we observed that members of the Gammaproteobacteria and Deltaproteobacteria dominate

surficial communities (Figure 5, Supplemental Information). In contrast to past work, this study reveals

that the MG-I Thaumarchaeota are the second most abundant class detected in the Gulf seafloor. This

group likely went unreported in previous studies due to recent advances in sequencing technology and

primer design (9,10). The majority of studies have focused on bacterial groups, whereas when universal

primers are used, Marine Group I Thaumarchaea are shown to be very abundant (47). Although

previous work in the Gulf of Mexico did not incorporate Archaea, our results on bacterial distributions

are further corroborated by clone libraries of bacterial communities constructed from surficial

sediments that were sampled prior to oil impacts which showed similar dominant classes (24).

Class level distributions along increasing sediment depths were generally consistent across core

profiles at different water depths. However, a main difference was the visible inflection point in class

relative abundances that coincided with oxygen depletion (Figure 5). These inflection points shifted

down the sediment column with increasing water depth while becoming broader. This pattern was

clearly evidenced by comparing the changes in relative abundance of the aerobic Marine Group I

(Thaumarchaeota) in cores from sites DSH08 (1110 m), DSH10 (1500 m) and DSH09 (2293 m) (Figure 5).

In addition, the transition between MG-I Thaumarchaeota and Phycisphaerae typically coincided with

the oxygen gradient and the oxic-anoxic interface, which increased in sediment column with increasing

water depth. Furthermore, this oxic-anoxic transition zone coincided with highest relative abundances

of Planctomycetacia. Similar patterns, although with different methods and at different scales, have

been elucidated in shallow, organic rich sediments in Aarhus Bay, where similar delineations in

transition zones were linked to bioturbation activity coinciding with oxygen penetration and increased

labile organic matter availability (60).

Our comprehensive dataset reveals distinct and predictable patterns in numerous OTU on a

basin wide scale (Figure 6, 7c). The majority of these OTUs are not closely affiliated with characterized

taxa. For example, the dataset reveals that the most abundant archaeal OTU in the Gulf seafloor

microbiome is distinct from previously characterized archaea, which are mainly chemolithoautotrophic

ammonia-oxidizers (Figure 6, Figure S6). The most abundant OTU within the Gammaproteobacteria is

also uncharacterized and is consistently distributed at the oxic-anoxic interface (Figure S7). The most

abundant and ubiquitously distributed OTU in the whole dataset is affiliated with *Syntrophobacter*, at

low sequence identity (Figure S8). Given that the function of these microbial groups is not yet known,

we discuss the distribution of these and many other dominant OTU further in the supplementary

material. Thus, the discussion is focused on taxa putatively associated with the marine nitrogen cycle

(Figure S11).

The four most abundant Marine Group I OTU (Thaumarchaeota) were all phylogenetically

related to *Nitrosopumilus maritimus* (3 OTU >97% sequence identity and 1 OTU at 93% identity).

Distributions of these four abundant *Nitrosopumilus*-like OTU were not driven by depth within the

sediment column as was found by Durbin and Teske, and instead were related to the overlying water

depth, showing putative niche differences due to organic matter inputs (2010) (Figure S6). Marine

Group I Thaumarchaeota are well known members of oxic and suboxic sediments (1,47,61–65). All

characterized isolates similar to these 4 OTU are aerobic autotrophs capable of oxidizing ammonia at

very low concentrations. Increases in their relative abundances have been shown to be concurrent with

decreases in the quality and quantity of organic carbon within sediments (47). It has also been shown

that in deep, low energy pelagic environments, a large portion of *Nitrosopumilus*-like populations were

capable of using urea as a carbon and ammonia source to fuel autotrophic nitrification (66).

All dominant OTU affiliated with the Planctomycetacia showed high sequence identity to

Candidatus Scalindua spp (99-100%). These dominant OTU are expected to be anaerobic ammonium

oxidizing bacteria (anammox), which are chemoautotrophic, ubiquitously found in anoxic systems, and

couple the oxidation of ammonium with nitrite as the electron acceptor (67,68). Members of this genus

tend to be psychrophilic, and are inhibited by low sulfide concentrations (4-100 µM), which potentially

explains their depth limits in our study, although it may also be due to ammonium or nitrite availability

(67). The Planctomycetes were nearly absent in the shallowest sites, and absent in surficial sediments; in

agreement with previous studies that indicate ca. *Scalindua* is more abundant in oligotrophic systems

(47,69). All dominant Planctomycetes OTU tended to reach a maximal relative abundance at mid-

sediment depths below the oxic-anoxic interface with an overlapping distribution to the *Nitrosopumulis-*

like OTUs (Figures 6, S4, S10). Due to their overlapping distributions, ammonium oxidizing archaea may

provide a nitrite source for ca. *Scalindua* (70).

In parallel with the distribution of ca. *Scalindua*-like microbial populations, putative bacterial

ammonium oxidizers (*Nitrosococcus-like*, and an unclassified Nitrosomonadaceae OTU) were also absent

in nearshore sediments or in surficial sediments from deep ocean sites and exhibited an increase in

relative abundance with increasing water depth (Figure S11). Abundant putative nitrite oxidizers

affiliated with *Nitrospira* and *Nitropina* groups were detected that were also only present in deep water

sediments below the aerobic-anaerobic transition zone, similar to the *Scalindua*-like population

distributions. While circumstantial, the relatively high abundances of these putative groups in deep

ocean sediments provides evidence of inorganic nitrogen species being important electron acceptors

and donors in deep Gulf of Mexico sediments, particularly for fueling autotrophic metabolisms. These

observations are supported by previous studies showing the complexity of nitrogen cycling in similar

sediments as well as the growing body of knowledge towards the importance of chemolithoautotrophs

in oligotrophic deep ocean sediments (1,71–73)

Construction of a predictive microbial community composition model throughout the Gulf of

Mexico

After visualizing the consistent biogeographic patterns portrayed in Figure 2, it was recognized

that a regression-based model may effectively reconstruct the relative abundances of microbial groups

across these regions. A machine learning method, random forest regression, was employed that has

previously been successful in population distribution models (11,74–79). As stated in the methods

section, the model was constructed using the variables water depth, sediment depth, latitude, and

longitude as proxies for carbon input, redox state, and geographic variations due to riverine inputs and

differing geochronology across the Gulf of Mexico (57).

This analysis was run for every abundantly detected OTU (9588 OTU). All OTU with greater than

50% variance explained by the full model were retained (4129 OTU). Overall, the models fit the

observed data well (R^2 = 0.88). Residuals were weakly correlated with OTU abundance as the model

tends to underestimate the very abundant OTU. The model was further tested by calculating the root

mean squared error (RMSE) compared to 1000 bootstrapped randomly permuted RMSE and was found

to be highly significant ($p < 2e^{-16}$, Z-test).

To verify the utility of the model, model predictions were compared with observed counts for

the dominant OTU (Figure S5, S6-S10). We anticipate that the model can be used as a tool to assess

future impacted sediments that may be poorly characterized and to test this, we hindcasted the

microbial community composition at previously impacted sediments. This was accomplished by

calculating the total root mean squared error (RMSE) for each sample using observed OTU abundance

and predicted OTU abundances, including the Mason et al. (2014) surficial sample set (Figure 7 C). The

higher the RMSE value for each sample, the great the observed community composition deviated from

the model prediction. The most impacted sites had higher RMSE values than the "below EPA limits"

sites, although both were skewed to the right of the model predictions for our data (Figure 7 C).

Therefore, while controlling for experimental based variation, this model can still be effective at

determining impacts.

Impacts of the DWH and internal testing of the predictive model

The earliest samples included in this study were collected in August 2012, over two years after

the Deepwater Horizon well blowout. No evidence of a large oil spill signature was detected in

sediments sampled from areas known to be impacted by sedimentary oil deposition related to the DWH

blowout (18,80–82). At the broad biogeographic scale visualized in Figure 2, the lack of an "oil-impacted

cluster" indicated that any potential impact on community composition was small relative to the strong

environmental gradients influenced primarily by water depth and sediment depth. To compare our data

explicitly with sediments that were shown to be strongly impacted by oil, a meta-analysis based

approach was employed. We observe that all surficial sediment samples from this study, including those

collected close to the DWH wellhead, clustered adjacent to samples classified as below EPA limits and

were distinct from those classified as most impacted by oiling (23) (Figure 7 B). Furthermore, the

similarities between the below EPA limit samples and our dataset indicated a temporally stable and

Gulf-wide conserved microbial community relative to a strong pulse disturbance. Finally, previously

impacted sediments that were collected and analyzed during this study did not deviate from model

predictions, unlike those collected in 2010 (Figure7 C).

These results are corroborated by a previous study that detailed a microbial community

succession event in oil impacted surficial sediments immediately following the oiled marine snow

sedimentation event, through the generation of full length SSU rRNA gene clone libraries (24). The

authors suggested that by November 2010 the microbial community composition once again resembled

pre-impacted surficial microbial communities (24). Our first samples were collected in August 2012 and

much of our dataset comes from 2013 and 2014 for the NGoM, years following a potential return to

baseline condition. Over much larger scales, our results do not show a disturbance signature in cores

collected at the Ixtoc-I wellhead or within the PEMEX (Mexican state-owned petroleum company)

restricted area that has extensive oil extraction pressures. The lack of an Ixtoc-I signature in the

microbial community is expected considering the resilience observed in the sedimentary communities in

the northern Gulf of Mexico, and the lack of a PEMEX oil signature indicates we did not sample

sediments that were recently exposed to oil.

Conclusions

The primary objectives of this study were to define the core microbiome across the Gulf of

Mexico seafloor, characterize biogeographic patterns in microbial populations in Gulf sediments, and

use these results to constrain the impacts of petroleum hydrocarbons from major oil spills to

sedimentary microbial communities. Using the largest marine sediment microbial community

composition dataset generated to date, we show that the distribution of microbial communities is

broadly consistent across the entire region in terms of the OTU detected and their relative abundances.

We further demonstrate that the distribution of seafloor microbial communities is well approximated by

the overlying water depth and depth within the sediment column, which together explain 38% of our

observed variation, regardless of geographic sampling region. The small differences between the regions

are potentially related to the amount of organic carbon reaching the seafloor, with the southern Gulf

showing greater depths of oxygen penetration as well as microbial communities that were more similar

to deeper-water counterparts in the northern Gulf. Microbial community composition across the Gulf

revealed similar taxonomic groups as observed in other marine sediment biogeography studies across

the globe and is dominated primarily by groups that are poorly characterized with unknown ecological

functions within the sediment. Putative ammonium oxidizing archaea (*Nitrosopumilus*-like OTU) were

among the most relatively abundant groups detected in the dataset across oxygenated sediments in the

Gulf of Mexico, while other putative chemolithoautotrophic nitrogen-transforming groups were

relatively abundant only in the deep ocean sediments. Overall, the dominant OTU detected from each of

the major classes, generally had low sequence identities to characterized taxa.

The lack of a strongly perturbed community in our dataset, along with the incorporation of

previous studies demonstrating an oil related perturbation signal, suggests that impacted sediment

communities had returned or are returning to baseline conditions. Finally, the biogeographical

distributions of seafloor microbial populations were used to generate a predictive model for over 4000

dominant populations that relies on easy-to-obtain geospatial variables. With this model, it is possible to

predict community compositions across the Gulf of Mexico, and use deviations from such a prediction as

a tool to assess future impacted sediments that may be poorly characterized along with natural

ecosystems that are conspicuously different from oligotrophic ecosystems such as cold seeps.

Acknowledgements

This work was made possible in part by a grant from The Gulf of Mexico Research Initiative to the C-

IMAGE II consortium and in part by a National Science Foundation Graduate Research Fellowship (WAO)

under grant no. 2013172310. The authors would like to thank the ship crews on the R/V Weatherbird III

and the R/V Justo Sierra that made the sample collection possible. Data are publicly available through

the Gulf of Mexico Research Initiative Information and Data Cooperative (GRIIDC) at

https://data.gulfresearchinitiative.org (doi: 10.7266/N70G3HN3). Any opinions, findings, and

conclusions or recommendations expressed in this material are those of the authors and do not

necessarily reflect the views of the National Science Foundation.

Competing Interests

414   The authors declare that they have no conflict of interest.

References

References

1. Danovaro R, Molari M, Corinaldesi C, Dell'Anno A. Macroecological drivers of archaea and

bacteria in benthic deep-sea ecosystems. Sci Adv [Internet]. 2016;2(4):1–12. Available from:

http://advances.sciencemag.org/content/2/4/e1500961.abstract

2. Burdige DJ. Geochemistry of marine sediments. Princeton University Press; 2006.

3. Jørgensen BB, Boetius A. Feast and famine — microbial life in the deep-sea bed. Nat Rev

Microbiol [Internet]. 2007;5(10):770–81. Available from:

http://www.nature.com/doifinder/10.1038/nrmicro1745

4. Arndt S, Jørgensen BB, LaRowe DE, Middelburg JJ, Pancost RD, Regnier P. Quantifying the

degradation of organic matter in marine sediments: A review and synthesis. Earth-Science Rev

[Internet]. 2013 Aug [cited 2014 May 28];123:53–86. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0012825213000512

5. Canfield D, Thamdrup B, E Kristensen. Aquatic Geomicrobiology. San Diego, CA: Elsevier; 2006.

6. Teske A. Marine deep sediment microbial communities. In: Rosenberg E, DeLong EF, Lory S,

Stackebrandt E, Thompson F, editors. The Prokaryotes [Internet]. Berlin, Heidelberg: Springer

Berlin Heidelberg; 2013 [cited 2014 May 23]. p. 123–33. Available from:

http://link.springer.com/10.1007/978-3-642-30123-0

7. Lozupone C a, Knight R. Global patterns in bacterial diversity. Proc Natl Acad Sci U S A.

2007;104(27):11436–40.

8. Bienhold C, Boetius A, Ramette A. The energy–diversity relationship of complex bacterial

communities in Arctic deep-sea sediments. ISME J [Internet]. 2012;6(4):724–32. Available from:

http://dx.doi.org/10.1038/ismej.2011.140

9.   Zinger L, Amaral-Zettler L a, Fuhrman J a, Horner-Devine MC, Huse SM, Welch DBM, et al. Global

patterns of bacterial beta-diversity in seafloor and seawater ecosystems. PLoS One [Internet].

2011 Jan [cited 2013 Nov 11];6(9):e24570. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3169623&tool=pmcentrez&rendert

ype=abstract

10.   Bienhold C, Zinger L, Boetius A, Ramette A. Diversity and biogeography of bathyal and abyssal

seafloor bacteria. PLoS One. 2016;11(1):1–20.

11.   Wei C-L, Rowe GT, Escobar-Briones E, Boetius A, Soltwedel T, Caley MJ, et al. Global Patterns and

Predictions of Seafloor Biomass Using Random Forests. PLoS One [Internet]. 2010;5(12):e15323.

Available from: http://dx.plos.org/10.1371/journal.pone.0015323

12.   Billett DSM, Lampitt RS, Rice AL, Mantoura RFC. Seasonal sedimentation of phytoplankton to the

deep-sea benthos. Vol. 302, Nature. 1983. p. 520–2.

13.   Smith CR, Baco AR. Ecology of whale falls at the deep-sea floor. Oceanogr Mar Biol. 2003;41:311–

54.

14.   Aller JY. Quantifying sediment disturbance by bottom currents and its effect on benthic

communities in a deep-sea western boundary zone. Deep Sea Res Part A, Oceanogr Res Pap.

1989;36(6):901–34.

15.   Joye SB, MacDonald IR, Leifer I, Asper V. Magnitude and oxidation potential of hydrocarbon gases

released from the BP oil well blowout. Nat Geosci [Internet]. 2011 Feb 13 [cited 2013 Jun

2];4(3):160–4. Available from: http://www.nature.com/doifinder/10.1038/ngeo1067

16.   Lubchenco J, McNutt M, Lehr B, Sogge M, Miller M, Hammond S, et al. Deepwater Horizon/BP Oil

458   Budget: What happened to the oil? National Oceanic and Atmospheric Administration Report.

459   Silver Spring, MD; 2010.

460   17.   King GMM, Kostka JE, Hazen TCC, Sobecky PAA. Microbial Responses to the Deepwater Horizon

461   oil spill: From Coastal Wetlands to the Deep Sea. Ann Rev Mar Sci. 2014;

462   18.   Valentine DL, Fisher GB, Bagby SC, Nelson RK, Reddy CM, Sylva SP, et al. Fallout plume of

463   submerged oil from Deepwater Horizon. Proc Natl Acad Sci [Internet]. 2014 Oct 27 [cited 2014

464   Oct 28];1–6. Available from: http://www.pnas.org/cgi/doi/10.1073/pnas.1414873111

465   19.   Joye SB. Deepwater Horizon, 5 years on. Science (80-). 2015;349(6248):592–3.

466   20.   Biddle JF, White JR, Teske AP, House CH. Metagenomics of the subsurface Brazos-Trinity Basin

467   (IODP site 1320): comparison with other sediment and pyrosequenced metagenomes. ISME J

468   [Internet]. 2011 Jun [cited 2014 Jun 9];5(6):1038–47. Available from:

469   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3131849&tool=pmcentrez&rendert

470   ype=abstract

471   21.   Nunoura T, Soffientino B, Blazejak A, Kakuta J, Oida H, Schippers A, et al. Subseafloor microbial

472   communities associated with rapid turbidite deposition in the Gulf of Mexico continental slope

473   (IODP Expedition 308). FEMS Microbiol Ecol [Internet]. 2009 Sep [cited 2014 Jun 25];69(3):410–

474   24. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19583789

475   22.   Orcutt BN, Joye SB, Kleindienst S, Knittel K, Ramette A, Reitz A, et al. Impact of natural oil and

476   higher hydrocarbons on microbial diversity, distribution, and activity in Gulf of Mexico cold-seep

477   sediments. Deep Sea Res Part II Top Stud Oceanogr. 2010;57(21):2008–21.

478   23.   Mason OU, Scott NM, Gonzalez A, Robbins-Pianka A, Bælum J, Kimbrel J, et al. Metagenomics

479   reveals sediment microbial community response to Deepwater Horizon oil spill. ISME J [Internet].

2014 Jan 23 [cited 2014 Jan 23];(October 2013):1–12. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24451203

24.   Yang T, Speare K, McKay L, MacGregor BJ, Joye SB, Teske A. Distinct Bacterial Communities in

Surficial Seafloor Sediments Following the 2010 Deepwater Horizon Blowout. Front Microbiol

[Internet]. 2016;7(September):1–18. Available from:

http://journal.frontiersin.org/article/10.3389/fmicb.2016.01384

25.   Kimes NE, Callaghan A V, Aktas DF, Smith WL, Sunner J, Golding B, et al. Metagenomic analysis

and metabolite profiling of deep-sea sediments from the Gulf of Mexico following the Deepwater

Horizon oil spill. Front Microbiol [Internet]. 2013 Jan [cited 2013 Jun 25];4(March):50. Available

from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3598227&tool=pmcentrez&rendert

ype=abstract

26.   Liu Z, Liu J. Evaluating bacterial community structures in oil collected from the sea surface and

sediment in the northern Gulf of Mexico after the Deepwater Horizon oil spill. Microbiologyopen

[Internet]. 2013 Apr 9 [cited 2013 Jun 3]; Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23568850

27.   Jernelöv A, Lindén O. Ixtoc I: a case study of the world's largest oil spill. Ambio. 1981;299–306.

28.   Schrope M. The lost legacy of the last great oil spill. Nature Publishing Group; 2010.

29.   Atlas RM. Fate of oil from two major oil spills: Role of microbial degradation in removing oil from

the Amoco Cadiz and IXTOC I spills. Environ Int. 1981;5(1):33–8.

30.   Schwing PT, Romero IC, Larson RA, O'Malley BJ, Fridrik EE, Goddard EA, et al. Sediment Core

Extrusion Method at Millimeter Resolution Using a Calibrated, Threaded-rod. Journal of

Visualized Experiments : JoVE. 2016.

31.   Soto LA, Botello A V., Licea-Durán S, Lizárraga-Partida ML, Yáñez-Arancibia A. The

environmental legacy of the Ixtoc-I oil spill in Campeche Sound, southwestern Gulf of Mexico.

Front Mar Sci [Internet]. 2014;1(November):1–9. Available from:

http://journal.frontiersin.org/article/10.3389/fmars.2014.00057/abstract

32.   Patton JS, Rigler MW, Boehm PD, Fiest DL. Ixtoc 1 oil spill: flaking of surface mousse in the Gulf of

Mexico. Nature [Internet]. 1981 Mar 19;290:235. Available from:

http://dx.doi.org/10.1038/290235a0

33.   Schlitzer R. Ocean data view [Internet]. 2012. Available from: http://odv.awi.de/

34.   Caporaso JG, Lauber CL, Walters W a, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-

throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J

[Internet]. 2012 Aug [cited 2013 May 21];6(8):1621–4. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3400413&tool=pmcentrez&rendert

ype=abstract

35.   Green SJ, Venkatramanan R, Naqib A. Deconstructing the polymerase chain reaction:

Understanding and correcting bias associated with primer degeneracies and primer-template

mismatches. PLoS One. 2015;10(5):1–21.

36.   Rognes T, Flouri T, Nichols B, Quince C, Mahe F. VSEARCH : a versatile open source tool for

metagenomics (# 13057). PeerJ Prepr. 2016;1–30.

37.   Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M. Swarm: robust and fast clustering method

for amplicon-based studies. PeerJ [Internet]. 2014;2:e593. Available from:

http://dx.doi.org/10.7717/peerj.593

38.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene

database project: improved data processing and web-based tools. Nucleic Acids Res. 2012

Nov;41(D1):D590–6.

39.   Paulson JN, Stine OC, Bravo HC, Pop M. Differential abundance analysis for microbial marker-

gene surveys. Nat Methods [Internet]. 2013 Dec [cited 2014 Apr 28];10(12):1200–2. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/24076764

40.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, et al. The Vegan Package.

Community Ecol Packag. 2007 Sep;10:631–7.

41.   R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,

Austria; 2014. Available from: http://www.r-project.org/

42.   Wickham H. ggplot2: elegant graphics for data analysis [Internet]. Springer New York; 2009.

Available from: http://had.co.nz/ggplot2/book

43.   Liaw A, Wiener M. Classification and Regression by randomForest. R News [Internet].

2002;2(3):18–22. Available from: http://cran.r-project.org/doc/Rnews/

44.   Divins DL, Metzger D. NGDC coastal relief model. Natl Geophys Data Center, Natl Ocean Atmos

Adm US Dep Commer www ngdc noaa gov/mgg/(last accessed December 2007). 2008;

45.   Jacob M, Soltwedel T, Boetius A, Ramette A. Biogeography of Deep-Sea Benthic Bacteria at

Regional Scale (LTER HAUSGARTEN, Fram Strait, Arctic). PLoS One [Internet]. 2013;8(9):e72779.

Available from: http://dx.plos.org/10.1371/journal.pone.0072779

46.   Schauer R, Bienhold C, Ramette A, Harder J. Bacterial diversity and biogeography in deep-sea

surface sediments of the South Atlantic Ocean. ISME J [Internet]. 2010;4(2):159–70. Available

from: http://www.nature.com/doifinder/10.1038/ismej.2009.106

47.   Learman DR, Henson MW, Thrash JC, Temperton B, Brannock PM, Santos SR, et al.

547   Biogeochemical and microbial variation across 5500 km of Antarctic surface sediment implicates

548   organic matter as a driver of benthic community structure. Front Microbiol. 2016;7(MAR):1–11.

549   48.   Wenzhöfer F, Glud RN. Benthic carbon mineralization in the Atlantic : a synthesis based on in situ

550   data from the last decade. Deep Res I. 2002;49:1255–79.

551   49.   Glud RN, Wenzhöfer F, Middelboe M, Oguri K, Turnewitsch R, Canfield DE, et al. High rates of

552   microbial carbon turnover in sediments in the deepest oceanic trench on Earth. Nat Geosci

553   [Internet]. 2013;6(4):284–8. Available from:

554   http://dx.doi.org/10.1038/ngeo1773%5Cnpapers2://publication/doi/10.1038/ngeo1773

555   50.   Honjo S, Manganini SJ, Krishfield RA, Francois R. Particulate organic carbon fluxes to the ocean

556   interior and factors controlling the biological pump: A synthesis of global sediment trap programs

557   since 1983. Prog Oceanogr. 2008;76(3):217–85.

558   51.   Moss JA, McCurry C, Schwing P, Jeffrey WH, Romero IC, Hollander DJ, et al. Molecular

559   characterization of benthic foraminifera communities from the Northeastern Gulf of Mexico shelf

560   and slope following the Deepwater Horizon event. Deep Res Part I Oceanogr Res Pap [Internet].

561   2016;115:1–9. Available from: http://dx.doi.org/10.1016/j.dsr.2016.04.010

562   52.   Goñi MA, Ruttenberg KC, Eglinton TI. Sources and contribution of terrigenous organic carbon to

563   surface sediments in the Gulf of mexico. Lett To Nat. 1997;389(SEPTEMBER):275–9.

564   53.   Dunn DE. Trends in nutrient inflows to the Gulf of Mexico from streams draining the

565   conterminous United States, 1972-93. 1996.

566   54.   Meybeck M, Ragu A. GEMS-GLORI world river discharge database [Internet]. PANGAEA; 2012.

567   Available from: https://doi.org/10.1594/PANGAEA.804574

568   55.   Turner RE, Rabalais NN, Justic D. Gulf of Mexico hypoxia: aleternate states and a legacy. Environ

569   Sci Technol. 2008;42(7):2323–7.

570   56.   Bianchi TS, DiMarco SF, Cowan JH, Hetland RD, Chapman P, Day JW, et al. The science of hypoxia

571   in the Northern Gulf of Mexico: a review. Sci Total Environ. 2010;408(7):1471–84.

572   57.   Brooks GR, Larson RA, Schwing PT, Romero I, Moore C, Reichart GJ, et al. Sedimentation pulse in

573   the NE Gulf of Mexico following the 2010 DWH blowout. PLoS One. 2015;10(7):1–24.

574   58.   Passow U, Ziervogel K, Asper V, Diercks a. Marine snow formation in the aftermath of the

575   Deepwater Horizon oil spill in the Gulf of Mexico. Environ Res Lett [Internet]. 2012 Sep 1 [cited

576   2014 Jul 23];7(3):035301. Available from: http://stacks.iop.org/1748-

577   9326/7/i=3/a=035301?key=crossref.eb1f8af2dbf39b3445e4d441894a2922

578   59.   Day JW, Day RH, Barreiro MT, Ley-Lou F, Madden CJ. Primary production in the Laguna de

579   Terminos, a tropical estuary in the southern Gulf of Mexico. Oceanol Acta, Spec issue. 1982;

580   60.   Chen X, Andersen TJ, Morono Y, Inagaki F, Jørgensen BB, Lever MA. Bioturbation as a key driver

581   behind the dominance of Bacteria over Archaea in near-surface sediment. Sci Rep [Internet].

582   2017;7(1):2400. Available from: https://doi.org/10.1038/s41598-017-02295-x

583   61.   Durbin AM, Teske A. Sediment-associated microdiversity within the Marine Group I

584   Crenarchaeota. Environ Microbiol Rep [Internet]. 2010 Oct [cited 2013 Jun 26];2(5):693–703.

585   Available from: http://www.ncbi.nlm.nih.gov/pubmed/23766257

586   62.   Pester M, Schleper C, Wagner M. The Thaumarchaeota: An emerging view of their phylogeny and

587   ecophysiology. Curr Opin Microbiol. 2011;14(3):300–6.

588   63.   Gillan DC, Danis B. The archaebacterial communities in Antarctic bathypelagic sediments. Deep

589   Sea Res Part II Top Stud Oceanogr. 2007;54(16):1682–90.

590   64.   Durbin AM, Teske A. Microbial diversity and stratification of South Pacific abyssal marine

591   sediments. Environ Microbiol [Internet]. 2011 Dec [cited 2014 May 31];13(12):3219–34. Available

592   from: http://www.ncbi.nlm.nih.gov/pubmed/21895908

593   65.   Durbin AM, Teske A. Archaea in organic-lean and organic-rich marine subsurface sediments: an

594   environmental gradient reflected in distinct phylogenetic lineages. Front Microbiol [Internet].

595   2012 Jan [cited 2013 Jun 10];3(May):168. Available from:

596   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3364523&tool=pmcentrez&rendert

597   ype=abstract

598   66.   Alonso-Saez L, Waller a. S, Mende DR, Bakker K, Farnelid H, Yager PL, et al. Role for urea in

599   nitrification by polar marine Archaea. Proc Natl Acad Sci. 2012;109(44):17989–94.

600   67.   Oshiki M, Satoh H, Okabe S. Ecology and physiology of anaerobic ammonium oxidizing bacteria.

601   Environ Microbiol. 2016;18(9):2784–96.

602   68.   Penton CR, Devol AH, Tiedje JM. Molecular evidence for the broad distribution of anaerobic

603   ammonium-oxidizing bacteria in freshwater and marine sediments. Appl Environ Microbiol.

604   2006;72(10):6829–32.

605   69.   Canion A, Overholt WA, Kostka JE, Huettel M, Lavik G, Kuypers MMM. Temperature response of

606   denitrification and anaerobic ammonium oxidation rates and microbial community structure in

607   Arctic fjord sediments. Environ Microbiol [Internet]. 2014;16(10):3331–44. Available from:

608   http://dx.doi.org/10.1111/1462-2920.12593

609   70.   Lipsewers YA, Bale NJ, Hopmans EC, Schouten S, Sinninghe Damst?? JS, Villanueva L. Seasonality

610   and depth distribution of the abundance and activity of ammonia oxidizing microorganisms in

611   marine coastal sediments (North Sea). Front Microbiol. 2014;5(SEP):1–12.

71.   Glud RN, Thamdrup B, Stahl H, Wenzhoefer F, Glud A, Nomaki H, et al. Nitrogen cycling in a deep

ocean margin sediment (Sagami Bay, Japan). Limnol Oceanogr. 2009;54(3):723–34.

72.   Corinaldesi C. New perspectives in benthic deep-sea microbial ecology. Front Mar Sci [Internet].

2015;2:17. Available from: https://www.frontiersin.org/article/10.3389/fmars.2015.00017

73.   Swan BK, Martinez-Garcia M, Preston CM, Sczyrba A, Woyke T, Lamy D, et al. Potential for

chemolithoautotrophy among ubiquitous bacteria lineages in the dark ocean. Science (80-).

2011;333(6047):1296–300.

74.   Bradter U, Kunin WE, Altringham JD, Thom TJ, Benton TG. Identifying appropriate spatial scales of

predictors in species distribution models with the random forest algorithm. Methods Ecol Evol.

2013;4(2):167–74.

75.   Cutler DR, Edwards TC, Beard KH, Cutler A, Hess KT, Gibson J, et al. Random forests for

classification in ecology. Ecology. 2007;88(11):2783–92.

76.   Oppel S, Huettmann F. Using a Random Forest Model and Public Data to Predict the Distribution

of Prey for Marine Wildlife Management. In: Cushman SA, Huettmann F, editors. Spatial

Complexity, Informatics, and Wildlife Conservation [Internet]. Tokyo: Springer Japan; 2010. p.

151–63. Available from: https://doi.org/10.1007/978-4-431-87771-4_8

77.   Smith A, Sterba-Boatwright B, Mott J. Novel application of a statistical technique, Random

Forests, in a bacterial source tracking study. Water Res [Internet]. 2010;44(14):4067–76.

Available from: http://dx.doi.org/10.1016/j.watres.2010.05.019

78.   Lima-Mendez G, Faust K, Henry N, Decelle J, Colin S, Carcillo F, et al. Determinants of community

structure in the grobal plankton interactome. Science (80-). 2015;348(6237):1262073.

79.   Hannigan GD, Duhaime MB, Koutra D, Schloss PD. Biogeography and environmental conditions

shape bacteriophage-bacteria networks across the human microbiome. PLOS Comput Biol

[Internet]. 2018 Apr 18;14(4):e1006099. Available from:

https://doi.org/10.1371/journal.pcbi.1006099

80.   Romero IC, Schwing PT, Brooks GR, Larson RA, Hastings DW, Ellis G, et al. Hydrocarbons in deep-

sea sediments following the 2010 Deepwater Horizon blowout in the northeast Gulf of Mexico.

PLoS One. 2015;10(5):1–23.

81.   Romero IC, Toro-Farmer G, Diercks AR, Schwing P, Muller-Karger F, Murawski S, et al. Large-scale

deposition of weathered oil in the Gulf of Mexico following a deep-water oil spill. Environ Pollut

[Internet]. 2017;228:179–89. Available from: http://dx.doi.org/10.1016/j.envpol.2017.05.019

82.   Chanton JP, Cherrier J, Wilson RM, Sarkodee-Adoo J, Bosman S, Mickle A, et al. Radiocarbon

evidence that carbon from the Deepwater Horizon spill entered the planktonic food web of the

Gulf of Mexico. Environ Res Lett [Internet]. 2012 Dec 1 [cited 2014 May 8];7(4):045303. Available

from: http://stacks.iop.org/1748-

9326/7/i=4/a=045303?key=crossref.bb52d0d288c12eb2c6f2ac2b8f8facea

Figure 1. Map of the sampling sites used in this study. All yellow points are sites that were profiled for oxygen concentrations, while sites with black points are those that only have microbial community data. Sites in red are represented by both oxygen and microbial community data. All sites sampled for microbial data are labeled with the site name, which can be seen in Table S1.

Figure 2. Large-scale patterns in microbial biogeography. All plots are generated from the same
NMDS ordination of Bray-Curtis dissimilarity. The stress for each plot is 0.152. (A) Samples are
colored based on increasing water depth. The shallow water samples (< 100m) clearly separate
from the deeper water samples (> 400m). (B) Samples are colored darker with increasing sediment
depths and there is a clear and consistent downcore progression from the "top-right" of the NMDS
plot towards the bottom left in every core sampled. (C) Samples are colored based on collection
date. (D) Cores were collected from the Southern Gulf of Mexico (colored in green) and the
Northern Gulf of Mexico (colored in pink).

Figure 3. Oyxgen profiles collected throughout the Gulf of Mexico. (A) Interpolated plot showing predicted oxygen penetration depths (mm) across our study site. (B) Oxygen penetration plotted against water column depth.

Figure 4. Differences in microbial community and oxygen profiles between the Northern and Southern Gulf of Mexico. The SGoM sites sampled have deeper depths of oxygen penetration and correspond to a community composition more similar to shallower sites in the NGoM. (A) NMDS of Bray-Curtis dissimilarity values for cores collected in 2014 (NGoM, pink) or 2015 (SGoM, green) from 500m – 2300m water depths. Down core progression is indicated by the connecting lines from surficial sediments on the left towards deeper samples on the right. (B) Corresponding oxygen profiles to all cores presented in A. (C, D) Same as plots A & B except only displaying sites collected between 1000-1300m.

Figure 5. Core profiles for the 12 most abundant classes detected from deep water transects. (A-
C) Representative profiles from the Northern Gulf of Mexico from the DSH core transect going
offshore towards increasing water depth. (D-F) Representative profiles from the Southern Gulf of
Mexico along a westward transect from the Ixtoc wellhead, also organized from shallowest to
deepest site in the transect. In all plots, blue lines represent the percent oxygen concentration
relative to the initial overlying water oxygen concentration.

 Figure 6 Core profiles representing deep water sites in the Northern Gulf of Mexico showing
 population (OTU) level distributions with increasing sediment and water depths. Core profiles are
 taken from the DSH transect and are arranged in order of increasing water depth. See Figure S4
 for similar trends within the Southern Gulf of Mexico.

Figure 7. Meta-study revealing localized impact and subsequent recovery. Data generated by Mason et al., 2014 from surficial samples (red, green) was re-analyzed with surficial samples from this study (blue). (A) Geographic map plotting each sampling location. Samples outlined in red

from Mason et al., exceeded EPA benchmarks while samples outlined in green were below EPA
benchmarks. The fill color for each point represents distance from well head, going from red to
green with increasing distance. (B) NMDS of Bray-Curtis dissimilarity values, plotted using the
same color scheme as in A. The wellhead samples (DWH01) are indicated by yellow outlines and
were collected in 2012 and 2013. (C) Deviations of every sample collected, including the full
sediment core for samples from this study, from the OTU distribution models. Samples are colored
by the sampling date or by oil contamination state. High RMSE (root mean squared error) values
represent samples that strongly differ from OTU relative abundance predictions.