1  **Transcriptomic characterization of 20 organs and tissues from mouse at single cell resolution**
2  **creates a *Tabula Muris***
3
4  The *Tabula Muris* Consortium
5

**We have created a compendium of single cell transcriptome data from the model**
**organism *Mus musculus* comprising nearly 100,000 cells from 20 organs and tissues.**
**These data reveal many new aspects of cell biology, including gene expression in**
**poorly characterized cell populations and the characterization of new populations of**
**cells in many tissues. Furthermore, they allow for direct and controlled comparison**
**of gene expression in cell types shared between tissues, such as immune cells from**
**distinct anatomical locations. Two distinct technical approaches were used for most**
**organs: one approach, microfluidic droplet-based 3'-end counting[1,2], enabled the**
**survey of thousands of cells at relatively low coverage, while the other, FACS-based**
**full length transcript analysis[3], enabled characterization of cell types with high**
**sensitivity and coverage.  The cumulative resource provides the foundation for a**
**comprehensive resource of transcriptomic cell biology.**

18
The cell is a fundamental unit of structure and function in biology, and multicellular
organisms have evolved a wide variety of different cell types with specialized roles.
Although cell types have historically been characterized on the basis of morphology and
phenotype, the development of molecular methods has enabled ever more precise
definition of their properties, typically by measuring protein or mRNA expression
patterns[4].  Technological advances have enabled increasingly greater degrees of
multiplexing of these measurements[5,6], and it is now possible to use highly parallel
sequencing to enumerate nearly every mRNA molecule in a given single cell[3,7]. This
approach has provided many novel insights into cell biology and the composition of
organs and tissues from a variety of organisms[8–17].  However, while these reports provide
valuable characterization of individual organs, it is challenging to compare data taken
with varying experimental techniques in independent labs from different animals. It
therefore remains an open question whether data from individual organs can be
synthesized and used as a more general resource for biology.
33
Here we report a compendium of cell types from the mouse *Mus musculus*. The
compendium is comprised of single cell transcriptome sequence data from 97,029 cells
isolated from 20 organs and tissues (Fig. 1). These organs were collected from 3 female
and 4 male, C57BL/6 NIA, 3 month old mice (10-15 weeks), whose developmental age is
roughly analogous to humans at 20 years of age. We analyzed multiple organs and tissues
from the same animal, thereby generating a data set controlled for age, environment and
epigenetic effects.  This enables the direct comparison of cell type composition between
organs as well as comparison of shared cell types across the entire organism. All data,
protocols, and analysis scripts from the compendium are shared as a public resource.
Gene counts and metadata from all single cells are accessible via Digital Object Identifier
(DOI) at 10.6084/m9.figshare.5715040 for the FACS data and
10.6084/m9.figshare.5715025 for the microfluidic droplet data. While these data are by
no means a complete representation of all mouse organs and cell types, they provide a

first draft attempt to create an organism-wide representation of cellular diversity and a
comparative framework for future studies using the large variety of murine disease
models.
50
We developed a specific procedure to collect 20 organs and tissues from the same mouse
(see Methods).  Briefly, each mouse was anesthetized with 2.5% v/v Avertin, followed by
transcardial perfusion with 20 ml PBS. Aorta, bladder, bone marrow, brain (cerebellum,
cortex, hippocampus, striatum), colon, diaphragm, fat (brown, gonadal, mesenteric,
subcutaneous), heart, kidney, liver, lung, mammary gland, pancreas, skin, skeletal
muscle, spleen, thymus, tongue, and trachea were immediately dissected and processed
into single cell suspensions (see Extended Data). Organs were dissociated into single
cells and either sorted by FACS into 384-well plates (with the exception of liver
hepatocytes that were sorted into 96 well plates and heart derived cardiomyocytes that
were hand-picked into 96-well plates), and in many cases also loaded into a microfluidic
droplet emulsion generating device for cell isolation followed by transcriptome capture
(Fig. 1a).  Once prepared into sequencing libraries, transcriptomes were sequenced to an
average depth of 685,500 reads per cell for the plate data and 7,709 unique molecular
identifiers (UMI) per cell for the microfluidic emulsion data.  After quality control
filtering, 44,879 FACS sorted cells and 55,656 microfluidic droplet processed cells were
retained for further analysis.  A comparison of the two methods showed differences for
each organ in the number of cells analyzed (Fig. 1b,c), reads per cell (Fig. 1d,f) and genes
per cell (Fig. 1e,g).
69
Using two distinct measurement approaches on the same samples yielded several
insights.  We compared the methods to understand their relative strengths and
weaknesses.  Importantly, our results here show that such comparisons can be misleading
when performed on a single organ type, as there is substantial variation in performance
across organs (Supp. Fig. 1). For example, the lung, trachea and thymus showed nearly
twice as many genes detected per cell in the FACS data as compared to the microfluidic
data, whereas heart, kidney and marrow show comparable numbers of expressed genes
detected by both methods (Supp. Fig. 1-3).  Number of genes detected is a fairly crude
approach to sensitivity and one must also consider other metrics such as dynamic range
for gene expression.  The FACS based approach has much higher dynamic range per
gene than the microfluidic droplets (Supp. Fig. 4) and enables the analysis of full length
transcripts. While the FACS approach generally resulted in higher sensitivity and
dynamic range, it also is more laborious and time consuming to perform.  The
microfluidic droplet approach is faster and enables analysis of larger numbers of cells,
albeit at generally lower sensitivity and dynamic range, and with reads only from the 3'
end of the transcript.  We also expect this study to facilitate the development of
computational tools to compare sequencing methods, and our data set will provide a
training and validation set for such algorithms.  The creation of these tools will crucial for
comparing independent data generated by labs around the world as various tissue atlas
studies begin generating results.
90
We performed unbiased graph-based clustering of the pooled set of transcriptomes across
all organs, and visualized them using tSNE (Fig. 2a). The majority of clusters contain

cells from only one organ, but a substantial minority contained cells from many organs.
To further dissect these clusters, we separately performed dimensional reduction and
clustering on cellular transcriptomes from each individual organ (Fig. 2b).  The resulting
clusters were manually annotated using *a priori* biological knowledge about cell-type
specific gene expression, which confirmed the presence of many specific cell types
within organs (Fig. 3). Many of these cell types have not previously been obtained in
pure populations and our data provide a wealth of new information on their characteristic
gene expression profiles. Initial annotation of each organ and tissue can be found in the
extended data, and a detailed discussion each cell types on an organ by organ basis can be
found in the supplement.  Some unexpected discoveries include: 1) novel immune cell
types in the lung, 2) a putative neuroendocrine cell population in the trachea, 3) new
differentiated keratinocyte cell types of the tongue, 4) strong sexual dimorphism in
hepatocytes with differential gene expression of male pheromones, 5) the suggestions of
new roles for genes such as *Neurog3*, *Hhex*, and *Prss53* in the adult pancreas, 6) sexual
dimorphism in epithelial and mesenchymal cells of the bladder, and 7) a novel cell
population expressing *Chodl* in skeletal muscle.
109
In order to better understand the relationships between cell types across organs, we
mapped the biologically informed annotations of organ-specific cell types onto the
clusters resulting from the unbiased analysis of the complete set of all cell transcriptomes.
This analysis revealed that the clusters with cells from many organs generally represent
shared cell types common to those organs (Figure 3).  For example, B cells from fat, limb
muscle, diaphragm, lung, spleen and marrow cluster together, as do T cells from spleen,
marrow, lung, limb muscle, fat and thymus.  Endothelial cells from fat, heart and lung
cluster together, but form a distinct grouping from endothelial cells from mammary,
kidney, trachea, limb muscle, aorta, diaphragm and pancreas. There are two clusters
containing myeloid cells from limb muscle, brain, diaphragm, aorta, pancreas, kidney,
trachea, heart, liver, fat and marrow. Lastly, a cluster containing mesenchymal stem cells
from fat, diaphragm, and limb muscle suggests these cells are share similar
transcriptomes with stromal cells in mammary, trachea, and lung. These findings show
that gene-specific manual annotation of cell types is consistent with unbiased clustering
based on whole transcriptome profiles, and that cell type identity is strong enough across
tissues to enable their unbiased identification at the whole transcriptome level.
126
To further investigate a common cell type across organs, we pooled all T-cells and
performed an unbiased clustering analysis, revealing 5 clusters (Figure 4) grouped into
two distinct sub-groups: clusters 0,4 and clusters 1,2,3.  Cluster 0 comprises cells from
the Thymus that are undergoing VDJ recombination characterized by the expression of
RAG (*Rag1* and *Rag2*) and TdT (*Dntt*), and includes uncommitted double positive T-
cells (*Cd4* and *Cd8a* positive).  Cluster 4 contains proliferating T-cells, predominantly
from the thymus, and we hypothesize that these are pre-T cells that are expanding after
having completed VDJ recombination. In contrast, Clusters 1,2,3 contain mature T-cells.
The cells in Cluster 3 are also predominantly from the thymus and show high *Cd5*
expression, suggesting that they are undergoing positive selection. Cluster 2 cells are
mostly peripheral cells and are most likely activated T-cells expressing the high affinity
IL2 receptor (*Il2ra* and *Il2rb)*. Interestingly, they also express MHC type II genes (*H2-Aa*

and *H2-Ab1*). Finally, Cluster 1 also represents mature T-cells, but primarily from the
spleen.
141
A key challenge for many single cell studies to date is understanding the potential
changes to the transcriptome caused by handling, dissociation and other experimental
manipulation.  A previous study in limb muscle showed that quiescent satellite cells tend
to become activated by dissociation and consequently express immediate early genes
among other genes.[18] We found that expression of these dissociation-related markers was
also clearly observed in our limb muscle data, as well as in mammary and bladder (Supp.
Fig. 6), but that many organs showed little evidence of cellular activation with this panel.
Therefore the dissociation-related activation markers found in limb muscle are not
universal across all organs.  This is not to say that other tissues lack dissociation-related
gene expression changes, but that some of the genes involved are specific to a given
tissue.  It is important to note that the presence of such gene expression changes does not
prevent the identification of cell type or the comparison of cell types across organs.
154
To understand the extent to which transcription factor (TF) expression determines cell
type, we performed a correlation analysis of TF expression[14] across the entire data set for
TFs that were significantly enriched ($p < 10^{-5}$, log10 fold change $> 0.3$) in at least one
unique combination of cell type and tissue (e.g., endothelial cells from liver or basal cell
of the bladder). (Fig 5A). The off-diagonal elements of the correlation matrix indicate
groups of transcription factors with correlated cell type expression.  We measured co-
expression of these groups of transcription factors and discovered that typically only one
of a few cell types used any given combination of two to four TFs.  (Supp. Fig. 7) The
largest cluster of correlated genes enriched for a broad swath of immune cells across most
tissues.  Similarly, another cluster of genes enriched for endothelial cells. We also
observed groups of TFs specific to several organs; for example within colon, goblet cells
were highly enriched for a cluster of TFs that are either known or potential new targets
for specification of goblet cells (Fig 5A, Supplementary Figure 7), and we also saw
clusters specific to muscle cells, pancreatic cells, neurons, microglia and astrocytes.
Another group of genes showed enrichment in several epithelial subtypes of tongue and
bladder.
171
172
We further analyzed tissue-specific TFs within the endothelial, epithelial and stromal cell
types found in several different organs (Fig 5B-D). Within epithelial cells, we found
genes grouped by tissue: tongue (Pax9), bladder (Ets1), skin keratinocyte stem cells
(Lhx2), and lung (Aebp1) (Fig 5B). Pax9 is enriched in tongue basal cells and is known
to be necessary for formation of filiform papillae. Foxq1 is enriched in bladder basal
cells, but its role in bladder basal cell differentiation is heretofore unexplored. Within
fibroblast/stromal cells, we found TFs that separated heart from kidney fibroblasts, and
specified the stromal cells of lung, trachea, or mammary gland (Fig 5C). Within
endothelial cells, we found less strong signatures for tissue specificity; however, liver,
brain, kidney, and lung-specific groups of TFs were distinguishable (Fig 5D).  These
results illustrate how single cell data taken across many organs and tissues can be used in

combination to identify which transcriptional regulatory programs are specific to cell
types of interest.
186
In conclusion, we have created a compendium of single-cell transcriptional
measurements across 20 organs and tissues of the mouse. This *Tabula Muris*, or "Mouse
Atlas", has many uses, including the discovery of new putative cell types, the discovery
of novel gene expression in known cell types, and the ability to compare cell types across
organs.  It will also serve as a reference of healthy young adult organs which can be used
as a baseline for current and future mouse models of disease. While it is not an
exhaustive characterization of all organs of the mouse, it does provide a rich data set of
the most highly studied tissues in biology. The *Tabula Muris* provides a framework and
description of many of the most populous and important cell populations within the
mouse, and represents a foundation for future studies across a multitude of diverse
physiological disciplines.
198
199

200 **Acknowledgements**
201
This work was funded by the Chan Zuckerberg Biohub, the National Institute on Aging
(DP1AG053015, T.W.-C), the Veterans Administration (T.W.-C.), and the Stanford
Diabetes Research Center (NIH DK116074, S.K.K.).  We thank Sony Biotechnology for
making an SH800S instrument available for this project.  Some cell sorting/flow
cytometry analysis for this project was done on a Sony SH800S instrument in the
Stanford Shared FACS Facility. Some fluorescence activated cell sorting (FACS) was
done with instruments in the VA Flow Cytometry Core, which is supported by the US
Department of Veterans Affairs (VA), Palo Alto Veterans Institute for Research
(PAVIR), and the National Institutes of Health (NIH).

**The *Tabula Muris* Consortium:**
213
214 **Overall Coordination:** Nicholas Schaum[b], Jim Karkanias[r], Norma F Neff[r], Andrew P.
215 May[r], Stephen R. Quake[o,r]\*, Tony Wyss-Coray[a,c,d]\*, and Spyros Darmanis[r]\*
216
217 \* to whom correspondence should be addressed: quake@stanford.edu, twc@stanford.edu,
218 spyros.darmanis@czbiohub.org
219
**Tissue collection and processing:** Ankit S. Baghel[b], Isaac Bakerman[b,h,i], Ishita Bansai[r],
Daniela Berdnik[a], Biter Bilen[a], Douglas Brownfield[l], Corey Cain[s], Michelle B. Chen[o],
Steven Chen[r] , Min Cho[r], Giana Cirolia[r], Stephanie G. Conley[b], Spyros Darmanis[r], Aaron
Demers[r], Kubilay Demir[b,n], Tessa Divita[r], Haley du Bois[a], Laughing Bear Torrez
Dulgeroff[b], Hamid Ebadi[r], Hernan Espinoza[l], Matt Fish[b,m,n], Benson M. George[b], Astrid
Gillich[l], Foad Green[r], Geraldine Genetiano[r], Xueying Gu[m], Gunsagar S. Gulati[b], Yan
Hang[m], Shayan Hosseinzadeh[r], Tal Iram[a], Taichi Isobe[b], Feather Ives[r], Robert Jones[o],
Kevin S. Kao[b], Guruswamy Karnam[q], Aaron M. Kershner[b], Bernhard Kiss[b,x], William
Kong[b], Jonathan Lam[m], Davis P. Lee[d], Song E. Lee[a], Benoit Lehallier[a], Guang Li[j],
Qingyun Li[w], Ling Liu[a,c], Annie Lo[r], Wan-Jin Lu[b,l], Anoop Manjunath[b], Andrew P. May[r],

Kaia L. May[r], Oliver L. May[r], Ashley Maynard[r], Marina McKay[r], Marco Mignardi[o],
Dullei Min[k], Antoine de Morree[a], Ahmad N. Nabhan[l], Norma F Neff[r], Katharine M. Ng[o],
Joseph Noh[b], Rasika Patkar[q], Weng Chuan Peng[m], Lolita Penland[r], Robert Puccinelli[r],
Eric J. Rulifson[m], Nicholas Schaum[b], Shaheen S. Sikandar[b], Rahul Sinha[b,e,f,g], Rene V
Sit[r], Krzysztof Szade[b,y], Weilun Tan[r], Cristina Tato[r], Krissie Tellez[m], Kyle J. Travaglini[l],
Carolina Tropini[p], Lucas Waldburger[r], Linda J. van Weele[b], Michael N. Woszczyna[a], Jinyi
Xiang[b], Soso Xue[o], Justin Youngyunpipatkul[r], Hanadie Yousef[a], Fabio Zanini[o], Macy E.
Zardeneta[d], Lu Zhou[w]
238
239  **Library preparation and sequencing:** Ishita Bansai[r], Steven Chen[r], Min Cho[r], Giana
240  Cirolia[r], Spyros Darmanis[r], Aaron Demers[r], Tessa Divita[r], Hamid Ebadi[r], Geraldine
241  Genetiano[r], Foad Green[r], Shayan Hosseinzadeh[r], Feather Ives[r], Annie Lo[r], Andrew P.
242  May[r], Ashley Maynard[r], Marina McKay[r], Norma F Neff[r], Lolita Penland[r], Rene V Sit[r],
243  Weilun Tan[r], Lucas Waldburger[r], Justin Youngyunpipatkul[r]
244
245  **Computational Data Analysis:** Joshua Batson[r], Olga Botvinnik[r], Paola Castro[r], Spyros
246  Darmanis[r], Joseph L. DeRisi[r,t], Jim Karkanias[r], Geoffrey M. Stanley[o], James T. Webber[r]
247
**Cell Type Annotation:** Ankit S. Baghel[b], Isaac Bakerman[b,h,i], Ben A. Barres[w], Joshua
Batson[r], Phil Beachy[b,l,m,n], Biter Bilen[a], Olga Botvinnik[r], Douglas Brownfield[l], Charles
K.F. Chan[v], Michelle B. Chen[o], Michael F. Clarke[b], Spyros Darmanis[r], Kubilay Demir[b,n],
Hamid Ebadi[r], Hernan Espinoza[l], Matt Fish[b,m,n], Benson M. George[b], Astrid Gillich[l],
Xueying Gu[m], Gunsagar S. Gulati[b], Yan Hang[m], Kerwyn Casey Huang[o,p,r], Tal Iram[a],
Taichi Isobe[b], Guruswamy Karnam[q], Aaron M. Kershner[b], Seung K. Kim[m,u], Bernhard M.
Kiss[b,x], William Kong[b], Mark A. Krasnow[l,n], Christin S. Kuo[k,l,n], Jonathan Lam[m], Benoit
Lehallier[a], Guang Li[j], Qingyun Li[w], Ling Liu[a,c], Wan-Jin Lu[b,l], Dullei Min[k], Antoine de
Morree[a], Ahmad N. Nabhan[l], Katharine M. Ng[o], Patricia K. Nguyen[b,h,i,j], Roel Nusse[b,m,n],
Rasika Patkar[q], Weng Chuan Peng[m], Lolita Penland[r],Thomas A. Rando[a,c,d], Eric J.
Rulifson[m], Nicholas Schaum[b], Shaheen S. Sikandar[b], Rahul Sinha[b,e,f,g], Krzysztof
Szade[b,z], Krissie Tellez[m], Kyle J. Travaglini[l],Carolina Tropini[p], Linda J. van Weele[b],
Bruce M. Wang[q], Kenneth Weinberg[k], Irving L. Weissman[b,e,f,g], Michael N. Woszczyna[a],
Sean Wu[b,h,j], Tony Wyss-Coray[a,c,d] , Jinyi Xiang[b], Hanadie Yousef[a], Macy E. Zardeneta[d],
Lu Zhou[w]
263
264  **Writing Group:**  Joshua Batson[r], Olga Botvinnik[r], Steven Chen[r], Spyros Darmanis[r],
265  Foad Green[r], Andrew P. May[r], Ashley Maynard[r], Stephen R. Quake[o,r], Nicholas Schaum[r],
266  James T. Webber[r], Tony Wyss-Coray[a,c,d]
267
268  **Principal Investigators:** Ben A. Barres[w], Phil Beachy[b,l,m,n], Charles Chan[b], Michael F.
269  Clarke[b], Spyros Darmanis[r], Kerwyn Casey Huang[o,p,r], Jim Karkanias[r], Seung K. Kim[m,u],
270  Mark A. Krasnow[l,n], Christin S. Kuo[k,l,n], Andrew P. May[r], Norma Neff[r], Roel Nusse[b,m,n],
271  Patricia K. Nguyen[b,h,i,j], Thomas A. Rando[a,c,d], Bruce M. Wang[q], Kenneth Weinberg[k],
272  Irving L. Weissman[b,e,f,g], Sean Wu[b,h,j], Stephen R. Quake[o,r], Tony Wyss-Coray[a,c,d]
273
[a]Department of Neurology and Neurological Sciences, Stanford University School of
Medicine, Stanford, California, USA

[b]Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School
of Medicine, Stanford, California, USA
[c]Paul F. Glenn Center for the Biology of Aging, Stanford University School of Medicine,
Stanford, California, USA
[d]Center for Tissue Regeneration, Repair, and Restoration, V.A. Palo Alto Healthcare
System, Palo Alto, California, USA
[e]Department of Pathology, Stanford University School of Medicine, Stanford, California,
USA
[f]Ludwig Center for Cancer Stem Cell Research and Medicine, Stanford University
School of Medicine, Stanford, California, USA
[g]Stanford Cancer Institute, Stanford University School of Medicine, Stanford, California,
USA.
[h]Stanford Cardiovascular Institute, Stanford University School of Medicine, Stanford,
California, USA
[i]Department of Medicine, Division of Cardiology, Stanford University School of
Medicine, Stanford, California, USA
[j]Department of Medicine, Division of Cardiovascular Medicine, Stanford University,
Stanford, California, USA
[k]Department of Pediatrics, Stanford University school of Medicine, Stanford, California,
USA
[l]Department of Biochemistry, Stanford University School of Medicine, Stanford,
California, USA
[m]Department of Developmental Biology, Stanford University School of Medicine,
Stanford, California, USA
[n]Howard Hughes Medical Institute, USA
[o]Department of Bioengineering, Stanford University, Stanford, California, USA
[p]Department of Microbiology & Immunology, Stanford University School of Medicine,
Stanford, California, USA
[q] Department of Medicine and Liver Center, University of California San Francisco, San
Francisco, California, USA
[r]Chan Zuckerburg Biohub, San Francisco, California, USA
[s]Flow Cytometry Core, V.A. Palo Alto Healthcare System, Palo Alto, California, USA
[t] Department of Biochemistry and Biophysics, University of California San Francisco,
San Francisco, California USA
[u] Department of Medicine and Stanford Diabetes Research Center, Stanford University,
Stanford, California USA
[v] Department of Surgery, Division of Plastic and Reconstructive Surgery, Stanford
University, Stanford, California USA
[w] Department of Neurobiology, Stanford University School of Medicine, Stanford, CA
USA
[x] Department of Urology, Stanford University School of Medicine, Stanford, California,
USA
[y] Department of Medical Biotechnology, Faculty of Biophysics, Biochemistry and
Biotechnology , Jagiellonian University, Poland
320

**Methods**

**Mice and Tissue Collection**
Four 10-15 week old male and four virgin female C57BL/6 mice were shipped from the National Institute on Aging colony at Charles River to the Veterinary Medical Unit (VMU) at the VA Palo Alto (VA). At both locations, mice were housed on a 12-h light/dark cycle, and provided food and water *ad libitum*. The diet at Charles River was NIH-31, and Teklad 2918 at the VA VMU. Littermates were not recorded or tracked, and mice were housed at the VA VMU for no longer than 2 weeks before euthanasia. Prior to tissue collection, mice were placed in sterile collection chambers for 15 minutes to collect fresh fecal pellets. Following anesthetization with 2.5% v/v Avertin, mice were weighed, shaved, and blood drawn via cardiac puncture before transcardial perfusion with 20 ml PBS. Mesenteric adipose tissue (MAT) was then immediately collected to avoid exposure to the liver and pancreas perfusate, which negatively impacts cell sorting. Isolating viable single cells from both pancreas and liver of the same mouse was not possible, therefore, 2 males and 2 females were used for each. Whole tissues were then dissected in the following order: colon, spleen, thymus, trachea, tongue, brain, heart, lung, kidney, gonadal adipose tissue (GAT), bladder, diaphragm, skeletal muscle (*tibialis anterior*), skin (dorsal), subcutaneous adipose tissue (SCAT, inguinal pad), mammary glands (fat pads 2, 3, and 4), brown adipose tissue (BAT, interscapular pad), aorta, and bone marrow (spine and limb bones). Following single cell dissociation as described below, cell suspensions were either used for FACS sorting of individual cells into 384-well plates, or for microfluidic droplet library preparation. All animal care and procedures were carried out in accordance with institutional guidelines approved by the VA Palo Alto Committee on Animal Research.

**Tissue dissociation and sample preparation:** Specific protocols for each tissue are described in the supplement.

**Single Cell Methods**

**Lysis plate preparation**
Lysis plates were created by dispensing 0.4 μl lysis buffer (0.5U Recombinant RNase Inhibitor (Takara Bio, 2313B), 0.0625% Triton$^{TM}$ X-100 (Sigma, 93443-100ML), 3.125 mM dNTP mix (Thermo Fisher, R0193), 3.125 μM Oligo-dT$_{30}$VN (IDT, 5'AAGCAGTGGTATCAACGCAGAGTACT$_{30}$VN-3') and 1:600,000 ERCC RNA spike-in mix (Thermo Fisher, 4456740)) into 384-well hard-shell PCR plates (Biorad HSP3901) using a Tempest or Mantis liquid handler (Formulatrix). 96 well lysis plates were also prepared with 4 μl lysis buffer. All plates were sealed with AlumaSeal CS Films (Sigma-Aldrich Z722634) or Microseal 'F' PCR plate seal (Biorad MSF 1001) and then spun down for 1 minute at 3220 xg and snap frozen on dry ice. Plates were stored at -80°C until used for sorting.

**FACS sorting**

After dissociation, single cells from each tissue were isolated in 384 or 96-well plates via
Fluorescence Activated Cell Sorting (FACS). Most tissues were sorted into 384-well
plates using SH800S (Sony) sorters. Heart and liver were sorted into 96-well plates and
cardiomyocytes were hand-picked into 96-well plates. Skeletal muscle and diaphragm
were sorted into 384-well plates on an Aria III (Becton Dickinson) sorter. Most tissues
used combinations of fluorescent antibodies to enrich for the presence of known rare cell
populations (see tissue section below), but some tissues were simply sorted into plates
after removal of dead cells, small and large debris. For all samples, an initial gate was set
to exclude small debris, platelets and cell aggregates, and most tissues included a forward
scattering gate to select for single cells. One color channel was used in most tissues to
stain and exclude dead cells and high prevalence cells that would otherwise dominate the
cell population. Color compensation was used whenever necessary.  Cells were sorted
using the highest purity setting on the sorter ("Single cell" in the case of the SH800s) for
all but the rarest cell types, for which the "Ultrapure" setting was used. Sorters were
calibrated using FACS buffer before collecting cells from any tissue and after every 8
sorted plates to ensure accurate dispensing into plate wells. For a typical sort, a tube
containing 1-3ml the pre-stained cell suspension was filtered, vortexed gently and loaded
onto the FACS machine. A small number of cells were flowed at low pressure to check
cell concentration and amount of debris. Then the pressure was adjusted, flow was
paused, the first destination plate was unsealed and loaded, and single-cell sorting started.
If a cell suspension was too concentrated, it was diluted it using FACS buffer or 1X PBS.
For some cell types (e.g. hepatocytes), 96 well plates were used when it was not possible
to sort individual cells accurately into 384 well plates. Immediately after sorting, plates
were sealed with a pre-labeled aluminium seal, centrifuged and flash frozen on dry ice.
On average, each 384-well plate took around 8 minutes to sort.
391
**cDNA synthesis and library preparation**
cDNA synthesis was performed using the Smart-seq2 protocol [1,2]. Briefly, 384-well
plates containing single-cell lysates were thawed on ice followed by first strand synthesis.
0.6 µl of reaction mix (16.7 U/µl SMARTScribe Reverse Transcriptase (Takara Bio,
639538), 1.67 U/µl Recombinant RNase Inhibitor (Takara Bio, 2313B), 1.67X First-
Strand Buffer (Takara Bio, 639538), 1.67 µM TSO (Exiqon, 5'-
AAGCAGTGGTATCAACGCAGACTACATrGrG+G-3'), 8.33 mM DTT (Bioworld,
40420001-1), 1.67 M Betaine (Sigma, B0300-5VL), and 10 mM $MgCl_2$ (Sigma, M1028-
10X1ML)) was added to each well using a Tempest liquid handler. Reverse transcription
was carried out by incubating wells on a ProFlex 2x384 thermal-cycler (Thermo Fisher)
at 42°C for 90 min and stopped by heating at 70°C for 5 min.
403
Subsequently, 1.5 µl of PCR mix (1.67X KAPA HiFi HotStart ReadyMix (Kapa
Biosystems, KK2602), 0.17 µM IS PCR primer (IDT, 5'-
AAGCAGTGGTATCAACGCAGAGT-3'), and 0.038 U/µl Lambda Exonuclease (NEB,
M0262L)) was added to each well with a Mantis liquid handler (Formulatrix), and second
strand synthesis was performed on a ProFlex 2x384 thermal-cycler by using the
following program: 1. 37°C for 30 minutes, 2. 95°C for 3 minutes, 3. 23 cycles of 98°C
for 20 seconds, 67°C for 15 seconds, and 72°C for 4 minutes, and 4. 72°C for 5 minutes.
411

The amplified product was diluted with a ratio of 1 part cDNA to 10 parts 10mM Tris-
HCl (Thermo Fisher, 15568025), and concentrations were measured with a dye-
fluorescence assay (Quant-iT dsDNA High Sensitivity kit; Thermo Fisher, Q33120) on a
SpectraMax i3x microplate reader (Molecular Devices). Sample plates were selected for
downstream processing if the mean concentration of blanks - ERCC-containing, non-cell
wells - was greater than 0 ng/μl, and, after linear regression of the values obtained from
the  Quant-iT dsDNA standard curve, the $R^2$ value was greater than 0.98.  Sample wells
were then selected if their cDNA concentrations were at least one standard deviation
greater than the mean concentration of the blanks. These wells were reformatted to a new
384-well plate at a concentration of 0.3 ng/μl and final volume of 0.4 μl using an Echo
550 acoustic liquid dispenser (Labcyte).
423
Illumina sequencing libraries were prepared as described in Darmanis et al. 2015.[13]
Briefly, tagmentation was carried out on double-stranded cDNA using the Nextera XT
Library Sample Preparation kit (Illumina, FC-131-1096). Each well was mixed with 0.8
μl Nextera tagmentation DNA buffer (Illumina) and 0.4 μl Tn5 enzyme (Illumina), then
incubated at 55°C for 10 min. The reaction was stopped by adding 0.4 μl "Neutralize
Tagment Buffer" (Illumina) and spinning at room temperature in a centrifuge at 3220xg
for 5 min. Indexing PCR reactions were performed by adding 0.4 μl of 5 μM i5 indexing
primer, 0.4 μl of 5 μM i7 indexing primer, and 1.2 μl of Nextera NPM mix (Illumina).
PCR amplification was carried out on a ProFlex 2x384 thermal cycler using the following
program: 1. 72°C for 3 minutes, 2. 95°C for 30 seconds, 3. 12 cycles of 95°C for 10
seconds, 55°C for 30 seconds, and 72°C for 1 minute, and 4. 72°C for 5 minutes.
435
**Library pooling, quality control, and sequencing**
Following library preparation, wells of each library plate were pooled using a
Mosquito  liquid handler (TTP Labtech). Pooling was followed by two purifications using
0.7x AMPure beads (Fisher, A63881). Library quality was assessed using capillary
electrophoresis on a Fragment Analyzer (AATI), and libraries were quantified by qPCR
(Kapa Biosystems, KK4923) on a CFX96 Touch Real-Time PCR Detection System
(Biorad). Plate pools were normalized to 2 nM and equal volumes from 10 or 20 plates
were mixed together to make the sequencing sample pool.  PhiX control library was
spiked in at 0.2% before sequencing.
445
446     **Sequencing libraries from 384-well and 96-well plates**
447     Libraries were sequenced on the NovaSeq 6000 Sequencing System (Illumina) using 2 x
448     100bp paired-end reads and 2 x 8bp or 2 x 12bp index reads with either a 200 or 300-
449     cycle kit (Illumina, 20012861 or 20012860).
450
451
**Microfluidic droplet single cell analysis**
Single cells were captured in droplet emulsions using GemCode Single-Cell Instrument
(10x Genomics, Pleasanton, CA, USA) and SC RNA-seq libraries were constructed as
per 10X Genomics protocol using GemCode Single-Cell 3′ Gel Bead and Library V2
Kit.  Briefly, single cell suspensions were examined using an inverted microscope and if
sample quality was deemed satisfactory, sample was diluted in PBS/2% FBS to achieve a

target concentration of 1000 cells/µl.  If cell suspensions contained cell aggregate or
debris, two additional washes in PBS/2% FBS at 300 x g for 5min at 4C were
performed.  Cell concentration was measured either with a Moxi GO II (Orflo
Technologies) or a hemocytometer.  Cells were loaded in each channel with a target
output of 5,000 cells per sample.  All reactions were performed in Biorad C1000 Touch
Thermal cycler with 96-Deep Well Reaction Module.  12 cycles were used for cDNA
amplification and sample index PCR.  Amplified cDNA and final library were evaluated
on a Fragment Analyzer using a High Sensitivity NGS Analysis Kit (Advanced
Analytical).  10x cDNA libraries were quantitated for average fragment length using a 12
or 96 capillary Fragment Analyzer (Advanced Analytical) and by qPCR with Kapa
Library Quantification kit for Illumina.  Each library was diluted to 2nM and equal
volumes of 16 libraries were pooled to fill each NovaSeq sequencing run.  Pools were
sequenced with 100 cycle run kits with 26 bases Read 1, 8 bases Index 1 and 90 bases
Read 2 (Illumina 20012862).  PhiX control library was spiked in at 0.2 to 1%.  Libraries
were sequenced on the NovaSeq 6000 Sequencing System (Illumina)
473
474    **Data Processing**
475    Sequences from the NovaSeq were demultiplexed using bcl2fastq version 2.19.0.316.
476    Reads were aligned using to a copy of the mm10 genome augmented with ERCC
477    sequences, using STAR version 2.5.2b with the following options:
478
479    ```
       --outFilterType BySJout --outFilterMultimapNmax 20 --alignSJoverhang
480    Min 8 --alignSJDBoverhangMin 1 --outFilterMismatchNmax 999 --outFilt
481    erMismatchNoverLmax 0.04 --alignIntronMin 20 --alignIntronMax 100000
482    0 --alignMatesGapMax 1000000 --outSAMstrandField intronMotif --outSA
483    Mtype BAM Unsorted --outSAMattributes NH HI NM MD --genomeLoad LoadA
484    ndKeep --outReadsUnmapped Fastx --readFilesCommand zcat
       ```
485
486    Gene counts were produced using HTSEQ version 0.6.1p1 with default parameters
487    except stranded was set to false and mode was set to intersection-nonempty.
488
489    Sequences from the microfluidic platform were demultiplexed and aligned using
490    CellRanger with default parameters.
491
492    **Clustering**
493    Standard procedures for filtering, variable gene selection, dimensionality reduction, and
494    clustering were performed using the Seurat package. For each tissue and each sequencing
495    method (FACS and microfluidic), the following steps were performed. (The parameters
496    used were tuned on a per-tissue basis. Values appear in the supplement.)
497
498        1.  Genes appearing in fewer than 5 cells were excluded.
499        2.  Cells with fewer than 500 genes were excluded. Cells with fewer than 50,000
500            reads (FACS) or 1000 UMI (microfluidic) were excluded.  (In some organs, cells
501            with more than 2 million reads were also excluded as a conservative measure to
502            avoid doublets.)
503        3.  Counts were log-normalized  (log(1 + counts per N)), then scaled by linear
504            regression against the number of reads (or UMIs), the percent of reads mapping to
505            Rn45s, and the percent of reads to ribosomal genes.

506    4.  Variable genes were selected using a threshold for dispersion (log of
507        variance/mean). The distribution of dispersion in each expression bin was
508        computed, and variable genes were those with a z-score of at least 0.5-1..
509    5.  The variable genes were projected onto a low-dimensional subspace using
510        principal component analysis. The number of principal components was selected
511        based on inspection of the plot of variance explained.
512    6.  A shared-nearest-neighbors graph was constructed based on the Euclidean
513        distance metric in the low-dimensional subspace. (The SNN is a weighted graph
514        where $w_{ij}$ is the Jaccard similarity of the k-neighborhood of i with the k-
515        neighborhood of j, where $k = 30$.) Cells were clustered using the Louvain method
516        to maximize modularity.
517    7.  Cells were visualized using a 2-dimensional t-distributed Stochastic Neighbor
518        Embedding on the same distance metric.
519    8.  Cell types were assigned to each cluster using the abundance of known marker
520        genes. (Plots showing the expression of the markers for each tissue appear in the
521        supplement).
522    9.  When clusters appeared to be mixtures of cell types, they were refined either by
523        increasing the resolution parameter for Louvain clustering or subsetting the data
524        and rerunning steps 3-7.
525

526   A similar analysis was done globally for all FACS processed cells and for all microfluidic
527   processed cells.
528

529   To attempt to remove the effect of technical variation from the gene expression matrix, it
530   is common practice to regress out statistics associated with quality, like the number of
531   reads or the percent of ribosomal or mitochondrial RNA. If those statistics differ
532   systematically between cell types or tissues, this can have the unwanted effect of
533   compressing the true variation between those groups. For example, the percent of
534   ribosomal RNA varies from 1% in the Liver and Pancreas to 5% in the Spleen and
535   Tongue. Hence we regress out these factors separately in the analysis of each tissue.
536

537   For the "dynamic range" comparison in Supplementary Figure 4, the gene expression was
538   scaled to log of reads per million for FACS and log of UMI per 10,000 for microfluidic
539   emulsions. Genes with nonzero expression were rank ordered and plotted.
540
541
542
543

**References**

1.  Thorsen, T., Roberts, R. W., Arnold, F. H. & Quake, S. R. Dynamic pattern formation in a vesicle-generating microfluidic device. *Phys. Rev. Lett.* **86,** (2001).
2.  Macosko, E. Z. *et al.* Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161,** 1202–1214 (2015).
3.  Ramsköld, D. *et al.* Full-length mRNA-Seq from single-cell levels of RNA and individual circulating tumor cells. *Nat. Biotechnol.* **30,** 777–782 (2012).
4.  Alberts, B. *et al. Molecular biology of the cell*. (Garland Science, 2002).
5.  Guo, G. *et al.* Resolution of Cell Fate Decisions Revealed by Single-Cell Gene Expression Analysis from Zygote to Blastocyst. *Dev. Cell* **18,** 675–685 (2010).
6.  Dalerba, P. *et al.* Single-cell dissection of transcriptional heterogeneity in human colon tumors. *Nat. Biotechnol.* **29,** (2011).
7.  Wu, A. R. *et al.* Quantitative assessment of single-cell RNA-sequencing methods. (2013). doi:10.1038/nmeth.2694
8.  Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature* **509,** (2014).
9.  Enge, M. *et al.* Single-Cell Analysis of Human Pancreas Reveals Transcriptional Signatures of Aging and Somatic Mutation Patterns. *Cell* **171,** (2017).
10. Halpern, K. B. *et al.* Single-cell spatial reconstruction reveals global division of labour in the mammalian liver. *Nature* **542,** 352–356 (2017).
11. Haber, A. L. *et al.* A single-cell survey of the small intestinal epithelium. *Nature* **551,** 333–339 (2017).
12. Villani, A.-C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science* **356,** eaah4573 (2017).
13. Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell level. *Proc. Natl. Acad. Sci. U. S. A.* **112,** (2015).
14. Gokce, O. *et al.* Cellular Taxonomy of the Mouse Striatum as Revealed by Single-Cell RNA-Seq. *Cell Rep.* **16,** (2016).
15. Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat. Neurosci.* **18,** 145–153 (2014).
16. Zeisel, A. *et al.* Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* **347,** 1138–42 (2015).
17. Li, H. *et al.* Classifying Drosophila Olfactory Projection Neuron Subtypes by Single-Cell RNA Sequencing. *Cell* **171,** 1206–1220.e22 (2017).
18. van den Brink, S. C. *et al.* Single-cell sequencing reveals dissociation-induced gene expression in tissue subpopulations. *Nat. Methods* **14,** 935–936 (2017).

583
584 **Figure captions**
585
586 **Figure 1.** a) 19 tissues from 4 male and 3 female mice were analyzed. After tissue
587 dissociation, cells were either sorted by FACS or manipulated in microfluidic emulsions,
588 after which they were lysed and their transcriptomes amplified, followed by sequencing,
589 read mapping, and data analysis. b) Barplot showing number of sequenced cells prepared
590 by FACS sorting from each tissue (n=20). c) Barplot showing number of sequenced cells
591 prepared by microfluidic emulsion from each tissue (n=12). d) Histogram of number of
592 reads per cell for each tissue from FACS sorted cells. e) Histogram of number of genes
593 detected per cell for each tissue from FACS sorted cells. f) Histogram of number of
594 unique molecular identifiers (UMI) sequenced per cell for each tissue from cells prepared
595 by microfluidic emulsion. g) Histogram of number of genes detected per cell for each
596 tissue for cells prepared by microfluidic emulsion.
597
598 **Figure 2**. a) Dimensionally reduced tSNE plot of all cells sorted by FACS color coded
599 by tissue of origin. b) Dimensionally reduced tSNE plots for each tissue of cells sorted by
600 FACS. Color coding indicates distinct clusters. c) Barplots of manually annotated cell
601 types based on differential gene expression across all tissues.
602
603 **Figure 3**. Comparison of cell type determination as done by unbiased whole
604 transcriptome comparison versus manual annotation by organ-specific experts. The x-
605 axis represents clusters from Figure 2a with multiple organs contributing, while the y-
606 axis represents manual expert annotation of cell types in an organ-specific fashion based
607 on the data in Figure 2b and 2c. The unbiased method discovers relationships between
608 similar cell types found in different organs (highlighted regions); in particular it groups T
609 cells from different organs into a single cluster, B cells from different organs into a
610 different single cluster, and endothelial cells from different organs into a single cluster.
611
612 **Figure 4**. Analysis of all T cells sorted by FACS. a) Dimensionally reduced tSNE plot of
613 all T cells colored by cluster membership. Five clusters were identified. b) Dotchart
614 showing level of expression (color scale) and number of expressing cells (point diameter)
615 within each cluster of T cells. c) Barplot showing fraction of T cells originating from Fat,
616 Lung, Marrow, Muscle, Spleen or Thymus for each of the T cell clusters. d) Barplot
617 showing fraction of $Cd4^+$/ $Cd8^+$/ $Cd4^+Cd8^+$ / $Cd4^-Cd8^-$ T cells for each of the T cell
618 clusters.
619 **Figure 5.** Transcription factor expression analysis. A. Gene-gene correlation (correlogram) of top
620 cell type-tissue specific transcription factors (TFs) (selected by minimum differential
621 expression of 0.3 and p-value of $10^{-6}$; immediate early genes excluded). Columns are
622 colored by the cell ontology (left column) and tissue (right column) of the cell
623 type+tissue with the highest expression of that gene. Insets show clusters of TFs enriched
624 in selected cell types and tissues. B. Correlogram of top tissue-specific TFs for epithelial
625 cells (cells ontologies containing "epithelial", "basal", "keratinocyte", or "epidermis").
626 Row colors correspond to tissue (right) and cell ontology (left) of most-enriched cell
627 type. tSNE plots are calculated using all variable genes and show tissue origin (right) and

628     gene expression of select TFs (bottom).  C. Correlogram and tSNE of tissue-specific TFs
629     within stromal cells (cells ontologies containing "stromal", "fibroblast", or
630     "mesenchymal").  D. Correlogram and tSNE of tissue-specific TFs within endothelial
631     cells.

633 **Supplementary Figure Captions**

635 **Supplementary Figure 1** Histograms of number of genes detected across tissues (Red =

636 FACS sorted cells, Cyan = cells processed in microfluidic emulsion)

638 **Supplementary Figure 2** Plots across three cell types which compare number of cells

639 expressing a given gene for measurements made by FACS (x axis) compared to

640 microfluidic emulsion (y axis). Each data point is a gene and the axes represent fraction

641 of cells which express that gene.

643 **Supplementary Figure 3** Correlation of number of genes detected in more cells across

644 all common cell types. X-axis represents number of genes present in cell types measured

645 by FACS, whereas Y-axis represents number of genes present in the same cell types

646 measured by microfluidic emulsion. Most but not all of the cell types have more genes

647 measured by FACS.

649 **Supplementary Figure 4** Plots showing examples of dynamic range for the two

650 measurements approaches; FACS generally has many more reads per gene.

652 **Supplementary Figure 5** a) Dimensionally reduced tSNE plot of all cells manipulated

653 by microfluidic emulsions color coded by tissue of origin. b) Dimensionally reduced

654 tSNE plots for each tissue of cells sorted by microfluidic emulsions. Color coding

655 indicates distinct clusters. c) Barplots of manually annotated cell types based on

656 differential gene expression across all tissues.

658 **Supplementary Figure 6** Analysis of dissociation induced gene expression scores

659 across tissues.

661 **Supplementary Figure 7** Cell type- and tissue-specific enrichment of gene expression

662 in TF gene clusters. Gene-gene correlations were calculated as in Figure 5 and clustered

663 by cutting a hierarchical clustering tree at a height of 1.5 (Euclidean distance of

664 correlations). To calculate scores, log10 cpm values were scaled to [0,1] for each gene.

665 The score on the x-axis is the log of the geometric mean of all of the TFs in the cluster

666 across all the cells of a cell type and tissue. The TFs in the cluster are indicated at the top

667 of the plot; for clusters with more than 5 genes, only the first 5 are shown. The

668 distribution of scores across all unique cell types and tissues is displayed, with the bar

669 plots colored by tissue origin of the cell type (colors as in Fig. 5). Gene clusters that are

670 highly specific show a strongly bimodal plot, with most cells expressing very low levels

671 of the gene and one or a small number expressing high levels (for example, the "Spdef

672 Atoh1 Creb3l4" cluster is highly specific to colon goblet cells). The names of the top

673 three cell types are displayed as a subtitle below the genes, with the highest-scored at the

674 top. Note that even though some gene clusters only separate one or two cell types, the top

675 3 by rank are still included in the subtitle.

# Cell type- and tissue-specific transcription factor signatures