**Uterine progesterone signaling is a target for metformin therapy in PCOS-like rats**

Min Hu[1,2,+]; Yuehui Zhang[1,3,+]; Jiaxing Feng[3,+]; Xue Xu[3]; Jiao Zhang[4]; Wei Zhao[3]; Xiaozhu

Guo[3]; Juan Li[1,2]; Edvin Vestin[1]; Peng Cui[1,5,6]; Xin Li[1,7,8]; Xiao-ke Wu[3]; Mats Brännström[9];

Linus R Shao[1]; and Håkan Billig[1]

[1] Department of Physiology/Endocrinology, Institute of Neuroscience and Physiology, The

Sahlgrenska Academy, University of Gothenburg, 40530 Gothenburg, Sweden; [2] Department

of Traditional Chinese Medicine, The First Affiliated Hospital of Guangzhou Medical

University, 510120 Guangzhou, China; [3] Department of Obstetrics and Gynecology, Key

Laboratory and Unit of Infertility in Chinese Medicine, First Affiliated Hospital, Heilongjiang

University of Chinese Medicine, 150040 Harbin, China; [4] Department of Acupuncture and

Moxibustion, Second Affiliated Hospital, Heilongjiang University of Chinese Medicine,

150040 Harbin, China; [5] Department of Integrative Medicine and Neurobiology, State Key

Lab of Medical Neurobiology, Shanghai Medical College and Institute of Acupuncture

Research (WHO Collaborating Center for Traditional Medicine), Institute of Brain Science,

Fudan University, 200032 Shanghai, China; [6] Institute of Integrative Medicine of Fudan

University, 200032 Shanghai, China; [7] Department of Gynecology, Obstetrics and

Gynecology Hospital of Fudan University, 200011 Shanghai, China; [8] Shanghai Key

Laboratory of Female Reproductive Endocrine Related Diseases, 200011 Shanghai, China; [9]

Department of Obstetrics and Gynecology, Sahlgrenska University Hospital, Sahlgrenska

Academy, University of Gothenburg, 41345 Gothenburg, Sweden

[+] The authors consider that the first three authors should be regarded as joint first authors.

**\* Present address, to where correspondence and reprint requests should be addressed**

Linus R. Shao, MD, PhD., Tel: +46 31-7863408, Fax: +46 31-7863512, E-mail:
linus.r.shao@fysiologi.gu.se
26

**Short title:** Metformin and uterine progesterone signaling

Word count of the full article, excluding references and figure legends: **4388**

**Abstract**

Impaired progesterone (P4) signaling is linked to endometrial dysfunction and infertility in

women with polycystic ovary syndrome (PCOS). Here we report for the first time that

elevated expression of progesterone receptor (PGR) isoforms A and B parallels increased

estrogen receptor (ER) expression in PCOS-like rat uteri. The aberrant PGR-targeted gene

expression in PCOS-like rats before and after implantation overlaps with dysregulated

expression of *Fkbp52* and *Ncoa2,* two genes that contribute to the development of uterine P4

resistance. *In vivo* and *in vitro* studies of the effects of metformin on the regulation of the

uterine P4 signaling pathway under PCOS conditions showed that metformin directly inhibits

the expression of PGR and ER along with the regulation of several genes that are targeted

dependently or independently of PGR-mediated uterine implantation. Functionally, metformin

treatment corrected the abnormal expression of cell-specific PGR and ER and some PGR-

target genes in PCOS-like rats with implantation. Additionally, we documented how

metformin contributes to the regulation of the PGR-associated MAPK/ERK/p38 signaling

pathway in the PCOS-like rat uterus. Our data provide novel insights into how metformin

therapy regulates uterine P4 signaling molecules under PCOS conditions.

**Key words:** Metformin, progesterone receptor, MAPK/ERK/p38 signaling pathway,

implantation, polycystic ovary syndrome

## Introduction

Polycystic ovary syndrome (PCOS) is a clinically and etiologically heterogeneous hormone-imbalance disorder that is associated with multiple reproductive and metabolic abnormalities (Rosenfield and Ehrmann 2016). Women suffering from PCOS present with arrested folliculogenesis and chronic anovulation-linked infertility (Azziz *et al.* 2016; Rosenfield and Ehrmann 2016), and they also have more adverse reproductive risk as evidenced by an increase in the prevalence of implantation failure, recurrent miscarriage, spontaneous abortion, premature delivery, endometrial carcinoma (Goodarzi *et al.* 2011; Palomba *et al.* 2015; Shao *et al.* 2014b). In addition to the ovarian dysfunction (Azziz *et al.* 2016; Rosenfield and Ehrmann 2016), it is assumed that the impairment of endometrial function also contributes to PCOS-associated infertility (Evans *et al.* 2016). Although progesterone (P4)-based oral contraceptive therapy is often efficacious (Lopes *et al.* 2014; Vrbikova and Cibula 2005), perturbations in endometrial P4 signaling that result from attenuated responsiveness and resistance to P4 are common in the endometrium of a PCOS patient (Li *et al.* 2014a; Piltonen 2016). P4 resistance is a condition in which tissues and cells do not respond appropriately to P4 (Chrousos *et al.* 1986), and this is evidenced by endometriosis and endometrial hyperplasia that may progress to endometrial carcinoma despite supplementation with P4 or its analogs (Gunderson *et al.* 2012; Shao *et al.* 2014a). Gene profiling experiments have shown that different endometrial genes are likely to act in concert in this abnormal condition in PCOS patients (Kim *et al.* 2009; Savaris *et al.* 2011); however, how changes in the expression of P4 signaling molecules contribute to the P4 resistance in a PCOS patient's endometrium is poorly understood.

P4 is an essential contributing factor in female reproductive tissues that regulates multiple physiological processes such as the menstrual cycle, implantation, pregnancy maintenance, and labor initiation (Evans *et al.* 2016). There are two major progesterone receptor (PGR)

isoforms, PGRA and PGRB, both of which are involved in a common P4 signaling pathway

for uterine cell-specific proliferation and differentiation (Li *et al.* 2014a; Patel *et al.* 2015). P4

binding activates both PGR isoforms and leads to translocation from the cytosol to the

nucleus followed by binding to the P4-responsive elements of the target genes, resulting in

alterations of PGR-targeted gene expression depending on the recruitment of co-regulators

(Patel *et al.* 2015). It has been reported that endometrial PGR expression is elevated in PCOS

patients who have anovulation compared to PCOS patients who still ovulate and to non-PCOS

patients (Margarit *et al.* 2010; Quezada *et al.* 2006). Additionally, PGR activity is also

modulated by the cytoplasmic mitogen-activated protein kinase (MAPK)/extracellular signal-

regulated kinase (ERK) signaling pathway (Gellersen and Brosens 2014; Patel *et al.* 2015).

While high levels of ERK1/2 expression and activation reflect the P4-PGR signaling-induced

decidualization status in human and rodent uteri (Lee *et al.* 2013; Tapia-Pizarro *et al.* 2017;

Thienel *et al.* 2002), it remains to be determined whether suppression of MAPK/ERK

signaling occurs in the endometrium and whether such dysregulation can negatively impact

uterine function under PCOS conditions.

Metformin is an anti-diabetic drug that is a clinically approved treatment in PCOS patients

worldwide (Naderpoor *et al.* 2015). Several diverse molecular mechanisms of metformin have

been demonstrated in human endometrial carcinoma tissues *in vivo* and in different

endometrial cancer cells *in vitro* (Shao *et al.* 2014b), and metformin's therapeutic effects on

endometrial function are evidenced by improvement of endometrial receptivity, enhancement

of endometrial vascularity and blood flow, and reversion of endometrial hyperplasia and

carcinoma into normal endometria in some women with PCOS (Jakubowicz *et al.* 2001; Li *et*

*al.* 2014b; Palomba *et al.* 2006). Our recent studies using a PCOS-like rat model found that

chronic treatment with metformin has significant anti-androgenic and anti-inflammatory

impacts in the uterus (Zhang *et al.* 2017). Given the central role of P4 signaling in uterine

implantation (Evans *et al.* 2016; Patel *et al.* 2015) and the ability of metformin to rescue

implantation failure in some PCOS-like rats by modulating the expression of multiple

implantation-related genes in the uterus *in vivo* (Zhang *et al.* 2017), we speculated that the

beneficial effects of metformin might be mechanistically linked to the uterine P4 signaling

pathway under pathological conditions such as PCOS. To address this hypothesis, we

analyzed PCOS-associated PGR isoform expression and the MAPK signaling network in

human and rat uterine tissues. By combining a PCOS-like rat model (Zhang *et al.* 2016) and

*in vitro* tissue culture approach (Li *et al.* 2015), we aimed to determine whether metformin

directly reverses aberrant PGR-targeted and implantation-related gene expression in the

PCOS-like rat uterus.

**Materials and Methods**

*Study approval*

All animal experiments were performed according to the National Institutes of Health

guidelines on the care and use of animals and were approved and authorized by the Animal

Care and Use Committee of the Heilongjiang University of Chinese Medicine, China (HUCM

2015-0112).

*Experimental animals and tissue preparations*

Adult female Sprague–Dawley rats (n = 134) were obtained from the Laboratory Animal

Centre of Harbin Medical University, Harbin, China (License number SCXK 2013-001).

Animals were housed in the animal care facility with free access to food and water and a

controlled temperature of 22°C ± 2°C with a 12 h light/dark cycle. Estrous cycles were

monitored daily by vaginal lavage according to a standard protocol (Feng *et al.* 2010). All rats

(70 days old) with the different stages of estrous cycle used in this study were confirmed by

examination of vaginal smears under a light microscope for two sequential cycles (about 8–10

days). Any PCOS-like (insulin+hCG-treated) rats that exhibited prolonged estrous cycles

(more than 5 days) were excluded from the study.

Experiment 1: Rats were randomly divided into control (saline treatment, n = 20) and

experimental (PCOS-like, n = 20) groups. The experimental group was treated with insulin

plus hCG to induce a PCOS-like metabolic and reproductive phenotype, and the control rats

were treated with an equal volume of saline (Zhang *et al.* 2018; Zhang *et al.* 2016). In brief,

insulin was started at 0.5 IU/day and gradually increased to 6.0 IU/day between day 1 and the

day 22 to induce hyperinsulinemia and insulin resistance, and 3.0 IU/day hCG was given on

all 22 days to induce hyperandrogenism. Animals were treated with twice-daily subcutaneous

injections until the end of the experiment. Rats with repeated insulin injections have not

shown any hypoglycemic episodes (Bogovich *et al.* 1999; Damario *et al.* 2000; Poretsky *et al.*

1992; Zhang *et al.* 2018). Detailed analysis of endocrine and metabolic parameters as well as

the uterine morphology in these animals has been reported previously (Zhang *et al.* 2016). On

day 23, each group of rats was divided into two subgroups of 10 rats each (Supplemental Fig.

1A). For treatment, metformin was dissolved in saline and given as a daily oral dose of 500

mg/kg by a cannula. The treatment time and tissue collection are described in our previous

study (Zhang *et al.* 2017).

Experiment 2: Rats were randomly divided into control (saline treatment, n = 21) and

experimental (PCOS-like, n = 15) groups and treated as described in Experiment 1

(Supplemental Fig. 1B). After metformin treatment, control and PCOS-like rats were mated

with fertile males of the same strain to induce implantation, which was determined by the

presence of a vaginal plug (day 1 of pregnancy). The rats were sacrificed between 0800 and

0900 hours on day 6 of pregnancy. To identify the implantation sites, rats were injected

intravenously with a Chicago Blue B dye solution (1% in saline) and sacrificed 10 min later.

Uteri were dissected and assessed for clearly delineated blue bands as evidence of early

implantation sites as described previously (Zhang *et al.* 2017).

Experiment 3: Rats were divided into control (saline treatment, n = 9) and experimental

(PCOS-like, n = 39) groups and treated as described in the Experiment 1. On the 23rd day, the

PCOS-like rats were divided into four subgroups and treated daily with P4 (4 mg/kg), RU486

(6 mg/kg), or both for 3 days. For treatment, P4 and RU486 were dissolved in 100% ethanol

and resuspended in sesame oil. All subcutaneous injections were in a volume of 100 μl. An

equal volume of 100% ethanol and sesame oil was injected into both healthy control rats and

PCOS-like rats as experimental controls (Supplemental Fig. 1C). The pharmacological doses

and treatment time intervals of P4 and RU486 were chosen on the basis of previous studies

(Kim *et al.* 2006; Knox *et al.* 1996).

After dissection, the uterine horns were trimmed free of fat and connective tissue. One side

of the uterus in each animal was fixed in 10% neutral formalin solution for 24 h at 4ºC and

embedded in paraffin for histochemical analysis. The other side was immediately frozen in

liquid nitrogen and stored at −70ºC for Western blot and quantitative real-time PCR (qRT-

PCR) analysis.

Detailed description of the methods including the primary *in vitro* tissue culture and

treatment, morphological assessment and immunostaining, protein isolation and Western blot

analysis, RNA extraction and qRT-PCR analysis, and measurement of biochemical

parameters used in this study are provided in Supplemental files.

***Statistical analysis***

GraphPad Prism was used for statistical analysis and graphing. For all experiments, n-

values represent the number of individual animals. Data are represented as the means ± SEM.

Statistical analyses were performed using SPSS version 24.0 statistical software for Windows

(SPSS Inc., Chicago, IL). The normal distribution of the data was tested with the Shapiro–

Wilk test. Differences between groups were analyzed by one-way ANOVA or two-way

ANOVA, and this was followed by Tukey's post-hoc test for normally distributed data or the

Kruskal–Wallis test followed by the Mann–Whitney U-test for skewed data. All $p$-values less

than 0.05 were considered statistically significant.

**Results**

***Metformin alters PGR isoform and PGR-targeted gene expression in PCOS-like rats***

The insulin+hCG-treated rats exhibit reproductive disturbances that mimic human PCOS

(Zhang *et al.* 2017; Zhang *et al.* 2016). Prompted by these findings, we set out to investigate

the impact of P4 signaling in this model. First, we showed that although the ratio of PGRA to

PGRB was not significantly different between control and PCOS-like rats, the PCOS-like rats

had increased levels of uterine PGRA and PGRB (Fig. 1A). While PGR immunoreactivity

was primarily evidenced in control rat uterine luminal and glandular epithelia as well as in the

stroma, the immunoreactivity of luminal epithelial PGR expression was associated with

increased numbers of luminal epithelial cells and increased immunoreactivity of PGR in the

stroma in PCOS-like rats (Fig. 1B). Metformin treatment did not significantly affect PGR

isoform expression in control rats and PCOS-like rats compared to those rats treated with

saline (Fig. 1A). However, we found that PGR immunoreactivity was decreased in the

luminal and glandular epithelia by metformin treatment in both control rats and PCOS-like

rats compared to those treated with saline (Fig. 1B). Conversely, intense immunoreactivity of

PGR expression was detected in the stroma located close to the luminal epithelia in control

and PCOS-like rats treated with metformin (Fig. 1B). In contrast to the epithelia and stroma,

no significant changes in PGR expression in the myometrium were found in any of the groups

(data not shown). Because a large body of evidence indicates that regulation of P4 signaling

results in changes in the expression of several PGR-targeted genes in the uterus (Bhurke *et al.*

2016), we profiled the expression of genes that are indicators for PGR activity in the rat uterus

by qRT-PCR. Quantitative data indicated that *Smo*, and *Nr2f2* mRNA levels were increased in

PCOS-like rats compared to control rats treated with saline. In contrast, the *Fkbp52* mRNA

level was decreased in PCOS-like rats compared to control rats (Fig. 1C). We next determined

the actions of metformin treatment on PGR-targeted gene expression and showed that *Ptch*,

*Fkbp52*, and *Ncoa2* levels were increased in PCOS-like rats treated with metformin compared

to PCOS-like rats treated with saline, while *Smo* and *Nr2f2* mRNA levels were decreased on

PCOS-like rats treated with metformin compared to those treated with saline (Fig. 1C).

***Metformin partially prevents implantation failure in parallel with regulation of PGR***

***isoform and PGR-targeted gene expression in PCOS-like rats***

Metformin has been shown to partially rescue the disruption of the implantation process in

PCOS-like rats (Zhang *et al.* 2017), and the altered endocrine and metabolic parameters in

these animals are shown in Supplemental Table 4. After metformin treatment, total

testosterone levels, the ratio of total testosterone to androstenedione, and fasting insulin levels

were all significantly higher in PCOS-like rats where implantation did not occur compared to

those with implantation, as was insulin resistance as assessed by the homeostasis model

assessment of insulin resistance, mirroring the endocrine and metabolic abnormalities in

PCOS patients (Azziz *et al.* 2016; Rosenfield and Ehrmann 2016). Of note, PCOS-like rats

that failed to implant embryos also exhibited decreased P4 levels. These data suggest that

implantation failure in PCOS-like rats treated with metformin is due not only to

hyperandrogenism and insulin resistance, but also to impairment of P4 signaling in the uterus.

Further morphological characterization of metformin-treated PCOS-like rats with no

implantation revealed the infiltration of immune cells into the glandular epithelial cell layer in

a similar manner to when hormone imbalances were studied in a previous report (Wira *et al.*

2005) (Supplemental Fig. 3, black arrowheads). To determine how impairment of P4

signaling causes implantation failure, we subsequently analyzed PGR isoform and PGR-

targeted gene expression in PCOS-like rats with no implantation. Although treatment with

metformin increased PGR isoform expression in control and PCOS-like rats, neither the

PGRA nor PGRB protein level was altered between PCOS-like rats with implantation and

with failed implantation (Fig. 2A). As shown in Figure 2B, while PGR protein was expressed

in the decidualizing stroma at the site of implantation in all groups, PGR immunoreactivity

was increased in the stroma of the inter-implantation region in control rats treated with

metformin. Furthermore, we found that the immunoreactivity of PGR was increased in the

epithelia in PCOS-like rats without implantation despite metformin treatment. Thus,

metformin appeared to participate in the regulation of uterine PGR expression in a cell type-

specific manner in PCOS-like rats before and after implantation. qRT-PCR data indicated that

*Ihh* and *Ncoa2* mRNAs were increased and that *Ptch* and *Fkbp52* mRNAs were decreased in

metformin-treated PCOS-like rats with no implantation compared to control rats treated with

saline or metformin and to metformin-treated PCOS-like rats with implantation (Fig. 2C).

***Metformin directly regulates PGR isoform, PGR-target, and implantation-related gene***

***expression in vitro***

Based on these *in vivo* observations, we asked whether the effect of metformin was direct

or indirect in the PCOS-like rat uterus. *In vitro* uterine tissue culture experiments revealed that

*Pgr* and *Pgrb*, mRNA levels were higher in PCOS-like rats compared to control rats, in

agreement with alteration of PGR isoform protein expression (Fig. 1C). Furthermore,

metformin treatment increased *Pgr* and *Pgrb* mRNA levels in control and PCOS-like rats in a

time-dependent manner (Fig. 3A). Consistent with the *in vivo* effects of metformin in PCOS-

like rats (Fig. 1C), *Ihh*, *Smo*, and *Nr2f2* mRNA levels were increased in the PCOS-like rat

uterus compared to the control rat uterus and were down-regulated by metformin treatment *in*

*vitro*. While the *Hand2* mRNA level was upregulated by metformin treatment at 48 h and 72 h,

we detected the upregulation of *Ptch*, *Fkbp52*, and *Ncoa2* mRNA levels in the PCOS-like rat

uterine tissues over a 72-h course after metformin treatment (Fig. 3A).

The expression of a number of implantation-related genes has been reported to be

regulated by metformin treatment in PCOS-like rats during implantation (Zhang *et al.* 2017).

These previous observations prompted further analysis of implantation-related gene

expression by metformin treatment *in vitro*. In contrast to the different regulation patterns of

*Spp1*, *Lrh1*, *Sgk1*, and *Krt13* mRNAs under *in vivo* and *in vitro* conditions, the *in vitro*

responses of uterine *Prl*, *Igfbp1*, *Il11*, *Pc6*, *Maoa*, *Ednrb*, *Hoxa10*, *Hoxa11*, and *Hbegf* mRNA

levels to metformin (Fig. 3B) were coincident with the *in vivo* regulation of the expression

pattern of these genes (Zhang *et al.* 2017). Our data indicated that metformin directly up-

regulates uterine *Prl*, *Maoa*, *Ednrb*, and *Hbegf* mRNA levels in PCOS-like rats during

implantation *in vivo*.

To ascertain whether the modulation of uterine gene expression is P4-mediated and PGR-

dependent in PCOS-like rats, insulin+hCG-treated rats were injected subcutaneously with P4

and/or RU 486 for three days. As shown in Figure 4, the increased PGR isoform protein levels

(Fig. 1A) were confirmed by analysis of *Pgr* and *Pgrb* mRNA expression in the PCOS-like

rat uterus. Although treatment with P4 and/or RU486 did not significantly affect *Pgr* mRNA

expression, we found that *Pgrb* mRNA levels were decreased in PCOS-like rats compared to

those rats with no treatment (Fig. 4). Among seven PGR-targeted genes (Fig. 1C), we found

that *Ptch*, *Hand2*, and *Fkbp52* mRNA levels were increased and that *Ihh*, *Smo*, and *Nr2f2*

mRNA levels were decreased in PCOS-like rats treated with P4 compared to those rats with

no treatment. We also observed that treatment with RU486 alone or combined with P4

reversed the changes in *Smo*, *Hand2*, and *Fkbp52* mRNA levels in PCOS-like rats (Fig. 4). No

significant differences of uterine *Ncoa2* mRNA expression were observed in PCOS-like rats

regarding the different treatments. Based on our current experimental approaches, it is likely

that another regulatory mechanism contributes to the metformin-induced up-regulation of

*Ncoa2* mRNA levels in PCOS-like rats.

***Metformin regulates the MAPK signaling pathway in PCOS-like rats before and after***

***implantation***

In an attempt to understand the changes in PGR activation and function observed in PCOS

patients (Patel *et al.* 2015), we performed a Western blot analysis to measure the expression

of several proteins that are involved in the MAPK signaling pathway in the uterus after

metformin treatment. As shown in Figure 5A, there was no significant difference in p-c-Raf,

p-MEK1/2, p-ERK1/2, p-p38 MAPK, or p38 MAPK expression between saline-treated and

metformin-treated rats. Quantitative protein data indicated that the expression of p-p38

MAPK and p38 MAPK was significantly decreased in PCOS-like rats compared to control

rats. Nevertheless, metformin treatment only reversed p-p38 MAPK protein expression in

PCOS-like rats.

We next assessed whether the MAPK/ERK signaling pathway contributes to uterine

implantation in control and PCOS-like rats treated with metformin. As shown in Figure 5B,

although the p-MEK1/2 level was decreased in control rats treated with metformin compared

to control rats treated with saline, no significant difference in p-c-Raf, p-ERK1/2, ERK1/2, p-

p38 MAPK, or p38 MAPK expression between these two groups was found. Furthermore, our

data showed that p-c-Raf, p-MEK1/2, and p-ERK1/2 protein levels were down-regulated in

PCOS-like rats treated with metformin regardless of the occurrence of implantation. We also

found that after metformin treatment PCOS-like rats with implantation exhibited decreased p-

p38 MAPK, but not p38 MAPK, expression.

***Up-regulation of estrogen receptor (ER) expression in PCOS-like rats can be suppressed by***

***metformin***

Because estrogen-ER signaling regulates uterine PGR expression and activity (Li *et al.*

2014a; Patel *et al.* 2015) and because increased circulating E2 in PCOS-like rats can be

inhibited by metformin treatment (Zhang *et al.* 2017), we sought to determine whether ER

subtypes (ERα and ERβ) are involved in the regulation of aberrant PGR expression in PCOS-

like rats and, if so, if metformin possibly alters ER subtype expression. Our data showed that

PCOS-like rats exhibited increased *Esr1* (ERα) and *Esr2* (ERβ) mRNA levels, which were

suppressed by metformin treatment. As shown in Figure 6A, while nuclear ERα

immunoreactivity was detected in the epithelia and stroma in control rats treated with saline

(Fig. 6B1), immunoreactivity of ERα was increased in the glandular epithelia and stroma in

PCOS-like rats (Fig. 6D1). Furthermore, treatment with metformin led to decreased ERα

immunoreactivity in control (Fig. 6C1) and PCOS-like rats (Fig. 6E1). No obvious difference

in ERα immunoreactivity was observed in the myometrium in any of the groups (Fig. 6B2-

E2). We also found that ERβ was mainly co-localized with ERα in the epithelia and stroma

but not in the myometrium in control and PCOS-like rats regardless of the different

treatments. Furthermore, with metformin treatment, we noted a significant increase in uterine

*Esr1* and *Esr2* mRNAs in PCOS-like rats without implantation (Fig. 7A).

Immunofluorescence staining revealed that, overall, immunoreactivities of both ERα and ERβ

were diminished in the decidualizing stroma at the site of implantation (Fig. 7B1–D1), in the

epithelia and stroma of the inter-implantation region (Fig. 7B2–D2) in control rats treated

with saline or metformin (Fig. 7B2–C2), and in the inter-implantation site of PCOS-like rats

treated with metformin (Fig. 7D2) compared to those rats before implantation (Fig 6. B1–E2).

Interestingly, PCOS-like rats with no implantation exhibited sustained nuclear ERα

immunoreactivity in the glandular epithelia and stroma (Fig. 7E1).

***Differential cell-specific expression of phospho-histone H3 in PCOS-like rats treated with***

***metformin***

As previously demonstrated (Avellaira *et al.* 2006), p-histone H3 is of special interest

because the endometrium of PCOS patients displays high levels of p-histone H3, which is

associated with cellular processes such as mitosis (Brenner *et al.* 2003). Quantitative

assessment of p-histone H3 indicated that no significant change in p-histone H3

immunoreactivity was present in the epithelia or stroma in any of groups (Supplemental Fig.

4E); however, metformin treatment decreased p-histone H3 immunoreactivity in the

myometrium in PCOS-like rats compared to those treated with saline. Of note, intensely p-

histone H3-positive stromal cells close to the luminal and glandular epithelia were found in

PCOS-like rats treated with metformin (Supplemental Fig. 4D2). Similarly, p-histone H3

immunoreactivity was significantly increased in the stroma at the inter-implantation sites in

PCOS-like rats treated with metformin independently of implantation (Supplemental Fig. 5E).

In PCOS-like rats without implantation, p-histone H3 immunoreactivity was often detected in

the luminal epithelia (Supplemental Fig. 5D1), although this was not statistically significant

compared to PCOS-like rats with implantation (Supplemental Fig. 5E). It is thus likely that

the regulation of mitotic activity by metformin is cell type-dependent in the uterus.

**Discussion**

Reproductive dysfunction and infertility manifest noticeably in PCOS patients (Evans *et al.*

2016). In striking contrast to the attention given to hyperandrogenism and insulin resistance in

women with PCOS, the aberrant P4 signaling pathway resulting in uterine P4 resistance has

received much less attention (Li *et al.* 2014a; Piltonen 2016). This study is the first to show

that the therapeutic effects of metformin on the regulation of uterine function in PCOS-like

rats is mediated through P4 signaling.

Elucidating the regulation of endometrial PGR levels under PCOS conditions is important

clinically. Our data show that increased PGR expression is paralleled with elevated ER

expression in PCOS-like rats. This expression pattern is associated with an increased

circulating E2 level (Zhang *et al.* 2016), suggests that E2-ER signaling contributes to the up-

regulation of PGR under PCOS conditions *in vivo*. Similar to PCOS patients (Hu *et al.*

submitted), PCOS-like rats also displayed high levels of PGR isoforms and ER subtypes in

the uterus. The induction of implantation is required for the activation of PGR, and

implantation subsequently alters gene expression in the endometrium (Gellersen and Brosens

2014; Patel *et al.* 2015); however, PGR-targeted gene expression in PCOS patients and

PCOS-like rats has only been demonstrated to a limited degree. The current study shows that

significantly decreased *Fkbp52* gene expression parallels increased expression of *Ihh*, *Smo*,

and *Nr2f2* mRNAs without changes in *Ncoa2* mRNA in PCOS-like rats. In addition, we also

found that abnormal expression of PGR-target genes, including *Fkbp52* and *Ncoa2*, is

retained in PCOS-like rats with implantation failure. This is supported by *in vivo* studies

showing that mice lacking *Fkbp52* (Tranguch *et al.* 2007; Yang *et al.* 2006) or *Ncoa2*

(Mukherjee *et al.* 2007; Mukherjee *et al.* 2006) demonstrate the absence of decidualization

after P4 supplementation due to diminished P4 responsiveness. Previous studies have reported

that women with endometriosis and endometrial hyperplasia/carcinoma who develop P4

resistance have low levels of PGR expression (Gunderson *et al.* 2012; Shao *et al.* 2014a).

Although it is currently unclear why differences exist in the regulation of uterine PGR

expression between different diseases with P4 resistance, it is likely that uterine P4 resistance

in PCOS-like rats is due to impaired PGR activity rather than PGR expression.

Defects in PGR isoform-specific P4 signaling in the mouse uterus can give rise to distinct

phenotypes of uterine impairment and implantation failure (Li *et al.* 2014a). Here we

observed no changes in total *Pgr* mRNA levels but a reduction of *Pgrb* mRNA levels in

PCOS-like rat uterus after treatment with P4 and/or RU486. This suggests that uterine *Pgra*

mRNA levels are increased. Meanwhile, several PGR target genes (e.g., *Ihh*, *Smo*, *Ptch*,

*Nr2f2*, *Hand2*, and *Fkbp52*) are significantly altered in PCOS-like rats after P4 treatment.

Thus, we speculate that changes in these P4-dependent PGR target gene expression in

PCOS-like rat uterus might be accounted for by an increase in *Pgra* mRNA expression.

Studies of mutant mice lacking specific PGR isoform will clarify the functional differences

between the two PGR isoforms in the progression of PCOS-induced uterine dysfunction.

P4-mediated and PGR-dependent regulation of ERK1/2 expression plays a critical role in

humans and rodents during endometrial decidualization and implantation (Lee *et al.* 2013;

Tapia-Pizarro *et al.* 2017; Thienel *et al.* 2002), but such regulation under PCOS conditions

has not previously been reported. The inhibition of ERK1/2 expression and activation has

been reported in ovarian granulosa and thecal cells in PCOS patients (Lan *et al.* 2015; Nelson-

Degrave *et al.* 2005), and we have previously shown that the expression and activation of

uterine ERK1/2 is suppressed in rats treated with insulin and hCG to induce the PCOS

phenotype (Zhang *et al.* 2016). The present study supports and extends this work. Here we

observed no changes in p-ERK1/2 or ERK1/2 expression in the rat uterus after prolonged

treatment with insulin and hCG. However, we observed that the levels of p-c-Raf and p-

MEK1/2, two upstream regulators of ERK1/2, were significantly decreased in PCOS-like rats

after uterine implantation, establishing a tight link between different MAPK/ERK signaling

molecules. Our data suggest that regulation of uterine ERK1/2 expression *in vivo* is time-

dependent (Lee *et al.* 2013; Tapia-Pizarro *et al.* 2017; Thienel *et al.* 2002), which is similar to

the regulation of PGR isoforms and PGR-targeted gene expression. The MAPK/ERK/p38

signaling pathway contributes to the regulation of inflammation and cytokine production

(Arthur and Ley 2013; Cuadrado and Nebreda 2010), and the dysregulation of inflammation-

related molecules is associated with PCOS conditions (Matteo *et al.* 2010; Orostica *et al.*

2016; Piltonen *et al.* 2013; Piltonen *et al.* 2015). Furthermore, like the activation of NFκB

signaling that induces the transcriptional levels of inflammation-related gene expression in

ovarian granulosa cells and in serum in PCOS patients (Liu *et al.* 2015; Zhao *et al.* 2015), our

previous study has shown that the sustained metformin treatment markedly suppresses uterine

inflammatory gene expression, especially the *Il-6* and *TNFα* mRNAs that are associated with

inhibition of nuclear NFκB translocation in PCOS-like rats (Zhang *et al.* 2017). Importantly,

p38 can antagonize ERK1/2 signaling mediated by protein phosphatase 2A and consequently

down-regulate inflammatory cytokine and chemokine production (Cuadrado and Nebreda

2010), and the anti-inflammatory effects of MAPK/p38 are involved in the regulation of

NFκB activity (Arthur and Ley 2013). These observations further indicate that metformin

inhibits NFκB-driven inflammatory processes through p38 activation rather than through

ERK1/2 inhibition in the PCOS-like rat uterus.

The results of the present study permit us to draw the following conclusions (Figure 8). 1)

With sustained low levels of P4, the expressions of both uterine PGR isoforms are elevated in

PCOS-like rats *in vivo*. This is positively associated with the high levels of ERs in PCOS-like

rats. Consistent with mouse knockout studies, altered expression of *Fkbp52* and *Ncoa2*, two

genes that contribute to uterine P4 resistance, is seen in PCOS-like rats before and after

implantation. 2) Metformin directly suppresses uterine PGR isoform expression along with

the correction of aberrant expression of PGR-targeted and implantation-related genes in

PCOS-like rats. Abnormal cell-specific regulation of PGR and ER, paralleling the aberrant

expression of PGR-targeted and implantation-related genes, is retained in PCOS-like rats with

implantation failure. 3) Increased PGR expression is associated with inhibition of the

MAPK/ERK/p38 signaling pathway, and the primary effect of metformin treatment is to

restore the MAPK/p38 signaling pathway in the PCOS-like rat uterus. Taken together, our

findings provide support for metformin therapy in the improvement of P4 signaling in PCOS-

like rats with uterine dysfunction and for its clinical relevance in the treatment of PCOS

patients with P4 resistance.

**Declaration of interest**

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

**Funding**

Research reported in this publication was supported by the Swedish Medical Research Council (grant number 10380), the Swedish federal government under the LUA/ALF agreement (grant number ALFGBG-147791), the Jane and Dan Olsson's Foundation, the Hjalmar Svensson Foundation, and the Adlerbert Research Foundation (to HB and LRS) as well as the National Natural Science Foundation of China (grant number 81774136), the Project of Young Innovation Talents in Heilongjiang Provincial University (grant number UNPYSCT-2015121), the Project of Innovation Talents (Young Reserve Talents) in Harbin city (grant number 2015RAQYJ089), and the Project of Excellent Innovation Talents by Heilongjiang University of Chinese Medicine (to YZ).

**Acknowledgements**

The authors thank the Centre for Cellular Imaging at the University of Gothenburg and the National Microscopy Infrastructure (grant number VR-RFI 2016-00968) for providing assistance in microscopy.

**REFERENCES**

Arthur JS & Ley SC 2013 Mitogen-activated protein kinases in innate immunity. *Nat Rev*
*Immunol* **13** 679-692.
Avellaira C, Villavicencio A, Bacallao K, Gabler F, Wells P, Romero C & Vega M 2006
Expression of molecules associated with tissue homeostasis in secretory endometria from
untreated women with polycystic ovary syndrome. *Hum Reprod* **21** 3116-3121.
Azziz R, Carmina E, Chen Z, Dunaif A, Laven JS, Legro RS, Lizneva D, Natterson-Horowtiz
B, Teede HJ & Yildiz BO 2016 Polycystic ovary syndrome. *Nat Rev Dis Primers* **2** 16057.
Bhurke AS, Bagchi IC & Bagchi MK 2016 Progesterone-Regulated Endometrial Factors
Controlling Implantation. *Am J Reprod Immunol* **75** 237-245.
Bogovich K, Clemons J & Poretsky L 1999 Insulin has a biphasic effect on the ability of
human chorionic gonadotropin to induce ovarian cysts in the rat. *Metabolism* **48** 995-1002.
Brenner RM, Slayden OD, Rodgers WH, Critchley HO, Carroll R, Nie XJ & Mah K 2003
Immunocytochemical assessment of mitotic activity with an antibody to phosphorylated
histone H3 in the macaque and human endometrium. *Hum Reprod* **18** 1185-1193.
Chrousos GP, MacLusky NJ, Brandon DD, Tomita M, Renquist DM, Loriaux DL & Lipsett
MB 1986 Progesterone resistance. *Adv Exp Med Biol* **196** 317-328.
Cuadrado A & Nebreda AR 2010 Mechanisms and functions of p38 MAPK signalling.
*Biochem J* **429** 403-417.
Damario MA, Bogovich K, Liu HC, Rosenwaks Z & Poretsky L 2000 Synergistic effects of
insulin-like growth factor-I and human chorionic gonadotropin in the rat ovary. *Metabolism*
**49** 314-320.
Evans J, Salamonsen LA, Winship A, Menkhorst E, Nie G, Gargett CE & Dimitriadis E 2016
Fertile ground: human endometrial programming and lessons in health and disease. *Nat Rev*
*Endocrinol* **12** 654-667.
Feng Y, Weijdegard B, Wang T, Egecioglu E, Fernandez-Rodriguez J, Huhtaniemi I, Stener-
Victorin E, Billig H & Shao R 2010 Spatiotemporal expression of androgen receptors in the
female rat brain during the oestrous cycle and the impact of exogenous androgen
administration: a comparison with gonadally intact males. *Mol Cell Endocrinol* **321** 161-174.
Gellersen B & Brosens JJ 2014 Cyclic decidualization of the human endometrium in
reproductive health and failure. *Endocr Rev* **35** 851-905.
Goodarzi MO, Dumesic DA, Chazenbalk G & Azziz R 2011 Polycystic ovary syndrome:
etiology, pathogenesis and diagnosis. *Nat Rev Endocrinol* **7** 219-231.
Gunderson CC, Fader AN, Carson KA & Bristow RE 2012 Oncologic and reproductive
outcomes with progestin therapy in women with endometrial hyperplasia and grade 1
adenocarcinoma: a systematic review. *Gynecol Oncol* **125** 477-482.
Jakubowicz DJ, Seppala M, Jakubowicz S, Rodriguez-Armas O, Rivas-Santiago A, Koistinen
H, Koistinen R & Nestler JE 2001 Insulin reduction with metformin increases luteal phase
serum glycodelin and insulin-like growth factor-binding protein 1 concentrations and
enhances uterine vascularity and blood flow in the polycystic ovary syndrome. *J Clin*
*Endocrinol Metab* **86** 1126-1133.
Kim HJ, Lee GS, Ji YK, Choi KC & Jeung EB 2006 Differential expression of uterine
calcium transporter 1 and plasma membrane Ca2+ ATPase 1b during rat estrous cycle. *Am J*
*Physiol Endocrinol Metab* **291** E234-241.
Kim JY, Song H, Kim H, Kang HJ, Jun JH, Hong SR, Koong MK & Kim IS 2009
Transcriptional profiling with a pathway-oriented analysis identifies dysregulated molecular
phenotypes in the endometrium of patients with polycystic ovary syndrome. *J Clin*
*Endocrinol Metab* **94** 1416-1426.

Knox KL, Ringstrom SJ, Szabo M, Perlyn CA, Sutandi S & Schwartz NB 1996 RU486 on an
estrogen background blocks the rise in serum follicle-stimulating hormone induced by
antiserum to inhibin or ovariectomy. *Endocrinology* **137** 1226-1232.

Lan CW, Chen MJ, Tai KY, Yu DC, Yang YC, Jan PS, Yang YS, Chen HF & Ho HN 2015
Functional microarray analysis of differentially expressed genes in granulosa cells from
women with polycystic ovary syndrome related to MAPK/ERK signaling. *Sci Rep* **5** 14994.

Lee CH, Kim TH, Lee JH, Oh SJ, Yoo JY, Kwon HS, Kim YI, Ferguson SD, Ahn JY, Ku BJ
*et al.* 2013 Extracellular signal-regulated kinase 1/2 signaling pathway is required for
endometrial decidualization in mice and human. *PLoS One* **8** e75282.

Li X, Cui P, Jiang HY, Guo YR, Pishdari B, Hu M, Feng Y, Billig H & Shao R 2015
Reversing the reduced level of endometrial GLUT4 expression in polycystic ovary syndrome:
a mechanistic study of metformin action. *Am J Transl Res* **7** 574-586.

Li X, Feng Y, Lin JF, Billig H & Shao R 2014a Endometrial progesterone resistance and
PCOS. *J Biomed Sci* **21** 2.

Li X, Guo JR, Lin JF, Feng Y, Billig H & Shao R 2014b Combination of Diane-35 and
metformin to treat early endometrial carcinoma in PCOS women with insulin resistance *J*
*Cancer* **5** 173-181.

Liu M, Gao J, Zhang Y, Li P, Wang H, Ren X & Li C 2015 Serum levels of TSP-1, NF-
kappaB and TGF-beta1 in polycystic ovarian syndrome (PCOS) patients in northern China
suggest PCOS is associated with chronic inflammation. *Clin Endocrinol (Oxf)* **83** 913-922.

Lopes IM, Maganhin CC, Oliveira-Filho RM, Simoes RS, Simoes MJ, Iwata MC, Baracat EC
& Soares JM, Jr. 2014 Histomorphometric Analysis and Markers of Endometrial Receptivity
Embryonic Implantation in Women With Polycystic Ovary Syndrome During the Treatment
With Progesterone. *Reprod Sci* **21** 930-938.

Margarit L, Taylor A, Roberts MH, Hopkins L, Davies C, Brenton AG, Conlan RS,
Bunkheila A, Joels L, White JO *et al.* 2010 MUC1 as a discriminator between endometrium
from fertile and infertile patients with PCOS and endometriosis. *J Clin Endocrinol Metab* **95**
5320-5329.

Matteo M, Serviddio G, Massenzio F, Scillitani G, Castellana L, Picca G, Sanguedolce F,
Cignarelli M, Altomare E, Bufo P *et al.* 2010 Reduced percentage of natural killer cells
associated with impaired cytokine network in the secretory endometrium of infertile women
with polycystic ovary syndrome. *Fertil Steril* **94** 2222-2227, 2227 e2221-2223.

Mukherjee A, Amato P, Allred DC, DeMayo FJ & Lydon JP 2007 Steroid receptor
coactivator 2 is required for female fertility and mammary morphogenesis: insights from the
mouse, relevance to the human. *Nucl Recept Signal* **5** e011.

Mukherjee A, Soyal SM, Fernandez-Valdivia R, Gehin M, Chambon P, Demayo FJ, Lydon
JP & O'Malley BW 2006 Steroid receptor coactivator 2 is critical for progesterone-dependent
uterine function and mammary morphogenesis in the mouse. *Mol Cell Biol* **26** 6571-6583.

Naderpoor N, Shorakae S, de Courten B, Misso ML, Moran LJ & Teede HJ 2015 Metformin
and lifestyle modification in polycystic ovary syndrome: systematic review and meta-analysis.
*Hum Reprod Update* **21** 560-574.

Nelson-Degrave VL, Wickenheisser JK, Hendricks KL, Asano T, Fujishiro M, Legro RS,
Kimball SR, Strauss JF, 3rd & McAllister JM 2005 Alterations in mitogen-activated protein
kinase kinase and extracellular regulated kinase signaling in theca cells contribute to
excessive androgen production in polycystic ovary syndrome. *Mol Endocrinol* **19** 379-390.

Orostica L, Astorga I, Plaza-Parrochia F, Vera C, Garcia V, Carvajal R, Gabler F, Romero C
& Vega M 2016 Proinflammatory environment and role of TNF-alpha in endometrial function
of obese women having polycystic ovarian syndrome. *Int J Obes (Lond)* **40** 1715-1722.

Palomba S, de Wilde MA, Falbo A, Koster MP, La Sala GB & Fauser BC 2015 Pregnancy
complications in women with polycystic ovary syndrome. *Hum Reprod Update* **21** 575-592.

Palomba S, Russo T, Orio F, Jr., Falbo A, Manguso F, Cascella T, Tolino A, Carmina E,
Colao A & Zullo F 2006 Uterine effects of metformin administration in anovulatory women
with polycystic ovary syndrome. *Hum Reprod* **21** 457-465.
Patel B, Elguero S, Thakore S, Dahoud W, Bedaiwy M & Mesiano S 2015 Role of nuclear
progesterone receptor isoforms in uterine pathophysiology. *Hum Reprod Update* **21** 155-173.
Piltonen TT 2016 Polycystic ovary syndrome: Endometrial markers. *Best Pract Res Clin*
*Obstet Gynaecol* **37** 66-79.
Piltonen TT, Chen J, Erikson DW, Spitzer TL, Barragan F, Rabban JT, Huddleston H, Irwin
JC & Giudice LC 2013 Mesenchymal stem/progenitors and other endometrial cell types from
women with polycystic ovary syndrome (PCOS) display inflammatory and oncogenic
potential. *J Clin Endocrinol Metab* **98** 3765-3775.
Piltonen TT, Chen JC, Khatun M, Kangasniemi M, Liakka A, Spitzer T, Tran N, Huddleston
H, Irwin JC & Giudice LC 2015 Endometrial stromal fibroblasts from women with polycystic
ovary syndrome have impaired progesterone-mediated decidualization, aberrant cytokine
profiles and promote enhanced immune cell migration in vitro. *Hum Reprod* **30** 1203-1215.
Poretsky L, Clemons J & Bogovich K 1992 Hyperinsulinemia and human chorionic
gonadotropin synergistically promote the growth of ovarian follicular cysts in rats.
*Metabolism* **41** 903-910.
Quezada S, Avellaira C, Johnson MC, Gabler F, Fuentes A & Vega M 2006 Evaluation of
steroid receptors, coregulators, and molecules associated with uterine receptivity in secretory
endometria from untreated women with polycystic ovary syndrome. *Fertil Steril* **85** 1017-
1026.
Rosenfield RL & Ehrmann DA 2016 The Pathogenesis of Polycystic Ovary Syndrome
(PCOS): The Hypothesis of PCOS as Functional Ovarian Hyperandrogenism Revisited.
*Endocr Rev* **37** 467-520.
Savaris RF, Groll JM, Young SL, DeMayo FJ, Jeong JW, Hamilton AE, Giudice LC &
Lessey BA 2011 Progesterone resistance in PCOS endometrium: a microarray analysis in
clomiphene citrate-treated and artificial menstrual cycles. *J Clin Endocrinol Metab* **96** 1737-
1746.
Shao R, Cao S, Wang X, Feng Y & Billig H 2014a The elusive and controversial roles of
estrogen and progesterone receptors in human endometriosis. *Am J Transl Res* **6** 104-113.
Shao R, Li X, Feng Y, Lin JF & Billig H 2014b Direct effects of metformin in the
endometrium: a hypothetical mechanism for the treatment of women with PCOS and
endometrial carcinoma. *J Exp Clin Cancer Res* **33** 41.
Tapia-Pizarro A, Archiles S, Argandona F, Valencia C, Zavaleta K, Cecilia Johnson M,
Gonzalez-Ramos R & Devoto L 2017 hCG activates Epac-Erk1/2 signaling regulating
Progesterone Receptor expression and function in human endometrial stromal cells. *Mol Hum*
*Reprod* **23** 393-405.
Thienel T, Chwalisz K & Winterhager E 2002 Expression of MAPkinases (Erk1/2) during
decidualization in the rat: regulation by progesterone and nitric oxide. *Mol Hum Reprod* **8**
465-474.
Tranguch S, Wang H, Daikoku T, Xie H, Smith DF & Dey SK 2007 FKBP52 deficiency-
conferred uterine progesterone resistance is genetic background and pregnancy stage specific.
*J Clin Invest* **117** 1824-1834.
Vrbikova J & Cibula D 2005 Combined oral contraceptives in the treatment of polycystic
ovary syndrome. *Hum Reprod Update* **11** 277-291.
Wira CR, Fahey JV, Sentman CL, Pioli PA & Shen L 2005 Innate and adaptive immunity in
female genital tract: cellular responses and interactions. *Immunol Rev* **206** 306-335.

Yang Z, Wolf IM, Chen H, Periyasamy S, Chen Z, Yong W, Shi S, Zhao W, Xu J, Srivastava
A *et al.* 2006 FK506-binding protein 52 is essential to uterine reproductive physiology
controlled by the progesterone receptor A isoform. *Mol Endocrinol* **20** 2682-2694.
Zhang Y, Hu M, Meng F, Sun X, Xu H, Zhang J, Cui P, Morina N, Li X, Li W *et al.* 2017
Metformin Ameliorates Uterine Defects in a Rat Model of Polycystic Ovary Syndrome.
*EBioMedicine* **18** 157-170.
Zhang Y, Meng F, Sun X, Sun X, Hu M, Cui P, Vestin E, Li X, Li W, Wu XK *et al.* 2018
Hyperandrogenism and insulin resistance contribute to hepatic steatosis and inflammation in
female rat liver. *Oncotarget* **In Press**.
Zhang Y, Sun X, Sun X, Meng F, Hu M, Li X, Li W, Wu XK, Brännström M, Shao R *et al.*
2016 Molecular characterization of insulin resistance and glycolytic metabolism in the rat
uterus. *Sci Rep* **6** 30679.
Zhao Y, Zhang C, Huang Y, Yu Y, Li R, Li M, Liu N, Liu P & Qiao J 2015 Up-regulated
expression of WNT5a increases inflammation and oxidative stress via PI3K/AKT/NF-kappaB
signaling in the granulosa cells of PCOS patients. *J Clin Endocrinol Metab* **100** 201-211.

**Figure legends**

**Figure 1. Chronic treatment with metformin alters PGR isoform protein expression and**

**PGR-target gene expression in the rat uterus *in vivo*.** A, Western blot analysis of protein

expression in the rat uterus was performed. Representative images and quantification of the

densitometric data (n = 8–9/group) of PGR isoforms are shown. B, Immunohistochemical

detection of PGRA/B in control rats treated with saline or metformin and in insulin+hCG-

treated rats treated with saline or metformin. Representative images (n = 5/group) are shown.

Lu, lumen; Le, luminal epithelial cells; Ge, glandular epithelial cells; Str, stromal cells. Scale

bars (100 µm) are indicated in the photomicrographs. High magnification images are shown

in the bottom panels. C, Uterine tissues from control rats treated with vehicle or metformin

and insulin+hCG-treated rats treated with saline or metformin (n = 6/group) were analyzed for

mRNA levels of *Ihh*, *Ptch*, *Smo*, *Nr2f2*, *Hand2*, *Fkbp52*, and *Ncoa2* by qRT-PCR. The

mRNA level of each gene relative to the mean of the sum of the *Gapdh* and *U87* mRNA

levels in the same sample is shown. Values are expressed as means ± SEM. Statistical tests

are described in the Material and Methods. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

**Figure 2. Chronic treatment with metformin alters PGR isoform protein expression and**

**PGR-target gene expression in the rat uterus after implantation.** A, Western blot analysis

of protein expression in the rat uterus was performed. Representative images and

quantification of the densitometric data of PR isoforms are shown (n = 5–7/group). B,

Immunohistochemical detection of PGRA/B in the uterine implantation and inter-implantation

sites. Representative images are shown (n = 5/group). DS, decidualized stroma; Lu, lumen;

Le, luminal epithelial cells; Ge, glandular epithelial cells; Str, stromal cells. Scale bars (100

µm) are indicated in the photomicrographs. C, Uterine tissues (n = 5–6/group) were analyzed

for mRNA levels of *Ihh*, *Ptch*, *Smo*, *Nr2f2*, *Hand2*, *Fkbp52*, and *Ncoa2* by qRT-PCR. The

mRNA level of each gene is shown relative to the mean of the sum of the *Gapdh* and *U87*

mRNA levels in the same sample. Values are expressed as means ± SEM. Statistical tests are

described in the Materials and Methods. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

**Figure 3. Specific regulation of uterine PGR isoforms, PGR-targeted and implantation-**

**related gene expression by metformin treatment *in vitro*.** Quantitative RT-PCR analysis of

(A) *Pgr*, *Pgrb*, *Ihh*, *Ptch*, *Smo*, *Nr2f2*, *Hand2*, *Fkbp52*, and *Ncoa2* and (B) *Prl*, *Igfbp1*, *Lif*,

*Il11*, *Pc6*, *Spp1*, *Maoa*, *Ednrb*, *Hoxa10*, *Hoxa11*, *Lrh1*, *Sgk1*, *Hbegf*, and *Krt13* mRNA levels

in rat uterine tissues treated with either saline or 10 mM metformin for the indicated culture

times (n = 6/group). mRNA levels were normalized to the average levels of *Gapdh* and *U87*

mRNA in the same sample. Values are expressed as means ± SEM. Statistical tests are

described in the Materials and Methods. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

**Figure 4. Specific regulation of uterine PGR isoforms and PGR-targeted gene expression**

**by treatment with P4 and/or RU486 *in vivo*.** Uterine tissues (n = 5–6/group) were analyzed

for mRNA levels of *Pgr*, *Pgrb*, *Ihh*, *Ptch*, *Smo*, *Nr2f2*, *Hand2*, *Fkbp52*, and *Ncoa2* by qRT-

PCR. mRNA levels were normalized to the average levels of *Gapdh* and *U87* mRNA in the

same sample. Values are expressed as means ± SEM. Statistical tests are described in the

Materials and Methods. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

**Figure 5. Chronic treatment with metformin alters the MAPK signaling pathway in the**

**rat uterus before and after implantation.** Western blot analysis of protein expression in the

rat uterus was performed. Representative images and quantification of the densitometric data

for p-c-Raf, p-MEK1/2, p-ERK1/2, ERK1/2, p-p38 MAPK, and p38 MAPK are shown (n =

6–9/group before implantation; n = 5–6/group after implantation). Values are expressed as

means ± SEM. Statistical tests are described in the Materials and Methods. * $p < 0.05$; ** $p <$

$0.01$.

**Figure 6. Chronic treatment with metformin alters ER subtype mRNA and protein**

**expression in the rat uterus *in vivo*.** A, Uterine tissues from control rats treated with saline

vehicle or metformin and insulin+hCG-treated rats treated with saline or metformin (n =

6/group) were analyzed for mRNA levels of *Esr1* (ERα) and *Esr2* (ERβ) by qRT-PCR. The

mRNA level of each gene relative to the mean of the sum of the *Gapdh* and *U87* mRNA

levels in the same sample is shown. Values are expressed as means ± SEM. Statistical tests

are described in the Material and Methods. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. B,

Immunofluorescence detection of ERα (red) and ERβ (green) in control rats treated with

saline (B1-2) or metformin (C1-2) and in insulin+hCG-treated rats treated with saline (D1-2)

or metformin (E1-2). Representative images are shown (n = 5/group). Cell nuclei were

counterstained with DAPI (blue, lower panel). Lu, lumen; Le, luminal epithelial cells; Ge,

glandular epithelial cells; Str, stromal cells. Scale bars (100 μm) are indicated in the

photomicrographs.

**Figure 7. Chronic treatment with metformin alters ER subtype mRNA and protein**

**expression in the rat uterus after implantation.** A, Uterine tissues from control rats treated

with saline or metformin and insulin+hCG-treated rats treated with saline or metformin (n =

5/group) were analyzed for mRNA levels of *Esr1* (ERα) and *Esr2* (ERβ) by qRT-PCR. The

mRNA level of each gene relative to the mean of the sum of the *Gapdh* and *U87* mRNA

levels in the same sample is shown. Values are expressed as means ± SEM. Statistical tests

are described in the Materials and Methods. * $p < 0.05$; ** $p < 0.01$. B, Immunofluorescence

detection of ERα (red) and ERβ (green) in control rats treated with saline (B1-2) or metformin

(C1-2) and in insulin+hCG-treated rats treated with metformin with implantation (D1-2) or

without implantation (E1). Representative images are shown (n = 5/group). Cell nuclei were

counterstained with DAPI (blue, lower panel). DS, decidualized stroma; Lu, lumen; Le,

luminal epithelial cells; Ge, glandular epithelial cells; Str, stromal cells. Scale bars (100 μm)

are indicated in the photomicrographs.

**Figure 8.** Schematic representation of the actions of metformin on the uterine progesterone

signaling in the PCOS-like rats. Note that bold symbol indicates metformin-regulated genes

and proteins and asterisk indicates that treatment with metformin or progesterone share the

same targeted genes and proteins.

**Supplementary Figure 1. Schematic representation of experimental groups and**

**treatment protocol.** n, the final number of animals.

**Supplementary Figure 2. Photographs and illustration of the procedures for rat uterine**

**tissue preparation and culture with and without metformin.** Culture media with and

without metformin treatment were changed daily.

**Supplementary Figure 3. Effects of metformin on uterine structure in rats after**

**implantation.** Female rats were mated individually with fertile male rats according to their

estrous cycle stage. Representative photomicrographs of uteri collected on day 6 after

pregnancy with implantation sites visualized by Chicago Blue B dye injection. Because the

insulin+hCG-treated rats without a normal estrous cycle display implantation failure, these

rats treated with saline were excluded from the analysis. Higher-magnification images of the

different areas are shown on the rightmost three panels of each row. MD, mesometrial

decidua; AD, antimesometrial decidua; En, endometrium; Myo, myometrium. Scale bars (100

μm) are indicated in the photomicrographs.

**Supplementary Figure 4. Chronic treatment with metformin alters p-histone H3 (Ser10)**

**protein expression in the rat uterus *in vivo*.** Immunofluorescence detection of p-histone H3

(red) in control rats treated with saline (A1-4) or metformin (B1-4) and in insulin+hCG-

treated rats treated with saline (C1-4) or metformin (D1-4). Cell nuclei were counterstained

with DAPI (blue). Lu, lumen; Le, luminal epithelial cells; Ge, glandular epithelial cells; Str,

stromal cells; M, myometrium. Scale bars (100 μm) are indicated in the photomicrographs.

The number of all p-histone H3-positive cells in whole luminal epithelial cells, glandular

epithelial/stromal cells and myometrium)/uterine section/animal was counted. Semi-

quantitative immunofluorescence of p-histone H3 (Q-H score) in the different uterine cell

types (n = 5/group) is shown in E. Values are expressed as means ± SEM. Statistical tests are

described in the Material and Methods. * $p < 0.05$.

**Supplementary Figure 5. Chronic treatment with metformin alters p-histone H3 (Ser10)**

**protein expression in the rat uterus after implantation.** Immunofluorescence detection of

p-histone H3 (red) in control rats treated with saline (A1-4) or metformin (B1-4) and in

insulin+hCG-treated rats treated with metformin with implantation (C1-4) or without

implantation (D1-2). Cell nuclei were counterstained with DAPI (blue). DS, decidualized

stroma; Le, luminal epithelial cells; Ge, glandular epithelial cells; Str, stromal cells; M,

myometrium. Scale bars (100 μm) are indicated in the photomicrographs. The number of all

p-histone H3-positive cells in whole luminal epithelial cells, glandular epithelial/stromal cells

and myometrium)/uterine section/animal was counted. Semi-quantitative immunofluorescence

of p-histone H3 (Q-H score) in the different uterine cell types is shown in E. Uterine tissues

were analyzed (n = 8/group; n = 5 for insulin+hCG-treated rats with metformin but without

implantation). Values are expressed as means ± SEM. Statistical tests are described in the

Materials and Methods. * $p < 0.05$; ** $p < 0.01$.

**A**

**B**   **PGRA/B**

**C**

**A**

**B** PGRA/B

**C**

# Figure 5

**A**

**B**

**A**

Fold changes in mRNA

Legend:
- □ Control + Saline
- ■ Control + Met
- ■ Ins + hCG + Saline (red)
- ■ Ins + hCG + Met (blue)

*Esr1*   *Esr2*

Control — Saline / Met

Ins + hCG — Saline / Met

B1 ERαERβ    C1    D1    E1
DAPI  Le  Lu  Ge  Ge  Str    Lu  Le  Ge  Str    Le  Lu  Ge  Str    Lu  Le  Str  Ge  Ge

B2 ERαERβ    C2    D2    E2
DAPI  Myo    Myo    Myo    Myo

Increased expression    Decreased expression    Unchanged expression

Suppl MM

**Materials and Methods**

*Primary in vitro tissue culture and treatment*

Uterine tissue culture and treatment was essentially carried out as described previously (Cui *et al.* 2015; Li *et al.* 2015). After rinsing in cold PBS, uterine tissues obtained from control (n = 5) and PCOS-like (n = 5) rats were divided into equal-sized explants and placed in 24-well tissue culture plates (Sarstedt, Newton, MA) containing RPMI-1640 medium with charcoal-stripped 10% fetal calf serum and 100 IU/ml penicillin/streptomycin (GIBCO-BRL, San Francisco, CA). Cultured tissues treated with sterile saline or metformin (10 mM in sterile saline) were incubated in a humidified incubator (37ºC, 95% $O_2$, 5% $CO_2$) and separately collected at 0 h, 24 h, 48 h, and 72 h after treatment (Supplemental Fig. 2). Each culture condition was performed in five replicates (five wells), and tissues from a minimum of five control rats and five PCOS-like (insulin+hCG-treated) rats were used. At the end of the experiments, cultured tissues were snap-frozen in liquid nitrogen and stored at –70°C.

*Morphological assessment and immunostaining*

Uterine tissues were fixed in 10% formalin, embedded in paraffin, and sectioned for hematoxylin and eosin staining according to standard procedures. Immunohistochemistry and immunofluorescence were performed according to previously described methods (Zhang *et al.* 2017; Zhang *et al.* 2016). The endogenous peroxidase and nonspecific binding were removed by incubation with 3% $H_2O_2$ for 10 min and with 10% normal goat serum for 1 h at room temperature. After incubation with the primary antibody (Supplemental Table 1) overnight at 4°C in a humidified chamber, the sections were stained using the avidin-biotinylated-peroxidase complex detection system (Vector Laboratories Inc., Burlingame, CA) followed by treatment with 3-amino-9-ethyl carbazole developing reagent plus High Sensitivity Substrate (SK-4200, Vector Laboratories). The sections were imaged on a Nikon E-1000

Suppl MM

microscope (Japan) and photomicrographed using Easy Image 1 (Bergström Instrument AB, Sweden).

The other half of the uterine sections were incubated with primary antibody in 0.01 M Tris-buffered saline supplemented with Triton X-100 (TBST) containing 5% nonfat milk overnight at 4°C, and a secondary antibody was applied at room temperature for 1 h. After the sections were washed with TBST, they were re-suspended in mounting medium containing DAPI (4′,6′-diamidino-2-phenylindole; Vector Laboratories). Sections were examined under an Axiovert 200 confocal microscope (Zeiss, Jena, Germany) equipped with a laser-scanning confocal imaging LSM 710 META system (Carl Zeiss) and photomicrographed. Background settings were adjusted from the examination of negative control specimens. Images of positive staining were adjusted to make optimal use of the dynamic range of detection. The immune staining was quantified by semi-quantitative histogram scoring (Q-H score) as described previously (Mariee *et al.* 2012). All morphological and immunohistochemical assays were performed by at least two researchers in an operator-blinded manner.

### *Protein isolation and Western blot analysis*

A detailed explanation of the Western blot analysis protocol has been published (Zhang *et al.* 2017; Zhang *et al.* 2016). Total protein was isolated from whole uterine tissue by homogenization in RIPA buffer (Sigma-Aldrich) supplemented with cOmplete Mini protease inhibitor cocktail tablets (Roche Diagnostics, Mannheim, Germany) and PhosSTOP phosphatase inhibitor cocktail tablets (Roche Diagnostics). After determining total protein by Bradford protein assay, equal amounts (30 μg) of protein were resolved on 4–20% TGX stain-free gels (Bio-Rad Laboratories GmbH, Munich, Germany) and transferred onto PVDF membranes. The membranes were probed with the primary antibody (Supplemental Table 1) in TBST containing 5% non-fat dry milk followed by HRP-conjugated secondary antibody. When necessary, the PVDF membranes were stripped using Restore PLUS Western blot

Suppl MM

stripping buffer (Thermo Scientific, Rockford, IL) for 15 minutes at room temperature, washed twice in TBST, and then re-probed. Ultraviolet activation of the Criterion stain-free gel on a ChemiDoc MP Imaging System (Bio-Rad) was used to control for proper loading. Band densitometry was performed using Image Laboratory (Version 5.0, Bio-Rad).

In the present study, we used a novel stain-free technology – ultraviolet activation of the Criterion stain-free gel on a ChemiDoc MP Imaging System (Bio-Rad) – which was used to control proper loading (total protein normalization). This technology represents a significant advancement over existing stain-based total protein quantitation approaches such as Coomassie blue, Ponceau S, and others and gives accurate protein loading control data in a standardized manner without requiring lengthy optimization. The detection of proteins on stain-free gels is based upon trihalocompound modification of tryptophan residues in proteins run on stain-free gels, which are exposed to UV. The modified tryptophans give a fluorescent signal that can be readily detected by a CCD camera. In our study, band densitometry was performed using Image Laboratory. When quantified, the intensity of each protein band was normalized to the total protein in individual sample. This method has been commonly used in both human and animal tissues to semi-quantify concentrations of specific proteins in many studies. Please visit the link given below for review of previous publications using the same method and technology:

(http://www.biorad.com/webroot/web/pdf/lsr/literature/Bulletin_6351.pdf).

### *RNA extraction and qRT-PCR analysis*

For RNA isolation, tissues from each rat were lysed using TRIzol Reagent (Life Technologies), and RNA was isolated following standard protocols. qRT-PCR was performed with a Roche Light Cycler 480 sequence detection system (Roche Diagnostics Ltd., Rotkreuz, Switzerland) as previously described (Zhang *et al.* 2017; Zhang *et al.* 2016). The PCR amplifications were performed with a SYBR green qPCR master mix (#K0252, Thermo

Suppl MM

Scientific, Rockford, IL). Total RNA was prepared from the frozen whole uterine tissues, and single-stranded cDNA was synthesized from each sample (2 μg) with M-MLV reverse transcriptase (#0000113467, Promega Corporation, Fitchburg, WI) and RNase inhibitor (40 U) (#00314959, Thermo Scientific). cDNA (1 μl) was added to a reaction master mix (10 μl) containing 2× SYBR green qPCR reaction mix (Thermo Scientific) and gene-specific primers (5 μM each of forward and reverse primers). All primers are indicated in Supplemental Table 2. All reactions were performed six times, and each reaction included a non-template control. The results for target genes were expressed as the amount relative to the average CT values of *GAPDH + U87* in each sample. Relative gene expression was determined with the $2^{-\Delta\Delta CT}$ method, and the efficiency of each reaction – as determined by linear regression – was incorporated into the equation.

## *Measurement of biochemical parameters*

Concentrations of gonadotropins (follicle stimulating hormone and luteinizing hormone), steroid hormones (17β-estradiol, progesterone, testosterone, 5α-dihydrotestosterone, and androstenedione), sex hormone-binding globulin, glucose, and insulin in rat serum samples were measured using commercially available assays (Cloud-Clone Corp., Houston, TX) as described previously (Zhang et al. 2017). All samples and standards were measured in duplicate. The intra- and inter-assay coefficients of variation are listed in Supplemental Table 3.

## References

Cui P, Li X, Wang X, Feng Y, Lin JF, Billig H & Shao R 2015 Lack of cyclical fluctuations of endometrial GLUT4 expression in women with polycystic ovary syndrome: Evidence for direct regulation of GLUT4 by steroid hormones. *BBA Clinical* **4** 85-91. (https:// doi: 10.1016/j.bbacli.2015.08.004)

Li X, Cui P, Jiang HY, Guo YR, Pishdari B, Hu M, Feng Y, Billig H & Shao R 2015 Reversing the reduced level of endometrial GLUT4 expression in polycystic ovary syndrome:

Suppl MM

a mechanistic study of metformin action. *American Journal of Translational Research* **7** 574-586.

Mariee N, Li TC & Laird SM 2012 Expression of leukaemia inhibitory factor and interleukin 15 in endometrium of women with recurrent implantation failure after IVF; correlation with the number of endometrial natural killer cells. *Human Reproduction* **27** 1946-1954. (https://doi: 10.1093/humrep/des134)

Zhang Y, Hu M, Meng F, Sun X, Xu H, Zhang J, Cui P, Morina N, Li X, Li W, et al. 2017 Metformin Ameliorates Uterine Defects in a Rat Model of Polycystic Ovary Syndrome. *EBioMedicine* **18** 157-170. (https://doi: 10.1016/j.ebiom.2017.03.023)

Zhang Y, Sun X, Sun X, Meng F, Hu M, Li X, Li W, Wu XK, Brännström M, Shao R, et al. 2016 Molecular characterization of insulin resistance and glycolytic metabolism in the rat uterus. *Scientific Report* **6** 30679. (https://doi: 10.1038/srep30679)

**Supplementary Table 1.** Antibodies: species, clone/catalog number, method, dilution, and source.

| Antibody | Species | Cat. No. | kDa | Method | Dilution | Source |
|---|---|---|---|---|---|---|
| PGR | Mouse | MA1-410 | B120 | WB | 1:500 | Thermo Fisher (Rockford, IL) |
| | | | A94 | IHC | 1:50 | |
| p-c-Raf | Rabbit | 9427 | 74 | WB | 1:500 | Cell Signaling Technology (Danver, MA) |
| p-MEK1/2 | Rabbit | 9154 | 45 | WB | 1:500 | Cell Signaling Technology |
| p-ERK1/2 | Rabbit | 4370 | 42,44 | WB | 1:1000 | Cell Signaling Technology |
| ERK1/2 | Rabbit | 4695 | 42,44 | WB | 1:2000 | Cell Signaling Technology |
| p-p38 MAPK | Rabbit | 4511 | 43 | WB | 1:300 | Cell Signaling Technology |
| p38 MAPK | Rabbit | 8690 | 40 | WB | 1:1000 | Cell Signaling Technology |
| p-JNK | Rabbit | 4668 | 46,54 | WB | 1:500 | Cell Signaling Technology |
| JNK | Rabbit | 9252 | 46,54 | WB | 1:1000 | Cell Signaling Technology |
| ERα | Mouse | 6F11 | | IHC, IF | 1:50 | Novocastra Laboratories Ltd. (Newcastle upon Tyne, UK) |
| ERβ | Rabbit | Ab3577 | | IF | 1:50 | Abcam (Cambridge, UK) |
| p-histone H3 (Ser10) | Rabbit | 3377 | | IF | 1:100 | Cell Signaling Technology |

PGR, progesterone receptor; p-MEK1/2, phosphorylation-MAP2K1/2; ERK1/2, extracellular signal-regulated kinases 1/2; p38 MAPK, p38 mitogen-activated protein kinase; JNK, jun-amino-terminal kinase; ERα, estrogen receptor alpha; WB, western blot; IHC, immunohistochemistry; IF, immunofluorescence.

**Supplementary Table 2.** Sequences of primer pairs used for qRT-PCR measurement.

| Gene | | Primer | Size |
|---|---|---|---|
| *Ihh* | Forward | CCTCGTAAACTCGTGCCTCT | 222 bp |
| | Reverse | CAGTGAGTTCAGACGGTCCT | |
| *Ptch* | Forward | GAACAAGCAACTTCCCCAAA | 190 bp |
| | Reverse | AATGTCGATGGGCTTGTCTC | |
| *Smo* | Forward | CAATGCGTGTTTCTTTGTGG | 253 bp |
| | Reverse | CGAGAGAGGCTGGTAAGTGG | |
| *Nr2f2* | Forward | GTCGCCTTTATGGACCACAT | 145 bp |
| | Reverse | CGTGGGCTACATCAGACAGA | |
| *Hand2* | Forward | CAGCTACATCGCCTACCTCA | 162 bp |
| | Reverse | TTCTTGTCGTTGCTGCTCAC | |
| *Fkbp5* | Forward | GCTGCCATCGAAAGCTGTAA | 102 bp |
| | Reverse | GTCAAAGTCATTCACGGCCA | |
| *Ncoa2* | Forward | GCGAATGTCACAGAGCACTT | 238 bp |
| | Reverse | ACTGCCAATCATTCCTGTGC | |
| *Pgr* | Forward | GGTCTAAGTCTCTGCCAGGTTTCC | 182 bp |
| | Reverse | CAACTCCTTCATCCTCTGCTCATTC | |
| *Pgrb* | Forward | GCATCGTCTGTAGTCTCGCCAATAC | 176 bp |
| | Reverse | GCTCTGGGATTTCTGCTTCTTCG | |
| *Prl* | Forward | CAAGAAGAAGGGGCCAACCT | 100 bp |
| | Reverse | CTGGTGGTGACTGTCCCTTC | |
| *Igfbp1* | Forward | TGTACTAGAACCTGCCGCAC | 126 bp |
| | Reverse | AGCAGCTGTTCCTCTGTCAT | |
| *Lif* | Forward | CGCCCAACATGACGGATTTC | 226 bp |
| | Reverse | TTGTTGCACAGACGGCAAAG | |
| *Il11* | Forward | GACTCCCTACCTACCTTGGC | 100 bp |
| | Reverse | GCAACCACTGTACATGTCGG | |

| Gene | | Primer | Size |
|---|---|---|---|
| Pc6 | Forward | GTGTGAGAATGGGTCGGAGA | 149bp |
| | Reverse | TTTCTTTCCACTTTCGGCCG | |
| Spp1 | Forward | CTGAAGCCTGACCCATCTCA | 143 bp |
| | Reverse | TCGTCGTCATCATCGTCCAT | |
| Maoa | Forward | AGGAGCTAGGCATAGAGACCT | 219 bp |
| | Reverse | TCTTGTCCCATTCCTGAGCG | |
| Ednrb | Forward | GGGTCTGCATGCTTAATCCC | 212 bp |
| | Reverse | CTTGGCCACTTCTCGTCTCT | |
| Hoxa10 | Forward | TCCGAAAACAGTAAAGCCTCTC | 127 bp |
| | Reverse | GCGTCTGGTGCTTCGTGTAA | |
| Hoxa11 | Forward | GACTCCCTACCTACCTTGGC | 100 bp |
| | Reverse | GCAACCACTGTACATGTCGG | |
| Lrh1 | Forward | CTGTGAAAGCTGCAAGGGTT | 154 bp |
| | Reverse | CAGCTTCATTCCAACGTCGA | |
| Sgk1 | Forward | GGGCCTTCACTTCTCTTTCC | 163 bp |
| | Reverse | GTGCAGATAACCCAAGGCAC | |
| Hbegf | Forward | GCTCTTCCACCTGGCTCAAT | 120 bp |
| | Reverse | CACAACCCACCCTGGGATAC | |
| Krt13 | Forward | GTTTCGGAGCTGGTTCTTGC | 277 bp |
| | Reverse | AGGCGGTCATTGAGGTTCTG | |
| Esr1 | Forward | ACGCTCTGCCTTGATCACAC | 132 bp |
| | Reverse | CCTGCTGGTTCAAAAGCGTC | |
| Esr2 | Forward | AAAGTAGCCGGAAGCTGACA | 138 bp |
| | Reverse | GCCTGACGTGAGAAAGAAGC | |
| Gapdh | Forward | TCTCTGCTCCTCCCTGTTCTA | 121 bp |
| | Reverse | GGTAACCAGGCGTCCGATAC | |
| U87 | Forward | CCAGGTGCAACAAAACCTGT | 188 bp |
| | Reverse | GCTGGACCCAAAACAACGAG | |

*Pgr*, progesterone receptor; *Ihh*, Indian hedgehog; *Ptch*, patched; *Smo*, smoothened frizzled class receptor; *Nr2f2*, nuclear receptor subfamily 2 group F member 2; *Hand2*, heart and neural crest derivatives expressed 2; *Fkbp5*, FK506 binding protein 5; *Ncoa2*, nuclear receptor coactivator 2; *Pgr*, progesterone receptor; *Pgrb*, progesterone receptor B; *Prl*, prolactin; *Igfbp1*, insulin-like growth factor binding protein 1; *Lif*, leukemia inhibitory factor; *Pc6*, subtilisin/kexin type 6; *Spp1*, osteopontin/secreted phosphoprotein 1; *Maoa*, monoamine oxidase; *Ednrb*, endothelin receptor B; *Hoxa10*, homeobox A10; *Lrh1*, liver receptor homolog-1; *Sgk1*, serum- and glucocorticoid-regulated kinase; *Hbegf*, heparin-binding EGF-like growth factor; *Krt13*, keratin 13; *Esr1*, estrogen receptor 1; *Gapdh*, glyceraldehyde-3-phosphate dehydrogenase; *U87*, small nucleolar RNA, C/D box 87(Snord87).

**Supplementary Table 3.** The intra-assay and inter-assay % CV for the rat gonadotropins, steroid hormones, and insulin.

|         | Intra-assay % CV | Inter-assay % CV |
|---------|------------------|------------------|
| FSH     | 6.4              | 6.8              |
| LH      | 6.4              | 6.6              |
| E2      | 6.5              | 6.8              |
| P4      | 6.4              | 6.6              |
| Total T | 6.2              | 6.6              |
| A4      | 6.7              | 6.9              |
| DHT     | 6.2              | 6.7              |
| SHBG    | 6.4              | 6.8              |
| Insulin | 6.3              | 6.6              |

FSH, follicle-stimulating hormone; LH, luteinizing hormone; E2, 17β-estradiol; P4, progesterone; T, testosterone; A4, androstenedione; DHT, 5α-dihydrotestosterone; SHBG, sex hormone-binding globulin.

**Supplementary Table 4.** Effects of implantation on the endocrine and metabolic alterations in control rats treated with saline or metformin, and insulin+hCG-treated (PCOS-like) rats with metformin.

| | Saline (n = 8) | Met (n = 10) | Insulin + hCG + Met (implantation) (n = 10) | Insulin + hCG + Met (no implantation) (n = 5) |
|---|---|---|---|---|
| FSH (ng/ml) | 2.13 ± 0.04 | 2.06 ± 0.04 | 2.14 ± 0.04 | 2.14 ± 0.10 |
| LH (ng/ml) | 2.00 ± 0.24 | 1.93 ± 0.11 | 2.31 ± 0.17 | 2.73 ± 0.36 |
| LH / FSH | 0.94 ± 0.11 | 0.94 ± 0.05 | 1.08 ± 0.08 | 1.28 ± 0.18 |
| E2 (ng/ml) | 1.31 ± 0.04 | 1.34 ± 0.03 | 1.23 ± 0.03 | 1.32 ± 0.05 |
| P4 (ng/ml) | 6.88 ± 0.35 | 6.51 ± 0.37 | 8.95 ± 0.44 [a, c] | 4.80 ± 0.11 [b, d, e] |
| Total T (ng/ml) | 5.11 ± 0.71 | 3.83 ± 0.22 | 3.78 ± 0.44 | 7.40 ± 0.09 [a, c, e] |
| A4 (ng/ml) | 0.142 ± 0.004 | 0.142 ± 0.006 | 0.162 ± 0.006 | 0.181 ± 0.016 [b, c] |
| DHT (pg/ml) | 102.97 ± 5.66 | 97.92 ± 9.28 | 102.00 ± 5.52 | 154.58 ± 8.39 [a, c, e] |
| E2 / Total T | 0.29 ± 0.04 | 0.36 ± 0.02 | 0.38 ± 0.05 | 0.18 ± 0.01 [d, f] |
| Total T / DHT | 0.051± 0.007 | 0.044 ± 0.006 | 0.038 ± 0.005 | 0.049 ± 0.003 |
| Total T / A4 | 36.37 ± 5.36 | 27.28 ± 1.77 | 23.26 ± 2.49 | 42.20 ± 3.52 [d, e] |
| SHBG (ng/ml) | 11.08 ± 0.33 | 10.89 ± 0.35 | 9.67 ± 1.16 | 10.91 ± 0.56 |
| Fasting glucose (mmol/l) | 4.19 ± 0.09 | 3.92 ± 0.10 | 4.53 ± 0.19 [d] | 4.98 ± 0.12 [a, c] |
| Fasting insulin (pg/ml) | 3.26 ± 0.05 | 3.17 ± 0.02 | 3.22 ± 0.04 | 3.42 ± 0.03 [b, c, f] |
| HOMA-IR | 0.61 ± 0.01 | 0.55 ± 0.01 | 0.65 ± 0.03 [c] | 0.76 ± 0.02 [a, c, f] |

Met, metformin; FSH, follicle-stimulating hormone; LH, luteinizing hormone; E2, 17α-estradiol; P4, progesterone; T, testosterone; A4, androstenedione; DHT, 5α-dihydrotestosterone; SHBG, sex hormone-binding globulin; HOMA-IR, Homeostasis model assessment of insulin resistance, HOMA-IR= fasting blood glucose (mmol/l) × fasting serum insulin (μU/ml) / 22.5.

Values are Mean ±SEM. The multiple comparisons between data were performed using one-way ANOVA and Tukey's post hoc test. A P-value less than 0.05 was considered statistically significant.

[a] $P < 0.01$ versus Saline group.

[b] $P < 0.05$ versus Saline group.

[c] $P < 0.01$ versus Met group.

[d] $P < 0.05$ versus Met group.

[e] $P < 0.01$ versus Insulin + hCG + Met (implantation) group.

[f] $P < 0.05$ versus Insulin + hCG + Met (implantation) group.

Inter-implantation site
Implantation site
Postimplantation

Control + Saline

A Implantation site
MD
AD
Myo

Intra-implantation site
En
Myo

Control + Metformin

B Implantation site
MD
AD
Myo

Inter-implantation site
Myo
En

(Ins + hCG) + Metformin

C Implantation site
AD
Myo
MD

Inter-implantation site
Myo
En

En
Myo

No implantation