Staphylococcus aureus wall teichoic acid is a pathogen-associated molecular

pattern that is recognized by langerin (CD207) on skin Langerhans cells

Rob van Dalen[1], Jacinto S. De La Cruz Diaz[2], Matevž Rumpret[1,#], Felix F. Fuchsberger[1,3,#],

Nienke H. van Teijlingen[4], Jonas Hanske[3], Christoph Rademacher[3], Theunis B.H.

Geijtenbeek[4], Jos A.G. van Strijp[1], Christopher Weidenmaier[5,6], Andreas Peschel[5,6], Daniel H.

Kaplan[2], and Nina M. van Sorge[1,*]

[1] Medical Microbiology, University Medical Center Utrecht, Utrecht University; Heidelberglaan

100, 3584 CX Utrecht, The Netherlands

[2] Departments of Dermatology and Immunology, University of Pittsburgh; 200 Lothrop Street,

W1043 BST Pittsburgh, PA 15261, USA

[3] Department of Biomolecular Systems, Max Planck Institute of Colloids and Interfaces; Am

Mühlenberg 1, 14424 Potsdam, Germany

[4] Department of Experimental Immunology, Academic Medical Center, University of

Amsterdam; Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

[5] Interfaculty Institute of Microbiology and Infection Medicine, University of Tübingen; Auf der

Morgenstelle 28, 72076 Tübingen, Germany

[6] German Center for Infection Research (DZIF), Partnersite Tübingen; Tübingen, Germany

# Authors contributed equally

* Correspondence: nsorge3@umcutrecht.nl

Abstract

Staphylococcus aureus is a major cause of skin and soft tissue infections and aggravator of the inflammatory skin disease atopic dermatitis (AD). Epicutaneous exposure to *S. aureus* induces Th17 responses through skin Langerhans cells (LCs), which paradoxically contribute to host defense but also to AD pathogenesis. The underlying molecular mechanisms of the association between *S. aureus* and skin inflammation are poorly understood. Here, we demonstrate that human LCs directly interact with *S. aureus* through the pattern-recognition receptor langerin (CD207). Human, but not mouse, langerin interacts with *S. aureus* through the conserved β-*N*-acetylglucosamine (GlcNAc) modifications on wall teichoic acid (WTA), thereby discriminating *S. aureus* from other staphylococcal species. Importantly, the specific *S. aureus* WTA glycoprofile strongly influences the level of Th1- and Th17-polarizing cytokines that are produced by *in vitro* generated LCs. Finally, in a murine epicutaneous infection model, *S. aureus* induced a more pronounced influx of inflammatory cells and pro-inflammatory cytokine transcripts in skin of human langerin transgenic mice compared to wild-type mice. Our findings provide molecular insight into the unique pro-inflammatory capacities of *S. aureus* in relation to inflammatory skin disease.

Keywords

Langerhans cell, langerin, *Staphylococcus aureus*, wall teichoic acid, glycosylation, atopic dermatitis

Introduction

The inflammatory skin disease atopic dermatitis (AD, also known as eczema) affects up to

20% of children and 3% of adults worldwide (1). An important characteristic of AD is the

disturbed microbiota composition with dominant presence of *Staphylococcus aureus*, but not

of other staphylococcal species (2,3). In particular, the *S. aureus* CC1 lineage is

overrepresented in AD isolates and was proposed to have particular yet-unidentified features

that enable colonization of AD skin (4). Langerhans cells (LCs) are key sentinel cells in the

skin epidermis and are implicated in *S. aureus*-induced skin inflammation. LCs are equipped

with a diverse set of pattern-recognition receptors (PRRs) to sense intruders, including the

LC-specific C-type lectin receptor (CLR) langerin (CD207) (5). LCs can phagocytose

microbes and initiate adaptive immune responses by activating skin-resident immune

memory cells or naïve immune cells in the lymph nodes (6,7). In response to *S. aureus*, LCs

induce Th17 responses that help to contain *S. aureus* infection but paradoxically also

aggravate AD (8,9). Despite the functional importance of LCs in *S. aureus*-mediated skin

pathology, the molecular interaction between LCs and *S. aureus* and the functional response

of LCs have received little attention.

A dominant and evolutionarily conserved component of the *S. aureus* surface is wall

teichoic acid (WTA), which is important in nasal colonization, *S. aureus*-induced endocarditis,

beta-lactam resistance and phage-mediated horizontal gene transfer (10-14). In the majority

of *S. aureus* lineages, WTA is composed of 20-40 ribitol phosphate (RboP) repeating units

modified with *D*-alanine and *N*-acetylglucosamine (GlcNAc). GlcNAc is linked to the

anomeric C4 of RboP in either α or β configuration by glycosyltransferases TarM and TarS,

respectively (12,15). Several *S. aureus* WTA glycoprofiles can be discriminated: WTA β-

GlcNAcylation is conserved in almost all *S. aureus* strains, whereas WTA α-GlcNAcylation is

only present in about one-third of the *S. aureus* isolates. A small selection of isolates even

completely lack WTA glycosylation (10,16). Finally, WTA of *S. aureus* lineage ST395 is

composed of a glycerol phosphate (GroP) backbone modified by *N*-acetylgalactosamine

(GalNAc) (14). WTA glycosylation is an important determinant in host-pathogen interactions,

which includes attachment to scavenger receptor SREC-1 in the nasal epithelium, and

opsonization by antibodies and mannose-binding lectin (17-19).

We demonstrate an important role of the PRR langerin in sensing the β-GlcNAc

epitope on *S. aureus* WTA, which explains the lack of binding to other non-AD associated

staphylococcal species. Interestingly, simultaneous decoration of WTA with α-GlcNAc

impairs langerin interaction and dampens cytokine responses of LCs, implying that *S. aureus*

can modulate immune detection and subsequent inflammation in the epidermis. Murine

infection experiments confirmed that langerin contributes to enhanced skin inflammation. In

conclusion, we identify WTA as a pathogen-associated molecular pattern (PAMP) of *S.*

aureus, which is recognized by langerin on LCs.

Results

Langerin is a receptor for *S. aureus* on human LCs

The molecular interaction between LCs and *S. aureus* has received little attention. We

therefore investigated whether LCs and *S. aureus* interact directly by incubating primary LCs

isolated from human skin with GFP-expressing *S. aureus*. LCs from four different donors

bound *S. aureus* in a dose-dependent manner (Figure 1A). The levels at which the

interaction was saturated varied between the donors from approximately 40% (donor 1) to

80% (donor 3) of *S. aureus*-positive LCs. To investigate the nature of interacting receptors on

LCs, we pre-incubated LCs with mannan, a ligand for many PRRs of the CLR family.

Depending on the bacteria-to-cell ratio, *S. aureus* binding was reduced by 35-70% compared

to non-blocking conditions in all donors (Figure 1A). Similarly, the interaction was inhibited by

approximately 35% by pre-incubation of the LCs with the monosaccharide GlcNAc (Figure

1A). Langerin is a mannan- and GlcNAc-specific CLR that is exclusively expressed on LCs.

We therefore investigated whether langerin would be involved in interaction with *S. aureus*.

Indeed, pre-incubation with an anti-langerin blocking antibody reduced binding of (*spa* and

sbi-deficient, to prevent aspecific antibody binding) *S. aureus* in donors 3 and 4 by 25-50%

compared to control, depending on the infective dose (Figure 1A). To confirm involvement of

langerin in the interaction between *S. aureus* and LCs, we introduced langerin in the THP1

cell line, which normally does not express langerin. Transduction of langerin, but not of

empty vector (EV), conferred *S. aureus* binding to THP1 cells, which could be completely

inhibited by addition of mannan or anti-langerin blocking antibody (Figure 1B).

It was previously demonstrated that *S. aureus*-exposed LCs initiate T cell proliferation

(20). However, the functional response of LCs was not assessed. Therefore, we stimulated

MUTZ-3-derived LCs (muLCs), a well-established cell model for human LCs (21,22), with *S.*

aureus and measured muLC activation through expression of co-stimulatory molecules and

cytokine production after 24 hours. Indeed, muLCs upregulated expression of co-stimulatory

molecules CD80 and CD86 and produced significant amounts of IL-8 and IL-12p70 in a

dose-dependent response to *S. aureus* (Figure 1C). Together, these data demonstrate that

LCs respond to *S. aureus* and that langerin is an important innate PRR for *S. aureus* on

human LCs.

Langerin recognizes *S. aureus* in a *tarS*-dependent manner through the conserved

WTA β-GlcNAc epitope

To further investigate langerin interaction with staphylococci, we tested binding of a FITC-

labeled trimeric construct of the extracellular domain of human langerin (langerin-FITC) to a

broader collection of 18 *S. aureus* strains from 11 different clonal complexes, as well as

several coagulase-negative staphylococci (CoNS). Langerin-FITC bound to most tested *S.*

aureus strains but to none of the CoNS species (Figure 2A), indicating that langerin interacts

with a ligand that is specific for and highly conserved in *S. aureus*. The three tested S.

aureus strains that showed no or low-level binding of langerin-FITC (ED133, Lowenstein and

PS187; Figure 2A), differ from the other tested *S. aureus* strains in the structural composition

of WTA. ED133 and Lowenstein completely lack WTA GlcNAcylation, whereas PS187

belongs to the ST395 lineage that expresses GroP-GalNAc WTA (14,16,23). Given the high

density of WTA on the *S. aureus* surface and apparent correlation between langerin

interaction and WTA structure, we hypothesized that WTA GlcNAc modifications are likely

candidates for the interaction with langerin.

To test this hypothesis, we assessed binding of langerin-FITC to a panel of *S. aureus*

knockout strains, which lack glycosyltransferases TarM and TarS that are required to modify

WTA with α-GlcNAc and β-GlcNAc, respectively. Loss of both glycosyltransferases (Δ*tarMS*)

reduced langerin-FITC binding to *S. aureus* to background levels in three different *S. aureus*

backgrounds (Figures 2B and S1A, B), demonstrating that WTA GlcNAc is the target for

langerin. To investigate whether langerin specifically recognized either α-GlcNAc or β-

GlcNAc, we tested the individual TarM and TarS knockout strains as well as Δ*tarMS*

complemented with either *tarM* or *tarS* on an expression plasmid (Δ*tarMS* p*tarM* and Δ*tarMS*

p*tarS*). Langerin-FITC only bound to *S. aureus* strains that express β-GlcNAc, whereas α-

GlcNAc was dispensable for binding (Figures 2B and S1A, B). Similarly, langerin-FITC

binding to *S. aureus* strains 82086 and PS66, which are naturally deficient for WTA α-

GlcNAc, was completely abrogated in isogenic Δ*tarS* strains (Figure 2C). These results show

that langerin interacts with *S. aureus* in a *tarS*-dependent manner and provide the first

demonstration of an anomeric-specific interaction of a human innate receptor with a Gram-

positive surface polysaccharide.

Although α-GlcNAc is not the target of langerin, its presence influences the level of

langerin-FITC binding: mutant strains lacking *tarM* (Δ*tarM* and Δ*tarMS* p*tarS*) showed

significantly increased binding compared to wild-type (Figures 2B and S1A, B). Possibly,

enhanced binding results from loss of steric hindrance by α-GlcNAc, since chemical analysis

of the WTA composition by Kurokawa *et al.* suggests that WTA of strain RN4220 Δ*tarM* does

not have increased β-GlcNAcylation (18).

As *S. aureus* expresses many human-specific adhesion or immune evasion factors

(24), we investigated the interaction with murine langerin-FITC, which has 76% identity with

the human langerin-FITC construct (25). Binding of murine langerin-FITC to *S. aureus* was

detectable, but was 10 to 100-fold lower than human langerin (Figure S1C). The EC50 of

human langerin-FITC for *S. aureus* USA300 was 9.7 (8.3 – 11.3) µg/ml, while binding of

murine langerin-FITC was not yet saturated at 50 µg/ml. Despite low level and non-saturable

binding, murine langerin interaction with *S. aureus* could be blocked by addition of mannan

(data not shown), suggesting that the interaction is specific. Altogether, this indicates that the

langerin-*S. aureus* interaction has a certain degree of species-specificity.

***S. aureus* induces a Th1 and Th17 cytokine profile in LCs, which is affected by the**

WTA glycoprofile

Given the importance of langerin for interaction between *S. aureus* and LCs, we investigated

whether distinct WTA GlcNAc glycoprofiles influenced the muLC response at the level of co-

stimulatory molecules and cytokine expression. In line with our initial observations (Figure

1D), stimulation of muLCs with wild-type *S. aureus* upregulated expression of activation

markers CD80, CD83 and CD86 (Figure 3A). Stimulation with β-GlcNAc-deficient *S. aureus*

Δ*tarS* reduced expression of these markers compared to wild-type, whereas stimulation with

α-GlcNAc deficient *S. aureus* Δ*tarM* enhanced expression (Figure 3A). In addition, muLCs

secreted significant levels of cytokines IL-6, IL-8 IL-12p70, IL-23p19 and TNFα (Figure 3B),

but not anti-inflammatory cytokine IL-10, in response to *S. aureus*. Cytokine levels were

significantly reduced after stimulation of muLCs with *S. aureus* Δ*tarS* compared to WT,

whereas stimulation with *S. aureus* Δ*tarM* significantly enhanced secretion of these cytokines

(Figure 3B). Interestingly, the level of muLC activation correlated with the interaction levels of

recombinant langerin-FITC to *S. aureus* WT, Δ*tarM* and Δ*tarS* strains (Figure 2B). These

data suggest that the previously described Th1 and Th17-polarizing response initiated by

LCs in response to *S. aureus* is strongly influenced by the specific glycoprofile of WTA.

Human langerin transgenic mice show enhanced inflammation to epicutaneous *S.*

***aureus* infection**

Given the observed species specificity of langerin for *S. aureus* WTA β-GlcNAc (Figure S1A),

we used human langerin - diphtheria toxin receptor (huLangerin-DTR) mice, which

constitutively express human langerin on mouse LCs, as a huLangerin transgenic mouse

model (26). Wild-type (WT) and huLangerin mice were epicutaneously inoculated with 1 x

10^7 colony forming units (CFU) of *S. aureus* Δ*tarM* (Figure S2A) (8,27). These genetically

stable mutant bacteria are unable to modulate WTA glycosylation through regulation of *tarM*,

thereby maximizing the interaction with human langerin. At the time of sacrifice and skin

collection (40 hours post-infection), the lesions of the huLangerin mice were clinically

different from those of WT mice (Figure S2B), although bacterial burden in the skin did not

differ between the groups (Figure S2C). Histological examination of the skin revealed more

extensive influx of inflammatory cells in the dermis of huLangerin mice compared to WT mice

(Figure 4A). Correspondingly, we observed significantly higher expression of the mouse IL-8

homolog CXCL1 (KC), but not of CXCL2 (MIP-2), in the huLangerin group as opposed to WT

controls (Figures 4B and S2D). In addition, we determined the transcript levels of the Th17

cytokines IL-6 and IL-17 and the anti-inflammatory cytokine IL-10. Though not significant,

both IL-6 and IL-17 showed a trend towards higher production in the huLangerin group, while

IL-10 was not induced in either group, corroborating the observed *in vitro* responses of

muLCs to *S. aureus* stimulation (Figures 4B and S2D). These results provide a first *in vivo*

demonstration of the involvement of human langerin in the skin immune response to *S.*

aureus.

Discussion

Despite the emerging role of LCs in *S. aureus*-mediated skin inflammation, there is limited

information on the molecular pathways and functional consequences of LC - *S. aureus*

interaction. We observe that LCs respond to *S. aureus* with a Th1 and Th17-polarizing

cytokine response, which corroborates findings by others, who have demonstrated that LCs

internalize *S. aureus* and subsequently polarize T cells towards Th17 (8,9,20,28).

Furthermore, we elucidate that detection of *S. aureus* WTA β-GlcNAc is of critical importance

for the induced cytokine response and can also be modified by co-decoration with α-GlcNAc,

a characteristic of approximately one-third of *S. aureus* isolates. The ability of *S. aureus* to

carefully regulate its WTA glycoprofile was also previously suggested in the context of lytic

podophage infection (16). Likely, *tarM* is regulated as part of the GraRS regulon, known to

control *S. aureus* susceptibility to antimicrobial defenses (29,30). However, whether and how

GraRS and WTA GlcNAcylation are affected during skin colonization remains to be

determined.

In addition to regulation of glycosylation, WTA abundance can be regulated through

tarH, the ATPase required for WTA transport across the membrane (31). High WTA

expression increases the ability to induce skin abscesses in mice (31). These results cannot

be compared directly to our study, since the mice were infected subcutaneously, thereby

bypassing the LCs. In addition, the species specificity of langerin should be taken into

account. We demonstrate that mouse langerin shows significantly reduced binding to *S.*

aureus compared to human langerin, underlining previous studies that reported differences in

ligand specificity of these orthologs (25).

LCs and langerin were previously implicated in host defense against various other

pathogens. LCs internalize and degrade HIV-1 viral particles in a langerin-dependent manner

to prevent infection of deeper layers of the mucosa (32,33). Langerin has also been identified

as a major receptor for fungal pathogens on LCs through recognition of mannose and beta-

glucan structures (34). The Gram-negative bacterium *Yersinia pestis* is the only other

bacterium known to interact with langerin and does so through its LPS core oligosaccharide

(35). We identify *S. aureus* WTA β-GlcNAc as a new ligand for langerin. WTA is an abundant

evolutionarily conserved feature of the surface of Gram-positive bacteria, making it

advantageous for the host to recognize such structures in a timely manner. Although several

receptors for *S. aureus* WTA have been described, langerin is the first human innate receptor

to discriminate between the α-GlcNAc and β-GlcNAc modifications.

As an opportunistic microbial resident of the skin *S. aureus* is involved in the

development of skin disease. Therefore, the recognition of *S. aureus* WTA by langerin on

epidermal LCs that are strategically localized at mucosal surfaces may be key to maintaining

skin homeostasis and preventing the development of infection or chronic inflammation.

Alternatively, it is also possible that *S. aureus* exploits langerin interaction to intentionally

elicit inflammation to perturb the skin barrier and release nutrients. Our epicutaneous

infection experiments in WT and huLangerin transgenic mice do not support the latter

hypothesis since we did not observe differences in bacterial burden despite increased

inflammation in huLangerin transgenic mice. It remains to be determined why the clearly

altered skin immune response in huLangerin mice does not result in differential bacterial

survival in the skin. Potentially, 40 hours post-infection is too early to observe such an effect

or the expression of immune evasion molecules allows survival of *S. aureus* in this hostile

environment.

The identification of *S. aureus* as a new langerin-interacting pathogen is especially

interesting in the context of AD. First, *S. aureus* is a driver of AD disease progression, which

is mediated by LCs (9). Second, genome-wide association studies (GWAS) identified *CD207*,

the gene encoding for langerin, as an AD susceptibility locus (36,37). In these studies,

polymorphisms in a putative enhancer region of *CD207*, which likely increase expression of

langerin, were protective for AD. Our data now functionally link langerin to *S. aureus*. Since

S. aureus is largely resistant to host defenses but most of the other commensals are not, this

could explain the strong association between *S. aureus* and AD, as well as the described

driver function of *S. aureus* in AD disease progression. Also our observation that WTA α-

GlcNAc attenuates LC activation can be important in the context of AD. The CC1 lineage is

particularly overrepresented in isolates from AD skin and was suggested to have unidentified

features that enable colonization by and proliferation of *S. aureus* on AD skin (4).

Interestingly, all CC1 strains are *tarM*-positive (38), providing the potential to regulate WTA

glycoprofile by co-decoration with α-GlcNAc. This could enable the bacteria to skew the

inflammatory status of the skin and gain an advantage to colonize AD skin. Our data may

provide molecular insight into the association between AD and *S. aureus* from two different

angles: on the immunological side we show how langerin and LCs are involved in the

immune response to *S. aureus*, while on the microbiological side the involvement of langerin

could explain the association of *S. aureus* but not CoNS species with AD, and possibly also

the overrepresentation of *tarM*-bearing CC1 strains in AD.

In conclusion, we identify *S. aureus* WTA as a PAMP and pinpoint langerin as a

molecular trigger for *S. aureus*-induced skin immune responses. Our findings give a deeper

understanding of the specific association of *S. aureus* with skin inflammation and can help in

the development of new treatment strategies for *S. aureus*-associated skin and soft tissue

infections and inflammatory skin diseases.

Methods

Bacterial strains and culture conditions

S. aureus, S. capitis, S. carnosus, S. epidermidis, S. lugdunensis, S. pseudintermedius, S.

saprophyticus and *S. simulans* strains (Supplementary Table S1) were grown overnight at

37°C with agitation in 5 ml Todd-Hewitt broth (THB; Oxoid). For *S. aureus* strains that were

plasmid complemented THB was supplemented with 10 μg/ml chloramphenicol (Sigma

Aldrich). A fresh five ml THB culture was inoculated by 150 μl overnight culture and grown to

an optical density at 600 nm (OD_{600nm}) of 0.4 for *S. capitis* and to OD_{600nm}=0.6-0.7 for all other

bacteria, which corresponds to mid-exponential growth phase.

Cell culture and muLC differentiation

MUTZ-3 cells (ACC-295, DSMZ) were cultured in a 12-well tissue culture plates (Corning) at

a density of $0.5\text{-}1.0\text{x}10^6$ cells/ml in MEM-alpha (Gibco) with 20% fetal bovine serum (FBS,

Hyclone, GE Healthcare), 1% GlutaMAX (Gibco), 10% conditioned supernatant from renal

carcinoma cell line 5637 (ACC-35, DSMZ), 100 U/ml penicillin and 100 μg/ml streptomycin

(Gibco) at 37°C with 5% CO_2. We obtained MUTZ-3 derived Langerhans cells (muLCs) by

differentiation of MUTZ-3 cells for 10 days in 100 ng/ml Granulocyte-Macrophage Colony

Stimulating Factor (GM-CSF, GenWay Biotech), 10 ng/ml Transforming Growth Factor-beta

(TGFβ; R&D Systems) and 2.5 ng/ml Tumor Necrosis Factor-alpha (TNFα; R&D Systems) as

described previously (21,22). The phenotype of differentiated muLCs was verified by surface

staining of CD34 (clone 581, BD Biosciences), CD1a (clone HI149, BD Biosciences) and

CD207 (clone DCGM4, Beckman Coulter) using the respective antibodies and analysis by

flow cytometry.

THP1 cells (TIB-202, ATCC) transduced with a lentiviral langerin construct or empty vector

(EV) were cultured in RPMI (Lonza) supplemented with 5% FBS (Biowest), 1% GlutaMAX

100 U/ml penicillin and 100 μg/ml streptomycin (Gibco) at 37°C with 5% CO_2.

Isolation of primary human Langerhans cells

Human skin tissue was collected from otherwise healthy anonymous donors undergoing

corrective breast or abdominal surgery. This study, including the tissue harvesting

procedures, were approved by the Medical Ethics Review Committee of the Academic

Medical Center Amsterdam, The Netherlands. Human Langerhans cells were isolated as

described previously (33). In short, skin grafts were obtained using a dermatome (Zimmer)

and incubated in medium supplemented with Dispase II (1 U/ml, Roche Diagnostics) after

which epidermal sheets were separated from the dermis and cultured for three days. After

incubation, migrated LCs were harvested and further purified using a Ficoll gradient (Axis-

shield). Isolated LCs were routinely 90% pure (CD1a+ Langerin+) and were frozen in Iscoves

Modified Dulbeccos's Medium (IMDM, Thermo Fisher) supplemented with 20% FBS and

10% DMSO. Before use, LCs were thawed by drop-wise addition of cold IMDM with 10%

FBS, washed twice and incubated in IMDM with FBS for 2 hours at 37°C with 5% CO_2 to

recover.

Creation of GFP-expression *S. aureus*

To create GFP-expressing bacteria, *S. aureus* Newman wild-type and *S. aureus* Newman

Δ*spa*Δ*sbi* were transformed as described previously with pCM29, which encodes

superfolded green fluorescent protein (sGFP) driven by the *sarA*P1 promoter (39,40). In

short, competent *S. aureus* were electroporated with pCM29 isolated from *E. coli* DC10B

with a Gene Pulser II (BioRad; 100 Ohm, 25uF, 2.5kV). After recovery, bacteria were

selected on TH agar supplemented with 10 μg/ml chloramphenicol. A single colony was

grown in THB with 10 μg/ml chloramphenicol under the usual growth conditions. Bacterial

expression of GFP was verified by confocal laser scanning microscopy (SP5, Leica).

Gamma-irradiation of *S. aureus*

Gamma-irradiated stocks of *S. aureus* strains were made by harvesting cultures in mid-

exponential growth phase by centrifugation (4,000 rpm, 8 min), which were concentrated 10x

in phosphate-buffered saline (PBS; Lonza) with 17% glycerol (VWR), frozen at -70°C and

exposed to 10 kGy of γ-radiation (Synergy Health, Ede, The Netherlands). Loss of viability of

S. aureus was verified by plating of the irradiated bacteria. A non-irradiated aliquot that

underwent the same freezing procedure was used to determine the concentration of colony

forming units (CFU) of the irradiated stocks.

Lentiviral transduction

A TrueORF sequence-validated cDNA clone of human CD207 (OriGene Technologies) was

amplified by PCR using Phusion polymerase (Thermo Fisher) and primers hLangerin-Fw and

hLangerin-FLAG-Rv (IDT, Supplementary Table S2). The PCR amplicon was cloned in a

BIC-PGK-Zeo-T2a-mAmetrine;EF1A construct by Gibson assembly (NEB) according to the

manufacturer's instructions. The langerin-encoding vector and an empty vector (EV) control

were introduced into THP1 cells by lentiviral transduction, as described by Van de Weijer *et*

al. (41). In short, lentivirus was produced by HEK293T cells (CRL-3216, ATCC) in 24-well

plates using standard lentiviral production protocols and third-generation packaging vectors.

After 3-4 days the supernatant containing the viral particles was harvested and stored at -

70ºC to kill any remaining cells. Approximately 50,000 THP1 cells were transduced by spin

infection (1000xg, 2 h, 33°C) using 100 µl supernatant supplemented with 8 µg/ml polybrene

(Santa Cruz Biotechnology). Complete medium was added after centrifugation and cells

were selected three days post-infection by 100 µg/ml zeocin (Gibco). Cellular expression of

langerin was verified by antibody staining of langerin (clone DCGM4, Beckman Coulter) and

measured using flow cytometry.

Bacterial binding assays

To test binding of bacteria to cells, 10^5 LCs, THP1-EV or THP1-langerin were incubated with

GFP-expressing *S. aureus* Newman or GFP-expressing *S. aureus* Newman Δ*spa* Δ*sbi* at

bacteria-to-cell ratios from 1 to 8 in TSM buffer (2.4 g/L Tris (Roche), 8.77 g/L NaCl (Sigma

Aldrich), 294 mg/L CaCl$_2$·2H2O (Merck), 294 mg/L MgCl$_2$·6H2O (Merck), pH=7.4) with 0.1%

bovine serum albumin (BSA; Merck) for 30 minutes at 4°C. Binding was blocked by 15

minutes pre-incubation with 10 µg/ml mannan (Sigma Aldrich), 50 mM GlcNAc (Serva) or 20

µg/ml anti-langerin blocking antibody (clone 10E2, Sony Biotechnology). Cells were washed

once with TSM 1% BSA, fixed in 1% formaldehyde (Brunschwig Chemie) in PBS and

measured by flow cytometry.

Production of recombinant langerin extracellular domains

The extracellular domains of truncated human langerin (residues 148–328) and mouse

langerin (residues 150–331) were recombinantly expressed from codon-optimized constructs

containing a C-terminal TEV cleavage site followed by a Strep-tag II cloned into pUC19 and

pET30a (EMD Millipore) expression vectors as described previously (25). Recombinant

human and murine ECDs were insolubly expressed in *E. coli* BL21(DE3), solubilized in 6 M

guanidinium hydrochloride in 100 mM Tris (pH 8) with 1 mM DTT, refolded by dialyisis

against Tris-buffered saline (pH 7.5) containing 10 mM $CaCl_2$ and purified via mannan-

coupled sepharose beads (Sigma Aldrich). Bound protein was eluted with Tris-buffered

saline (pH 7.5) containing 5 mM EDTA. Protein concentrations were determined by A280 nm

using the calculated molar extinction coefficients of 56,170 M^{-1} cm^{-1} for the human langerin

ECD and 56,170 M^{-1} cm^{-1} for the murine ECD. The proteins were fluorescently labeled with

fluorescein isothiocyanate (FITC, Thermo Fisher) by adding slowly 100 µL of the dye solution

(1 mg/ml in DMSO) to 2 ml of a 2 mg/ml protein solution in HEPES-buffered saline (pH 7.2)

containing 20 mM D-mannose (Sigma Aldrich) and 5 mM $CaCl_2$. After stirring for 90 min at

room temperature, the reaction was quenched by addition of 50 mM ethanolamine (pH 8.5,

Sigma Aldrich). Unreacted dye molecules were removed by buffer exchange using a Zeba

spin column (Thermo Fisher) and active protein was purified over mannan affinity column as

described above. All chemicals used for the production of recombinant langerin extracellular

domains were obtained from Carl Roth if not indicated otherwise.

Langerin binding assay

Bacteria in mid-exponential growth phase were harvested by centrifugation (4,000 rpm, 8

minutes) and resuspended at $OD_{600nm}=0.4$ in TSM buffer with 0.1% BSA. Bacteria were

incubated with 1-50 µg/ml recombinant langerin-FITC (human or mouse) for 30 minutes at

37°C with agitation, washed once with TSM 1% BSA, fixed in 1% formaldehyde and

analyzed by flow cytometry.

muLC stimulation

We stimulated 5×10^4 muLCs with *S. aureus* USA300 WT, USA300 Δ*tarM* or USA300 Δ*tarS*

at bacteria-to-cell ratios of 0, 1, 10 and 50 in IMDM with 10% FBS. After 24 hours,

supernatants were collected by centrifugation (300xg, 10 min, 4°C) and stored at -150°C until

further analysis, and cells were washed once in PBS 0.1% BSA. Expression levels of the

activation and maturation markers were determined by flow cytometry using the following

antibodies: CD80 (clone 2D10), CD83 (clone HB15e) and CD86 (clone IT2.2, all from Sony

Biotechnology) and their corresponding isotype controls (BD Biosciences).

Cytokine assays

The IL-8 and IL12p70 concentrations were initially determined by ELISA (Sanquin and

Thermo Fisher, respectively) according to the manufacturer's instructions. Concentrations of

IL-6, IL-8, IL-10, IL-12p70, IL-23p19 and TNFα cytokines were determined by Luminex xMAP

assay (Luminex Corporation), performed by the Multiplex Core Facility UMC Utrecht, The

Netherlands.

Flow cytometry

Flow cytometry was performed on FACSVerse (BD Biosciences), per sample 10.000 events

within the set gate were collected. Data were analyzed using FlowJo 10 (FlowJo, LLC).

Epicutaneous murine infection model

We used 6- to 10 week-old sex-matched wild type C57BL/6 mice (obtained from Jackson

laboratories) and huLangerin-DTR mice (26), provided by D. H. Kaplan (University of

Pittsburgh, Pennsylvania USA). All mice were housed in a specific pathogen-free facility

under standard conditions at the University of Pittsburgh. The mouse infection protocols were

approved beforehand by the Institutional Animal Care and Use of Committee of the

University of Pittsburgh. As described previously, mice were first anesthetized with a mixture

of ketamine and xylazine (100/10 mg/kg body weight), shaved on the back with electric

clippers, chemically depilated with Nair hair removal cream (Church & Dwight) according to

the manufacturer's instructions, and the stratum corneum was removed by 15 strokes of 220

grit sandpaper (3M) (8,27). After 24 hours, the mice were epicutaneously inoculated with

PBS or *S. aureus* USA300 Δ*tarM*, which was grown overnight at 37°C in THB, in 50 µl of

sterile PBS. Forty hours post-infection the mice were sacrificed and skin sections of 1 cm^2

were collected. The sections were either 1) homogenized, serially diluted in sterile PBS,

grown overnight on THB-agar plates at 37°C and colony forming units were counted, 2)

homogenized and processed for RNA extraction or 3) fixed in 1% formalin in PBS. The fixed

tissue sections were embedded in paraffin, cut, stained with hematoxylin and eosin, and

digitalized (Hamamatsu NanoZoomer) by the Department of Pathology, UMC Utrecht, The

Netherlands, and subsequently analyzed using NDP.view2.6.13 (Hamamatsu).

Gene expression analysis

Whole skin was homogenized and processed for extraction and isolation of RNA, using

TRIzol reagents (Thermo Fisher), following manufacturer's instructions. RNA was quantified

using a standard Nanodrop and cDNA was obtained using High-Capacity cDNA Reverse

Transcriptase (Thermo Fisher). Quantitive PCR on cDNA was accomplished by using

Taqman Gene Expression Master Mix and Taqman Gene Expression Assays for *IL-17*, *IL-6*,

CXCL1, *CXCL2*, *IL-10* and *GAPDH* (Thermo Fisher) on a StepOnePlus Real Time PCR

System (Applied Biosystems). Fold upregulation of transcripts was calculated from ΔΔCt

values relative to *GAPDH* expression and normalized for PBS mock infection.

Statistical analysis

Statistical analyses were performed using Graphpad Prism 7.02 (GraphPad Software). We

used unpaired two-tailed *t*-tests for comparisons between two groups and one-way ANOVAs

with a common control group followed by Dunnett's multiple comparisons test. The THP1-

langerin dose-response curves were tested using a two-way ANOVA followed by Dunnett's

multiple comparison test and the langerin-FITC concentration curves were tested against

wild-type langerin-FITC using a two-way ANOVA followed by Tukey's multiple comparison

test. Data are presented as the geometric mean or percentage positive cells (flow cytometry),

mean concentration (cytokine arrays) or fold upregulation (real-time PCR) + standard error of

the mean (SEM).

Data availability

The data that support these findings are available from the corresponding author upon

request.

Acknowledgements

The authors would like to thank: the Multiplex Core Facility (UMC Utrecht, The Netherlands)

for performing the Luminex assay; David Gerlach, Guoqing Xia and Volker Winstel (currently

or previously at University of Tübingen, Germany) for advice and communication regarding

the shipment of strains; Samantha van der Beek, Eline van Yperen and Małgorzata Mnich

(UMC Utrecht, The Netherlands) for technical assistance; Tanja de Gruijl (VU University

Medical Center, Amsterdam, The Netherlands) for the protocol detailing MUTZ-3 culture and

differentiation. This work was supported by a VIDI grant (91713303) from the Dutch Scientific

Organization (NWO) to N.M.v.S and R.v.D. A.P. is supported by German Research Council

(DFG) grants TRR34, TRR156 and the German Center for Infection Research (DZIF). J.H.

thanks the DFG Collaborative Research Centre 765 "Multivalency" for a fellowship. C.R.

thanks the DFG for funding (RA1944/2-1).

Author Contributions

R.v.D., M.R. and N.M.v.S. planned the experiments. R.v.D., J.S.D.L.C.D. and M.R.

performed the experiments and prepared the figures. F.F.F., J.H. and C.R. supplied the

langerin-FITC constructs, N.H.v.T. and T.B.H.G. provided the primary LCs, C.W. and A.P.

provided the bacterial strains, D.H.K. provided the mice. R.v.D. and N.M.v.S. wrote the

manuscript, N.H.v.T., J.A.G.v.S., C.W. and A.P provided critical feedback.

Competing interests

The authors declare no competing financial interests.

References

1. Nutten S (2015) Atopic dermatitis: global epidemiology and risk factors. Ann Nutr Metab 66 Suppl 1:8-16. Doi:10.1159/000370220

2. Totte JE, van der Feltz WT, Hennekam M, van Belkum A, van Zuuren EJ, et al. (2016) Prevalence and odds of *Staphylococcus aureus* carriage in atopic dermatitis: a systematic review and meta-analysis. Br J Dermatol. Doi:10.1111/bjd.14566

3. Nakatsuji T, Chen TH, Narala S, Chun KA, Two AM, et al. (2017) Antimicrobials from human skin commensal bacteria protect against *Staphylococcus aureus* and are deficient in atopic dermatitis. Sci Transl Med 9(378). Doi:10.1126/scitranslmed.aah4680

4. Geoghegan JA, Irvine AD, Foster TJ (2017) *Staphylococcus aureus* and Atopic Dermatitis: A Complex and Evolving Relationship. Trends Microbiol. Doi:10.1016/j.tim.2017.11.008

5. Valladeau J, Ravel O, Dezutter-Dambuyant C, Moore K, Kleijmeer M, et al. (2000) Langerin, a novel C-type lectin specific to Langerhans cells, is an endocytic receptor that induces the formation of Birbeck granules. Immunity 12(1):71-81.

6. Kissenpfennig A, Henri S, Dubois B, Laplace-Builhe C, Perrin P, et al. (2005) Dynamics and function of Langerhans cells in vivo: dermal dendritic cells colonize lymph node areas distinct from slower migrating Langerhans cells. Immunity 22(5):643-54. Doi:10.1016/j.immuni.2005.04.004

7. Seneschal J, Clark RA, Gehad A, Baecher-Allan CM, Kupper TS (2012) Human epidermal Langerhans cells maintain immune homeostasis in skin by activating skin resident regulatory T cells. Immunity 36(5):873-84. Doi:10.1016/j.immuni.2012.03.018

8. Igyártó BZ, Haley K, Ortner D, Bobr A, Gerami-Nejad M, et al. (2011) Skin-resident murine dendritic cell subsets promote distinct and opposing antigen-specific T helper cell responses. Immunity 35(2):260-72. Doi:10.1016/j.immuni.2011.06.005

9. Kobayashi T, Glatz M, Horiuchi K, Kawasaki H, Akiyama H, et al. (2015) Dysbiosis and *Staphylococcus aureus* Colonization Drives Inflammation in Atopic Dermatitis. Immunity 42(4):756-66. Doi:10.1016/j.immuni.2015.03.014

10. Winstel V, Kuhner P, Salomon F, Larsen J, Skov R, et al. (2015) Wall Teichoic Acid Glycosylation Governs *Staphylococcus aureus* Nasal Colonization. mBio 6(4):e00632. Doi:10.1128/mBio.00632-15

11. Weidenmaier C, Peschel A, Xiong YQ, Kristian SA, Dietz K, et al. (2005) Lack of wall teichoic acids in *Staphylococcus aureus* leads to reduced interactions with endothelial cells and to attenuated virulence in a rabbit model of endocarditis. The Journal of infectious diseases 191(10):1771-7. Doi:10.1086/429692

12. Brown S, Xia G, Luhachack LG, Campbell J, Meredith TC, et al. (2012) Methicillin resistance in *Staphylococcus aureus* requires glycosylated wall teichoic acids. Proceedings of the National Academy of Sciences of the United States of America 109(46):18909-14. Doi:10.1073/pnas.1209126109

13. Xia G, Corrigan RM, Winstel V, Goerke C, Grundling A, et al. (2011) Wall teichoic Acid-dependent adsorption of staphylococcal siphovirus and myovirus. Journal of bacteriology 193(15):4006-9. Doi:10.1128/JB.01412-10

14. Winstel V, Liang C, Sanchez-Carballo P, Steglich M, Munar M, et al. (2013) Wall teichoic acid structure governs horizontal gene transfer between major bacterial pathogens. Nature communications 4:2345. Doi:10.1038/ncomms3345

15. Xia G, Maier L, Sanchez-Carballo P, Li M, Otto M, et al. (2010) Glycosylation of wall teichoic acid in *Staphylococcus aureus* by TarM. The Journal of biological chemistry 285(18):13405-15. Doi:10.1074/jbc.M109.096172

16. Li X, Gerlach D, Du X, Larsen J, Stegger M, et al. (2015) An accessory wall teichoic acid glycosyltransferase protects *Staphylococcus aureus* from the lytic activity of Podoviridae. Sci Rep 5:17219. Doi:10.1038/srep17219

17. Baur S, Rautenberg M, Faulstich M, Grau T, Severin Y, et al. (2014) A nasal epithelial receptor for *Staphylococcus aureus* WTA governs adhesion to epithelial cells and modulates nasal colonization. PLoS pathogens 10(5):e1004089. Doi:10.1371/journal.ppat.1004089

18. Kurokawa K, Jung DJ, An JH, Fuchs K, Jeon YJ, et al. (2013) Glycoepitopes of staphylococcal wall teichoic acid govern complement-mediated opsonophagocytosis via human serum antibody and mannose-binding lectin. The Journal of biological chemistry 288(43):30956-68. Doi:10.1074/jbc.M113.509893

19. Park KH, Kurokawa K, Zheng L, Jung DJ, Tateishi K, et al. (2010) Human serum mannose-binding lectin senses wall teichoic acid Glycopolymer of *Staphylococcus aureus*, which is restricted in infancy. The Journal of biological chemistry 285(35):27167-75. Doi:10.1074/jbc.M110.141309

20. van der Aar AM, Picavet DI, Muller FJ, de Boer L, van Capel TM, et al. (2013) Langerhans cells favor skin flora tolerance through limited presentation of bacterial antigens and induction of regulatory T cells. The Journal of investigative dermatology 133(5):1240-9. Doi:10.1038/jid.2012.500

21. Masterson AJ, Sombroek CC, De Gruijl TD, Graus YM, van der Vliet HJ, et al. (2002) MUTZ-3, a human cell line model for the cytokine-induced differentiation of dendritic cells from CD34+ precursors. Blood 100(2):701-3.

22. Santegoets SJ, Masterson AJ, van der Sluis PC, Lougheed SM, Fluitsma DM, et al. (2006) A CD34(+) human cell line model of myeloid dendritic cell differentiation: evidence for a CD14(+)CD11b(+) Langerhans cell precursor. Journal of leukocyte biology 80(6):1337-44. Doi:10.1189/jlb.0206111

23. Lee JH, Kim NH, Winstel V, Kurokawa K, Larsen J, et al. (2015) Surface Glycopolymers Are Crucial for In Vitro Anti-Wall Teichoic Acid IgG-Mediated Complement Activation and Opsonophagocytosis of *Staphylococcus aureus*. Infect Immun 83(11):4247-55. Doi:10.1128/IAI.00767-15

24. Koymans KJ, Vrieling M, Gorham RD, Jr., van Strijp JA (2016) Staphylococcal Immune
Evasion Proteins: Structure, Function, and Host Adaptation. Curr Top Microbiol Immunol.
Doi:10.1007/82_2015_5017

25. Hanske J, Schulze J, Aretz J, McBride R, Loll B, et al. (2017) Bacterial Polysaccharide
Specificity of the Pattern Recognition Receptor Langerin Is Highly Species-dependent. The
Journal of biological chemistry 292(3):862-71. Doi:10.1074/jbc.M116.751750

26. Bobr A, Olvera-Gomez I, Igyarto BZ, Haley KM, Hogquist KA, et al. (2010) Acute ablation
of Langerhans cells enhances skin immune responses. J Immunol 185(8):4724-8.
Doi:10.4049/jimmunol.1001802

27. Kashem SW, Igyarto BZ, Gerami-Nejad M, Kumamoto Y, Mohammed J, et al. (2015)
Candida albicans morphology and dendritic cell subsets determine T helper cell
differentiation. Immunity 42(2):356-66. Doi:10.1016/j.immuni.2015.01.008

28. Iwamoto K, Moriwaki M, Niitsu Y, Saino M, Takahagi S, et al. (2017) *Staphylococcus*
aureus from atopic dermatitis skin alters cytokine production triggered by monocyte-derived
Langerhans cell. J Dermatol Sci. Doi:10.1016/j.jdermsci.2017.08.001

29. Kraus D, Herbert S, Kristian SA, Khosravi A, Nizet V, et al. (2008) The GraRS regulatory
system controls *Staphylococcus aureus* susceptibility to antimicrobial host defenses. BMC
Microbiol 8:85. Doi:10.1186/1471-2180-8-85

30. Falord M, Mader U, Hiron A, Debarbouille M, Msadek T (2011) Investigation of the
Staphylococcus aureus GraSR regulon reveals novel links to virulence, stress response and
cell wall signal transduction pathways. PLoS One 6(7):e21323.
Doi:10.1371/journal.pone.0021323

31. Wanner S, Schade J, Keinhorster D, Weller N, George SE, et al. (2017) Wall teichoic
acids mediate increased virulence in *Staphylococcus aureus.* Nat Microbiol 2:16257.
Doi:10.1038/nmicrobiol.2016.257

32. de Witte L, Nabatov A, Pion M, Fluitsma D, de Jong MA, et al. (2007) Langerin is a
natural barrier to HIV-1 transmission by Langerhans cells. Nature medicine 13(3):367-71.
Doi:10.1038/nm1541

33. Ribeiro CM, Sarrami-Forooshani R, Setiawan LC, Zijlstra-Willems EM, van Hamme JL, et
al. (2016) Receptor usage dictates HIV-1 restriction by human TRIM5alpha in dendritic cell
subsets. Nature 540(7633):448-52. Doi:10.1038/nature20567

34. de Jong MA, Vriend LE, Theelen B, Taylor ME, Fluitsma D, et al. (2010) C-type lectin
Langerin is a beta-glucan receptor on human Langerhans cells that recognizes opportunistic
and pathogenic fungi. Mol Immunol 47(6):1216-25. Doi:10.1016/j.molimm.2009.12.016

35. Yang K, Park CG, Cheong C, Bulgheresi S, Zhang S, et al. (2015) Host Langerin
(CD207) is a receptor for *Yersinia pestis* phagocytosis and promotes dissemination.
Immunology and cell biology 93(9):815-24. Doi:10.1038/icb.2015.46

36. Paternoster L, Standl M, Waage J, Baurecht H, Hotze M, et al. (2015) Multi-ancestry
genome-wide association study of 21,000 cases and 95,000 controls identifies new risk loci
for atopic dermatitis. Nat Genet 47(12):1449-56. Doi:10.1038/ng.3424

37. Cai XY, Zheng XD, Fang L, Zhou FS, Sheng YJ, et al. (2017) A variant on chromosome
2p13.3 is associated with atopic dermatitis in Chinese Han population. Gene 628:281-5.
Doi:10.1016/j.gene.2017.07.059

38. Winstel V, Xia G, Peschel A (2014) Pathways and roles of wall teichoic acid glycosylation
in *Staphylococcus aureus*. International journal of medical microbiology : IJMM 304(3-4):215-
21. Doi:10.1016/j.ijmm.2013.10.009

39. Pang YY, Schwartz J, Thoendel M, Ackermann LW, Horswill AR, et al. (2010) agr-
Dependent interactions of *Staphylococcus aureus* USA300 with human polymorphonuclear
neutrophils. J Innate Immun 2(6):546-59. Doi:10.1159/000319855

40. Schenk S, Laddaga RA (1992) Improved method for electroporation of *Staphylococcus*
aureus. FEMS Microbiol Lett 73(1-2):133-8.

41. van de Weijer ML, van Muijlwijk GH, Visser LJ, Costa AI, Wiertz EJ, et al. (2016) The E3
Ubiquitin Ligase TMEM129 Is a Tri-Spanning Transmembrane Protein. Viruses 8(11).
Doi:10.3390/v8110309

42. Sibbald MJ, Winter T, van der Kooi-Pol MM, Buist G, Tsompanidou E, et al. (2010)
Synthetic effects of secG and secY2 mutations on exoproteome biogenesis in
Staphylococcus aureus. Journal of bacteriology 192(14):3788-800. Doi:10.1128/JB.01452-09

43. Kreiswirth BN, Lofdahl S, Betley MJ, O'Reilly M, Schlievert PM, et al. (1983) The toxic
shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. Nature
305(5936):709-12.

44. CDC (1999) From the Centers for Disease Control and Prevention. Four pediatric deaths
from community-acquired methicillin-resistant *Staphylococcus aureus*--Minnesota and North
Dakota, 1997-1999. JAMA 282(12):1123-5.

45. Mwangi MM, Wu SW, Zhou Y, Sieradzki K, de Lencastre H, et al. (2007) Tracking the in
vivo evolution of multidrug resistance in *Staphylococcus aureus* by whole-genome
sequencing. Proceedings of the National Academy of Sciences of the United States of
America 104(22):9451-6. Doi:10.1073/pnas.0609839104

46. Guinane CM, Ben Zakour NL, Tormo-Mas MA, Weinert LA, Lowder BV, et al. (2010)
Evolutionary genomics of *Staphylococcus aureus* reveals insights into the origin and
molecular basis of ruminant host adaptation. Genome Biol Evol 2:454-66.
Doi:10.1093/gbe/evq031

47. Rosenstein R, Nerz C, Biswas L, Resch A, Raddatz G, et al. (2009) Genome analysis of
the meat starter culture bacterium *Staphylococcus carnosus* TM300. Appl Environ Microbiol
75(3):811-22. Doi:10.1128/AEM.01982-08

48. Mack D, Siemssen N, Laufs R (1992) Parallel induction by glucose of adherence and a
polysaccharide antigen specific for plastic-adherent *Staphylococcus epidermidis*: evidence
for functional relation to intercellular adhesion. Infection and immunity 60(5):2048-57.

49. Chassain B, Lemee L, Didi J, Thiberge JM, Brisse S, et al. (2012) Multilocus sequence
typing analysis of *Staphylococcus lugdunensis* implies a clonal population structure. J Clin
Microbiol 50(9):3003-9. Doi:10.1128/JCM.00988-12

50. Ben Zakour NL, Bannoehr J, van den Broek AH, Thoday KL, Fitzgerald JR (2011)
Complete genome sequence of the canine pathogen *Staphylococcus pseudintermedius*.
Journal of bacteriology 193(9):2363-4. Doi:10.1128/JB.00137-11

51. Monk IR, Shah IM, Xu M, Tan MW, Foster TJ (2012) Transforming the untransformable:
application of direct transformation to manipulate genetically *Staphylococcus aureus* and
Staphylococcus epidermidis. mBio 3(2). Doi:10.1128/mBio.00277-11

Figures

Figure 1. Langerin is a receptor for *S. aureus* on human LCs
(A) Binding of *S. aureus* to isolated primary human LCs. LCs from donors 1 and 2 were
incubated with GFP-expressing *S. aureus* Newman and LCs from donors 3 and 4 with GFP-
expressing *S. aureus* Newman Δ*spa*Δ*sbi* and binding was assessed by flow cytometry. The
interaction was blocked by addition of mannan, GlcNAc or anti-langerin blocking antibody
(donors 3 and 4 only).
(B) Binding of *S. aureus* to THP1-langerin cells. Human langerin-transduced or empty vector
(EV)-transduced THP1 cells were incubated with different amounts of GFP-expressing *S.*
aureus Newman Δ*spa*Δ*sbi*. The interaction was blocked by addition of mannan or anti-
langerin blocking antibody. Within each ratio, THP1-langerin was compared to the other
conditions by two-way ANOVA followed by Dunnett's multiple comparison test.
(C) Expression of co-stimulatory molecules CD80 and CD86 and production of cytokines IL-8
and IL12p70 by muLCs after incubation with γ-irradiated *S. aureus* USA300 (24 h). Values
were compared to unexposed control by one-way ANOVA followed by Dunnett's multiple
comparison test.
Data are presented as percentage GFP+ cells (A, B), geometric mean fluorescent intensity or
mean concentration (C) +/- standard error of mean (SEM) from three independent
experiments, except for (A) (four donors with single measurements). $*P < 0.05$, $**P < 0.01$,
$***P < 0.001$.

Figure 2. Langerin recognizes *S. aureus* in a *tarS*-dependent manner through the conserved WTA β-GlcNAc epitope

Binding of recombinant human langerin-FITC to (A) 18 wild-type *S. aureus* strains (11 different clonal complexes, indicated above the bars and by different color) and a selection of coagulase-negative staphylococcal species (CoNS); (B) *S. aureus* USA300 wild-type (WT) and WTA biosynthesis mutants Δ*tarMS*, Δ*tarM*, Δ*tarS*, Δ*tarMS* p*tarM* and Δ*tarMS* p*tarS*; and (C) two representative *S. aureus* isolates (82086 and PS66) that naturally lack *tarM* and their isogenic Δ*tarS* mutants. For B, C: all strains were grown to mid-exponential phase and incubated with langerin-FITC (blue) or buffer (black). Binding was assessed by flow cytometry. Data were compared by one-way ANOVA followed by Dunnett's multiple comparison test (B) or by unpaired two-tailed *t*-test (C) and are presented as geometric mean fluorescence intensity + SEM from three independent experiments. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 3. *S. aureus* induces a Th1 and Th17 cytokine profile in LCs, which is affected by the WTA glycoprofile
Expression of (A) co-stimulatory molecules CD80 and CD86 and maturation marker CD83 and (B) cytokines IL-6, IL-8, IL12p70, IL23p19 and TNFα by muLCs. muLCs were incubated with γ-irradiated *S. aureus* USA300 wild-type (WT), Δ*tarM* and Δ*tarS* for 24 h. muLCs stimulated with WT *S. aureus* were compared to the unstimulated control and muLCs stimulated with Δ*tarM* and Δ*tarS* were compared to their respective WT controls within the same ratio by one-way ANOVA followed by Dunnett's multiple comparison test. Data are presented as geometric mean fluorescence intensity or mean concentration + SEM from three independent experiments. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Figure 4. Human langerin transgenic mice show an enhanced inflammation to epicutaneous *S. aureus* infection

(A) Representative images of hematoxylin and eosin staining of skin biopsies and (B) transcript abundance of *CXCL1*, *IL-6* and *IL-17* from the lesions of WT (n=3) and huLangerin (n=4) mice 40 hours post-epicutaneous inoculation with *S. aureus* Δ*tarM*. The scale bars in (A) represent 100 μm. Data in (B) are presented as fold upregulation + SEM relative to *GAPDH* from three or four technical replicates and normalized for the WT/PBS control. The groups were compared by unpaired two-tailed *t*-test. **$P < 0.01$.