Transcription-dependent spreading of canonical yeast GATA factor across the body of highly

expressed genes

Aria Ronsmans[1,3], Maxime Wéry[2,3], Camille Gautier[2], Marc Descrimes[2], Evelyne Dubois[1], Antonin

Morillon[2,4] and Isabelle Georis[1,4]

[1] Metabolism of Model Microorganisms, Institut de Recherches Microbiologiques J.M. Wiame - Labiris,

Laboratoire de Microbiologie, Université Libre de Bruxelles, avenue Emile Gryson 1, 1070 Brussels,

Belgium

[2] ncRNA, epigenetic and genome fluidity, Institut Curie, PSL Research University, CNRS UMR 3244,

Université Pierre et Marie Curie, 26 rue d'Ulm, 75248 Paris Cedex 05, France

[3] Equal contribution

[4] Corresponding authors: igeoris@ulb.ac.be (lead contact), antonin.morillon@curie.fr

Abstract

GATA transcription factors are highly conserved among eukaryotes and play roles in transcription of

genes implicated in cancer progression and hematopoiesis. However, although their consensus binding

sites have been well defined *in vitro*, the *in vivo* selectivity for recognition by GATA factors remains

poorly characterized.

Using ChIP-Seq, we identified the Dal80 GATA factor targets in yeast. Our data reveal Dal80 binding to

a large set of promoters, sometimes independently of GATA sites. Strikingly, Dal80 was also detected

across the body of promoter-bound genes, correlating with high, Dal80-sensitive expression.

Mechanistic single-gene experiments showed that Dal80 spreading across gene bodies is independent

of intragenic GATA sites but requires transcription elongation. Consistently, Dal80 co-purified with the

post-initiation form of RNA Polymerase II.

Our work suggests that GATA factors could play dual, synergistic roles during transcription initiation

and elongation, promoting efficient remodeling of the gene expression program in response to

environmental changes.

Introduction

In eukaryotes, gene transcription by RNA polymerase II (PolII) is initiated by the binding of specific

transcription factors to double stranded DNA. The yeast transcription factors target regulatory regions

called UAS or URS (for *U*pstream *A*ctivating/*R*epressing *S*equences), generally directly adjacent to the

core promoter. The generated regulatory signals converge at the core promoter where they permit

the regulation of PolII recruitment via the 'TATA box-binding protein' and associated general

transcription factors (Hahn & Young, 2011; Levine & Tjian, 2003). The transcription factor binding sites

are usually short sequences ranging from 8 to 20 bp (Zambelli, Pesole, & Pavesi, 2013). They are most

often similar but generally not identical, differing by some nucleotides from one another (Zambelli et

al., 2013), making it sometimes difficult to predict whether a given UAS will function as such *in vivo*.

GATA factors constitute a family of transcription factors highly conserved among eukaryotes and

characterized by the presence of a DNA binding domain which consists of four cysteines (fitting the

consensus sequence $CX_2CX_{17-18}CX_2C$) coordinating a zinc ion followed by a basic carboxy-terminal tail

(Omichinski et al., 1993). *In vitro*, this conserved binding domain allows the recognition of consensus

GATA DNA sequences (GATAA, GATAAG or GATTAG), found in the promoter of numerous genes, which

was demonstrated in various organisms either with direct or indirect methods (T. S. Cunningham,

Dorrington, & Cooper, 1994; Horak et al., 2002; Omichinski et al., 1993; Stanbrough & Magasanik,

1996; Starich, Wikström, Arst, Clore, & Gronenborn, 1998). Structure elucidation of GATA factor

binding domains in complex with their DNA target suggest an important role for the basic tail in

sequence specificity determination (Omichinski et al., 1993; Starich et al., 1998).

Since its discovery 40 years ago in chicken cells, the family of GATA factors was extended in human

cells and represents master regulators of hematopoiesis and cancer (Bresnick, Katsumura, Lee, Johnson, & Perkins, 2012). However, although approximately 7 million GATA motifs can be found in the human genome, the GATA factors occupy only 0.1–1% of them. Conversely, other regions are occupied by GATA factors despite lacking the consensus motif (Cheng et al., 2009; Tripic et al., 2009). In addition, GATA factors can swap among them for the same motif and switch from active or repressive transcriptional activity. All these observations raised the main paradigm on understanding the selective recruitment and activity of GATA factors and what the determinants of their residency on the chromatin are (Bresnick, Lee, Fujiwara, Johnson, & Keles, 2010; Grass et al., 2003).

In yeast, the family of GATA transcription factors contains over 10 members (Lowry & Atchley, 2000). Four of them are implicated in the regulation of *N*itrogen *C*atabolite *R*epression- (NCR-) sensitive genes, the expression of which is repressed in the presence of a preferred nitrogen source (glutamine, asparagine, ammonia) and derepressed when only poor nitrogen sources (e.g. proline, leucine, urea) are available (Magasanik & Kaiser, 2002). These key GATA factors involved in NCR signaling are two activators (Gln3 and Gat1/Nil1) and two repressors (Gzf3/Nil2/Deh1 and Dal80/Uga43) (J A Coffman, Rai, Cunningham, Svetlov, & Cooper, 1996; Coornaert, Vissers, Andre, & Grenson, 1992; T S Cunningham & Cooper, 1991; Minehart & Magasanik, 1991; Rowen, Esiobu, & Magasanik, 1997; Soussi-Boudekou, Vissers, Urrestarazu, Jauniaux, & André, 1997). Moreover, the expression of *DAL80* and *GAT1* is also NCR-sensitive, which implies cross- and autogenous regulations of the GATA factors in the NCR mechanisms (Jonathan A. Coffman et al., 1997; I. Georis, Feller, Vierendeels, & Dubois, 2009; Isabelle Georis, Tate, Vierendeels, Cooper, & Dubois, 2015; Soussi-Boudekou et al., 1997). Under nitrogen limitation, expression of *DAL80* is highly induced (T S Cunningham & Cooper, 1991), and Dal80 enters the nucleus where it competes with the two GATA activators for the same binding sites (André et al., 1995; Jonathan A. Coffman et al., 1997; T. S. Cunningham et al., 1994). Although initially described as being active under nitrogen abundance (Rowen et al., 1997; Soussi-Boudekou et al., 1997), the Gzf3 repressor also localizes to NCR-sensitive promoters in conditions of activation (I. Georis et al., 2009).

In vitro, the Gln3 and Gat1 activators bind, presumably as monomers, to single GATA sequences

(Thomas S. Cunningham, Svetlov, Rai, Smart, & Cooper, 1996), like their orthologous vertebrate

counterparts. On the other hand, the Dal80 repressor can form homo-and heterodimers with Gzf3,

thanks to the presence of an N-terminal leucine zipper (Svetlov & Cooper, 1998) and, consistently,

Dal80 binds *in vitro* to two GATA sequences (Jonathan A. Coffman et al., 1997; T. S. Cunningham &

Cooper, 1993; T. S. Cunningham et al., 1994; Svetlov & Cooper, 1998). *In vivo*, GATA factor binding site

recognition also appears to require repeats of GATA motifs within promoters. Indeed, synthetic

constructions containing a portion of the NCR-sensitive *DAL5* promoter have shown in a β-

galactosidase assay that at least two copies of the consensus are required to support efficient gene

activation. This led to the actual fuzzy definition of UAS_{NTR}, consisting in two GATA sites located

sufficiently close to one another to present a binding platform for GATA factors (Bysani, Daugherty, &

Cooper, 1991; Cooper, Rai, & Yoo, 1989; Rai, Genbauffe, Sumrada, & Cooper, 1989).

In vivo, GATA ChIP analyses have been carried out genome-wide in yeast (Harbison et al., 2004; Lee,

2002; MacIsaac et al., 2006; Venters & Pugh, 2009). They allowed gaining insights into the GATA factor

gene network through the identification of direct targets and characterization of the DNA elements

required to render a gene NCR-sensitive. However, these studies were not performed in activating

conditions, when all GATA factors are expressed, localized in the nucleus and active, so that the current

list of GATA factor targets are likely to be underestimated. On another hand, bioinformatic analyses

have shown that, since GATA sequences are short, they can be found almost everywhere throughout

the genome. Therefore, based on the sole criteria of the presence of repeated GATA sequences in

yeast promoters, a third of the yeast genes could hypothetically be NCR regulator targets (van Helden,

André, & Collado-Vides, 1998). Such GATA motif repetitions have been found in the promoter of 91

genes, inducible by GATA activators in absence of a good nitrogen source, supposed to be directly

targeted by the GATA activators (Scherens, Feller, Vierendeels, Messenguy, & Dubois, 2006).

Nevertheless, the functionality of these supposed UAS still needs to be directly demonstrated *in vivo*

(Hahn & Young, 2011).

Here, we provide the first genome-wide identification of Dal80 targets in yeast, in physiological

conditions where Dal80 is fully expressed and active. Using a ChIP-Seq approach combined to a

bioinformatic peak-calling procedure, we defined the exhaustive set of Dal80-bound promoters, which

turned out to be much larger than anticipated. Our data indicate that at some promoters, Dal80

recruitment occurs independently of GATA sites. Strikingly, Dal80 was also detected across the body

of a subset of genes bound at the promoter, globally correlating with high and Dal80-sensitive

expression. Mechanistic single-gene experiments confirmed the Dal80 binding profiles, further

indicating that Dal80 spreading across gene bodies occurs independently of intragenic GATA sites but

requires transcription elongation. Finally, co-immunoprecipitation experiments revealed that Dal80

physically interacts with both intitiating and elongating form of PolII. Our work suggests that in addition

to their well-established role at the (pre)inititation stage, GATA factors could be involved in

transcription elongation, promoting efficient remodeling of the gene expression program in response

to environmental changes.

Results

Genome-wide identification of Dal80-bound promoters

In order to determine the genome-wide occupancy of a GATA factor in yeast, our rationale was to

choose Dal80 as it is known to be highly expressed in derepressing conditions and forms chromosome

foci when tagged by GFP (Distler, Kulkarni, Rai, & Cooper, 2001). We thus grew yeast cells in proline-

containing media and performed a ChIP-Seq analysis using a Dal80-Myc[13] tagged strain and the

isogenic untagged strain, as a control (Fig. 1a). Dal80-bound regions were identified using a peak-

calling algorithm (see Material & Methods). A promoter was defined as bound by Dal80 when >75% of

the -350 to -100 region (relative to the downstream ORF start site) was overlapped by a peak (Fig. 1b).

We chose to use as the reference coordinate the translation initiation codon rather than the

transcription start site (TSS) because the latter has not been accurately determined for all genes. Then,

our arbitrary definition of the promoter as the -350 to -100 region relative to the ATG codon was based

on the distribution of the TSS-ATG distance for genes with an annotated TSS (median and average

distance = 58 and 107 bp, respectively; see Supp. Fig. 1a).

Strikingly, Dal80 was found to bind to 1269 gene promoters (Fig. 1c-d and Supp. Table 3a). Figure 1E

shows a snapshot of ChIP-Seq signals at the promoter of the *FUR4* gene, identified as bound by Dal80.

The number of Dal80-bound promoters, corresponding to 22% of all protein-coding genes promoters,

is much higher than expected given the hundred target genes that are generally cited for the GATA

transcriptional activators (Godard et al., 2007; Scherens et al., 2006), that are presumed to bind to the

same sites as Dal80. However, we noted that some peaks (221) overlapped with several promoters

(471), mainly of divergent genes (442). In such cases, it is therefore possible that only one of the two

divergent promoters is indeed targeted by Dal80, so that the number of Dal80-bound promoters could

actually be a little less important than 1269.

As expected, genes related to previous nitrogen regulation screens were present within our list (Airoldi

et al., 2016; Boer et al., 2007; Boer, De Winde, Pronk, & Piper, 2003; Cox, Pinchak, & Cooper, 1999;

Daugherty, Rai, El Berry, & Cooper, 1993; Godard et al., 2007; Scherens et al., 2006; Tai et al., 2005;

Wu, Zhang, Hayes, Panoutsopoulou, & Oliver, 2004)(Supp. Table 3b). Among the genes showing Dal80-

binding to the promoter, we also noticed a strong enrichment for cytoplasmic translation genes, as

well as genes involved in small molecule biosyntheses, including amino acids (Supp. Table 3c).

Surprisingly, analysis of GATA site occurrence over Dal80-bound and unbound promoters revealed no

major difference between the two classes (Fig. 1f), 48.2% and 51.3% of Dal80-bound and unbound

promoters containing at least two GATA sites, respectively (see Supp. Table 3d). Intriguingly, 20% of

Dal80-bound promoters do not contain any GATA site (Fig. 1f and Supp. Table 3d), indicating that Dal80

recruitment can also occur independently of the presence of consensus GATA sites (see Supp. Fig. 1b

for visualization of Dal80 binding to a GATA-less promoter).

In summary, our ChIP-Seq analysis revealed that Dal80 binds to a set of promoters larger than

previously expected, targeting biosynthetic functions and protein synthesis in addition to nitrogen

catabolite repression.

Dal80 binds the intragenic region of a subset of genes

Metagene analysis revealed that the genes bound by Dal80 at the promoter also display a signal along the gene body, although this intragenic signal remains globally lower than in the promoter-proximal region (Fig. 1d). This observation prompted us to investigate the possibility that Dal80 also occupies the gene body, at least for a subset of genes.

We identified 189 genes showing Dal80 intragenic occupancy, according to a >75% overlap of the ORF by a Dal80-Myc[13] peak (Fig. 2a-b). Among them, 144 (76%) were also bound at the promoter (Fig. 2b). Conversely, only 11% of the Dal80-bound promoters (144/1269) showed downstream intragenic binding (Fig. 2b). On the other hand, 45 genes showing Dal80 intragenic binding were not bound at the promoter (Fig. 2b). Hence, we distinguished four classes of genes (Supp. Table 4): (*i*) those bound by Dal80 at the promoter only ("P" class; Fig. 2c), (*ii*) those showing both promoter and intragenic binding ("P & O" class; Fig. 2d), (*iii)* those bound across the ORF only ("O" class; Fig. 2e), (*iv*) the unbound genes (Fig. 2f). Interestingly, we noted that the global Dal80-Myc[13] signal at the promoter was higher for the "P & O" class in comparison to the "P" class (compare Fig. 2c-d).

In conclusion, a subset of genes displays intragenic Dal80 binding, and in most cases, this correlates with strong Dal80 recruitment at the promoter.

Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels

In order to correlate Dal80 binding with gene expression and Dal80-sensitivity in similar culture conditions, we performed total RNA-seq analysis in wild type and *dal80Δ* cells grown in medium with proline as the nitrogen source.

The expression of 546 genes was significantly altered in proline-grown *dal80Δ* cells compared to wild type (Fig. 3a, Supp. Table 5a), among which 232 and 314 genes were negatively and positively regulated by Dal80, respectively. The first group was enriched for genes involved in small molecule catabolic processes (Supp. Table 5b), while the positively regulated genes were mostly involved in amino acid biosynthetis (Supp. Table 5c). Again, we noticed an overlap between Dal80-repressed genes and nitrogen regulated genes that were identified in other genome-wide screens (Supp. Table 3b).

Analysis of mRNA levels in proline-grown cells revealed a significant correlation between Dal80 binding

and regulation at the mRNA level: 39.6% (216/546) of Dal80-regulated genes were bound by Dal80 at

their promoter and/or across the ORF, which is significantly more than expected if Dal80 binding and

regulation were independent (P<0.00001, Chi-square test; Supp. Fig. 2a). More importantly, additional

differences are emerging between the different classes of genes. First, gene expression was found to

be globally similar between the unbound genes and those of the "O" class (P=0.15, Wilcoxon rank-sum

test; Fig. 3b). Second, Dal80 binding at the promoter of genes correlated with significantly increased

expression levels, in comparison to the unbound genes ($P < 2.2e^{-16}$, Wilcoxon rank-sum test; Fig. 3b).

Third, the genes bound by Dal80 both at the promoter and across the ORF ("P & O" class) were even

more expressed than the genes showing promoter binding only ($P = 1.3e^{-14}$, Wilcoxon rank-sum test;

Fig. 3b). Together with the observation that genes of the "P & O" class globally showed higher Dal80-

Myc[13] ChIP-Seq signal at the promoter than those of the "P" class (Fig. 2c-d), our results indicate that

Dal80 spreading across gene bodies correlates with a strong recruitment at the promoter, higher

sensitivity to Dal80 and high expression in proline-containing medium. Consistently, metagene analysis

of Dal80-regulated genes is very similar to that of "P & O" class (Supp. Fig. 2b). Noticeably, 15% (170)

and 2% (75) of the genes belonging to the "P" and "unbound" classes respectively display expression

levels higher than median levels of expression of the genes belonging to the "P&O" class (Supp. Fig.

2c). These proportions raise even further when the first quartile (25%) of the "P&O" class is considered

(Supp. Fig. 2c), suggesting that elevated expression levels does not automatically imply binding at ORF.

In conclusion, our genome-wide analyses revealed a significant correlation between Dal80 recruitment

to the chromatin and Dal80-sensitive expression. We also showed that Dal80 binding both at the

promoter and across the gene body correlates with high levels of gene expression.

Dal80 binding across the ORF of well-characterized NCR-sensitive genes

In order to validate our genome-wide observations at the level of known NCR-sensitive genes, we

characterized the binding profiles of Dal80-Myc[13] along four NCR-sensitive genes of the "P & O" class

(see snapshots of ChIP-Seq signals in Fig. 4a-d). We chose the NAD$^+$-dependent glutamate

dehydrogenase gene (*GDH2*) and three permease-coding genes (*GAP1*, *DAL5* and *MEP2*), aware that

their regulation could differ (Isabelle Georis, Tate, Cooper, & Dubois, 2011; Godard et al., 2007;

Scherens et al., 2006) and that GATA sites are differently represented throughout these genes (Fig. 4a-

d).

As expected from our genome-wide observations, Dal80 bound not only the promoter, but also the

coding region of *DAL5*, *GAP1*, *GDH2* and *MEP2* in proline-grown cells (Fig. 4e and Supp. Fig. 3a). Binding

of Dal80 to the *GDH2* ORF was reduced (Fig. 4e), which is consistent with the ChIP-Seq profile (Fig. 4d).

We ruled out any erroneous interpretation of our data upon showing that, using the same

experimental design, the Gal4-Myc[13] protein only occupied the *GAL10* promoter, and not the coding

region (Supp. Fig. 3b). Moreover, intragenic GATA factor binding did not rely on an intermediate RNA

molecule, since the Dal80-Myc[13] signal across the *MEP2* gene body was unaffected upon RNAse

treatment of the chromatin extracts before the immunoprecipitation (Fig. 4f and Supp. Fig. 3c).

The results of the ChIP-qPCR show that the level of Dal80 binding also varied from one gene to another.

As striking example, Dal80 was more present at the *GAP1* and *MEP2* ORFs, which are more highly

expressed than *DAL5* in proline-grown cells (Supp. Table 5a). This is consistent with the correlation

observed at the genomic level between Dal80 binding both to the promoter and across gene bodies

and high expression levels (Fig. 3b). Altogether, these observations prompted an important

mechanistic question: how can Dal80 be localized to gene bodies ?

Dal80 binding within gene bodies occurs independently of intragenic GATA sites

Like most NCR-sensitive genes, the four genes described above are all characterized by stretches of

GATA sequences in their promoters (Fig. 4a-d). A possible explanation for Dal80 binding across gene

bodies could be that functional, intragenic GATA sequences, directly recruit Dal80. Given the low

complexity of the GATA consensus, the probability that a gene is devoid of intragenic GATA site is very

low. Indeed, only 10% of all yeast ORFs have no GATA site, and most genes contain an average of 5

intragenic GATA sites (Supp. Fig. 3c). However, according to current models, GATA sites should be

present in stretches of at least two in order to be functional *in vivo*, which is not the case within *DAL5*,

GDH2, *GAP1* and *MEP2* ORFs, that contain some GATA sites, but relatively distant from each other.

Interestingly, 23 of the 144 P&O-bound genes did not contain any GATA site in their ORF, hence

showing that Dal80 spreading across the body of genes does not necessarily require intragenic GATA

sites, at least for these genes.

In order to test if the presence of a NCR-sensitive promoter could confer intragenic Dal80 binding

across the body of a non-NCR-sensitive gene, devoid of any GATA cluster, we placed the *URA3* ORF

under the control of the *MEP2* promoter, at the *MEP2 locus* (P_{MEP2}-*URA3*; Fig. 5a). Only two GATA sites

are present in the *URA3* ORF, and separated by 300bp. In this strain, the expression of *URA3* became

NCR-sensitive and followed wild type *MEP2* expression (Supp. Fig. 4a), correlating with PolII

recruitment over the *URA3* ORF (Supp. Fig. 4b). In these conditions, we observed Dal80-Myc[13] binding

at the promoter of *MEP2* and also across *URA3* (Fig. 5b). Importantly, Dal80 was not detected across

URA3 when it was expressed from its native *locus*, under the control of its promoter (Fig. 5b). These

results show that significant recruitment of Dal80 at an ORF region must be preceded by promoter

recruitment.

Finally, we analyzed Dal80 binding at the *UGA4 locus*, another well-characterized NCR-sensitive gene,

which is devoid of intragenic GATA site clusters (Fig. 5c). *UGA4* expression is induced by GABA (γ-

aminobutyric acid) and is strongly repressed by Dal80 in the absence of the inducer (André, Hein,

Grenson, & Jauniaux, 1993). Therefore, to derepress *UGA4* without inducer and to analyze Dal80

binding along *UGA4*, we deleted the repressive C-terminal leucine zipper domain of Dal80 (Dal80ΔLZ-

Myc[13]), known for its homodimerization and heterodimerization with the Gzf3 transcriptional

repressor (Coornaert et al., 1992; Svetlov & Cooper, 1998)(Fig. 5d). As a control experiment, *UGA4*

expression was quantified in Dal80-Myc[13] and Dal80ΔLZ-Myc[13] cells. As expected, *UGA4* mRNA level

was very low in Dal80-Myc[13] cells, whereas deletion of Dal80 leucine zipper led to a strong

derepression of *UGA4* (Supp. Fig. 4c). Analysis of PolII occupancy across *UGA4* using ChIP confirmed

that elevated mRNA levels correlate with increased transcription (Supp. Fig. 4d). In these conditions,

full-length Dal80-Myc[13] binding was restricted to the *UGA4* promoter, while Dal80ΔLZ-Myc[13] binding

was detected at the promoter and across the body of *UGA4* (Fig. 5e), correlating with the elevated

transcript and PolII levels. Surprisingly, the leucine zipper of Dal80 and consequently, its dimerization,

needed for *UGA4* repression, were not required for its spreading across *UGA4* gene body. Importantly,

these results show that promoter binding is not sufficient to confer intragenic binding, but suggest that

transcription activation is required.

Together, these data indicate that Dal80 binding at the promoter and across the body of NCR-sensitive

genes correlated with expression levels and required transcription activation.

Transcription elongation is necessary for Dal80 binding across *MEP2*

We hypothesized that the elongating PolII complex could be responsible for Dal80 spreading beyond

Dal80-bound promoters.

To assess whether the transcriptionally active PolII complex is necessary for Dal80 binding across the

MEP2 gene, we used an *rpb1-1* strain, known to rapidly involve a cessation of PolII activity when cells

are shifted to a restrictive temperature of 37°C (T. S. Kim et al., 2010; Nonet, Scafe, Sexton, & Young,

1987). We analyzed Dal80-Myc[13] binding along *MEP2* in WT and *rpb1-1* cells. When *rpb1-1* cells were

shifted at 37°C for 1hr, *MEP2* mRNA and PolII levels showed a 2-fold (Supp. Fig. 5a) and >10-fold

decrease (Supp. Fig. 5b), respectively, reflecting the expected elongation defect when *rpb1-1* cells are

shifted in non-permissive conditions. In the same conditions, we observed a significant >5-fold

reduction of Dal80-Myc[13] levels across the *MEP2* ORF, while the binding at the promoter was not

affected (Fig. 6a-b). This result suggests that Dal80 spreading across the body of NCR-sensitive genes

would be a consequence of the PolII-driven elongation process.

To get insights into the mechanism by which Dal80 associates to gene bodies, we tested whether it

physically interacts with the transcriptionnaly engaged form of PolII (Fig. 6c). Total protein extracts

from Dal80-Myc[13] cells were immunoprecipitated with antibodies directed against the PolII CTD and

its phospho-forms Ser2P and Ser5P, respectively characteristic of elongating and initiating PolII forms.

All three antibodies enabled effective immunoprecipitation, whereas no antibody and nonspecific

antibody controls generated a lower or no signal at all. Thus, Dal80 physically interacts with

phosphoforms of the PolIII, suggesting a strong association with PolII engaged in active transcription

from initiating to elongating polymerase.

Together, our data indicate that Dal80 spreading across the body of NCR-sensitive genes depends on

active transcription and that Dal80 interacts with the transcriptioanlly active form of PolII, supporting

a model where Dal80 spreading across the body of highly expressed, NCR-sensitive genes might be the

result of Dal80-PolII association at post-initiation transcription phases.

Discussion

Eukaryotic GATA factors belong to a small but important family of DNA binding proteins involved in

development and response to environmental changes in multicellular organisms in unicellular

organisms, respectively. In yeast, four GATA factors are involved in Nitrogen Catabolite Repression

(NCR), controlling gene expression in response to nitrogen source availability. One of them, the Dal80

repressor, itself NCR-sensitive, acts to modulate the intensity of NCR responses.

Over the past decade, a number of studies have screened the genome aiming at gathering an inventory

of genes regulated by the nitrogen source. Although >500 genes have been shown to be differentially

expressed upon change of the nitrogen source (Airoldi et al., 2016; Godard et al., 2007), the list of NCR-

sensitive genes was reduced to about 100, based on their sensitivity to GATA factors (Boer et al., 2007,

2003; Godard et al., 2007; Scherens et al., 2006), suggesting that the number of Dal80 targets would

be situated in that range. Here, our ChIP-Seq analysis identified more than 1200 Dal80-bound

promoters, which considerably extends the list of potential Dal80 targets. In fact, the number of Dal80-

bound promoters could even have been more important. Indeed, the GATA consensus binding site is

rather simple and short, so that in yeast, a total number of 10,000 putative binding sites can be found

in all protein-coding gene promoters, 2930 promoters having at least 2 GATA sites, which is thought

to be a prerequisite for *in vivo* binding and function of the GATA factors. The difference between the

number of promoters with ≥2 GATA sites and the number of Dal80-bound promoters suggests the

existence of a selectivity for Dal80 recruitment. This selectivity could rely on promoter architecture

and/or chromatin structure, conditioning the requirement for auxiliary DNA binding factors that would

stabilize Dal80 at some promoters. Moreover, although we observed a significant correlation between

Dal80 binding and regulation, the expression of most of the Dal80-bound genes was not affected in a

dal80Δ mutant strain. Again, Dal80-dependence for transcribing these genes, as well as their NCR

sensitivity, could require the presence of cofactors which are not produced or inactive under the tested

growth conditions. In mammals, GATA factors also display an extraordinary complexity in the

relationships between binding and expression regulation. Like Dal80, GATA-1 and GATA-2 only occupy

a small subset of their abundant binding motif throughout the genome, and the presence of the

conserved binding site is insufficient to cause GATA-dependent regulation in most instances

(Hasegawa et al., 2016). GATA-1 binding kinetics, stoichiometry and heterogenous complex

formations, conditioned by composite promoter architecture, influence its transcriptional activity and

hence diversify gene expression profiles (Hasegawa et al., 2016).

Given the high conservation at the amino acid level between the DNA binding domain of the four yeast

NCR GATA factors, it is likely that they all recognize identical sequences (GATAA, GATAAG or GATTAG).

This consensus has been largely validated in the past using gene reporter experiments, mutational

analyses and *in vitro* binding experiments on naked DNA. Nonetheless, of the 1269 bound promoters,

48% contained at least two GATA sites, a proportion that is not different from that observed among

unbound promoters, and the amount of GATA sites per promoter was not different between the two

groups either. In addition, Dal80 recruitment was found to occur independently of the presence of

GATA sites in 20% of Dal80-bound promoters, as also previously observed in mammalian cells (Cheng

et al., 2009; Fujiwara et al., 2009). Future experiments will be required to decipher the molecular

mechanism by which Dal80 can be recruited to these GATA-less promoters.

Unexpectedly, although Dal80 has always been described as a repressor, we identified 314 genes that

are positively regulated by Dal80 (their expression is significantly decreased upon Dal80 deletion).

These genes are distinct from the NCR-sensitive genes : only 4 belong to the NCR-sensitive list of

(Godard et al., 2007) and are significantly enriched in amino acid biosynthetic processes, resembling

the amino acid starvation response mediated by the Gcn4 transcriptional activator. Interestingly, the

promoter of 122/314 Dal80-activated genes contain Gcn4-binding sites (Supp. Fig. 3c), and this group

of 314 Dal80-activated genes is significantly enriched for genes regulated by the General Amino Acid

Control (GAAC; YeastMine Gene List, Publication Enrichment, $p<1.6e-13$), through the Gcn4 activator.

Interconnections between NCR and GAAC have already been demonstrated, mostly at the level of

nitrogen catabolism control: 1-a large number of non-preferential nitrogen sources leads to increased

transcription of GAAC targets (Godard et al., 2007); and 2- Gcn4 contributes, with Gln3, to the

expression of NCR-sensitive genes (Staschke et al., 2010; Valenzuela, Aranda, & González, 2001).

However, this is the first time that evidence are provided suggesting a positive function for the negative

GATA factor at the level biosynthetic gene expression.

The most striking and unexpected finding of this work is the observation that for 189 genes, Dal80

occupied the gene body ("O" and "P & O" classes). The 45 genes displaying Dal80 intragenic occupancy

with no recruitment at the promoter ("O" class) were globally neither Dal80-sensitive nor NCR-

sensitive, similar to the unbound genes. Furthermore, we also noted that a substantial fraction of these

genes correspond to small dubious ORFs, near or even overlapping the promoter, Dal80-bound, of the

adjacent gene. In these cases, the limited resolution of the ChIP-Seq technique, combined to the small

size of these genes, might have allowed them to pass through the filters of the procedure to identify

the genes showing Dal80 intragenic binding. Overall, these observations suggest that the existence of

the "O" class is likely to be physiologically irrelevant. Dal80 binding at the promoter and spreading

sacross the body of the 144 genes belonging to the "P & O" class correlated with high expression levels

and sensitivity to Dal80-dependent repression. The observation that a considerable fraction of "P"

class genes have similar or higher expression levels than most of the "P&O" class genes suggest that

the well-known artifact of hyper-chippable regions is not applicable here (Teytelman, Thurtle, Rine, &

van Oudenaarden, 2013). Our observations were also experimentally confirmed using ChIP

experiments, at the level of well-characterized NCR-sensitive genes. Our data indicate that Dal80

spreading across genes bodies does not rely on intragenic GATA sites. Rather, promoter binding

appears to be required but not sufficient: Dal80 spreading across genes depends on active

transcription. Indeed, the observation that impaired PolII elongation correlates with decreased

intragenic binding further indicates that Dal80 binding across gene bodies depends on ongoing

transcription elongation. Consistently, we detected a physical interaction between Dal80 and

transcriptionnaly active forms of PolII. Together, our data leads us to propose a model where Dal80

could travel from the promoter of highly expressed, NCR-sensitive genes through the gene body by

accompanying the elongating PolII complex (Figure 7). Additional investigations will be required to

define which domain of Dal80 is responsible for the interaction with the transcription machinery, to

determine whether there is any causal relationship between Dal80 intragenic binding and high

expression levels, and to decipher the potential role of Dal80 during transcription elongation.

Whereas the binding of elongation factors across gene bodies has been thoroughly documented (Sims,

Belotserkovskaya, & Reinberg, 2004), it has also been described for few specific transcription factors.

For example, Gal4 was reported to bind to its consensus DNA target within the *ACC1* ORF, but the

authors concluded that the observed transcriptional repression of the *ACC1* gene was most likely

resulting from random *GAL4* binding "noise" over the genome, thus having no physiological

explanation for this ORF-bound transcription factor (Li & Johnston, 2001). Likewise, Gcn4 was detected

across the *PHO8* ORF, with concomitant recruitment of the SAGA complex, but without any impact on

gene expression (Topalidou & Thireos, 2003). More recently, binding of the Gcn4 transcription factor

to its consensus site at some ORFs, when located in proximity of the transcriptional start site, was

found to play a consistent role in controlling embedded cryptic promoters in yeast, thereby affecting

Gcn4-dependent transcription of some genes (Rawal et al., 2018).

A recent study has identified CTD phosphorylation of polII as a hub that optimizes transcriptome

changes to adequately balance optimal growth and stress tolerance responses (Chasman et al., 2014).

The addition of nitrogen to nitrogen-limited cells rapidly results in the transient overproduction of

transcripts required for protein translation (stimulated growth) whereas accelerated mRNA

degradation favours rapid clearing of the most abundant transcripts, like those involved in high affinity

permease production, that are highly expressed NCR-sensitive genes, for example (Airoldi et al., 2016).

The involvement of the Nrd1-Nab3-Sen1 (NNS) and TRAMP complexes in these regulatory responses

has been envisioned very recently (Bresson, Tuck, Staneva, & Tollervey, 2017; Merran & Corden, 2017).

Thus, a possible role of Dal80 (and possibly of the other GATA factors) binding along highly expressed

genes could be to transmit nutritional signals to the RNA decay machinery for rapid and efficient

remodeling of gene expression in response to changing nitrogen supplies. Here, an attractive model

(Figure 7) would hypothesize that Dal80 binding to polII could prevent the decay of the dedicated

stress-response mRNAs to enhance their cellular life-cycle by regulating/repressing RNA decay factors

that might associate otherwise on the polymerase. Other elongation-related processes, like histone

modification and chromatin remodeling (Henikoff & Shilatifard, 2011; Tessarz & Kouzarides, 2014),

mRNA export (Meinel et al., 2013) or roadblock termination (Candelli et al., 2018) can also be

proposed.

In this respect, in human cells, GATA factors are also reported to occupy non-canonical sites within the

genome, further reinforcing that they can be recruited to the chromatin independently of their motif

(Cheng et al., 2009; Fujiwara et al., 2009). In addition, 43% of the GATA1 peaks were collected among

exon, introns and 3'UTR of coding genes in human erythroleukemia cells (Fujiwara et al., 2009). It is

tempting to hypothesize that GATA factors could have a dual or synergistic role during transcription,

i.e. recruiting/stabilizing the PIC complex as for any classical transcription factor in the

promoter/enhancer regions and promoting elongation at a post initiation step interacting with the

RNAPII. Interestingly, GATA1 in erythropoid cells interacts with the mediator (Stumpf et al., 2006), a

complex also known to travel with PolII (Allen & Taatjes, 2015).

Materials and Methods

Experimental Model and Subject Details

Experiments were conducted using *S. cerevisiae* strains of the FY genetic background. The strains used

are listed in Supp. Table 1. Dal80 was tagged with 13 copies of the *c-myc* epitope (Myc[13]) as described

(Longtine et al., 1998) using primers listed in Supp. Tables 1 and 2. The P_{MEP2}-*URA3* allele in strains

FV806-808 was created by amplification of the *URA3* gene using the same strategy, with primers listed

in Supp. Tables 1 and 2.

Cultures were grown at 29°C to mid-log phase (A_{660nm} = 0.5) in YNB (without amino acids or ammonia)

minimal medium containing the indicated nitrogen source at a 0.1% final concentration, glucose (3%)

and the appropriate supplements (20 µg/ml uracil, histidine and tryptophan) to cover auxotrophic

requirements.

Chromatin immunoprecipitation

Cell extracts and immunoprecipitations were conducted as described (I. Georis et al., 2009) using

primers listed in Supp. Table 2. The cells (100-ml cultures grown to an absorbance (A660 nm = 0.6)

corresponding to 6 × 106 cells/ml) were treated with 1% formaldehyde for 30 min at 25 °C and mixed

by orbital shaking. Glycine was then added to a final concentration of 500 mm and incubation

continued for 5 min. The cells were collected, washed once with cold 10 mm Tris-HCl, pH 8, washed

once with cold FA-SDS buffer (50 mm HEPES-KOH, pH 7.5, 150 mm NaCl, 1 mm EDTA, 1% Triton X-100,

0.1% sodium deoxycholate, 0.1% SDS, 1 mm phenylmethylsulfonyl fluoride), and resuspended in 1 ml

of cold FA-SDS buffer. An equal volume of glass beads (0.5 mm in diameter) was added, and the cells

were disrupted by vortexing for 30 min in a cold room. The lysate was diluted into 4 ml of FA-SDS

buffer, and the glass beads were discarded. The cross-linked chromatin was then pelleted by

centrifugation (17,000 × g for 35 min), washed for 60 min with FA-SDS buffer, resuspended in 1.6 ml

of FA-SDS buffer for 15 min at 4 °C, and sonicated three times for 30 s. each (Bioruptor, Diagenode) to

yield an average DNA fragment size of 700 base pairs. Finally, the sample was clarified by centrifugation

at 14,000 × g for 30 min and diluted 4-fold in FA-SDS buffer, and aliquots of the resultant chromatin

containing solution were stored at −80 °C. Myc[13]-tagged proteins were immunoprecipitated by

incubating 100 µl of the chromatin containing solution for 180 min at 4°C with 2 µl of mouse anti-Myc

antibodies (SC-40, Santa Cruz) prebound to 10 µl of Dynabeads Pan Mouse IgG (Dynal) according to

the manufacturer's instructions. Immune complexes were washed six times in FA-SDS buffer and

recovered by treating with 50 µl of Pronase Buffer (25 mm Tris, pH 7.5, 5 mm EDTA, 0.5% SDS) at 65

°C with agitation. Input (IN) and immunoprecipitated (IP) fractions were then subjected to Pronase

treatment (0.5 mg/ml; Roche Applied Science) for 60 min at 37°C, and formaldehyde cross-links were

reversed by incubating the eluates overnight at 65°C. Finally, the samples were treated with RNase (50

µg/ml) for 60 min at 37°C. DNA from the IP fractions was purified using the High Pure PCR Product

Purification Kit (Roche Applied Science) and eluted in 50 µl of 20 mm Tris buffer, pH 8. IN fractions

were boiled 10 min and diluted 500-fold with no further purification prior to quantitative PCR analysis.

Quantitative RT-PCR

Quantitative RT-PCR was performed as described previously (I. Georis et al., 2009) using primers listed

in Supp. Table 2. Total RNA was extracted from 4-ml cultures and cDNA was generated from 100 to

500 ng of total RNA using a RevertAid H Minus first-strand cDNA synthesis kit with oligo(dT)$_{18}$ primers

from Fermentas using the manufacturer's recommended protocol. cDNAs were subsequently

quantified by RT-PCR using the Maxima SYBR green qPCR master mix from Fermentas.

Co-Immunoprecipitation

Cultures (100 ml) were harvested, washed once in 50 mM Tris, pH 8, and resuspended in 1ml of buffer

(50 mM Tris, pH 8, 150 mM NaCl, 5 mM EDTA, 0.05% NP-40, 1 mM phenylmetyhlsulfonyl fluoride, and

complete protease inhibitor cocktail tablets [Roche]). Lysis was performed by shaking with 425-600

µm acid-washed glass beads (Sigma) on an IKA Vibrax VXR orbital shaker at maximum speed for 30 min

at 4°C. Cell debris and glass beads were removed by centrifugation. Immunoprecipitation was

performed by incubating 200 µl of total cell extracts with 20 µl of Dynabeads PAN mouse

immunoglobulin G (Invitrogen) that were preinucbated with anti-HA (SCBT, SC-7392), anti-CTD (SCBT,

CTD4H8), anti-Ser2P (BioLegend, H5) or anti-Ser5P (BioLegend, H14) antibodies and 20 µl of 1%

phosphate-buffered saline-bovine serum albumin for 2 h under orbital shaking (800 rpm) at 30°C.

Immune complexes were washed three times in lysis buffer, eluted by boiling in sodium dodecyl sulfate

(SDS) sample buffer, and loaded on SDS-polyacrylamide gel for anti-Myc Western blotting.

ChIP-Seq analysis and Peak-calling

ChIP-Seq analysis was performed from two biological replicates of proline-grown 25T0b (no tag) and

FV078 (Dal80-Myc[13]) cells. For each condition, libraries were prepared from 10 ng of "input" or "IP"

DNA using the TruSeq ChIP Sample Preparation Kit (Illumina). Single-read sequencing (50 nt) of the

libraries was performed on a HiSeq 2500 sequencer.

Reads were uniquely mapped to the *S. cerevisiae* S288C reference genome using Bowtie2 v2.1.0

(Langmead & Salzberg, 2012), with a tolerance of 1 mismatch in seed alignment. Tags densities were

normalized on the total number of uniquely reads mapped.

Dal80-bound regions were identified through a peak-calling procedure using version 2.0.9 of MACS

(Zhang et al., 2008), with a minimum false discovery rate (FDR) of 0.001.

Total RNA-Seq

For each strain and condition, total RNA was extracted from two biological replicates using standard

hot phenol procedure, ethanol-precipitated, resuspended in nuclease-free H_2O (Ambion) and

quantified using a NanoDrop 2000c spectrophotometer. Ribosomal RNAs were depleted from 1 μg of

total RNA using the RiboMinus™ Eukaryote v2 Kit (Life Technologies). After concentration using the

Ribominus™ Concentration Module (Life Technologies), rRNA-depleted RNA was quantified using the

Qubit RNA HS Assay kit (Life Technologies). In parallel, rRNA depletion efficiency and integrity of both

total and rRNA-depleted RNA were checked by analysis in a RNA 6000 Pico chip, in a 2100 bioanalyzer

(Agilent). Strand-specific total RNA-Seq libraries were prepared from 125 ng of rRNA-depleted RNA

using the TruSeq® Stranded Total RNA Sample Preparation Kit (Illumina), following manufacturer's

instructions. Paired-end sequencing (2 x 50 nt) of the libraries was performed on a HiSeq 2500

sequencer. Sequenced reads were mapped to the reference genome using version 2.0.6 of TopHat (D.

Kim et al., 2013), as described (Wery et al., 2016). Tags densities were normalized on the total number

of reads uniquely mapped on ORFs. Differential expression analysis was performed using DESeq

(Anders & Huber, 2010).

Quantification and Statistical Analysis

Statistical details are to be found in the figure legends. Error bars correspond to standard error.

Statistical significance tests were carried out using the Student's t test (Microsoft Excel) when

indicated.

Availability of data and materials

Sequence data can be accessed at the NCBI Gene Expression Omnibus using accession numbers

GSE86307 and GSE86325. Genome browsers for visualization of processed ChIP-Seq and RNA-Seq data

are accessible at http://vm-gb.curie.fr/dal80.

Further information and requests for resources and reagents should be directed to and will be fulfilled

by the lead contact, Isabelle Georis (igeoris@ulb.ac.be). Bioinformatics and genome wide dataset

requests could also be addressed to antonin.morillon@curie.fr for rapid processing.

Acknowledgements

This work has benefited from the facilities and expertise of the NGS platform of Institut Curie,

supported by the Agence Nationale de la Recherche (ANR-10-EQPX-03, ANR10-INBS-09-08) and the

Canceropôle Ile-de-France. A. Morillon's lab is supported by the Agence Nationale de la Recherche

(DNA-Life) and the European Research Council (EpincRNA starting grant, DARK consolidator grant).

Work by IG and ED is funded by the CoCoF and FNRS (FRFC 2.4547.11). Work by AR was funded by a

FRIA fellowship.

We are grateful to all members of our labs for discussions and critical reading of the manuscript. We

thank F. Vierendeels for excellent technical assistance.

Competing financial interest

The authors declare that they have no competing interests.

References

Airoldi, E. M., Miller, D., Athanasiadou, R., Brandt, N., Abdul-Rahman, F., Neymotin, B., … Gresham,

D. (2016). Steady-state and dynamic gene expression programs in Saccharomyces cerevisiae in

response to variation in environmental nitrogen. *Molecular Biology of the Cell*, *27*, 1–38.

https://doi.org/10.1091/mbc.E14-05-1013

Allen, B. L., & Taatjes, D. J. (2015). The Mediator complex: a central integrator of transcription.

Nature Reviews Molecular Cell Biology, *16*(3), 155–166. https://doi.org/10.1038/nrm3951

Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data. *Genome*

Biology, *11*(10), R106. https://doi.org/10.1186/gb-2010-11-10-r106

André, B., Hein, C., Grenson, M., & Jauniaux, J. C. (1993). Cloning and expression of the UGA4 gene

coding for the inducible GABA-specific transport protein of Saccharomyces cerevisiae.

Molecular & General Genetics, *237*(1–2), 17–25. https://doi.org/10.1007/BF00282779

André, B., Talibi, D., Boudekou, S. S., Hein, C., Vissers, S., & Coornaert, D. (1995). Two mutually

exclusive regulatory systems inhibit UASGATA, a cluster of 5'-GAT(A/T)A-3' upstream from the

UGA4 gene of Saccharomyces cerevisiae. *Nucleic Acids Research*, *23*(4), 558–564.

https://doi.org/10.1093/nar/23.4.558

Boer, V. M., De Winde, J. H., Pronk, J. T., & Piper, M. D. W. (2003). The genome-wide transcriptional

responses of Saccharomyces cerevisiae grown on glucose in aerobic chemostat cultures limited

for carbon, nitrogen, phosphorus, or sulfur. *Journal of Biological Chemistry*, *278*(5), 3265–3274.

https://doi.org/10.1074/jbc.M209759200

Boer, V. M., Tai, S. L., Vuralhan, Z., Arifin, Y., Walsh, M. C., Piper, M. D. W., … Daran, J. M. (2007).

Transcriptional responses of Saccharomyces cerevisiae to preferred and nonpreferred nitrogen

sources in glucose-limited chemostat cultures. *FEMS Yeast Research*, *7*(4), 604–620.

https://doi.org/10.1111/j.1567-1364.2007.00220.x

Bresnick, E. H., Katsumura, K. R., Lee, H. Y., Johnson, K. D., & Perkins, A. S. (2012). Master regulatory

GATA transcription factors: Mechanistic principles and emerging links to hematologic

malignancies. *Nucleic Acids Research*, *40*(13), 5819–5831. https://doi.org/10.1093/nar/gks281

Bresnick, E. H., Lee, H. Y., Fujiwara, T., Johnson, K. D., & Keles, S. (2010). GATA switches as

developmental drivers. *Journal of Biological Chemistry*, *285*(41), 31087–31093.

https://doi.org/10.1074/jbc.R110.159079

Bresson, S., Tuck, A., Staneva, D., & Tollervey, D. (2017). Nuclear RNA Decay Pathways Aid Rapid

Remodeling of Gene Expression in Yeast. *Molecular Cell*, *65*(5), 787–800.

https://doi.org/10.1016/j.molcel.2017.01.005

Bysani, N., Daugherty, J. R., & Cooper, T. G. (1991). Saturation mutagenesis of the UAS(NTR) (GATAA)

responsible for nitrogen catabolite repression-sensitive transcriptional activation of the

allantoin pathway genes in Saccharomyces cerevisiae. *Journal of Bacteriology*, *173*(16), 4977–

4982.

Candelli, T., Challal, D., Briand, J., Boulay, J., Porrua, O., Colin, J., & Libri, D. (2018). High-resolution

transcription maps reveal the widespread impact of roadblock termination in yeast. *The EMBO*

Journal, *37*(4), e97490. https://doi.org/10.15252/embj.201797490

Chasman, D., Ho, Y.-H., Berry, D. B., Nemec, C. M., MacGilvray, M. E., Hose, J., … Gasch, A. P. (2014).

Pathway connectivity and signaling coordination in the yeast stress-activated signaling network.

Molecular Systems Biology, *10*(11), 759. https://doi.org/10.15252/msb.20145120

Cheng, Y., Wu, W., Kumar, S. A., Yu, D., Deng, W., Tripic, T., … Hardison, R. C. (2009). Erythroid GATA1

function revealed by genome-wide analysis of transcription factor occupancy, histone

modifications, and mRNA expression. *Genome Research*, *19*(12), 2172–2184.

https://doi.org/10.1101/gr.098921.109

Coffman, J. A., Rai, R., Cunningham, T., Svetlov, V., & Cooper, T. G. (1996). Gat1p, a GATA family

protein whose production is sensitive to nitrogen catabolite repression, participates in

transcriptional activation of nitrogen-catabolic genes in Saccharomyces cerevisiae. *Molecular*

and Cellular Biology, *16*(3), 847–858.

Coffman, J. A., Rai, R., Loprete, D. M., Cunningham, T., Svetlov, V., & Cooper, T. G. (1997). Cross

regulation of four GATA factors that control nitrogen catabolic gene expression in

Saccharomyces cerevisiae. *Journal of Bacteriology*, *179*(11), 3416–3429.

Cooper, T. G., Rai, R., & Yoo, H. S. (1989). Requirement of upstream activation sequences for nitrogen

catabolite repression of the allantoin system genes in Saccharomyces cerevisiae. *Molecular and*

Cellular Biology, *9*(12), 5440–5444.

Coornaert, D., Vissers, S., Andre, B., & Grenson, M. (1992). The UGA43 negative regulatory gene of

Saccharomyces cerevisiae contains both a GATA-1 type zinc finger and a putative leucine zipper.

Current Genetics, *21*(0172–8083 (Print)), 301–307.

Cox, K. H., Pinchak, A. B., & Cooper, T. G. (1999). Genome-wide transcriptional analysis in S.

cerevisiae by mini-array membrane hybridization. *Yeast*, *15*(8), 703–713.

https://doi.org/10.1002/(SICI)1097-0061(19990615)15:8<703::AID-YEA413>3.0.CO;2-Z

Cunningham, T. S., & Cooper, T. G. (1991). Expression of the DAL80 gene, whose product is

homologous to the GATA factors and is a negative regulator of multiple nitrogen catabolic

genes in Saccharomyces cerevisiae, is sensitive to nitrogen catabolite repression. *Molecular and*

Cellular Biology, *11*(12), 6205–15. https://doi.org/10.1128/MCB.11.12.6205

Cunningham, T. S., & Cooper, T. G. (1993). The Saccharomyces cerevisiae DAL80 repressor protein

binds to multiple copies of GATAA-containing sequences (URS(GATA)). *Journal of Bacteriology*,

175(18), 5851–5861. https://doi.org/10.1128/JB.175.18.5851-5861.1993

Cunningham, T. S., Dorrington, R. A., & Cooper, T. G. (1994). The UGA4 UAS(NTR) site required for

GLN3-dependent transcriptional activation also mediates DAL80-responsive regulation and

DAL80 protein binding in Saccharomyces cerevisiae. *Journal of Bacteriology*, *176*(15), 4718–

4725.

Cunningham, T. S., Svetlov, V. V., Rai, R., Smart, W., & Cooper, T. G. (1996). Gln3p Is capable of

binding to UASNTR elements and activating transcription in Saccharomyces cerevisiae. *Journal*

of Bacteriology, *178*(12), 3470–3479.

Daugherty, J. R., Rai, R., El Berry, H. M., & Cooper, T. G. (1993). Regulatory circuit for responses of

nitrogen catabolic gene expression to the GLN3 and DAL80 proteins and nitrogen catabolite

repression in Saccharomyces cerevisiae. *Journal of Bacteriology*, *175*(1), 64–73.

Descrimes, M., Zouari, Y. Ben, Wery, M., Legendre, R., Gautheret, D., & Morillon, A. (2015). VING: a

software for visualization of deep sequencing signals. *BMC Research Notes*, *8*(1), 419.

https://doi.org/10.1186/s13104-015-1404-5

Distler, M., Kulkarni, A., Rai, R., & Cooper, T. G. (2001). Green fluorescent protein-Dal80p illuminates

up to 16 distinct foci that colocalize with and exhibit the same behavior as chromosomal DNA

proceeding through the cell cycle of saccharomyces cerevisiae. *Journal of Bacteriology*, *183*(15),

4636–4642. https://doi.org/10.1128/JB.183.15.4636-4642.2001

Fujiwara, T., O'Geen, H., Keles, S., Blahnik, K., Linnemann, A. K., Kang, Y. A., … Bresnick, E. H. (2009).

Discovering Hematopoietic Mechanisms through Genome-wide Analysis of GATA Factor

Chromatin Occupancy. *Molecular Cell*, *36*(4), 667–681.

https://doi.org/10.1016/j.molcel.2009.11.001

Georis, I., Feller, A., Vierendeels, F., & Dubois, E. (2009). The Yeast GATA Factor Gat1 Occupies a

Central Position in Nitrogen Catabolite Repression-Sensitive Gene Activation. *Molecular and*

Cellular Biology, *29*(13), 3803–3815. https://doi.org/10.1128/MCB.00399-09

Georis, I., Tate, J. J., Cooper, T. G., & Dubois, E. (2011). Nitrogen-responsive regulation of GATA

protein family activators Gln3 and Gat1 occurs by two distinct pathways, one inhibited by

rapamycin and the other by methionine sulfoximine. *Journal of Biological Chemistry*, *286*(52),

44897–44912. https://doi.org/10.1074/jbc.M111.290577

Georis, I., Tate, J. J., Vierendeels, F., Cooper, T. G., & Dubois, E. (2015). Premature termination of

GAT1 transcription explains paradoxical negative correlation between nitrogen-responsive

mRNA, but constitutive low-level protein production. *RNA Biology*, *12*(8), 824–837.

https://doi.org/10.1080/15476286.2015.1058476

Godard, P., Urrestarazu, A., Vissers, S., Kontos, K., Bontempi, G., van Helden, J., & Andre, B. (2007).

Effect of 21 Different Nitrogen Sources on Global Gene Expression in the Yeast Saccharomyces

cerevisiae. *Molecular and Cellular Biology*, *27*(8), 3065–3086.

https://doi.org/10.1128/MCB.01084-06

Grass, J. A., Boyer, M. E., Pal, S., Wu, J., Weiss, M. J., & Bresnick, E. H. (2003). GATA-1-dependent

transcriptional repression of GATA-2 via disruption of positive autoregulation and domain-wide

chromatin remodeling. *Proceedings of the National Academy of Sciences of the United States of*

America, *100*(15), 8811–8816. https://doi.org/10.1073/pnas.1432147100

Hahn, S., & Young, E. T. (2011). Transcriptional regulation in Saccharomyces cerevisiae: Transcription

factor regulation and function, mechanisms of initiation, and roles of activators and

coactivators. *Genetics*, *189*(3), 705–736. https://doi.org/10.1534/genetics.111.127019

Harbison, C. T., Gordon, D. B., Lee, T. I., Rinaldi, N. J., Macisaac, K. D., Danford, T. W., … Young, R. A.

(2004). Transcriptional regulatory code of a eukaryotic genome. *Nature*, *431*(7004), 99–104.

https://doi.org/10.1038/nature02800

Hasegawa, A., Kaneko, H., Ishihara, D., Nakamura, M., Watanabe, A., Yamamoto, M., … Shimizu, R.

(2016). GATA1 binding kinetics on conformation-specific binding sites elicit differential

transcriptional regulation. *Molecular and Cellular Biology*, *36*(16), 2151–2167.

https://doi.org/10.1128/MCB.00017-16

Henikoff, S., & Shilatifard, A. (2011). Histone modification: Cause or cog? *Trends in Genetics*, *27*(10),

389–396. https://doi.org/10.1016/j.tig.2011.06.006

Horak, C. E., Mahajan, M. C., Luscombe, N. M., Gerstein, M., Weissman, S. M., & Snyder, M. (2002).

GATA-1 binding sites mapped in the beta-globin locus by using mammalian chIp-chip analysis.

Proceedings of the National Academy of Sciences of the United States of America, *99*(5), 2924–

2929. https://doi.org/10.1073/pnas.052706999

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., & Salzberg, S. L. (2013). TopHat2: accurate

alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome*

Biology, *14*(4), R36. https://doi.org/10.1186/gb-2013-14-4-r36

Kim, T. S., Liu, C. L., Yassour, M., Holik, J., Friedman, N., Buratowski, S., & Rando, O. J. (2010). RNA

polymerase mapping during stress responses reveals widespread nonproductive transcription in

yeast. *Genome Biology*, *11*(7), R75. https://doi.org/10.1186/gb-2010-11-7-r75

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,

9(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lee, T. I. (2002). Transcriptional Regulatory Networks in Saccharomyces cerevisiae. *Science*,

298(5594), 799–804. https://doi.org/10.1126/science.1075090

Levine, M., & Tjian, R. (2003). Transcription regulation and animal diversity. *Nature*, *424*(6945), 147–

151. https://doi.org/10.1038/nature01763

Li, Q., & Johnston, S. A. (2001). Are All DNA Binding and Transcription Regulation by an Activator

Physiologically Relevant? *Molecular and Cellular Biology*, *21*(7), 2467–2474.

https://doi.org/10.1128/MCB.21.7.2467-2474.2001

Longtine, M. S., McKenzie, A., Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., … Pringle, J. R.

(1998). Additional modules for versatile and economical PCR-based gene deletion and

modification in Saccharomyces cerevisiae. *Yeast*, *14*(10), 953–961.

https://doi.org/10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

Lowry, J. A., & Atchley, W. R. (2000). Molecular Evolution of the GATA Family of Transcription

Factors: Conservation Within the DNA-Binding Domain. *Journal of Molecular Evolution*, *50*(2),

103–115. https://doi.org/10.1007/s002399910012

MacIsaac, K. D., Wang, T., Gordon, D. B., Gifford, D. K., Stormo, G. D., & Fraenkel, E. (2006). An

improved map of conserved regulatory sites for Saccharomyces cerevisiae. *BMC Bioinformatics*,

7, 113. https://doi.org/10.1186/1471-2105-7-113

Magasanik, B., & Kaiser, C. A. (2002). Nitrogen regulation in Saccharomyces cerevisiae. *Gene*, *290*(1–

2), 1–18. https://doi.org/10.1016/S0378-1119(02)00558-9

Meinel, D. M., Burkert-Kautzsch, C., Kieser, A., O'Duibhir, E., Siebert, M., Mayer, A., … Sträßer, K.

(2013). Recruitment of TREX to the Transcription Machinery by Its Direct Binding to the

Phospho-CTD of RNA Polymerase II. *PLoS Genetics*, *9*(11), e1003914.

https://doi.org/10.1371/journal.pgen.1003914

Merran, J., & Corden, J. L. (2017). Yeast RNA-Binding Protein Nab3 Regulates Genes Involved in

Nitrogen Metabolism. *Mol. Cell. Biol.*, *37*(18), 1–11.

Minehart, P. L., & Magasanik, B. (1991). Sequence and expression of GLN3, a positive nitrogen

regulatory gene of Saccharomyces cerevisiae encoding a protein with a putative zinc finger

DNA-binding domain. *Molecular and Cellular Biology*, *11*(12), 6216–6228.

https://doi.org/10.1128/MCB.11.12.6216

Nonet, M., Scafe, C., Sexton, J., & Young, R. (1987). Eucaryotic RNA polymerase conditional mutant

that rapidly ceases mRNA synthesis. *Molecular and Cellular Biology*, *7*(5), 1602–1611.

https://doi.org/10.1128/MCB.7.5.1602.Updated

Omichinski, J. G., Clore, G. M., Schaad, O., Felsenfeld, G., Trainor, C., Appella, E., … Gronenborn, A. M.

(1993). NMR structure of a specific DNA complex of Zn-containing DNA binding domain of

GATA-1. *Science (New York, N.Y.)*, *261*(5120), 438–446.

https://doi.org/10.1126/science.8332909

Rai, R., Genbauffe, F. S., Sumrada, R. A., & Cooper, T. G. (1989). Identification of sequences

responsible for transcriptional activation of the allantoate permease gene in Saccharomyces

cerevisiae. *Molecular and Cellular Biology*, *9*(2), 602–608.

Rawal, Y., Chereji, R., Valabhoju, V., Qiu, H., Ocampo, J., Clark, D., & Hinnebusch, A. G. (2018). Gcn4

Binding in Coding Regions Can Activate Internal and Canonical 5' Promoters in Yeast. *Molecular*

Cell, *70*(2), 297–311. https://doi.org/10.1016/j.molcel.2018.03.007

Rowen, D. W., Esiobu, N., & Magasanik, B. (1997). Role of GATA factor Nil2p in nitrogen regulation of

gene expression in Saccharomyces cerevisiae. *Journal of Bacteriology*, *179*(11), 3761–3766.

Scherens, B., Feller, A., Vierendeels, F., Messenguy, F., & Dubois, E. (2006). Identification of direct

and indirect targets of the Gln3 and Gat1 activators by transcriptional profiling in response to

nitrogen availability in the short and long term. *FEMS Yeast Research*, *6*(5), 777–791.

https://doi.org/10.1111/j.1567-1364.2006.00060.x

Sims, R. J., Belotserkovskaya, R., & Reinberg, D. (2004). Elongation by RNA polymerase II: The short

and long of it. *Genes and Development*, *18*(20), 2437–2468.

https://doi.org/10.1101/gad.1235904

Soussi-Boudekou, S., Vissers, S., Urrestarazu, A., Jauniaux, J. C., & André, B. (1997). Gzf3P, a fourth

GATA factor involved in nitrogen-regulated transcription in Saccharomyces cerevisiae.

Molecular Microbiology, *23*(6), 1157–1168. https://doi.org/10.1046/j.1365-

2958.1997.3021665.x

Stanbrough, M., & Magasanik, B. (1996). Two transcription factors, Gln3p and Nil1p, use the same

GATAAG sites to activate the expression of GAP1 of Saccharomyces cerevisiae. *Journal of*

Bacteriology, *178*(8), 2465–2468.

Starich, M. R., Wikström, M., Arst, H. N., Clore, G. M., & Gronenborn, A. M. (1998). The solution

structure of a fungal AREA protein-DNA complex: an alternative binding mode for the basic

carboxyl tail of GATA factors. *Journal of Molecular Biology*, *277*(3), 605–620.

https://doi.org/10.1006/jmbi.1998.1625

Staschke, K. A., Dey, S., Zaborske, J. M., Palam, L. R., McClintick, J. N., Pan, T., … Wek, R. C. (2010).

Integration of general amino acid control and Target of Rapamycin (TOR) regulatory pathways

in nitrogen assimilation in yeast. *Journal of Biological Chemistry*, *285*(22), 16893–16911.

https://doi.org/10.1074/jbc.M110.121947

Stumpf, M., Waskow, C., Krötschel, M., van Essen, D., Rodriguez, P., Zhang, X., … Borggrefe, T. (2006).

The mediator complex functions as a coactivator for GATA-1 in erythropoiesis via subunit

Med1/TRAP220. *Proceedings of the National Academy of Sciences of the United States of*

America, *103*(49), 18504–18509. https://doi.org/10.1073/pnas.0604494103

Svetlov, V. V., & Cooper, T. G. (1998). The Saccharomyces cerevisiae GATA factors Dal80p and Deh1p

can form homo- and heterodimeric complexes. *Journal of Bacteriology*, *180*(21), 5682–5688.

Tai, S. L., Boer, V. M., Daran-Lapujade, P., Walsh, M. C., de Winde, J. H., Daran, J. M., & Pronk, J. T.

(2005). Two-dimensional transcriptome analysis in chemostat cultures. *J Biol Chem*, *280*(1),

437–447.

Tessarz, P., & Kouzarides, T. (2014). Histone core modifications regulating nucleosome structure and

dynamics. *Nature Reviews Molecular Cell Biology*, *15*(11), 703–708.

https://doi.org/10.1038/nrm3890

Teytelman, L., Thurtle, D. M., Rine, J., & van Oudenaarden, A. (2013). Highly expressed loci are

vulnerable to misleading ChIP localization of multiple unrelated proteins. *Proceedings of the*

National Academy of Sciences, *110*(46), 18602–18607.

https://doi.org/10.1073/pnas.1316064110

Topalidou, I., & Thireos, G. (2003). Gcn4 occupancy of open reading frame regions results in the

recruitment of chromatin-modifying complexes but not the mediator complex. *EMBO Reports*,

4(9), 872–876. https://doi.org/10.1038/sj.embor.embor931

Tripic, T., Deng, W., Cheng, Y., Zhang, Y., Vakoc, C. R., Gregory, G. D., … Blobel, G. A. (2009). SCL and

associated proteins distinguish active from repressive GATA transcription factor complexes.

Blood, *113*(10), 2191–2201. https://doi.org/10.1182/blood-2008-07-169417

Valenzuela, L., Aranda, C., & González, A. (2001). TOR modulates GCN4-dependent expression of

genes turned on by nitrogen limitation. *Journal of Bacteriology*, *183*(7), 2331–2334.

https://doi.org/10.1128/JB.183.7.2331-2334.2001

van Helden, J., André, B., & Collado-Vides, J. (1998). Extracting regulatory sites from the upstream

region of yeast genes by computational analysis of oligonucleotide frequencies. *Journal of*

Molecular Biology, *281*(5), 827–842. https://doi.org/10.1006/jmbi.1998.1947

Venters, B. J., & Pugh, B. F. (2009). A canonical promoter organization of the transcription machinery

and its regulators in the Saccharomyces genome. *Genome Research*, *19*(3), 360–371.

https://doi.org/10.1101/gr.084970.108

Wery, M., Descrimes, M., Vogt, N., Dallongeville, A. S., Gautheret, D., & Morillon, A. (2016).

Nonsense-Mediated Decay Restricts LncRNA Levels in Yeast Unless Blocked by Double-Stranded

RNA Structure. *Molecular Cell*, *61*(3), 379–392. https://doi.org/10.1016/j.molcel.2015.12.020

Wu, J., Zhang, N., Hayes, A., Panoutsopoulou, K., & Oliver, S. G. (2004). Global analysis of nutrient

control of gene expression in Saccharomyces cerevisiae during growth and starvation. *Proc Natl*

Acad Sci USA, *101*(9), 3148–3153.

Zambelli, F., Pesole, G., & Pavesi, G. (2013). Motif discovery and transcription factor binding sites

before and after the next-generation sequencing era. *Briefings in Bioinformatics*, *14*(2), 225–

237. https://doi.org/10.1093/bib/bbs016

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., … Liu, X. S. (2008).

Model-based Analysis of ChIP-Seq (MACS). *Genome Biology*, *9*(9), R137.

https://doi.org/10.1186/gb-2008-9-9-r137

Figure Legends

Figure 1. Genome-wide identification of Dal80-bound promoters.

a. Overview of the ChIP-Seq analysis. Biological duplicates of FV78 (Dal80-Myc[13]) and 25T0b (no tag) cells were grown to mid-log phase in proline-containing medium, and then harvested. After chromatin extraction and sonication, Dal80-Myc[13] was immunoprecipitated using α-Myc antibody. Co-precipitated DNA fragments were purified and used to construct ChIP-Seq libraries. After sequencing of the libraries, signals were computed using uniquely mapped reads. Dal80-bound regions were identified using a peak-calling procedure using MACS2.

b. Identification of Dal80-bound promoters. After peak-calling, Dal80-bound promoters were identified on the basis of a >75% overlap of the -100 to -350 regions (relative to the downstream ORF start site) by the peak (represented in as a red dashed line).

c. Heatmap view of the ChIP-Seq signal in the ATG +/- 500 bp region for the 1269 genes identified as bound by Dal80 at the promoter, in untagged (left) and Dal80-Myc[13] cells (right). The colour goes from yellow to red as the signal increases.

d. Metagene view of the ChIP-Seq signal along the ATG +/- 600 bp region for the 1269 genes identified as bound by Dal80 at the promoter (solid lines) and for the unbound genes (dashed lines), in untagged

(black) and Dal80-Myc[13] cells (blue). For each group of genes, normalized coverage (tag/nt) for each

gene was piled up, and the average signal was plotted. The shading surrounding each line denotes the

95% confidence interval.

e. Snapshot of ChIP-Seq signals along the *FUR4 locus*. Densities (tag/nt) are shown for the untagged

(black line) and Dal80-Myc[13] (blue line) strains. ORFs and the Dal80-Myc[13] ChIP-Seq peak are

represented by grey arrows and the red dashed line, respectively. The position (and orientation) of

each GATA site is represented by vertical segments above (sense GATA sites) or below (antisense GATA

sites) the *locus* line. The snapshot was produced using the VING software (Descrimes *et al.*, 2015).

f. Number of GATA sites in the promoter of Dal80-unbound (4522) and promoter-bound (1268) genes.

The analysis was performed using RSAT (Medina-Rivera *et al.*, 2015), across the -500 to -1 region

(relative to the ATG codon of the downstream ORF).

Figure 2. Dal80 binds the intragenic region of a subset of genes.

a. Identification of genes showing Dal80 intragenic binding.

b. Venn diagram showing the number of genes that are bound by Dal80 at the promoter and/or across

the ORF.

c. Metagene view of the ChIP-Seq signal along the ATG +/- 600 bp region for the 1125 genes identified

as bound by Dal80 at the promoter only ("P"), in untagged (black) and Dal80-Myc[13] cells (blue).

MetaSignal computation was as described in Fig. 1d. The shading surrounding each line denotes the

95% confidence interval.

d. Same as above for the 144 genes bound by Dal80 at the promoter and across the ORF ("P & O").

e. Same as above for the 45 genes bound by Dal80 across the ORF only ("O").

f. Same as above for the 4484 genes that are not bound by Dal80.

Figure 3. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels.

a. Scatter plot of densities (tag/nt, $\log_2$ scale) for genes in WT and *dal80Δ* cells grown in proline-

containing medium. Total RNA was extracted from exponentially growing biological replicates of 25T0b

(WT) and FV080 (*dal80Δ*). After rRNA depletion, strand-specific RNA-Seq libraries were constructed

and then sequenced. Tag densities were computed using uniquely mapped reads. Dal80-regulated

genes were identified on the basis of a min. 2-fold change (up or down) in the *dal80Δ* mutant relative

to the WT control, with a *P* <0.05 upon differential expression analysis using DESeq (Anders & Huber,

2010). Unaffected (n=5252), Dal80-repressed (n=232) and Dal80-activated (n=314) genes are shown

as grey, red and blue dots, respectively.

b. Box-plot of densities (tag/nt, $\log_2$ scale) in the WT strain grown in proline-containing medium, for

genes of the unbound, "O", "P" and "P & O" classes. *P*-values obtained upon Wilcoxon rank-sum test

are indicated.

Figure 4. Dal80 binding across the ORF of well-characterized NCR-sensitive genes.

a-d. Position of GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites-

the *locus* line) and qPCR probes (black and grey lettered segments) along 4 NCR-sensitive loci: *DAL5*

(A), *GAP1* (B), *GDH2* (C) and *MEP2* (D).

e. Cells of FV078 (wt *DAL80-MYC[13]*) strain were grown in proline as unique nitrogen source. Anti-Myc

ChIP was performed as described in Experimental procedures. Each histogram represents the average

(x10 000) of the value IP/IN (input or total chromatin) of at least 3 independent cultures on which two

IPs were performed. The associated error bars correspond to the standard error (DAL5$_{P1-P2}$, n=6;

DAL5$_{O3-O4}$, n=5; GAP1$_{P1-P2}$, n=4; GAP1$_{O9-O10}$, n=4; GDH2$_{P1-P2}$, n=4; GDH2$_{O5-O6}$, n=4; MEP2$_{P9-P10}$, n=11;

MEP2$_{O1-O2}$, n=7). Black histograms represent qPCR performed on promoter regions whereas grey ones

indicate qPCR data obtained within genes. Dotted line represents the highest no tag control levels.

Primers are described in Supp. Table 2: a, DAL5$_{P1-P2}$; b, DAL5$_{O3-O4}$; c, GAP1$_{P1-P2}$; d, GAP1$_{O9-O10}$; e, GDH2$_{P1-}$

$_{P2}$; f, GDH2$_{O5-O6}$; g, MEP2$_{P9-P10}$; h, MEP2$_{O1-O2}$.

f. ChIP analysis was conducted as in Panel E, with or without RNase treatment before

immunoprecipitation (n=3).

Figure 5. Dal80 binding within gene bodies occurs independently of intragenic GATA sites.

a. Schematic representation of the P_{MEP2}-*URA3 locus*, GATA sites (vertical segments above -sense GATA

sites- or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, *MEP2*

promoter; grey segments, *URA3* ORF).

b. Occupancy of Dal80-Myc[13] at the *MEP2* promoter and *URA3* ORF at the *MEP2* and *URA3 loci* in wild

type and P_{MEP2}-*URA3* strains. Cells of *DAL80-MYC[13]* wild type (FV078) and P_{MEP2}-*URA3* (FV808) strains

were cultivated at 29°C in the presence of proline (Pro) as unique nitrogen source. ChIP analysis was

conducted as in Fig. 4. The first histogram summarizes the data obtained by qPCR probing the promoter

of *MEP2* (a, $MEP2_{P9\text{-}P10}$), whereas the two following histograms are generated by qPCR performed with

oligos positioned in the ORF of *URA3* (b, $URA3_{O1\text{-}O2}$; c, $URA3_{O3\text{-}O4}$), either at the *URA3 locus* (in wild type

strain), or both at the *URA3* and *MEP2 loci*. Dotted line represents the highest no tag control levels.

The associated error bars correspond to the standard error (n=6).

c. Schematic representation of the *UGA4 locus*, GATA sites (vertical segments above -sense GATA sites-

or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, promoter;

grey segments, ORF).

d. Wild type and truncated (ΔLZ) versions of Dal80-Myc[13] used for the ChIP experiment at the *UGA4*

locus.

e. Occupancy of the *UGA4 locus* by Dal80-Myc[13]. Cells of FV078 (*DAL80-MYC[13]*; wt) and FV136

(*DAL80ΔLZ-MYC[13]*; ΔLZ) strains were grown at 29°C in the presence of proline (Pro) as unique nitrogen

source. ChIP analysis was conducted as described in Fig. 4. Black histograms represent qPCR performed

with promoter-specific primers (a, $UGA4_{P1\text{-}P2}$; b, $UGA4_{P9\text{-}P10}$) whereas grey ones indicate qPCR data

obtained with ORF-specific primers (c, $UGA4_{O1\text{-}O2}$; d, $UGA4_{O3\text{-}O4}$) at the *UGA4 locus*. The associated error

bars correspond to the standard error (n=6).

Figure 6. Transcription elongation is necessary for Dal80 binding across *MEP2*.

a. Schematic representation of the *MEP2 locus*, GATA sites (vertical segments above -sense GATA sites-

or below -antisense GATA sites- the *locus* line), and qPCR probe positions (black segments, promoter;

grey segments, ORF).

b. Occupancy of Dal80-Myc[13] at the *MEP2 locus* when transcription elongation is impaired. Wild type

(FV673) or *rpb1-1* (FV675) *DAL80-MYC[13]* cells were grown at 29°C in the presence of proline (Pro) as

unique nitrogen source, and shifted at 37°C for one hour at mid-log phase. ChIP analysis was conducted

as in Fig. 4. The first 2 black columns represent qPCR with 2 primer pairs within the *MEP2* promoter (a,

MEP2$_{P5-P6}$; b, MEP2$_{P13-P14}$) and the 2 following grey ones correspond to qPCR data obtained with 2

primer pairs within the *MEP2* ORF (c, MEP2$_{O11-O12}$ d, MEP2$_{O9-O10}$). The associated error bars correspond

to the standard error (n=6).

c. Coimmunoprecipitation of Dal80-Myc[13] with different phosphoforms of PolII. Total proteins were

extracted, immunoprecipitated with the indicated antibodies, and subjected to anti-Myc western blot

analysis.

Figure 7. Model summarizing the correlations observed between Dal80 binding, Dal80 dependency

and expression levels.

Supplementary Information

Supplementary Table 1.xlsx : Strains used in this study. (Nonet et al., 1987)

Supplementary Table 2.xlsx : Primers used in this study.

Supplementary Table 3.xlsx - Related to Figure 1 : Genome-wide identification of Dal80-bound

promoters. *Sheet a.* List of gene promoters bound by Dal80. *Sheet b.* Overlap of genes identified in our

screens and in previous genome-wide expression screens. *Sheet c.* GO term analysis of Dal80-bound

promoters. *Sheet d.* Whole genome GATA site contents of promoters and ORFs, correlated with Dal80

binding, regulation, and NCR status and presence of Gcn4-binding site.

Supplementary Table 4.xlsx - Related to Figure 2 : Dal80 binds the intragenic region of a subset of

genes. Inventory of gene promoters and ORFs bound by Dal80 and their intersection.

Supplementary Table 5.xlsx - Related to Figure 3 : Dal80 binding across gene bodies correlates with

Dal80-sensitive, high expression levels. *Sheet a.* List of Dal80-regulated genes on proline. *Sheet b.*

Gene ontology term analysis of the 232 genes repressed by Dal80 on proline. *Sheet c.* Gene ontology

term analysis of the 314 genes activated by Dal80 on proline.

Supplementary Figure 1.pdf - Related to Figure 1 : Genome-wide identification of Dal80-bound

promoters.

a. Box-plot of the distance between the annotated TSS and ORF start site (translation initiation codon,

ATG) for protein-coding genes.

b. Snapshot of ChIP-Seq signals along a GATA-less *locus* (*ALD6*). Densities (tag/nt) are shown for the

untagged (black line) and Dal80-Myc[13] (blue line) strains. ORFs and the Dal80-Myc[13] ChIP-Seq peak are

represented by grey arrows and the red dashed line, respectively. The snapshot was produced using

the VING software (Descrimes et al., 2015).

Supplementary Figure 2.pdf – Related to Figure 3 : Dal80 binding across gene bodies correlates with

Dal80-sensitive, high expression levels.

a. Contingency table showing the number of Dal80-regulated or unaffected genes (defined upon RNA-

Seq analysis in WT and *dal80Δ* cells) among the Dal80-bound and unbound genes (identified upon

ChIP-Seq analysis in untagged and Dal80-Myc[13] cells). Results that were experimentally observed or

that were expected in case of independence are indicated in bold and in brackets, respectively.

b. Metagene view of the ChIP-Seq signal along the 216 Dal80-regulated and -bound genes (solid lines),

and the 330 Dal80-regulated and -unbound genes (dashed lines), in the untagged (black) and Dal80-

Myc[13] (blue) strains. The shading surrounding each line denotes the 95% confidence interval.

c. Density-plot of RNA-Seq signal (tag/nt, log2 scale) in the WT strain grown in proline-containing , for

genes of the "unbound" (blue, n=4484), "P" (red, n=1125) and P&O" (black, n=144) classes. Y-axis:

proportion of genes for each class. The highlighted areas correspond to the 75 (2%) and 170 (15%)

genes of the "unbound" and "P" classes, respectively, showing a signal higher than the median of the

"P&O" class. A box-plot representation of the same RNA-Seq signals is shown on the top of the density-

plot.

d. Same as c., highlighting the 949 (21%) and 632 (56%) genes of the "unbound" and "P" classes,

respectively, showing a signal higher than the first quartile value for the "P&O" class.

Supplementary Figure 3.pdf – Related to Figure 4 : Dal80 binding across the ORF of well-

characterized NCR-sensitive genes – Control experiments.

a. Cells of FV078 (wt *DAL80-MYC[13]*) strain were grown in glutamine (Gln) or proline (Pro) as unique

nitrogen sources. Anti-Myc ChIP was performed as described in Experimental procedures. Each

histogram represents the average (x10 000) of the value IP/IN (input or total chromatin) of at least 3

independent cultures on which two IPs were performed. The associated error bars correspond to the

standard error (DAL5$_{P1-P2}$, n=6; DAL5$_{O3-O4}$, n=5; GAP1$_{P1-P2}$, n=4; GAP1$_{O9-O10}$, n=4; GDH2$_{P1-P2}$, n=4; GDH2$_{O5-}$

$_{O6}$, n=4; MEP2$_{P9-P10}$, n=11; MEP2$_{O1-O2}$, n=7). Black histograms represent qPCR performed on promoter

regions whereas grey ones indicate qPCR data obtained within genes. Primers are described in Supp.

Table 2: DAL5$_{P1-P2}$, DAL5$_{O3-O4}$, GAP1$_{P1-P2}$, GAP1$_{O9-O10}$, GDH2$_{P1-P2}$, GDH2$_{O5-O6}$, MEP2$_{P9-P10}$, MEP2$_{O1-O2}$.

b. Gal4-Myc[13] binding at the *GAL1-10* promoter. Schematic representation of the *GAL1-10 locus*.

Position of the Gal4-binding sites (vertical segments across the *locus* line) and qPCR probes (black and

grey lettered segments) are indicated. Cells of FV437 (wt *GAL4-MYC[13]*) strain were grown in YP with

glucose or galactose as unique carbon sources. Anti-Myc ChIP was performed as described in

Experimental procedures. Each histogram represents the average (x10 000) of the value IP/IN (input

or total chromatin) of 2 independent cultures. The associated error bars correspond to the standard

error (n=2). Black histograms represent qPCR performed on promoter regions whereas grey ones

indicate qPCR data obtained within genes. Primers are described in Supp. Table 2.

c. Binding of Dal80-Myc[13] at the *MEP2 locus*, in untreated or RNase-treated cells. Schematic

representation of the *MEP2 locus*. Position of the GATA sites (vertical segments above -sense GATA

sites- or below -antisense GATA sites- the *locus* line) and qPCR probes (black and grey lettered

segments) are indicated. Cells of untagged (25T0b) or FV078 (*DAL80-MYC[13]*; wt) strains were grown at

29°C in the presence of proline (Pro) as unique nitrogen source. ChIP analysis was conducted with or

without RNase treatment before immunoprecipitation. The first 3 black columns represent qPCR with

3 primer pairs within the *MEP2* promoter MEP2$_{P5-P6}$; MEP2$_{P13-P14}$; MEP2$_{P9-P10}$) and the 3 following grey

ones correspond to qPCR data obtained with 3 primer pairs within the *MEP2* ORF (MEP2$_{O1-O2}$; MEP2$_{O11-}$

$_{O12}$ MEP2$_{O9-O10}$). Each histogram represents the average (x10 000) of the value IP/IN (input or total

chromatin) of independent cultures. The associated error bars correspond to the standard error (n=3).

Supplementary Figure 4.pdf – Related to Figure 5 : Dal80 binding within gene bodies occurs

independently of intragenic GATA sites – Control experiments.

a. Expression of the non-NCR-sensitive gene *URA3* under the control of P$_{MEP2}$. Cells of wild type (25T0b)

and P$_{MEP2}$-*URA3* (FV806) strains were cultivated at 29°C in the presence of proline (Pro) as unique

nitrogen source. The expression of *URA3* is normalized to the expression of the *TBP1* gene. Each

column corresponds to the mean value of data obtained for at least 2 independent cultures and the

error bars correspond to the standard error (n=2).

b. PolII occupancy at the *URA3* and *P$_{MEP2}$-URA3 loci*. Cells of *DAL80-MYC[13]* wild type (FV078) and P$_{MEP2}$-

URA3 (FV808) strains were cultivated at 29°C in the presence of glutamine (Gln) or proline (Pro) as

unique nitrogen sources. Anti-polII (CTD4H8) ChIP was performed as described in Experimental

procedures. Each histogram represents the average (x10 000) of the value IP/IN (input or total

chromatin) of at least 2 independent cultures on which two IPs were performed. Histograms represent

qPCR performed with URA3$_{O1-O2}$ primers (n=2).

c. *UGA4* expression in a *DAL80ΔLZ* mutant strain. Cells of FV078 (*DAL80-MYC[13]*; wt) and FV136

(*DAL80ΔLZ-MYC[13]*; ΔLZ) strains were cultivated at 29°C in the presence of proline (Pro) as unique

nitrogen source. Total RNA was isolated and *TBP1*-normalized *UGA4* mRNA levels were quantified by

qRT-PCR using UGA4$_{O1-O2}$ primers. Histograms represent the averages of at least 2 independent

experiments and the associated error bars correspond to the standard error (n=3).

d. Occupancy of the *UGA4 locus* by PolII. Cells of FV078 (*DAL80-MYC[13]*; wt) and FV136 (*DAL80ΔLZ-*

MYC[13]; ΔLZ) strains were grown at 29°C in the presence of proline (Pro) as unique nitrogen source.

Anti-polII (CTD4H8) ChIP analysis was conducted as in Panel B using UGA4$_{O1-O2}$ primers (n=2).

Supplementary Figure 5.pdf – Related to Figure 6 : Transcription elongation is necessary for Dal80

binding across *MEP2* – Control experiments.

a. Effect of PolII elongation defects on *MEP2* expression. Wild type (FV673) or *rpb1-1* (FV675) *DAL80-*

MYC[13] cells were grown at 29°C in the presence of glutamine (Gln) or proline (Pro) as unique nitrogen

sources, and shifted at 37°C for one hour at mid-log phase. Total RNA was isolated and *TBP1-*

normalized *MEP2* mRNA levels were quantified by qRT-PCR using MEP2$_{O9-O10}$ primers. Histograms

represent the average of at least 2 independent experiments and the associated error bars correspond

to the standard error (n=2).

b. PolII occupancy at the *MEP2 locus*. Wild type (FV673) or *rpb1-1* (FV675) *DAL80-MYC[13]* cells were

grown at 29°C in the presence of glutamine (Gln) or proline (Pro) as unique nitrogen sources, and

shifted at 37°C for one hour at mid-log phase. ChIP analysis was conducted as in Supp. Fig. 4b using

MEP2$_{O9-O10}$ primers (n=4).

Figure 1. Genome-wide identification of Dal80-bound promoters. a. Overview of the ChIP-Seq analysis. **b.** Identification of Dal80-bound promoters. **c.** Heatmap view of the ChIP-Seq signal). **d.** Metagene view of the ChIP-Seq signal. **e.** Snapshot of ChIP-Seq signals along the *FUR4 locus*. **f**. Number of GATA sites in the promoter of Dal80-unbound (4522) and promoter-bound (1268) genes.

Figure 2. Dal80 binds the intragenic region of a subset of genes. a. Identification of genes showing Dal80 intragenic binding. **b.** Venn diagram showing the number of genes that are bound by Dal80 at the promoter and/or across the ORF. **c.** Metagene view of the ChIP-Seq signal along the ATG +/- 600 bp region for the 1125 genes identified as bound by Dal80 at the promoter only ("P"), in untagged (black) and Dal80-Myc[13] cells (blue). **d.** Same as above for the 144 genes bound by Dal80 at the promoter and across the ORF ("P & O"). **e.** Same as above for the 45 genes bound by Dal80 across the ORF only ("O"). **f.** Same as above for the 4484 genes that are not bound by Dal80.

Figure 3. Dal80 binding across gene bodies correlates with Dal80-sensitive, high expression levels. a. Scatter plot of densities (tag/nt, $\log_2$ scale) for genes in WT and $dal80\Delta$ cells grown in proline-containing medium. **b.** Box-plot of densities (tag/nt, $\log_2$ scale) in the WT strain grown in proline-containing medium, for genes of the unbound, "O", "P" and "P & O" classes.

Figure 4. Dal80 binding across the ORF of well-characterized NCR-sensitive genes. a-d. Position of GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line) and qPCR probes (black and grey lettered segments) along 4 NCR-sensitive loci: *DAL5* (A), *GAP1* (B), *GDH2* (C) and *MEP2* (D). **e.** Cells of FV078 (wt *DAL80-MYC*[13]) strain were grown in proline as unique nitrogen source. Anti-Myc ChIP was performed as described in Experimental procedures. **f.** ChIP analysis was conducted as in Panel E, with or without RNase treatment before immunoprecipitation.

Figure 5. Dal80 binding within gene bodies occurs independently of intragenic GATA sites. a. Schematic representation of the P*MEP2*-*URA3 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, *MEP2* promoter; grey segments, *URA3* ORF). **b.** Occupancy of Dal80-Myc[13] at the *MEP2* promoter and *URA3* ORF at the *MEP2* and *URA3 loci* in wild type and P*MEP2*-*URA3* strains. **c.** Schematic representation of the *UGA4 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line) and qPCR probe positions (black segments, promoter; grey segments, ORF). **d.** Wild type and truncated (ΔLZ) versions of Dal80-Myc[13] used for the ChIP experiment at the *UGA4 locus*. **e.** Occupancy of the *UGA4 locus* by Dal80-Myc[13].

a

b

c

Figure 6. Transcription elongation is necessary for Dal80 binding across *MEP2*. a. Schematic representation of the *MEP2 locus*, GATA sites (vertical segments above -sense GATA sites- or below -antisense GATA sites- the *locus* line), and qPCR probe positions (black segments, promoter; grey segments, ORF). **b.** Occupancy of Dal80-Myc[13] at the *MEP2 locus* when transcription elongation is impaired. **c.** Coimmunoprecipitation of Dal80-Myc[13] with different phosphoforms of PolII.

Figure 7. Model summarizing the correlations observed between Dal80 binding, Dal80 dependency and expression levels.