The bovine alveolar macrophage DNA methylome is resilient to infection with

Mycobacterium bovis

Running Title: Impact of mycobacterial infection on DNA methylation

O'Doherty, A.M.[1,*], Rue-Albrecht, K.[2,*], Magee, D.A.[1], Ahting, S.[3], Irwin, R.[4], Hall, T.J.[1],

Browne, J.A.[1], Nalpas, N.C.[5], Walsh, C.P.[4], Gordon, S.V.[6,7], Wojewodzic, M.W.[8] and

MacHugh, D.E.[1,7]

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University

College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[2] Kennedy Institute of Rheumatology, Nuffield Department of Orthopaedics, Rheumatology

and Musculoskeletal Sciences, University of Oxford, Headington, Oxford OX3 7FY, UK.

[3] Institute of Molecular Medicine, Trinity College Dublin, Dublin, D08 W9RT, Ireland.

[4] Genomic Medicine Research Group, Biomedical Sciences Research Institute, Centre for

Molecular Biosciences, University of Ulster, Coleraine, BT52 1SA UK.

[5] Quantitative Proteomics and Proteome Centre Tübingen, Interfaculty Institute for Cell

Biology, University of Tübingen, 72076 Tübingen, Germany.

[6] UCD School of Veterinary Medicine, University College Dublin, Belfield, Dublin, D04

V1W8, Ireland.

[7] UCD Conway Institute of Biomolecular and Biomedical Research, University College

Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[8] University of Birmingham, School of Biosciences, Edgbaston, Birmingham, B15 2TT, UK.

* These authors contributed equally to this work.

Grant Support:

The study was supported by Science Foundation Ireland (SFI) Investigator Programme

Awards (grant nos. SFI/08/IN.1/B2038 and SFI/15/IA/3154), an SFI Technology Innovation

Development Award (grant no. 13/TIDA/B2675), a European Union Framework 7 Project

Grant (No: KBBE-211602-MACROSYS), and the UCD Wellcome Trust funded

Computational Infection Biology PhD Programme (Grant no: 097429/Z/11/Z).

Keywords: tuberculosis, DNA methylation, whole genome bisulfite sequencing, WGBS,

Mycobacterium bovis

Abstract

Epigenetic modifications, such as DNA methylation, are pivotal in orchestrating gene expression patterns in various mammalian biological processes. They represent compelling mechanisms for conveying cellular response to external environmental stimuli more durably than transient transcriptional bursts/regulation events. Perturbation of the bovine alveolar macrophage (bAM) transcriptome due to *Mycobacterium bovis* infection on transcriptomic patterns has been well documented; however, the impact of this intracellular pathogen on the bAM epigenome has not been determined. For the current investigation, whole genome bisulfite sequencing (WGBS) was used to assess the effect of *M. bovis* infection on the bAM DNA methylome (5-methylcytosine). The methylomes of bAM infected with *M. bovis* ($n = 8$) were compared to those of non-infected control bAM ($n = 8$) at 24 hours post-infection (hpi). No differences in DNA methylation (CpG or non-CpG) were observed between control and infected bAM. Analysis of DNA methylation at proximal promoter regions uncovered >250 genes harbouring intermediately methylated (IM) promoters (average methylation = 33–66%). Gene ontology (GO) analysis, focusing on genes with low, intermediate or highly methylated promoters, revealed that genes with IM promoters were enriched for immune-related GO categories; this enrichment was not observed for genes in the high or low methylation groups. Targeted analysis of two non-imprinted genes in the IM category, *C1QB* and *IL2RA*, confirmed the WGBS observation. This study is the first in cattle to examine genome-wide DNA methylation at single nucleotide resolution in an important bovine cellular host-pathogen interaction model and provides evidence for intermediate promoter methylation in bAM.

Introduction

Infection with *Mycobacterium bovis*, the causative agent of bovine tuberculosis (BTB), accounts annually for more than $3 billion of losses to global agriculture through lost productivity and disease control costs. [1] There is also evidence suggesting that the burden of *M. bovis* as the cause of zoonotic tuberculosis in humans may be underestimated, [2] which highlights the need for a more detailed understanding of the impact of *M. bovis* in both cattle and humans. Unravelling host cellular processes that are perturbed or manipulated by intracellular pathogens is an important area of research in infection biology, particularly for disease control and the development of next-generation diagnostics and prognostics. In this regard, host cell epigenetic modifications induced, either as a component of the response to *M. bovis* infection, or as an immunoevasion strategy by the pathogen itself, remain to be fully elucidated. [3]

Epigenetic modifications to the genome, such as DNA methylation and histone tail modifications, in combination with RNA-mediated regulatory mechanisms are fundamental in modulating tissue-specific gene expression. [4-6] Epigenetic gene regulation represents an important framework for understanding how environmental stimuli are disseminated to the transcriptome and preserved through subsequent somatic cell divisions. [5] DNA methylation (5-methylcytosine), the most widely studied epigenetic modification, is involved in a variety of cellular processes including genomic imprinting, X-chromosome inactivation, chromosome stability and gene transcription [7] and has been proposed to be influenced by external stimuli across a wide range of biological contexts. [8-12] Therefore, we hypothesised that DNA methylation may be involved with the bovine host response to infection with *M. bovis*; this mechanism has previously been proposed for human tuberculosis caused by infection with *Mycobacterium tuberculosis*. [13]

Host epigenomic plasticity to *M. tuberculosis* has been reported previously. [14, 15] Sharma

and colleagues showed that non-CpG loci in the host genome were hypermethylated following

reduced representation bisulfite sequencing (RRBS) analysis of THP-1 macrophages (a human

monocytic cell line) infected with *M tuberculosis*. [14] In addition, Zheng *et al*. [15] demonstrated

that interleukin gene promoter sequences, and their receptors, were associated with

hypermethylation following analysis of THP-1 cells infected with clinical strains of *M.*

tuberculosis, using the human inflammatory response methyl-profiler DNA methylation PCR

array. Furthermore, it has been demonstrated that DNA methylation is associated with hypoxic

survival of *M. tuberculosis*. [16] Most recently, Doherty *et al*. reported that there were in excess of

750 differentially methylated regions between *M. bovis*-infected and healthy cattle in a study

using RRBS to examine CD4$^+$ T lymphocytes isolated from circulating blood samples. [17]

A range of other studies have highlighted the impact of invading microorganisms on host

DNA methylation patterns. For example, distinct DNA methylation changes have been observed

in macrophages infected with the intracellular protozoan *Leishmania donovani*, the causative

agent of the tropical disease visceral leishmaniasis. [18] In addition, global DNA methylation

changes have been detected in human neutrophils infected with *Anaplasma phagocytophilum*,

which causes granulocytic anaplasmosis. [19] Finally, it has been proposed that, during chronic

Helicobacter pylori infection in humans, functional *H. pylori* DNA methyltransferases enter host

epithelial cells and methylate their recognition sequences in chromosomal DNA, potentially

contributing to the pathogenesis of gastric adenocarcinoma or lymphoma of the mucosa-

associated lymphoid tissue. [20]

Our group has previously revealed the impact of *M. bovis* infection on the mammalian

alveolar macrophage gene expression, demonstrating that the bAM transcriptome is substantially

reprogrammed as a consequence of both host-driven defence responses and mycobacterial-

induced perturbation and manipulation of cellular processes. [21, 22] However, the effect of

M. bovis on the bovine host epigenome, specifically the DNA methylome of bAM, remains

unexplored. Recent work has shown that intracellular microbial infection can lead to alterations

of the host DNA methylome; therefore, for the present study we used WGBS to test the

hypothesis that bAM DNA methylation patterns are altered during the earliest stage of *M. bovis*

infection in cattle.

Materials and Methods

Ethics statement

All animal procedures were performed according to the provisions of the Cruelty to Animals Act of 1876 and EU Directive 2010/63/EU. Ethical approval was obtained from the University College Dublin Animal Ethics Committee (protocol number AREC-13-14-Gordon).

Isolation and infection of bovine alveolar macrophages

Isolation and purification of bAM from cattle was performed as previously described by our group [21, 22] and is summarized in Fig. 1. Briefly, total lung cells were harvested by pulmonary lung lavage with Hank's Balanced Salt Solution (Invitrogen, Life Technologies) following the removal of lungs from eight unrelated Holstein-Friesen male calves. Total lung cells were washed and cultured for 24 h at 37 °C in R10$^+$ media (RPMI 1640 medium supplemented with antibiotics [Invitrogen]). After incubation, cells were prepared for infection by dissociation and seeding at 5×10^5, for each biological replicate. The purity of the seeded macrophages was confirmed by flow cytometry using anti-CD14 antibody. bAM were infected with *M.bovis* strain AF2122/97 at a multiplicity of infection (MOI) of 10 bacilli per alveolar macrophage as described in detail previously. [21, 22] These studies used comparative RNA-seq-based transcriptomics and targeted quantitative assays (RT-qPCR and multiplex ELISA) of several NF-κB-inducible pro- and anti-inflammatory cytokines and chemokines, including CCL-4, IL-1β, IL-6, IL-10 and IL-12, to verify that at 24 hpi *M. bovis*-treated bAM cells were infected and had internalised bacilli. [21, 22]

Isolation of DNA and library preparation

DNA was extracted from *M. bovis*-infected and control bAM, isolated from eight animals, using the DNeasy kit (Qiagen) according to the manufacturer's recommendations. DNA was quantified using the Qubit dsDNA HS Assay Kit (ThermoFisher Scientific). Libraries were prepared for WGBS using the post–bisulfite conversion library preparation method for methylation analysis (EpiGnome™ Methyl-Seq Kit, Epicentre, Illumina) according to the manufacturer's instructions. Genomic DNA (50 ng) isolated from *M. bovis*-infected or non-infected bAM (24 hpi) was bisulfite-modified (EZ Methylation-Direct Kit, Zymo) according to the manufacturer's guidelines. DNA synthesis was performed by mixing bisulfite-converted DNA with 2 μl DNA synthesis primer, incubating at 95 °C for 5 min, cooling on ice, followed by addition of 4 μl EpiGnome DNA Synthesis Premix 0.5 μl, 100 mM DTT and 0.5 μl EpiGnome Polymerase. Reactions were incubated at 25 °C for 5 min followed by 42 °C for 30 min, then cooled to 37 °C for 2 min before addition of 1 μl Exonuclease I to each reaction. Following this, reaction mixtures were incubated at 37 °C for 10 min, 95 °C for 3 min and then held at 25 °C. DNA was di-tagged by adding 7.5 μl EpiGnome TT Premix and 0.5 μl DNA polymerase to each reaction and incubating at 25 °C for 30 min, 95 °C for 3 min and cooling to 4 °C. Tagged DNA was purified using the using the AMPure XP (1.6× beads, 40 μl) system. A PCR step was performed to generate the second strand of DNA, complete the addition of the Illumina adaptor sequences and incorporate an index sequence. 22.5 μl of di-tagged DNA was mixed with 25 μl FailSafe PCR PreMix E, 1 μl EpiGnome Forward PCR Primer, 1 μl EpiGnome Index PCR Primer and 0.5 μl FailSafe PCR Enzyme (1.25 U) and subjected to an initial denaturation of ds DNA at 95 °C for 1 min followed by 10 cycles of 95 °C for 30 sec, 55 °C for 30 sec and 68 °C for 3 min. Following PCR, the reactions were incubated at 68 °C for 7 min. EpiGnome libraries

were purified using the AMPure (1× beads, 50 µl) system to remove primer dimers. Libraries

were quantified by Qubit using the Qubit dsDNA HS Assay Kit (ThermoFisher Scientific) and

library quality was assessed on an Agilent BioAnalyzer using the High sensitivity DNA assay kit

(Agilent Technologies).

Pyrosequencing

Genomic DNA was extracted from *M. bovis*-infected and control bAM (isolated from a

parallel set of four animals to those used for WGBS) and quantified with the High sensitivity

DNA assay kit (Agilent Technologies). DNA (200 ng) was bisulfite-modified using the EZ

Methylation-Direct Kit (Zymo) and eluted in 50 µl elution buffer. Bisulfite PCR reactions were

performed in 25 µl consisting of 0.2 µm each primer, 2 mM $MgCl_2$, 1× PCR buffer (minus

magnesium), 0.2 mM dNTPs, Platinum *Taq* DNA polymerase (Invitrogen), and 3 µl bisulfite-

modified DNA. Primer sequences are detailed in Table 1. PCR cycling conditions were as

follows: 95 °C for 5 min followed by 40 cycles of 30 sec each at 95 °C; either 55 °C (*TNF,*

NFKB2 and *IL12A*) or 58 °C (*TLR2*) for 30 sec; 72 °C for 30 sec, and a final elongation step of

5 min at 72 °C. PCR products were verified by electrophoresis on a 2 % w/v agarose gel before

pyrosequencing (Pyromark Q24, Qiagen). Pyrosequencing assays were designed in-house and

carried out as previously described. [23, 24] Only pyrosequencing reactions that passed Pyromark

Q24 internal controls for bisulfite modification were included in the analysis. Two-tailed paired

sample *t*-tests were used to assess statistically significant DNA methylation between control and

M. bovis-infected samples.

Bisulfite PCR, cloning, sequencing and combined bisulfite restriction analysis

Bisulfite-converted DNA from control and infected bAM was amplified in 25 µl

reactions containing 0.2 µM primers, 1× buffer, 0.2 mM dNTPs, 2.5 U Platinum *Taq* DNA

polymerase and 3 mM $MgCl_2$. Primer sequences are detailed in Table 2. PCR cycling conditions

were as follows: 95 °C for 3 min followed by 35 cycles of 30 sec each at 95 °C; 58 °C, 72 °C for

30 sec, and a final elongation step of 5 min at 72 °C. PCR products were purified using the

Wizard clean up kit (Promega) and cloned into the pJET1.2/blunt vector (Fermentas). Insertion

of PCR products was verified by digestion with *Bgl*II and positive clones were sequenced using

conventional Sanger sequencing (Eurofins Genomics). Combined bisulfite restriction analysis

(COBRA) was carried out using *Taqα*I, and/or *Aci*I as outlined in. [25] Sequence analysis and

alignment was performed using DNAStar EditSeq, MegAlign (www.dnastar.com) and BiQ Meth

Analyzer (http://biq-analyzer.bioinf.mpi-inf.mpg.de). During this analysis, sequences with low

C-T conversion rate (< 95 %) and with a high number of sequencing errors (sequence identity

with genomic sequence less than 80%) were excluded from the alignment. Identical clones were

also excluded from the analysis.

Illumina sequencing and initial quality control

Pooled libraries were sequenced at the Michigan State University Research Technology

Support Facility. Paired-end reads (2×150 bp) were obtained by Illumina sequencing of each

pooled library on four lanes of a HiSeq 2500 sequencer, in rapid run mode. After pooling data

from all lanes, bisulfite-treated (BS) libraries yielded 45.3–67.4 million read pairs per sample;

comparably, non-bisulfite (NON-BS) libraries yielded a total of 40–50 million read pairs per

sample across all lanes (Supp. table 1).

Quality   control   of   raw   read   pairs   using   *FastQC*

(www.bioinformatics.babraham.ac.uk/projects/fastqc) revealed similar QC metrics for both

infected and control samples (Supp. table 2). Although samples from animals 1, 4, 5, and 6

raised warnings of *Over-Represented Sequences*, this warning was systematically triggered by N-

polymers in the second mate, a technical issue resolved by quality trimming. As expected, BS

libraries raised significantly more *Per Base Sequence Content* and *Per Base GC Content* than

NON-BS libraries, due to the nature of the bisulfite treatment (Supp. table 2).

Adapter and quality trimming

Stringent adapter trimming (overlap $\geq$1 bp at the 3' end of each read), and quality

trimming (Phred $\leq$ 20 from the 3' end of each read) using *Trim Galore!* [version 0.4.1]

(www.bioinformatics.babraham.ac.uk/projects/trim_galore) left 97.5–98.4 % of raw read pairs in

bisulfite-treated samples, and 93.1–94.3% of raw read pairs for NON-BS samples. However, this

trend was reversed at the nucleotide level, with 77.5–86.9 % of sequenced bases left in BS

libraries, against 84.8–87.3 % in NON-BS libraries. Notably, BS libraries generally showed

higher levels of adapter contamination (54.2–77.3 % of raw reads) relative to NON-BS libraries

(48.6–58.2 % of raw reads), based on the stringent detection rule described above. Second read

mates displayed a larger proportion of low-quality sequenced bases trimmed (7.6–13.5 % of raw

sequenced bases) relative to first mates (1.9–4.5 % of raw sequenced bases), in both BS and

NON-BS libraries (Supp. table 1).

Notably, quality control of trimmed libraries revealed a significant improvement of *Over-

Represented Sequences*, and full resolution of *Adapter Content* warnings (data not shown). As a

result of stringent adapter trimming (even a single trailing A at the 3' end was trimmed;

following Trim Galore! default settings), all samples raised warnings of *Per Base Sequence*

Content caused by the severe under-representation of A nucleotides at the 3' end of reads, a

known artefact of the stringent trimming process with no notable repercussion on the subsequent

alignments and methylation calls.

Alignment of bisulfite-treated libraries

BS libraries were aligned using *Bismark* [version 0.15.2] [26] and the *Bowtie2* aligner

[version 2.2.6] [27] in strand-specific (directional) mode to computationally bisulfite-converted

copies of the top and bottom strands of the *Bos taurus* UMD3.1 genome assembly. [28] Alignment

efficiency (*i.e.*, read pairs aligned to a unique locus) reached 59.7–68.3 %, for a total of 28.8–

42.7 million read pairs aligned uniquely per sample. Aligned reads were found evenly distributed

between top and bottom strand of the BS-converted genome (Supp. Table 3). Bismark

methylation calls revealed methylation levels in the range of 69.2–73.8 % in CpG context, for a

total of 113–156 million methylation calls per sample. In contrast, non-CpG context displayed

markedly low methylation levels (0.7–1.7 %), with orders of magnitude larger counts of

methylation calls, owing to their broader definition of methylation context (392–543 million

calls in CHG context; 1.1–1.5 billion calls in CHH context).

Deduplication of aligned bisulfite-treated libraries

Paired-end alignments where both mates aligned to the same position in the genome were

removed from the Bismark alignment output using the *deduplicate_bismark* script to mitigate the

impact of duplicate DNA fragments sequenced. This procedure discarded 6.1–18.8 % aligned

read pairs, leaving 25.5–38.7 million aligned read pairs for subsequent methylation calls (Supp.

Table 4).

Methylation calls

244   The *bismark_methylation_extractor* script was used in a two-pronged approach. First,

245 methylation calls extracted from the full sequence of aligned read pairs were used to evaluate *M*-

246 bias across the aligned mates. *M*-bias plots show the methylation proportion across each possible

247 position in the read, and reveal anomalies at any position of the sequenced reads, often found

248 toward ends of the sequenced reads. After analysis of the *M*-bias plots generated in the first pass,

249 the second call to the *bismark_methylation_extractor* script was set to ignore the first seven

250 bases at the 5' end of both read mates. Collated reports of the Bismark pipeline leading to the

251 final methylation calls are available as HTML files in Supplementary file 1.

252 *Statistical analyses*

253   Methylation calls in CpG context were imported from the individual Bismark CpG

254 reports, combined and processed in a *BSseq* container of the *bsseq* Bioconductor package

255 [version 1.9.2; www.bioconductor.org/packages/bsseq]. [29] Genomic coordinates of CpG islands

256 on all sequences (*i.e.*, unmasked sequence CpG island track) in the UMD3.1/bosTau6 assembly

257 were obtained from the UCSC Table Browser. [30]

258 *Non CpG methylation*

259   Non-CpG methylation (CHH and CHG) present in the WGBS reads were analysed using

260 the methylKit R package. [31] De-duplicated bam files produced during alignment with Bismark

261 were sorted and saved as sam formatted files. Individual CHH or CHG were imported separately.

262 These files were imported to the methylKit object using strict criteria: at least $10\times$ coverage per

263 feature and the feature must be present across all samples. This resulted in a total of 685,311 and

264 284,641 features for CHH and CHG respectively, with a median coverage of $40\times$ for CHH and

265 $41\times$ for CHG. Median coverage between all samples was used to calculate a scaling factor to

normalize the coverage across samples. Differential methylation was determined for individual

features with an overdispersion parameter included (shrinkMN) and Benjamini-Hochberg (B-H)

FDR adjustment for multiple comparisons. [32] All downstream analyses were carried out

separately for CHH and CHG methylation. Average methylation between groups was tested

using a paired *t*-test.

Expression dynamics of genes associated with chromatin configuration and DNA methylation

Differentially expressed genes in *M. bovis*-infected bAM, relative to non-infected bAM at

2, 6, 24 and 48 hpi, were previously identified by our group using RNA-seq. [22] A comprehensive

list (EPI-list) of 151 genes previously identified as being involved with histone modifications or

DNA methylation was generated from the literature (Supp. File 2). RNA-seq transcriptomics

data was mined, at each time point, using the EPI-list. Ultimately, 86 genes were identified from

the RNA-seq data (Supp. File 2). These 86 genes were denoted as genes of interest (GOI) and

their expression was determined at 2, 6, 24 and 48 hpi using the previously published lists of

differentially expressed genes ($P < 0.05$, B-H FDR-adjusted).

Gene ontology enrichment analyses

Gene ontology enrichment analyses were performed using the Bioconductor *GOseq*

software package [33] and the annotation package *org.Bt.eg.db*

(https://bioconductor.org/packages/org.Bt.eg.db). Notably, the probability weighting function

(PWF) supplied to *GOseq* was calculated without length bias for the analysis of promoters, as

those were defined in this study to a constant width of 2 kb (1.5 kb upstream and 500 bp

downstream of TSS).

Results

WGBS summary statistics

In summary, 16 individually barcoded WGBS libraries, prepared using bAM DNA extracted from eight *M. bovis*-infected and eight non-infected samples, were sequenced on an Illumina HiSeq 2500 sequencer in rapid run mode. This generated 45.3–67.4 million read pairs per sample and ~32× sequencing depth per condition (*M. bovis*-infected and non-infected bAM). These data satisfy previously defined criteria for WGBS, with respect to the number of independent biological replicates and the sequencing depth. [34] An assessment of bisulfite conversion rates was performed using non-CpG methylation according to Clark *et al.* [35] Based on this approach, conversion efficiencies were > 99% using CHH methylation values and > 98% using CHG methylation values.

When all CpG dimers in the reference bovine genome (UMD3.1) were considered— approximately 55.1 million stranded loci, including potentially unmappable CpG dimers—the 25.5–38.7 million aligned read pairs used for methylation calls led to an average 1.1–1.4 methylation call per individual strand-specific CpG dimer in each individual sample. As a result of collapsing methylation calls as unstranded CpG loci, the even distribution of aligned read pairs on both strands of the reference genome (Supp. Table 3) doubled coverage to 2.12–2.78× per unstranded CpG dimer (~27.5 million unstranded loci). Unstranded methylation calls were used from this point onwards. While the mean coverage of CpG dimers covered in at least one sample was similar (2.13–2.81×; ~27.3 million loci), CpG dimers covered in all samples was larger for each sample (4.6–5.9×; 4.7 million loci). Notably, the average coverage of all CpG dimers in known CpG islands (CGIs)—including CpG dimers with null coverage—was similar

to those latter values (3.9–5.4×; 2.9 million loci), suggesting a consistent coverage of CGIs

across all samples.

Genome-wide scan for differentially methylated regions 24 hpi

An unbiased genome-wide scan was performed to identify potential DMRs, including

only CpG loci with at least two methylation calls for at least six of the eight biological replicates

in each sample group, thereby ensuring at least 12× coverage for any CpG dimer in both sample

groups. As a comparison, the analysis was repeated after randomising samples from both

infection groups to produce a distribution of *t*-statistics under the null hypothesis. The *bsseq*

package was used to calculate *t*-statistics in a paired design for both original and randomised sets

of sample (Fig. 2A and 2B).

Potential DMRs were identified as genomic regions including at least three loci with

absolute *t*-statistics greater than 4.6 and a mean difference in methylation level (across samples

and loci) greater than 10 % between the two groups. This analysis did not reveal significant

differences in methylation between infected and non-infected bAM; regions identified in the

original data were comparable to randomised data in number of regions identified and their

properties (*e.g.*, width, number of methylation loci, sum of *t*-statistics, proportion of regions

showing increased and decreased methylation level) (Supp. Table 5).

Distribution of methylation across different genomic regions

Following the genome-wide scan, DNA methylation was determined at the following

defined functional genomic elements: gene bodies, intergenic sequences and proximal promoters.

Genomic elements were defined as described previously by Peat and colleagues [36] and the

number of regions compared in each category are outlined in Table 2. As expected for a

differentiated/somatic cell type, the majority of CpGs within intragenic sequences, gene bodies

and CpG-deficient promoters were widely methylated. [37, 38] CpG-rich promoters containing a

CpG island (CGI) or overlapping a CGI (Promoter CGI and CGI promoter) were mostly

hypomethylated (Fig. 3). Interestingly, CGIs remote from annotated gene promoters (non-

promoter CGIs) showed variable methylation—most were hypermethylated (>75% methylated,

18,581 CGIs) with 9,103 non-promoter CGIs hypomethylated (<25%).

Pyrosequencing validation of WGBS at key immune function genes

To confirm the WGBS observation that DNA methylation was not different between

control and infected bAM, 24 hpi, a small panel of key immune genes, *TNF*, *IL12A*, *TLR2* and

NFKB2, were selected for targeted analysis by pyrosequencing. Transcription of these genes has

previously been shown to be upregulated in bAM 24 hpi with *M. bovis*; [21, 22] the specific loci that

were analysed by pyrosequencing are detailed in Figure 4A. Using DNA isolated from a parallel

set of control ($n = 4$) and infected ($n = 4$) bAM samples average methylation levels at the

proximal promoter regions of *NFKB2*, *TLR2*, *IL12A* and *TNF* was determined. Methylation at all

four genes was found to be low ($< 15\%$) in both infected and non-infected bAM samples.

Statistical analysis using a paired *t*-test did not reveal significant differences ($P \geq 0.05$) in mean

methylation levels between proximal promoter regions of infected and non-infected bAM

samples (Fig. 4B), supporting the WGBS observation that *M. bovis* does not have an effect on

the CpG methylation in bAM.

Promoter methylation level and gene ontology in bovine alveolar macrophages

Leveraging the absence of significant DMRs between infected and non-infected bAM

samples, methylation calls were pooled across all sixteen samples (eight infected and eight

controls) to analyse methylation levels in bAM gene promoters, with maximal coverage. In this

analysis, promoters were defined as regions spanning 1.5 kb upstream and 500 bp downstream of

each transcription start site (TSS), with a minimum of 10 CpGs each associated with at least five

methylation calls were included (26.8 million loci). Of the 24,616 genes annotated in the bovine

genome (Ensembl BioMart March 2016 archive), 22,964 were retained for this analysis, on the

basis that their promoter contained at least ten loci, each locus having at least five methylation

calls. For those genes, mean promoter methylation was estimated and summarised, alongside

average gene body methylation, in Figure 2C.

Notably, the vast majority of gene promoters were found at either extreme of the

methylation range. Indeed, 18,438 promoters (80.3 %) display methylation levels greater than

75 % or lower than 25 % ($8,145 \geq 75\%$ methylated; $10,293 \leq 25\%$ methylated). However, 2,580

promoters (9.7 %) displayed an average intermediate methylation level (IM, 33–66 %).

Strikingly, gene ontology (GO) analysis of the genes associated with IM promoters (33–66 %)

using the *GOseq* package [33] revealed a marked enrichment for immune-related GO categories

including "defense response" ($P < 10^{-08}$), "defense response to bacterium" ($P < 10^{-07}$), "response

to bacterium" ($P < 10^{-07}$), "chemokine-mediated signaling pathway" ($P < 10^{-06}$) and "chemokine

activity" ($P < 10^{-04}$), among others (Supp. Table 6). In contrast, no significant enrichment for

immune-related GO categories was found for promoters with methylation levels 0–1 % (759

promoters), 0–10 % (5,605), 10–20% (3,255), or 90–99 % (1,997) (Supp. Table 6). Instead, the

latter only suggested enrichment for generic GO categories (*e.g.*, "intracellular organelle",

"transcription regulatory region DNA binding").

A hallmark of some imprinted genes is that they contain a 5' differentially methylated

region that is IM (resulting from parent-of-origin specific methylation patterns); therefore, the

IM   promoter   list   was   interrogated   for   known   bovine   imprinted   genes

(www.geneimprint.com/site/genes-by-species.Bos+taurus). This analysis confirmed IM at the

promoters of the following imprinted genes; *PLAGL1*, *SNRPN*, *MEST*, *PEG10*, *GNAS* and *NNAT*

(Supp. Fig. 1).

Targeted analysis of intermediately methylated (IM) gene promoters

To confirm the presence of IM at immune gene promoters, COBRA and clonal analysis

of bisulfite PCR products was performed. Firstly, proximal promoter alignment plots for the IM

group were visually screened to remove promoters that were included due to averaging of

sequences with high and low methylation (example of this in figure 5). This analysis was

restricted to IM promoters containing a minimum of 30 CpGs (1,034 loci), to ensure sufficient

CpG coverage during COBRA and clonal bisulfite sequencing analysis. Of the 1,034 IM

promoters, 267 promoters remained in the IM group and 60/267 (22.5%) of them had a promoter

CGI (Supp. File 3). Two immune-related gene promoters with the highest CpG content, *C1QB*

and *IL2RA*, were selected for further analysis (Fig. 6). Clonal analysis revealed that, although

there are clearly hypermethylated and hypomethylated *C1QB* and *IL2RA* alleles, the prominent

allelic methylation pattern is mosaic (Fig. 7 and 8); suggesting that the IM promoters analysed

are methylated in an allele-independent as opposed to an allele-specific pattern, an observation

that has been previously reported. [39] To further confirm our WGBS and clonal bisulfite

sequencing results we carried out COBRA on *C1QB* and *IL2RA* IM regions, using *M. bovis*-

infected and non-infected bAM (Fig. 7 and 8). Results from COBRA support our observation

that the *C1QB* and *IL2RA* proximal promoters were IM. Additionally, bovine sperm, kidney,

liver and heart samples were assessed using COBRA to determine whether IM might be tissue-

specific. COBRA indicated that *IL2RA* was almost completely methylated in sperm and

predominantly methylated in the kidney, liver and heart; suggesting a potential tissue-specific IM

in bAM (Fig. 8). This possible tissue-specific IM pattern was not observed at the *C1QB* locus

(Fig. 7).

Non-CpG methylation analysis

Overall, we found a low level of methylation in the context of CHH: mean values of 0.98

% and 0.96 % were estimated for control and *M. bovis*-infected bAM, respectively (Fig. 9). CHH

methylation was not different between control and infected-bAM (*t*-statistic 1.32, df = 12.7,

$P > 0.05$). Similarly, for CHG methylation we found an overall low mean methylation of 1.53 %

and 1.49 % for control and *M. bovis*-infected bAM, respectively (Fig. 9). There was no

difference for this mean methylation in CHH context between groups (*t*-statistic 1.26, df = 13.5,

$P > 0.05$). Neither clustering on all data nor top 5,000 most variable features revealed any

patterns in these data sets for CHH and CHG methylation. This was also concordant with

differential methylation tests showing no loci as significantly differentially methylated between

groups by a methylation difference greater than 1 % and q-value = 0.01.

Relationship between M. bovis *infection and expression of chromatin and DNA modifiers*

Based on our WGBS results, bAM DNA methylation is not affected by infection with

M. bovis; therefore, we next determined whether *M. bovis* infection has an effect on chromatin.

To do this, transcription analysis of chromatin and DNA modifying enzymes was carried out

using our previously published RNA-seq data from *M. bovis*-infected bAM [22] and a similar

approach to that detailed by Nestorov and colleagues. [40] A list of 151 genes (EPI-list) that encode

chromatin and DNA modifying enzymes was assembled from the literature (Supp. File 2). To

identify chromatin and DNA modifying-associated genes that were detected by RNA-seq,

differentially expressed genes ($P < 0.05$, B-H FDR-adjusted) at each time point were compiled

and searched using the list of 151 known genes. This identified a list of 86 genes of interest

(GOI). The number of GOIs was determined at each time point and the results were as follows:

2 hpi 0/86, 6 hpi 8/86 (3 upregulated, 5 downregulated), 24 hpi 37/86 (16 upregulated, 21

downregulated) and 48 hpi 48/86 (19 upregulated, 29 downregulated) (Supp. File 2). *HDAC5*,

KDM2B, *EZH1*, *PRDM2*, *SETMAR*, *SMYD4* and *USP12* were differentially expressed at all time

points post-infection (excluding 2 hpi).

Discussion

Here we present genome-wide DNA methylation profiles of bAM infected with *M. bovis* versus non-infected controls. We show that CpG methylation in bAM is not altered in response to *M. bovis*, 24 hpi. Since previous studies suggest that DNA methylation changes are established relatively late in the silencing pathway and are preceded by alterations to histone modifications and chromatin packing [41] this may reflect the early post-infection time point examined in this study. Examination of the WGBS data, focusing on the relationship between DNA methylation and proximal promoters, revealed an enrichment of gene promoters that were intermediately methylated.

Global methylation patterns were first analysed using an unbiased genome-wide scan to identify differentially methylated loci between *M. bovis*-infected and control bAM. Following this, we examined the impact of infection on the bAM methylome in greater detail by assessing DNA methylation at specific genomic features. Given the relationship between promoter methylation, gene body methylation and transcription [42] these genomic features comprised the main focus of these analyses. Promoters and CGIs were separated into the following categories as previously described: [36] CGI promoters (*i.e.*, gene promoters overlapping a CpG island), non-CGI promoters, promoter CGIs and non-promoter CGIs. None of these approaches revealed any differentially methylated loci between *M. bovis*-infected and non-infected control bAM. Therefore, it is unlikely that the substantial transcriptomic perturbation observed in bAM during the first 24 h of *M. bovis* infection [22] is due to reconfiguration of CpG methylation patterns. On the other hand, the results presented here indicate that cell signalling and transcription factor-driven gene regulatory transduction cascades lead to the rapid transcriptional activation of immune- and other genes. This observation is supported by previous work showing that DNA

methylation changes in THP-1 macrophages infected with *M. tuberculosis* do not occur at CpGs.

[14] In this study Sharma and colleagues demonstrated that methylation was perturbed at non-

CpGs. Similarly, Lyu *et al.* recently demonstrated that infection of human THP-1 macrophages

with virulent and avirulent *M. tuberculosis* is not associated with host DNA methylation

changes. [43] Unlike THP-1 macrophages infected with *M. tuberculosis*, we show that non-CpG

methylation is not altered in bAM infected with *M. bovis*. The lack of differences in non-CpG

methylation may be explained, in part, by the differences in cell types used – comparing a

macrophage-like human cell line (THP-1) to a primary, differentiated bovine macrophage.

Mycobacteria have recently been reported to modulate the host immune response through

chromatin modifications. [44, 45] Given the absence of differential CpG methylation between non-

infected and *M. bovis*-infected bAM and the differential expression of genes encoding chromatin

modifiers observed in the current study, it is reasonable to hypothesise that chromatin

reconfiguration may have a role in regulating host gene expression in response to infection with

M. bovis.

To comprehensively annotate gene promoter methylation in bAM we quantified average

DNA methylation at proximal promoter regions spanning the TSS (1,500 bp upstream and

500 bp downstream). As expected, the majority of promoters containing or overlapping a CGI

were hypomethylated and those promoters not associated with CGIs were, generally, highly

methylated. [46] However, a large number of promoters (2,580) exhibited mean methylation levels

ranging between 33–66% (intermediately methylated; IM). Interestingly, in addition to this, gene

ontology analysis of the genes proximal to these promoters indicated a marked enrichment for

immune-function related categories. Further analysis of the IM promoter group revealed that

most promoters were included due to averaging of methylated and unmethylated CpGs within

the 2 kb promoter regions. After removing these promoters, 267 promoters remained that exhibited IM. Validation experiments, using clonal analysis and COBRA, confirmed intermediate DNA methylation at the proximal promoter of two of these non-imprinted IM genes, *C1QB* and *IL2RA*. Six of the 267 promoters were proximal to known bovine imprinted genes, displaying predominant intermediate methylation of 5' CGIs; as expected for imprinted genes in an adult somatic cell type. [47] The remaining promoters are IM non-imprinted genes. Previous work by Weber and co-workers demonstrated that, in somatic cells, the concentration of CpGs within a gene promoter is related to the level of DNA methylation; promoters with a high frequency of CpGs (HCP) tend to be unmethylated and promoters with a lower CpG content (LCP) tend to be methylated. [46] Sixty of the 267 IM promoters identified in this study contained high frequencies of CpGs (CGIs) normally associated with unmethylated HCPs, suggesting that promoter IM in bAM is occurring irrespective of CpG density. It has been suggested that intermediate DNA methylation is a conserved signature of genome regulation associated with intermediately active rather than suppressed gene expression. [39] It is possible that these intermediately methylated promoters are a signature of bAM and functionally associated with this particular cell type. However, Elliot and colleagues demonstrated that different tissues and cell types are intermediately methylated equally;[39] therefore, the function of intermediate methylation at these genomic loci remains to be fully elucidated.

Conclusion

This is the first comprehensive analysis of the mammalian alveolar macrophage DNA methylome in response to infection with a mycobacterial pathogen. Although the epigenome of host bAM was not perturbed by a 24 h exposure to the pathogenic bacterium, *M. bovis*, this work provides the first annotation of genome-wide DNA methylation patterns in the bovine genome

497   and is directly aligned with the goal of the Functional Annotation of Animal Genomes (FAANG)

498   project to '*produce comprehensive maps of functional elements in the genomes of domesticated*

499   *animal species*'. [48] Furthermore, this work also provides evidence for differential methylation at

500   the proximal promoter regions of more than 200 non-imprinted genes.

Acknowledgments

We would like to thank Felix Krueger of the Babraham Institute for advice on WGBS data processing and quality control. In addition, we would like to thank Han Haige for assistance with translation of a Chinese journal article.

Disclosure of interest The authors report no conflict of interest.

Figure Legends

Figure 1. Schematic representation of sample preparation. Bovine alveolar macrophages (bAM) were isolated, post-mortem, from the lungs of age-matched male Holstein-Friesian calves by lavage. Purity of the cells was confirmed using flow cytometry with anti-CD14. Isolated cells were washed and seeded for 24 h prior to infection. Infected bAM were exposed to *M. bovis* at a multiplicity of infection ratio of 10:1 for 2 h. After 2 h the media was replaced in control and infected samples and cells were harvested after 24 h for analysis of DNA methylation.

Figure 2. (A) Distribution of paired *t*-statistics between *M. bovis*-infected and control non-infected bAM samples based on smoothed WGBS data. **(B)** Distribution of paired *t*-statistics between randomised samples based on smoothed WGBS data. **(C)** Distribution of average methylation level (%) in promoters and gene bodies across all samples. Only CpG loci with coverage greater or equal to 10 were considered. Only genes where gene body and promoter both had 10 or more sufficiently covered CpG loci were considered.

Figure 3. Distribution of DNA methylation in different genomic contexts in non-infected and *M. bovis*-infected bovine alveolar macrophages. Analysis of WGBS data from *M. bovis*-infected and non-infected bovine alveolar macrophages (bAM) revealed that genomic

methylation, in the context of CpGs, was not altered at any of the sequence features outlined

(intergenic regions, gene bodies, or promoters with or without CpG islands (CGIs) in the host

following infection. Blue and red violins represent non-infected and *M. bovis*-infected bAM,

respectively.

Figure 4. Pyrosequencing validation of whole-genome bisulfite sequencing (WGBS) results

for a panel of key immune gene promoters. **(A)** WGBS proximal promoter plots.

Representative plots showing the average methylation spanning a 10 kb region at the 5' end of

the *TNF*, *IL12A*, *TLR2* and *NFKB2* genes. **(B)** Pyrosequencing validation of WGBS. Locations

of the pyrosequencing assays are denoted by 'PCR' in Figure 4A. Methylation was not different

(paired *t*-test $P \geq 0.05$) between *M. bovis*-infected and non-infected control bAM. The number of

CpG dinucleotides analysed at each loci were *TNF* (9 CpGs), *IL12A* (11 CpGs), *TLR2* (9 CpGs)

and *NFKB2* (8 CpGs).

Figure 5. Analysis of promoters with highly methylated and unmethylated sequence. 1,034

proximal promoters, with a minimum of 30 CpGs, shown to be intermediately methylated (IM)

in the WGBS analysis were visually inspected to remove false positives. 767 were eliminated

from the IM group due to averaging of highly methylated and unmethylated CpGs in the

proximal promoter region (green dashed box). Four examples are presented here: the *GUCY2D*,

SYN1, *RRP1B* and *FOXRED2* genes.

Figure 6. Gene promoters for combined bisulfite restriction analysis (COBRA) and clonal

analysis. WGBS aligments at the *C1QB* and *IL2RA* gene loci. Each panel represents a 10 kb

region at the 5' end of the gene. Green dashed boxes illustrate the *C1QB* and *IL2RA* proximal

promoter regions (the TSS minus 1.5 kb, plus 500 bp) identified as intermediately methylated

(IM) during WGBS data analysis (average methylation 33−66%). PCR: region analysed using

bisulfite PCR, cloning and Sanger sequencing; CGIs: CpG islands; Gene: transcribed region;

Exons: shows the location of the first exon; Red line: *M. bovis*-infected bAM; Blue line: non-

infected control bAM.

Figure 7. Confirmation of an intermediately methylated promoter region at the *C1QB* gene

locus. (A) Clonal analysis of seven CpG dinucleotides in a 269 bp fragment of the bovine *C1QB*

5' promoter region, *a–d* represent sequencing of four biological replicates. Closed and open

circles denote methylated and unmethylated CpGs, respectively.

Figure 8. **Confirmation of an intermediately methylated promoter region at the *IL2RA***

gene locus. (A) Clonal analysis of 10 CpG dinucleotides in a 378 bp fragment of the bovine

IL2RA 5' promoter region, *a–d* represent sequencing of four biological replicates. Closed and

open circles denote methylated and unmethylated CpGs, respectively.

Figure 9. Non-CpG methylation levels differ but not significantly in bovine alveolar

macrophages (bAM) infected with *Mycobacterium bovis*. Mean methylation at the non-CpG

contexts CHGH (left) and CHHG (right) for control and *M. bovis*-infected bAMs, respectively:

differences were not significant by *t*-statistic.

567 **Table 1** Primers use for targeted DNA methylation analysis.

Gene Name	Gene Symbol & Ensembl ID	Forward primer 5'-3'	Reverse Primer 5'-3'	Sequencing Primer 5'-3'
Tumour necrosis factor alpha	**TNF** ENSBTAG00000025471	AGTAATTGGTTTAGAGAAGTTTATTTAGAA	CTTCCTTAATAAAAAAACCCATAAACTCAT **BIOTIN***	GGTTTAGAGAAGTTTATTTAGAAT
Interleukin 12A	**IL12A** ENSBTAG00000015150	TAATTAGAGAGTTAGGTTGGTTATTTATTG **BIOTIN***	ATAAAAATATAACCCCTAATTTAACCTCC	CAACCACCACCCTCA
Toll-like receptor 2	**TLR2** ENSBTAG00000008008	GGGGATGTTAGAGGATTTTAATTTTTGAT **BIOTIN***	ACCCCAACCCCCTCCTCC	CTAAACCACAAAAATTAC
Nuclear Factor Kappa-B, Subunit 2	**NFKB2** ENSBTAG00000006017	TTTGGTGGTGGGAGAGGT **BIOTIN***	CCTCCTCCCACCCTTACC	ACCACCCAAAAATCTAA
Complement C1q B Chain	**C1QB** ENSBTAG00000011196	AGAATTTGAATTAGGGTTTTTGAT T	AAACACTTTCAAATCCCATTTCTA	n/a
Interleukin 2 receptor subunit alpha	**IL2RA** ENSBTAG00000020892	TTAGGGTATTATGGTGAGAGAATTAAG	AAAAAAACAAAAAAATTCCCACTAC	n/a

* Denotes which primer in the pyrosequencing assay that was HPLC purified and 5'Biotinylated

Table 2 Number of tiles/regions in each category.

Region	Control	*M. bovis*
Intergenic	347,561	347,561
Gene Body	166,466	166,466
CGI Promoters	12,047	12,047
Non-CGI Promoters	13,413	13,413
Promoter CGIs	11,222	11,222
Non-Promoter CGIs	30,284	30,284

References

1. Garnier T, Eiglmeier K, Camus JC, Medina N, Mansoor H, Pryor M, Duthoy S, Grondin S, Lacroix C, Monsempe C, et al. The complete genome sequence of Mycobacterium bovis. Proc Natl Acad Sci U S A 2003; 100:7877-82; PMID:12788972; http://dx.doi.org/10.1073/pnas.1130426100;

2. Olea-Popelka F, Muwonge A, Perera A, Dean AS, Mumford E, Erlacher-Vindel E, Forcella S, Silk BJ, Ditiu L, El Idrissi A, et al. Zoonotic tuberculosis in human beings caused by *Mycobacterium bovis*-a call for action. Lancet Infect Dis 2017; 17:e21-e5; PMID:27697390; http://dx.doi.org/10.1016/S1473-3099(16)30139-6;

3. Kathirvel M, Mahadevan S. The role of epigenetics in tuberculosis infection. Epigenomics 2016; 8:537-49; PMID:27035266; http://dx.doi.org/10.2217/epi.16.1;

4. Holoch D, Moazed D. RNA-mediated epigenetic regulation of gene expression. Nat Rev Genet 2015; 16:71-84; PMID:25554358; http://dx.doi.org/10.1038/nrg3863;

5. Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. Nat Genet 2003; 33 Suppl:245-54; PMID:12610534; http://dx.doi.org/10.1038/ng1089;

6. O'Doherty AM, McGettigan PA. Epigenetic processes in the male germline. Reprod Fertil Dev 2015; 27:725-38; PMID:25200708; http://dx.doi.org/10.1071/rd14167;

7. Yong WS, Hsu FM, Chen PY. Profiling genome-wide DNA methylation. Epigenetics Chromatin 2016; 9:26; PMID:27358654; http://dx.doi.org/10.1186/s13072-016-0075-3;

8. Gonzalez RM, Ricardi MM, Iusem ND. Epigenetic marks in an adaptive water stress-responsive gene in tomato roots under normal and drought conditions. Epigenetics 2013; 8:864-72; PMID:23807313; http://dx.doi.org/10.4161/epi.25524;

9. Kucharski R, Maleszka J, Foret S, Maleszka R. Nutritional control of reproductive status in honeybees via DNA methylation. Science 2008; 319:1827-30; PMID:18339900; http://dx.doi.org/10.1126/science.1153069;

10. Navarro-Martin L, Vinas J, Ribas L, Diaz N, Gutierrez A, Di Croce L, Piferrer F. DNA methylation of the gonadal aromatase (cyp19a) promoter is involved in temperature-dependent sex ratio shifts in the European sea bass. PLoS Genet 2011; 7:e1002447; PMID:22242011; http://dx.doi.org/10.1371/journal.pgen.1002447;

11. O'Doherty AM, O'Gorman A, al Naib A, Brennan L, Daly E, Duffy P, Fair T. Negative energy balance affects imprint stability in oocytes recovered from

postpartum dairy cows. Genomics 2014; 104:177-85; PMID:25084396; http://dx.doi.org/10.1016/j.ygeno.2014.07.006;

12. Waterland RA, Kellermayer R, Laritsky E, Rayco-Solon P, Harris RA, Travisano M, Zhang W, Torskaya MS, Zhang J, Shen L, et al. Season of conception in rural gambia affects DNA methylation at putative human metastable epialleles. PLoS Genet 2010; 6:e1001252; PMID:21203497; http://dx.doi.org/10.1371/journal.pgen.1001252;

13. Esterhuyse MM, Linhart HG, Kaufmann SH. Can the battle against tuberculosis gain from epigenetic research? Trends in microbiology 2012; 20:220-6; PMID:22464289; http://dx.doi.org/10.1016/j.tim.2012.03.002;

14. Sharma G, Sowpati DT, Singh P, Khan MZ, Ganji R, Upadhyay S, Banerjee S, Nandicoori VK, Khosla S. Genome-wide non-CpG methylation of the host genome during M. tuberculosis infection. Scientific reports 2016; 6:25006; PMID:27112593; http://dx.doi.org/10.1038/srep25006;

15. Zheng L, Leung ET, Wong HK, Lui G, Lee N, To KF, Choy KW, Chan RC, Ip M. Unraveling methylation changes of host macrophages in Mycobacterium tuberculosis infection. Tuberculosis 2016; 98:139-48; PMID:27156630; http://dx.doi.org/10.1016/j.tube.2016.03.003;

16. Shell SS, Prestwich EG, Baek SH, Shah RR, Sassetti CM, Dedon PC, Fortune SM. DNA methylation impacts gene expression and ensures hypoxic survival of Mycobacterium tuberculosis. PLoS pathogens 2013; 9:e1003419; PMID:23853579; http://dx.doi.org/10.1371/journal.ppat.1003419;

17. Doherty R, Whiston R, Cormican P, Finlay EK, Couldrey C, Brady C, O'Farrelly C, Meade KG. The CD4(+) T cell methylome contributes to a distinct CD4(+) T cell transcriptional signature in Mycobacterium bovis-infected cattle. Scientific reports 2016; 6:31014; PMID:27507428; http://dx.doi.org/10.1038/srep31014;

18. Marr AK, MacIsaac JL, Jiang R, Airo AM, Kobor MS, McMaster WR. Leishmania donovani infection causes distinct epigenetic DNA methylation changes in host macrophages. PLoS pathogens 2014; 10:e1004419; PMID:25299267; http://dx.doi.org/10.1371/journal.ppat.1004419;

19. Sinclair SH, Yegnasubramanian S, Dumler JS. Global DNA methylation changes and differential gene expression in Anaplasma phagocytophilum-infected human neutrophils. Clinical epigenetics 2015; 7:77; PMID:26225157; http://dx.doi.org/10.1186/s13148-015-0105-1;

20. Sitaraman R. Helicobacter pylori DNA methyltransferases and the epigenetic field effect in cancerization. Frontiers in microbiology 2014; 5:115; PMID:24723914; http://dx.doi.org/10.3389/fmicb.2014.00115;

21. Magee DA, Conlon KM, Nalpas NC, Browne JA, Pirson C, Healy C, McLoughlin KE, Chen J, Vordermeier HM, Gormley E, et al. Innate cytokine profiling of bovine alveolar macrophages reveals commonalities and divergence in the response to Mycobacterium bovis and Mycobacterium tuberculosis infection. Tuberculosis 2014; 94:441-50; PMID:24882682; http://dx.doi.org/10.1016/j.tube.2014.04.004;

22. Nalpas NC, Magee DA, Conlon KM, Browne JA, Healy C, McLoughlin KE, Rue-Albrecht K, McGettigan PA, Killick KE, Gormley E, et al. RNA sequencing provides exquisite insight into the manipulation of the alveolar macrophage by tubercle bacilli. Scientific reports 2015; 5:13629; PMID:26346536; http://dx.doi.org/10.1038/srep13629;

23. O'Doherty AM, Magee DA, O'Shea LC, Forde N, Beltman ME, Mamo S, Fair T. DNA methylation dynamics at imprinted genes during bovine pre-implantation embryo development. BMC Dev Biol 2015; 15:13; PMID:25881176; http://dx.doi.org/10.1186/s12861-015-0060-2;

24. O'Doherty AM, Rutledge CE, Sato S, Thakur A, Lees-Murdock DJ, Hata K, Walsh CP. DNA methylation plays an important role in promoter choice and protein production at the mouse *Dnmt3L* locus. Dev Biol 2011; 356:411-20; PMID:21645502; http://dx.doi.org/10.1016/j.ydbio.2011.05.665;

25. O'Doherty AM, O'Shea LC, Fair T. Bovine DNA methylation imprints are established in an oocyte size-specific manner, which are coordinated with the expression of the DNMT3 family proteins. Biol Reprod 2012; 86:67; PMID:22088914; http://dx.doi.org/biolreprod.111.094946 [pii]

10.1095/biolreprod.111.094946;

26. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. Bioinformatics 2011; 27:1571-2; PMID:21493656; http://dx.doi.org/10.1093/bioinformatics/btr167;

27. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods 2012; 9:357-9; PMID:22388286; http://dx.doi.org/10.1038/nmeth.1923;

28. Zimin AV, Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D, Hanrahan F, Pertea G, Van Tassell CP, Sonstegard TS, et al. A whole-genome assembly of the domestic

cow, *Bos taurus*. Genome Biol 2009; 10:R42; PMID:19393038; http://dx.doi.org/gb-2009-10-4-r42 [pii]

10.1186/gb-2009-10-4-r42;

29.   Hansen KD, Langmead B, Irizarry RA. BSmooth: from whole genome bisulfite sequencing reads to differentially methylated regions. Genome Biol 2012; 13:R83; PMID:23034175; http://dx.doi.org/10.1186/gb-2012-13-10-r83;

30.   Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ. The UCSC Table Browser data retrieval tool. Nucleic Acids Res 2004; 32:D493-6; PMID:14681465; http://dx.doi.org/10.1093/nar/gkh103;

31.   Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, Mason CE. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. Genome Biol 2012; 13:R87; PMID:23034086; http://dx.doi.org/10.1186/gb-2012-13-10-r87;

32.   Benjamini Y, Hochberg Y. Controlling the false discovery rate - a practical and powerful approach to multiple testing. J R Stat Soc Ser B Method 1995; 57:289-300

33.   Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for RNA-seq: accounting for selection bias. Genome Biol 2010; 11:R14; PMID:20132535; http://dx.doi.org/10.1186/gb-2010-11-2-r14;

34.   Ziller MJ, Hansen KD, Meissner A, Aryee MJ. Coverage recommendations for methylation analysis by whole-genome bisulfite sequencing. Nat Methods 2015; 12:230-2, 1 p following 2; PMID:25362363; http://dx.doi.org/10.1038/nmeth.3152;

35.   Clark SJ, Smallwood SA, Lee HJ, Krueger F, Reik W, Kelsey G. Genome-wide base-resolution mapping of DNA methylation in single cells using single-cell bisulfite sequencing (scBS-seq). Nature protocols 2017; 12:534-47; PMID:28182018; http://dx.doi.org/10.1038/nprot.2016.187;

36.   Peat JR, Dean W, Clark SJ, Krueger F, Smallwood SA, Ficz G, Kim JK, Marioni JC, Hore TA, Reik W. Genome-wide bisulfite sequencing in zygotes identifies demethylation targets and maps the contribution of TET3 oxidation. Cell reports 2014; 9:1990-2000; PMID:25497087; http://dx.doi.org/10.1016/j.celrep.2014.11.034;

37.   Chen ZX, Riggs AD. DNA methylation and demethylation in mammals. J Biol Chem 2011; 286:18347-53; PMID:21454628; http://dx.doi.org/10.1074/jbc.R110.205286;

38. Deaton AM, Webb S, Kerr AR, Illingworth RS, Guy J, Andrews R, Bird A. Cell type-specific DNA methylation at intragenic CpG islands in the immune system. Genome Res 2011; 21:1074-86; PMID:21628449; http://dx.doi.org/10.1101/gr.118703.110;

39. Elliott G, Hong C, Xing X, Zhou X, Li D, Coarfa C, Bell RJ, Maire CL, Ligon KL, Sigaroudinia M, et al. Intermediate DNA methylation is a conserved signature of genome regulation. Nature communications 2015; 6:6363; PMID:25691127; http://dx.doi.org/10.1038/ncomms7363;

40. Nestorov P, Hotz HR, Liu Z, Peters AH. Dynamic expression of chromatin modifiers during developmental transitions in mouse preimplantation embryos. Scientific reports 2015; 5:14347; PMID:26403153; http://dx.doi.org/10.1038/srep14347;

41. Smith ZD, Meissner A. DNA methylation: roles in mammalian development. Nat Rev Genet 2013; 14:204-20; PMID:23400093; http://dx.doi.org/10.1038/nrg3354;

42. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. Nat Rev Genet 2012; 13:484-92; PMID:22641018; http://dx.doi.org/10.1038/nrg3230;

43. Lyu LN, Jia HY, Li ZH, Liu ZQ, Zhang ZD. [Changes and differences of DNA methylation in human macrophages infected with virulent and avirulent Mycobacterium tuberculosis]. Zhonghua jie he he hu xi za zhi = Zhonghua jiehe he huxi zazhi = Chinese journal of tuberculosis and respiratory diseases 2017; 40:509-14; PMID:28728275; http://dx.doi.org/10.3760/cma.j.issn.1001-0939.2017.07.006;

44. Sharma G, Upadhyay S, Srilalitha M, Nandicoori VK, Khosla S. The interaction of mycobacterial protein Rv2966c with host chromatin is mediated through non-CpG methylation and histone H3/H4 binding. Nucleic Acids Res 2015; 43:3922-37; PMID:25824946; http://dx.doi.org/10.1093/nar/gkv261;

45. Yaseen I, Kaur P, Nandicoori VK, Khosla S. Mycobacteria modulate host epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3. Nature communications 2015; 6:8922; PMID:26568365; http://dx.doi.org/10.1038/ncomms9922;

46. Weber M, Hellmann I, Stadler MB, Ramos L, Paabo S, Rebhan M, Schubeler D. Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome. Nat Genet 2007; 39:457-66; PMID:17334365; http://dx.doi.org/10.1038/ng1990;

47. Pervjakova N, Kasela S, Morris AP, Kals M, Metspalu A, Lindgren CM, Salumets A, Magi R. Imprinted genes and imprinting control regions show predominant

intermediate methylation in adult somatic tissues. Epigenomics 2016; 8:789-99; PMID:27004446; http://dx.doi.org/10.2217/epi.16.8;

48.   Andersson L, Archibald AL, Bottema CD, Brauning R, Burgess SC, Burt DW, Casas E, Cheng HH, Clarke L, Couldrey C, et al. Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes project. Genome Biol 2015; 16:57; PMID:25854118; http://dx.doi.org/10.1186/s13059-015-0622-4;

A

C

B

Supp table 1

Click here to access/download
Supplementary Material - for review
Supp Table 1_22-12-17.xlsx

Supp table 2

Click here to access/download
Supplementary Material - for review
Supp Table 2 22-12-17.xlsx

Supp table 3

Click here to access/download
Supplementary Material - for review
Supp Table 3 22-12-17.xlsx

Click here to access/download
Supplementary Material - for review
Supp Table 4 22-12-17.xlsx

Click here to access/download
Supplementary Material - for review
Supp Table 5 22-12-17.xlsx

Click here to access/download
Supplementary Material - for review
Supp Table 6 22-12-17.xlsx

Click here to access/download
Supplementary Material - for review
Supp Figure 1 22-12-17.pdf

Supp file 1a

Click here to access/download
Supplementary Material - for review
C11_CAGATC_R1_merged_val_1.fq.gz_bismark_bt2_P
E_report.html

Supp file 1b

Click here to access/download
Supplementary Material - for review
C12_TAGCTT_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Supp file 1c

Supp file 1d

Click here to access/download
Supplementary Material - for review
C4_TGACCA_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html

Supp file 1e

Click here to access/download
Supplementary Material - for review
C5_CAGATC_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html

Supp file 1f

Click here to access/download
Supplementary Material - for review
C6_TAGCTT_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html

Supp file 1g

Click here to access/download
Supplementary Material - for review
C7_ATCACG_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html

Click here to access/download
**Supplementary Material - for review
C8_TGACCA_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html**

Supp file 1i

Click here to access/download
Supplementary Material - for review
M11_ACTTGA_R1_merged_val_1.fq.gz_bismark_bt2_P
E_report.html

Supp file 1j

Click here to access/download
Supplementary Material - for review
M12_GGCTAC_R1_merged_val_1.fq.gz_bismark_bt2_P
E_report.html

Supp file 1k

Click here to access/download
Supplementary Material - for review
M1_CGATGT_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Click here to access/download
Supplementary Material - for review
M4_ACAGTG_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Supp file 1m

Click here to access/download
Supplementary Material - for review
M5_ACTTGA_R1_merged_val_1.fq.gz_bismark_bt2_PE_
report.html

Supp file 1n

Click here to access/download

Supplementary Material - for review

M6_GGCTAC_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Supp file 1o

Click here to access/download
Supplementary Material - for review
M7_CGATGT_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Click here to access/download
Supplementary Material - for review
M8_ACAGTG_R1_merged_val_1.fq.gz_bismark_bt2_PE
_report.html

Supp file 2

Click here to access/download
Supplementary Material - for review
Supp File 2 22-12-17.xlsx

Click here to access/download
Supplementary Material - for review
Supp File 3 22-12-17.pdf