**A divergent CheW confers plasticity to nucleoid-associated chemosensory arrays**

Annick Guiseppi[1*], Juan Jesus Vicente[2*], Julien Herrou[1], Deborah Byrne[3], Aurelie Barneoud[1],

Audrey Moine[1], Leon Espinosa[1], Marie-Jeanne Basse[4], Virginie Molle[5], Tâm Mignot[1], Philippe

Roche[4] and Emilia M.F. Mauriello[1#]

*[1] Aix Marseille Univ, CNRS, LCB, Marseille, France; [2]Physiology & Biophysics, University of*

*Washington, Seattle, WA, USA; [3]Protein Purification Platform, Institut de Microbiologie de la*

*Méditerranée, CNRS, Marseille, France; [4]CNRS, INSERM, Aix Marseille Univ, Institute Paoli-*

*Calmettes, CRCM, Marseille, France; [5]Montpellier II et I University, CNRS, Laboratoire de*

*Dynamique des Interactions Membranaires Normales et Pathologiques, France.*

*These authors contributed equally

[#]Correspondence to:

Emilia M.F. Mauriello

Aix Marseille Univ, CNRS, LCB

31 Chemin Joseph Aiguier, Marseille, 13402, France

Tel. +33491164321

Email : emauriello@imm.cnrs.fr

**Running Title: The formation of multiple Frz clusters requires FrzB**

**ABSTRACT**

Chemosensory systems are highly organized signaling pathways that allow bacteria to adapt to environmental changes. The Frz chemosensory system from *M. xanthus* possesses two CheW-like proteins, FrzA (the core CheW) and FrzB. We found that FrzB does not interact with FrzE (the cognate CheA) as it lacks the amino acid region responsible for this interaction. FrzB, instead, acts upstream of FrzCD in the regulation of *M. xanthus* chemotaxis behaviors and activates the Frz pathway by allowing the formation and distribution of multiple chemosensory clusters on the nucleoid. These results, together, show that the lack of the CheA-interacting region in FrzB confers new functions to this small protein.

**AUTHOR SUMMARY**

Chemosensory systems are signaling complexes that are widespread in bacteria and allow

the modulation of different cellular functions, such as taxis and development, in response to

the environment. We show that the *Myxococcus xanthus* FrzB is a divergent CheW lacking

the region involved in the interaction with the histidine kinase FrzE. Instead, it acts upstream

of FrzCD to allow the formation of multiple distributed Frz chemosensory arrays at the

nucleoid. The loss of the CheA-interacting region in FrzB might have been selected to confer

plasticity to nucleoid-associated chemosensory systems. By unraveling a new accessory

protein and its function, this work opens new insights into the knowledge of the regulatory

potentials of bacterial chemosensory systems.

**INTRODUCTION**

Chemosensory systems are specialized regulatory pathways that allow bacteria to perceive

their external environment and respond with various cellular behaviors (1–3). In these

systems, environmental signals are transduced inside the cells, initially by receptors called

Methyl-accepting Chemotaxis Proteins (MCP). Most MCPs possess a transmembrane

domain, but 14% are soluble proteins (4). Ligand-bound MCPs regulate the

autophosphorylation of the histidine kinase CheA that, in turn, transfers phosphoryl groups

to at least two response regulators: CheY which is the most downstream component of the

pathway and CheB, which, together with CheR, constitutes the adaptation module. Che

systems also include one or more versions of the docking protein CheW, which interacts

with the C-terminal cytoplasmic tip of the MCP and with the P5 domain of CheA. CheW and

CheA[P5] are paralogs and are topologically similar to SH3 domains from eukaryotic scaffold

proteins that also play a role in signal transduction. Such domains are characterized by two

β-sheets (each composed of five β-strands) designated subdomain 1 and 2 (5–7). MCP:

CheA[P5]: CheW interactions all involve subdomains 1 and 2. In particular, the β-strands 3 and

4 of subdomain 1 of CheA[P5] interact with the β-strands 4 and 5 of subdomain 2 of CheW (8).

The receptor cytoplasmic tip binds CheW or CheA at the hydrophobic junction between

subdomain 1 and 2 (8).

High-resolution microscopy revealed that, intracellularly, Che proteins are organized in

complex arrays running parallel to the bacterial inner membrane, with the MCP layer linked

to a CheA/W baseplate (9,10). Transverse views of these Che arrays show ordered MCP

hexagons networked by CheA-CheW rings. CheA-CheW rings contain alternating and

interacting CheA[P5] domains and CheWs. MCPs hexagons are constituted by six MCP trimers

of dimers. One MCP in the dimer is connected with either a CheA[P5] domain or a CheW,

which allows connections between the MCP hexagons and the CheW-CheA[P5] rings (11) (Fig.

1A). The presence of each of these three proteins is essential for the integrity of the system

(12,13). This structure provides the typical signaling properties of chemotaxis systems,

including signal amplification and sensitivity (14–17).

The gliding bacterium *M. xanthus* uses the cytoplasmic Frz signal transduction pathway to

bias the frequency at which cells change their direction of movement in response to

environmental signals (18). This behavior allows cells to glide towards favorable directions or

away from toxic compounds. The Frz system induces cell reversal by promoting the

coordinated inversion of the cell polarity of the two *M. xanthus* motility systems: the Social

(S) motility system powered by polar Type IV Pili and the Adventurous (A) motility powered

by polarly assembled Agl-Glt complexes (19–21). The regulation of the motility systems by

the Frz pathway is essential for cells to enter a developmental program that leads to the

formation of multicellular fruiting bodies. The Frz core is composed of a cytoplasmic MCP

(FrzCD), a CheA (FrzE) and a CheW (FrzA) (22) (Fig. 1B), encoded by a single operon (Fig. 1C).

In the absence of any of these three proteins, cells display drastically reduced reversal

frequencies and are no longer able to respond to isoamyl alcohol (IAA), a Frz activator

(20,22). The Frz system also includes a second CheW-like protein, FrzB, described as an

accessory because while in its absence cells show phenotypes similar to those caused by the

deletion of core proteins, *ΔfrzB* cells are still able to respond to IAA with increased reversal

frequencies (20,22) (Fig. 1B).

In this paper, we show that FrzB is a divergent CheW lacking the β-strands involved in the

interaction with the histidine kinase FrzE (β4 and β5). Strikingly, the introduction of β4 and

β5 from FrzA to FrzB converts the latter into a canonical CheW. Instead of mediating signal

transduction from the FrzCD receptor to FrzE, FrzB acts upstream of FrzCD to allow the

formation of multiple distributed Frz chemosensory clusters at the nucleoid. These clusters

collapse in a unique and less active array in the absence of FrzB. Finally, the loss of the CheA-

interacting region in FrzB might have been selected to confer plasticity to nucleoid-

associated chemosensory systems.

# RESULTS

## FrzB is upstream of the FrzCD receptor in the Frz signaling pathway

The first two genes of the *frz* operon encode the CheW-like proteins, FrzA and FrzB. *ΔfrzA* cells show the same motility and fruiting body formation phenotypes as *ΔfrzCD* and *ΔfrzE*, suggesting that FrzA is the core CheW of the Frz pathway (22,23) (Fig. 2A). The aberrant behaviors at the colony level are due to the inability of *ΔfrzA* strains to modulate single-cell reversal frequencies even in the presence of isoamyl alcohol (IAA), a known Frz activator (Fig. 2B) (20,22).

While a *frzB* deletion causes the same defects as *ΔfrzA* in the absence of IAA, IAA completely rescues the reversal frequency defect of *ΔfrzB* cells (Fig. 2B). This effect is detectable at both single-cell and colony level (Fig. 2) (22). These results show that signals from the FrzCD receptor can still be transduced to FrzE in the absence of FrzB and, thus, FrzB is not part of the Frz core. Moreover, a double *ΔfrzA ΔfrzB* phenocopies *ΔfrzA*, further suggesting the central role of FrzA and the accessory role of FrzB (Figure S1).

In order to determine the functional position of FrzB in the Frz pathway, we tested the epistasis of FrzB with respect to FrzCD. Because FrzCD is a central component of the Frz pathway, *ΔfrzB ΔfrzCD* and *ΔfrzA ΔfrzCD* double mutants showed the same aberrant motility and developmental phenotypes as *ΔfrzCD* (and *ΔfrzA*) at both single-cells and colony levels (Figure S1 and Fig. 2) (22). We thus took advantage of a *frzCD* allele, *frzCDc*, whose expression confers a constitutively hyper-reversing phenotype (Fig. 2) (Bustamante et al., 2004) and constructed *ΔfrzA frzCDc* and *ΔfrzB frzCDc* double mutants. The *ΔfrzA frzCDc* reversal frequencies and colony phenotypes were indistinguishable from those of *ΔfrzA* cells, confirming that FrzA is located downstream of the receptor in the signaling pathway (Fig. 2). On the other hand, *ΔfrzB frzCDc* phenotypes resembled those of *frzCDc* (Fig. 2), indicating

that FrzB is upstream of FrzCD in the Frz signaling pathway and that its requirement is

bypassed by a constitutive *frzCDc* mutation.

**FrzB does not interact with the downstream histidine kinase FrzE**

Genetic data show that FrzB is upstream of FrzCD in the Frz regulatory pathway. We thus

wanted to verify if FrzB could interact with FrzE. For this, we heterologously expressed and

co-purified, using GST-affinity chromatography, GST-FrzA or GST-FrzB with either 6His-FrzCD

or 6His-FrzE$^{CheA}$. Both 6His-FrzCD and 6His-FrzE$^{CheA}$ could be co-purified with GST-FrzA,

suggesting that FrzA interacts with both FrzCD and FrzE$^{CheA}$ (Fig. 3A). On the other hand, only

6His-FrzCD, but not 6His-FrzE$^{CheA}$, could be co-purified with GST-FrzB, suggesting that FrzB

only interacted with FrzCD but not with FrzE$^{CheA}$ (Fig. 3A). We then wanted to test whether,

by using high amounts of GST-FrzB, we could detect an interaction with immobilized 6His-

FrzE$^{CheA}$ in  Biolayer interferometry (BLi) (24). While low amounts of GST-FrzB still produced

an interaction with immobilized 6His-FrzCD (Kd = 0.7 $\mu$M), no interactions could be detected

with immobilized 6His-FrzE1-658, even when we used high amounts of GST-FrzB (Fig. 3B).

If FrzB does not interact with FrzE, it should not be able to promote the phosphorylation of

the FrzE kinase domain (FrzE$^{CheA}$) *in vitro*. In fact, it has been previously shown that the

presence of both FrzCD and FrzA is strictly required to achieve the transfer of a phosphoryl

group from ATP to the Histidine at position 49 of FrzE$^{CheA}$ *in vitro* (25) (Fig. 3C). As expected,

when we replaced FrzA with FrzB in the mixture, no phosphorylation was detected,

suggesting that FrzB cannot mediate the FrzE$^{CheA}$ autophosphorylation *in vitro* (Fig. 3C).

Next, we asked what prevented FrzB from interacting with FrzE. To address this question, we

decided to generate 3D homology models of the FrzE$^{P4-P5}$:FrzA and FrzE$^{P4-P5}$:FrzB complexes

based on the *T. maritima* CheA$^{P4-P5}$:CheW crystal structure (Fig. 3D) (8). The structural

models predict that FrzB cannot interact with FrzE due to the lack of a 20 amino acid region

forming the β-strands 4 and 5 of subdomain 2. These strands correspond exactly to the

CheW region important for the CheW: CheA$^{P5}$ interaction (Fig. 3D and E) (8). On the other

hand, and consistent with the two-hybrid system results, FrzB is predicted to contain all the

hydrophobic amino acids involved in its interaction with FrzCD (Fig. 3E).

To verify if the lack of the FrzE-interacting region in FrzB was directly associated with the loss

of the FrzB-FrzE interaction, we designed a chimeric FrzB protein carrying the β-strands 4

and 5 from FrzA (FrzB$^{β4-β5}$). First, we tested if purified FrzB$^{β4-β5}$ was able to replace FrzA in our

FrzE$^{CheA}$ phosphorylation *in vitro* assay. Strikingly, when FrzB$^{β4-β5}$ was the only CheW in the

reaction mixture, it promoted the FrzE$^{CheA}$ phosphorylation similarly to FrzA, thus acting as a

CheW (Fig. 3C). The FrzE$^{CheA}$ phosphorylation levels were a direct consequence of the FrzCD

activation state, because when we used the hyperactive FrzCD$^c$ allele in the place of its wild-

type version, we observed a higher level of phosphorylated FrzE$^{CheA}$ (Fig. 3C).

Then, to test if FrzB$^{β4-β5}$ could mediate FrzE autophosphorylation *in vivo* as *in vitro*, we

transferred the gene coding for FrzB$^{β4-β5}$ into a *ΔfrzA ΔfrzB* strain (Figure S1) at the

endogenous locus. FrzB$^{β4-β5}$ was stably expressed (Figure S2), and cells could reverse in the

presence of IAA similarly to *ΔfrzB* (Fig. 2). These results suggest that *in vivo,* like *in vitro*,

FrzB$^{β4-β5}$ can connect FrzCD to FrzE.

**FrzB might be a stabilizing factor of multiple Frz signaling arrays**

It has been previously shown that Frz proteins form nucleoid-associated signaling complexes

(26,27). To verify whether FrzB was part of the Frz clusters, we constructed a *mCherry-frzB*

fusion that replaced *frzB* at the endogenous locus and encoded a functional protein (Figure

S2 and S3). mCherry-FrzB formed multiple nucleoid-associated clusters that resembled those

formed by FrzCD-GFP and FrzE-mCherry (Fig. 4A) (27).

Then, we decided to check how the absence of FrzA and FrzB affected the formation of these

complexes. For that, we transferred a *frzCD-gfp* gene fusion (27) into *ΔfrzA* and *ΔfrzB*

mutants. *M. xanthus* cells of the resulting strains were then imaged by fluorescence

microscopy. While FrzCD-GFP was completely dispersed in the cytoplasm in the absence of

FrzA, this fusion protein formed a single nucleoid-associated cluster in the absence of FrzB

(Fig. 4B). These results suggest that, even if FrzB is dispensable for the assembly of Frz

chemosensory arrays, it is important for the formation of multiple distributed clusters on the

nucleoid. These clusters collapse into a single cluster in the absence of FrzB. Of 230 cells

examined, 82% have a single cluster, whereas 18% have no cluster. It is possible that, in the

absence of FrzB and during cell division, only one daughter cell inherits the single Frz array

and the other one will synthesize a new one. This would explain the lack of clusters in a small

population of cells. In the wild type, the multiple Frz clusters would be expected to be

distributed to both daughter cells (27). In the absence of both FrzA and FrzB, FrzCD-GFP was

diffused in the cytoplasm, as in *ΔfrzA* (Fig. 4B). The aberrant localization patterns were not

due to a change in the FrzCD-GFP protein levels nor in a change of the nucleoid structure

(Figure S5 and Figure S6). Interestingly, and by yet unknown mechanisms, FrzA not only is

essential for Frz cluster formation, but it also allows the association of FrzCD to the nucleoid

(Fig. 4).

**DISCUSSION**

Chemosensory systems are modified two-component systems in which the addition of accessory proteins confers unique properties like signal amplification and sensitivity. In this work, we found that the CheW-like protein FrzB lacks a 20 amino acid sequence corresponding to two β-strands that are important for CheA$^{P5}$:CheW interactions (8). Thus, FrzB cannot interact with FrzE and mediate its autophosphorylation. The loss of this protein region might have been selected to confer to this small protein the unique function of allowing the formation and distribution of multiple nucleoid Frz chemosensory arrays. How would FrzB accomplish these tasks? It has been shown that Che arrays are constituted by hexagons of MCP trimers of dimers networked by CheA$^{P5}$-CheW rings (11). By homology, we believe that Frz proteins can have such hexagonal organization using the nucleoid as scaffold. However, if FrzB cannot interact with FrzE$^{P5}$, it cannot participate to the formation of CheA$^{P5}$-CheW rings either. In addition to CheA$^{P5}$-CheW rings, the formation of Che networks implies the formation of six-member CheW rings that only interact with the MCPs (Fig. 1) (17,28,29). The 6-CheW rings have been visualized by cryoEM, and their suggested function is to stabilize the Che arrays by favoring allosteric interactions, like CheW-CheA rings, but without a direct effect on signaling. We can imagine that 6-FrzB rings (in addition to FrzE$^{P5}$-FrzA rings and 6-FrzA rings) might further stabilize small Frz clusters, which would otherwise collapse into one single cluster (Fig. 5). Alternatively FrzB allows the formation of small Frz clusters by breaking the networking of MCP trimers of dimers with Che$^{P5}$-CheW rings (Fig. 5).

Interestingly, the FrzCD receptor does not contain an obvious ligand-binding domain. Instead, it contains an N-terminal region that has been recently shown to be involved in DNA binding (27). Thus, FrzB could also function to deliver signals to the Frz pathway. FrzB might

bind directly to some cytoplasmic molecules and transduce signals to FrzCD, thus functioning

as a sensor of the cellular metabolic state. However, FrzB could also connect FrzCD to

another MCP or sensor kinase. *M. xanthus* encodes a total of 21 MCPs (30). Among them,

only Mcp7 is cytoplasmic like FrzCD, but its localization in a single cluster randomly localized

within cells, strongly excludes an association with the Frz pathway (30). We could imagine

that FrzCD, which forms clusters on the nucleoid (27), could associate with a transmembrane

MCP or sensor kinase at regions where the nucleoid surface is particularly close to the inner

membrane (31). In fact, previous work shows the association of FrzCD with membrane

fractions (32) as well as interactions with envelope-associated A-motility proteins (33,34).

The hypothesis that FrzB is a Frz input does not rule out that FrzCD itself can somehow bind

signals. Indeed, FrzCD responds to IAA independently of FrzB (Fig. 2).

In recent years, the discovery of different accessory Che proteins allowed the identification

of new functions and degrees of complexity in the regulation of the cellular responses

operated by chemosensory systems. By characterizing a new accessory protein and its

function, this work opens new insights on the knowledge of the regulatory potentials of

bacterial chemosensory systems.

**MATERIALS AND METHODS**

**Bacterial strains, plasmids and growth**

*S*trains and plasmids are listed in Table S1 and S2, respectively. *M. xanthus* strains were

grown at 32°C in CYE rich media as previously described.

Plasmids were introduced into *M. xanthus* cells by electroporation. Deletions and GFP

fusions were inserted in frame, using the pBJ113 or pBJ114 vectors, to avoid polar effects on

the downstream gene expression (22,30).

Strains EM736 and EM743 were obtained by deleting *frzA* from codon 15 to 145 and *frzB*

from codon 11 to 92, respectively. In both strains we also deleted *frzCD* from codon 6 to

182. Strain EM742 was obtained by deleting *frzA* and *frzB* as above.

To construct plasmid pEM399 to generate EM515, a DNA fragment was generated by PCR

overlapping a DNA region amplified with primers CCCAAGCTTCTTCAGCTCGAGGCTGGCAA

and TGCCCCTTGCTCACCATCTCCTCCTCCTTCTCATGGAA and *mCherry* amplified with primers

CATGAGAAGGAGGAGGAGATGGTGAGCAAGGGCGAGGA                                      and

CGGGATCCTCTAGACTTGCACAGCTCGTCCATGC from pEM147 (33). This fusion product was

digested     with     HindIII     and     BamHI.     *frzB*     was     amplified     with     primers

CGGGATCCGGAGCAGGAGACCTCCTCTTCTTCGACAT                                        and

GGAATTCGCGGACGAGGACAGCCTCAA   and   digested   with   BamHI   and   EcoRI.   The   two

fragments were then cloned into pBJ113 previously digested with HindIII and BamHI in a

three-partner ligation reaction.

To obtain strains EM440 and EM512, plasmid pDPA20 was transferred into strains DZ4479

and DZ4478 respectively.

For the construction of *frzB^{84-85}* we replaced codon 27 to 36 of FrzB with codon 41 to 75 of

FrzA. Plasmids pEM463 and was synthesized by Biomatik.

*Escherichia coli* cells were grown under standard laboratory conditions in LB broth

supplemented with antibiotics if necessary.

**Phenotypic analysis**

Developmental assays were performed by spotting 10 µl of cells at 5 $OD_{600}$ directly onto CF

1.5 % agar plates. Vegetative swarming phenotypes were analyzed by spotting the same

amount of cells onto CYE plates containing a concentration of 0.5 % agar. The IAA response

of cell colonies was analyzed by spotting cells (10 µl of cells at 5 $OD_{600}$) on CYE plates

supplied with 0.15 % of IAA and containing a concentration of 1.5% agar. After incubation at

32°C the swarming colonies and the fruiting bodies were photographed at an Olympus SZ61

stereoscope with a Nikon DXM1200 Digital Camera.

**Reversal frequencies**

5 µl of cells from 4 x $10^8$ cfu $ml^{-1}$ vegetative CYE cultures were spotted on a thin fresh TPM

agar supplied or not with 0.15 % IAA. Time-lapse movies were shot for 1 hour with frames

captured every 30 seconds. Movies were obtained and reversal frequencies calculated with

Image J (Rasband, W.S., ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA,

http://imagej.nih.gov/ij/, 1997-2012) and FIJI as previously described (20). For non-reversing

strains, the number of reversals for each cell was plotted against time using the R software

(https://www.r-project.org/). For strains that frequently reversed, the number of periods

between two reversals were plotted against time using the R software. Reversal frequencies

were measured from cells issued from two to six biological replicates. Statistical significance

was obtained with a Wilcox Test from R.

**Protein purification and co-purification**

BL21(DE3) [F– ompT hsdSB(rB– mB–) gal dcm (DE3)] cells were grown in LB broth

supplemented 100 μg/ml ampicillin to mid-exponential phase at 37°C. Overexpression was

induced by adding 0,2 mM IPTG. Cells were then grown at 16°C over night. Cells were

washed and resuspended in lysis buffer (50 mM TrisHCl, pH 8; 300 mM NaCl; 100 μg/ml

PMSF; 30 U/mL Benzonase) and lysed at the French press. The cell lysates were centrifuged

at 4°C for 20 min at 18000× rpm.

For BLi experiments, soluble 6His-tagged proteins were purified using a NiNTA resine (GE

Healthcare) and imidazole was removed by dialysis into a buffer containing 50 mM TrisHCl,

pH 8 and 300 mM NaCl. GST-proteins were purified by GST-affinity chromatography using

the same lysing buffer supplied of 10 mM reduced glutathione to elute.

For co-purification experiments, *E. coli* cells expressing each of GST-FrzA, GST-FrzB, 6His-

FrzCD or 6His-FrzECheA were mixed at one to one ratio to obtain the following

combinations: GST-FrzA: 6His-FrzCD, GST-FrzA: 6His- FrzECheA, GST-FrzB: 6His-FrzCD, GST-

FrzB: 6His- FrzECheA. Mixed cell cultures were resuspended in lysis buffer (50 mM TrisHCl,

pH 8; 300 mM NaCl; 100 μg/ml PMSF; 30 U/mL Benzonase) and lysed at the French press.

The cell lysates were centrifuged at 4°C for 20 min at 18000× rpm.

Proteins were, then, co-purified by GST-affinity chromatography as described above.

**Biolayer Interpherometry**

Protein-protein interaction experiments were conducted at 25°C with the BLItz instrument

from ForteBio (Menlo Park, CA, USA). The BLI consists in a real time optical biosensing

technique exploits the interference pattern of white light reflected from two surfaces to

measure biomolecular interactions (24). Purified 6His-FrzCD or 6His-FrzE$^{CheA}$ protein ligands

were immobilized onto two different Ni-NTA biosensors (ForteBio) in duplicate at 4 µM

concentrations. GST-FrzB was used as the analyte throughout the study at concentrations

ranging from 0.2 to 20 µM. The assay was conducted in buffer containing Tris HCl 50 mM

pH8 and NaCl 300 mM. The binding reactions were performed with an initial baseline during

30 seconds, an association step at 120 seconds and a dissociation step of 120 seconds with

lateral shaking at 2200rpm.  A double reference subtraction (sensor reference and GST-FrzB

at 10 µM) was applied to account for non-specific binding, background, and signal drift to

minimize sensor variability.

**Sequence alignment and homology modeling**

Predictions of secondary structures and protein sequence alignments were obtained with

Jpred (35) and Clustal Omega (36), respectively. Homology models of FrzA (Uniprot P43498)

or FrzB (Uniprot P43499) in the presence of FrzE (Uniprot P18769) were built with

MODELLER 9v12 (37). Crystal structure of the complex between bacterial chemotaxis

histidine kinase CheA$^{P4-P5}$ and receptor-adaptor protein CheW (PDB codes 2CH4) was used as

template (38). The modeling was performed in a two-step manner: First, each individual

domain was modeled separately using the 3D structure of its *T. maritima* counterpart as a

template. CheW (2CH4 chain W) was used to build models of FrzA and FrzB. The N-terminal

domain of CheA (2CH4 chain A resid 355-540) and the C-terminal domain of CheA (2CH4

chain A resid 541-671) were used to build N- and C-terminal domains of FrzE. In a second

step, each model was superimposed to its counterpart in the x-ray structure of the CheW-

CheA complex (chains W and A). The Frz proteins exhibit high enough identities to their Che

counterparts to build reasonable 3D models. FrzB vs CheW (26.1% identity, 45% similarity),

FrzA vs CheW (26.6% identity, 46.7% similarity), FrzE$^{p4}$ vs. CheA$^{p4}$ (46.5% identity, 71.7%

similarity), FrzE$^{p5}$ vs. CheA$^{p5}$ (26.1% identity, 53.8% similarity). Quality of the models was

assessed with Prosess server (http://www.prosess.ca). Structural alignments were

performed with Protein structure comparison service PDBeFold at European Bioinformatics

Institute (http://www.ebi.ac.uk/msd-srv/ssm) (39). Fig. 3C showing the lowest energy model

was generated with pymol (http://pymol.org). Fig. 3D was generated with ESPript web

server (40).

**Fluorescence microscopy and image analysis**

For fluorescence microscopy analyses, 5 µl of cells from 4 x 10$^8$ cfu ml$^{-1}$ vegetative CYE

cultures were spotted on a thin fresh TPM agar pad at the top a slide (41). A cover slip was

added immediately on the top of the pad, and the obtained slide was analyzed by

microscopy using a Nikon Eclipse TE2000 E PFS inverted epifluorescence microscope (100 x

oil objective NA 1.3 Phase Contrast) (42).

To study the co-localization with the DNA, the TPM agar pads were supplied with 1µg ml$^{-1}$

DAPI stain as shown by Moine et al. (27). FrzB clusters, nucleoid areas and cell areas were

automatically detected and verified manually with the "MicrobeJ" Fiji/ImageJ plugin created

by A. Ducret, Brun Lab (http://www.indiana.edu/~microbej/index.html). All data plots and

statistical tests were obtained with the R software (https://www.r-project.org/).

***In vitro* autophosphorylation assay**

*In vitro* phosphorylation assays were performed with *E. coli* purified recombinant proteins. 4

µg of FrzE$^{CheA}$ was incubated with 1 µg of FrzA, FrzB or FrzB$^{β4-β5}$ and FrzCD or FrzCD$^c$ in 25 µl

of buffer P (50 mM Tris-HCl, pH 7.5; 1 mM DTT; 5 mM MgCl$_2$; 50 mM KCl; 5 mM EDTA; 50 µM

ATP, 10% (v/v) glycerol) supplemented with 200 µCi ml$^{-1}$ (65 nM) of [γ-33P]ATP (PerkinElmer,

3000 Ci mmol$^{-1}$) for 10 minutes at room temperature in order to obtain the optimal

FrzE$^{CheA}$ autophosphorylation activity. Each reaction mixture was stopped by addition of 5 ×

Laemmli and quickly loaded onto SDS-PAGE gel. After electrophoresis, proteins were

revealed using Coomassie Brilliant Blue before gel drying. Radioactive proteins were

visualized by autoradiography using direct exposure to film (Carestream).

**Acknowledgements**

We thank Romain Mercier, Dorothée Murat and Laetitia My for their support and helpful discussions. We also thank Anne-Valerie Le Gall for her technical support.  We thank the anonymous reviewers, whose comments helped to greatly improve the manuscript.

References

1.     Sourjik V, Berg HC. Binding of the Escherichia coli response regulator CheY to its
target measured in vivo by fluorescence resonance energy transfer. Proc Natl Acad Sci U S A.
2002 Oct 1;99(20):12669–74.
2.     Berleman JE, Bauer CE. Involvement of a Che-like signal transduction cascade in
regulating cyst cell development in Rhodospirillum centenum. Mol Microbiol. 2005
Jun;56(6):1457–66.
3.     Bible A, Russell MH, Alexandre G. The Azospirillum brasilense Che1 chemotaxis
pathway controls swimming velocity, which affects transient cell-to-cell clumping. J
Bacteriol. 2012 Jul;194(13):3343–55.
4.     Collins KD, Lacal J, Ottemann KM. Internal sense of direction: sensing and signaling
from cytoplasmic chemoreceptors. Microbiol Mol Biol Rev MMBR. 2014 Dec;78(4):672–84.
5.     Bilwes AM, Alex LA, Crane BR, Simon MI. Structure of CheA, a signal-transducing
histidine kinase. Cell. 1999 Jan 8;96(1):131–41.
6.     Griswold IJ, Zhou H, Matison M, Swanson RV, McIntosh LP, Simon MI, et al. The
solution structure and interactions of CheW from Thermotoga maritima. Nat Struct Biol.
2002 Feb;9(2):121–5.
7.     Reebye V, Frilling A, Hajitou A, Nicholls JP, Habib NA, Mintz PJ. A perspective on non-
catalytic Src homology (SH) adaptor signalling proteins. Cell Signal. 2012 Feb;24(2):388–92.
8.     Li X, Fleetwood AD, Bayas C, Bilwes AM, Ortega DR, Falke JJ, et al. The 3.2 Å
resolution structure of a receptor: CheA:CheW signaling complex defines overlapping
binding sites and key residue interactions within bacterial chemosensory arrays.
Biochemistry. 2013 Jun 4;52(22):3852–65.
9.     Khursigara CM, Wu X, Subramaniam S. Chemoreceptors in Caulobacter crescentus:
trimers of receptor dimers in a partially ordered hexagonally packed array. J Bacteriol. 2008
Oct;190(20):6805–10.
10.    Briegel A, Ortega DR, Tocheva EI, Wuichet K, Li Z, Chen S, et al. Universal architecture
of bacterial chemoreceptor arrays. Proc Natl Acad Sci U S A. 2009 Oct 6;106(40):17181–6.
11.    Briegel A, Li X, Bilwes AM, Hughes KT, Jensen GJ, Crane BR. Bacterial chemoreceptor
arrays are hexagonally packed trimers of receptor dimers networked by rings of kinase and
coupling proteins. Proc Natl Acad Sci U S A. 2012 Mar 6;109(10):3766–71.
12.    Sourjik V, Berg HC. Localization of components of the chemotaxis machinery of
Escherichia coli using fluorescent protein fusions. Mol Microbiol. 2000 Aug;37(4):740–51.
13.    Wadhams GH, Martin AC, Warren AV, Armitage JP. Requirements for chemotaxis
protein localization in Rhodobacter sphaeroides. Mol Microbiol. 2005 Nov;58(3):895–902.
14.    Sourjik V, Berg HC. Functional interactions between receptors in bacterial
chemotaxis. Nature. 2004 Mar 25;428(6981):437–41.
15.    Ames P, Parkinson JS. Conformational suppression of inter-receptor signaling defects.
Proc Natl Acad Sci U S A. 2006 Jun 13;103(24):9292–7.
16.    Li M, Hazelbauer GL. Selective allosteric coupling in core chemotaxis signaling
complexes. Proc Natl Acad Sci U S A. 2014 Nov 11;111(45):15940–5.
17.    Piñas GE, Frank V, Vaknin A, Parkinson JS. The source of high signal cooperativity in
bacterial chemosensory arrays. Proc Natl Acad Sci U S A. 2016 Mar 22;113(12):3335–40.
18.    Blackhart BD, Zusman DR. "Frizzy" genes of Myxococcus xanthus are involved in
control of frequency of reversal of gliding motility. Proc Natl Acad Sci U S A. 1985
Dec;82(24):8767–70.

19.    Treuner-Lange A, Macia E, Guzzo M, Hot E, Faure LM, Jakobczak B, et al. The small G-
protein MglA connects to the MreB actin cytoskeleton at bacterial focal adhesions. J Cell
Biol. 2015 Jul 20;210(2):243–56.
20.    Guzzo M, Agrebi R, Espinosa L, Baronian G, Molle V, Mauriello EMF, et al. Evolution
and Design Governing Signal Precision and Amplification in a Bacterial Chemosensory
Pathway. PLoS Genet. 2015 Aug 20;11(8):e1005460.
21.    Chang Y-W, Rettberg LA, Treuner-Lange A, Iwasa J, Søgaard-Andersen L, Jensen GJ.
Architecture of the type IVa pilus machine. Science. 2016 Mar 11;351(6278):aad2001.
22.    Bustamante VH, Martinez-Flores I, Vlamakis HC, Zusman DR. Analysis of the Frz signal
transduction system of Myxococcus xanthus shows the importance of the conserved C-
terminal region of the cytoplasmic chemoreceptor FrzCD in sensing signals. Mol Microbiol.
2004 Sep;53(5):1501–13.
23.    McBride MJ, Weinberg RA, Zusman DR. "Frizzy" aggregation genes of the gliding
bacterium Myxococcus xanthus show sequence similarities to the chemotaxis genes of
enteric bacteria. Proc Natl Acad Sci U S A. 1989 Jan;86(2):424–8.
24.    Concepcion J, Witte K, Wartchow C, Choo S, Yao D, Persson H, et al. Label-free
detection of biomolecular interactions using BioLayer interferometry for kinetic
characterization. Comb Chem High Throughput Screen. 2009 Sep;12(8):791–800.
25.    Inclán YF, Vlamakis HC, Zusman DR. FrzZ, a dual CheY-like response regulator,
functions as an output for the Frz chemosensory pathway of Myxococcus xanthus. Mol
Microbiol. 2007 Jul;65(1):90–102.
26.    Kaimer C, Zusman DR. Regulation of cell reversal frequency in Myxococcus xanthus
requires the balanced activity of CheY-like domains in FrzE and FrzZ. Mol Microbiol. 2016
Apr;100(2):379–95.
27.    Moine A, Espinosa L, Martineau E, Yaikhomba M, Jazleena PJ, Byrne D, et al. The
nucleoid as a scaffold for the assembly of bacterial signaling complexes. PLOS Genet. 2017
Nov 21;13(11):e1007103.
28.    Liu J, Hu B, Morado DR, Jani S, Manson MD, Margolin W. Molecular architecture of
chemoreceptor arrays revealed by cryoelectron tomography of Escherichia coli minicells.
Proc Natl Acad Sci U S A. 2012 Jun 5;109(23):E1481-1488.
29.    Cassidy CK, Himes BA, Alvarez FJ, Ma J, Zhao G, Perilla JR, et al. CryoEM and computer
simulations reveal a novel kinase conformational switch in bacterial chemotaxis signaling.
eLife. 2015 Nov 19;4.
30.    Moine A, Agrebi R, Espinosa L, Kirby JR, Zusman DR, Mignot T, et al. Functional
organization of a multimodular bacterial chemosensory apparatus. PLoS Genet. 2014
Mar;10(3):e1004164.
31.    Adams DW, Wu LJ, Errington J. Nucleoid occlusion protein Noc recruits DNA to the
bacterial cell membrane. EMBO J. 2015 Feb 12;34(4):491–501.
32.    McBride MJ, Köhler T, Zusman DR. Methylation of FrzCD, a methyl-accepting taxis
protein of Myxococcus xanthus, is correlated with factors affecting cell behavior. J Bacteriol.
1992 Jul;174(13):4246–57.
33.    Mauriello EMF, Nan B, Zusman DR. AglZ regulates adventurous (A-) motility in
Myxococcus xanthus through its interaction with the cytoplasmic receptor, FrzCD. Mol
Microbiol. 2009 May;72(4):964–77.
34.    Nan B, Mauriello EMF, Sun I-H, Wong A, Zusman DR. A multi-protein complex from
Myxococcus xanthus required for bacterial gliding motility. Mol Microbiol. 2010
Jun;76(6):1539–54.

35. Cole C, Barber JD, Barton GJ. The Jpred 3 secondary structure prediction server.
Nucleic Acids Res. 2008 Jul 1;36(suppl 2):W197–201.
36. Sievers F, Higgins DG. Clustal omega. Curr Protoc Bioinforma Ed Board Andreas
Baxevanis Al. 2014;48:3.13.1-16.
37. Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Curr
Protoc Bioinforma Ed Board Andreas Baxevanis Al. 2016;54:5.6.1-5.6.37.
38. Park S-Y, Borbat PP, Gonzalez-Bonet G, Bhatnagar J, Pollard AM, Freed JH, et al.
Reconstruction of the chemotaxis receptor-kinase assembly. Nat Struct Mol Biol. 2006
May;13(5):400–7.
39. Krissinel E, Henrick K. Secondary-structure matching (SSM), a new tool for fast
protein structure alignment in three dimensions. Acta Crystallogr D Biol Crystallogr. 2004
Dec;60(Pt 12 Pt 1):2256–68.
40. Gouet P, Robert X, Courcelle E. ESPript/ENDscript: extracting and rendering sequence
and 3D information from atomic structures of proteins. Nucleic Acids Res. 2003 Jul
1;31(13):3320–3.
41. Mignot T, Merlie JP, Zusman DR. Regulated pole-to-pole oscillations of a bacterial
gliding motility protein. Science. 2005 Nov 4;310(5749):855–7.
42. Ducret A, Maisonneuve E, Notareschi P, Grossi A, Mignot T, Dukan S. A microscope
automated fluidic system to study bacterial processes in real time. PloS One.
2009;4(9):e7282.
43. Mauriello EMF, Jones C, Moine A, Armitage JP. Cellular targeting and segregation of
bacterial chemosensory systems. FEMS Microbiol Rev. 2018 Jul 1;42(4):462–76.
44. Ducret A, Quardokus EM, Brun YV. MicrobeJ, a tool for high throughput bacterial cell
detection and quantitative analysis. Nat Microbiol. 2016 20;1(7):16077.
25

**FIGURE LEGEND**

**Fig. 1.** Schematic representation of the supramolecular organization of Che proteins. (A)

MCP form trimers of dimers (each dimer is shown as a green circle), which, in turn, form

hexagons connected with rings composed of the CheA-P5 domain (dark blue bars) and CheW

(white bars). The light blue circles represent the CheA-P4 domain and, the red circles the

interface between the β-strands 3 and 4 of subdomain 1 of CheA-P5 and the β-strands 4 and

5 of subdomain 2 of CheW. Rings containing six CheW proteins (shown at the center of the

array) might serve to modulate the stability and activation of the system. A signaling unit is

represented in the red box (Adapted from (43)). (B) FrzCD, FrzE$^{CheA}$, FrzA and FrzB proteins

organization depicted by homology with Che proteins. (C) Schematic representation of the

*frz* operon.

**Fig. 2.** The effect of FrzB on the IAA response and the location of the *frzB* gene in the Frz

pathway. (A) Motility and fruiting body formation phenotypes were photographed at 48h

and 72h, respectively. (B) Box plots of reversal frequencies of single cells moving on agar

pads supplemented or not with 0.15% IAA. The lower and upper boundaries of the boxes

correspond to 25th and 75th percentiles, respectively. The median is shown as a line at the

center of each box, and whiskers represents the 10th and 90th percentiles. For the reversal

frequency measurements, approximately 500 cells from three to six biological replicates,

were used. *frzCDc* corresponds to the allelic variant *frzCDΔ6-182*.

**Fig. 3.** Interaction of FrzB with FrzCD and FrzE. (A) GST affinity chromatography co-

purification of the indicated pairs. A, B, CD and E stand for GST-FrzA, GST-FrzB, 6His-FrzCD

and 6His-FrzE$^{CheA}$, respectively. In the first gel, lanes are issued from the same gel and were

grouped for simplicity. (B) In the left panel, BLi sensograms show the binding of different

amounts of GST-FrzB with immobilized 6His-FrzCD. BlitZ ProTM was used to calculate the

affinity constant (Kd = 0.7 µM), using a global 1:1 fit. The central panel shows the responses

at 145 s (nm, x axis) (average of two biological replicates) of the binding of immobilized 6His-

FrzCD with different concentrations of GST-FrzB (µM, y axis). Data were analyzed by

GraphPad Prism 5.0, based on steady-state levels of the responses. In the right panel, BLi

sensograms show the binding of different amounts of GST-FrzB with immobilized 6His-FrzCD

or 6His-FrzE$^{CheA}$ for comparison. A reference subtraction (sensor reference and GST-FrzB at

10 µM) was applied to account for non-specific binding, background, and signal drift to

minimize sensor variability. (C) The effect of the FrzB$^{β4-β5}$ chimera on the FrzE

phosphorylation. The FrzE kinase domain (FrzE$^{CheA}$) auto-phosphorylation was tested *in*

*vitro* by incubation of FrzE$^{CheA}$ in the presence of FrzA, FrzB or FrzB$^{β4-β5}$, the indicated

different forms of FrzCD and ATPγP$^{33}$ as a phosphate donor. (D) 3D models of the FrzE$^{P4-}$

$^{P5}$:FrzA and FrzE$^{P4-P5}$:FrzB complexes based on the *T. maritima* CheA$^{P4-P5}$:CheW crystal

structure (PDB 2ch4). The 20 amino acid region forming the β-strands 4 and 5 of subdomain

2 and corresponding to the CheW region important for the CheW:CheA$^{P5}$ interaction are

highlighted in red in the FrzA structure. These residues are missing in FrzB. (E) Protein

structural alignments between the *T. maritima* CheW x-ray structure and *M. xanthus* FrzA

and FrzB theoretical 3D models. CheW residues in atomic contact with CheA and MCP are

shown as red stars and blue squares, respectively. Secondary structures determined from

CheW x-ray structure are shown on top of the alignment.

**Fig. 4**. Localization of FrzB-mCherry in *M. xanthus* cells. (A) (Left) Micrographs of *mCherry-*

*frzB* cells stained with the DNA DAPI stain. (Middle) mCherry (red) and DAPI fluorescence

(blue) profiles are shown with the fluorescence intensity (arbitrary units) represented on the

*y*-axis and the cell length positions with -1 and +1 indicating the poles, on the *x*-axis. The cell

boundaries were drawn manually from the phase-contrast images. (Right) Plot of correlation

coefficients of DAPI and mCherry localization. $R^2$ values > 0.5 indicate significant

correlations. The table compares the $R^2$ values between the indicated fluorescent fusion and

DAPI staining. (B) (Left) Fluorescence micrographs of the indicated *M. xanthus* strains

carrying FrzCD-gfp fusions. The cell boundaries were drawn manually from the phase-

contrast images. (Right) For each indicated strain, more than 120 cells (*x* axis) from at least

two biological replicates are represented as lines and ordered according to their length

(pixels) in demographs. The GFP fluorescence intensity along the cell body is represented as

colored pixels at the corresponding cell position (from -1 to +1 on the *y* axis). "0" is the cell

center. On the right, a scale indicates the fluorescence intensity and the corresponding

colors.

**Fig. 5.** Proposed model on how FrzB might confer plasticity to Frz clusters. (A) FrzB

participates to the formation of 6-CheW rings to stabilize small Frz clusters, which would

otherwise collapse in one single cluster (C). (B) Alternatively FrzB allows the formation of

small Frz clusters by breaking the networking of MCP trimers of dimers with Che[P5]-CheW

rings.

22
23

**SUPPLEMENTARY FIGURE AND TABLE LEGEND**

**Figure S1.** (A) Motility and fruiting body formation phenotypes were photographed at 48h

and 72h, respectively. (B) Box plots of reversal frequencies of single cells moving on agar

pads supplemented or not with 0.15% IAA. The lower and upper boundaries of the boxes

correspond to 25% and 75% percentiles, respectively. The median is shown as a line at the

center of each box and whiskers represents the 10% and 90% percentiles. For the reversal

frequency measurements, approximately 120 cells, issued from two biological replicates,

were used. For WT approximately 500 cells, issued from six biological replicates, were used.

**Figure S2.** Western blot with anti-FrzB antibodies on the cell extracts of the indicated *M.*

*xanthus* strains. The *pilA::frzB* strain, used as positive control, expresses *frzB* under the

control of the *pilA* promoter for high expression.  The white line is used to indicate that two

lanes from the same gel were separated by other lanes in the original gel.

**Figure S3.** Motility and fruiting body formation phenotypes of the indicated strains were

photographed at 48h and 72h, respectively.

**Figure S4.** Western blot with anti-FrzCD antibodies on the cell extracts of the indicated *M.*

*xanthus* strains.

**Figure S5**. (A) Cells of the indicated strains were incubated 30 minutes with DAPI and then

imaged at the fluorescence microscope. The nucleoid and cell surfaces were measured

automatically with Microbe J (44). (B) The ratio between the nucleoid and cell surface was

then calculated for each cell and averages values plotted. The numbers of analyzed cells are

indicated per each strain. Cells were issued from two independent biological replicates.

**Table S1.** List of strains used in this work.

**Table S2.** List of plasmids used in this work.

**A**

Signaling unit

**B**

FrzCD (MCP)

FrzA
(CheW)

FrzB
(CheW)

P5

P3

P4

FrzE^CheA
(CheA)

P2

P1

P

**C**

*frzZ
(cheY-cheY)*

*frzA
(cheW)*

*frzB
(cheW)*

*frzCD
(mcp)*

*frzE
(cheA-cheY)*

*frzG
(cheB)*

*frzF
(cheR)*

**A**

| | wt | ΔfrzA | ΔfrzB | ΔfrzA ΔfrzB frzB$^{Aβ4β5}$ | frzCDc | frzCDc ΔfrzA | frzCDc ΔfrzB |
|---|---|---|---|---|---|---|---|
| **Swarming** | | | | | | | |
| **Fruiting bodies** | | | | | | | |
| **IAA response** | | | | | | | |

**B**

**A**

| Protein fusion | R² * | Source |
|---|---|---|
| FrzCD-GFP | 0,76 | Moine et al., 2016 |
| FrzE-mCherry | 0,80 | Moine et al., 2016 |
| mCherry-FrzB | 0,62 | This study |
| FrzCDΔ6-130-GFP | 0,08 | Moine et al., 2016 |

* Average of 10 cells

**B**

FrzCD-GFP

**(A) WT**

NUCLEOID

CYTOPLASM

FrzCD
FrzE-P5
FrzE-P4
FrzA
FrzB

**(B) WT**

**(C) ΔfrzB**

|  | **S-motility** | **Fruiting body formation** |

*wt*

*frzB-mCherry*

Δ*frzB*

**Table S1. Strain list**

| Strains | Genotype | Reference or source |
|---------|----------|---------------------|
| DZ2 | *wt* | Zusman et al., 1982 |
| DZ4478 | *ΔfrzA* | Bustamante et al ; 2004 |
| DZ4479 | *ΔfrzB* | Bustamante et al ; 2004 |
| DZ4487 | *frzCDΔ6-182* | Bustamante et al ; 2004 |
| DZ4480 | *ΔfrzCD* | Bustamante et al ; 2004 |
| EM538 | *ΔfrzA frzCDΔ6-182* | This study |
| EM539 | *ΔfrzB frzCDΔ6-182* | This study |
| DZ4620 | *frzCD-gfp* | Mauriello et al., 2009 |
| EM440 | *ΔfrzB frzCD-gfp* | This study |
| EM512 | *ΔfrzA frzCD-gfp* | This study |
| EM515 | *frzB-mCherry* | This study |
| EM605 | *ΔfrzA ΔfrzB frzB$^{β4-β5}$* | This study |
| EM736 | *ΔfrzA ΔfrzCD* | This study |
| EM742 | *ΔfrzA ΔfrzB* | This study |
| EM743 | *ΔfrzB ΔfrzCD* | This study |
| EM699 | *ΔfrzB frzB$^{β4-β5}$* | This study |
| EM700 | *ΔfrzA frzB$^{β4-β5}$* | This study |
| EM773 | *ΔfrzA ΔfrzB frzCD-gfp* | This study |

**Table S2. Plasmid list**

| Plasmid | Expression plasmid | Source |
|---|---|---|
| pEM411 | pBJ113 with construct for *ΔfrzA frzCDΔ6-182* | This study |
| pEM412 | pBJ113 with construct for *ΔfrzB frzCDΔ6-182* | This study |
| pEM463 | pBJ113 with construct *ΔfrzA ΔfrzB frzB$^{β4-β5}$* (Biomatik) | This study |
| pEM524 | pBJ114 with construct *ΔfrzA ΔfrzB* | This study |
| pEM525 | pBJ114 with construct *ΔfrzA* in DZ4480 | This study |
| pEM526 | pBJ114 with construct *ΔfrzB ΔfrzCD* | This study |
| pEM536 | pBJ114 with construct *ΔfrzA frzB$^{β4-β5}$* | |
| pEM537 | pBJ114 with construct *ΔfrzB frzB$^{β4-β5}$* | |
| pDPA20 | pKY480 with construct for *frzCD-gfp* | Mauriello et al., 2009 |
| pEM399 | pBJ113 with construct for *frzB-mCherry* | This study |
| pETPhos *frzCD* | pETPhos with *frzCD* tagged with 6-his inducible with IPTG | Guzzo et al., 2015 |
| pETPhos *frzE$^{CheA}$* | pETPhos with *frzE$^{kinase}$* tagged with 6-his inducible with IPTG | Guzzo et al., 2015 |
| pETPhos *frzCD$^c$* | pETPhos with *frzCD$^c$* tagged with 6-his inducible with IPTG | Guzzo et al., 2015 |
| pGEX(M) *frzA* | pGEX (M) with *frzA* tagged with GST | Guzzo et al., 2015 |
| pEM464 | pETPhos with *frzB$^{β4-β5}$* tagged with 6-his inducible with IPTG (Biomatik) | This study |