**Letter**

**An unconventional NOI/RIN4 domain of a rice NLR protein binds host EXO70 protein to confer fungal immunity**

**Koki Fujisaki[1], Yoshiko Abe[1], Eiko Kanzaki[1], Kazue Ito[1], Hiroe Utsushi[1], Hiromasa Saitoh[2], Aleksandra Białas[3], Mark J Banfield[4] Sophien Kamoun[3], Ryohei Terauchi[1,5]**

**Affiliation**

  **1** Iwate Biotechnology Research Center, Kitakami, Iwate, Japan, **2** Department of Molecular Microbiology, Tokyo University of Agriculture, Setagaya, Tokyo, Japan, **3** The Sainsbury Laboratory, Norwich Research Park, Norwich, UK**, 4** Dept. of Biological Chemistry, John Innes Centre, Norwich Research Park, Norwich, UK, **5** Laboratory of Crop Evolution, Kyoto University, Muko, Kyoto, Japan

A subset of plant nucleotide-binding domain and leucine-rich repeat-containing

(NLR) proteins carry extraneous integrated domains that have been proposed to mediate

pathogen effector recognition. The current view is that these unconventional domains

function by directly binding or serving as substrates for pathogen effectors, yet only a

few domains have been functionally characterized to date. Here we report that the

integrated NOI domain of the rice NLR protein Pii-2, together with its partner Pii-1,

mediates immunity to the rice blast fungus *Magnaporthe oryzae* by indirect recognition

of the AVR-Pii effector. We discovered that the Pii-2 NOI domain does not physically

interact with the effector itself but instead binds the host protein OsExo70-F3, which is

a target of AVR-Pii. We further identified mutations within the NOI core motif

(PxFGxW) of Pii-2 that abolish both OsExo70-F3 binding and *Pii*-mediated resistance

to *M. oryzae* expressing AVR-Pii. This led us to propose a novel conceptual model in

which an NLR-integrated domain functions to detect host proteins targeted by pathogen

effectors, in a framework that extends classical indirect recognition models.

(174 words)

Plants have evolved immune receptors to defend against pathogen infection. These

include cytosolic nucleotide-binding domain and leucine-rich repeat-containing (NLR)

proteins that recognize pathogen avirulence (AVR) effectors and activate a complex

immune response that includes hypersensitive cell death. Plant NLRs have a modular

architecture comprising either an N-terminal coiled-coil (CC) or a Toll/interleukin-1

receptor (TIR) homology domain, followed by nucleotide binding (NB) and leucine rich

repeat (LRR) domains. To date, several reports have shown that genetically-linked NLR

pairs (e.g., *RPS4/RPS1*, *RGA4/RGA5* and *Pikp-1/Pikp-2*) are required to act together for

the recognition of cognate AVRs. Of these pairs, one NLR is a sensor NLR that

perceives pathogen effectors while the other is a helper NLR that is necessary to

activate immune signaling (ref. 1,2). A number of sensor NLRs, perhaps as many as

10%, have recently been reported to contain additional unconventional domains called

integrated domains (IDs) that appear to have originated from host targets of effectors

(ref. 3,4). The current view is that NLR-integrated domains function as baits or decoys

that bind pathogen effectors or serve as substrates of effectors, in order to detect

invading pathogens. One example is *Arabidopsis* RRS1, an NLR that carries a

C-terminal domain with similarity to WRKY transcriptional factors. RRS1 is required

for the perception of bacterial effectors PopP2 and AvrRps4, which are bacterial

effectors that target *Arabidopsis* WRKY proteins to enhance virulence. Therefore, Le

Roux *et al*. (2015) concluded that a WRKY domain targeted by pathogen effectors has

integrated into RRS1 to facilitate recognition of the bacterial effectors (ref. 5,6). Other

examples include the rice NLR pairs Pia (RGA4/RGA5) and Pikp (Pikp-1/Pikp-2),

which recognize the effectors AVR-Pia and AVR-PikD of the rice blast fungus,

*Magnaporthe oryzae* (syn. *Pyricularia oryzae*), respectively. Remarkably, both RGA5

and Pikp-1 include a Heavy Metal-Associated (HMA) domain integrated in different

positions in these NLRs. Binding of the integrated HMA domain to the corresponding

AVR effectors triggers the hypersensitive response and immunity to the blast fungus

(ref. 7,8). These findings let to the proposal that unconventional integrated domains of

NLRs have evolved to directly bind pathogen effectors or serve as a substrate for the

effectors's activity (ref. 9). As yet, the degree to which IDs associate with host proteins

is largely unknown.

The *Pii* locus in rice encodes a pair of CC-NLR proteins, Pii-1 and Pii-2, that detect

the *M. oryzae* effector AVR-Pii and mount an effective immune response against races

of this pathogen that express AVR-Pii (ref. 10,11). Despite repeated attempts using

various protein-protein interaction methods, we did not detect direct interaction between

AVR-Pii and either one of Pii-1 and Pii-2 (ref. 12). Instead, we discovered that AVR-Pii

binds two members of the large EXO70 protein family of rice, OsExo70F-2 and

OsExo70F-3 (ref. 12). EXO70 is a subunit of the exocyst, an evolutionarily conserved

vesicle tethering complex that functions in the last stage of exocytosis (ref. 13,14).

Although we do not understand the degree to which AVR-Pii binding to EXO70

contributes to virulence, we found that *OsExo70-F3* is genetically required for

*Pii*-mediated immunity to *M. oryzae* expressing *AVR-Pii* (ref. 12). This led us to

propose that Pii response to AVR-Pii follows the indirect recognition model, in which

the NLR protein recognizes modifications of a host protein targeted by the effector (ref.

12).

Here we report that Pii recognition of AVR-Pii follows a novel conceptual model of

NLR perception of plant pathogens. We found that the 280 C-terminal amino acids of

Pii-2 (Pii-2-CT) bind the rice protein OsExo70-F3 that is targeted by the AVR-Pii

effector (Fig. 1, Supplementary Fig. 1, 2 and 3). We further delineated the interaction

region to the 103 amino acids at the C-terminus of the protein (Pii-2-CTC)

(Supplementary Fig. 1, 2 and 3). Interestingly, this region of Pii-2 contains a short

sequence with similarity to a core motif of nitrate-induced (NOI) domain (PxFGxW)

that is also conserved in the *Pi5-2* which is allelic to *Pii-2* (Supplementary Fig. 3 and ref.

3,15). The NOI domain also defines the *Arabidopsis* RPM1-INTERACTING

PROTEIN4 (RIN4), which is known to negatively regulate plant defense and is targeted

by bacterial effectors AvrRpt2, AvrB and AvrRpm1 (ref. 16). In fact, the protease

effector AvrRpt2 cleaves RIN4 protein at the NOI motif PxFGxW. To further

investigate the importance of the Pii-2 NOI motif, we introduced multiple mutations in

this region and tested the resulting mutants for binding to OsExo70-F3 (Fig.1,

Supplementary Fig. 4). Mutations in the NOI conserved residues, Pro(P), Phe(F) and

Trp(W) abolished binding to OsExo70-F3 (Fig. 1, Supplementary Fig. 4 and 5). This

indicates that the Pii-2 NOI core motif is required for OsExo70-F3 binding.

Next, we investigated the role of the NOI motif in *Pii*-mediated immunity. We

determined the effect of mutations within the NOI core motif on *Pii*-mediated resistance

to *M. oryzae* using genetic complementation of a *pii-2* loss-of-function rice line with

*Pii-2* transgenes carrying the mutations in the NOI motif described above. To obtain a

*pii-2* loss-of-function mutant rice, we screened 5,600 ethyl methanesulfonate (EMS)

mutant lines of rice cultivar Hitomebore, which carries the *Pii* gene (ref. 17). This led to

the identification of two *pii-2* loss-of-function mutants, Hit5882 and Hit13701, carrying

an amino acid substitution (V371D) and a nonsense mutation (Q247stop) in the Pii-2

NB-domain, respectively (Supplementary Fig. 3; ref. 11). A full-length cDNA of

wild-type *Pii-2* under the control of the 35*S* promoter (Hit5882+WT-Pii-2) transformed

in Hit5882 complemented *Pii*-mediated resistance (Fig. 2). In contrast, transformation

of Hit5882 with a full-length cDNA of *Pii-2* carrying the NOI-m5 mutation failed to

confer *Pii*-dependent resistance in all transgenic lines tested (Hit5882+Pii-2-NOI-m5;

Fig. 2). The Pii-2-NOI-m5 mutant also failed to complement the second

*Pii*-loss-of-function rice line Hit13701 (Supplementary Fig. 6). Overall, our data

genetically link Pii-2 NOI motif binding to OsExo70-F3 with *Pii*-mediated resistance to

*M. oryzae* expressing the AVR-Pii effector. This suggests that NOI binding to EXO70

is crucial for Pii function.

The current view is that integrated domains of NLR immune receptors function by

directly binding or serving as substrates for pathogen effectors. Our findings suggest

that NLR-integrated domains may function by detecting host proteins targeted by

pathogen effectors in a novel model in which an integrated domain of an NLR mediates

indirect recognition of a pathogen effector (ref. 18,19) (Fig. 3). We hypothesize that the

rice NLR Pii-2 has integrated a NOI domain originating from a host NOI-EXO70

complex targeted by AVR-Pii. This NOI integration has enabled Pii to monitor

OsExo70-F3 *via* physical interaction, which allowed indirect detection of AVR-Pii (Fig.

3). Recent reports showing RIN4-EXO70 interactions (ref. 20,21) support this

hypothesis.

Our work also points to the NOI-EXO70 complex as a major target of plant pathogen

effectors. These domains have therefore been acquired as sensor domains by NLR

immune receptors, and both NOI and EXO70 are commonly seen integrated into NLRs

(ref. 22, 23). Whether integrated NOI and EXO70 domains can mediate direct

recognition of effectors remains to be determined. Nonetheless, our finding that

NLR-integrated domains can indirectly detect AVR effectors expands the mechanistic

view of pathogen detection by immune receptors—that the interactors of

NLR-integrated domains can be either pathogen effectors, or host proteins that are

targeted by effectors. This work shows the first example of indirect pathogen

recognition *via* an integrated NLR, highlighting how the arms race between plants and

pathogens has driven the emergence of a wealth of molecular interactions and

mechanisms in the fight for survival.

**Methods**

**Plasmids and Plants**

The primers and their sequences used for plasmid constructions are listed in

Supplementary Table 1. For Y2H assay, OsExo70-F3 in pGADT7 (prey vector),

*Pii-1-CC*, *Pii-1-NB*, *Pii-1-LRR+CT*, *Pii-2-CC*, *Pii-2-NB* and *Pii-2-LRR+CT* in

pGBKT7 (bait vector) have been previously constructed (ref. 17). To construct

pGBKT7-Pii-2-CT and pGBKT7-Pii-2-CTC, *Pii-2-CT* and *Pii-2-CTC* fragments were

amplified by PCR using primer sets: KF825f/KF826r and KF827f/KF826r, respectively,

and inserted into pGBKT7 vector by utilizing *Eco*RI and *Bam*HI sites. To prepare bait

vectors of the *Pii-2-CT* mutant, *Pii-2-CT-m5*, the N-terminal half and the C-terminal

half of *Pii-2-CT* fragment were amplified by using primer sets (KF825f/KF831r and

KF830f/KF826r, respectively), and resulting PCR products were mixed and used as

DNA template for second PCR using a primer set (KF825f /KF826r) to amplify

*Pii-2-CT-m5* fragment. Similarly, to construct other pGBKT7-Pii-2-CT mutants, primer

sets (1$^{st}$ PCR : KF825f/KF833r for *Pii-2-CT-VA-N* and KF832f/KF826r for

*Pii-2-CT-VA-C*; KF825f/KF835r for *Pii-2-CT-PA-N* and KF834f/KF826r for

*Pii-2-CT-PA-C*; KF825f/KF837r for *Pii-2-CT-FA-N* and KF836f/KF826r for

*Pii-2-CT-FA-C*; KF825f/KF839r for *Pii-2-CT-GA-N* and KF838f/KF826r for

*Pii-2-CT-GA-C*; KF825f/KF841r for *Pii-2-CT-WA-N* and KF840f/KF826r for

*Pii-2-CT-WA-C*) were used for PCR to amplify the N- and C-terminal half fragments of

*Pii-2-CT* mutants. The N- and C-terminal half fragments of each *Pii-2-CT* mutants were

mixed and used as DNA template for second PCR using a primer set (KF825f /KF826r).

After the second PCR, the resulting 0.85 kb PCR products were inserted into pGBKT7

vector by utilizing *Eco*RI and *Bam*HI sites.

For preparing the expression vectors in rice, full-length coding sequences of *Pii-2*

was amplified from cDNA by PCR using primer set, KF828f/KF829r, and was

introduced into pCAMBIA1300S vector (Cambia, www.cambia.org) by utilizing *Kpn*I

and *Xba*I sites (pCAMBIA-Pii-2). To construct expression vector for Pii-2 mutant

(Pii-2-NOI-m5), two *Pii-2* fragments were amplified by using primer sets

(KF828f/KF831r and KF830f/KF829r, respectively), and resulting PCR products were

mixed and used as DNA template for the second PCR using a primer set (KF828f

/KF829r) to amplify *Pii-2-NOI-m5* fragment. The *Pii-2-NOI-m5* fragment was inserted

into pCAMBIA1300S vector (Cambia, www.cambia.org) by utilizing *Kpn*I and *Xba*I

sites (pCAMBIA-Pii-2-NOI-m5).

To prepare transgenic rice plants expressing wild-type *Pii-2* and mutant *Pii-2*

(Pii-2-NOI-m5), pCAMBIA-Pii-2 and pCAMBIA-Pii-2-NOI-m5 were introduced into

*Agrobacterium tumefaciens*, which were used for transformation of Hit5882 and

Hit13701 that are pii-2 deficient mutants of the rice cultivar Hitomebore (ref. 11).

**Yeast two-hybrid assay**

Y2H assay was performed as described previously (ref. 24). Ten-times dilution series

[OD600 = 3.0 (x1), 0.3 (x10$^{-1}$) and 0.03 (x10$^{-2}$)] of yeast cells were prepared and spotted

onto quadruple dropout medium (QDO); basal medium lacking Trp, Leu, Ade and His

but containing 5-Bromo-4-Chloro-3- indolyl a-D-galactopyranoside (X-a-gal)

(Clontech). To detect interactions, both QDO medium with or without 10 mM

3-amino-1,2,4-triazole (3AT) (Sigma) was used. Yeast cells were also spotted onto

double dropout medium (DDO); basal medium lacking Trp, Leu to test cell viability. To

check protein accumulation in yeast cells, yeast cells were propagated in liquid DDO at

30°C   overnight. Forty mg yeast cells were collected and resuspended with 160 μl GTN

+ DC buffer [10% glycerol, 25mM Tris-HCl (pH 7.5), 150mM NaCl, 1 mM DTT and 1

tablet of complete EDTA-free (Roche, Basel   Switzerland)]. Then, 160 μl of 0.6 N

NaOH was added, mixed gently, and incubated at room temperature for 10 min. Next,

160 μl of gel sample buffer [40%(w/v)  glycerol,  240 mM  Tris-HCl  pH  6.8,  8%  (w/v)

SDS,  0.04%  (w/v)  bromophenol  blue,  400  mM  DTT] was added and incubated at 95 °C

for 5 min. After 20,000 x g centrifugation for 5 min, the supernatant was subjected to

SDS-PAGE. Proteins expressed from bait and prey vectors were immunologically

detected by using anti-Myc antibody (MBL, Nagoya, Japan) and anti-HA 3F10 (Roche),

respectively.

**Assays for fungal pathogenicity and gene expression**

Wild-type *M. oryzae* isolate Sasa2 (no AVR-Pii) is stored at Iwate Biotechnology

Research Center, and a transgenic *M. oryzae* strain (Sasa2+AVR-Pii-s) was previously

established as Sasa2(+22p:pex33) (ref. 17). Rice leaf blade spot inoculation with

conidial suspension ($5 \times 10^5$ conidia/ml) was performed as described previously (ref.

24). Disease lesions were photographed at 10 days after inoculation, and vertical length

of the lesions was measured. To check *Pii-2* expression in transgenic rice plants,

semi-quantitative RT-PCR was performed as described previously (ref.12). Messenger

RNA from *Pii-2* transgene but not from intrinsic gene was specifically detected by

using    a    primer    set    (5'-CTTTCGCGAGCTCGGTACCAACAATG-3'    and

5'-ATCCACATCATTTTGTAACAATAG-3'). As a control, rice *Actin* was detected by

using    a    primer    set    (5'-    CTGAAGAGCATCCTGTATTG    -3'    and    5'-

GAACCTTTCTGCTCCGATGG -3').

**References**

1.  Baggs, E., Dagdas, G. and Krasileva, KV. (2017) NLR diversity, helpers and

integrated domains: making sense of the NLR IDentity. *Curr. Opin. Plant Biol.* 38,

59-67.

2.  Cesari, S.(2017) Multiple strategies for pathogen perception by plant immune

receptors. *New Phytol.* doi:10.1111/nph.14877.

3.  Cesari, S., Bernoux, M., Moncuquet, P., Kroj, T. and Dodds, P.N. (2014) A novel

conserved mechanism for plant NLR protein pairs: the "integrated decoy"

hypothesis. *Front Plant Sci.* 5, 606.

4.  Wu, C.H., Krasileva, K, V., Banfield, M. J., Terauchi, R., and Kamoun, S. (2015)

The "sensor domains" of plant NLR proteins: more than decoys. *Front Plant Sci.*

6,134.

5.  Le Roux, C., Huet, G., Jauneau, A., Camborde, L., Trémousaygue, D., Kraut, A.,

Zhou, B., Levaillant, M., Adachi, H., Yoshioka, H., Raffaele, S., Berthomé, R.,

Couté, Y., Parker, J.E. and Deslandes, L. (2015) A receptor pair with an integrated

decoy converts pathogen disabling of transcription factors to immunity. *Cell.* 161,

1074-1088.

6.  Sarris, P.F., Duxbury, Z., Huh, S.U., Ma, Y., Segonzac, C., Sklenar, J., Derbyshire,

P., Cevik, V., Rallapalli, G., Saucet, S.B., Wirthmueller, L., Menke, F.L., Sohn, K.H.

and Jones, J.D. (2015) A Plant Immune Receptor Detects Pathogen Effectors that

Target WRKY Transcription Factors. *Cell.* 161, 1089-1100.

7.  Cesari, S., Thilliez, G., Ribot, C., Chalvon, V., Michel, C., Jauneau, A., Rivas, S.,

Alaux, L., Kanzaki, H., Okuyama, Y., Morel, J.B., Fournier, E., Tharreau, D.,

Terauchi, R. and Kroj, T. (2013) The rice resistance protein pair RGA4/RGA5

recognizes the Magnaporthe oryzae effectors AVR-Pia and AVR1-CO39 by direct

binding. *Plant Cell* 25, 1463-1481.

8.  Maqbool, A., Saitoh, H., Franceschetti, M., Stevenson, C.E., Uemura, A., Kanzaki,

H., Kamoun, S., Terauchi, R. and Banfield, M.J. (2015) Structural basis of pathogen

recognition by an integrated HMA domain in a plant NLR immune receptor. *Elife*.

25, 4.

9.  Kroj, T., Chanclud, E., Micel-Romiti, C., Grand, X. and Morel, JB. (2016)

Integration of decoy domains derived from protein targets of pathogen effectors into

plant immune receptors is widespread. *New Phytol*. 210, 618-626.

10. Takagi, H., Uemura, A., Yaegashi, H., Tamiru, M., Abe, A., Mitsuoka, C., Utsushi,

H., Natsume, S., Kanzaki, H., Matsumura, H., Saitoh, H., Yoshida, K., Cano, L.M.,

Kamoun, S. and Terauchi, R. (2013) MutMap-Gap: whole-genome resequencing of

mutant F2 progeny bulk combined with de novo assembly of gap regions identifies

the rice blast resistance gene Pii. *New Phytol*. 200, 276-283

11. Takagi, H., Abe, A., Uemura, A., Oikawa, K., Utsushi, H., Yaegashi, H., Shimizu,

M., Abe, Y., Kanzaki, H., Saitoh, H., Terauchi, R. and Fujisaki, K. (2017) Rice blast

resistance gene *Pii* is controlled by a pair of NBS-LRR genes *Pii-1* and *Pii-2*.

*bioRXiv* doi. https://doi.org/10.1101/227132

12. Fujisaki, K., Abe, Y., Ito, A., Saitoh, H., Yoshida, K., Kanzaki, H., Kanzaki, E.,

242 Utsushi, H., Yamashita, T., Kamoun, S. and Terauchi, R. (2015) Rice Exo70

243 interacts with a fungal effector, AVR-Pii, and is required for AVR-Pii-triggered

244 immunity. *Plant J*. 83, 875-887.

245 13. Heider, MR. and Muson, M. (2012) Exorcising the exocyst complex. *Traffic* 13,

246 898-907.

247 14. Zhang, Y., Liu, CM., Emons, AM. and Ketelaar, T. (2010) The plant exocyst. *J.*

248 *Integr. Plant Biol*. 52, 138-146.

249 15. Nishimura, M.T., Monteiro, F. and Dangl, J.L. (2015) Treasure your exceptions:

250 unusual domains in immune receptors reveal host virulence targets. *Cell*. 161,

251 957-60.

252 16. Kim, M.G., da Cunha, L., McFall, A.J., Belkhadir, Y., DebRoy, S., Dangl, J.L. and

253 Mackey, D. (2005) Two Pseudomonas syringae type III effectors inhibit RIN4-

254 regulated basal defense in Arabidopsis. *Cell*. 121, 749-759.

255 17. Yoshida, K., Saitoh, H., Fujisawa, S., Kanzaki, H., Matsumura, H., Yoshida, K.,

256 Tosa, Y., Chuma, I., Takano, Y., Win, J., Kamoun, S. and Terauchi, R. (2009)

257 Association genetics reveals three novel avirulence genes from the rice blast fungal

258 pathogen Magnaporthe oryzae. *Plant Cell* 21, 1573-1591

259 18. Dangl, J.L. and Jones, J.D. (2001) Plant pathogens and integrated defence responses

260 to infection. *Nature* 411, 826-833.

261 19. van der Hoorn, R.A. and Kamoun, S. (2008) From Guard to Decoy: a new model for

262 perception of plant pathogen effectors. *Plant Cell* 20, 2009-2017

20. Afzal,, A.J., Kim, J.H. and Mackey, D. (2013) The role of NOI-domain containing

proteins in plant immune signaling. *BMC Genomics*. 14, 327.

21. Sabol, P., Kullich, I. and Zarsky, V. (2017) RIN4 recruits the exocyst subunit

EXO70B1 to the plasma membrane. *J. Exp. Bot.* 68, 3253-3265.

22. Sarris, PF., Cevik, V., Dagdas, G., Jones, JDG. And Krasileva, KV. (2016)

Comparative analysis of plant immune receptor architectures uncovers host proteins

likely targeted by pathogens. *BMC Biol*. 14, 8.

23. Brabham, HJ., Hernández-Pinzón I., Hoden, S., Lorang, J. and Moscou, MJ. (2017)

An ancestral integration at Mla is maintained as a *trans*-species polymorphism.

BioRxiv.

24. Kanzaki, H., Yoshida, K., Saitoh, H., Fujisaki, K., Hirabuchi, A., Alaux, L., Fournier,

E., Tharreau, D. and Terauchi, R. (2012) Arms race co-evolution of *Magnaporthe*

*oryzae AVR-Pik* and rice *Pik* genes driven by their physical interactions. *Plant J*. 72,

894–907.

25. Lee, SK., Song, MY., Seo, YS., Kim, HK., Ko, S., Cao, PJ., Suh, JP., Yi, G., Roh,

JH., Lee, S., An, G., Hahn, TR., Wang, GL., Ronald, P. and Jeon, JS. (2009) Rice

Pi5-mediated resistance to *Magnaporthe oryzae* requires the presence of two

coiled-coil-nucleotide-binding-leucine-rich repeat genes. *Genetics*. 181, 1627–1638.

**Acknowledgements**

We thank members of our lab for discussions and general assistance. We also thank

Matt Moscow and Helen Brabham, The Sainsbury Laboratory, UK for sharing their

unpublished results. This work was supported by Science and Technology Research

Promotion Program for Agriculture, Forestry, Fisheries and Food Industry, Japan,

Grant-in-aid for MEXT (Scientific Research on Innovative Areas 23113009), JSPS

KAKENHI (Grant No. 26850029) and JSPS KAKENHI 15H05779, the Biotechnology

and Biological Sciences Research Council (BBSRC, UK, grants BB/J00453 and

BB/P012574).

**Contributions**

K.F. S.K. and R.T. initiated, conceived and coordinated the project; K.F. identified and

characterized NOI core motif in Pii-2; E.K. generated transgenic rice plants; K.F., Y.A.

and H.U. prepared plamid vectors and performed fungal infection assay and RT-PCR.

K.F. and K.I. performed yeast two hybrid assay. H.S., M.B., S.K. and R.T. supervised

the project; K.F., A.B., S.K., and R.T. wrote the manuscript

**Competing interests**

The authors declare no competing financial interests.

**Corresponding author**

Co-correspondence to Ryohei Terauchi and Koki Fujisaki.

**Figure 1. OsExo70-F3 binds the NOI domain of Pii-2 C-terminal region (Pii-2-CT).**

(A) A schematic diagram of Pii-1 and Pii-2 NLRs. Coiled-coil (CC), nucleotide-binding (NB) and leucine rich repeat (LRR) domains are shown in light grey. OsExo70-F3 binds the C-terminus of the Pii-2 (Pii-2-CT) shown in dark grey (Pii-2-CT) that contains the core motif (PxFGxW) of NOI domain.

(B) Alignment of amino acid sequences around the NOI core motif of *Arabidopsis* RIN4, rice Pi5-2 and Pii-2 proteins. Residues of the NOI core motif are shown in green. Asterisks indicate conserved amino acids among *Arabidopsis* RIN4, Pi5-2 and Pii-2 proteins, which are substituted for Ala (A) in the Pii-2-CT-m5 mutant.

(C) Yeast two hybrid (Y2H) assay showing the interaction of OsExo70-F3 with the wild-type Pii-2-CT but not with the Pii-2-CT-m5 mutant. Protein interactions were tested in the yeast grown on Quadruple dropout (QDO) + 3AT medium (Trp⁻Leu⁻Ade⁻His⁻ X$\alpha$ gal⁺ +10 mM 3AT : right panels). Viability of all transformed yeast cells were demonstrated by growing them on double dropout (DDO) medium (Trp⁻Leu⁻ : left panels). Yeast cells were spotted onto the media as tenfold dilution series (x1, x10⁻¹, x10⁻²). Empty vectors (labeled as "empty") were used as negative controls.

**A**

**B**

**C**

**Figure 2. *AVR-Pii* recognition of *Pii* requires Pii-2 NOI core motif.**

(A) Results of rice leaf blade spot inoculation assay of *M. oryzae*. Two *M. oryzae* isolates, with (transgenic Sasa2+AVR-Pii; top) or without (Sasa2; bottom) *AVR-Pii* were spot inoculated onto the leaves of rice lines Moukoto (Pii-), Hitomebore (Pii+), Hit5882, a *pii-2* Hitomebore mutant, and transgenic Hit5882 lines expressing a full-length cDNA of the wild-type *Pii-2* (Hit5882+WT-Pii-2) and those expressing the mutant version of *Pii-2* with m5 mutation in the NOI core motif (Hit5882+Pii-2-NOI-m5). Representative data of two line each of Hit5882+WT-Pii-2 (W1 and W2) and Hit5882+Pii-2-NOI-m5 (m2 and m3) are shown.

(B) Bar graphs showing disease lesion lengths after inoculation of an *M. oryzae* isolate Sasa2+AVR-Pii onto rice lines from panel A, labelled on the bottom. Here, five lines each from Hit5882+WT-Pii-2 (W1 to W5) and Hit5882+Pii-2-NOI-m5 (m1 to m5) were tested. The graph illustrates average lesion lengths and standard deviations from six inoculated spots per line.

(C) Expression of *Pii-2* transgenes in the tested rice plants. Messenger RNA from *Pii-2* transgene but not from intrinsic gene was specifically detected by semi-quantitative RT-PCR using a combination of primers derived from *Pii-2* gene and the binary vector sequences. Rice *Actin* was used as control.

**Figure 3. A scheme showing possible roles of NLR-integrated domains in the effector recognition.**

(A) A widely accepted model of NLR-ID function. NLR directly interacts with pathogen effector *via* its integrated domain, and subsequently activates immune response. An *Arabidopsis* sensor NLR, RRS1 (top), has the WRKY domain at the C-terminus, and bacterial effectors AvrRps4/PopP2 directly bind to the RRS1 WRKY ID (ref. 5,6). The rice sensor NLRs, Pikp-1 (middle) and RGA5 (bottom), carry integrated HMA domain. Pathogen AVR effectors AVR-PikD and AVR-Pia, respectively, directly bind to their cognate HMA (ref. 7,8). In the current model (ref. 1,2), these sensor NLRs trigger ETI together with their NLR helpers.

(B) A novel model of ID function in indirect detection of pathogen effectors. Pii-2 is an NLR with an ID, NOI core motif (PxFGxW), residing in its C-terminal region. Pii-2-ID directly binds OsExo70-F3, which is necessary for AVR-Pii recognition. OsExo70-F3 also binds AVR-Pii. We propose that Pii-2 monitors OsExo70-F3, the status of which is altered upon binding of AVR-Pii. Thus, Pii-2 ID guards a host protein, OsExo70-F3, that is targeted by a pathogen effector, AVR-Pii.