A complex performance landscape for suction-feeding reveals constraints and

adaptations in a population of reef damselfish

Tal Keren[1,2], Moshe Kiflawi[2,3], Christopher H Martin[4], Victor China[1,2], Ofri Mann[2,3],

Roi Holzman[1,2]

[1] Department of Zoology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv

69978, Israel.

[2] The Inter-University Institute for Marine Sciences, POB 469, Eilat 88103, Israel.

[3] Department of life Sciences, Ben Gurion University, Beer Sheva, Israel.

[4] Biology Department, University of North Carolina at Chapel Hill, 120 South

Rd., Chapel Hill, NC 27599, United States

Keywords: Suction feeding, adaptive peaks, trait diversification, intraspecific

variation, predator-prey dynamics

Correspondence should be sent to: Roi Holzman, holzmaban@post.tau.ac.il,

telephone 972-8-6360111, Coral Beach, Eilat 88103 Israel, P.O.B 469.

Abstract

The ability to predict how multiple traits interact in determining performance is

key to understanding the evolution of complex functional systems. Similar to

Simpson's adaptive landscape, which describes the fitness consequences of varying

morphological traits, performance landscapes depict the performance consequences of

varying morphological traits. Mapping the population's location with respect to the

topographic features of the landscape could inform us on the selective forces

operating on the traits that underlie performance. Here, we used a mechanistic model

derived from first principles of hydrodynamics to construct a hypothetical

performance landscape for zooplankton prey capture using suction feeding. We then

used the landscape to test whether a population of *Chromis viridis*, a coral reef

zooplanktivore, is located on a performance peak or ridge based on measurements of

kinematic variables recorded *in-situ* during undisturbed foraging. Observed trait

combinations in the wild population closely matched regions of high feeding

performance in the landscape, however the population was not located on a local

performance peak. This sub-optimal performance was not due to constraints

stemming from the observed trait correlations. The predominant directions of

variation of the phenotypic traits was tangent to the 'path of steepest ascent' that

points towards the local peak, indicating that the population does not reside on a

"performance ridge". Rather, our analysis suggests that feeding performance is

constrained by stabilizing selection, possibly reflecting a balance between selection

on feeding performance and mechanical or genetic constraints.

Introduction

Understanding the mechanisms that shape adaptive diversification is a major goal in

evolutionary biology. Because morphology can affect performance, it should respond

most strongly to selection on performance (Arnold, 1983; Arnold, 2003; Lande and

Arnold, 1983). The earliest framework to assess the fitness consequences of changing

phenotype was Simpson's "adaptive landscape" (Simpson, 1944; Simpson, 1955),

which complemented Wright's adaptive landscape for genotypes (Wright, 1932). This

approach consists of mapping all possible phenotypes or genotypes onto fitness,

producing a rugged multi-dimensional space in which peaks reflect adaptive

combinations surrounded by maladaptive "valleys". The adaptive landscape was an

attempt to conceptualize the complex relationship between multivariate trait

combinations and their fitness consequences. In practice, linking phenotypes and

fitness has proved difficult, especially when considering multiple traits. As a way to

circumvent these difficulties, Arnold (Arnold, 1983) suggested viewing performance

as a mediator between morphology and fitness. He further suggested that a

performance landscape (i.e. the mapping of all trait combinations to their respective

performance values) should provide valuable insights regarding both the selective and

non-selective forces involved in trait evolution. This is done through an analysis of

the topography of the landscape and the distribution of phenotypic diversity in a

population relative to the landscape (Arnold, 2003; Arnold et al., 2001). Conceptually,

the performance landscape is similar to the adaptive landscape, with the caveat that

performance does not necessarily map linearly to fitness, just as phenotype does not

map linearly to fitness. Therefore, the individual performance surface (defined as the

derivative of performance with respect to trait value; (Arnold, 2003; Arnold et al.,

2001)) is not necessarily equal or proportional to the individual fitness surface (Fear

and Price, 1998; Lande and Arnold, 1983)

The topographic features of the performance landscape reflect the selective

regimes imposed on performance-determining traits. Stabilizing, disruptive, and

correlational selection are reflected by curved landscapes, where performance peaks

correspond to local adaptive optima (Arnold, 2003; Arnold et al., 2001). When

mapping a natural population onto such a landscape, the distance between the

population's mean and the theoretical peak can provide evidence for constraint,

assuming sufficient time for phenotypic evolution (Collar et al., 2009; Hansen, 1997).

For example, Arnold and Bennett (Arnold and Bennett, 1988) reconstructed the

performance surface that relates crawling performance to body and tail vertebral

numbers, and concluded that these two parts of the vertebral column evolve on a

performance ridge, defined as an axis on which morphological evolution will result in

little change to performance (Arnold, 2003; Arnold and Bennett, 1988). However,

generating the landscape by collecting performance data for individuals within the

population has a limited ability to infer performance away from the population means,

and cannot provide predictions on performance in areas outside the trait space of the

observed population. As Arnold put it "It is not clear…that a rugged topography, and

the quandries resulting from local peaks, apply in the case of phenotypic characters.

More importantly, we have no device that can tell us about landscape features far

from the population's average. Mt. Improbable is, after all, Mt. Unknowable. This is

the sense in which the local vision of the landscape is the most tractable." (Arnold,

2003).

An alternative to reconstructing the landscape is to use mechanistic models of

performance. Over the last decades, biomechanical theory and computational methods

were used to mechanistically model many aspects of performance such as swimming

(Fish and Lauder, 2006; Liao, 2007; Sfakiotakis et al., 1999), running (Barasuol et al.,

2013; Kingma et al., 1996; Minetti, 1998), slithering (Hu et al., 2009), flying and

gliding (Paranjape et al., 2012; Wu, 2011) among numerous other examples. These

models are based on first principles, and produce reliable estimates of performance

given a set of phenotypic (often kinematics and morphological) traits that are used as

input variables. Because a mechanistic model can (mathematically) solve performance

for any input values, it is not necessary to obtain input values from a specific

organism, and it is possible to predict performance outside the trait ranges and

combinations observed within the population. Furthermore, the data used to generate

the landscape is independent of the population, enabling one to generate hypotheses

regarding the location of the population relative to the landscape. However,

applications of biomechanical models to understand the distribution of performance-

determining traits on the performance landscape are rare.

Our goal was to construct and visualize a performance landscape using an

empirically-verified mechanistic model of suction-feeding, and map upon it the

distribution of individuals within a wild population of *Chromis viridis*, a Coral Reef

damselfish which feeds predominantly on zooplankton using suction-feeding.

Mapping the population's location with respect to the topographic features of the

performance landscape could inform us on the selective forces operating on the traits

that underlie performance. Specifically, we considered three mutually exclusive

scenarios (Fig. 1). Under scenario I, selection for suction performance has a week

effect on morphological evolution of the traits that underlie performance, leading to a

distribution of phenotypic traits that is unrelated to the performance landscape. In that

case, we would expect that the distribution of performance values in the observed

population would be similar to that of a population placed at random on the landscape

(Fig. 1A). An alternative scenario (II) would be that selection for suction performance

has a dominant effect on morphological evolution, and the population sits at the top of

a performance peak (Fig. 1B). In that case, we would expect that the distribution of

performance values in the observed population will be higher than that of a population

placed at random on the landscape, and that the mean individual performance surface

is zero. Lastly, under scenario III, the population could be under stabilizing selection,

with selection for suction performance countered by trade-offs, genetic of mechanical

constraints (Fig. 1B). In that case, we would expect that the distribution of

performance values in the observed population will be higher than that of a population

placed at random on the landscape and that the selection gradient for each phenotypic

trait is different than zero (Fig. 1C).

Methods

Study system

Chromis viridis is a common Indo-pacific coral reef species , found at depths of

up to 30 m. *C. viridis* live in schools of a few dozen to a few hundred, with adult size

ranging from 27 to 59 mm long (Allen and Randall, 1980). Schools are found

inhabiting branching corals, feeding on drifting zooplankton in the vicinity of their

home coral during the day and sleeping among its branches during the night

(Goldshmid et al., 2004).

Chromis viridis is an exclusive zooplanktivore, using suction- feeding to capture

drifting prey over the coral reefs (Carassou et al., 2008; Coughlin and Strickler, 1990).

Prey capture using suction feeding is a widespread and evolutionarily conserved prey

capture mechanism among teleost fishes (Day et al., 2015; Wainwright et al., 2015).

Suction feeding can be treated as a hydrodynamic interaction between a solid particle

(the prey) and unsteady flows (the suction flow), and can be modeled by tracking the

forces exerted on the prey and its trajectory during the strike (Day et al., 2015;

Holzman et al., 2012). This hydrodynamic modeling allows us to quantitatively

estimate the performance consequences of any given trait combination and to

reconstruct a performance landscape independent of the observed population for any

phenotype (i.e. any combination of values for the input traits).

The mechanism of suction feeding consists of a swift opening of the mouth and

expansion of the oral cavity, which generates a flow of water into the predator's

mouth. Prey that cannot withstand the force of the flow are sucked into the fish's

mouth with the surrounding water (Day et al., 2015; Holzman et al., 2012).

Observations, experiments, and hydrodynamic modeling reveal that feeding

performance is determined by multiple morphological and biomechanical traits. These

traits include the size of the mouth, the speed of mouth opening, and the speed of

closing the distance between the predator and prey through forward swimming, and

jaw protrusion (Day et al., 2015; Holzman et al., 2008a; Holzman et al., 2012). Many

zooplankton species evade those predation attempts by sensing the hydrodynamic

disturbance caused by the approaching predator and the suction flows (Buskey et al.,

2002; Fields and Yen, 1997; Heuch et al., 2007). In copepods, rapid swimming away

from the predator is triggered by the deflection of sensory setae on the antennae in

response to the hydrodynamic disturbance. Highly sensitive prey will react to a

hydrodynamic disturbance when the predator is still too far to capture it, and could

escape (Kiorboe and Visser, 1999; Visser, 2001). Across zooplankton species,

sensitivity thresholds differ between species and developmental stages (Green et al.,

2003; Kiorboe and Visser, 1999; Visser, 2001). Thus, high feeding performance, from

the predator's standpoint, is its ability to capture sensitive prey by minimizing

hydrodynamic disturbance (Gemmell et al., 2014; Holzman and Wainwright, 2009;

Viitasalo et al., 1998), which in turn, increases the range of available prey.

Data acquisition

We filmed prey-acquisition strikes (hereafter "strikes") of *C. viridis* feeding on

naturally occurring drifting zooplankton in their natural habitat. All video sequences

were recorded over a period of three months, from November 2013 to January 2014.

Videos were recorded during daytime, using natural light, as these fish are visual

predators and are only active during the day. Filming was conducted using

synchronized high-speed cameras in a waterproof housing (500 frames per second at

1.3 megapixels; Hispec1, Fastec Imaging inc. San-Diego). The underwater video

system (Fig. 2) was moved between five schools, each consisting of tens to hundreds

of individuals. The cameras were positioned at an angle of ~30° with respect to one

another, and were located ~1 m from the coral inhabited by the focal fish school. The

two cameras were aimed at the same point in space, ensuring multiple views on

objects in the visualized volume. Cameras were calibrated using DLTdv5 (Hedrick,

2008), providing a three-dimensional reconstruction of feeding strikes and enabling

time-resolved measurements of distances and locations in a volume of ~20×25×10 cm.

The system was re-calibrated every filming session, and measurement error (< 1%)

was estimated by filming a ruler that was moved within the visualized volume. Live

video feed from the camera was viewed onshore through a tethered cable. A trained

observer manually triggered the system to record voluntary prey-acquisition strikes of

the naturally foraging fish each time such a strike was observed.

For each recorded strike, we digitized four landmarks on the body of the fish: the

tip of the upper jaw, the tip of the lower jaw and the base of the pectoral fin (Fig 2,

Fig 3). From these landmarks, we calculated the size of the mouth, the location of the

mouth center and the location of the fish's body at each frame. From these

measurements we then derived the values of six kinematic traits that are known to be

important in determining suction feeding performance: *I)* max gape, defined as the

maximal diameter of the fish's mouth; *II)* time to peak gape, defined as the time it

took the fish to open its mouth from 20% to 95% of peak gape; *III)* maximal jaw

protrusion measured during the strike, at the fish's frame of reference; *IV)* time to

peak jaw protrusion, defined as the time it took the fish to protrude its jaws from 20%

to 95% of maximal protrusion distance; *V)* ram speed, defined as the fish's swimming

speed during the strike; and *VI)* timing of peak protrusion, defined as the time

difference between the time of maximal gape and time of maximal jaw protrusion

(Fig 3A). Calculations were made as described in (Holzman et al., 2008b). Overall,

we analyzed a total of 110 strikes in which we could clearly identify and digitize the

landmarks.

The high ratio (~1:20) between the number of fish and the number of sequences

recorded in each school translated to a low probability of repeatedly filming the same

individual. To further verify data independence, we digitized for each fish the base of

the caudal, dorsal, and pelvic fins and used these data to estimate the length and body

depth of each individual. Using the measurement error obtained by repeatedly

measuring the same fish in several points in space, we calculated the confidence

intervals for the measurements of the length and body depth ($\pm$ 2mm). We then

assessed the overlap in size (length and body depth) between all the fish observed in

our recordings. We assumed that when an individual is recorded twice, the measures

of its length and body depth in the one movie would be within the confidence

intervals of these measurements in the other movie. This analysis is conservative, as

there could be multiple individuals with the same length and body depth. Overall, <

10% of the strikes in each school were performed by individuals of the same size.

Therefore, we regarded the strikes as true replicates.

Using SIFF to estimate feeding performance

Estimating feeding performance from each set of six phenotypic traits was done

using the SIFF model (Holzman et al., 2012). SIFF uses a set of parameters (Fig. 3B)

that characterize the prey and describe the kinematics of the mouth and flow speed

during the strike to predict the motion of the prey relative to the mouth during suction

feeding (Holzman et al., 2007; Holzman et al., 2012; Wainwright and Day, 2007). In

this study, we varied SIFF input parameter values based on measured morphology and

strike kinematics of *C. viridis* to estimate feeding performance in 3291 phenotypic

combinations of the 6 aforementioned traits. These combinations were generated by

sampling from the range of phenotypic traits observed in the population, allowing us

to construct the performance landscape independently of the observed strikes. As such,

we were unconstrained by the distribution or interdependence between traits

manifested in the observed strikes. Prey capture performance was defined based on

the sensitivity level of the simulated prey that enabled its escape; trait combinations

that resulted in capture of highly sensitive prey were thus defined as "high

performance strikes" (see below; Fig. 3B,C; (Holzman et al., 2007; Holzman et al.,

2012; Wainwright and Day, 2007)).

SIFF is described in detail in (Holzman et al., 2007; Holzman et al., 2012;

Wainwright and Day, 2007). In brief, SIFF uses the flow field realized during the

strike and a set of parameters that characterize the prey, the body, and the mouth to

predict the motion of prey relative to the mouth during suction-feeding, given the

suction flow generated by the fish and the ability of the prey to move away from it

(Fig 3B). According to SIFF, the movement of the prey relative to the predator's

mouth determines whether prey is captured or not. The total force exerted on the prey

is the sum of five component forces: Drag, acceleration reaction force, the force

resulting from the pressure gradient across the prey, prey swimming forces, and

gravitational forces [the latter will be ignored in the current discussion as most aquatic

organisms are approximately neutrally buoyant]. These forces result from the

differential in speeds and accelerations between the prey and the water around it, as

well as from the gradient of flow across the prey (Holzman et al., 2007; Holzman et

al., 2012; Wainwright and Day, 2007).

Flow visualization studies (Day et al., 2015; Holzman et al., 2008b; Staab et al.,

2012) reveal that the flow field in front of the mouth can be estimated based on mouth

kinematics. We used the linear speed of mouth opening Gs (ms^{-1}; the derivative of

gape with respect to time during mouth opening) to estimate peak flow speed at the

center of the mouth Fs (ms^{-1}) as follows:

$Fs = 1.55 * Gs$ (equation 1)

based on the relationships observed in (Holzman et al., 2008b). The temporal pattern

of the flow was assumed to follow the gape cycle, with flow starting at 20% and

peaking at 95% of maximal gape (Day et al., 2015; Higham et al., 2006; Holzman et

al., 2008b). Based on these flow visualization studies, flow speed at any given point in

space can be estimated given the flow speed at the mouth center and the instantaneous

gape diameter (Day et al., 2015; Higham et al., 2006; Holzman et al., 2008b). From

this data, we could derive temporal and spatial flow gradients, and calculate the force

exerted on the prey.

A widespread predatory strategy across teleosts to rapidly reduce the distance to

the prey is to advance the mouth towards the prey, realized by swimming (ram) and

jaw protrusion (Longo et al., 2016). In the context of feeding on evasive

zoooplanktonic prey, swimming incurs a cost, because the advancing body produces a

hydrodynamic disturbance (sometimes referred to as "bow wave"). Similarly to the

suction flows, the bow wave can be sensed by the prey to trigger an escape response

(Fields and Yen, 1997; Stewart et al., 2013; Yen et al., 1992). Hydrodynamic

disturbance was quantified as strain rate, which can be understood as the rate of

deformation of the flow with respect to time. Strain rate is a useful metric for

hydrodynamic disturbance because higher strain rates cause faster deflection of

sensory antennae of copepods (and other marine invertebrates), ultimately triggering

their escape response (Fields and Yen, 1997; Yen et al., 1992). We used observed

movements of the body and mouth center as SIFF inputs for the distance between the

prey and predator and the properties of the bow wave. The relationship between

swimming speed and the hydrodynamic disturbance generated by the body was

estimated by visualizing the flow field in front of a formalin-fixed *C. viridis* specimen

(standard length = 50 mm) held at different flow speeds (2.5 – 20 cm s^{-1}; n= 10

speeds) in a recirculating flume. Flow visualization and analysis followed the protocol

described in (Holzman and Wainwright, 2009), except that here the fish was mounted

in the flume and not free swimming. These measurements showed that strain rate

increased with increasing swimming speed and decreased with the distance to the prey,

following the equation:

$Strain = 10^{-0.564*d+3.366*ram+0.77}$ (equation 2)

where *strain* is the strain rate (s^{-1}) at a distance *d* (mm) from the mouth, and ram

is the swimming speed (m s^{-1}).

Prey was modeled as a naturally buoyant prolate spheroid (0.1 mm in peak

diameter, 2 mm length) located on the centerline across from the mouth. The initial

location of the prey at the onset of simulation was determined as the point in space

where the mouth center would be at the time of peak gape. The prey was modeled to

escape directly away from the predator. Peak escape force ($2x10^{-5}$ N), escape duration

(10 ms) and latency (1 ms) were adopted from (Buskey and Hartline, 2003; Buskey et

al., 2002). The prey was modeled to escape once it sensed a hydrodynamic

disturbance that exceeded a given strain rate threshold. Strain at the location of the

prey was calculated as the sum of strain rates resulting from the advancing body and

the suction flows (i.e we did not account for a possible interaction between the two

sources of flow).

For each set of parameters, SIFF determines whether the prey was captured or

escaped. To estimate feeding performance, we iteratively ran SIFF for each set of

parameters starting with an unrealistically sensitive prey, which escaped when strain

rate exceeded $10^{-2}\,s^{-1}$. This threshold ensured that all prey will escape in the first

iteration. We then increased strain rate threshold by 5% (thus decreasing prey

sensitivity) and re-ran SIFF until the prey was captured, and recorded the strain rate

threshold in the last simulation (Fig 3C). This value was used as an estimate of the

most sensitive prey that can be captured under a given set of traits, thus representing

feeding performance.

A trait set of 3291 phenotypic combinations was generated by randomly sampling

from the range of phenotypic traits observed in the population (under a uniform

probability distribution). Each trait was sampled independently of the other 5 traits,

thus producing sets of combinations that were not observed in the population studied.

This sampling ensured dense uniform coverage of the phenotypic space, regardless of

the distance from the population mean (Fig S1). Thus, our theoretical calculation of

the performance landscape is independent of the trait covariance relationships in the

focal *Chromis* population.

Estimating the contours of the performance landscape.

Phenotypic data and performance estimates of this simulated population were

used to construct a continuous performance landscape, statistically modeling the

relationship between phenotypic traits and performance in the complex, multivariate,

mechanism of suction-feeding. To estimate the topography of the performance

landscape we used multivariable generalized additive model (GAM) with penalized

cubic regression splines, and shrinkage smoothing to further penalize highly

smoothed terms (Wood et al., 2015). Thus, shrinkage smoothing reduces the

estimated effect of parameters with little explanatory power. Cubic regression splines

were chosen over thin plate regression and simple multiple linear regressions as they

provided the best fit in terms of deviance explained and prediction of SIFF results.

We used GAM to fit splines that represent, in the multivariate space, the linkage

between the six kinematic traits across 3291 simulated individuals with their feeding

performance. Given a model structure specified by a GAM model formula, the GAM

function attempts to find the appropriate smoothness for each spline term (Wood et al.,

2015). The fitted model had an adjusted R-square of 0.899, and explained 90.4% of

the deviance. Of the six traits and their paired interactions, only two interactions

(maximum jaw protrusion with time to peak gape and with gape-protrusion time

difference) were non-significant (and thus were assigned a nearly zero coefficient).

To further test the predictive ability of the GAM surface, we divided the dataset

into training/test datasets, containing randomly sampled 3181 trait sets used to

reconstruct the landscape (training set) and 110 randomly-sampled trait sets used to

test it (test set). We then calculated, for the test set only, a linear regression of strain

rate thresholds derived from SIFF vs. those derived from the landscape. The process

was repeated 100 times. The mean ($\pm$ 95% confidence interval) intercept and slope of

that regression were -0.004 ($\pm$0.33) and 1.009 ($\pm$ 0.015), respectively, and the R^2 was

0.88 ($\pm$ 0.05). Given that the 110 trait combinations of the test set were not used to

compile the performance landscape, we concluded that this model is suitable to

predict feeding performance for any given phenotypic combination within the

observed phenotypic ranges.

Scenario I: the distribution of phenotypic traits is unrelated to the performance

landscape.

To test this scenario, we tested whether the feeding performance of the observed

population is significantly higher than that expected by chance. We generated a set of

simulated populations, by allowing 110 simulated individuals to evolve each trait

separately under a Brownian motion model using the observed mean as the starting

point. Simulation of Brownian motion was conducted using the package "Geiger" in

R statistics (Harmon et al., 2008) using a star phylogeny with branch lengths of 1, and

the Brownian rate parameter estimated for the observed population. Thus, this

simulation (hereafter "uncorrelated BM") assumed no trait correlation and no

phylogenetic structure within our population. We generated 1000 simulated

populations, each with 110 trait sets (simulated "individuals"), and calculated their

feeding performance by projecting their trait values on the performance landscape. If

the distribution of phenotypic traits is unrelated to the performance landscape, we

would expect the median performance of the observed population to fall within the

95% confidence interval of the median performance of the simulated population.

Scenario II: does the population sit at the top of an adaptive peak?

If the observed median feeding performance is significantly higher than that

expected based on the simulated populations, it would be consistent with selection to

maximize suction-feeding performance. If the population is located on the

performance peak, then a small change in the traits that determine performance should

result in a decrease (or no increase) in performance. We mapped the 110 observed

sets of traits onto the performance landscape, and estimated the feeding performance

for each set. We then calculated the "individual performance surface" following

(Arnold, 1983; Arnold, 2003). These performance surfaces are the slope of the

performance surface at the individual's observed location in performance space.

Slopes were calculating numerically by adding a small value (0.1 standard deviation

of the mean) to one of the six traits, and calculating the difference in performance

between the original and shifted set. The process was repeated to calculate the slope

for each trait. We further used response surface modeling in the RSM package in R

(Lenth, 2009) to locate the stationary point of the response surface, which indicated a

performance peak (or trough, depending on the mapped performance; (Brodie et al.,

1995)).

Scenario III: high-performance population resides off the performance peak

If the mean individual performance surface are greater than zero, it would mean

that the population is located off the performance peak. In this case, trait variance and

covariance could be indicative of the processes generating the off-peak location. For

example, trait correlations could limit the realized distribution of performance-

determining traits to off-peak locations. To test this hypothesis, we randomized the

order of values in each trait, thereby breaking the correlations between this trait and

all the other traits. We then compared the distribution of performance (strain rate

threshold) to that in the observed population. This was done sequentially for each trait.

Additionally, we generated a second set of simulated populations by allowing 110

simulated individuals to move in a correlated Brownian motion manner (using the

package "Geiger" in R (Harmon et al., 2008), using the observed mean as the starting

point and the observed variance-covariance matrix of the observed population, again

assuming no phylogenetic structure within our population (hereafter "correlated BM)".

We generated 1000 simulated populations, each with 110 trait sets (simulated

"individuals"), and calculated their feeding performance by projecting their trait

values on the performance landscape.

Trait correlations does not necessarily constrain performance. Arnold (Arnold,

2003) suggested that phenotypic traits will evolve along performance ridges that

transverse the multidimensional space. If such a ridge in the landscape guides the

distribution of the traits in the population, we would expect that the major axis of trait

variation in the population to align with the direction of the ridge in the performance

space, i.e. parallel (180° or 0°) to the direction to the peak. We therefore used a

Response-Surface Method (RSM) (Lenth, 2009) to detect the path of steepest descent.

This path is the vector, in the multi-dimension space, which will result in the largest

performance increase for a 0.1 standard deviations change in the phenotypic traits

(results were identical for 0.5 and 1 SD). We quantified the primary axis of variation

in each phenotypic trait using the eigenvectors of PC1 from a correlation based PCA

analysis of the 110 observed strikes. We then calculated the angle in the 6-

dimentional space, between the two vectors as:

$$cos^{-1}\left(\frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\|\|\vec{v}\|}\right)$$

where $\vec{u}$ is the vector of coefficients (eigenvectors) for PC1 and $\vec{v}$ is the vector of

steepest descents from the RSM analysis, the numerator is the dot product of the two

vectors, and the denominator is the product of the vectors' norm.

Results

Performance and phenotypic attributes of the observed population

Overall, the morphological and kinematic characteristics of suction-feeding in the

observed population of *C. viridis* could be described as a swift opening of the mouth

(median time to peak gape = 20 ms), to a relatively small maximal diameter (median

= 2.6 mm), with a fast jaw protrusion (median time to peak protrusion = 23 ms),

reaching its maximal distance (median = 2 mm) 2 milliseconds after the time of max

gape, while maintaining a ram speed of ~2 body length per second (median = 77 mm

s^{-1}). With the exception of maximal jaw protrusion distance, all phenotypic traits were

skewed and significantly differed from normal (Shapiro-Wilk's p<0.05;

Supplementary Table S1). Several significant correlations were observed between

traits (Spearman rho r > |0.2|; Table 1), including a correlation between maximal gape

diameter and max jaw protrusion (r=0.59, p<0.001), between time to peak gape and

time to peak jaw protrusion (r=0.73, p<0.001), between time to peak gape and max

jaw protrusion (r=–0.4, p<0.001), and between time to peak jaw protrusion and max

jaw protrusion (r=–0.37, p<0.001).

The constructed performance landscape revealed a complex, non-monotonic,

performance surface (Figure 4; Figure S2) featuring ridges, peaks and flat projections.

The density of the contours was different for the different trait combinations,

indicating variable performance gradients for each set of traits. Of the six traits and

their paired interactions, all but two interactions were significant in the GAM model

(Table S2), indicating the importance of trait combinations in determining

performance. The range of strain rate thresholds estimated across the landscape is

ecologically plausible, ranging from 1-10 s^{-1} (Green et al., 2003; Kiorboe and Visser,

1999; Viitasalo et al., 1998; Visser, 2001), reinforcing the validity of this model in

predicting feeding performance based on phenotypic data.

Scenario I: the distribution of phenotypic traits is unrelated to the performance

landscape:

The median strain rate threshold of the observed population (2.16 s^{-1}) was

significantly lower by ~25% than the mean median of the 1000 "uncorrelated BM"

simulated populations (mean = 2.86; p < 0.001; 95% confidence intervals = 2.58 -

3.09 s^{-1}), and was outside the distribution of median performance in the simulated

populations (p<0.001; Figure 5A), refuting the hypothesis that the distribution of

phenotypic traits is unrelated to the performance landscape.

Scenario II: the population sits at the top of an 'adaptive peak'

While the performance of the observed population was higher than that of the

simulated population, the observed population was located off (local and global)

performance peaks. This was evident by the individual performance surface

calculated for the six traits (Fig 6). Five of the selection surfaces were significantly

larger than zero, indicating that a change equivalent to 0.1 sd of the trait value would

significantly increase mean performance (decrease threshold strain rate). TTPJP was

the only trait for which individual performance surface was not significantly different

than 0 (Fig 6D). The RSM analysis identified a local performance peak at peak gape

diameter of 5.1 mm, TTPG of 0.019 s, jaw protrusion distance of 6.6 mm, TTPJP of

0.007s (TTPG -0.012), ram speed of 61.8 mm/s and a timing differential of 0.012s.

None of the individuals in the observed population had such a combination, however

they are within the range observed for similar-sized fish.

Scenario III: the population evolves toward the 'adaptive peak', but has not

reached there

We calculated the angle in the 6-dimentional space between the eigenvector of

PC1 of the observed data (explaining ~35% of the total variance), and the vector of

the steepest descents, which depicts the direction of change, on the landscape, that

will result in maximal performance increase. Contrary to the "performance ridge"

hypothesis, that angle was 103.7°, indicating that the major axis of trait variation is

tangent to the direction to the peak. A similar result (an angle of 84.5°) was obtained

when we calculated the angle between the vector of steepest descents and the vector

sum of the coeficients for PC1-2.

Are trait correlations responsible for the off-peak location of the population?

Trait correlations can limit the distribution of traits in the trait space and therefor

constrain performance, providing a possible mechanism for the off-peak location of

our population. However, in contrast to this expectation, observed trait correlations

contributed to feeding performance. Simulating populations with the observed trait

correlations resulted in a higher performance (lower strain rate threshold) compared to

the "uncorrelated BM" populations. However, the median strain rate threshold of the

"correlated BM" populations was still significantly higher by ~15% than the observed

population (mean 2.55 s^{-1}; 95% confidence intervals for = 2.30 - 2.81 s^{-1}; P=0.005;

Figure 5B). The positive effect of trait correlations was consistent for all 6 traits;

removal of trait correlations for each of the traits (by randomizing the values of one

trait with respect to the others; see methods) produced similar distributions and

significantly reduced performance compared to the observed combinations (Figure 7).

Thus, the observed trait correlations are associated with higher feeding performance

than the non-correlated case, but it is unlikely that they result from trait distribution

along a performance ridge.

Discussion

In this study, we refine the "performance landscape" framework suggested by

Arnold (Arnold, 2003) by mapping the phenotypic distribution of performance-

determining traits to a landscape derived from a theoretical model of a fitness-

determining performance. Our approach allows a robust interpretation of the

performance landscape beyond the ranges observed in the sampled population, and

curtails "edge effects" that originally resulted in higher inaccuracy of the landscape

near the population edge. Mapping the observed population on the landscape

indicated that the population was located on the "upper slopes" of an adaptive peak,

but not on the peak itself. Nevertheless, trait correlations did not seem to limit

performance, as breaking trait correlations reduced performance. The major axes of

phenotypic variation were tangent to the direction to the local performance peak,

inconsistent with the scenario of trait distribution along a performance ridge (Fig 1).

We conclude that our approach can help uncover the relationships between

performance and trait evolution in complex functional systems.

The Chromis viridis population is located off a performance peak

The performance landscape for suction feeding reported here was complex,

rugged, and featured performance troughs and ridges (Figure 4). The landscape

clearly showed that different phenotypic traits have a different effect on performance

as indicated by the different slopes of the performance surfaces (Fig 4), and the

different selection gradients for the different traits (Fig 6). Moreover, these slopes

often change in relation to the values of the other traits. For example, for low values

of time to peak jaw protrusion (TTPJP < 0.02 s), the effect of time to peak gape

(TTPG) on performance is much stronger than at high values (TTPJP > 0.06 s; Fig 4).

This is probably responsible for some of the variation in individual performance

gradients (Fig 6). These trends demonstrate the complexity of the relationship

between form and function in this functional system, and the importance of an

integrative analysis to accurately model it.

Functional systems are often considered to be 'optimized' for performance, i.e.

the distribution of phenotypic traits within a species or a population is assumed to

correspond to performance peak (Bishop et al., 2008; van Leeuwen and Muller, 1984).

This notion is often supported by using the measured performance data to construct a

putative landscape (Arnegard et al., 2014; Arnold and Bennett, 1988). More often,

and especially in an inter-specific context, it is the distribution of phenotypic traits

themselves that is used to infer the existence and location of a "peak" (Collar et al.,

2009; Ingram and Mahler, 2013; Shoval et al., 2012). However, the relationship's

correspondence (or lack thereof) between the performance landscape and the

distribution of species traits are poorly demonstrated.

As pointed out by Arnold, (Arnold, 2003) it is problematic to infer the location of

performance peaks from such data. Therefore, the location of the population with

respect to performance peaks and ridges is difficult to resolve and is largely unclear.

Here we show that suction feeding performance of *Chromis viridis* is not optimal,

although it is significantly higher than that expected under a model of random motion

across the landscape. Such a distribution is often explained by trade-offs with other

performance axes. For example, Polly et al (Polly et al., 2016) argued that shell shape

in turtles is determined by the trade-off between shell strength and form drag.

However *C. viridis* is a zooplankton specialist, feeding mainly on copepods (Allen

and Randall, 1980), and it is highly likely that the performance landscape can be

reduced to one prey escape strategy (i.e. strain-sensitive prey). Unlike in mouth-

brooding, nest building or biting/scraping fish species (Ostlund-Nilsson and Nilsson,

2004; Smith and Wootton, 1995), the mouth in *C. viridis* is used mainly for suction

feeding and respiration. Therefore, it is unlikely that other fitness-determining

functions impose strong direct selection on the traits we measured.

Instead, we speculate that suction feeding performance is constrained by

mechanical limitations. Our model is purely functional, accounting for the effects of

strike kinematics on performance. However, it does not account for energetic, genetic,

developmental, or biomechanical constraints on the suction feeding mechanism

(Carroll and Wainwright, 2009; Van Wassenbergh et al., 2006; Wainwright et al.,

2007). Therefore, high-performance trait combinations identified by the model may

be unrealistic due to such constraints. For example, biomechanical principles dictate

that increases in gape size are likely to increase TTPG (Hulsey and Wainwright, 2002;

Oufiero et al., 2012) and increase the energetic cost of the strike (Carroll and

Wainwright, 2009). Such couplings are not accounted for by SIFF and are therefore

not implicit in the landscape. Similarly, faster time to peak jaw protrusion might

permit capturing more sensitive prey (Figure 4), however in reality it could be

disfavored because it will increase the energetic cost of the strike (Carroll and

Wainwright, 2009). Likewise, the relative timing of peak gape and peak jaw

protrusion could be a consequence of the biomechanical lever system (Westneat,

2005), bearing little effect on performance within the observed range of trait values.

Genetic linkages are also expected to constrain the distribution of traits in the

observed population (Lande and Arnold, 1983; Schluter, 1996). In our simulated

populations, we accounted for such correlations, which had a strong effect on the

phenotypic distribution of our populations and on performance.

Alternatively, it could be that that the location of the population off the

performance peak could result from a misalignment of the performance and fitness

peaks. For example, it could be that the performance peak allows the capture of highly

strain-sensitive prey. However, the energetic cost of feeding with a morphology that

sits upon that peak is high. If the abundance of such highly evasive copepods is very

low compared to other (less evasive) copepod species, the fitness peak (or peak

energetic gain) will be shifted towards the current location of the population.

However, we are unaware of community-wide data on the distribution of

hydrodynamic performance of copepods that would enable testing this idea.

The performance landscape is multidimensional, and it is therefore only possible

to visualize it in two dimensional projections (Fig 4, Fig S2). Qualitatively examining

the distribution of the observed population in these 2-D maps can be a useful tool in

generating hypotheses regarding the distribution of the phenotypic traits on the

landscape. For example, Figure 4D and 4F may support the intuition that some traits

follow "performance ridges" (Arnold, 2003; Arnold et al., 2001; Ghalambor et al.,

2003; Whibley et al., 2006), along which morphological variation can accumulate

without performance consequences. However, it is important to remember that the 2-

D visualization only represents the behavior of the displayed variables at median (or

another pre-defined) values of the other phenotypic traits, and cannot accurately

represent trait distributions in the multidimensional space (Blonder, 2016; Gavrilets,

1997). The intuition that traits follow "performance ridges" is not supported by the

individual selection gradients (Fig 6), which should have followed a hyperbolic shape

(i.e. lower selection differential close to the population mean) if traits were located on

a ridge. Additionally, the primary axis of variation was tangential to the path of

steepest performance ascent, which is inconsistent with the "performance

ridge" hypothesis.

Generality of the framework

Following the work of Hansen and Martins (Hansen and Martins, 1996), Arnold

et al. (Arnold et al., 2001) suggested that using performance landscapes can be helpful

in understanding micro and macro-evolutionary processes. However, such

applications of the performance landscape were rare. For example, Tseng (Tseng,

2013) used finite element analysis to map two traits (skull width-to-length and depth-

to-length ratios) to two functional properties, the mechanical advantage and strain

energy of the Carnivora skull. That study calculated the strength of theoretical skull

shapes to estimate the functional properties of traits that lay outside the distribution of

observed skull shapes, concluding that skull shapes evolved towards higher

performance, but also discovering high-performance regions that are not occupied by

extant species. Similarly, Polly et al (Polly et al., 2016) used finite element analysis

and hydrodynamic theory to predict the trade-off between shell strength and form

drag, and mapped the distribution of turtle species with respect to the resulting Pareto

front. By using the equations yielded from the landscape it is possible to relate

mechanistic processes to divergence in trait values across intra- and interspecific

levels: our results provide evidence for the evolution of feeding morphology on a

complex performance landscape as an important process that drives the present intra-

specific phenotypic distribution. It is possible to scale this method to higher

phylogenetic levels, and test the role of the performance landscape in shaping inter-

specific phenotypic distributions. Current methods for inferring adaptive regimes

from morphological data use inter-specific trait variance to deduce the locations of

"adaptive peaks" (Ingram and Mahler, 2013; Pennell and Harmon, 2013; Uyeda and

Harmon, 2014). These methods do not require *a priori* assumptions regarding the

manifestation of form to function, and the phylogenetic position, optimum, and

relative "heights" of the adaptive peaks are inferred from the data. Using theoretical

performance landscapes to detect adaptive peaks can be advantageous over such

methods because (1) it can easily account for interactions between traits in

multivariate space, (2) the accuracy of the landscape is independent of the sample size

of the focal population, is unaffected by incomplete taxon sampling, and is uniform

across the morphospace and (3) can estimate performance ridges and more complex

nonlinear features of empirical landscapes, which cannot be generated by simple

regime-shifting models such as the Ornstein-Uhlenbeck model (Hansen and Martins,

1996). Indeed, field experiments now demonstrate that complex fitness landscapes

drive adaptive radiation in rapidly speciating fish groups (Arnegard et al., 2014;

Martin and Wainwright, 2013). It will be interesting to compare the predictions of

such methods with the topography of empirical performance landscapes.

Importantly, while our study focuses on suction feeding in fish, the framework

we present here is general and widely applicable across taxa and functional systems.

Over the last decades, biomechanical theory and computational methods were used to

mechanistically model many aspects of performance such as swimming (Fish and

Lauder, 2006; Liao, 2007; Sfakiotakis et al., 1999), running (Barasuol et al., 2013;

Kingma et al., 1996; Minetti, 1998), slithering (Hu et al., 2009), flying and gliding

(Paranjape et al., 2012; Wu, 2011) among numerous other examples. These models

can be used to generate performance landscapes and map the distribution of functional

traits on these landscapes in a variety of functional systems, and in many species.

Because performance is fundamentally connected to viability selection in the wild,

this framework now enables us to test how these complex functional systems shape

evolutionary trajectories across species.

The performance landscape can be a useful tool to generate predictions regarding

the functional consequences of shifting trait means in response to ecological (e.g.

predation or competition) and evolutionary (e.g. co-evolution) processes (Arnold,

2003; Arnold et al., 2001; Ghalambor et al., 2003). For example, the performance

landscape can predict how removing large individuals with larger gape (e.g. through

predation or fishing) will change the distribution of performance in the population.

Likewise, performance landscapes can be used to predict the morphological response

to evolutionary changes in prey escape capabilities such as increased sensitivity or the

ability to accelerate and swim faster. However, the complex nature of the landscape

warrants caution when inferring selection, because selection may operate differently

on the same traits in different areas of morphospace, depending on the values of other

traits of the system. For example, two individuals that have the same trait value for

maximal jaw protrusion can be subjected to different selection regimes, depending on

the values of their five other traits. The selective force operating on each point in the

multi-dimensional space can be calculated as the sum (for all traits) of the derivatives

of performance with respect to changing trait values. It also follows that simple

mathematical functions (Arnold et al., 2001) fail to capture the complex nonlinear

features of the landscape (Schluter, 1996).

References

Allen, G. R. and Randall, J. E. (1980). A review of the damselfisches (Teleostei:

Pomacentridae) of the Red Sea. *Isr. J. Zool.* **29**, 1–98.

Arnegard, M. E., McGee, M. D., Matthews, B., Marchinko, K. B., Conte, G. L.,

Kabir, S., Bedford, N., Bergek, S., Chan, Y. F., Jones, F. C., et al. (2014).

Genetics of ecological divergence during speciation. *Nature* **511**, 1–17.

Arnold, S. J. (1983). Morphology, performance and fitness. *Integr. Comp. Biol.* **23**,

347–361.

Arnold, S. J. (2003). Performance surfaces and adaptive landscapes. *Integr. Comp.*

Biol. **43**, 367–375.

Arnold, S. J. and Bennett, A. F. (1988). Behavioural variation in natural

populations. V. Morphological correlates of locomotion in the garter snake

(Thamnophis radix). *Biol. J. Linn. Soc.* **34**, 175–190.

Arnold, S. J., Pfrender, M. E. and G., J. A. (2001). The Apaptive Landscape as a

conceptual bridge between mico- and macroevolution. *Genetica* **112–113**,

9–32.

Barasuol, V., Buchli, J., Semini, C., Frigerio, M., De Pieri, E. R. and Caldwell, D.

G. (2013). A reactive controller framework for quadrupedal locomotion on

challenging terrain. In *Proceedings - IEEE International Conference on*

Robotics and Automation, pp. 2554–2561. IEEE.

Bishop, K. L., Wainwright, P. C. and Holzman, R. (2008). Anterior-to-posterior

wave of buccal expansion in suction feeding fishes is critical for optimizing

fluid flow velocity profile. *J. R. Soc. Interface* **5**, 1309–16.

Blonder, B. (2016). Do Hypervolumes Have Holes? *Am. Nat.* **187**, E93-105.

Brodie, E. D., Moore, A. J. and Janzen, F. J. (1995). Visualizing and quantifying

natural selection. *Trends Ecol. Evol.* **10**, 313–318.

Buskey, E. J. and Hartline, D. K. (2003). High-speed video analysis of the escpae

response of the copepod Acartia tonsa to shadows. *Biol Bull* **204**, 28–37.

Buskey, E. J., Lenz, P. H. and Hartline, D. K. (2002). Escape behavior of

planktonic copepods in response to hydrodinamic disturbances: high speed

video analysis. *Mar. Ecol. Prog. Ser.* **235**, 135–146.

Carassou, L., Kulbicki, M., Nicola, T. J. R. and Polunin, N. V. C. (2008).

Assessment of fish trophic status and relationships by stable isotope data in

the coral reef lagoon of New Caledonia, southwest Pacific. *Aquat. Living*

Resour. **21**, 1–12.

Carroll, A. M. and Wainwright, P. C. (2009). Energetic limitations on suction

feeding performance in centrarchid fishes. *J. Exp. Biol.* **212**, 3241–3251.

Collar, D. C., O'Meara, B. C., Wainwright, P. C. and Near, T. J. (2009). Piscivory

limits diversification of feeding morphology in centrarchid fishes. *Evolution*

(N. Y). **63**, 1557–1573.

Coughlin, D. J. and Strickler, J. R. (1990). Zooplankton capture by a coral reef

fish: an adaptive response to evasive prey. *Environ. Biol. Fishes* **29**, 35–42.

Day, S. W., Higham, T. E., Holzman, R. and Van Wassenbergh, S. (2015).

Morphology, Kinematics, and Dynamics: The Mechanics of Suction Feeding

in Fishes. *Integr. Comp. Biol.* **55**, 21–35.

Fear, K. K. and Price, T. (1998). The Adaptive Surface in Ecology. *Oikos* **82**, 440.

Fields, D. M. and Yen, J. (1997). The escape behavior of marine copepods in

response to a quantifiable fluid mechanical disturbance. *J. Plankton Res.* **19**,

1289–1304.

Fish, F. and Lauder, G. V. (2006). Passive and active flow control by swimming

fishes and mammals. *Annu. Rev. Fluid Mech.* **38**, 193–224.

Gavrilets, S. (1997). Evolution and speciation on holey adaptive landscapes.

Trends Ecol. Evol. **12**, 307–312.

Gemmell, B. J., Adhikari, D., Longmire, E. K., Gemmell, B. J., Adhikari, D. and

Longmire, E. K. (2014). Volumetric quantification of fluid flow reveals fish '

s use of hydrodynamic stealth to capture evasive prey Volumetric

quantification of fluid flow reveals fish ' s use of hydrodynamic stealth to

capture evasive prey. *J. R. Soc. Interface* **11**, 20130880.

Ghalambor, C. K., Walker, J. A. and Reznick, D. N. (2003). Multi-trait Selection,

Adaptation, and Constraints on the Evolution of Burst Swimming

Performance. *Integr. Comp. Biol.* **43**, 431–438.

Goldshmid, R., Holzman, R., Weihs, D. and Genin, A. (2004). Aeration of corals

by sleep-swimming fish. *Limnol. Oceanogr.* **49**, 1832–1839.

Green, S., Visser, A. W., Titelman, J. and Kiorboe, T. (2003). Escape responses

of copepod nauplii in the flow field of the blue mussel, Mytilus edulis. *Mar.*

Biol. **142**, 727–733.

Hansen, T. F. (1997). Stabilizing selection and the comparative analysis of

adaptation: Evolution. *Evolution (N. Y).* **51**, 1341–1351.

Hansen, T. F. and Martins, E. P. (1996). Translating between microevolutionary

process and macroevolutionary patterns: the correlation structure of

interspecific data. *Evolution (N. Y).* **50**, 1404–1417.

Harmon, L. J., Weir, J. T., Brock, C. D., Glor, R. E. and Challenger, W. (2008).

GEIGER: Investigating evolutionary radiations. *Bioinformatics* **24**, 129–131.

Hedrick, T. L. (2008). Software techniques for two- and three-dimensional

kinematic measurements of biological and biomimetic systems. *Bioinspir.*

Biomim. **3**, 34001.

Heuch, P. A., Doall, M. H. and Yen, J. (2007). Water flow around a fish mimic

attracts a parasitic and deters a planktonic copepod. *J. Plankton Res.* **29**, i3–

i16.

Higham, T. E., Day, S. W. and Wainwright, P. C. (2006). Multidimensional

analysis of suction feeding performance in fishes: fluid speed, acceleration,

strike accuracy and the ingested volume of water. *J. Exp. Biol.* **209**, 2713–

2725.

Holzman, R. and Wainwright, P. C. (2009). How to surprise a copepod: Strike

kinematics reduce hydrodynamic disturbance and increase stealth of

suction-feeding fish. *Limnol. Oceanogr.* **54**, 2201–2212.

Holzman, R., Day, S. W. and Wainwright, P. C. (2007). Timing is everything:

coordination of strike kinematics affects the force exerted by suction feeding

fish on attached prey. *J. Exp. Biol.* **210**, 3328–3336.

Holzman, R., Day, S. W., Mehta, R. S. and Wainwright, P. C. (2008a). Jaw

protrusion enhances forces exerted on prey by suction feeding fishes. *J. R.*

Soc. Interface **5**, 1445–1457.

Holzman, R., Collar, D. C., Day, S. W., Bishop, K. L. and Wainwright, P. C.

(2008b). Scaling of suction-induced flows in bluegill: morphological and

kinematic predictors for the ontogeny of feeding performance. *J. Exp. Biol.*

211, 2658–68.

Holzman, R., Collar, D. C., Mehta, R. S. and Wainwright, P. C. (2012). An

integrative modeling approach to elucidate suction-feeding performance. *J.*

Exp. Biol. **215**, 1–13.

Hu, D. L., Nirody, J., Scott, T. and Shelley, M. J. (2009). The mechanics of

slithering locomotion. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 10081–5.

Hulsey, C. D. and Wainwright, P. C. (2002). Projecting mechanics into

morphospace: disparity in the feeding system of labrid fishes. *Proc. R. Soc. B*

Biol. Sci. **269**, 317–326.

Ingram, T. and Mahler, D. L. (2013). SURFACE: detecting convergent evolution

from comparative data by fitting Ornstein-Uhlenbeck models with stepwise

Akaike Information Criterion. *Methods Ecol. Evol.* **4**, 416–425.

Kingma, I., de Looze, M. P., Toussaint, H. M., Klijnsma, H. G. and Bruijnen, T.

B. M. (1996). Validation of a full body 3-D dynamic linked segment model.

Hum. Mov. Sci. **15**, 833–860.

Kiorboe, T. and Visser, A. W. (1999). Predator and prey preception in copepods

due to hydromechanical signals. *Mar. Ecol. Prog. Ser.* **179**, 81–95.

Lande, R. and Arnold, S. (1983). The measurement of selection on correlated

characters. *Evolution (N. Y).* **37**, 1210–1226.

Lenth, R. V. (2009). Response-Surface Methods in R, using rsm. *J. Stat. Softw.* **32**,

1–17.

Liao, J. C. (2007). A review of fish swimming mechanics and behaviour in altered

flows. *Philos. Trans. R. Soc. B Biol. Sci.* **362**, 1973–1993.

Longo, S. J., McGee, M. D., Oufiero, C. E., Waltzek, T. B. and Wainwright, P. C.

(2016). Body ram, not suction, is the primary axis of suction-feeding

diversity in spiny-rayed fishes. *J. Exp. Biol.* **219**, 119–128.

Martin, C. H. and Wainwright, P. C. (2013). Multiple fitness peaks on the

adaptive landscape drive adaptive radiation in the wild. *Science* **339**, 208–

11.

Minetti, A. E. (1998). The biomechanics of skipping gaits: a third locomotion

paradigm? *Proc. R. Soc. B Biol. Sci.* **265**, 1227–1233.

Ostlund-Nilsson, S. and Nilsson, G. E. (2004). Breathing with a mouth full of eggs: respiratory consequences of mouthbrooding in cardinalfish. *Proceedings. Biol. Sci.* **271**, 1015–22.

Oufiero, C. E., Holzman, R. A., Young, F. A. and Wainwright, P. C. (2012). New insights from serranid fishes on the role of trade-offs in suction-feeding diversification. *J. Exp. Biol.* **215**, 3845–3855.

Paranjape, A. A., Dorothy, M. R., Chung, S. J. and Lee, K. D. (2012). A flight mechanics-centric review of bird-scale flapping flight. *Int. J. Aeronaut. Sp. Sci.* **13**, 267–282.

Pennell, M. W. and Harmon, L. J. (2013). An integrative view of phylogenetic comparative methods: connections to population genetics, community ecology, and paleobiology. *Ann. N. Y. Acad. Sci.* **1289**, 90–105.

Polly, P. D., Stayton, C. T., Dumont, E. R., Pierce, S. E., Rayfield, E. J. and Angielczyk, K. D. (2016). Combining geometric morphometrics and finite element analysis with evolutionary modeling: towards a synthesis. *J. Vertebr. Paleontol.* **36**, e1111225.

Schluter, D. (1996). Adaptive Radiation Along Genetic Lines of Least Resistance. *Evolution (N. Y).* **50**, 1766–1774.

Sfakiotakis, M., Lane, D. M. and Davies, J. B. C. (1999). Review of fish swimming modes for aquatic locomotion. *IEEE J. Ocean. Eng.* **24**, 237–252.

Shoval, O., Sheftel, H., Shinar, G., Hart, Y., Ramote, O., Mayo, A., Dekel, E., Kavanagh, K. and Alon, U. (2012). Evolutionary trade-offs, Pareto optimality, and the geometry of phenotype space. *Science* **336**, 1157–60.

Simpson, G. (1944). *Tempo and mode in evolution.*

Simpson, G. (1955). Major features of evolution.

Smith, C. and Wootton, R. J. (1995). The costs of parental care in teleost fishes.

Rev. Fish Biol. Fish. **5**, 7–22.

Staab, K. L., Holzman, R., Hernandez, L. P. and Wainwright, P. C. (2012).

Independently evolved upper jaw protrusion mechanisms show convergent

hydrodynamic function in teleost fishes. *J. Exp. Biol.* **215**, 1456–1463.

Stewart, W. J., Cardenas, G. S. and Mchenry, M. J. (2013). Zebrafish larvae

evade predators by sensing water flow. *J. Exp. Biol.* **216**, 388–398.

Tseng, Z. J. (2013). Testing Adaptive Hypotheses of Convergence with Functional

Landscapes: A Case Study of Bone-Cracking Hypercarnivores. *PLoS One* **8**,

e65305.

Uyeda, J. C. and Harmon, L. J. (2014). A Novel Bayesian Method for Inferring

and Interpreting the Dynamics of Adaptive Landscapes from Phylogenetic

Comparative Data. *Syst. Biol.* **63**, 902–918.

van Leeuwen, J. L. and Muller, M. (1984). Optimum sucking techniques for

predatory fish. *Trans. Zool. Soc. London* **37**, 137–169.

Van Wassenbergh, S., Aerts, P. and Herrel, A. (2006). Hydrodynamic

modelling of aquatic suction performance and intra-oral pressures:

limitations for comparative studies. *J. R. Soc. Interface* **3**, 507–14.

Viitasalo, M., Kiorboe, T., Flinkman, J., Pedersen, L. W. and Visser, A. W.

(1998). Predation vulnerability of planktonic copepods: Consequences of

predator foraging strategies and prey sensory abilities. *Mar. Ecol. Prog. Ser.*

175, 129–142.

Visser, A. W. (2001). Hydromechanical signals in the plankton. *Mar. Ecol. Prog.*

Ser. **222**, 1–24.

Wainwright, P. C. and Day, S. W. (2007). The forces exerted by aquatic suction

feeders on their prey. *J. R. Soc. Interface* **4**, 553–60.

Wainwright, P., Carroll, A. M., Collar, D. C., Day, S. W., Higham, T. E. and

Holzman, R. A. (2007). Suction feeding mechanics, performance, and

diversity in fishes. *Integr. Comp. Biol.* **47**, 96–106.

Wainwright, P. C., McGee, M. D., Longo, S. J. and Patricia Hernandez, L.

(2015). Origins, Innovations, and Diversification of Suction Feeding in

Vertebrates. *Integr. Comp. Biol.* **55**, 134–145.

Westneat, M. W. (2005). Skull Biomechanics and Suction Feeding in Fishes. In

Fish Physiology (ed. Lauder, G. V) and Shadwick, R. E.), pp. 29–75. San Diego:

Elsevier Academic Press.

Whibley, A. C., Langlade, N. B., Andalo, C., Hanna, A. I., Bangham, A., Thébaud,

C. and Coen, E. (2006). Evolutionary paths underlying flower color

variation in Antirrhinum. *Science* **313**, 963–6.

Wood, S. N., Shaw, S. and Goude, Y. (2015). Generalized additive models for

large data sets. *J. R. Stat. Soc. Ser. C (Applied Stat.* **64**, 139–155.

Wright, S. (1932). *The roles of mutation, inbreeding, crossbreeding, and selection*

in evolution.

Wu, T. Y. (2011). Fish Swimming and Bird/Insect Flight. *Annu. Rev. Fluid Mech.*

43, 25–58.

Yen, J., Lenz, P. H., Gassie, D. V. and Hartline, D. K. (1992). Mechanoperception

in marine copepods: electrophysiological studies on the first antennae. *J.*

Plankton Res. **14**, 495–512.

Table 1: The correlation structure between phenotypic and kinematic traits. With
the exception of maximal jaw protrusion distance, all phenotypic traits were skewed
and significantly differed from normal (Shapiro-Wilk's $p<0.05$; Supplementary Table
S1). Therefore, we used Spearman rho for correlation between the traits in the
observed population.

Trait	Max gape	Time to peak gape	Max jaw protrusion	Time to peak jaw protrusion	Ram speed
Max gape					
Time to peak gape	-0.2*				
Max jaw protrusion	0.59***	-0.4***			
Time to peak jaw protrusion	-0.1	0.73***	-0.37***		
Ram speed	0.12	0.04	0.19*	0.12	
Gape-protrusion time difference	-0.11	0.11	0.001	-0.23*	-0.02

Significance level indicated by: * - $p<0.05$, ** - $p<0.01$, *** - $p<0.001$. N/S - $p>0.05$.

8
9

Figures

Fig 1: Possible scenarios depicting the distribution of the phenotypic traits with

respect to topography of a two-dimensional performance landscape. Trait values for

each individual in a population ('observed') are depicted as black circles. X and Y

axes represent two performance-determining traits, red colors represent higher

performance. Under scenario I, selection for suction performance has a weak effect on

the traits that underlie performance, leading to a distribution of phenotypic traits that

is unrelated to the performance landscape. Under scenario (II), selection for suction

performance has a dominant effect on morphological evolution, and the population

sits at the top of a performance peak Lastly, under scenario III, the population could

be under stabilizing selection, with selection for suction performance countered by

trade-offs, genetic of mechanical constraints, or could lay on a performance ridge. In

the latter case, the major axis of phenotypic variation (dashed black line) should align

with the path of steepest ascent (white arrow), pointing toward the local peak.

Figure 2: the underwater filming system at the Coral Reef in Eilat (upper panel),

was set up to film the zooplanktivorous *chromis viridis* during natural foraging

(middle panel). The system captured stereoscopic views of the feeding fish, observed

with the two cameras (lower panels). Landmarks (red circles 1-3 in lower panels)

were digitized on the fish's body and head to estimate strike kinematics at 500 frames

per second.

Figure 3: using SIFF to estimate prey capture performance from field-observed

and simulated kinematics. Using an underwater high-speed 3D camera system (Fig 2),

we recorded body and mouth kinematics of *Chromis viridis* (A). From these, we

calculated a set of 6 functionally-relevant traits (red circles in B; values for the

presented strike are indicated in boxes to the left and bottom of main panel in A).

These traits, in addition to prey's traits (blue circles in B) were used as input variables

for SIFF, a hydrodynamic model of prey capture in fish. The model calculates the

displacement of the prey in space and time, based on the forces exerted on it by the

suction flows (generated by the fish) and the prey's propulsive force. For each set of

input parameters, the outcome (prey escape/capture) is deterministic. To estimate

performance, the model was ran iteratively for each set of input parameters while

increasing the strain rate threshold for prey escape (C). At a sufficiently high

threshold (causing the prey to escape later in time) the prey is captured, and that

threshold is recorded. This process can be carried with observed (filled points in C; a

threshold of 1.5 s^{-1}) or simulated kinematics, for example using a larger gape diameter

(empty circles in C; yielding a threshold of 1.25 s^{-1}). To reconstruct the performance

landscape, we used ~3300 trait sets, generated computationally by sampling in

random from range of observed trait values in the field population.

Figure 4: 2D projections of the multidimensional performance landscape,

generated using a hydrodynamic model of predator-prey interactions. The

multidimensional performance landscape was generated based on 3300 simulated

prey-acquisition strikes, featuring random trait combinations selected from the

observed trait range. Colors (and white contours) represent feeding performance,

estimated as strain rate threshold (s^{-1}) of the most sensitive prey that can be captured

by a predator using that trait combination. Red colors represent high performance

(low strain rate thresholds), yellow represents low performance (high strain rate

thresholds). Observed trait combinations (black circles, n=110) are overlaid on the

performance surface. Note that the 2D visualizations only represent the performance

of the featured variables at median values of the other four phenotypic traits, and do

not represent trait distribution in the multidimensional space. The plot features six of

the 15 possible 2D projections (See Fig S2 for all projections).

Figure 5: Distribution of performance (strain rate thresholds) in 1000 populations,

each with 110 simulated individuals, *versus* performance in the observed population

A) Median performance in the observed population (vertical red line) and its

distribution in populations generated by an uncorrelated Brownian motion process

("uncorrelated BM" populations; blue bars). B) Median performance in the observed

population (vertical red line) and its distribution in populations generated by a

correlated Brownian motion process ("correlated BM" populations; red bars).

Sensitivity threshold median of the observed population is outside the 95%

confidence interval for both simulated populations (p<0.001 for both; n=1000).

Fig 6: Individual performance surface (Arnold, 2003), as a function of trait values.

These selection surfaces represent the slope of the decrease in strain rate threshold

(increased performance) following shift of 0.1 standard deviations in trait value. The

direction of the change in each trait is indicated by the arrow in the top right corner of

each panel. The length of the arrow is proportional to one standard deviations of the

trait. Mean (±sd) individual performance surface (ISSP) is depicted for each panel.

Vertical gray line depicts the trait mean. The horizontal red line represents ISSP=0;

negative values on the y-axis indicate that performance decreased (strain rate

threshold increased).

Figure 7: the effect of trait correlations on performance. We broke the correlation

structure between the trait depicted in each panel and the other 5 traits and asked how

the performance of these new combinations was affected. Mean denotes the average

of median strain rate threshold for 1000 simulated populations; p value was calculated

by comparing the distribution of median strain rate thresholds of the simulated

populations with that of the observed one. Note that breaking the correlation for each

single trait with the other five had similar effects across traits, reducing performance

by 7.5-20%. Thus, the observed trait correlations are associated in higher feeding

performance than the non-correlated case.

1   Supplementary tables and figures

3   Table S1: Summary statistics for the six morphological traits in the observed
4   population

Trait	Median	Mean	SD	CV	Deviation from normal distribution
Max gape	2.60 mm	2.73 mm	0.56 mm	0.205	**
Time to peak gape	20 msec	24 msec	11 msec	0.463	***
Max jaw protrusion	2 mm	2.15 mm	0.70 mm	0.328	N/S
Time to peak jaw protrusion	23 msec	25 msec	12 msec	0.485	***
Ram speed	76.8 mm/sec	86 mm/sec	36 mm/sec	0.414	***
Gape-protrusion time difference	-2 msec	-2.1 msec	2.4 msec	1.106	*

5   Deviation from normal distribution was tested using Shapiro-Wilk's test. Significance
6   level indicated by: * - $p<0.05$, ** -$p<0.01$, *** - $p<0.001$. N/S - $p>0.05$.
7

1　Table S2: Contribution of morphological traits and their interactions to the GAM-
2　smoothed performance landscape.

Trait	Effective degrees of freedom	Deviance from null (Wald statistic)	Significance level
Max gape (gape_max)	5.10	211.625	***
Time to peak gape (TTPG)	3.64	60.4	***
Max jaw protrusion (JP_max)	5.82	153.039	***
Time to peak jaw protrusion (TTPJP)	6.72	74.413	***
Ram speed (ram_spd)	5.58	2596.523	***
Gape-protrusion time difference (time_diff_peak)	4.02	7.095	***
gape_max:TTPG	2.48	4.854	**
gape_max:JP_max	7.17	3.981	***
gape_max:TTPJP	10.88	6.721	***
gape_max:ram_spd	10.53	45.606	***
gape_max:time_diff_peak	5.68	2.235	*
TTPG:JP_max	0.00	0.481	N/S
TTPG:TTPJP	11.47	23.131	***
TTPG:ram_spd	7.39	18.293	***
TTPG:time_diff_peak	13.44	12.051	***
JP_max:TTPJP	10.74	12.511	***
JP_max:ram_spd	7.49	33.797	***
JP_max:time_diff_peak	3.27	1.573	N/S
TTPJP:ram_spd	12.22	14.315	***
TTPJP:time_diff_peak	11.58	34.186	***
ram_spd:time_diff_peak	9.20	2.318	**

3　Wald statistics and p-values were calculated according to Wood (2013). Significance
4　level indicated by: * - $p<0.05$, ** -$p<0.01$, *** - $p<0.001$. N/S - $p>0.05$.
5

2   Fig S1: The distribution of 3291 phenotypic combinations in a 2D data projection

3   (time to peak gape and time to peak jaw protrusion). Trait combinations (blue circles)

4   were generated by randomly sampling from the range of phenotypic traits observed in

5   the population, and used to generate the landscape (black contours and red-yellow

6   colors).

Figure S2: 2D projections of the multidimensional performance landscape,

generated using a hydrodynamic model of predator-prey interactions. The plot

features all possible 2D projections available for the six traits measured. The

multidimensional performance landscape was generated based on 3300 simulated

prey-acquisition strikes, featuring random trait combinations selected from the

observed trait range. Colors (and white contours) represent feeding performance,

estimated as strain rate threshold (s^{-1}) of the most sensitive prey that can be captured

by a predator using that trait combination. Red colors represent high performance

(low strain rate thresholds), yellow represents low performance (high strain rate

thresholds). Observed trait combinations (black circles, n=110) are overlaid on the

performance surface. Note that the 2D visualizations only represent the performance

of the featured variables at median values of the other four phenotypic traits, and do

not represent trait distribution in the multidimensional space.

1   References for supplementary methods

2   Wood, S.N. (2013) On p-values for smooth components of an extended

3 generalized additive model. Biometrika 100:221-228