Modeling Spatio-temporal Dynamics of Chromatin Marks

Petko Fiziev[1,2,3], Jason Ernst[1,2,3,4,5,6,*]

[1] Bioinformatics Interdepartmental Program, University of California, Los Angeles

[2] Departments of Biological Chemistry, University of California, Los Angeles

[3] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at UCLA

[4] Computer Science Department, University of California, Los Angeles

[5] Jonsson Comprehensive Cancer Center, University of California, Los Angeles

[6] Molecular Biology Institute, University of California, Los Angeles, California, USA

* To whom correspondence should be addressed. Email: jason.ernst@ucla.edu

ABSTRACT

To model spatial changes of chromatin mark peaks over time we developed ChromTime, a computational method that identifies regions for which peaks either expand or contract significantly or hold steady between time points. We applied our method to chromatin marks in a range of biological systems. Predicted expanding and contracting peaks likely mark regulatory regions associated with transcription factor binding dynamics and gene expression changes. Spatial dynamics of peaks are informative about gene expression changes beyond localized signal changes. In proximity of gene starts, peaks preferentially expand in the same direction as transcription and contract in the opposite direction.

KEYWORDS: epigenomics, time course ChIP-seq, spatial dynamics, histone modifications

BACKGROUND

Genome-wide mapping of histone modifications (HMs) and related chromatin marks using chromatin

immunoprecipitation coupled with high throughput sequencing (ChIP-seq) has emerged as a powerful

approach to annotate genomes and study cell states[1–3]. Through the efforts of large consortia projects

such as the ENCODE[4], Roadmap Epigenomics[5] and BLUEPRINT[6] as well as individual labs[7–9]

multiple different chromatin marks have been mapped across more than a hundred different cell and

tissue types. These maps have yielded numerous insights into gene regulation and genetic and

epigenetic association with disease[10–14].

While many mapping efforts have largely focused on single or unrelated cell and tissue types[1, 4], a

growing number of biological processes have been studied with temporal epigenomic data using time

course ChIP-seq assays that map chromatin marks at consecutive stages during particular biological

processes. Such datasets have been generated for a wide range of biological settings including T cell

development[15], adipogenesis[16], hematopoiesis[17], macrophage differentiation[18], neural

differentiation[10], cardiac development[19, 20], somatic cell reprogramming[21–24], embryogenesis[25]

and many others[26–34]. The output of these experiments presents a unique opportunity to study the

spatio-temporal changes of epigenetic peaks and associated regulatory elements. However, almost all

computational methods designed or applied to epigenomic data have been developed based on single or

multiple unrelated samples. For example, continuous regions of enrichments of single marks are detected

by peak or domain calling methods[35–39]. In cases when multiple HMs are mapped in the same cell

type, methods such as ChromHMM[40] and Segway[41] can be used to produce genome-wide chromatin

state annotations. In addition, algorithms have been developed for pairwise comparisons of ChIP-seq

signal data by differential peak calling[42, 43].

In the context of time course ChIP-seq data, only a few methods have been proposed that consider

temporal dependencies between samples. One such method, TreeHMM[44], produces a chromatin state

genome annotation similar to ChromHMM and Segway, while taking into account a tree-like structure that

captures lineage relationships between the input cell types in order to potentially derive a more consistent

annotation across samples. Another method, GATE[27], produces a genome annotation based on

clustering fixed length genomic loci that can be modeled with the same switch from one chromatin state

to another over time.

One important limitation of methods for pairwise comparison of ChIP-seq data and time course modeling

of it is that they do not directly consider or model spatial changes in the genomic territory occupied by

chromatin marks over time. Spatial properties of genomic peaks continuously marked by HMs have

gained increasing attention as a potentially important characteristic of chromatin marks. For example,

long peaks of H3K27ac have been associated with active cell type specific locus control regions termed

super-enhancers or stretch enhancers in a number of cell types[45, 46]. Also, the length of H3K4me3

peaks has been associated with transcriptional elongation and consistency of cell identity genes[47]. In

the context of cancer, long H3K4me3 peaks have been linked to transcriptional elongation and enhancer

activity at tumor suppressor genes and have been observed to be significantly shortened in tumor

cells[48]. Long H3K4me3 domains have been implicated to mark loci involved in psychiatric disorders[49].

Expanded domains of H3K27me3 and H3K9me3 marks have been shown to be characteristic of

terminally differentiated cells compared to stem cells[50]. These studies suggest that length of epigenetic

peaks is a dynamic feature that can correlate with activity of putative functional elements regulating

specific genes. Computational methods that do not explicitly reason about the spatial changes of HMs

have significant limitations for studying the dynamics of these properties because they are unable to

detect territorial changes that might be associated with redistribution of signal or identify asymmetric

directional peak boundary movements.

In this work, we present ChromTime, a novel computational method for detection of expanding,

contracting and steady peaks, which can detect patterns of changes in the genomic territory occupied by

chromatin mark peaks from time course ChIP-seq data (**Fig 1A**). We applied ChromTime to a diverse set

of data from different developmental, differentiation and reprogramming time courses. Expansions and

contractions in general mark regulatory regions associated with changes in transcription factor binding

and gene expression. ChromTime enables studying the directionality of spatial dynamics of chromatin

mark peaks relative to other genomic features, which existing computational approaches do not directly

address. Our results show that the direction of expansions and contractions correlates with direction of

transcription near transcription start sites. ChromTime is a general method that can be used to analyze

time course ChIP-seq data from a wide range of biological systems to gain insights into the dynamics of

gene regulation.

RESULTS

Model for detecting expanding, contracting and steady peaks from temporal ChIP-seq data

We developed a computational method, ChromTime (https://github.com/ernstlab/ChromTime), designed

for systematic detection of expansions, contractions and steady peaks from time course ChIP-seq of

single chromatin marks (**Fig 1B**). ChromTime takes as input a set of genomic coordinates of aligned

sequencing reads from ChIP-seq and, optionally, control experiments over the time course. The method

consists of two stages – block finding and dynamics prediction. During the block finding stage,

ChromTime determines continuous genomic regions (blocks) that may contain ChIP-seq peaks

throughout the time course. To achieve this, ChromTime partitions the genome into fixed length bins and

counts the number of ChIP-seq and control reads that map to each bin at each time point. Nearby bins

that show significant enrichment are joined into continuous intervals, which subsequently are grouped into

blocks if they overlap across time points. As a result, large portions of the genome that are likely to

contain background noise at all time points are filtered out, so that peak boundary dynamics are

determined within a subset of the genome potentially enriched for the chromatin mark.

During the dynamics prediction stage, for each block ChromTime determines the most likely positions of

the peak boundaries at each time point and whether the peak expands, contracts or holds steady at each

boundary between consecutive time points. The method uses a probabilistic mixture model to partition the

signal within each block at each time point into background and peak components (**Fig 1C**) by reasoning

jointly about the data from all time points in the time course. The model assumes that central positions in

blocks have higher prior probability to be enriched for ChIP-seq reads on average and thus more likely to

associate with the peak component (see **Methods, Fig S1D**). The number of sequencing reads in bins

from each component at each time point is modelled with different negative binomial distributions that can

account for the local abundance of control reads. Furthermore, between any two consecutive time points

the boundaries of the peaks are assumed to follow one of three possible dynamics: steady, expand or

contract. For steady dynamics, the peak boundaries are enforced to have the same genomic position. For

expanding and contracting dynamics the number of genomic bins that the peak boundaries move

between the two time points is modelled with different negative binomial distributions which depend on

the pair of time points and the corresponding dynamic. Each dynamic is also assumed to have a prior

probability which captures its genome-wide frequency at each time point.

All model parameters are learned jointly from the whole time course. As a result, ChromTime can adapt to

different boundary movements, dynamics frequencies and noise levels across experiments and biological

systems. The estimated parameters are used to make a prediction for each block for the most likely

positions of the peak boundaries and the corresponding boundary dynamics that had generated the

signal within the block. The final output contains predicted peak boundaries annotated and colored by

their assigned dynamics, which can be used for downstream analysis with existing tools and visualized in

genome browsers (**Fig 2, S2,** https://github.com/ernstlab/ChromTime).

Reproducibility of ChromTime predictions and enrichment for TF binding and DNaseI
hypersensitivity sites

To investigate the reproducibility of ChromTime predictions, we applied ChromTime separately to two

biological replicate datasets for the H3K4me2 and the H3K(9/14)ac marks in T cell development in

mouse[15] and confirmed on average strong enrichment for the same ChromTime annotations co-

localizing across replicates (**Fig S3**). We then applied the method to data from pooled replicates for the

H3K4me2 and H3K(9,14)ac marks from the mouse T cell development study[15] and to data for the

H3K4me3 and H3K27ac marks from a study on stem cell reprogramming in human[21]. To investigate the

biological relevance of ChromTime predictions, we intersected their genomic positions with the dynamics

of orthogonal genomic annotations for transcription factor binding[4, 15] and DNaseI hypersensitivity

sites[5]. Predicted expanding and contracting peaks enriched for sites bound by important regulators in

each biological system as well as sites bound by generic TFs in a cell type specific manner (**Fig 3**).

Furthermore, predicted steady peaks associated with TF binding sites and DNaseI hypersensitive sites

that are shared between the first and the last time point in each time course, which mark potentially stable

regulatory elements.

Predicted spatial dynamics by ChromTime associate better with gene expression changes

compared to boundary position changes of peaks called from individual time points in isolation

To investigate whether ChromTime's approach for reasoning jointly about the whole time course

increases power to detect associations with gene expression compared to considering boundary

differences of peaks at consecutive time points called in isolation, we analyzed gene expression changes

of genes with TSSs overlapping ChromTime predictions in relation to posterior probabilities for

expansions and contractions compared to boundary differences of peaks called with ChromTime from

data from individual time points in isolation. We specifically investigated this in the context of H3K4me2

peaks in mouse T cell development[15] and for H3K4me3 peaks in stem cell reprogramming in

human[21] (**Fig 4**). In most cases, ranking block boundaries by their predicted ChromTime posterior

probabilities for expansions and contractions associated on average with larger gene expression changes

compared to ranking block boundaries directly based on the change in the genomic positions of the

boundaries of ChromTime peaks called at individual time points in isolation. These results also held when

using peaks from two different peak callers, MACS2[35] and SICER[37] applied on data from individual

time points (**Fig S4**).

Spatial dynamics contain information about gene expression changes in consecutive time points

not captured by corresponding pairwise HM signal changes

We next investigated whether there is additional information in ChromTime predictions beyond what can

be captured by pairwise ChIP-seq signal density changes or by differential peak calls. For this analysis,

we focused on H3K4me2 in mouse T cell development[15] and H3K4me3 in human stem cell

reprogramming[21]. At each pair of consecutive time points from the first to the last time points with

predicted peaks in a block, we computed the change in ChIP-seq signal density within the left most and

right most predicted peak boundaries in the block. We associated the signal density changes with gene

expression changes at the nearest transcription start site within 50kb of each block and computed the

average gene expression change as a function of the ChIP-seq signal density change within blocks (**Fig**

5). We found that locations with the same ChIP-seq signal density change can associate with significantly

different average gene expression changes of proximal genes depending on the predicted ChromTime

dynamics. Notably, bidirectional expansions, expansions occurring on both sides of a peak, associated

with greater average increase in gene expression than unidirectional expansions, those expansions

occurring on one side with steady on the other. These unidirectional expansions in turn associated with

greater expression change than steady regions, those regions with a steady call on both sides of a peak.

We observed a similar relationship for contractions and decrease of gene expression. These results were

replicated also after substituting ChIP-seq signal density changes with differential peak scores as

outputted by two different differential peak calling methods (SICER[37] and MACS2[35], **Fig S5**).

Therefore, ChromTime predictions can provide additional information about gene expression changes

beyond what is contained in the corresponding ChIP-seq signal density changes as measured directly or

by utilizing differential peak calling procedures.

Spatial dynamics are correlated between multiple histone marks

Previous studies have shown that the locations of different HMs can be correlated[1, 51]. In this context,

we tested whether multiple HMs can also exhibit jointly the same type of spatio-temporal dynamics. For

this purpose, we compared the genomic locations of predicted expansions, contractions and steady

peaks for different HMs within the same time course. We focused on two previously published time

courses – stem cell reprogramming in human[21] and adipogenesis in mouse[16], where multiple HMs

were mapped (**Fig 6**). In both datasets, we observed that expansions for H3K4me1, H3K4me2, H3K4me3

and H3K27ac co-localized preferentially and similarly for contractions and steady. In contrast, different

spatial dynamics for H3K36me3 and H3K27me3 tended to occupy distinct locations. These results

suggest that spatial dynamics of HMs are coordinated at least at a subset of genomic locations.

Direction of expansions and contractions is correlated with direction of transcription

ChromTime can predict unidirectional expansions and contractions, which enables analysis of

directionality of spatial dynamics of peaks, an aspect of chromatin regulation that has not been previously

systematically explored. To investigate this, we applied ChromTime on data from 11 previously published

studies from a variety of developmental, differentiation and reprogramming processes (**Table 1**) for nine

different HMs including narrow and broad marks and for Pol2. We observed that unidirectional

expansions and contractions are predicted in most cases on average to be the majority of all expansions

and contractions, respectively, at a given pair of consecutive time points (**Fig S6**). One hypothesis for the

prevalence of asymmetric boundary movements for the promoter associated chromatin marks is that the

direction of boundary movements is associated with the asymmetry of transcription initiation in promoter

regions. To test this hypothesis, for each dataset we compared the prevalence of each class of

unidirectional dynamics as a function of its distance to the nearest annotated transcription start site (TSS)

and the orientation of the corresponding gene (**Fig 7**). Consistent with our hypothesis, for H3K4me3,

H3K4me2, H3K(9/14)ac, H3K79me2, and for Pol2, we found that unidirectional expansions that expand

into the gene body (i.e. in the same direction as transcription) were substantially more frequently found in

proximity of TSSs compared to unidirectional expansions in the opposite direction. Moreover, this

difference was not observed for expansions distal from TSSs. Similarly, in most cases for these data

unidirectional contractions that contract towards the TSS of the nearest gene (i.e. in the opposite direction

of transcription) were substantially more frequently found compared to unidirectional contraction in the

opposite direction in proximity of TSSs, whereas their frequencies at distal sites showed much smaller

differences. HMs H3K27ac, H3K4me1 and H3K27me3 exhibited the same trend, but to a smaller degree.

DISCUSSION

In this work, we presented ChromTime, a novel computational method for systematic detection of

expanding, contracting and steady peaks of chromatin marks from time course ChIP-seq data.

ChromTime employs a probabilistic graphical model that directly models changes in the genomic territory

occupied by single chromatin marks over time. This approach allowed us to directly encode our modeling

assumptions about dependencies between variables in an interpretable and extendable framework.

We applied ChromTime on datasets for broad and narrow HMs and for Pol2 from a variety of

developmental, differentiation and reprogramming courses. Our results show that the method can identify

sets of expanding and contracting peaks that are biologically relevant to the corresponding systems. In

particular, expansions and contractions associate with up- and down-regulation of gene expression and

differential transcription factor binding, supporting the biological relevance of ChromTime predictions.

ChromTime gains power by both reasoning jointly about all time points in a time course and by explicitly

modeling the peak boundary movements. Supporting this we demonstrated that territorial changes

identified by ChromTime had better agreement with gene expression changes compared to considering

directly the boundary change of peaks called on data from individual time points in isolation. Additionally,

we found that expanding and contracting peaks associated on average with greater change in gene

expression compared to peaks with steady boundaries even after controlling for ChIP-seq signal density

changes. Some of the power that ChromTime gains from considering spatial information might be

explained by its ability to differentiate territorial expansions or contractions which can reflect changes in

the number sites of TF binding in close vicinity from changes in ChIP-seq signal within steady peak

boundaries. Changes in ChIP-seq signal without territorial expansions or contractions might reflect a

change in proportion of cells with the chromatin mark without large changes in activity in any one cell.

Additional power can come from the temporal and spatial information that allows the model to effectively

smooth over noise in the data enabling more biologically relevant inferences.

ChromTime enables novel analysis of directionality of spatial epigenetic dynamics. In this context, we

found that asymmetric unidirectional expansions and contractions for several marks correlate strongly

with direction of transcription in promoter proximal regions, which suggests that spatial dynamics at such

locations may be related to actions of the transcriptional machinery. One possible explanation for the

observed correlation between the direction of spatial dynamics of at least some HMs and transcription

can be provided in part by previous studies that have shown that the Pol2 elongation machinery can

recruit H3K4-methyltransferases such as members of the SET[52] and MLL[53] families at the promoters

of genes. Our findings are consistent with such models where the Pol2 complex itself may be facilitating

the attachment and removal of these marks[54].

One current limitation of the ChromTime method is that while the runtime of ChromTime still scales

linearly with the number of time points, T, the number of observed combinations of dynamics can scale

exponentially with T. This exponential growth can complicate downstream analyses that directly consider

each combination of dynamics, as there will be 3^{T-1} possible sequences of dynamics at each side of a

peak. Extensions of the ChromTime model could model the large number of combinations as being

instances of a smaller number of more distinct dynamic patterns.

CONCLUSIONS

The availability of time-series HM data provides an opportunity to understand chromatin dynamics in

many biological systems. To better leverage the information in these experiments we presented

ChromTime, a method to detect expansions, contractions and steady peaks of single chromatin marks

between time points from time course ChIP-seq data. Our method gains power by both reasoning about

data from all time points in the time course and by explicitly modeling movements of peak boundaries.

We showed that ChromTime predictions are reproducible across biological replicates and associate with

relevant genomic features such as changes in gene expression and transcription factor binding. We

demonstrated that territorial changes of peaks can contain additional information beyond ChIP-seq signal

changes with respect to gene expression of proximal genes. ChromTime allows for novel analysis of

directionality of spatial dynamics of chromatin marks. In this context, we showed for multiple chromatin

marks that the direction of asymmetric expansions and contractions of peaks correlates strongly with

direction of transcription in proximity of transcription start sites. ChromTime is generally applicable to

modeling time courses of chromatin marks, and thus should be a useful tool to gaining insights into

dynamics of epigenetic gene regulation in a range of biological systems.

METHODS

Overview of the ChromTime method

ChromTime takes as input a set of files in BED format with genomic coordinates of aligned sequencing

reads from ChIP-seq experiments for a single mark over the time course and, optionally, from a set of

control experiments. ChromTime consists of two stages (**Fig 1B-C**):

1) Detecting genomic intervals (blocks) potentially containing regions of ChIP-seq signal enrichment

(peaks)

2) Learning a probabilistic mixture model for boundary dynamics of peaks within blocks throughout

the time course and computing the most likely spatial dynamic and peak boundaries for each

block throughout the whole time course

Detecting genomic blocks containing regions of ChIP-seq signal enrichment

The aim of this stage is to determine approximately the genomic coordinates of regions with potential

ChIP-seq peaks at any time point in the time course. The signal within these blocks will be used as input

to build the mixture model in the next stage of ChromTime. ChromTime supports analysis of both narrow

marks and broad marks in two different modes. The method partitions the genome into non-overlapping

bins of predefined length, BIN_SIZE (200 bp in narrow mode, 500 bp in broad mode by default) and

counts for each bin and time point the number of sequencing reads whose alignment starting positions

after shifting by a predefined number of bases (100 bp in the direction of alignment by default) are within

its boundaries. Next, each bin at each time point is tested for enrichment based on a Poisson background

distribution at a predefined false discovery rate (5% by default). The expected number of reads for a bin

at position p and time point t, $\lambda_{t,p}$, in the Poisson test is computed conservatively as the maximum of:

1) If control reads are provided: for each of windows of size w=1,000bp, 5,000bp and 20,000bp the

average number of control reads in the window centered at the current bin, normalized by the

ratio of total reads in ChIP-seq and control experiments, that is:

$$\lambda_{t,p,w} = \frac{\# \,[\text{Total Foreground Reads}]}{\# \,[\text{Total Control Reads}]} \frac{\text{BIN\_SIZE}}{w} \text{Ctrl}_{t,p,w}$$

where $\text{Ctrl}_{t,p,w}$ is the total number of control reads in each of window of size w around the bin at

position p at time point t.

2) The average number of foreground ChIP-seq reads per genomic bin;

3) 1 read

Testing multiple different windows sizes for the background is a strategy we adopted from the MACS2

peak caller[35].

Within each time point, consecutive bins that are significantly enriched are merged into continuous

intervals. The intervals are further extended in both directions to include continuous stretches of bins

where each bin is significant based on a Poisson background distribution at a weaker P-value threshold

(0.15 by default). Extended intervals within a predefined number of non-significant bins, MAX_GAP (3

bins by default), are further joined together. This joining strategy has been previously implemented by

other peak callers for single datasets such as SICER[37]. Next, overlapping intervals across time points

are grouped into blocks. To capture more of the potential background signal and to increase the likelihood

that central bins within blocks contain higher ChIP-seq signal, the start and end positions of each block

are extended additionally by a predefined number of bins, BLOCK_EXTEND, (5 by default) upstream

from the left-most coordinate and downstream from the right-most coordinate of the intervals in the block,

respectively, or up to the middle point between the current block and its adjacent blocks if they are within

BLOCK_EXTEND bins apart. Restricting BLOCK_EXTEND to a relatively limited number of bins helps in

keeping the running time of the method within reasonable bounds.

In narrow mode, blocks that contain multiple intervals at the same time point separated by gaps of non-

significant bins longer than MAX_GAP are split into sub-blocks at each gap between those intervals. In

particular, all gaps within a block are intersected across the time points that have gaps. For each gap

intersection, the block is split at the position with the lowest average ChIP-seq signal across all time

points. In broad mode, no such splitting is performed in order to avoid excessive peak fragmentation.

Probabilistic mixture model for boundary dynamics of peaks within blocks across the time course

The ChIP-seq signal within the blocks is used as input to build a probabilistic mixture model for the

boundary dynamics of the peaks within blocks. One core assumption of the model is that each block

contains zero or one peak at each time point. This implies that at each time point, the bins within a block

can be partitioned into three continuous intervals: left-flanking background, foreground peak, and right-

flanking background. Let $O_{t,p}$ denote the number of observed ChIP-seq reads in a bin at position p at time

point t, and $V_{t,p}$ denote the label of that bin (one of Peak or Background). The conditional distribution of

$O_{t,p}$ conditioned on $V_{t,p}$ is modeled with different negative binomial distributions depending on the value of

$V_{t,p}$:

$$P\left(O_{t,p} = k \middle| V_{t,p} = \mathrm{Peak}\right) = \mathrm{NB}\left(k; \mu_{\mathrm{Peak},t,p}, \delta_t\right) = \frac{\Gamma(k + \delta_t)}{k!\,\Gamma(\delta_t)} \left(\frac{\delta_t}{\mu_{\mathrm{Peak},t,p} + \delta_t}\right)^{\delta_t} \left(\frac{\mu_{\mathrm{Peak},t,p}}{\mu_{\mathrm{Peak},t,p} + \delta_t}\right)^{k}$$

357   and

$$P\big(O_{t,p} = k \big| V_{t,p} = \text{Background}\big) = \text{NB}\big(k; \mu_{\text{Background},t,p}, \delta_t\big)$$

$$= \frac{\Gamma(k + \delta_t)}{k!\,\Gamma(\delta_t)}\left(\frac{\delta_t}{\mu_{\text{Background},t,p} + \delta_t}\right)^{\delta_t}\left(\frac{\mu_{\text{Background},t,p}}{\mu_{\text{Background},t,p} + \delta_t}\right)^{k}$$

360   Similarly to negative binomial regression models[55], ChromTime models the mean of each component

361   through the log link as a linear combination of a two-dimensional vector of covariates, $X_{t,p} = (1, \log \lambda_{t,p})$,

362   which includes a constant term and the logarithm of the expected number of reads as computed in the

363   previous section, $\lambda_{t,p}$:

$$\mu_{\text{Peak},t,p} = \exp\left[\alpha_t + \gamma_t \log \lambda_{t,p}\right]$$

$$\mu_{\text{Background},t,p} = \exp\left[\beta_t + \gamma_t \log \lambda_{t,p}\right]$$

where α_t, β_t and γ_t are time point specific scalar parameters. Negative binomial distributions have been

successfully employed in a similar manner to capture the over-dispersion of sequencing reads in ChIP-

seq experiments in peak callers for single samples such as ZINBA[56]. Of note however, ChromTime

requires that the dispersion parameter δ_t and the coefficient γ_t are shared between the two components

at each time point. The first requirement ensures that the distribution with the smaller mean value has

higher probabilities compared to the distribution with the larger mean value for the lowest values of the

support domain of the negative binomial distribution, and that the opposite holds for the largest values of

the support domain (See **Supplementary Methods**). Sharing the dispersion parameter here is analogous

to sharing the variance parameter in Gaussian mixture models. The second requirement to share the γ_t

parameter ensures that the input signal has equal importance in each component.

Formally, let $B_{L,t}$ and $B_{R,t}$ denote the bin indices relative to the beginning of the block of the first and the

last bin in the peak partition at time t, respectively, and N be the length of the block (i.e. $1 \leq B_{L,t} \leq N +$

1 and $0 \leq B_{R,t} \leq N$, with values of $B_{L,t} = N + 1$ and $B_{R,t} = 0$ corresponding to the special cases of starting

a peak after all positions and ending a peak before all positions in a block, respectively). For $B_{L,t}$ and $B_{R,t}$

to denote valid interval boundaries, ChromTime requires that $B_{L,t} \leq B_{R,t} + 1$ at each time point. This can

be formally encoded by introducing one auxiliary binary variable for each time point in the model, Z_t, such

that:

$$P(Z_t = 1 | B_{L,t} = l, B_{R,t} = r) = \begin{cases} 1 & \text{if } l \leq r + 1 \\ 0 & \text{otherwise} \end{cases}$$

385   and thus also

$$P(Z_t = 0 | B_{L,t} = l, B_{R,t} = r) = \begin{cases} 0 & \text{if } l \leq r + 1 \\ 1 & \text{otherwise} \end{cases}$$

387   ChromTime treats all Z_t variables as observed with values equal to 1 for all blocks and time points. Then,

388   the probability of a given partitioning of the ChIP-seq signal at time t under the model is:

$$P\big(B_{L,t} = l, B_{R,t} = r, Z_t = 1, O_t \big| X_t\big)$$

$$= P\big(Z_t = 1 | B_{L,t} = l, B_{R,t} = r\big)$$

$$\times \prod_{p=1}^{l-1} \text{NB}(O_{t,p}; \mu_{\text{Background},t} = \exp[\beta_t + \gamma_t \log \lambda_{t,p}], \delta_t)$$

$$\times \prod_{p=l}^{r} \text{NB}\big(O_{t,p}; \mu_{\text{Peak},t} = \exp[\alpha_t + \gamma_t \log \lambda_{t,p}], \delta_t\big)$$

$$\times \prod_{p=r+1}^{N} \text{NB}(O_{t,p}; \mu_{\text{Background},t} = \exp[\beta_t + \gamma_t \log \lambda_{t,p}], \delta_t)$$

391   An important special case of the above formulation when $B_{L,t} = B_{R,t} + 1$ corresponds to modelling the

392   whole signal at time point t as background, which enables ChromTime to accommodate time points with

393   no peaks.

395   ChromTime assumes uniform prior probabilities for the left and the right end boundaries at the first time

396   point:

$$P(B_{L,1} = l) = Unif(1, N + 1)$$

397 and

$$P(B_{R,1} = r) = Unif(0, N)$$

where $Unif(a, b)$ denotes the uniform distribution of integer numbers in the closed interval $[a, b]$.

Between any two time points the ChromTime model allows for one of three possible dynamics at both the

left and the right end boundaries of a peak: Steady, Expand or Contract. To capture the change of

boundary positions between consecutive time points t and $t+1$ we define the quantities $J_{L,t} = B_{L,t} - B_{L,t+1}$

and $J_{R,t} = B_{R,t+1} - B_{R,t}$ corresponding to the left and right boundaries respectively. Positive values of $J_{L,t}$

and $J_{R,t}$ indicate the number of bases a peak expanded, whereas negative values indicate the number of

bases a peak contracted, and a value of 0 indicates the peak held steady on the left and the right side

respectively. ChromTime models $J_{L,t}$ *and* $J_{R,t}$ with a different probability distribution for each of the three

dynamics. Let $D_{S,t}$ denote the dynamic between time points t and $t+1$ on boundary side S, where S is one

of L (left side) or R (right side). For Steady dynamic, ChromTime uses the Kronecker delta function:

$$P\big(J_{S,t}\big|D_{S,t} = \text{Steady}\big) = \begin{cases} 1, & \text{if } J_{S,t} = 0 \\ 0, & \text{otherwise} \end{cases}$$

412 For Expand and Contract, ChromTime employs negative binomial distributions to model the number of

413 genomic bins a peak boundary moves relative to the minimal movement of one bin required for peak

414 expansions and contractions:

$$P\big(J_{S,t} = j\big|D_{S,t} = \text{Expand}\big) = NB(j - 1; \mu_{\text{Expand,t}}, \delta_{\text{Expand,t}})$$

416 and

$$P\big(J_{S,t} = j\big|D_{S,t} = \text{Contract}\big) = NB(-j - 1; \mu_{\text{Conract,t}}, \delta_{\text{Contract,t}})$$

Furthermore, each distribution is parametrized with a mean and dispersion parameter depending on the

dynamic and the time point, t. $\mu_{\text{Expand,t}}, \delta_{\text{Expand,t}}$ for expansions, and $\mu_{\text{Contract,t}}, \delta_{\text{Contract,t}}$ for contractions.

Of note, in negative binomial distributions the probabilities for negative integers are defined to be 0.

Therefore, the above parametrization enforces that boundary movements of negative or zero length (i.e.

contracting or steady, respectively) are impossible for expansions and that boundary movements of

positive or zero length (i.e. expanding or steady) are impossible for contractions.

The ChromTime model additionally assumes that there is a prior probability to observe each dynamic

between time points t and $t+1$, $P(D_{S,t} = d) = \pi_{t,d}$, which is the same at each side (left and right). Thus,

the joint probability to observe a particular partition of the data into peaks and background and dynamics

$d_{L,t}$ and $d_{R,t}$ on the left and right side at two consecutive time points, respectively, can be expressed as:

$$P\big(B_{L,t} = l_t, B_{L,t+1} = l_{t+1}, D_{L,t} = d_{L,t}, B_{R,t} = r_t, B_{R,t+1} = r_{t+1}, D_{R,t} = d_{R,t}, Z_t = 1, Z_{t+1} = 1, O_t, O_{t+1} | X_t, X_{t+1}\big) =$$

$$= P\big(B_{L,t} = l_t, B_{R,t} = r_t, Z_t = 1, O_t | X_t\big) \times P\big(J_{L,t} = l_t - l_{t+1} | D_{L,t} = d_{L,t}\big) \times P\big(D_{L,t} = d_{L,t}\big)$$

$$\times P\big(J_{R,t} = r_{t+1} - r_t | D_{R,t} = d_{R,t}\big) \times P\big(D_{R,t} = d_{R,t}\big)$$

$$\times P\big(B_{L,t+1} = l_{t+1}, B_{R,t+1} = r_{t+1}, Z_{t+1} = 1, O_{t+1} | X_{t+1}\big)$$

431 In a block O with covariates X in a time course with T time points, the total probability of a particular

432 sequence of dynamics and boundary positions on the left side ($\mathcal{D}_L$ and $\mathcal{B}_L$ respectively) and on the right

433 side ($\mathcal{D}_R$ and $\mathcal{B}_R$ respectively) can be expressed as:

$$P(\mathcal{D}_L, \mathcal{B}_L, \mathcal{D}_R, \mathcal{B}_R, O|X) =$$

$$P\big(B_{L,1} = l_1, B_{R,1} = r_1, Z_1 = 1, O_1 | X_1\big) \prod_{t=2}^{T} \Big(P\big(B_{L,t} = l_t, B_{R,t} = r_t, Z_t = 1, O_t | X_t\big)$$

$$\times P\big(J_{L,t-1} = l_{t-1} - l_t | D_{L,t-1} = d_{L,t-1}\big) \times P\big(D_{L,t-1} = d_{L,t-1}\big)$$

$$\times P\big(J_{R,t-1} = r_t - r_{t-1} | D_{R,t-1} = d_{R,t-1}\big) \times P\big(D_{R,t-1} = d_{R,t-1}\big)\Big)$$

where $\mathcal{D}_S = (d_{S,t} | 1 \leq t \leq T - 1)$ are *T-1* dimensional vectors of dynamics labels for each pair of

consecutive time points on the left or the right side ($S = L$ or R), respectively, and $\mathcal{B}_S$ are the

corresponding *T* dimensional vectors of boundary positions on each side at each time point. The total

probability of the ChIP-seq signal in a block can be expressed as a sum over all possible sequences of

dynamics and peak boundary positions that can generate the block across all time points. Thus, the

probability of block $\boldsymbol{O}$ is:

$$P(\boldsymbol{O}|\boldsymbol{X}) = \sum_{\mathfrak{D}_L, \mathcal{B}_L, \mathfrak{D}_R, \mathcal{B}_R} P(\mathfrak{D}_L, \mathcal{B}_L, \mathfrak{D}_R, \mathcal{B}_R, \boldsymbol{O}|\boldsymbol{X})$$

where $\mathfrak{D}_L$ and $\mathfrak{D}_R$ each iterate over all possible 3^{T-1} combinations of peak boundary dynamics, and

$\mathcal{B}_L$ and $\mathcal{B}_R$ each iterate over all possible ways to place left and right end boundaries across all time points

that are consistent with the requirements that $B_{L,t} \leq B_{R,t} + 1$ at each time point. Let $\mathcal{O}$ be the total set of

blocks in the data, $\mathcal{X}$ be the set of two-dimensional vectors containing the constant term and the log of

the expected number of reads at each position and time point for each block, and *M* be the total number

of blocks. Then the total likelihood of all blocks is:

$$P(\mathcal{O}|\mathcal{X}) = \prod_{i=1}^{M} P(\boldsymbol{O}_i|\boldsymbol{X}_i)$$

452   **Model optimization**

454   The total set of parameters of the model consists of:

455   *1)* Prior probabilities of each dynamic *d* at each time point *t*: $\pi_{t,d}$

456   2) Parameters of the negative binomial distributions that model the Peak and the Background

457   components at each time point: α_t, β_t, γ_t and δ_t.

458   3) Parameters of the negative binomial distributions that model the boundary movements in Expand

459   and Contract dynamics at each time point: $\mu_{\text{Expand},t}$, $\delta_{\text{Expand},t}$ and $\mu_{\text{Contract},t}$, $\delta_{\text{Contract},t}$, respectively.

461 The optimal parameter values are attempted to be estimated by Expectation Maximization (EM). In

462 particular, ChromTime finds a local maximum of the expectation of the complete log-likelihood function

463 (for details, see **Supplementary Methods**):

$$LL\big(\theta;\tilde{\theta},\mathcal{O},\mathcal{X}\big) = \sum_{i=1}^{M} \log P(\boldsymbol{O}_i|\boldsymbol{X}_i) =$$

$$= \sum_{i=1}^{M}\sum_{l_1=1}^{N_i+1}\sum_{r_1=l_1-1}^{N_i} P\big(B_{L,1}=l_1,B_{R,1}=r_1|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)\big(\log P\big(B_{L,1}=l_1\big)+\log P\big(B_{R,1}=r_1\big)\big)$$

$$+ \sum_{i=1}^{M}\sum_{t=1}^{T}\sum_{p=1}^{N_i} \Big(P\big(V_{t,p}=\text{Peak}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)\log\Big(\text{NB}\big(O_{i,t,p};\mu_{\text{Peak},t}=\exp[\alpha_t+\gamma_t\log\lambda_{i,t,p}],\delta_t\big)\Big)$$

$$+ P\big(V_{t,p}=\text{Background}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)\log\Big(\text{NB}\big(O_{i,t,p};\mu_{\text{Background},t}=\exp[\beta_t+\gamma_t\log\lambda_{i,t,p}],\delta_t\big)\Big)\Big)$$

$$+ \sum_{i=1}^{M}\sum_{t=1}^{T-1}\sum_{S\in\{L,R\}}\sum_{d_{S,t}\in\left\{\substack{\text{Expand}\\\text{Contract}}\right\}}\sum_{j=1}^{N_i} P\big(J_{S,t}=(-1)^{\omega(d_{S,t})}j, D_{S,t}=d_{S,t}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)\log\Big(\text{NB}\big(j-1;\mu_{d_{S,t}},\delta_{d_{S,t}}\big)\Big)$$

$$+ \sum_{i=1}^{M}\sum_{t=1}^{T-1}\sum_{S\in\{L,R\}}\sum_{d_{S,t}\in\left\{\substack{\text{Steady}\\\text{Expand}\\\text{Contract}}\right\}} P\big(D_{S,t}=d_{S,t}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)\log\big(\pi_{t,d_{S,t}}\big)$$

466 where

$$\omega\big(d_{S,t}\big) = \begin{cases}1 & \text{if } d_{S,t}=\text{Contract}\\0 & \text{otherwise}\end{cases}$$

469 In the above equation, $\tilde{\theta}$ denotes the set of parameters at the current iteration, and θ denotes the set of

470 parameters to be estimated in the next iteration. Additionally, $P\big(V_{t,p}=\text{Peak}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)$ and

471 $P\big(V_{t,p}=\text{Background}|\boldsymbol{O}_i,\boldsymbol{X}_i;\tilde{\theta}\big)$ denote the posterior probabilities that the bin of block *i* at position *p* and

472 time point *t* is assigned to the peak and the background component, respectively:

$$P\left(V_{t,p} = \text{Peak}\middle|\boldsymbol{O}_i, \boldsymbol{X}_i; \tilde{\theta}\right) = \sum_{l,r\,|\,l \leq p \leq r} P\left(B_{L,t} = l, B_{R,t} = r, Z_t = 1\middle|\boldsymbol{O}_i, \boldsymbol{X}_i; \tilde{\theta}\right)$$

$$P\left(V_{t,p} = \text{Background}\middle|\boldsymbol{O}_i, \boldsymbol{X}_i; \tilde{\theta}\right) = 1 - P\left(V_{t,p} = \text{Peak}\middle|\boldsymbol{O}_i, \boldsymbol{X}_i; \tilde{\theta}\right)$$

Parameter initialization

The values of the model parameters before the first EM iteration are initialized as follows:

1) All dynamics priors, $\pi_{t,d}$, are set uniformly to $\frac{1}{3}$;

2) All parameters for the distributions modelling the Peak and Background components,

α_t, β_t, γ_t and δ_t, are set to 1;

3) All dispersion and mean parameters for the distributions modelling the boundary movements in

Expand and Contract dynamics, $\mu_{\text{Expand,t}}$, $\delta_{\text{Expand,t}}$, $\mu_{\text{Contract,t}}$, and $\delta_{\text{Contract,t}}$, are set to 1.

At each time point, ChromTime requires that the left boundary of each non-zero length peak is placed

before its right boundary. This requirement results in non-uniform prior probabilities for the $V_{t,p}$ variables,

which model whether a bin at position *p* at time point *t* belongs to the peak or to the background

component, making bins in the middle of the block more likely to be in a peak compared to flanking bins

on each side. In particular, at any single time point the prior probability for a bin at position p ($1 \leq p \leq N$)

to be in a peak is:

$$P\left(V_{t,p} = \text{Peak}|Z_t = 1\right) = \frac{\#\,[\text{peak intervals containing bin } p \text{ in a block of length } N]}{\#\,[\text{total peak intervals in a block of length } N]}$$

$$= \frac{p(N - p + 1)}{\sum_{i=1}^{N+1}(N - i + 2)} = \frac{2p(N - p + 1)}{(N + 1)(N + 2)}$$

which is symmetric with respect to the center of the block and has its maximum of $\frac{N}{2(N+1)} < \frac{1}{2}$ at $p = \frac{N}{2}$. This

prior probability plays an important role during the learning stage of ChromTime, because it directs the

model during the initial iterations of the EM to correctly associate the Peak component with high ChIP-seq

signal and the Background component with low ChIP-seq signal. The assumption that high ChIP-seq

signal will more likely be located in the middle of blocks is motivated by the procedure that determines the

block boundaries in the first phase of ChromTime, which naturally produces blocks with high signal in the

middle compared to their flanking regions (**Fig S1D**). As result, no further efforts from the initialization or

the training procedures are necessary in practice to identify correctly each component.

Expectation step

ChromTime provides an efficient implementation of the expectation step of EM based on a dynamic

programming algorithm similar to the Baum-Welch algorithm for hidden Markov models. In brief, at each

time point there are $O(N^2)$ ways to place the start and end positions of a peak, resulting in $O(N^4)$

combinations between any pair of consecutive time points. Thus, a standard forward-backward procedure

that caches intermediate results can compute all expectations in $O(T^*N^4)$ time and $O(T^*N^2)$ memory.

Since $O(T^*N^4)$ time complexity can result in very long running times even for moderate N, ChromTime

splits blocks that are longer than a predefined number of bins, MAX_BINS, (30 by default) into two halves

(left and right) and estimates all sufficient statistics in each half independently. If block i is longer than

MAX_BINS, the split is performed at the position with the highest average ChIP-seq signal across all time

points in the block, K_i. This splitting procedure corresponds to imposing an additional constraint on the

values of the boundary position variables that $B_{L,t} \leq K_i$ and $B_{R,t} \geq K_i - 1$ at each time point, t, while still

having all bins between the left and the right boundaries annotated as peak bins (i.e. $V_{t,p} = \text{Peak}$ for all p

such that $B_{L,t} \leq p \leq B_{R,t}$ and $V_{t,p} = \text{Background}$ for all other values of p). This heuristic reduces the time

complexity to $O(T^*N^2)$ and the memory footprint to $O(T^*N)$, thus making the whole EM procedure run in

feasible time and space. Since the $O(T^*N^4)$ algorithm is applied only to blocks shorter than MAX_BINS

bins, the total running time of ChromTime in a dataset of M blocks remains at most quadratic in the length

of longer peaks in the data, $O(M^*T^*N^2)$.

For computational efficiency, if there are more than 10,000 blocks, ChromTime randomly selects 10,000

as input for the EM procedure.

Maximization step

The form of the complete log-likelihood implies that each set of model parameters can be optimized

independently by solving for the roots of the respective partial derivatives. The dynamics prior

probabilities are updated after each EM iteration as:

$$P\big(D_{L,t} = d\big) = P\big(D_{R,t} = d\big) = \frac{1}{2M}\sum_{i=1}^{M}\sum_{S\in\{L,R\}} P(D_{S,t} = d|\boldsymbol{O}_i, \boldsymbol{X}_i; \tilde{\theta})$$

The parameters of all negative binomial distributions in the model are determined by using an iterative

numerical procedure that aims to find the roots of the partial derivatives of the complete log-likelihood

function with respect to each parameter. In brief, finding the maximum likelihood estimators for each

distribution at each EM iteration is equivalent to fitting weighted negative binomial regressions with the

corresponding posterior probabilities as weights. For this purpose, ChromTime re-implements a widely

used fitting method for weighted negative binomial regressions from the MASS package[57] in R except

that the Iterative Reweighted Least Squares part is substituted with the HYBRD method for root finding

from the MINPACK package[58] and the overall procedure is further extended to handle shared

parameters. More details are provided in the **Supplementary Methods**.

Computing the most likely spatial dynamic and peak boundaries for each block across the whole

time course

After the optimal values for all model parameters are estimated from the data, for each block the most

likely positions of the peak boundaries at each time point are calculated. This procedure consists of two

steps. First, ChromTime determines for each block all time points with significantly low probability of

containing a false positive non-zero peak. Second, conditioned on those time points, ChromTime

computes the most likely assignment of the peak boundary variables at each side and each time point.

During the first step, for each block and each time point ChromTime computes the posterior probability

that the whole time point is modelled as background, $\varphi_{t,i} = \prod_{p=1}^{N_i} P(V_{t,p} = \text{Background}|\boldsymbol{O}_i, \boldsymbol{X}_i)$. This

probability can be interpreted as the probability of making a false positive non-zero length peak call as

estimated by the model at time point t in block O_i. To determine significant time points with low false

positive probability, $\varphi_{t,i}$, ChromTime computes a time point specific threshold, τ_t, at a predefined false

discovery rate (0.05 by default) by applying the standard Benjamini-Hochberg procedure[59] on all values

of $\varphi_{t,i}$ from time point t.

In the second step, for each block ChromTime computes the most likely sequence of assignments of the

boundary positions, conditioned on the event that all time points that failed to pass the FDR threshold in

the previous step for the block are assigned to having no peaks. In particular, ChromTime executes a

dynamic programming algorithm similar to the Viterbi algorithm for hidden Markov models, which uses the

following recursive formula to find the most likely position for the peak boundaries at each side, *S* (Left or

Right) and enforces that time points that failed the FDR test contain no peaks:

$$
\mathrm{DP}_{t,l,r} = \begin{cases} \log P\big(B_{L,1} = l, B_{R,1} = r, O_1 \big| X_1\big) & , t = 1 \\[2em] \max\limits_{\substack{l_{t-1} \in [1, N+1] \\ r_{t-1} \in \begin{cases} [l_{t-1}-1, N] \text{ if } \varphi_{t,i} \leq \tau_t \\ \{l_{t-1}-1\} \text{ otherwise} \end{cases}}} \begin{pmatrix} \mathrm{DP}_{t,l_{t-1},r_{t-1}} + \\ \log P\big(B_{L,t} = l, B_{R,t} = r, O_t \big| X_t\big) + \\ \log P\big(J_{L,t-1} = l_{t-1} - l \big| D_{L,t-1} = \mathrm{DYN}(l_{t-1} - l)\big) + \\ \log P\big(D_{L,t-1} = \mathrm{DYN}(l_{t-1} - l)\big) + \\ \log P\big(J_{R,t-1} = r - r_{t-1} \big| D_{R,t-1} = \mathrm{DYN}(r - r_{t-1})\big) + \\ \log P\big(D_{R,t-1} = \mathrm{DYN}(r - r_{t-1})\big) \end{pmatrix} & , t > 1 \end{cases}
$$

$$
\text{where } \mathrm{DYN}(j) = \begin{cases} \text{Steady} & \text{if } j = 0 \\ \text{Expand} & \text{if } j \geq 1 \\ \text{Contract} & \text{if } j \leq -1 \end{cases}
$$

and DP denotes the dynamic programing cube of size $T(N+1)^2$ that stores the log likelihood for the best

assignment of the peak boundary variables up to time point t. Tracing the DP cube from the highest value

on row *T* back to row 1 retrieves the best assignment of the peak end variables. Similarly to the

expectation step of the EM phase, for blocks longer than MAX_BINS bins the best Viterbi path is chosen

among the splits at the top MAX_BINS positions in the block sorted by their average ChIP-seq signal

across all time points. Since MAX_BINS is a predefined constant, the whole procedure has the same time

and space complexity as computing the expectations in the EM phase of ChromTime. The dynamic

between any two time points is determined from the direction of the movement of the optimal positions of

the corresponding boundaries.

Finally, the software provides an option to parallelize all computations during the EM phase and the

Viterbi decoding over multiple CPU cores, which can further reduce the running time by up to a factor

equal to the number of allocated cores. In our tests, processing ChIP-seq for the H3K4me2 mark and

control input data from 5 time points in mouse T cell development[15] took 3 hours by using 4 CPU cores

on a MacBook Pro laptop with 2.7GHz Intel Core i7 and 16 GB RAM.

ChromTime parameters used in this study

In this work, we applied ChromTime in narrow mode on all data for H3K4me2, H3K4me3, H3K27ac and

H3K(9,14)ac marks. We applied ChromTime in broad mode on all data for H3K79me2, Pol2, H3K4me1,

H3K27me3, H3K9me3 and H3K36me3 marks. All other parameters were set to their default values.

Analyses with External Data

The procedures for analyses with external are described in **Supplementary Methods.**

ABBREVIATIONS

HM: Histone mark

TF: Transcription factor

TSS: Transcription start-site

ChIP-seq: Chromatin immunoprecipitation coupled with high throughput DNA sequencing

FDR: False Discovery Rate

EM: Expectation Maximization

DECLARATIONS

Authors' contributions

J.E. conceived and supervised the project, designed the method, proposed analyses and wrote the

manuscript. P.F. designed the method, implemented the software, proposed analyses, performed all

analyses, processed all data and wrote the manuscript. Both authors read and approved the final

manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

ChromTime software is freely available at: https://github.com/ernstlab/ChromTime

No new experimental datasets were generated within this study. All ChIP-seq datasets used as input for

ChromTime are publicly available from the references listed in **Table 1**. Links to all other datasets (gene

expression, TF binding and DNaseI hypersensitivity) used in this study are provided in the

Supplementary Methods.

Funding

This work was supported by the CIRM Training Grant TG2-01169, the Eli and Edythe Broad Center of

Regenerative Medicine and Stem Cell Research at UCLA Training Program (P.F.); NIH grants

R01ES024995, U01HG007912, DP1DA044371, an NSF CAREER Award #1254200 and an Alfred P.

Sloan Fellowship (J.E.).

Acknowledgements

We are grateful to Constantinos Chronis, Kathrin Plath and members of the Ernst lab for useful discussions.

Competing interests

None

REFERENCES

[1]   J. Ernst *et al.*, "Mapping and analysis of chromatin state dynamics in nine human cell types.," *Nature*, vol. 473, no. 7345, pp. 43–9, May 2011.

[2]   A. Barski *et al.*, "High-resolution profiling of histone methylations in the human genome.," *Cell*, vol. 129, no. 4, pp. 823–37, May 2007.

[3]   T. S. Mikkelsen *et al.*, "Genome-wide maps of chromatin state in pluripotent and lineage-committed cells.," *Nature*, vol. 448, no. 7153, pp. 553–60, Aug. 2007.

[4]   ENCODE_Project_Consortium, "An integrated encyclopedia of DNA elements in the human genome.," *Nature*, vol. 489, no. 7414, pp. 57–74, Sep. 2012.

[5]   Roadmap_Epigenomics_Consortium *et al.*, "Integrative analysis of 111 reference human epigenomes," *Nature*, vol. 518, no. 7539, pp. 317–330, Feb. 2015.

[6]   J. H. A. Martens and H. G. Stunnenberg, "BLUEPRINT: mapping human blood cell epigenomes.," *Haematologica*, vol. 98, no. 10, pp. 1487–9, Oct. 2013.

[7]   F. D. Lay *et al.*, "Reprogramming of the human intestinal epigenome by surgical tissue transposition.," *Genome Res.*, vol. 24, no. 4, pp. 545–53, Apr. 2014.

[8]   P. Fiziev *et al.*, "Systematic Epigenomic Analysis Reveals Chromatin States Associated with Melanoma Progression.," *Cell Rep.*, vol. 19, no. 4, pp. 875–889, Apr. 2017.

[9]   S. Mei *et al.*, "Cistrome Data Browser: a data portal for ChIP-Seq and chromatin accessibility data

in human and mouse," *Nucleic Acids Res.*, vol. 45, no. D1, pp. D658–D662, Jan. 2017.

[10]   M. J. Ziller *et al.*, "Dissecting neural differentiation regulatory networks through epigenetic

footprinting.," *Nature*, vol. 518, no. 7539, pp. 355–359, Dec. 2014.

[11]   A. M. Tsankov *et al.*, "Transcription factor binding dynamics during human ES cell differentiation,"

Nature, vol. 518, no. 7539, pp. 344–349, Feb. 2015.

[12]   K. K.-H. Farh *et al.*, "Genetic and epigenetic fine mapping of causal autoimmune disease variants,"

Nature, vol. 518, no. 7539, pp. 337–343, Oct. 2014.

[13]   E. Gjoneska *et al.*, "Conserved epigenomic signals in mice and humans reveal immune basis of

Alzheimer's disease," *Nature*, vol. 518, no. 7539, pp. 365–369, Feb. 2015.

[14]   A. Gusev *et al.*, "Partitioning heritability of regulatory and cell-type-specific variants across 11

common diseases.," *Am. J. Hum. Genet.*, vol. 95, no. 5, pp. 535–52, Nov. 2014.

[15]   J. A. Zhang, A. Mortazavi, B. A. Williams, B. J. Wold, and E. V Rothenberg, "Dynamic

transformations of genome-wide epigenetic marking and transcriptional control establish T cell

identity.," *Cell*, vol. 149, no. 2, pp. 467–82, Apr. 2012.

[16]   T. S. Mikkelsen *et al.*, "Comparative epigenomic analysis of murine and human adipogenesis.,"

Cell, vol. 143, no. 1, pp. 156–69, Oct. 2010.

[17]   D. Lara-Astiaso *et al.*, "Chromatin state dynamics during blood formation," *Science (80-.).*, p.

science.1256271-, Aug. 2014.

[18]   D. K. Goode *et al.*, "Dynamic Gene Regulatory Networks Drive Hematopoietic Specification and

Differentiation," *Dev. Cell*, Feb. 2016.

[19]   S. L. Paige *et al.*, "A temporal chromatin signature in human embryonic stem cells identifies

regulators of cardiac development.," *Cell*, vol. 151, no. 1, pp. 221–32, Sep. 2012.

[20]   J. A. Wamstad *et al.*, "Dynamic and coordinated epigenetic regulation of developmental transitions

in the cardiac lineage.," *Cell*, vol. 151, no. 1, pp. 206–20, Sep. 2012.

[21]   D. Cacchiarelli *et al.*, "Integrative Analyses of Human Reprogramming Reveal Dynamic Nature of

Induced Pluripotency," *Cell*, vol. 162, no. 2, pp. 412–424, Jul. 2015.

[22] J. Chen *et al.*, "Hierarchical Oct4 Binding in Concert with Primed Epigenetic Rearrangements

during Somatic Cell Reprogramming," *Cell Rep.*, Jan. 2016.

[23] R. P. Koche *et al.*, "Reprogramming factor expression initiates widespread targeted chromatin

remodeling.," *Cell Stem Cell*, vol. 8, no. 1, pp. 96–105, Jan. 2011.

[24] C. Chronis *et al.*, "Cooperative Binding of Transcription Factors Orchestrates Reprogramming.,"

Cell, vol. 168, no. 3, p. 442–459.e20, Jan. 2017.

[25] O. Bogdanovic *et al.*, "Dynamics of enhancer chromatin signatures mark the transition from

pluripotency to cell specification during embryogenesis.," *Genome Res.*, vol. 22, no. 10, pp. 2043–

53, Oct. 2012.

[26] N. Nègre *et al.*, "A cis-regulatory map of the Drosophila genome.," *Nature*, vol. 471, no. 7339, pp.

527–31, Mar. 2011.

[27] P. Yu *et al.*, "Spatiotemporal clustering of epigenome reveals rules of dynamic gene regulation.,"

Genome Res., p. gr.144949.112-, Oct. 2012.

[28] P. Arnold *et al.*, "Modeling of epigenome dynamics identifies transcription factors that mediate

Polycomb targeting.," *Genome Res.*, p. gr.142661.112-, Sep. 2012.

[29] N. Koike *et al.*, "Transcriptional architecture and chromatin landscape of the core circadian clock in

mammals.," *Science*, vol. 338, no. 6105, pp. 349–54, Oct. 2012.

[30] R. Ostuni *et al.*, "Latent Enhancers Activated by Stimulation in Differentiated Cells," *Cell*, vol. 152,

no. 1–2, pp. 157–71, Jan. 2013.

[31] S. K. Reilly *et al.*, "Evolutionary changes in promoter and enhancer activity during human

corticogenesis," *Science (80-.)*., vol. 347, no. 6226, pp. 1155–1159, Mar. 2015.

[32] A. Weiner *et al.*, "High-Resolution Chromatin Dynamics during a Yeast Stress Response.," *Mol.*

Cell, vol. 58, no. 2, pp. 371–386, Mar. 2015.

[33] J. Cotney *et al.*, "The evolution of lineage-specific regulatory activities in the human embryonic

limb.," *Cell*, vol. 154, no. 1, pp. 185–96, Jul. 2013.

[34]   J. Zhu *et al.*, "A time-series analysis of altered histone H3 acetylation and gene expression during

704   the course of MMAIII-induced malignant transformation of urinary bladder cells," *Carcinogenesis*,

705   vol. 38, no. 4, pp. 378–390, Apr. 2017.

706   [35]   Y. Zhang *et al.*, "Model-based analysis of ChIP-Seq (MACS).," *Genome Biol.*, vol. 9, no. 9, p.

707   R137, Jan. 2008.

708   [36]   S. Heinz *et al.*, "Simple combinations of lineage-determining transcription factors prime cis-

709   regulatory elements required for macrophage and B cell identities.," *Mol. Cell*, vol. 38, no. 4, pp.

710   576–89, May 2010.

711   [37]   S. Xu, S. Grullon, K. Ge, and W. Peng, "Spatial clustering for identification of ChIP-enriched

712   regions (SICER) to map regions of histone methylation patterns in embryonic stem cells.,"

713   *Methods Mol. Biol.*, vol. 1150, pp. 97–111, 2014.

714   [38]   H. Xing *et al.*, "Genome-Wide Localization of Protein-DNA Binding and Histone Modification by a

715   Bayesian Change-Point Method with ChIP-seq Data," *PLoS Comput. Biol.*, vol. 8, no. 7, p.

716   e1002613, Jul. 2012.

717   [39]   A. Harmanci, J. Rozowsky, and M. Gerstein, "MUSIC: identification of enriched regions in ChIP-

718   Seq experiments using a mappability-corrected multiscale signal processing framework.," *Genome

719   Biol.*, vol. 15, no. 10, p. 474, 2014.

720   [40]   J. Ernst and M. Kellis, "ChromHMM: automating chromatin-state discovery and characterization.,"

721   *Nat. Methods*, vol. 9, no. 3, pp. 215–6, Mar. 2012.

722   [41]   M. M. Hoffman, O. J. Buske, J. Wang, Z. Weng, J. A. Bilmes, and W. S. Noble, "Unsupervised

723   pattern discovery in human chromatin structure through genomic segmentation." *Nat. Methods*,

724   vol. 9, no. 5, pp. 473–6, May 2012.

725   [42]   M. Allhoff, K. Seré, H. Chauvistré, Q. Lin, M. Zenke, and I. G. Costa, "Detecting differential peaks

726   in ChIP-seq signals with ODIN.," *Bioinformatics*, vol. 30, no. 24, pp. 3467–75, Dec. 2014.

[43] H. Xu, C.-L. Wei, F. Lin, and W.-K. Sung, "An HMM approach to genome-wide identification of

differential histone modification sites from ChIP-seq data.," *Bioinformatics*, vol. 24, no. 20, pp.

2344–9, Oct. 2008.

[44] J. Biesinger, Y. Wang, and X. Xie, "Discovering and mapping chromatin states using a tree hidden

Markov model," *BMC Bioinformatics*, vol. 14, no. Suppl 5, p. S4, 2013.

[45] D. Hnisz *et al.*, "Super-enhancers in the control of cell identity and disease.," *Cell*, vol. 155, no. 4,

pp. 934–47, Nov. 2013.

[46] S. C. J. Parker *et al.*, "Chromatin stretch enhancer states drive cell-specific gene regulation and

harbor human disease risk variants," *PNAS*, p. 1317023110-, Oct. 2013.

[47] B. A. Benayoun *et al.*, "H3K4me3 breadth is linked to cell identity and transcriptional consistency.,"

Cell, vol. 158, no. 3, pp. 673–88, Jul. 2014.

[48] K. Chen *et al.*, "Broad H3K4me3 is associated with increased transcription elongation and

enhancer activity at tumor-suppressor genes.," *Nat. Genet.*, vol. 47, no. 10, pp. 1149–57, Oct.

2015.

[49] A. Dincer *et al.*, "Deciphering H3K4me3 broad domains associated with gene-regulatory networks

and conserved epigenomic landscapes in the human brain," *Transl. Psychiatry*, vol. 5, no. 11, p.

e679, Nov. 2015.

[50] R. D. Hawkins *et al.*, "Distinct epigenomic landscapes of pluripotent and lineage-committed human

cells.," *Cell Stem Cell*, vol. 6, no. 5, pp. 479–91, May 2010.

[51] Z. Wang *et al.*, "Combinatorial patterns of histone acetylations and methylations in the human

genome.," *Nat. Genet.*, vol. 40, no. 7, pp. 897–903, Jul. 2008.

[52] H. H. Ng, F. Robert, R. A. Young, and K. Struhl, "Targeted recruitment of Set1 histone methylase

by elongating Pol II provides a localized mark and memory of recent transcriptional activity.," *Mol.*

Cell, vol. 11, no. 3, pp. 709–19, Mar. 2003.

[53] E. Smith, C. Lin, and A. Shilatifard, "The super elongation complex (SEC) and MLL in

752   development and disease.," *Genes Dev.*, vol. 25, no. 7, pp. 661–72, Apr. 2011.

753   [54]   S. A. Lacadie, M. M. Ibrahim, S. A. Gokhale, and U. Ohler, "Divergent Transcription and

754   Epigenetic Directionality of Human Promoters," *FEBS J.*, Apr. 2016.

755   [55]   A. C. Cameron and P. K. Trivedi, *Regression analysis of count data*, Second edition. Cambridge

756   University Press, 2013.

757   [56]   N. U. Rashid, P. G. Giresi, J. G. Ibrahim, W. Sun, and J. D. Lieb, "ZINBA integrates local

758   covariates with DNA-seq data to identify broad and narrow regions of enrichment, even within

759   amplified genomic regions," *Genome Biol.*, vol. 12, no. 7, p. R67, 2011.

760   [57]   W. N. Venables and B. D. Ripley, *Modern Applied Statistics with S*, Fourth. New York: Springer,

761   2002.

762   [58]   Moré, B. Garbow, and K. Hillstrom, "User Guide for MINPACK-1," *ANL-80-74, Argonne Natl. Lab.*,

763   1980.

764   [59]   Y. Benjamini and Y. Hochberg, "Controlling the False Discovery Rate: A Practical and Powerful

765   Approach to Multiple Testing," *Journal of the Royal Statistical Society. Series B (Methodological)*,

766   vol. 57. WileyRoyal Statistical Society, pp. 289–300, 1995.

767   [60]   Z. Bar-Joseph, D. K. Gifford, and T. S. Jaakkola, "Fast optimal leaf ordering for hierarchical

768   clustering.," *Bioinformatics*, vol. 17 Suppl 1, pp. S22-9, 2001.

FIGURE LEGENDS

Figure 1: **Overview of the ChromTime method. (A)** Examples of locations of peaks of the H3K4me2

mark that exhibit steady, expanding and contracting boundary dynamics, shown from left to right

respectively, across five time points during T cell development in mouse[15]. Time points 1, 2 and 3

correspond to in vitro differentiated T cell precursors (FLDN1, FLDN2a, and FLDN2b), whereas time

points 4 and 5 correspond to in vivo purified thymocytes (ThyDN3 and ThyDP). Normalized ChIP-seq

signal and called MACS2[35] peaks are shown at each time point. The bottom part of each plot shows the

corresponding ChromTime output. A peak upstream of Zfp148 gene remains relatively steady with

fluctuations of MACS2 peak boundaries across time. In contrast, a peak at the Skap1/GM11529 promoter

expands gradually in both directions relative to its location at the first time point. Conversely, a peak

upstream of GPR141 gene contracts towards a narrow genomic interval at the last time point. **(B)**

Schematic workflow of the ChromTime method. During the block finding stage, the input ChIP-seq and,

optionally, control reads are used to determine genomic positions of blocks of signal enrichments across

time points. In the dynamics prediction stage, for each block, peak boundary positions are predicted at

each time point and peak boundary dynamics are predicted at each pair of consecutive time points. **(C)**

Schematic of the dynamics prediction stage for one block. Boxes represent genomic bins at each time

point. The corresponding ChIP-seq signal is depicted as blue bars for each bin whose height represents

the number of reads that map to the bin. During the dynamics prediction stage, the ChromTime method

learns a probabilistic mixture model from the input data in order to partition each block at each time point

into peak and background components. Genomic bins in the peak component (orange) mark ChIP-seq

peaks whereas those in the background component (white) mark flanking background signal. The

movement of the boundaries on the left and the right side of peaks between consecutive time points are

estimated by reasoning jointly about the input data from all time points.

Figure 2: **Sample output from ChromTime with contracting peaks.** Genome browser screenshot with

sample output of ChromTime for H3K4me2 from the T cell development time course in mouse[15] with 5

time points at the Esam/Vsig2/Nrgn locus. The input ChIP-seq signal and MACS2[35] peaks (black boxes

under each signal track) are shown in the upper panel of the screenshot. The ChromTime predicted

peaks colored by their boundary dynamics for each block at each time point are shown in the bottom

panel. The first peak in each block is colored in dark grey. Each subsequent peak is colored with respect

to the predicted dynamic relative to its previous time point. Peaks with steady boundaries on both sides

are shown in light grey, and those with at least one contracting boundary are shown in blue. Nearby

peaks that touch boundaries are visualized as one peak by the genome browser. Not shown in the figure

are expanding peaks, peaks at single time points and peaks with opposite dynamics (Expand on the left

and Contract on the right, or vice versa), which would be colored in red, orange and black, respectively.

See **Fig S2** for examples of predicted expanding peaks.

Figure 3: **Enrichments of TF binding events and DNaseI hypersensitivity sites in ChromTime**

predictions. For data from mouse T cell development[15] and human stem cell reprogramming[21], the

following three sets of ChromTime peaks are examined: 1) peaks present at all time points whose

boundaries hold steady on both sides of the peak throughout the whole time course (T1-Tn Steady); 2)

non-contracting peaks that expand between at least one pair of consecutive time points and have a peak

at the last time point of the time course (Tx-Tn Expand); and 3) non-expanding peaks that contract

between at least one pair of consecutive time points and have a peak at the first time point of the time

course (T1-Tx Contract). The first column in each table shows the percentage of bases covered by each

set out of all bases covered by predicted peaks. The last row shows the baseline percentage of bases

covered by each genomic annotation out of all bases covered by ChromTime peaks at any time point for

each mark. Percentages are color coded from 0 (white) to 100 (green). Fold enrichments in each column

are color coded from 1 (white) to the maximum value in the column (red). **(A)** Fold enrichments of cell

type specific and constitutive peaks of the GATA3 transcription factor, which is a master regulator in T cell

differentiation, in predicted H3K4me2 peaks during T cell development in mouse[15]. **(B)** As in **(A)** for

DNaseI hypersensitive sites, CEBP, POL2 and RAD21 binding in predicted H3K4me3 dynamics during

stem cell reprogramming of fibroblast cells in human[21]. **(C)** As in **(A-B)** for DNaseI hypersensitive sites,

CEBP, P300 in predicted H3K27ac dynamics in stem cell reprogramming in human[21]. In addition,

enrichments are shown for OCT4 and NANOG, which are key transcription factors responsible for

maintaining stem cell identity.

Figure 4: Predicted spatial dynamics by ChromTime associate better with gene expression

changes compared to boundary position changes of peaks from individual time points called in

isolation. For **(A)** H3K4me2 in mouse T cell development[15] and **(B)** H3K4me3 in human stem cell

reprogramming[21], the plots show differences of the average gene expression ranks of block boundaries

ranked based on the length of boundary movement of ChromTime peaks called at individual time points

in isolation and of block boundaries ranked based on ChromTime posterior probabilities computed by

using data from the whole time course (see **Supplementary Methods**) for expansions (left plots) and

contractions (right plots) overlapping annotated TSSs. To rank block boundaries based on the boundary

movements of ChromTime peaks called in isolation, block boundaries were sorted by the difference in the

boundary positions of peaks called in isolation between consecutive time points in descending order for

expansions and in ascending order for contractions. Positive values in the plots correspond to ranks for

which rankings based on ChromTime posterior probabilities associate on average with larger gene

expression changes compared to rankings based on boundary movements of peaks called in isolation

between consecutive time points, and vice versa for negative values. The black lines and the shaded

regions show the difference of the average ranks for a set of randomized rankings of the block

boundaries and their 95% confidence intervals, respectively.

Figure 5: **Spatial dynamics can contain additional information about gene expression changes**

beyond ChIP-seq signal density changes. Gene expression change is plotted as function of ChIP-seq

signal density change after loess smoothing for each predicted ChromTime dynamic for **(A)** H3K4me2

dynamics in T cell development in mouse[15]; and **(B)** in H3K4me3 dynamics in stem cell reprogramming

in human[21]. Peaks of each type of dynamics were pooled from all time points for this analysis. Peaks

with asymmetric dynamics E/S and S/E were pooled together in the "E-S" group. Similarly C/S and S/C

peaks were pooled in the "C-S" group. In both systems, peaks with the same signal density change can

associate with different gene expression changes depending on the predicted spatial dynamic. Shaded

regions represent 95% confidence intervals.

Figure 6: Spatial dynamics of multiple different HMs co-localize within a time course. Hierarchical

clustering with optimal leaf ordering[60] of the geometric average fold enrichments taken across all time

points of the overlap of every pair of predicted spatial dynamics for mapped HMs in **(A)** mouse

adipogenesis[16] and **(B)** human stem cell reprogramming[21]. At each pair of time points, "Expand" and

"Contract" dynamics are defined as all peaks that are predicted as either unidirectional or bidirectional

expansions and contractions, respectively, whereas "Steady" dynamics are defined as all peaks that have

predicted steady boundaries at both sides. Peaks with opposing dynamics at each side (i.e. E/C and C/E)

were excluded from this analysis. In both datasets, expansions, contractions and steady peaks of

H3K4me1, H3K4me2, H3K4me3 and H3K27ac tend to cluster together within each of the three classes,

whereas spatial dynamics of H3K27me3 and H3K36me3 peaks tend to occupy different locations. All

enrichments were capped at 50.

Figure 7: **Direction of asymmetric dynamics correlates with direction of transcription. (A) (i)** Left

panel shows a schematic representation of unidirectional expansions that expand in the same direction

as transcription and in the opposite direction of transcription. The adjacent plots show, for each mark, the

average $\log_2$ ratio between the fraction of unidirectional expansions that expand in the same directions as

transcription of the nearest gene and the fraction of unidirectional expansions that expand in the opposite

direction of transcription of the nearest gene for blocks that are within 1 kb of annotated TSSs and for

more distal blocks. Positive values correspond to enrichment of unidirectional expansions in the same

direction as transcription. For marks mapped in at least six time courses, a black line is plotted

representing the average across all data sets and significant differences are denoted with asterisks based

on a two-tailed Mann-Whitney test at a p-value threshold of 0.05. **(ii)** Left panel shows analogous

schematic for unidirectional contractions. Likewise, adjacent plots show, for each mark, the average ratio

between the fraction of unidirectional contractions that contract in the opposite direction of transcription of

the nearest TSS and unidirectional contractions that contract in the same direction as transcription of the

nearest TSS. **(B)** Left panel shows an example of unidirectional expansions between pairs of time points

that expand in the same direction as transcription at the Hs6st1 gene of the H3K4me2 mark in the T cell

development dataset[15]. Right panel shows an example of unidirectional contractions between pairs of

time points that contract in the opposite direction of transcription at the DNMT3B gene of the H3K4me2

mark in the T cell development dataset. The predicted ChromTime peaks colored by their boundary

dynamics are shown under the signal track for each time point.

System	Histone Marks	Species	# Time Points	Reference
Adipogenesis	H3K4me2 H3K4me3 H3K27ac H3K4me1 H3K36me3 H3K27me3	Mouse Human	4	[16]
Blood Formation	H3K4me2 H3K4me3 H3K27ac H3K4me1	Mouse	5-7	[17]
Cardiac Development	H3K4me3 H3K27ac H3K4me1 H3K27me3 H3K36me3 Pol2	Mouse	4	[20]
Cardiac Development	H3K4me3 H3K27me3 H3K36me3	Human	5	[19]
Embryogenesis	H3K4me3 H3K27ac H3K4me1	Zebrafish	4	[25]
Macrophage Differentiation	H3K4me3	Mouse	5	[18]

	H3K9ac			
	H3K27ac			
	H3K27me3			
Neural Differentiation	H3K4me3	Human	5	[10]
	H3K27ac			
	H3K27me3			
	H3K4me1			
Stem Cell Reprogramming	H3K4me2	Human	4-6	[21]
	H3K4me3			
	H3K27ac			
	H3K4me1			
	H3K27me3			
	H3K36me3			
Stem Cell Reprogramming	H3K4me2	Mouse	4	[24]
	H3K4me3			
	H3K9ac			
	H3K27ac			
	H3K27me3			
	H3K36me3			
	H3K4me1			
	H3K79me2			
	H3K9me3			
Stem Cell Reprogramming	H3K4me3	Mouse	9	[22]
	H3K27ac			
	H3K4me1			
	H3K27me3			
	Pol2			
T Cell Development	H3K4me2	Mouse	5	[15]

	H3K(9,14)ac H3K27me3			

Figure 1

Figure 2

Figure 3

A.

Dynamic	% all block bases	FLDN1-ThyDP shared GATA3	ThyDP-specific GATA3	FLDN1-specific GATA3
T1-Tn Steady	28	1.2	1.1	0.5
Tx-Tn Expand	13	0.6	3.4	0.2
T1-Tx Contract	55	1	0.3	1.5
Base %		0.1	0.2	0.1

B.

Dynamic	% all block bases	H1-IMR90 shared DNaseI	H1-IMR90 shared CEBPB	H1-IMR90 shared POL2	H1-IMR90 shared RAD21	H1-specific DNaseI	H1-specific CEBPB	H1-specific POL2	H1-specific RAD21	IMR90-specific CEBPB	IMR90-specific POL2	IMR90-specific RAD21	IMR90-specific DNaseI
T1-Tn Steady	26	1.5	1.5	1.8	1.3	0.6	0.9	1.2	0.9	1.2	1.2	1.2	0.8
Tx-Tn Expand	38	0.7	0.7	0.4	0.9	1.9	1.6	1.3	1.6	0.4	0.4	0.5	0.3
T1-Tx Contract	22	1.2	1.1	1.3	1	0.3	0.4	0.5	0.4	1.9	1.9	1.8	2.4
Base %		18	0.7	4.5	1.4	11	0.3	3	0.9	1.7	4.2	1	3.1

C.

Dynamic	% all block bases	H1-IMR90 shared DNaseI	H1-IMR90 shared CEBPB	H1-IMR90 shared P300	H1-specific DNaseI	H1-specific CEBPB	H1-specific P300	H1 NANOG	H1 OCT4	IMR90-specific DNaseI	IMR90-specific CEBPB	IMR90-specific P300
T1-Tn Steady	2	3.6	3.7	1.4	0.8	1.2	1.3	1.6	1.3	0.6	1.2	1.7
Tx-Tn Expand	14	1.4	1.4	1	3	3.5	2	3.6	3.8	0.2	0.3	0.3
T1-Tx Contract	26	0.5	0.7	1	0.2	0.2	0.3	0.2	0.2	2.3	2	2.2
Base %		10	0.5	31	4.9	0.2	7.9	0.4	0.3	7.8	2.5	1.5

Figure 4

A.
H3K4me2 in mouse T cell development

B.
H3K4me3 in human stem cell reprogramming

Figure 5

A.

B.

Figure 6

A.

	H3K27ME3 Steady	H3K4ME3 Expand	H3K27AC Expand	H3K4ME2 Expand	H3K4ME1 Expand	H3K4ME1 Contract	H3K27AC Contract	H3K4ME2 Contract	H3K4ME3 Contract	H3K4ME1 Steady	H3K4ME2 Steady	H3K27AC Steady	H3K4ME3 Steady	H3K27ME3 Contract	H3K36ME3 Steady	H3K36ME3 Contract	H3K36ME3 Expand	H3K27ME3 Expand	
	0	0	0	0	0	0	0	1	2	0	1	0	0	0	0	0	0	50	H3K27ME3 Expand
	0	3	4	5	5	1	1	1	0	2	1	1	1	1	0	0	50	0	H3K36ME3 Expand
	0	0	2	2	2	3	5	4	7	4	3	3	1	0	0	50	0	0	H3K36ME3 Contract
	0	3	2	2	3	1	1	1	1	2	2	1	1	0	11	0	0	0	H3K36ME3 Steady
	0	3	4	4	3	0	0	0	0	1	1	0	1	50	0	0	1	0	H3K27ME3 Contract
	0	0	11	13	3	3	14	11	0	6	34	50	50	1	1	1	1	0	H3K4ME3 Steady
	0	25	0	11	4	3	0	15	33	7	32	50	50	0	1	3	1	0	H3K27AC Steady
	0	20	16	0	4	3	18	0	27	8	44	32	34	1	2	3	1	1	H3K4ME2 Steady
	0	5	7	6	0	0	8	8	7	11	8	7	6	1	2	4	2	0	H3K4ME1 Steady
	0	0	1	1	1	10	50	50	50	7	27	33	0	0	1	7	0	2	H3K4ME3 Contract
	0	1	2	0	1	12	50	50	50	8	0	15	11	0	1	4	1	1	H3K4ME2 Contract
	0	4	0	1	2	12	50	50	50	8	18	0	14	0	1	5	1	0	H3K27AC Contract
	0	1	1	0	0	50	12	12	10	0	3	3	3	0	1	3	1	0	H3K4ME1 Contract
	0	11	13	16	50	0	2	1	1	0	4	4	3	3	3	2	5	0	H3K4ME1 Expand
	0	50	50	50	16	0	1	0	1	6	0	11	13	4	2	2	5	0	H3K4ME2 Expand
	0	50	50	50	13	1	0	2	1	7	16	0	11	4	2	2	4	0	H3K27AC Expand
	0	50	50	50	11	1	4	1	0	5	20	25	0	3	3	0	3	0	H3K4ME3 Expand
	16	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	H3K27ME3 Steady

B.

	H3K4ME2 Expand	H3K4ME3 Expand	H3K27AC Expand	H3K4ME1 Expand	H3K4ME1 Contract	H3K4ME2 Contract	H3K27AC Contract	H3K4ME3 Contract	H3K4ME1 Steady	H3K4ME2 Steady	H3K27AC Steady	H3K4ME3 Steady	H3K27ME3 Contract	H3K27ME3 Steady	H3K36ME3 Contract	H3K27ME3 Expand	H3K36ME3 Steady	H3K36ME3 Expand	
	3	5	2	3	1	1	1	0	1	1	1	0	1	1	0	0	0	50	H3K36ME3 Expand
	1	1	1	1	1	1	1	1	2	1	2	1	0	0	0	0	23	0	H3K36ME3 Steady
	2	7	1	3	1	2	2	7	2	3	1	6	0	0	0	50	0	0	H3K27ME3 Expand
	2	1	2	1	2	2	2	4	2	1	2	1	0	0	50	0	0	0	H3K36ME3 Contract
	2	10	1	3	0	0	1	2	2	3	1	4	0	50	0	0	0	1	H3K27ME3 Steady
	4	7	2	3	0	0	0	1	1	2	0	1	50	0	0	0	0	1	H3K27ME3 Contract
	8	0	9	3	3	7	11	0	11	25	38	50	1	4	1	6	1	0	H3K4ME3 Steady
	8	14	0	3	3	8	0	28	12	22	50	38	0	1	2	1	2	1	H3K27AC Steady
	0	17	10	3	3	0	12	19	10	34	22	25	2	3	1	3	1	1	H3K4ME2 Steady
	7	8	8	0	0	6	8	10	19	10	12	11	1	2	2	2	2	1	H3K4ME1 Steady
	3	0	2	0	6	23	40	50	10	19	28	0	1	2	4	7	1	0	H3K4ME3 Contract
	3	4	0	2	5	16	50	40	8	12	0	11	0	1	2	2	1	1	H3K27AC Contract
	0	2	3	1	7	50	16	23	6	0	8	7	0	0	2	2	1	1	H3K4ME2 Contract
	1	1	2	0	15	7	5	6	0	3	3	3	0	0	2	1	1	1	H3K4ME1 Contract
	10	13	11	50	0	1	2	0	0	3	3	3	3	3	1	3	1	3	H3K4ME1 Expand
	15	24	50	11	2	3	0	2	8	10	0	9	2	1	2	1	1	2	H3K27AC Expand
	36	50	24	13	1	2	4	0	8	17	14	0	7	10	1	7	1	5	H3K4ME3 Expand
	50	36	15	10	1	0	3	3	7	0	8	8	4	2	2	2	1	3	H3K4ME2 Expand

Color Key

0 10 20 30 40 50
Value

Figure 7