Modeling Spatio-temporal Dynamics of Chromatin Marks

Petko Fiziev[1,2,3], Jason Ernst[1,2,3,4,5,6,*]

[1] Bioinformatics Interdepartmental Program, University of California, Los Angeles

[2] Departments of Biological Chemistry, University of California, Los Angeles

[3] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at UCLA

[4] Computer Science Department, University of California, Los Angeles

[5] Jonsson Comprehensive Cancer Center, University of California, Los Angeles

[6] Molecular Biology Institute, University of California, Los Angeles, California, USA

* To whom correspondence should be addressed. Email: jason.ernst@ucla.edu

ABSTRACT

To model spatial changes of chromatin mark peaks over time we developed and applied ChromTime, a computational method that predicts regions for which peaks either expand or contract significantly or hold steady between time points. Predicted expanding and contracting peaks can mark regulatory regions associated with transcription factor binding and gene expression changes. Spatial dynamics of peaks provided information about gene expression changes beyond localized signal density changes. ChromTime detected asymmetric expansions and contractions, which for some marks associated with the direction of transcription. ChromTime facilitates the analysis of time course chromatin data in a range of biological systems.

KEYWORDS: epigenomics, time course, spatial dynamics, histone modifications, chromatin marks

BACKGROUND

Genome-wide mapping of histone modifications (HMs) and related chromatin marks using chromatin

immunoprecipitation coupled with high-throughput sequencing (ChIP-seq) and DNA accessibility through

assays for DNase I hypersensitivity[1] (DNase-seq) or transposase-accessible chromatin[2] (ATAC-seq)

assays have emerged as a powerful approach to annotate genomes and study cell states[3–5]. Through

the efforts of large consortia projects such as the ENCODE[6], Roadmap Epigenomics[7] and

BLUEPRINT[8] as well as individual labs[9–11] multiple different chromatin marks have been mapped

across more than a hundred different cell and tissue types. These maps have yielded numerous insights

into gene regulation and genetic and epigenetic association with disease[12–16].

While many mapping efforts have largely focused on single or unrelated cell and tissue types[3, 6], a

growing number of biological processes have been studied with temporal epigenomic data using assays

such as ChIP-seq, ATAC-seq or DNase-seq in a time course that map chromatin marks at consecutive

stages during particular biological processes. Such datasets have been generated for a wide range of

biological settings including T cell development[17], adipogenesis[18], hematopoiesis[19, 20],

macrophage differentiation[21], neural differentiation[12], cardiac development[22, 23], somatic cell

reprogramming[24–27], embryogenesis[28] and many others[7, 29–37]. The output of these experiments

presents a unique opportunity to study the spatio-temporal changes of epigenetic peaks and associated

regulatory elements. However, almost all computational methods designed or applied to epigenomic data

have been developed based on single or multiple unrelated samples. For example, continuous regions of

enrichments of single marks are detected by peak or domain calling methods[38–42]. In cases when

multiple chromatin marks are mapped in the same cell type, methods such as ChromHMM[43] and

Segway[44] can be used to produce genome-wide chromatin state annotations. In addition, methods

have been developed for pairwise comparisons of ChIP-seq signal data by differential peak calling[45,

46].

In the context of time course chromatin data, only a few methods have been proposed that consider

temporal dependencies between samples. One such method, TreeHMM[47], produces a chromatin state

genome annotation similar to ChromHMM and Segway, while taking into account a tree-like structure that

captures lineage relationships between the input cell types in order to potentially derive a more consistent

annotation across samples. Another method, GATE[30], produces a genome annotation based on

clustering fixed length genomic loci that can be modeled with the same switch from one chromatin state

to another over time.

One important limitation of methods for pairwise comparison or time course modeling of chromatin data is

that they do not directly consider or model spatial changes in the genomic territory occupied by chromatin

marks over time. Spatial properties of genomic peaks continuously marked by HMs have gained

increasing attention as a potentially important characteristic of chromatin marks. For example, long peaks

of H3K27ac have been associated with active cell type specific locus control regions termed super-

enhancers or stretch enhancers in a number of cell types[48, 49]. Also, the length of H3K4me3 peaks has

been associated with transcriptional elongation and consistency of cell identity genes[50]. In the context

of cancer, long H3K4me3 peaks have been linked to transcriptional elongation and enhancer activity at

tumor suppressor genes and have been observed to be significantly shortened in tumor cells[51]. Long

H3K4me3 domains have been implicated to mark loci involved in psychiatric disorders[52]. Expanded

domains of H3K27me3 and H3K9me3 marks have been shown to be characteristic of terminally

differentiated cells compared to stem cells[53]. These studies suggest that length of epigenetic peaks is a

dynamic feature that can correlate with activity of putative functional elements regulating specific genes.

Computational methods that do not explicitly reason about the spatial changes of chromatin marks have

significant limitations for studying the dynamics of these properties because they are unable to detect

territorial changes that might be associated with redistribution of signal or identify asymmetric directional

peak boundary movements.

In this work, we present ChromTime, a novel computational method for detection of expanding,

contracting and steady peaks, which can detect patterns of changes in the genomic territory occupied by

chromatin mark peaks from time course sequencing data (**Fig 1A**). We applied ChromTime to a diverse

set of data from different developmental, differentiation and reprogramming time courses. Predicted

expansions and contractions in general mark regulatory regions associated with changes in transcription

factor (TF) binding or gene expression. ChromTime enables studying the directionality of spatial dynamics

of chromatin mark peaks relative to other genomic features, which existing computational approaches do

not directly address. Our results show that the direction of predicted expansions and contractions

correlates with direction of transcription near transcription start sites (TSSs). ChromTime is a general

method that can be used to analyze time course chromatin data from high-throughput sequencing assays

such as from ChIP-seq, ATAC-seq and DNase-seq for a wide range of biological systems to gain insights

into the dynamics of gene regulation.

RESULTS

Model for detecting expanding, contracting and steady peaks from temporal chromatin data

We developed a computational method, ChromTime (https://github.com/ernstlab/ChromTime), designed

for systematic detection of expansions, contractions and steady peaks from time course chromatin data

for single chromatin features (**Methods, Fig 1B**). ChromTime takes as input a set of genomic coordinates

of aligned sequencing reads from foreground and, optionally, control experiments over the time course.

The foreground experiments are data from a chromatin sequencing assay such as ChIP-seq, ATAC-seq

or DNase-seq performed at a series of time points. The method consists of two stages – block finding and

dynamics prediction. During the block finding stage, ChromTime determines continuous genomic regions

(blocks) that may contain peaks of foreground signal enrichment during the time course (**Fig S1A-B**). To

achieve this, ChromTime partitions the genome into fixed length bins and counts the number of

foreground and control reads that map to each bin at each time point. Nearby bins that show significant

enrichment are joined into continuous intervals, which subsequently are grouped into blocks if they

overlap across time points. As a result, large portions of the genome that are likely to contain background

noise at all time points are filtered out, so that peak boundary dynamics are determined within a subset of

the genome potentially enriched for the chromatin mark.

During the dynamics prediction stage, for each block, ChromTime determines the most likely positions of

the peak boundaries at each time point and whether the peak expands, contracts or holds steady at each

boundary between consecutive time points. The method uses a probabilistic mixture model to partition the

signal within each block at each time point into background and peak components (**Fig 1C, S1C**) by

reasoning jointly about the data from all time points in the time course. The method assumes that central

positions in blocks are more likely to be enriched for foreground reads and thus the peak component is

flanked by the background components (**Fig S1D**). The number of sequencing reads in bins from each

component at each time point is modelled with different negative binomial distributions that can account

for the local abundance of control reads. Furthermore, between any two consecutive time points the

boundaries of the peaks are assumed to follow one of three possible dynamics: steady, expand or

contract. For steady dynamics, the peak boundaries are enforced to have the same genomic position. For

expanding and contracting dynamics, the number of genomic bins that the peak boundaries move

between the two time points is modelled with different negative binomial distributions which depend on

the pair of time points and the corresponding dynamic. ChromTime models time points that have no bins

in the peak component with zero length peaks. Thus, appearances of peaks, except at the first time point,

are modeled as expansions from zero length peaks and the disappearances of peaks are modeled as

contractions to zero length peaks. Each dynamic is also assumed to have a prior probability which

captures information about its genome-wide frequency at each time point.

All model parameters are learned jointly from the whole time course. As a result, ChromTime can adapt to

different boundary movements, dynamics frequencies and noise levels across experiments and biological

systems. The estimated parameters are used to make a prediction for each block for the most likely

positions of the peak boundaries and the corresponding boundary dynamics that had generated the

signal within the block. The final output contains predicted peak boundaries annotated and colored by

their assigned dynamics, which can be used for downstream analysis with existing tools and visualized in

genome browsers (**Fig 2, S2,** https://github.com/ernstlab/ChromTime).

Reproducibility of ChromTime predictions and association with changes in gene expression, TF binding and DNaseI hypersensitivity sites

To investigate the reproducibility of ChromTime predictions, we applied ChromTime separately to two biological replicate datasets for the H3K4me2 and the H3K(9/14)ac marks in T cell development in mouse[17] and confirmed on average strong enrichment for the same ChromTime annotations co-localizing across replicates (**Fig S3**). We then applied the method to data from pooled replicates for the H3K4me2 mark from the mouse T cell development study[17], to data for the H3K4me3 and H3K27ac marks from a study on stem cell reprogramming in human[24], to ATAC-seq data from a mouse stem cell reprogramming time course[27] and to a human fetal brain development time course that we constructed from DNase-seq datasets from Roadmap Epigenomics[7]. To investigate the biological relevance of ChromTime predictions, for blocks overlapping TSSs we examined changes in the corresponding gene expression. Peaks with predicted expanding and contracting boundaries that overlap annotated TSSs associated with increase and decrease, respectively, in gene expression (**Fig 3, S4**). Additionally for all chromatin marks, we examined enrichments of TF binding sites across all blocks[6, 17, 27], and in the case of HMs, also enrichments of DNaseI hypersensitivity sites (DHSs)[7]. Predicted peaks with expanding and contracting boundaries enriched for sites bound by important regulators in each biological system as well as sites bound by generic TFs in a cell type specific manner. Expanding and contracting HM peaks also enriched for cell type specific DHSs. Furthermore, peaks with predicted steady boundaries showed enrichment for TF binding sites that are shared between the first and the last time point in the corresponding time courses, which mark potentially stable regulatory elements. Similar enrichments in the case of HM peaks were also seen for shared DHSs.

Predicted spatial dynamics by ChromTime associate better with gene expression changes compared to boundary position changes of peaks called from individual time points in isolation

We next investigated whether ChromTime's approach for reasoning jointly about the whole time course increases power to detect associations with gene expression compared to considering boundary

differences of peaks at consecutive time points called in isolation. Specifically, we analyzed gene

expression changes of genes with TSSs overlapping ChromTime peaks in relation to posterior

probabilities for expansions and contractions compared to boundary differences of peaks called with

ChromTime from data from individual time points in isolation. We investigated this in the context of

H3K4me2 peaks in mouse T cell development[17] and for H3K4me3 peaks in stem cell reprogramming in

human[24]. In most cases, ranking boundary changes of peaks in blocks with at least one non-zero length

peak by their predicted ChromTime posterior probabilities for expansions and contractions associated on

average with larger gene expression changes compared to ranking boundaries directly based on the

change in the genomic positions of the boundaries of ChromTime peaks called at individual time points in

isolation (**Fig 4, S5A**). These results also held when using peaks from two different peak callers,

MACS2[38] and SICER[40] applied on data from individual time points (**Fig S5B-C**).

Spatial dynamics contain information about gene expression changes between consecutive time

points not captured by corresponding pairwise signal density changes

We next investigated whether there is additional information in ChromTime predictions beyond what can

be captured by pairwise signal density changes or by differential peak calls. For this analysis, we focused

on H3K4me2 in mouse T cell development[17] and H3K4me3 in human stem cell reprogramming[24]. For

pairs of consecutive time points, we computed the change in signal density in the region starting at the

left most and ending at the right most predicted peak boundary in the block (**Supplementary Methods**).

We associated the signal density changes with gene expression changes at the nearest TSS within 50kb

of each block and computed the average gene expression change as a function of the signal density

change within blocks (**Fig 5**). We found that locations with the same signal density change can associate

with significantly different average gene expression changes of proximal genes depending on the

predicted ChromTime dynamics. Notably, bidirectional expansions, expansions occurring on both sides of

a peak, associated for a range of signal density changes with greater average increase in gene

expression than unidirectional expansions, those expansions occurring on one side with steady on the

other, when controlling for the signal density change. These unidirectional expansions in turn associated

for a range of signal density changes with greater expression change than steady regions, those regions

with a steady call on both sides of a peak, when controlling for the signal density change. We observed a

similar relationship for contractions and decrease of gene expression. These results were replicated also

after substituting ChIP-seq signal density changes with differential peak scores from two differential peak

calling methods, SICER[40] and MACS2[38] (**Fig S6A-B**). Therefore, ChromTime predictions can provide

additional information about gene expression changes beyond what is contained in the corresponding

signal density changes as measured directly or by utilizing differential peak calling procedures.

Spatial dynamics are correlated between multiple chromatin marks

Previous studies have shown that the locations of different chromatin marks can be correlated[3, 54]. In

this context, we tested whether multiple chromatin marks can also exhibit jointly the same type of spatio-

temporal dynamics. For this purpose, we compared the genomic locations of predicted expansions,

contractions and steady peaks for different chromatin marks within the same time course. We focused on

three previously published time courses – stem cell reprogramming in human[24], stem cell

reprogramming in mouse[27] and adipogenesis in mouse[18], where multiple chromatin marks were

mapped (**Fig 6, S7**). In all three datasets, we observed that predicted expansions co-localized

preferentially for H3K4me2, H3K4me3 and H3K27ac and to a lesser extent for H3K4me1 and similarly for

predicted contractions and steady peaks. In contrast, different predicted spatial dynamics for H3K36me3

and H3K27me3 tended to occupy distinct locations. In addition, in mouse reprogramming[27], ChromTime

predicted dynamics of ATAC-seq, H3K4me2, H3K4me3, H3K27ac, H3K9ac and to a lesser extent of

H3K4me1 and H3K79me2 peaks co-localized preferentially (**Fig S7**). These results suggest that spatial

dynamics of chromatin marks are coordinated at least at a subset of genomic locations.

Direction of expansions and contractions is correlated with direction of transcription

ChromTime can predict unidirectional expansions and contractions, which enables analysis of

directionality of spatial dynamics of peaks, an aspect of chromatin regulation that has not been previously

systematically explored. To investigate this, we applied ChromTime on data from 13 previously published studies from a variety of developmental, differentiation and reprogramming processes (**Table 1**) for nine different HMs including narrow and broad marks, for Pol2, ATAC-seq and DNase-seq. We observed that unidirectional expansions and contractions are predicted in most cases on average to be the majority of all expansions and contractions, respectively, at a given pair of consecutive time points (**Fig S8**). One hypothesis for the prevalence of asymmetric boundary movements for the promoter associated chromatin marks is that the direction of boundary movements is associated with the asymmetry of transcription initiation in promoter regions. To test this hypothesis, for each dataset we compared the prevalence of each class of unidirectional dynamics as a function of its distance to the nearest annotated TSS and the orientation of the corresponding gene (**Fig 7**). Consistent with our hypothesis, for H3K4me3, H3K4me2, H3K(9/14)ac, H3K79me2, and for Pol2, we found that unidirectional expansions that expand into the gene body (i.e. in the same direction as transcription) were substantially more frequently found in proximity of TSSs compared to unidirectional expansions in the opposite direction. Moreover, this difference was not observed for expansions that are distal from TSSs. Similarly, in most cases for these data unidirectional contractions that contract towards the TSS of the nearest gene (i.e. in the opposite direction of transcription) were substantially more frequent compared to unidirectional contraction in the opposite direction in proximity of TSSs, whereas their frequencies at distal sites showed much smaller differences. HMs H3K27ac, H3K4me1 and H3K27me3, some ATAC-seq datasets and peaks in the DNase-seq time course exhibited the same trend, but to a smaller degree.

DISCUSSION

In this work, we presented ChromTime, a novel computational method for systematic detection of expanding, contracting and steady peaks of chromatin marks from time course high-throughput sequencing data. ChromTime employs a probabilistic graphical model that directly models changes in the genomic territory occupied by single chromatin marks over time. This approach allowed us to directly encode our modeling assumptions about dependencies between variables in an interpretable and extendable framework.

We applied ChromTime on ChIP-seq data for broad and narrow HMs and for Pol2, and on ATAC-seq and

DNase-seq data from a variety of developmental, differentiation and reprogramming courses. Our results

show that the method can identify sets of expanding and contracting peaks that are biologically relevant

to the corresponding systems. In particular, expansions and contractions associate with up- and down-

regulation of gene expression and differential TF binding, supporting the biological relevance of

ChromTime predictions.

ChromTime gains power by both reasoning jointly about all time points in a time course and by explicitly

modeling the peak boundary movements. Supporting this, in our analyses we observed that territorial

changes identified by ChromTime had better agreement with gene expression changes compared to

considering directly the boundary change of peaks called on data from individual time points in isolation.

Additionally, we also observed for a range of cases that expanding and contracting peaks associated on

average with greater change in gene expression compared to peaks with steady boundaries even after

controlling for signal density changes. Some of the power that ChromTime gains from considering spatial

information might be explained by its ability to differentiate territorial expansions or contractions which can

reflect changes in the number sites of TF binding in close vicinity from changes in signal density within

steady peak boundaries. Changes in signal density without territorial expansions or contractions might

reflect a change in the proportion of cells with the chromatin mark without large changes in activity in any

one cell. Additional power can come from the temporal and spatial information that allows the model to

effectively smooth over noise in the data, thus enabling more biologically relevant inferences.

ChromTime enables novel analysis of directionality of spatial epigenetic dynamics. In this context, we

found that asymmetric unidirectional expansions and contractions for several marks correlate strongly

with direction of transcription in promoter proximal regions, which suggests that spatial dynamics at such

locations may be related to actions of the transcriptional machinery. One possible explanation for the

observed correlation between the direction of spatial dynamics of at least some HMs and transcription

can be provided in part by previous studies that have shown that the Pol2 elongation machinery can

recruit H3K4-methyltransferases such as members of the SET[55] and MLL[56] families at the promoters

of genes. Our findings are consistent with such models where the Pol2 complex itself may be facilitating

the attachment and removal of these marks[57].

The ChromTime software is also relatively efficient particularly when using its option to parallelize all

computations during the parameter learning and prediction phases over multiple CPU cores. In our tests,

processing ChIP-seq data for the H3K4me2 mark and control data from 5 time points in mouse T cell

development[17] took 3 hours on a laptop computer using 4 CPU cores.

We applied ChromTime to a range of data types, however we found no single setting of the method

options to be preferred in all cases (**Methods**). We thus created three modes with different default

options: punctate mode used for ATAC-seq and DNase-seq, narrow mode used for ChIP-seq of narrow

HMs, and broad mode used for ChIP-seq of broad HMs and Pol2. In principle, ChromTime can also be

applied on ChIP-seq data of sequence specific TFs in punctate mode. However for this data, where

binding can often be associated with a single point source such as individual instances of DNA sequence

regulatory motifs, methods that predict the single point source across time points and the binding intensity

associated with the source at each time point may be a more natural way to model the data.

Another limitation of the ChromTime method is that while the runtime of ChromTime still scales linearly

with the number of time points, T, the number of observed combinations of dynamics can scale

exponentially with T. This exponential growth can complicate downstream analyses that directly consider

each combination of dynamics, as there will be 3^{T-1} possible sequences of dynamics at each side of a

peak. Extensions of the ChromTime model could model the large number of combinations as being

instances of a smaller number of more distinct dynamic patterns.

CONCLUSIONS

The increasing availability of time course chromatin data provides an opportunity to understand chromatin

dynamics in many biological systems. To facilitate reaching this goal we developed ChromTime, which

systematically detects expanding, contracting, and steady peaks allowing extraction of additional

information from this data. ChromTime gains power by both reasoning about data from all time points in

the time course and by explicitly modeling movements of peak boundaries. We showed that ChromTime

predictions associate with relevant genomic features such as changes in gene expression and TF

binding. We demonstrated that territorial changes of peaks can contain additional information beyond

signal density changes with respect to gene expression of proximal genes. ChromTime allows for novel

analysis of directionality of spatial dynamics of chromatin marks. In this context, we showed for multiple

chromatin marks that the direction of predicted asymmetric expansions and contractions of peaks strongly

associates with direction of transcription in proximity of TSSs. ChromTime is generally applicable to

modeling time courses of chromatin marks, and thus should be a useful tool to gaining insights into

dynamics of epigenetic gene regulation in a range of biological systems.

METHODS

Overview of the ChromTime method

ChromTime takes as input a set of files in BED format with genomic coordinates of aligned sequencing

reads from experiments for a single chromatin mark from a high-throughput sequencing experiment such

as ChIP-seq, ATAC-seq or DNase-seq over a time course and, optionally, from a set of control

experiments. ChromTime consists of two stages (**Fig 1B-C**):

1) Detecting genomic intervals (blocks) potentially containing regions of signal enrichment (peaks)

2) Learning a probabilistic mixture model for boundary dynamics of peaks within blocks throughout

the time course and computing the most likely spatial dynamic and peak boundaries for each

block throughout the whole time course

Detecting genomic blocks containing regions of signal enrichment

The aim of this stage is to determine approximately the genomic coordinates of regions with potential

peaks of signal enrichment at any time point in the time course (**Fig S1A-B**). The signal within these

blocks will be used as input to build the mixture model in the next stage of ChromTime. ChromTime

supports analysis of punctate, narrow and broad marks in three different modes, which are defined by

different default options. The method partitions the genome into non-overlapping bins of predefined

length, BIN_SIZE (by default, 200 bp in narrow and punctate modes, 500 bp in broad mode) and counts

for each bin and time point the number of sequencing reads whose alignment starting positions after

shifting by a predefined number of bases (SHIFT, 100 bp in the direction of alignment by default) are

within its boundaries. Next, each bin at each time point is tested for enrichment based on a Poisson

background distribution at a predefined false discovery rate (5% by default). The expected number of

reads for a bin at position p and time point t, $\lambda_{t,p}$, in the Poisson test is computed conservatively as the

maximum of:

1) If control reads are provided: for each of windows of size w=1,000bp, 5,000bp and 20,000bp the

average number of control reads in the window centered at the current bin, normalized by the

ratio of total reads in the foreground and control experiments, that is:

352
$$\lambda_{t,p,w} = \frac{\text{\# [Total Foreground Reads]}}{\text{\# [Total Control Reads]}} \frac{\text{BIN\_SIZE}}{w} \text{Ctrl}_{t,p,w}$$

where $\text{Ctrl}_{t,p,w}$ is the total number of control reads in each of window of size w around the bin at

position p at time point t.

2) The average number of foreground reads per genomic bin.

3) 1 read.

Testing multiple different windows sizes for the background is a strategy we adopted from the MACS2

peak caller[38].

Within each time point, consecutive bins that are significantly enriched are merged into continuous

intervals. The intervals are further extended in both directions to include continuous stretches of bins

where each bin is significant based on a Poisson background distribution at a weaker P-value threshold

(0.15 by default). Extended intervals within a predefined number of non-significant bins, MAX_GAP (3

bins by default), are further joined together. This joining strategy has been previously implemented by

other peak callers for single datasets such as SICER[40]. Next, overlapping intervals across time points

are grouped into blocks. To capture more of the potential background signal and to increase the likelihood

that central bins within blocks contain higher foreground signal, the start and end positions of each block

are extended additionally by a predefined number of bins, BLOCK_EXTEND, (5 by default) upstream

from the left-most coordinate and downstream from the right-most coordinate of the intervals in the block,

respectively, or up to the middle point between the current block and its adjacent blocks if they are within

BLOCK_EXTEND bins apart. Restricting BLOCK_EXTEND to a relatively limited number of bins helps in

keeping the running time of the method within reasonable bounds.

In narrow and punctate modes, blocks that contain multiple intervals at the same time point separated by

gaps of non-significant bins longer than MAX_GAP are split into sub-blocks at each gap between those

intervals. In particular, all gaps within a block are intersected across the time points that have gaps. For

each gap intersection, the block is split at the position with the lowest average foreground signal across

all time points. In broad mode, no such splitting is performed in order to avoid excessive peak

fragmentation.

Probabilistic mixture model for boundary dynamics of peaks within blocks across the time course

The foreground and the expected signal within the blocks are used as input to build a probabilistic mixture

model for the boundary dynamics of the peaks within blocks (**Fig S1C**). One core assumption of the

model is that each block contains at each time point exactly one peak, which can potentially have a length

of zero bins. This implies that at each time point, the bins within a block can be partitioned into three

continuous intervals: left-flanking background, foreground peak, and right-flanking background. For the

bin in block i, at time point t and position p, let $O_{i,t,p}$ denote the random variable that models the number

of observed foreground reads, and let $o_{i,t,p}$ denote the corresponding observed read counts. Let $V_{i,t,p}$

denote the random variable for the label of the corresponding bin, which can either have the value PEAK

or BACKGROUND. Let $X_{i,t,p}$ denote a random variable for the vector of covariates for the corresponding

bin, and $x_{i,t,p}$ their corresponding values. The distribution of $O_{i,t,p}$ conditioned on $V_{i,t,p}$ and $X_{i,t,p}$ is modeled

with different negative binomial distributions depending on the value of $V_{i,t,p}$ and $x_{i,t,p}$:

399
$$P\big(O_{i,t,p} = o_{i,t,p}\big|V_{i,t,p} = \text{PEAK}, X_{i,t,p} = x_{i,t,p}\big) = \text{NB}\big(o_{i,t,p}; \mu_{\text{PEAK},i,t,p}, \delta_t\big)$$

400
$$= \frac{\Gamma(o_{i,t,p} + \delta_t)}{o_{i,t,p}!\,\Gamma(\delta_t)}\left(\frac{\delta_t}{\mu_{\text{PEAK},i,t,p} + \delta_t}\right)^{\delta_t}\left(\frac{\mu_{\text{PEAK},i,t,p}}{\mu_{\text{PEAK},i,t,p} + \delta_t}\right)^{o_{i,t,p}}$$

402   and

404
$$P\big(O_{i,t,p} = o_{i,t,p}\big|V_{i,t,p} = \text{BACKGROUND}, X_{i,t,p} = x_{i,t,p}\big) = \text{NB}\big(o_{i,t,p}; \mu_{\text{BACKGROUND},i,t,p}, \delta_t\big)$$

405
$$= \frac{\Gamma(o_{i,t,p} + \delta_t)}{o_{i,t,p}!\,\Gamma(\delta_t)}\left(\frac{\delta_t}{\mu_{\text{BACKGROUND},i,t,p} + \delta_t}\right)^{\delta_t}\left(\frac{\mu_{\text{BACKGROUND},i,t,p}}{\mu_{\text{BACKGROUND},i,t,p} + \delta_t}\right)^{o_{i,t,p}}$$

407   where δ_t is the dispersion parameter. Similarly to negative binomial regression models[58], ChromTime

408   models the mean of each component through the log link as a linear combination of a two-dimensional

409   vector of covariates, $x_{i,t,p} = (1, \log \lambda_{i,t,p})$, which includes a constant term and the logarithm of the

410   expected number of reads in the bin as computed in the previous section, $\lambda_{i,t,p}$:

412
$$\mu_{\text{PEAK},i,t,p} = \exp[\alpha_t + \gamma_t \log \lambda_{i,t,p}]$$

$$\mu_{\text{BACKGROUND},i,t,p} = \exp[\beta_t + \gamma_t \log \lambda_{i,t,p}]$$

where α_t, β_t and γ_t are time point specific scalar parameters. Negative binomial distributions have been

successfully employed in a similar manner to capture the over-dispersion of sequencing reads in peak

callers for single samples such as ZINBA[59]. Of note however, ChromTime requires that the dispersion

parameter δ_t and the coefficient γ_t are shared between the two components at each time point. The first

requirement ensures that the distribution with the smaller mean value has higher probabilities compared

to the distribution with the larger mean value for the lowest values of the support domain of the negative

binomial distribution, and that the opposite holds for the largest values of the support domain

(**Supplementary Methods**). Sharing the dispersion parameter here is analogous to sharing the variance

parameter in Gaussian mixture models. The second requirement to share the γ_t parameter ensures that

the control signal has equal importance in each component.

Formally, let $B_{i,L,t}$ and $B_{i,R,t}$ denote the random variables corresponding to the first and the last bin,

respectively, in the peak partition at time t for block i relative to the beginning of the block, and let N_i be

the length of the block. We then have $1 \le B_{i,L,t} \le N_i + 1$ and $0 \le B_{i,R,t} \le N_i$, with values of $B_{i,L,t} = N_i + 1$

and $B_{i,R,t} = 0$ corresponding to the special cases of starting a peak after all positions and ending a peak

before all positions in a block, respectively. For $B_{i,L,t}$ and $B_{i,R,t}$ to denote valid interval boundaries,

ChromTime also requires that $B_{i,L,t} \le B_{i,R,t} + 1$ at each time point. These constraints can be formally

encoded by introducing one auxiliary binary variable for each time point in the model, $Z_{i,t}$, such that:

$$P(Z_{i,t} = 1 | B_{i,L,t} = l, B_{i,R,t} = r) = \begin{cases} 1 & \text{if } 1 \le l \le r + 1 \le N_i + 1 \\ 0 & \text{otherwise} \end{cases}$$

and thus also

$$P(Z_{i,t} = 0 | B_{i,L,t} = l, B_{i,R,t} = r) = \begin{cases} 0 & \text{if } 1 \le l \le r + 1 \le N_i + 1 \\ 1 & \text{otherwise} \end{cases}$$

ChromTime treats all $Z_{i,t}$ variables as observed with values equal to 1 for all blocks and time points.

The conditional probability of the bin labels, $V_{i,t,p}$, given the peak boundaries, $B_{i,L,t}$ and $B_{i,R,t}$, are defined

to be:

$$443 \qquad P(V_{i,t,p} = \text{PEAK}|B_{i,L,t} = l, B_{i,R,t} = r) = \begin{cases} 1 & \text{if } l \le p \le r \\ 0 & \text{otherwise} \end{cases}$$

444   and thus also

$$445 \qquad P(V_{i,t,p} = \text{BACKGROUND}|B_{i,L,t} = l, B_{i,R,t} = r) = \begin{cases} 0 & \text{if } l \le p \le r \\ 1 & \text{otherwise} \end{cases}$$

447   The probability of the observed read counts at time t, $\boldsymbol{o}_{i,t}$, and $Z_{i,t} = 1$, conditioned on the values of the

448   peak boundaries, $B_{i,L,t}$ and $B_{i,R,t}$, and the covariates at time point t, $\boldsymbol{x}_{i,t}$, under the model is then:

$$450 \qquad P\big(\boldsymbol{O}_{i,t} = \boldsymbol{o}_{i,t}, Z_{i,t} = 1 \big| B_{i,L,t} = l, B_{i,R,t} = r, \boldsymbol{X}_{i,t} = \boldsymbol{x}_{i,t}\big)$$

$$451 \qquad = P\big(Z_{i,t} = 1 | B_{i,L,t} = l, B_{i,R,t} = r\big)$$

$$452 \qquad \times \prod_{p=1}^{l-1} \text{NB}(o_{i,t,p}; \mu_{\text{BACKGROUND},i,t} = \exp[\beta_t + \gamma_t \log \lambda_{i,t,p}], \delta_t)$$

$$453 \qquad \times \prod_{p=l}^{r} \text{NB}\big(o_{i,t,p}; \mu_{\text{PEAK},i,t} = \exp[\alpha_t + \gamma_t \log \lambda_{i,t,p}], \delta_t\big)$$

$$454 \qquad \times \prod_{p=r+1}^{N_i} \text{NB}(o_{i,t,p}; \mu_{\text{BACKGROUND},i,t} = \exp[\beta_t + \gamma_t \log \lambda_{i,t,p}], \delta_t)$$

An important special case of the above formulation when $B_{i,L,t} = B_{i,R,t} + 1$ corresponds to modelling the

whole signal at time point t as background, which enables ChromTime to accommodate time points that

are all background by modelling them with zero length peaks. For this reason, ChromTime blocks

internally have the same number of peak boundaries at all time points even if some time points are

predicted as zero length peaks (i.e. all background). Boundaries of zero length peaks are treated by the

model in the same way as boundaries of non-zero length peaks.

ChromTime assumes uniform prior probabilities for the left and the right end boundaries at the first time

point:

465
$$P(B_{i,L,1} = l) = Unif(1, N_i + 1)$$

466   and

467
$$P(B_{i,R,1} = r) = Unif(0, N_i)$$

where $Unif(a, b)$ denotes the uniform distribution of integer numbers in the closed interval $[a, b]$.

Let $D_{i,s,t}$ denote the dynamic between time points *t* and *t+1* on boundary side *s*, where *s* is one of *L* (left

side) or *R* (right side). Between any two time points the ChromTime model allows for one of three

possible dynamics at both the left and the right end boundaries of a peak: STEADY, EXPAND or

CONTRACT. To capture the change of boundary positions between consecutive time points *t* and *t+1* we

define the quantities $J_{i,L,t} = B_{i,L,t} - B_{i,L,t+1}$ and $J_{i,R,t} = B_{i,R,t+1} - B_{i,R,t}$ corresponding to the left and right

boundaries respectively. Positive values of $J_{i,L,t}$ and $J_{i,R,t}$ indicate the number of bases a peak expanded,

whereas negative values indicate the number of bases a peak contracted, and a value of 0 indicates the

peak held steady on the left and the right side respectively. ChromTime models $J_{i,L,t}$ *and* $J_{i,R,t}$ with

different probability distributions for each of the three dynamics. For STEADY dynamic, ChromTime uses

the Kronecker delta function:

482
$$P(J_{i,s,t} | D_{i,s,t} = \text{STEADY}) = \begin{cases} 1, & \text{if } J_{i,s,t} = 0 \\ 0, & \text{otherwise} \end{cases}$$

484   For expanding and contracting dynamics, ChromTime employs negative binomial distributions to model

485   the number of genomic bins a peak boundary moves relative to the minimal movement of one bin

486   required for peak expansions and contractions:

488
$$P(J_{i,s,t} = j | D_{i,s,t} = \text{EXPAND}) = NB(j - 1; \mu_{\text{EXPAND,t}}, \delta_{\text{EXPAND,t}})$$

489   and

490
$$P\big(J_{i,s,t} = j \big| D_{i,s,t} = \text{CONTRACT}\big) = NB(-j-1; \mu_{\text{CONTRACT,t}}, \delta_{\text{CONTRACT,t}})$$

Furthermore, each distribution is parametrized with a mean and dispersion parameter depending on the

dynamic and the time point, t: $\mu_{\text{EXPAND,t}}, \delta_{\text{EXPAND,t}}$ for expansions, and $\mu_{\text{CONTRACT,t}}, \delta_{\text{CONTRACT,t}}$ for

contractions. Of note, in negative binomial distributions the probabilities for negative integers are defined

to be 0. Therefore, the above parametrization enforces that boundary movements of negative or zero

length (i.e. contracting or steady, respectively) are impossible for expansions and that boundary

movements of positive or zero length (i.e. expanding or steady) are impossible for contractions.

The ChromTime model additionally assumes that there is a prior probability to observe each dynamic

between time points t and $t+1$, $P(D_{i,s,t} = d) = \pi_{t,d}$, which is the same at each side (left and right). Users

have the option to set a minimum prior probability (MIN_PRIOR) for the dynamics for all time points. This

parameter can be used to avoid learning priors too close to zero, which in some cases can occur for more

punctate marks where the short length of the peaks can cause the prior to become a dominant influence

on the class assignment of the spatial dynamics. By default, MIN_PRIOR=0 in narrow and broad modes

and MIN_PRIOR=0.05 in punctate mode.

For a time course with T time points we can express for block i the probability of a particular sequence of

dynamics and boundary positions on the left side ($\boldsymbol{d}_L$ and $\boldsymbol{b}_L$ respectively) and on the right side ($\boldsymbol{d}_R$ and $\boldsymbol{b}_R$

respectively), and observing foreground counts $\boldsymbol{o}_i$ and $\boldsymbol{Z}_i = 1$ conditioned on the values of the covariates,

$\boldsymbol{x}_i$ as:

511
$$P\big(\boldsymbol{D}_{i,L} = \boldsymbol{d}_L, \boldsymbol{B}_{i,L} = \boldsymbol{b}_L, \boldsymbol{D}_{i,R} = \boldsymbol{d}_R, \boldsymbol{B}_{i,R} = \boldsymbol{b}_R, \boldsymbol{O}_i = \boldsymbol{o}_i, \boldsymbol{Z}_i = \boldsymbol{1} \big| \boldsymbol{X}_i = \boldsymbol{x}_i\big) =$$

$$P(B_{i,L,1} = l_1) \times P(B_{i,R,1} = r_1) \times P(\boldsymbol{O}_{i,1} = \boldsymbol{o}_{i,1}, Z_{i,1} = 1 | B_{i,L,1} = l_1, B_{i,R,1} = r_1, \boldsymbol{X}_{i,1} = \boldsymbol{x}_{i,1})$$

$$\times \prod_{t=2}^{T} \left(P(\boldsymbol{O}_{i,t} = \boldsymbol{o}_{i,t}, Z_{i,t} = 1 | B_{i,L,t} = l_t, B_{i,R,t} = r_t, \boldsymbol{X}_{i,t} = \boldsymbol{x}_{i,t}) \right.$$

$$\times P(J_{i,L,t-1} = l_{t-1} - l_t | D_{i,L,t-1} = d_{L,t-1}) \times P(D_{i,L,t-1} = d_{L,t-1})$$

$$\left. \times P(J_{i,R,t-1} = r_t - r_{t-1} | D_{i,R,t-1} = d_{R,t-1}) \times P(D_{i,R,t-1} = d_{R,t-1}) \right)$$

where $\boldsymbol{Z}_i = \boldsymbol{1}$ is used to denote $Z_{i,t} = 1$ for all t, $d_{s,t}$ for t=1,...,*T-1* is the dynamic label for the t^{th} pair of

consecutive time points on the left or the right side ($s = L$ or R), respectively. Also $\boldsymbol{b}_L$ and $\boldsymbol{b}_R$ are the

vectors of *T* boundary positions containing l_t and r_t for t=1,...,*T*, respectively.

The total probability of the signal in a block can be expressed as a sum over all possible sequences of

dynamics and peak boundary positions that can generate the block across all time points. Thus, the

probability of block i having observations $\boldsymbol{o}_i$ and $\boldsymbol{Z}_i = \boldsymbol{1}$ given the covariates $\boldsymbol{x}_i$:

$$P(\boldsymbol{O}_i = \boldsymbol{o}_i, \boldsymbol{Z}_i = \boldsymbol{1} | \boldsymbol{X}_i = \boldsymbol{x}_i) = \sum_{\boldsymbol{d}_L, \boldsymbol{b}_L, \boldsymbol{d}_R, \boldsymbol{b}_R} P(\boldsymbol{D}_{i,L} = \boldsymbol{d}_L, \boldsymbol{B}_{i,L} = \boldsymbol{b}_L, \boldsymbol{D}_{i,R} = \boldsymbol{d}_R, \boldsymbol{B}_{i,R} = \boldsymbol{b}_R, \boldsymbol{O}_i = \boldsymbol{o}_i, \boldsymbol{Z}_i = \boldsymbol{1} | \boldsymbol{X}_i = \boldsymbol{x}_i)$$

where $\boldsymbol{d}_L$ and $\boldsymbol{d}_R$ each iterate over all possible 3^{T-1} combinations of peak boundary dynamics, and

$\boldsymbol{b}_L$ and $\boldsymbol{b}_R$ each iterate over all possible ways to place left and right end boundaries across all time points

that are consistent with the requirements that $1 \le B_{i,L,t} \le B_{i,R,t} + 1 \le N_i + 1$ at each time point.

Let $\boldsymbol{o}$ be the total set of observed read counts in all blocks in the data, $\boldsymbol{x}$ be the set of the corresponding

two-dimensional vectors containing the constant term and the log of the expected number of reads at

each position and time point for each block, $\boldsymbol{Z} = \boldsymbol{1}$ denotes all $\boldsymbol{Z}_i = \boldsymbol{1}$, and *M* be the total number of

blocks. Then, the likelihood of all blocks conditioned on their covariates is:

$$P(\boldsymbol{O} = \boldsymbol{o}, \boldsymbol{Z} = \boldsymbol{1} | \boldsymbol{X} = \boldsymbol{x}) = \prod_{i=1}^{M} P(\boldsymbol{O}_i = \boldsymbol{o}_i, \boldsymbol{Z}_i = \boldsymbol{1} | \boldsymbol{X}_i = \boldsymbol{x}_i)$$

538 We note that in the above formulation allows ChromTime to model the appearance of a peak, if it occurs

539 after the first time point in the time course, as an expansion from a zero length peak at the previous time

540 point. Similarly, the disappearance of a peak is modelled as a contraction to a zero length peak at the

541 next time point.

543 **Model optimization**

545 The total set of parameters of the model consists of:

546  *1)* Prior probabilities of each dynamic d at each time point t: $\pi_{t,d}$.

547  2) Parameters of the negative binomial distributions that model the PEAK and the BACKGROUND

548   components at each time point: α_t, β_t, γ_t and δ_t.

549  3) Parameters of the negative binomial distributions that model the boundary movements in

550   EXPAND and CONTRACT dynamics at each time point: $\mu_{\text{EXPAND},t}, \delta_{\text{EXPAND},t}$ and

551   $\mu_{\text{CONTRACT},t}, \delta_{\text{CONTRACT},t}$, respectively.

553 The optimal parameter values are attempted to be estimated by Expectation Maximization (EM). In

554 particular, ChromTime attempts to optimize the conditional log-likelihood of the observed counts and

555 $Z_i = 1$ given the covariates (**Supplementary Methods**):

$$\sum_{i=1}^{M} \log P(\boldsymbol{O}_i = \boldsymbol{o}_i, \boldsymbol{Z}_i = \mathbf{1} | \boldsymbol{X}_i = \boldsymbol{x}_i)$$

559 **Computing the most likely spatial dynamic and peak boundaries for each block across the whole**

560 **time course**

562 After the optimal values for all model parameters are estimated from the data, for each block the most

563 likely positions of the peak boundaries at each time point are calculated. This procedure consists of two

steps. First, ChromTime determines for each block all time points with significantly low probability of

containing a false positive non-zero length peak. Second, conditioned on those time points, ChromTime

computes the most likely assignment of the peak boundary variables at each side and each time point

(**Supplementary Methods**).

ChromTime options used in this study

In this work, we applied ChromTime in narrow mode on all data for H3K4me2, H3K4me3, H3K27ac and

H3K(9,14)ac marks. We applied ChromTime in punctate mode on all ATAC-seq and DNase-seq data. No

control reads were used for ATAC-seq and DNase-seq. In addition, foreground reads for ATAC-seq were

shifted by 5bp in the direction of alignment (SHIFT=5), and for DNase-seq no shifting was applied

(SHIFT=0). We applied ChromTime in broad mode on all data for H3K79me2, Pol2, H3K4me1,

H3K27me3, H3K9me3 and H3K36me3 marks. All other options were set to their default values.

Timing Evaluation

The timing evaluation was conducted on a MacBook Pro laptop with 2.7GHz Intel Core i7 and 16 GB

RAM using 4 CPU cores.

Analyses with External Data

The procedures for analyses with external are described in **Supplementary Methods.**

ABBREVIATIONS

ATAC-seq: Assay for transposase accessible chromatin coupled with high-throughput DNA sequencing

ChIP-seq: Chromatin immunoprecipitation coupled with high-throughput DNA sequencing

DHS: DNase I Hypersensitive Site

DNase-seq: DNase I hypersensitivity assay followed by high-throughput DNA sequencing

EM: Expectation Maximization

FDR: False Discovery Rate

HM: Histone mark

TF: Transcription factor

TSS: Transcription start site

DECLARATIONS

Authors' contributions

J.E. conceived and supervised the project, designed the method, proposed analyses and wrote the

manuscript. P.F. designed the method, implemented the software, proposed analyses, performed all

analyses, processed all data and wrote the manuscript. Both authors read and approved the final

manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

ChromTime software is freely available at: https://github.com/ernstlab/ChromTime

No new experimental datasets were generated within this study. All datasets used as input for

ChromTime are publicly available from the references listed in **Table 1**. Links to all other datasets (gene

expression, TF binding and DHSs) used in this study are provided in the **Supplementary Methods**.

Funding

This work was supported by the CIRM Training Grant TG2-01169, the Eli and Edythe Broad Center of

Regenerative Medicine and Stem Cell Research at UCLA Training Program (P.F.); NIH grants

R01ES024995, U01HG007912, DP1DA044371, an NSF CAREER Award #1254200 and an Alfred P.

Sloan Fellowship (J.E.).

Acknowledgements

We are grateful to Constantinos Chronis, Kathrin Plath and members of the Ernst lab for useful

discussions.

Competing interests

None

REFERENCES

[1]   A. P. Boyle *et al.*, "High-resolution mapping and characterization of open chromatin

across the genome.," *Cell*, vol. 132, no. 2, pp. 311–22, Jan. 2008.

[2]   J. D. Buenrostro, P. G. Giresi, L. C. Zaba, H. Y. Chang, and W. J. Greenleaf,

"Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position.," *Nat. Methods*, vol. 10, no.

12, pp. 1213–8, Dec. 2013.

[3]   J. Ernst *et al.*, "Mapping and analysis of chromatin state dynamics in nine human cell

types.," *Nature*, vol. 473, no. 7345, pp. 43–9, May 2011.

[4]   A. Barski *et al.*, "High-resolution profiling of histone methylations in the human genome.,"

Cell, vol. 129, no. 4, pp. 823–37, May 2007.

[5]   T. S. Mikkelsen *et al.*, "Genome-wide maps of chromatin state in pluripotent and lineage-

committed cells.," *Nature*, vol. 448, no. 7153, pp. 553–60, Aug. 2007.

[6]   ENCODE_Project_Consortium, "An integrated encyclopedia of DNA elements in the

human genome.," *Nature*, vol. 489, no. 7414, pp. 57–74, Sep. 2012.

[7]   Roadmap_Epigenomics_Consortium *et al.*, "Integrative analysis of 111 reference human epigenomes," *Nature*, vol. 518, no. 7539, pp. 317–330, Feb. 2015.

[8]   J. H. A. Martens and H. G. Stunnenberg, "BLUEPRINT: mapping human blood cell epigenomes.," *Haematologica*, vol. 98, no. 10, pp. 1487–9, Oct. 2013.

[9]   F. D. Lay *et al.*, "Reprogramming of the human intestinal epigenome by surgical tissue transposition.," *Genome Res.*, vol. 24, no. 4, pp. 545–53, Apr. 2014.

[10] P. Fiziev *et al.*, "Systematic Epigenomic Analysis Reveals Chromatin States Associated with Melanoma Progression.," *Cell Rep.*, vol. 19, no. 4, pp. 875–889, Apr. 2017.

[11] S. Mei *et al.*, "Cistrome Data Browser: a data portal for ChIP-Seq and chromatin accessibility data in human and mouse," *Nucleic Acids Res.*, vol. 45, no. D1, pp. D658–D662, Jan. 2017.

[12] M. J. Ziller *et al.*, "Dissecting neural differentiation regulatory networks through epigenetic footprinting.," *Nature*, vol. 518, no. 7539, pp. 355–359, Dec. 2014.

[13] A. M. Tsankov *et al.*, "Transcription factor binding dynamics during human ES cell differentiation," *Nature*, vol. 518, no. 7539, pp. 344–349, Feb. 2015.

[14] K. K.-H. Farh *et al.*, "Genetic and epigenetic fine mapping of causal autoimmune disease variants," *Nature*, vol. 518, no. 7539, pp. 337–343, Oct. 2014.

[15] E. Gjoneska *et al.*, "Conserved epigenomic signals in mice and humans reveal immune basis of Alzheimer's disease," *Nature*, vol. 518, no. 7539, pp. 365–369, Feb. 2015.

[16] A. Gusev *et al.*, "Partitioning heritability of regulatory and cell-type-specific variants across 11 common diseases.," *Am. J. Hum. Genet.*, vol. 95, no. 5, pp. 535–52, Nov. 2014.

[17] J. A. Zhang, A. Mortazavi, B. A. Williams, B. J. Wold, and E. V Rothenberg, "Dynamic transformations of genome-wide epigenetic marking and transcriptional control establish T cell identity.," *Cell*, vol. 149, no. 2, pp. 467–82, Apr. 2012.

[18] T. S. Mikkelsen *et al.*, "Comparative epigenomic analysis of murine and human

adipogenesis.," *Cell*, vol. 143, no. 1, pp. 156–69, Oct. 2010.

[19] D. Lara-Astiaso *et al.*, "Chromatin state dynamics during blood formation," *Science.*, vol.

345, no. 6199, pp. 943–949, Aug. 2014.

[20] M. R. Corces *et al.*, "Lineage-specific and single-cell chromatin accessibility charts

human hematopoiesis and leukemia evolution.," *Nat. Genet.*, vol. 48, no. 10, pp. 1193–

203, 2016.

[21] D. K. Goode *et al.*, "Dynamic Gene Regulatory Networks Drive Hematopoietic

Specification and Differentiation," *Dev. Cell*, Feb. 2016.

[22] S. L. Paige *et al.*, "A temporal chromatin signature in human embryonic stem cells

identifies regulators of cardiac development.," *Cell*, vol. 151, no. 1, pp. 221–32, Sep.

2012.

[23] J. A. Wamstad *et al.*, "Dynamic and coordinated epigenetic regulation of developmental

transitions in the cardiac lineage.," *Cell*, vol. 151, no. 1, pp. 206–20, Sep. 2012.

[24] D. Cacchiarelli *et al.*, "Integrative Analyses of Human Reprogramming Reveal Dynamic

Nature of Induced Pluripotency," *Cell*, vol. 162, no. 2, pp. 412–424, Jul. 2015.

[25] J. Chen *et al.*, "Hierarchical Oct4 Binding in Concert with Primed Epigenetic

Rearrangements during Somatic Cell Reprogramming," *Cell Rep.*, Jan. 2016.

[26] R. P. Koche *et al.*, "Reprogramming factor expression initiates widespread targeted

chromatin remodeling.," *Cell Stem Cell*, vol. 8, no. 1, pp. 96–105, Jan. 2011.

[27] C. Chronis *et al.*, "Cooperative Binding of Transcription Factors Orchestrates

Reprogramming.," *Cell*, vol. 168, no. 3, p. 442–459.e20, Jan. 2017.

[28] O. Bogdanovic *et al.*, "Dynamics of enhancer chromatin signatures mark the transition

from pluripotency to cell specification during embryogenesis.," *Genome Res.*, vol. 22, no.

10, pp. 2043–53, Oct. 2012.

[29] N. Nègre *et al.*, "A cis-regulatory map of the Drosophila genome.," *Nature*, vol. 471, no.

7339, pp. 527–31, Mar. 2011.

[30]   P. Yu *et al.*, "Spatiotemporal clustering of epigenome reveals rules of dynamic gene

regulation.," *Genome Res.*, p. gr.144949.112-, Oct. 2012.

[31]   P. Arnold *et al.*, "Modeling of epigenome dynamics identifies transcription factors that

mediate Polycomb targeting.," *Genome Res.*, p. gr.142661.112-, Sep. 2012.

[32]   N. Koike *et al.*, "Transcriptional architecture and chromatin landscape of the core

circadian clock in mammals.," *Science*, vol. 338, no. 6105, pp. 349–54, Oct. 2012.

[33]   R. Ostuni *et al.*, "Latent Enhancers Activated by Stimulation in Differentiated Cells," *Cell*,

vol. 152, no. 1–2, pp. 157–71, Jan. 2013.

[34]   S. K. Reilly *et al.*, "Evolutionary changes in promoter and enhancer activity during human

corticogenesis," *Science (80-.).*, vol. 347, no. 6226, pp. 1155–1159, Mar. 2015.

[35]   A. Weiner *et al.*, "High-Resolution Chromatin Dynamics during a Yeast Stress

Response.," *Mol. Cell*, vol. 58, no. 2, pp. 371–386, Mar. 2015.

[36]   J. Cotney *et al.*, "The evolution of lineage-specific regulatory activities in the human

embryonic limb.," *Cell*, vol. 154, no. 1, pp. 185–96, Jul. 2013.

[37]   J. Zhu *et al.*, "A time-series analysis of altered histone H3 acetylation and gene

expression during the course of MMAIII-induced malignant transformation of urinary

bladder cells," *Carcinogenesis*, vol. 38, no. 4, pp. 378–390, Apr. 2017.

[38]   Y. Zhang *et al.*, "Model-based analysis of ChIP-Seq (MACS).," *Genome Biol.*, vol. 9, no.

9, p. R137, Jan. 2008.

[39]   S. Heinz *et al.*, "Simple combinations of lineage-determining transcription factors prime

cis-regulatory elements required for macrophage and B cell identities.," *Mol. Cell*, vol. 38,

no. 4, pp. 576–89, May 2010.

[40]   S. Xu, S. Grullon, K. Ge, and W. Peng, "Spatial clustering for identification of ChIP-

enriched regions (SICER) to map regions of histone methylation patterns in embryonic

stem cells.," *Methods Mol. Biol.*, vol. 1150, pp. 97–111, 2014.

[41] H. Xing *et al.*, "Genome-Wide Localization of Protein-DNA Binding and Histone

Modification by a Bayesian Change-Point Method with ChIP-seq Data," *PLoS Comput.*

Biol., vol. 8, no. 7, p. e1002613, Jul. 2012.

[42] A. Harmanci, J. Rozowsky, and M. Gerstein, "MUSIC: identification of enriched regions in

ChIP-Seq experiments using a mappability-corrected multiscale signal processing

framework.," *Genome Biol.*, vol. 15, no. 10, p. 474, 2014.

[43] J. Ernst and M. Kellis, "ChromHMM: automating chromatin-state discovery and

characterization.," *Nat. Methods*, vol. 9, no. 3, pp. 215–6, Mar. 2012.

[44] M. M. Hoffman, O. J. Buske, J. Wang, Z. Weng, J. A. Bilmes, and W. S. Noble,

"Unsupervised pattern discovery in human chromatin structure through genomic

segmentation.," *Nat. Methods*, vol. 9, no. 5, pp. 473–6, May 2012.

[45] M. Allhoff, K. Seré, H. Chauvistré, Q. Lin, M. Zenke, and I. G. Costa, "Detecting

differential peaks in ChIP-seq signals with ODIN.," *Bioinformatics*, vol. 30, no. 24, pp.

3467–75, Dec. 2014.

[46] H. Xu, C.-L. Wei, F. Lin, and W.-K. Sung, "An HMM approach to genome-wide

identification of differential histone modification sites from ChIP-seq data.,"

Bioinformatics, vol. 24, no. 20, pp. 2344–9, Oct. 2008.

[47] J. Biesinger, Y. Wang, and X. Xie, "Discovering and mapping chromatin states using a

tree hidden Markov model," *BMC Bioinformatics*, vol. 14, no. Suppl 5, p. S4, 2013.

[48] D. Hnisz *et al.*, "Super-enhancers in the control of cell identity and disease.," *Cell*, vol.

155, no. 4, pp. 934–47, Nov. 2013.

[49] S. C. J. Parker *et al.*, "Chromatin stretch enhancer states drive cell-specific gene

regulation and harbor human disease risk variants," *PNAS*, p. 1317023110-, Oct. 2013.

[50] B. A. Benayoun *et al.*, "H3K4me3 breadth is linked to cell identity and transcriptional

consistency.," *Cell*, vol. 158, no. 3, pp. 673–88, Jul. 2014.

[51] K. Chen *et al.*, "Broad H3K4me3 is associated with increased transcription elongation and

enhancer activity at tumor-suppressor genes.," *Nat. Genet.*, vol. 47, no. 10, pp. 1149–57,

Oct. 2015.

[52]   A. Dincer *et al.*, "Deciphering H3K4me3 broad domains associated with gene-regulatory

networks and conserved epigenomic landscapes in the human brain," *Transl. Psychiatry*,

vol. 5, no. 11, p. e679, Nov. 2015.

[53]   R. D. Hawkins *et al.*, "Distinct epigenomic landscapes of pluripotent and lineage-

committed human cells.," *Cell Stem Cell*, vol. 6, no. 5, pp. 479–91, May 2010.

[54]   Z. Wang *et al.*, "Combinatorial patterns of histone acetylations and methylations in the

human genome.," *Nat. Genet.*, vol. 40, no. 7, pp. 897–903, Jul. 2008.

[55]   H. H. Ng, F. Robert, R. A. Young, and K. Struhl, "Targeted recruitment of Set1 histone

methylase by elongating Pol II provides a localized mark and memory of recent

transcriptional activity.," *Mol. Cell*, vol. 11, no. 3, pp. 709–19, Mar. 2003.

[56]   E. Smith, C. Lin, and A. Shilatifard, "The super elongation complex (SEC) and MLL in

development and disease.," *Genes Dev.*, vol. 25, no. 7, pp. 661–72, Apr. 2011.

[57]   S. A. Lacadie, M. M. Ibrahim, S. A. Gokhale, and U. Ohler, "Divergent Transcription and

Epigenetic Directionality of Human Promoters," *FEBS J.*, Apr. 2016.

[58]   A. C. Cameron and P. K. Trivedi, *Regression analysis of count data*, Second edition.

Cambridge University Press, 2013.

[59]   N. U. Rashid, P. G. Giresi, J. G. Ibrahim, W. Sun, and J. D. Lieb, "ZINBA integrates local

covariates with DNA-seq data to identify broad and narrow regions of enrichment, even

within amplified genomic regions," *Genome Biol.*, vol. 12, no. 7, p. R67, 2011.

[60]   Z. Bar-Joseph, D. K. Gifford, and T. S. Jaakkola, "Fast optimal leaf ordering for

hierarchical clustering.," *Bioinformatics*, vol. 17 Suppl 1, pp. S22-9, 2001.

FIGURE LEGENDS

Figure 1: **Overview of the ChromTime method. (A)** Examples of H3K4me2 peaks with steady,
expanding and contracting boundary dynamics, shown from left to right respectively, across five time
points during mouse T cell development[17]. Time points 1, 2 and 3 correspond to in vitro differentiated T
cell precursors (FLDN1, FLDN2a and FLDN2b), whereas time points 4 and 5 correspond to in vivo
purified thymocytes (ThyDN3 and ThyDP). Normalized ChIP-seq signal, MACS2[38] peaks (black
rectangles) and ChromTime output are shown for each time point. Peaks upstream of Zfp148 gene are
called steady by ChromTime despite fluctuations of MACS2 peak boundaries. In contrast, ChromTime
calls a peak at the Skap1/GM11529 promoter to expand after time points 2 and 3. Conversely,
ChromTime calls a peak upstream of GPR141 gene to contract after time points 2, 3, and 4. **(B)** Overview
of the ChromTime method. During the block finding stage, input foreground and, optionally, control reads
are used to determine blocks of signal enrichment. In the dynamics prediction stage, for each block, peak
boundary positions are predicted at each time point and peak boundary dynamics are predicted at each
pair of consecutive time points. **(C)** Schematic of predicting dynamics for one block. Boxes represent
genomic bins at each time point. Foreground signal is depicted as blue bars for each bin whose height
represents the number of reads mapped to the bin. ChromTime learns a probabilistic mixture model from
the input data to partition each block at each time point into peak and background components. Bins in
the peak component (orange) mark peaks of signal enrichment whereas those in the background
component (white) mark flanking background signal. The movement of the boundaries on the left and the
right side of peaks between consecutive time points are estimated by reasoning jointly about the input
data from all time points.

Figure 2: **Sample output from ChromTime with contracting peaks.** Genome browser screenshot with
sample output of ChromTime for H3K4me2 from the T cell development time course in mouse[17] with 5
time points at the Esam/Vsig2/Nrgn locus. Time points 1, 2 and 3 correspond to in vitro differentiated T
cell precursors (FLDN1, FLDN2a, and FLDN2b), whereas time points 4 and 5 correspond to in vivo
purified thymocytes (ThyDN3 and ThyDP). The input ChIP-seq signal and MACS2[38] peaks (black boxes
under each signal track) are shown in the upper panel of the screenshot. The ChromTime predicted
peaks colored by their boundary dynamics for each block at each time point are shown in the bottom

panel. The first peak in each block is colored in dark grey. Each subsequent peak is colored with respect

to the predicted dynamic relative to its previous time point. Peaks with steady boundaries on both sides

are shown in light grey, and those with at least one contracting boundary are shown in blue. Nearby

peaks that touch boundaries are visualized as one peak by the genome browser. Not shown in the figure

are expanding peaks, peaks at single time points and peaks with opposite dynamics (EXPAND on the left

and CONTRACT on the right, or vice versa), which would be colored in red, orange and black,

respectively. See **Fig S2** for examples of predicted expanding peaks.

Figure 3: **Changes in GATA3 binding and gene expression at predicted H3K4me2 dynamics in T**

cell development. (**A**) Fold enrichments of cell type specific and shared peaks of GATA3, which is a

master regulator in T cell development[17], are shown for three sets of blocks with predicted H3K4me2

peaks: 1) blocks with peaks present at all time points whose boundaries hold steady on both sides

throughout the whole time course (T1-Tn Steady); 2) blocks with non-contracting peaks whose

boundaries expand between at least one pair of consecutive time points and have a peak at the last time

point (Tx-Tn Expand); and 3) blocks with non-expanding peaks whose boundaries contract between at

least one pair of consecutive time points and have a peak at the first time point (T1-Tx Contract). First

column shows the percentage of bases out of all bases covered by peaks of the set. Last row shows the

baseline percentage for each feature out of all bases covered by ChromTime peaks at any time point.

Percentages are colored from 0 (white) to 100 (green). Fold enrichments in each column are colored from

1 (white) to the maximum value in the column (red). FLDN1 and ThyDP denote differentiated T cell

precursors and purified thymocytes, which are the first and the last time point, respectively. **(B)**

Boundaries of predicted H3K4me2 peaks in blocks with at least one predicted non-zero length peak

overlapping annotated TSSs were sorted in decreasing order by their posterior probability for EXPAND

dynamic (left plots) and CONTRACT dynamic (right plots) at each pair of consecutive time points

(**Supplementary Methods**). Gene expression differences between consecutive time points were

calculated as the average difference across all genes with overlapping TSSs for each block. For each

posterior rank (X-axis) the plot shows the cumulative average gene expression difference (Y-axis).

Expanding boundaries associated with increase of gene expression and contracting boundaries

associated with decrease of gene expression. Shaded regions correspond to 95% confidence intervals.

Figure 4: ChromTime predictions associate better with expression changes than boundary

movements of peaks called in isolation. (A) For H3K4me2 in mouse T cell development[17]

ChromTime was applied once with data from all time points (ChromTime ALL), and once with single time

points in isolation (ChromTime SINGLE; **Supplementary Methods**). Time points 1, 2 and 3 correspond to

T cell precursors, whereas 4 and 5 to purified thymocytes. Peaks called by both procedures overlapping

annotated TSSs were analyzed for their relationship with gene expression changes. (**i**) (Left) Comparison

of agreement with expression for expansions when applying ChromTime ALL and ChromTime SINGLE

for the change between time points 3 and 4. Peak boundaries were sorted in decreasing order of their

EXPAND posterior probabilities from ChromTime ALL and compared to sorting them in decreasing order

of the difference of peak boundary positions in ChromTime SINGLE peaks with positive differences in

boundary positions indicating peaks expanding with time. Each boundary was also ranked by the average

gene expression difference of TSSs overlapping its block in decreasing order with positive expression

differences indicating gain with time. The cumulative average boundary rank of expression change (Y-

axis) is shown for the boundary change ranking for ChromTime ALL and ChromTime SINGLE (X-axis).

Low Y-values indicate stronger association with expression changes. Black line shows expected average

expression change rank. Shaded regions indicate 95% confidence intervals. Plots for other time points

can be found in **Fig S5**. (Right) Analogous to left plots for Contract posterior probabilities for ChromTime

ALL, increasing order of the difference of boundary change positions for ChromTime SINGLE, and

increasing order of expression changes. (**ii**) Differences between ChromTime ALL and ChromTime

SINGLE values shown in (**i**) between time points 3 and 4 as well as for all other pairs of time points.

Positive values correspond to boundary ranks for which ChromTime ALL posteriors better associate with

gene expression changes than boundary movements of ChromTime SINGLE peaks. Black lines show

expected difference of zero between random rankings. **(B)** As in (**A**) for H3K4me3 in human stem cell

reprogramming[24]. Time points correspond to human inducible and immortalized fibroblasts-like (hiF-T)

cells, hiF-T at 5, 10 and 20 days after induction, and human induced pluripotent stem cells (hIPSC).

Figure 5: **Spatial dynamics can contain additional information about gene expression changes beyond signal density changes.** Gene expression change is plotted as function of ChIP-seq signal density change after loess smoothing for each predicted ChromTime dynamic for **(A)** H3K4me2 dynamics in T cell development in mouse[17]; and **(B)** in H3K4me3 dynamics in stem cell reprogramming in human[24] (**Supplementary Methods**). Peaks of each type of dynamics were pooled from all time points for this analysis. Peaks with asymmetric dynamics E/S and S/E were pooled together in the "E-S" group. Similarly C/S and S/C peaks were pooled in the "C-S" group. The total number of peaks in each group is shown in parenthesis. In both systems, for a range of signal density changes, peaks with the same signal density change associated with different gene expression changes depending on the predicted spatial dynamic. Shaded regions represent 95% confidence intervals.

Figure 6: Spatial dynamics of multiple different chromatin marks co-localize within a time course. Hierarchical clustering with optimal leaf ordering[60] of the geometric average fold enrichments taken across all time points of the overlap of every pair of predicted spatial dynamics for mapped HMs in **(A)** mouse adipogenesis[18] and **(B)** human stem cell reprogramming[24]. At each pair of time points, "Expand" and "Contract" dynamics are defined as all peaks that are predicted as either unidirectional or bidirectional expansions and contractions, respectively, whereas "Steady" dynamics are defined as all peaks that have predicted steady boundaries at both sides. Peaks with "Expand" dynamic on one side and "Contract" dynamic on the other were excluded from this analysis. In both datasets, expansions, contractions and steady peaks of H3K4me2, H3K4me3 and H3K27ac and to a lesser extent of H3K4me1 tend to cluster together within each of the three classes, whereas spatial dynamics of H3K27me3 and H3K36me3 peaks tend to occupy different locations. All enrichments were capped at 50 before clustering.

Figure 7: **Direction of asymmetric dynamics correlates with direction of transcription. (A) (i)** Left panel shows a schematic representation of unidirectional expansions that expand in the same direction as transcription and in the opposite direction of transcription. The adjacent plots show, for each mark, the average $\log_2$ ratio across all time points in each time course between the fraction of unidirectional

expansions that expand in the same directions as transcription of the nearest gene and the fraction of

unidirectional expansions that expand in the opposite direction of transcription of the nearest gene for

blocks that are within 1 kb of annotated TSSs and for more distal blocks. Positive values correspond to

enrichment of unidirectional expansions in the same direction as transcription. For marks mapped in at

least six time courses, a black line is plotted representing the average across all data sets and significant

differences are denoted with asterisks based on a two-tailed Mann-Whitney test at a p-value threshold of

0.05. **(ii)** Left panel shows analogous schematic for unidirectional contractions. Likewise, adjacent plots

show, for each mark, the average $\log_2$ ratio between the fraction of unidirectional contractions that

contract in the opposite direction of transcription of the nearest TSS and unidirectional contractions that

contract in the same direction as transcription of the nearest TSS. **(B)** Left panel shows an example of

unidirectional expansions between pairs of time points that expand in the same direction as transcription

at the Hs6st1 gene of the H3K4me2 mark in the T cell development dataset[17]. Right panel shows an

example of unidirectional contractions in the opposite direction of transcription at the DNMT3B gene.

Time points 1, 2 and 3 correspond to in vitro differentiated T cell precursors, whereas time points 4 and 5

correspond to in vivo purified thymocytes. The predicted ChromTime peaks colored by their boundary

dynamics are shown under the signal track for each time point.

Table 1: Datasets used for analysis with ChromTime

System	Chromatin Marks	Species	# Time Points	Reference
Adipogenesis	H3K4me2 H3K4me3 H3K27ac H3K4me1 H3K36me3 H3K27me3	Mouse Human	4	[18]
Blood Formation	H3K4me2 H3K4me3	Mouse	5-7	[19]

	H3K27ac			
	H3K4me1			
	ATAC-seq			
Blood Formation	ATAC-seq	Human	5	[20]
Human Fetal Brain Development	DNase-seq	Human	3	[7]
Cardiac Development	H3K4me3	Mouse	4	[23]
	H3K27ac			
	H3K4me1			
	H3K27me3			
	H3K36me3			
	Pol2			
Cardiac Development	H3K4me3	Human	5	[22]
	H3K27me3			
	H3K36me3			
Embryogenesis	H3K4me3	Zebrafish	4	[28]
	H3K27ac			
	H3K4me1			
Macrophage Differentiation	H3K4me3	Mouse	5	[21]
	H3K9ac			
	H3K27ac			
	H3K27me3			
Neural Differentiation	H3K4me3	Human	5	[12]
	H3K27ac			
	H3K27me3			
	H3K4me1			
Stem Cell Reprogramming	H3K4me2	Human	4-6	[24]
	H3K4me3			

	H3K27ac			
	H3K4me1			
	H3K27me3			
	H3K36me3			
Stem Cell Reprogramming	H3K4me2	Mouse	4	[27]
	H3K4me3			
	H3K9ac			
	H3K27ac			
	H3K27me3			
	H3K36me3			
	H3K4me1			
	H3K79me2			
	H3K9me3			
	ATAC-seq			
Stem Cell Reprogramming	H3K4me3	Mouse	9	[25]
	H3K27ac			
	H3K4me1			
	H3K27me3			
	Pol2			
T Cell Development	H3K4me2	Mouse	5	[17]
	H3K(9,14)ac			
	H3K27me3			

Figure 1

A.

upstream Zfp148   SKAP1/Gm11529   upstream GPR141

T1
T2
T3
T4
T5

ChromTime output

B.

Input: Aligned sequencing reads for a single chromatin feature at multiple time points and control [optionally]

T1
T2
T3
T4
T5

1) Determine genomic coordinates of blocks of signal enrichment

Blocks

T1
T2
T3
T4
T5

2) Call peaks and classify peak boundaries as:
- Steady
- Expanding
- Contracting

Peaks

T1
T2
T3
T4
T5

■ Contract   ■ Expand   ■ Steady

C.

Background   Peak   Background

T1

Steady   Steady

T2

Expand   Expand

T3

Expand   Expand

T4

Steady   Steady

T5

Figure 2

H3K4me2 in mouse T cell development

A.

Dynamic	% all block bases	FLDN1-ThyDP shared GATA3	ThyDP-specific GATA3	FLDN1-specific GATA3
T1-Tn Steady	28	1.2	1.1	0.5
Tx-Tn Expand	13	0.6	3.4	0.2
T1-Tx Contract	55	1	0.3	1.5
Base %		0.1	0.2	0.1

B.

EXPAND Dynamic

CONTRACT Dynamic

Cumulative Average Expression Change

Rank by EXPAND Posterior in Descending Order

Rank by CONTRACT Posterior in Descending Order

T 1/2
T 2/3
T 3/4
T 4/5

Figure 4

A.

H3K4me2 in mouse T cell development

i)

Expansions T3/4 — ChromTime ALL, ChromTime SINGLE T3 & T4

Contractions T3/4 — ChromTime ALL, ChromTime SINGLE T3 & T4

ii)

Expansions at All Time Points — T 1/2, T 2/3, T 3/4, T 4/5

Contractions at All Time Points — T 1/2, T 2/3, T 3/4, T 4/5

B.

H3K4me3 in human stem cell reprogramming

i)

Expansions T3/4 — ChromTime ALL, ChromTime SINGLE T3 & T4

Contractions T3/4 — ChromTime ALL, ChromTime SINGLE T3 & T4

ii)

Expansions at All Time Points — T 1/2, T 2/3, T 3/4, T 4/5

Contractions at All Time Points — T 1/2, T 2/3, T 3/4, T 4/5

Figure 5

A.

H3K4me2 in mouse T cell development

B.

H3K4me3 in human stem cell reprogramming

Figure 6

Figure 7

A.

i) Unidirectional Expansions

ii) Unidirectional Contractions

Dataset

- Mouse Reprogramming (Chronis et al., 2017)
- Human Adipogenesis
- Mouse Macrophage Differentiation
- Mouse T-cell Development
- Human Neural Differentiation
- Mouse Cardiac Development
- Human Brain Development
- Human Hematopoiesis
- Human Cardiac Development
- Mouse Reprogramming (Chen et al., 2016)
- Mouse Hematopoiesis
- Zebrafish Embryogenesis
- Human Reprogramming
- Mouse Adipogenesis

B.

Hs6st1

DNMT3B