1   **Methylation of *E. coli* Class I Release Factors Increases the Accuracy of**
2   **Translation Termination *in vitro***
3

6   Gürkan Korkmaz

7   Department of Cell and Molecular Biology, Uppsala University, Sweden

10   Correspondents to Gürkan Korkmaz via gurkankorkmaz@mail.com

Keywords:

Ribosome, stop codon recognition, kinetics, post-translational modification, accuracy

Abbreviations: RF - release factor, RC - release complex

ABSTRACT

Ribosomal protein synthesis (translation) is a highly accurate process. Translation

termination, in particular, must be accurate to prevent truncated proteins. How this

accuracy is achieved is not fully understood in all its details. Using an *E. coli in vitro*

system, I explore novel mechanisms that contribute to the high accuracy of translation

termination. By comparing the Michaelis-Menten parameters of methylated and non-

methylated release factors on cognate and non-cognate codons. Post-translational

methylation of a strictly conserved GGQ motif in class I release factors increases the

accuracy of termination by up to 5-fold. This happens by increasing both the maximum

rate of peptide release (k_{cat}) and Michaelis-Menten constant (K_M). Further, I

demonstrate here that a non-methylated release factor acts like an uncompetitive

inhibitor of enzyme reactions. Overall, this study shows that the methylation of class I

release factors is a novel mechanism contributing to highly accurate translation

termination.

INTRODUCTION

Translation termination is an essential component of protein production. mRNA translation is terminated when a stop codon is translocated into the ribosomal A site. This new configuration is now called a release complex (RC). The subsequent binding of the class I release factor induces peptide release. In nearly all life forms, the codons UAA, UAG, and UGA are used as termination signals, with some exceptions (Ivanova et al., 2014). In bacteria, the class I release factors are RF1 and RF2. Both class I release factors possess a strictly conserved glycine-glycine-glutamine (GGQ) motif, which is essential for the ester bond hydrolysis by which the peptide is released from the tRNA.

Stop codon recognition is highly accurate *in vitro* (Freistroffer et al., 2000) and *in vivo* (Jørgensen et al., 1993), but our understanding of how this accuracy is achieved is still rudimentary. The accuracy of a reaction is defined as the ratio of the efficiency of a cognate to a non-cognate reaction (Freistroffer et al., 2000). The efficiency is determined by the ratio of the maximum rate of the reaction (k_{cat}) to the concentration at half k_{cat} (K_M). When the ratio of cognate and non-cognate reaction is one, accuracy is non-existent, meaning no discrimination between the cognate and non-cognate substrate are observable.

Methylation of the glutamine in the GGQ motif has been shown to increase k_{cat} (Indrisiunaite et al., 2015). Methylation happens via the hemK protein, which modifies glutamine 253 (*E. coli* numbering) to N^5-methyl-glutamine (Heurgue-Hamard et al., 2002). However, its effect on K_M is unknown.

Highly conserved recognition motifs of RF1 and RF2 are located in the vicinity of the stop codons, but about 75Å distant from the catalytic center (Korostelev, 2011). These motifs (PxT in RF1 and SPF in RF2) were thought to be responsible for the specificity

of stop codon recognition by the release factors and therefore named 'tripeptide

anticodons', analogous to the anticodons of tRNAs (Ito et al., 1996, 2000).

Class II release factor RF3 was shown to affect accuracy partly (Freistroffer et al.,

2000). Further, it was shown that depletion of the ribosomal protein L11 from an RC

increases the accuracy of translation termination with RF1 but not RF2 (Bouakaz et

al., 2006). The accuracy of translation termination is mostly thought to be due to the

extensive interaction network between class I release factor, stop codon, and rRNA

(Sund et al., 2010).

The strictly conserved GGQ motif is essential for ester bond hydrolysis, as shown by

studies where mutations within the GGQ motif disrupted translation termination *in vitro*

(Zavialov et al., 2002) and *in vivo* (Frolova et al., 1999; Seit-Nebi et al., 2001).

However, in *E. coli* different mechanisms seem to be at play for several reasons. The

conserved methylation of the GGQ motif glutamine to N^5-methyl-glutamine is not

essential in *E. coli*, as the responsible enzyme for methylation RF1/RF2, hemK, is not

essential for cell viability (Heurgue-Hamard et al., 2002; Pannekoek et al., 2005). Also,

release factors where the GGQ motif is altered to GGA are still able to induce

translation termination *in vitro* (Shaw and Green, 2007; Zavialov et al., 2002). On the

other hand, methylation of the class I release factors is essential for cell viability in

minimal media (Mora et al., 2007). As mentioned earlier, methylation increases the

maximum rate of peptide release (k_{cat}) (Indrisiunaite et al., 2015), but the magnitude

of the k_{cat} increase was shown to depend on the nature of the amino acid on the tRNA

occupying the P site (Pierson et al., 2016).

The actual ester bond hydrolysis occurs via a coordination of a nucleophilic hydroxide

molecule by the GGQ motif into the PTC. Crystal structures of class I release factors

indicate a difference between the positioning of the methylated and non-methylated

glutamine residue within the PTC (Pierson et al., 2016).

Although RF1 and RF2 share high sequence and structural similarity, they have

different codon specificities. RF1 recognizes the stop codons UAA and UAG, whereas

RF2 recognizes the stop codons UAA and UGA (Scolnick et al., 1968).

RESULTS & DISCUSSION

In this study, I investigated the methylation of the GGQ glutamine as an additional factor that might enhance the accuracy of translation termination *in vitro*. The accuracy of a reaction is defined as the ratio of the efficiency (k_{cat} / K_M) of a cognate to a non-cognate reaction (Freistroffer et al., 2000). When the ratio of cognate and non-cognate reaction is one, accuracy is non-existent, meaning no discrimination between the cognate and non-cognate substrate are observable. I systematically determined the k_{cat} and K_M parameters for release factors on various purified release complexes (Korkmaz and Sanyal, 2017) containing cognate and non-cognate codons in A site and compared the resulting accuracy with the reactions involving methylated (mRF1 or mRF2) and non-methylated (RF1 or RF2) release factors (Table 1 and Figure 1 and 2). UAA was used as the cognate codon in all experiments, while UAG and UGA were used as the non-cognate substrates for RF1 and RF2 respectively since these codons allow discrimination similar to non-cognate sense codons (Freistroffer et al., 2000).

The determined cognate k_{cat} parameters were in agreement with previous data for non-methylated release factors (Bouakaz et al., 2006; Freistroffer et al., 2000) and also with recent studies that used methylated release factors (Indrisiunaite et al., 2015; Pierson et al., 2016). I observed a 4.5-fold increase in the maximum rate of peptide release for mRF1 on its cognate UAA codon and a 5-fold increase in the maximum rate of tritium-labeled fMet (^{3H}fMet) release on the non-cognate codon UGA. However, it should be noted that for the cognate reactions, the K_M values for reactions with either mRF1 or RF1 were below the detection limit of 5 nM. Thus, the real difference in K_M might be much larger than what could be measured in these experiments. In Table 1 the estimated K_M parameters are listed. The K_M value form RF1 on UGA decreased by 3-fold, from 164 nM to 48 nM, when methylation was missing. These resulted in an

accuracy of 25 for RF1 on UGA. When RF1 was methylated, the accuracy increased

to 90 (an increase of over 3.5-fold) (Table 1).

The effect of methylation on the translation termination accuracy was even more

pronounced with RF2 on the non-cognate UAG codon. With methylation, the k_{cat}

increased 2.6-fold for the non-cognate substrate and nearly 16-fold for the cognate

substrate (UAA). The accuracy of translation termination increased 5-fold, from 13 to

68, when RF2 was methylated.

As mentioned earlier, it is not surprising that the maximum rate of ^{3H}fMet release

increased in all cases; however, unexpectedly, the K_M parameters were also affected

by release factor methylation. For RF1 and mRF1, the change in K_M (Table 1) could

not be unambiguously determined in the experimental setup as both values were

below the detection limit, but an approximately 3-fold difference in K_M was observed

for the non-cognate reading of UGA; methylation of RF1 resulted in an increase of the

K_M value. More pronounced changes in K_M were observed for RF2 and mRF2. For

cognate stop codon recognition, a nearly 5-fold increase in K_M was observed when

RF2 was methylated (26 nM for RF2 and 123 nM for mRF2); a 4-fold increase was

seen in the case of the non-cognate substrate UAG (251 nM for RF2 and 1045 nM for

mRF2).

When hemK is knocked out, cells have compromised fitness, yet they are viable (Mora

et al., 2007). The knockout of hemK consequently leads to the lack of methylation in

class I RFs and as shown here. The protein hemK becomes essential when cells are

grown in minimal media (Mora et al., 2007). In the light of the data presented here,

one could explain this phenotype by the fact of the lower accuracy of RFs than the

slowed down peptide release. In minimal media, ribosomes are likely to pause on

codons due to the amino acid limitation (Li et al., 2012). Inaccurate RFs could bind to

these complexes and induce premature termination.

Interestingly, the Lineweaver-Burk plots of the methylated and non-methylated release

factor kinetics data revealed that the non-methylated release factors behave similarly

to uncompetitive inhibitors in enzyme reactions (Figure 3). However, the mechanism

of the inhibition or details of any mechanistic similarity to uncompetitive enzyme

inhibition remains obscure.

Overall, I herein presented evidence that methylation of class I release factors

contribute to the accuracy of translation termination. This is surprising by the fact that

a post-translational modification, far from the initial binding site, effects the binding

constant K_M.

Further studies, unraveling the potential *in vivo* aspects of this finding are imminent,

especially in light of the ribosome rescue phenomenon. This study demonstrates that

release factor methylation plays an additional biological role beyond the known

increase of the maximum rate of peptide release. This previously unknown variable

can explain how the outstanding accuracy of translation termination is achieved in the

absence of any proofreading steps.

ACKNOWLEDGEMENTS

GK was funded partly by a scholarship from the Sven and Lilly Lawski Foundation for Scientific Research. GK would like to acknowledge Marina Fridman from the Champalimaud Centre for the Unknown, Lisbon, Portugal, for proofreading an early version of the manuscript. Further Suparna Sanyal from the Department of Cell and Molecular Biology, Uppsala University, Sweden, is acknowledged for grant acquisition that supported the experimental part of this work (Swedish Research Council 2013-8778, 2014-4423, 2008-6593 [Linnaeus grant to Uppsala RNA Research Center], and the Knut and Alice Wallenberg Foundation to RiboCORE [KAW2011.0081]).

CONFLICT OF INTEREST

No conflict of interest is stated.

REFERENCES

Bouakaz, L., Bouakaz, E., Murgola, E.J., Ehrenberg, M., and Sanyal, S. (2006). The
Role of Ribosomal Protein L11 in Class I Release Factor-mediated Translation
Termination and Translational Accuracy. J. Biol. Chem. *281*, 4548–4556.

Freistroffer, D.V., Kwiatkowski, M., Buckingham, R.H., and Ehrenberg, M. (2000).
The accuracy of codon recognition by polypeptide release factors. Proc Natl Acad
Sci USA *97*, 2046–2051.

Frolova, L.Y., Tsivkovskii, R.Y., Sivolobova, G.F., Oparina, N.Y., Serpinsky, O.I.,
Blinov, V.M., Tatkov, S.I., and Kisselev, L.L. (1999). Mutations in the highly conserved
GGQ motif of class 1 polypeptide release factors abolish ability of human eRF1 to
trigger peptidyl-tRNA hydrolysis. RNA *5*, 1014–1020.

Heurgue-Hamard, V., Champ, S., Engstrom, A., Ehrenberg, M., and Buckingham,
R.H. (2002). The hemK gene in Escherichia coli encodes the N5-glutamine
methyltransferase that modifies peptide release factors. EMBO J *21*, 769–778.

Indrisiunaite, G., Pavlov, M.Y., Heurgué-Hamard, V., and Ehrenberg, M. (2015). On
the pH Dependence of Class-1 RF-Dependent Termination of mRNA Translation. J.
Mol. Biol. *427*, 1848–1860.

Ito, K., Ebihara, K., Uno, M., and Nakamura, Y. (1996). Conserved motifs in
prokaryotic and eukaryotic polypeptide release factors: tRNA-protein mimicry
hypothesis. Proc. Natl. Acad. Sci. U. S. A. *93*, 5443–5448.

Ito, K., Uno, M., and Nakamura, Y. (2000). A tripeptide "anticodon" deciphers stop
codons in messenger RNA. Nature *403*, 680–684.

Ivanova, N.N., Schwientek, P., Tripp, H.J., Rinke, C., Pati, A., Huntemann, M., Visel,
A., Woyke, T., Kyrpides, N.C., and Rubin, E.M. (2014). Stop codon reassignments in
the wild. Science *344*, 909–913.

Jelenc, P.C., and Kurland, C.G. (1979). Nucleoside triphosphate regeneration
decreases the frequency of translation errors. Proc. Natl. Acad. Sci. *76*, 3174–3178.

Jørgensen, F., Adamski, F.M., Tate, W.P., and Kurland, C.G. (1993). Release Factor-
dependent False Stops are Infrequent in Escherichia coli. J. Mol. Biol. *230*, 41–50.

Korkmaz, G., and Sanyal, S. (2017). R213I mutation in release factor 2 (RF2) is one
step forward for engineering an omnipotent release factor in bacteria Escherichia
coli. J. Biol. Chem. *292*, 15134–15142.

Korostelev, A.A. (2011). Structural aspects of translation termination on the
ribosome. RNA N. Y. N *17*, 1409–1421.

Li, G.-W., Oh, E., and Weissman, J.S. (2012). The anti-Shine-Dalgarno sequence
drives translational pausing and codon choice in bacteria. Nature *484*, 538–541.

Mora, L., Heurgué-Hamard, V., de Zamaroczy, M., Kervestin, S., and Buckingham,
R.H. (2007). Methylation of Bacterial Release Factors RF1 and RF2 Is Required for
Normal Translation Termination in Vivo. J. Biol. Chem. *282*, 35638–35645.

Pannekoek, Y., Heurgue-Hamard, V., Langerak, A.A.J., Speijer, D., Buckingham,
R.H., and van der Ende, A. (2005). The N5-Glutamine S-Adenosyl-L-Methionine-
Dependent Methyltransferase PrmC/HemK in Chlamydia trachomatis Methylates
Class 1 Release Factors. J Bacteriol *187*, 507–511.

Pierson, W.E., Hoffer, E.D., Keedy, H.E., Simms, C.L., Dunham, C.M., and Zaher,
H.S. (2016). Uniformity of Peptide Release Is Maintained by Methylation of Release
Factors. Cell Rep. *17*, 11–18.

Scolnick, E., Tompkins, R., Caskey, T., and Nirenberg, M. (1968). Release factors
differing in specificity for terminator codons. Proc Natl Acad Sci USA *61*, 768–774.

Seit-Nebi, A., Frolova, L., Justesen, J., and Kisselev, L. (2001). Class-1 translation
termination factors: invariant GGQ minidomain is essential for release activity and
ribosome binding but not for stop codon recognition. Nucleic Acids Res *29*, 3982–
3987.

Shaw, J.J., and Green, R. (2007). Two Distinct Components of Release Factor
Function Uncovered by Nucleophile Partitioning Analysis. Mol. Cell *28*, 458–467.

Sund, J., Ander, M., and Aqvist, J. (2010). Principles of stop-codon reading on the
ribosome. Nature *465*, 947–950.

Zavialov, A.V., Mora, L., Buckingham, R.H., and Ehrenberg, M. (2002). Release of
Peptide Promoted by the GGQ Motif of Class 1 Release Factors Regulates the
GTPase Activity of RF3. Mol. Cell *10*, 789–798.

Figure & Tables

Table 1 Comparison of Michaelis-Menten parameters and accuracy measurements of methylated and non-methylated class I release factors. The *E. coli* K12 genes encoding RF1 (*prfA*) and RF2 (*prfB*) were already available as clones in the pTRC and pET11a vectors, respectively. The final construct pET11a-prfB encoded a prfB[T246A] variant to enable the overexpression of an otherwise toxic RF2 (Uno et al., 1996). Both constructs contained C-terminal His$_6$-tags. The release factor methyltransferase *hemK* (gene: *prmC*), was available as a clone in pACDuet1; the protein contained no tag. All constructs were under the control of an Isopropyl β-D-1-thiogalactopyranoside (IPTG) inducible *lac* promoter. Plasmids pTRC-prfA and pET11a-prfB were introduced separately into *E. coli* BL21(DE3)Gold cells (Agilent Technologies), already harbouring pACDuet1-prmC, via chemical transformation. Protein production was induced with 1 mM IPTG in exponential phase cultures grown at 37 °C in LB medium. To allow full methylation of the class I release factors, cells were further grown for 4 h. The bacteria were lysed using a French press and purified using HisTrap nickel affinity chromatography (GE Healthcare) per manufacturer's instructions. Purified proteins were concentrated, dialyzed against 1 L of Polymix Buffer (pH 7.5) (Indrisiunaite et al., 2015; Jelenc and Kurland, 1979), with one buffer exchange after 3 h, and stored at -80 °C until use. Methylation of the release factors was verified by mass spectrometry as described earlier (Heurgue-Hamard et al., 2002). For the *in vitro* assays, the experiments were prepared and performed as described earlier (Korkmaz and Sanyal, 2017) with the following mRNAs 5′ GGG AAU UCG GGC CCU UGU UAA CAA UUA AGG AGG UAU ACU <u>AUG</u> STOP CUG CAG (A)$_{21}$ 3′ (the start codon is underlined, and STOP indicates the position of the stop codon).

		Cognate UGA		Non Cognate UGA		Accuracy Ratio
mRF1	kcat (s-1)	1.7	+/- 0.03	0.620	+/- 0.05	
	KM (nM)	5	/	164	+/- 3.0E-8	
	MM (s-1 nM-1)	0.340		0.004		
	Accuracy		**89.9**			3.63
RF1	kcat (s-1)	0.372	+/- 0.005	0.120	+/- 0.008	
	KM (nM)	5	/	40	+/- 1.8E-8	
	MM (s-1 nM-1)	0.074		0.003		
	Accuracy		**24.8**			
		UAA		UAG		
mRF2	kcat (s-1)	12*	+/- 1.1	1.490	+/- 0.1	
	KM (nM)	123*	+/- 2.9E-8	1045	+/- 3.42E-7	
	MM (s-1 nM-1)	0.097*		0.001		
	Accuracy		**68.4**			5.23
RF2	kcat (s-1)	0.75	+/- 0.03	0.560	+/- 0.03	
	KM (nM)	26	+/- 5.5E-9	254	+/- 3.5E-8	
	MM (s-1 nM-1)	0.029		0.002		
	Accuracy		**13.1**			

235 *Data from (Korkmaz and Sanyal, 2017).

Figure 1 Michaelis-Menten graphs of class I release factors on cognate and non-cognate substrates. (A) mRF1 on UAA (B) RF1 on UAA (C) mRF1 on UGA (D) RF1 on UGA

Figure 2 Michaelis-Menten graphs of class I release factors on cognate and non-

cognate substrates.(A) mRF2 (Korkmaz and Sanyal, 2017) on UAA (B) RF2 on UAA

(C) mRF2 on UAG (D) RF2 on UAG

Figure 3 Lineweaver-Burk plots of kinetic data from Table 1. (A) Lineweaver-Burk plot comparison of methylated and non-methylated RF1 kinetic data on UAA. For illustration purposes, the K_M value for RF1 was evaluated as observed in Table 1. (B) Comparison of mRF1 and RF1 on UAG. (C) Comparison of mRF2 and RF2 on UAA. (D) Comparison of mRF2 and RF2 on UAG. All K_M values are given in nM.