Title: **Frequent Non-random Shifts in the Temporal Sequence of Developmental**

Landmark Events during Fish Evolutionary Diversification

Running head: **Evolution of fish developmental sequence**

Authors

Fumihiro Ito[1,2] *, Tomotaka Matsumoto[2,3], Tatsumi Hirata[2,4]

[1]Mammalian Genetics Laboratory, Genetic Strains Research Center, National Institute of

Genetics, Mishima 411-8540, Shizuoka, Japan

[2]Department of Genetics, Graduate University for Advanced Studies (SOKENDAI), Mishima

411-8540, Shizuoka, Japan

[3]Division of Evolutionary Genetics, Department of Population Genetics, National Institute of

Genetics, Yata 1111, Mishima, Shizuoka 411-8540, Japan.

[4]Division of Brain Function, National Institute of Genetics, Yata 1111, Mishima 411-8540,

Shizuoka, Japan.

*Correspondence author

Authors' current address:

Fumihiro Ito

Mammalian Genetics Laboratory, Genetic Strains Research Center, National Institute of

Genetics, Mishima, Shizuoka 411-8540, Japan. Tel: +81-55-981-6816, FAX:+81-55-981-6817.

Email address: fumihiro.0301@gmail.com

Abstract

Morphology is a consequence of sequentially occurring developmental events, termed

developmental sequence, and evolutionary changes in the sequence can generate morphological

diversities. In this study, we examined evolutionary dynamics of the developmental sequence at

a macro-evolutionary scale using the teleost fish. From the previous literature describing

development of 31 fish species, we extracted the developmental sequences of 20 landmark

events involving the whole body plan, and by using them, reconstructed ancestral

developmental sequences. The phylogenetic comparisons of these sequences revealed

event-dependent heterogeneity in the frequency of changing the sequences. We then determined

potential event shifts that can parsimoniously explain the sequence changes on each node of the

fish phylogenetic tree. These "heterochronic shifts" are widely distributed on almost of all the

branches across the fish phylogeny. The simulation-based analysis indicated that this

distribution of heterochronic shifts is not the result of random accumulation over phylogenetic

time, but exhibits a curious constant trend so that individual phylogenetic branches harbor

similar numbers of heterochronic shifts regardless of length. It is of great interest to know how

these findings are related to morphological divergence in animals during evolution.

1. Introduction

The morphology of each multicellular organism is constructed by a fixed temporal sequence of

developmental events, termed developmental sequence. Because development is an inherently

step-by-step process, one might assume that the temporal sequence is not readily changeable

and is phylogenetically conserved among closely related species that share morphological

characteristics. Along these lines, if an evolutionary change occurs in the developmental

sequence, it could bring about a significant impact on animal body plan and lead to

morphological diversities. Indeed, previous comparisons of developmental sequences have

detected rare epoch-making changes that can provide morphological uniqueness to one species

that is different from the others (Strauss, 1990; Jeffery, Richardson, Coates & Bininda-Emonds,

2002; Maxwell, Harrison & Lasson, 2010) supporting the idea that the developmental sequence

is basically a conserved trait in the phylogenetic history.

56   Regarding evolution of the developmental sequences, another influential factor would

be the phylotypic period (Duboule, 1994). This is the developmental time frame during which

evolutionally distant animal species resemble each other in appearance. The well-accepted

hourglass-like model defines the phylotypic period as the middle phase of ontogenic

development, typically known as the pharyngulal stage. Recent transcriptome analyses have

indeed confirmed that inter-species diversities are kept to the minimum during this embryonic

stage (Kalinka et al., 2010; Irie & Kurtani, 2011), suggesting some unknown biological reasons

underlying this curious regularity. Although the phylotypic period was originally proposed by

the morphological resemblance, very few morphological analyses have actually been conducted

on species similarities during the period.

66   To explore the role for the developmental sequence in animal morphological evolution,

the critically missing information is empirical evaluation of evolutionary changes that had

actually occurred in the developmental sequences. In particular, very few systematic

comparisons have been made on the sequences of a wide range of developmental events that

cover the whole body plan in any class of animals. Therefore, we have few clues about how

commonly or rarely the developmental sequences of these animals had changed during their

evolutionary history. In the last several decades, comparative methods for developmental

sequences have been developed by several groups (Nunn & Smith, 1998; Jeffery,

Bininda-Emonds, Coates & Richardson, 2002; Jeffery, Bininda-Emonds, Coates & Richardson,

2005; Harrison & Larsson, 2008; Germain & Laurin, 2009). These methods compare the

relative order of developmental events among different species and successfully detected

potential evolutionary shifts of the events in a parsimonious manner, namely "heterochronic

shifts" in developmental sequences (Schoch, 2006; Smirthwaitet, Rundle, Bininda-Emonds &

Spicer, 2007; Sanchez-Villagra, Goswami, Weisbecker, Mock & Kuratani, 2008; Laurin, 2014;

Carril & Tambussi, 2017). Although most of these analyses have so far focused on the

developmental sequences for a particular organ or limited body part, we considered that the

methods themselves are similarly applicable to a global analysis of developmental sequences for

the whole body plan.

In this study, we conducted a comprehensive survey of developmental sequences using

teleost fish. The teleost fish is the largest group of vertebrates. Its abundant group members are

characterized by great morphological diversities (Nelson, Grande & Wilson, 2016) and, at the

same time, share the common characteristics of the fish body plan such as vertebrae, eyes,

medial fins and swim bladders (Romer & Parsons, 1986). Owing to the popularity as

developmental research materials, there are well-established staging tables for many fish species

that cover common clear-cut developmental landmarks. Hence, the teleost fish can provide an

ideal dataset for systematic analyses of the early developmental sequences. Among the

widely-used developmental landmarks, we chose 20 events that individually contribute to

distinct body parts across the whole body plan. Using the dataset of 31 different fish species, we

compared the developmental sequences and reconstructed their ancestral sequences over the

phylogenetic tree. Our analysis indicated that the developmental sequences are in fact frequently

changeable during the course of evolution, and that these changes are associated with the two

following characteristics. (1) The frequency of sequence changes differs widely depending on

the developmental event. (2) Heterochronic shifts are not simply accumulated along the

phylogenetic time, but their number is kept more or less constant in individual phylogenetic

branches regardless of length. We will discuss the potential implications of these findings.

2. Materials and Methods

2.1 Construction of fish phylogenetic tree

The overall topology of the phylogenetic tree followed the molecular phylogenetic relationship

reported previously by Near et al. (2012, 2013). The minor branches missing in the tree were

inserted based on the phylogenetic data obtained from Saitoh et al. (2011) and Yang et al.

(2015) for Cypriniformes, Perez et al. (2007) and Friedman et al. (2013) for Cichliformes, and

Pohl, Milvertz, Meyer & Vences. (2015) for Cyprinodontiformes. The divergent times were

determined using the public database TIMETREE, *The Timescale of Life* (Hedes & Kumar,

2009) (Supplemental file.1).

 2.2 Data sampling

Information about the temporal sequence of developmental events was extracted from 31

published research articles that describe normal fish development (Figure 1). The 20

developmental events used in this study were the first recognitions of blood circulation (bc),

caudal fin ray (cfr), eye pigmentation (ep), embryonic shield (es), first somite (fs), hatch (h),

heart beat/pulsing (hb), Kupffer's vesicle (kv), lens or lens placode/primodium (le), medial

finfold (mff), mouth opening (mo), olfactory vesicle/pit/placode (olf), otolithes (oto), otic

vesicle/placode/primodium (ot), optic vesicle/placode/primodium (op), pectoral fin bud (pfb),

swim bladder (sw), tail bud (tb), three brain regionalization (tbr), and tail lift from yolk (tl).

According to information in the articles, temporal orders of the developmental events were

ranked (Figure 1). When the article did not describe a developmental event, the event was

treated as a missing datum.

2.3 Rank analyses of developmental events

The raw ranks of individual developmental events were determined for the developmental

sequences of extant fish (Figure 1) and the ancestral developmental sequences reconstructed as

described below. The raw ranks were then normalized by the total number of the ranked events

(r_{max}) in each species, resulting in the relative scaling of the ranks in the range between $1/r_{max}$

and 1 in all the species (Weisbecker, Goswami, Wroe & Sanchez-Villagra, 2008). To quantify

variation of the ranks among the developmental sequences, pairwise distances in the ancestral

ranks between all pairs of the sequences were summed and averaged for each pair of

combinations.

2.4 Reconstruction of ancestral developmental sequences

We used the event-pairing method developed by Jeffery et al. (2002). In brief, all of the 190

pairs of developmental events in each species were scored based on the relative timing; when

one event occurs earlier, simultaneously or later compared with another event, the timing was

rated as 0, 1, or 2, respectively. By comparing the scored event-pairing matrices of different

species, the ancestral event-pairing matrix was reconstructed at each node of the fish

phylogenetic tree in a parsimonious manner under accelerated transformation (acctran) and

delayed transformation (deltran) optimizations using PAUP* software (Swofford, 2002). The

ancestral event-pairing matrices were then re-converted to the ancestral developmental

sequences (Supplemental file.2).

2.5 Detection of heterochronic shifts in fish phylogenetic tree

The heterochronic shifts between two developmental sequences at each phylogenetic node were

detected using the Parsimov algorithm developed by Jeffery et al. (2005). This parsimony-based

algorithm determines the minimum number of event shifts that can explain the difference

between two developmental sequences. Following the instructions, we implemented a Perl

script, Parsimv7g.pl, with PAUP* output log file in both acctran and deltran optimizations, and

mapped the detected heterochronic shifts onto the fish phylogeny (Supplemental files 3).

2.6 Simulation-base analyses

We examined whether the estimated number of heterochronic shifts in each branch can be

simply explained by random accumulation in the phylogenetic tree. The simulation was based

on a simple assumption that a heterochronic shift occurs at a constant rate per unit time and

therefore, accumulates in proportion to branch length in the phylogenetic tree. In this simulation,

we did not consider the event-dependent differences in the shift frequencies. The simulation

randomly distributed the estimated heterochronic shifts over the fish phylogenic branches solely

depending on their branch lengths. The simulation was replicated 100,000 times to obtain the

expected distribution of heterochronic shifts in each branch under the assumption of random

accumulation. The distribution of heterochronic shifts was then compared with the actual

distribution of the hetrochronic shifts in the fish phylogenetic branches. In this study, we used

year as the time scale of the branch length. However, in some analyses, we converted the time

scale to generation by considering the average generation times of individual fish species

(Supplemental file 4) and confirmed the consistency of the results.

3. Results

3.1 Phylogenetic relationship of 31 fish

For the present analyses, we used 30 teleost fish belonging to 13 distinct orders as the in-group,

because the developmental sequences of these fish have been well-documented in previous

articles (Figure 1). As an out-group, the amiadae fish, *Amia calva*, was used because it retains

ancestral morphological characteristics and because a recent molecular analysis confirmed its

location as the out-group of teleost fish (Near et al., 2012). In the constructed teleost

phylogenetic tree, the examined fish species were widely distributed and represented distinct

branches of teleost clade in a fairly unbiased manner (Figure 2). Because fish development in

the marine environment has rarely been documented, the fish species covered in this study were

basically freshwater fish, but also included several anadromous fish, such as *Gasterosteus*

aculeatus, which develop in freshwater but migrate between the sea and freshwater in their adult

life cycles.

3.2 Comparison of temporal orders of developmental events among fish

We selected 20 developmental events that consistently appeared as landmarks in the

developmental staging of many fish species (Figure 1). For this selection, in the hope to gain a

global picture of developmental sequences for the whole body plan, we intentionally included

events that belong to substantially different biological systems and contexts; e.g., the ones that

originate from different germ layers or that give rise to different cell types or separate body

parts. Additionally, the list also included a small number of seemingly interrelated events such

as formations of optic vesicle/placode/primodium (op), lens/lens placode (le) and eye

pigmentation (ep). We gathered information about these 20 events from the articles reporting

the development of 31 fish, and ranked the orders of individual events in the temporal sequence

for each species (Figure 1).

We first compared rank orders of each event among 30 in-group fish species. To

minimize effects of simultaneous occurrence of events and missing data on the comparison, the

raw ranks (Figure 1) were rescaled to normalized ranks that fit within the same range in all the

fish species (see Methods). Figure 3a shows distribution of the normalized ranks for individual

developmental events, which were horizontally arranged according to the average values.

Interestingly, the ranges of variations in the rank widely differed depending on the event. One

extreme case was embryonic shield (es), which always appeared first in the developmental

sequences obtained from the 29 fish species with no variation, excluding one missing

description in *Galaxias maculatus* (Figure 1). In contrast, relatively large variations in the rank

were observed for the appearance of Kupffer's vesicle (kv), hatch (h), medial finfold (mff) and

swim bladder (sb), suggesting that these events can more easily change their temporal orders in

the developmental sequence (Figure 3a).

To explore the evolutionary history of developmental sequences, we next

reconstructed ancestral developmental sequences at each node of the phylogenetic branches by

using the event-pairing method (Jeffery et al., 2002). This algorithm compares the relative

orders of all the event pairs between two different developmental sequences and generates the

ancestral sequences determined as a parsimonious solution under acctran and deltran

optimizations (Supplemental file 2). Using the obtained ancestral developmental sequences, we

compared the normalized ranks of individual events as we did in Figure 3a. Overall, the rank

orders of individual events in the ancestral developmental sequences (Figure 3b and 3c) were

quite similar to those in the extant fish sequences (Figure 3a); when the developmental events

were horizontally aligned in the same order of the extant average ranks, there were only a few

inversions in the order of two successive events at the average level (e.g., the order between first

somite (fs) and tail bud (tb)). Because this sequence reconstitution was based on parsimony, the

variations of estimated ranks were kept to nearly minimum. Still, individual events exhibited a

similar trend of rank variations to that observed in the extant fish sequences, further confirming

the idea that some developmental events change their orders more frequently than the others

during evolution.

Because the rank seemed to fluctuate depending on the event, we more

systematically analyzed the size of variations of the ranks. As an index of rank changeability,

the pairwise rank distances between all pairs of the ancestral developmental sequences were

measured and represented as the average value for each pair (Figure 4). Comparable values

were obtained by acctran and deltran optimizations (Spearman's rank correlation for the two

optimizations; r = 0.839). When the events were temporally arranged along the standard

ontogenic time frame defined as the average rank orders in the extant fish, the rank

changeability was found to be squeezed in the middle phase of the developmental sequence,

involving three brain regionalizations (tbr), otic vesicle/placode/primodium (ot) and lens or lens

placode/primodium (le) (Figure 4). The medial finfold (mff) around late-tail bud stage, in

contrast, recorded the largest rank changeability.

We then focused on the actual sequence of developmental events. Figure 5 shows the

percentage of the sequences in which one event (shown in row) occurs later than another

(shown in column) among the 30 extant in-group fish. In general, the sequence of two

temporally distant events was quite conservative, with no reversal in the order in many

combinations, whereas the neighboring events more frequently change their orders. If a closer

look was given to the sequence of anatomically interrelated events, the temporal order of the

optic vesicle/placode/primodium (op) and the lens/lens placode (le) was fixed in all the fish

species, and that of the lens/lens placode (le) and the eye pigmentation (ep) was almost fixed

except for one sequence reversal in *Heterobranchus bidorsalis*. Another interesting trend was

about the timing of hatch (h), which often changed the orders with the three late events, mouth

opening (mo), swim bladder (sb) and caudal fin ray (cfr). Similar results were obtained from the

comparison of event orders in ancestral developmental sequences (Figure S1).

3.3 Distribution of heterochronic shifts across the fish phylogenetic tree

Using the widely-used Parsimov algorithm (Jeffery et al., 2005), we next searched for

heterochronic shifts of the events that can explain the changes from one sequence to another at

every node of the fish phylogenetic tree. Although this is a parsimony-based algorithm and

therefore estimates the minimum number of event shifts, we detected 184 (acctran), 179

(deltran) and 94 (conserved between acctran and deltran) heterochronic shifts in total

(Supplemental file 3). The fish phylogenetic tree has 60 branches in total. When the detected

shifts were mapped over the phylogenetic tree, multiple heterochronic shifts were observed on

almost all the branches (Figure 2, Supplemental file 3). For example, in the relatively long

branch of order Salmoniformes, five and three heterochronic shifts were detected by acctran and

deltran optimizations, respectively.

Because a substantial number of heterochronic shifts were detected widely across the

fish phylogeny, we wondered whether these shifts might happen rather frequently and be

randomly accumulated over the evolutionary history. To address this question, we took a

simulation-based approach. Given that a heterochronic shift occurs at a random stochastic

manner and is neutrally accumulated, we simulated the expected distribution of the number of

heterochronic shifts, of which the number was nearly proportional to the phylogenetic branch

length (white circles in Figure 6a and 6b). By contrast, the actual distribution of heterochronic

shifts detected by the Parsimov analysis was much more constant, regardless of the branch

length in both acctran and deltran optimizations (black circles in Figure 6a and 6b). Coefficient

of variation of the number of heterochronic shifts across the branches also showed smaller

values for the experimental dataset than for the simulation data (Figure 6c), indicating that

branch-by-branch fluctuations of the number of heterochronic shifts are actually more limited

compared with the values expected under simulation. In addition, the number of the

phylogenetic branches that harbored no heterochronic shifts was significantly smaller for the

experimental dataset than that for the simulation data (Figure 6d). Because inclusion of an

extremely long branch could skew the statistical results, we performed the same statistical

comparison using only relatively short branches ($\leq$ 50Mya and $\leq$ 20Mya). These analyses again

showed similar results, indicating that the number of heterochronic shifts per branch is more

constant than the expectation under the assumption of random accumulation (Figure 6c, 6d, and

Figure S2). Replacing the phylogenic time scale with the generation number basically did not

qualitatively affect the results of the analyses (Figures S3 and S4).

The heterochronic shifts of developmental events are sometimes associated to

differentiation of terminal phenotypes (Gunter, Koppermann & Meyer, 2014). Thus, we

examined the topological distribution of the heterochronic shifts by separately examining

internal and terminal branches. In both of the branch types, the numbers of actual heterochronic

shifts were basically in the range of the expected numbers in the simulation (Figure 6e and 6f).

Significant differences were only exceptionally observed in the all branch category under the

acctran optimization; however, we cannot rule out the possibility that the inclusion of extremely

long branches in this category affected the results. In conclusion, this analysis did not positively

support a preferential occurrence of heretochronic shifts in either the external or internal

branches.

4. Discussion

The present study provides empirical evidence that developmental sequences are in fact

changeable during evolution; the extant fish species clearly involve historic signs showing that

their ancestors had experienced dynamic and frequent rearrangement of the developmental

sequences. This finding may not be exactly concordant with the traditional view that the

developmental sequence is a phylogenetically conserved trait, which provides the blueprint for

the common body plan among related species. One reason is probably our wide selection of

developmental events; we intentionally took up the events that cover a whole variety of

embryonic origins, cell types, body parts and biological systems, aiming for understanding the

global body plan. In contrast, the major focus of previous studies was in-depth understanding of

developmental sequences for a restricted body part or organ (Schlosser, 2008; Hautier et al.,

2011; Workman, Charvet, Clancy, Darlington & Finlay, 2013). Therefore, even though we only

analyzed one group of species that share the highly conserved body plan, rather frequent shifts

of the events could be observed. There is increasing evidence for modular control of formation

of different body parts (Klingenberg, Badyaev, Sowry & Beckwith 2001; Kawanishi et al.,

2013; Schmidt & Starck, 2010). This modular nature of individual body parts can underlie the

large fluctuations of developmental sequences observed in this study, and possibly contribute to

individual evolution of different body parts toward morphological diversification.

Although the heterochronic shifts detected in this study are widespread across fish

phylogeny, our simulation-based analyses uncovered certain regularity in the distribution.

Namely, the shifts are not randomly accumulated over the evolutionary time, but there appears

to be some force making the number of shifts constant in individual phylogenetic branches.

Currently, we cannot effectively interpret this perplexing result, and only note two potential

scenarios from different angles, which are not necessarily mutually exclusive. First, it may be

possible that the heterochronic shift is a branching-related process. If a new branch often comes

with new shifts, then the shift number should be more correlated with the branch number rather

than its length. If speciation processes are concerned, this is actually a very attractive scenario,

because the heterochronic shift can support differentiation of branch-specific phenotypes in

each species. Second, the seeming constancy of the shift number might be related to the limited

configuration of acceptable developmental sequences. Our event sequence analyses indeed

showed that only certain types of changes are acceptable in the developmental sequences

(Figure 5). This limitation probably stems from both developmental and evolutionary

constraints in order to construct a fit functional body plan. Yet, for the moment, we cannot

determine how the limitation of sequence configurations can shape the distribution of potential

heterochronic shifts, because they are limited, but still a great many acceptable sequences exist.

One interesting finding of this study is that some developmental events change their

temporal orders more drastically than others during evolution. Of particular note is the

emergence of medial finfold (mff), of which rank changeability was the highest among all the

events (Figure 4). The medial finfold is a morphogenetic field for fins. A recent study reports

that this single morphogenetic field actually is a mosaic composed of three distinct fin primodia

(Larouch, Zelditch & Cloutier, 2017). Thus, it is possible that the three primodia behaved as

independent modules during evolution, thereby expanding the temporal range of this event.

Another interesting example is the timing of hatch (h), which is relatively easily changeable

with the three following developmental events, mouth opening (mo), swim bladder (sb) and

caudal fin ray (cfr). All these events are directly related to the life strategy of how a fish

survives during the larval stage, and therefore changing the orders may be advantageous under

some circumstances, particularly when fish have to adapt to a new environment (Miller &

Kendall, 2009). Indeed, the temporal shift of birth timing has been regarded as a symbolic

example of "heterochrony", an evolutionary force based on maneuvering developmental

machinery (Keyte & Smith, 2012; Dail, Reznick & Brainerd, 2017; Botelho, Smith-Paredes &

Vargas, 2015; Werneburg, Laurin, Koyabu & Sanchez-Villagra, 2016).

When the developmental events were aligned along the ontogenetic sequence, the rank

changeability was squeezed in the middle phase of the early development involving three brain

regionalization (tbr), otic placode/primodium (ot), and lens formation (le) (Figure 4). These

events are typical characteristics of the conserved phylotypic stage defined by the hourglass

model (Richardson, 1995; Irie, 2017). The hourglass model has been gaining increasing support

from the recent transcriptome analyses but still lacks sufficient evidence from objective

morphological analyses. Although the relationship between the developmental sequences and

morphological similarity is not very straightforward, our results seem to provide another

support for the hourglass model from the morphological point of view.

There is a common observation that the external temperature affects developmental

time frames (Schmidt & Starck, 2010; Mabee, Olmstead & Cubbage, 2000). Because most fish

reproduce by external fertilization and the embryos develop under fluctuating temperatures,

temporal shifts of individual developmental events might sometimes occur in fish under the

natural environment. Indeed, a study reports that the developmental sequence is polymorphic

even in one fish species (de Jong, Colbert, Witte & Richardson, 2009). Thus, fish may be

relatively tolerant to sporadic shifts of developmental events in the natural ontogeny, and

frequent encounters with such situations somehow increase the chances that some fish adopt a

shift in a persistent manner. Currently, it is not clear whether such fish-specific circumstances

are reflected in the present results. Future systematic analyses using other groups of animals

will address this issue.

Acknowledgements

We thank Dr. Toshihiko Shiroishi, Dr. Yasushi Hiromi, Dr. Naoki Irie, Dr. Takanori Amano, Dr.

Yuuta Moriyama and Dr. Kousuke Mouri for comments that helped greatly improve the

manuscript. We also thank Dr. Erin E. Maxwell and Dr. Luke B. Harrison for their advice on

parsimony analyses.

Reference

Arakawa, T., Kanno, Y., Akiyama, N., Kitano, T., Nakatsuji, N., & Nakatsuji, T. (1999). Stages

of embryonic development of the ice goby (shiro-uo), *Leucopsarion petersii*. *Zoological*

Science, 16, 761-773. doi.org/10.2108/zsj.16.761

Armstrong, P. B., & Child J. S., (1965). Stages in normal development of *Fundulus heroclitus*.

The Biological Bulletin, 128, 143-168. doi:10.2307/1539545

Ballard, W. W. (1969). Normal embryonic stages of *Gobius niger jozo*. *Pubbl Staz Zool Napoli*,

37, 1-17.

Ballard, W. W. (1973). Normal embryonic stages for salmonid fishes, based on Salmo gairdneri

Richardson and *Salvelinus fontinalis* (Mitchill). *Journal of Experimental Zoology Part A:*

Ecological and Integrative Physiology, 184, 7-26. doi: 10.1002/jez.1401840103

Ballard, W. W., (1986). Morphogenetic movements and a provisional fate map of development

in the holostean fish, *Amia calva*. *Journal of Experimental Zoology Part A: Ecological and*

Integrative Physiology, 238, 355-372. doi:10.1002/jez.1402380308

Balon,, E. K. (1977). Early ontogeny of *Labeotropheus Ahl*, 1927 (Mbuna, Cichlidae, Lake

Malawi), with a discussion on advanced protective styles in fish reproduction and development.

Environmental Biology of Fishes, 2, 147-176.

Basak, S. K., Basak, B., Gupta, N., Haque, M. M., & Amin, R. (2014). Embryonic and larval

development of silver barb (*Barodes gonionotus*) in a mobile hatchery under laboratory

Condition. *European Scientific Journal, Special ed*, 3, 258-270.

Benzie, V. (1968). Stages in the normal development of *Galaxias maculatus attenuatus*

(Jenyns). *New Zealand Journal of Marine and Freshwater Research*, 2, 606-627.

Botelho, J. F., Smith-Paredes, D., & Vargas A. O. (2015). Altriciality and the Evolution of Toe

Orientation in Birds. *Journal of Evolutionary Biology*, 42,

502-510. doi: .org/10.1007/s11692-015-9334-7

Carril, J., & Tambussi, C. P. (2017). Skeletogenesis of *Myiopsitta monachus* (Psittaciformes)

and sequence heterochronies in Aves. *Evolution & Development*, 19, 17-28.

doi:10.1111/ede.12211

Cunningham, J. E. R., & Balon, E. K. (1985). Early ontogeny of *Adinia xenica* (Pisces,

Cyprinodontiformes): 1. The development of embryos in hiding. *Environmental Biology of*

Fishes, 14, 115-166.

de Jong, I. M., Colbert, M. W., Witte, F., & Richardson, M. K. (2009). Polymorphism in

developmental timing: intraspecific heterochrony in a Lake Victoria cichlid. *Evolution &*

Development, 11, 625-635. doi:10.1111/j.1525-142X.2009.00370.x

de Jong, I. M., Witte, F., & Richardson, M. K. (2009). Developmental stages until hatching of

the Lake Victoria cichlid *Haplochromis piceatus* (Teleostei: Cichlidae). *Journal of Morphology*,

270, 519-535. doi:10.1002/jmor.10716

Dial, T. R., Reznick, D. N., & Brainerd, E. L. (2017). Heterochrony in the evolution of

Trinidadian guppy offspring size: maturation along a uniform ontogenetic trajectory.

Proceedings of the Royal Society B: Biological Sciences, 284, doi:10.1098/rspb.2017.1319

Duboule, D. (1994). Temporal colinearity and the phylotypic progression: a basis for the

stability of a vertebrate Bauplan and the evolution of morphologies through heterochrony.

Development (suppl.), 135-142.

Friedman, M., Keck, B. P., Dornburg, A., Eytan, R. I., Martin, C. H., Hulsey, C. D., . . . Near, T.

J. (2013). Molecular and fossil evidence place the origin of cichlid fishes long after Gondwanan

rifting. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20131733.

doi:10.1098/rspb.2013.1733

Fujimura, K., & Okada, N. (2007). Development of the embryo, larva and early juvenile of Nile

tilapia *Oreochromis niloticus* (Pisces: Cichlidae). Developmental staging system. *Development*

Growth & Differentiation, 49, 301-324. doi:10.1111/j.1440-169X.2007.00926.x

Germain, D., & Laurin, M. (2009). Evolution of ossification sequences in salamanders and

urodele origins assessed through event-pairing and new methods. *Evolution & Development*, 11,

170-190. doi:10.1111/j.1525-142X.2009.00318.x

Gunter, H. M., Koppermann, C., & Meyer A. (2014). Revisiting de Beer's textbook example of

heterochrony and jaw elongation in fish: calmodulin expression reflects heterochronic growth,

and underlies morphological innovation in the jaws of belonoid fishes. *Evodevo*, 5, 8.

doi:10.1186/2041-9139-5-8

Hall, T. E., Smith, P., & Johnston, I. A. (2004). Stages of Embryonic Developmsnt in the

Atlantic Cod *Gadus morhua*. *Journal of Morphology*, 259, 255-270. doi: 10.1002/jmor.10222

Harrison, L. B., & Larsson, H. C. (2008). Estimating evolution of temporal sequence changes: a

practical approach to inferring ancestral developmental sequences and sequence heterochrony.

Systematic Biology, 57, 378-387. doi:10.1080/10635150802164421

Hautier, L., Weisbecker, V., Goswami, A., Knight, F., Kardjilov, N., & Asher, R. J. (2011).

Skeletal ossification and sequence heterochrony in xenarthran evolution. *Evolution &*

Development, 13, 460-476. doi:10.1111/j.1525-142X.2011.00503.x

Hedges, S. B. & Kumar, S. (2009). The timetree of life. New York, NY: Oxford University

Press.

Humphrey, C., Klumpp, D. W., & Pearson, R. (2003). Early development and growth of the

east rainbowfish, *Melanotaenia splendida splendida. Marine and Freshwater Research*, 54,

17-25. doi: org/10.1071/MF02037

Irie, N. (2017). Remaining questions related to the hourglass model in vertebrate evolution.

Current Opinion in Genetics & Development, 45, 103-107. doi:10.1016/j.gde.2017.04.004

Irie, N., & Kuratani, S. (2011). Comparative transcriptome analysis reveals vertebrate

phylotypic period during organogenesis. *Nature Communications*, 2, 248.

doi:10.1038/ncomms1248

Iwamatsu, T. (2004). Stages of normal development in the medaka *Oryzias latipes*. *Mechanisms*

of Development, 121, 605-618. doi:10.1016/j.mod.2004.03.012

Iwamatsu, T., & Hirata, K. (1984). Normal course of development of the Java medaka, *Oryzias*

javanicus. *Bull Aichi Univ Edu* (Nat Sci), 33, 87-109.

Jeffery, J. E., Bininda-Emonds, O. R., Coates, M. I., & Richardson, M. K. (2002). Analyzing

evolutionary patterns in amniote embryonic development. *Evolution & Development*, 4,

292-302. doi: 10.1046/j.1525-142X.2002.02018.x

Jeffery, J. E., Bininda-Emonds, O. R., Coates, M. I., & Richardson, M. K. (2005). A new

technique for identifying sequence heterochrony. *Systematic Biology*, 54, 230-240.

doi:10.1080/10635150590923227

Jeffery, J. E., Richardson, M. K., Coates, M. I., & Bininda-Emonds, O. R. (2002). Analyzing

developmental sequences within a phylogenetic framework. *Systematic Biology*, 51, 478-491.

doi:10.1080/10635150290069904

480 Kalinka, A. T., Varga, K. M., Gerrard, D. T., Preibisch, S., Corcoran, D. L., Jarrells, J., . . .

481 Tomancak, P. (2010). Gene expression divergence recapitulates the developmental hourglass

482 model. *Nature*, 468, 811-814. doi:10.1038/nature09634

484 Kawanishi, T., Kaneko, T., Moriyama, Y., Kinoshita, M., Yokoi, H., Suzuki, T., . . . Takeda, H.

485 (2013). Modular development of the teleost trunk along the dorsoventral axis and *zic1/zic4* as

486 selector genes in the dorsal module. *Development*, 140, 1486-1496. doi:10.1242/dev.088567

488 Keyte, A., & Smith, K. K. (2012). Heterochrony in somitogenesis rate in a model marsupial,

489 *Monodelphis domestica*. *Evolution & Development*, 14, 93-103.

490 doi:10.1111/j.1525-142X.2011.00524.x

492 Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B., & Schilling, T.F. (1995). Stages of

493 embryonic development of the zebrafish. *Developmental Dynamics*, 203, 253-310.

494 doi:10.1002/aja.1002030302

496 Klingenberg, C. P., Badyaev, A. V., Sowry, S. M., & Beckwith, N. J. (2001). Inferring

497 developmental modularity from morphological integration: analysis of individual variation and

498 asymmetry in bumblebee wings. *American Naturalist*, 157, 11-23. doi:10.1086/317002

500 Kratochwil, C. F., Sefton, M. M., & Meyer, A. (2015). Embryonic and larval development in

501 the Midas cichlid fish species flock (*Amphilophus spp*.): a new evo-devo model for the

investigation of adaptive novelties and species differences. *BMC Developmental Biology*, 15, 12.

doi:10.1186/s12861-015-0061-1

Larouche, O., Zelditch, M. L., & Cloutier, R. (2017). Fin modules: an evolutionary perspective

on appendage disparity in basal vertebrates. *BMC Biology*, 15, 32.

doi:10.1186/s12915-017-0370-x

Laurin, M. (2014). Assessment of modularity in the urodele skull: an exploratory analysis using

ossification sequence data. *Journal of Experimental Zoology Part B: Molecular and*

Developmental Evolution, 322, 567-585. doi:10.1002/jez.b.22575

Long, W. L, & Ballard, W. W., (1976). Normal Embryonic Stages of the White Sucker,

Catostomus commersoni. Copeia, 1976, 342-351. doi. 10.2307/1443957

Mabee, P. M., Olmstead, K. L., & Cubbage, C.C. (2000). An experimental study of intraspecific

variation, developmental timing, and heterochrony in fishes. *Evolution*, 54, 2091-2106

Marimuthu, K., & Haniffa, M. A. (2007). Embryonic and Larval Development of the Striped

Snakehead *Channa striatus. Taiwania*, 52, 84-92. dio:10.6165/tai

Maxwell, E. E., Harrison, L. B., & Larsson, H. C. (2010). Assessing the phylogenetic utility of

sequence heterochrony: evolution of avian ossification sequences as a case study. *Zoology*

(Jena), 113, 57-66. doi:10.1016/j.zool.2009.06.002

McElman, J. F., & Balon, E. K. (1979). Early ontogeny of walleye, *Stizostedion vitreum*, with

steps of saltatory development. *Environmental Biology of Fishes*, 4, 309-348.

Meijide, F. J., & Guerrero, G. A., (2000). Embryonic and larval development of a

substrate-brooding cichlid *Cichlasoma dimerus* (Heckel, 1840) under laboratory conditions.

Journal of Zoology, 252, 481-493. doi: 10.1111/j.1469-7998.2000.tb01231.x

Miller, B. S., & Kendall A. W. (2009). Early life history of marine fishes. Berkeley, Calif.,

University of California Press; xii, 364 p. p.

Near, T. J., Dornburg, A., Eytan, R. I., Keck, B. P., Smith, W. L., Kuhn, K. L., . . . Wainwright,

P. C. (2013). Phylogeny and tempo of diversification in the superradiation of spiny-rayed fishes.

Proceedings of the National Academy of Sciences of the United States of America, 110,

12738-12743. doi:10.1073/pnas.1304661110

Near, T. J., Eytan, R. I., Dornburg, A., Kuhn, K. L., Moore, J. A., Davis, M. P., . . . Smith, W. L.

(2012). Resolution of ray-finned fish phylogeny and timing of diversification. *Proceedings of*

the National Academy of Sciences of the United States of America, 109, 13698-13703.

doi:10.1073/pnas.1206625109

Nelson, J. S, Grande, T. C, & Wilson, M. V. H. (2016). Fishes of the world (5th ed). New York:

Wiley.

Nunn, C. L., & Smith, K. K. (1998). Statistical analyses of developmental sequences: the

craniofacial region in marsupial and placental mammals. *American Naturalist*, 152, 82-101.

doi:10.1086/286151

Olaniyi, W. A., & Omitogun, O. G. (2014). Embryonic and larval developmental stages of

African giant catfish *Heterobranchus bidorsalis* (Geoffroy Saint Hilaire, 1809) (Teleostei,

Clariidae). *Springerplus*, 3, 677. doi:10.1186/2193-1801-3-677

Pelluet, D. (1944) Criteria for the recognition of development stages in the salmon (*Salmo*

salar). *Journal of Morphology*, 74, 395-407. doi: 10.1002/jmor.1050740305

Perez, G. A., Rican, O., Orti, G., Bermingham, E., Doadrio, I., & Zardoya, R. (2007).

Phylogeny and biogeography of 91 species of heroine cichlids (Teleostei: Cichlidae) based on

sequences of the cytochrome b gene. *Molecular Phylogenetics and Evolution*, 43, 91-110.

doi:10.1016/j.ympev.2006.08.012

Pohl, M., Milvertz, F. C., Meyer, A., & Vences, M. (2015). Multigene phylogeny of

cyprinodontiform fishes suggests continental radiations and a rogue taxon position of

Pantanodon. *Vertebrate Zoology*, 65, 37-44.

Puvaneswari, S., Marimuthu, K., Karuppasamy, R., & Haniffa, M. A. (2009). Early embryonic

and larval development of Indian catfish, *Heteropneustes fossilis. EurAsian Journal of*

BioSciences 3, 84-96. doi: 10.5053/ejobios.2009.3.0.12

Richardson, M. K. (1995). Heterochrony and the phylotypic period. *Developmental Biology,*

172, 412-421. doi:10.1006/dbio.1995.8041

Romer, A. S. & Parsons, T. S. (1986). The vertebrate body (6th ed). Philadelphia, PA: Saunders

College Publications.

Saitoh K, Sado T, Doosey MH, Bart Jr HL, Inoue JG, Nishida M., . . . Miya M. (2011).

Evidence from mitochondrial genomics supports the lower Mesozoic of South Asia as the time

and place of basal divergence of cypriniform fishes (Actinopterygii: Ostariophysi). *Zoological*

Journal of the Linnean Society, 161, 633 – 662. doi:10.1111/j.1096-3642.2010. 00651.x

Sanchez-Villagra, M. R., Goswami, A., Weisbecker, V., Mock, O., & Kuratani, S. (2008).

Conserved relative timing of cranial ossification patterns in early mammalian evolution.

Evolution & Development, 10, 519-530. doi:10.1111/j.1525-142X.2008.00267.x

Schlosser, G. (2008). Development of the retinotectal system in the direct-developing frog

Eleutherodactylus coqui in comparison with other anurans. *Frontiers in Zoology*, 5, 9.

doi:10.1186/1742-9994-5-9

Schmidt, K., & Starck, J. M. (2010). Developmental plasticity, modularity, and heterochrony

during the phylotypic stage of the zebra fish, *Danio rerio*. *Journal of Experimental Zoology*

Part B: Molecular and Developmental Evolution, 314, 166-178. doi:10.1002/jez.b.21320

Schoch, R. R. (2006). Skull ontogeny: developmental patterns of fishes conserved across major

tetrapod clades. *Evolution & Development*, 8, 524-536. doi:10.1111/j.1525-142X.2006.00125.x

Shardo, J. D. (1995). Comparative embryology of teleostean fishes. I. Development and staging

of the American shad, *Alosa sapidissima* (Wilson, 1811). *Journal of Morphology*, 225,125-167.

doi:10.1002/jmor.1052250202

Smirthwaite, J. J., Rundle, S. D., Bininda-Emonds, O. R., & Spicer J .I. (2007). An integrative

approach identifies developmental sequence heterochronies in freshwater basommatophoran

snails. *Evolution & Development*, 9, 122-130. doi:10.1111/j.1525-142X.2007.00143.x

Strauss, R. E. (1990). Heterochronic Variation in the Developmental Timing of Cranial

Ossifications in Poeciliid Fishes (Cyprinodontiformes). *Evolution*, 44, 1558-1567.

doi:10.1111/j.1558-5646.1990.tb03846.x

Swarup, H. (1958). Stages of development of the stickleback *Gasterosteus aculeatus*. *Journal of*

Embryology and Experimental Morphology, 6, 373-383.

Swofford, D. L. (2002). PAUP*. Phylogenetic Analysis Using Parsimony (*and other methods),

v. 4beta10. Sunderland, MA: Sinauer.

Tavolga, W. N., & Rugh, R. (1947). Development of the platyfish, Platypoecilus maculatus.

Zoologica; Scientific Contributions of the New York Zoological Society, 32, 1-15.

Tsai, H. Y., Chang, M., Liu, S. C., Abe, G., & Ota, K. G. (2013). Embryonic development of

goldfish (*Carassius auratus*): a model for the study of evolutionary change in developmental

mechanisms by artificial selection. *Developmental Dynamics,* 242, 1262-1283.

doi:10.1002/dvdy.24022

Verma, P., (1970). Normal stages in the development of *Cyprinus carpio* var. communis L. *Acta*

Biologica Academiae Scientiarum Hungaricae, 21, 207-218.

Weisbecker, V., Goswami, A., Wroe, S., & Sanchez-Villagra, M. R. (2008). Ossification

heterochrony in the therian postcranial skeleton and the marsupial-placental dichotomy.

Evolution, 62, 2027-2041. doi:10.1111/j.1558-5646.2008.00424.x

Werneburg, I., Laurin, M., Koyabu, D., & Sanchez-Villagra, M. R. (2016). Evolution of

organogenesis and the origin of altriciality in mammals. *Evolution & Development,* 18, 229-244.

doi:10.1111/ede.12194

Workman, A. D., Charvet, C. J., Clancy, B., Darlington, R. B., & Finlay, B. L. (2013).

Modeling transformations of neurodevelopmental sequences across mammalian species. Journal

of Neuroscience, 33, 7368-7383. doi:10.1523/JNEUROSCI.5746-12.2013

Wourms, J. P. (1972). Developmental biology of annual fishes. I. Stages in the normal

development of *Austrofundulus myersi* Dahl. *Journal of Experimental Zoology Part A:*

Ecological and Integrative Physiology, 182, 143-167. doi:10.1002/jez.1401820202

Yang, L., Sado, T., Vincent Hirt, M., Pasco-Viel, E., Arunachalam, M., Li, J., . . . Mayden, R. L.

(2015). Phylogeny and polyploidy: resolving the classification of cyprinine fishes (Teleostei:

Cypriniformes). *Molecular Phylogenetics and Evolution*, 85, 97-116.

doi:10.1016/j.ympev.2015.01.014

Zalina, I., Saad, C. R., Christianus, A., & Harmin, S. A. (2012). Induced breeding and

embryonic development of climbing perch (*Anabas testudineus*, Bloch). *Journal of Fisheries*

and Aquatic Science, 7, 291-306. doi.org/10.5567/ECOLOGY-IK.2013.8.14

Author contribution

F.I. designed the study, performed the majority of the experiments, analyzed data and wrote the

manuscript. T.M contributed to designing and conducting the simulation. T.M. and T.H.

supervised the study and helped F.I. to write the manuscript with input from other people.

Competing interests

The authors have no competing interests.

Figure Legends

FIGURE 1 Temporal orders of developmental events in the 31 fish

The temporal sequence of developmental events was extracted from the reference listed for each

species. Abbreviations of developmental events are; bc: blood circulation, cfr: caudal fin ray,

ep: eye pigmentation, es: embryonic shield, fs: first somite, h: hatch, hb: heart beat/pulsing, kv:

kupffer's vesicle, le: lens or lens placode/primodium, mff: medial finfold, mo: mouth opening,

olf: olfactory vesicle/pit/placode, oto: otolithes, ot: otic vesicle placode/primodium, op: optic

vesicle/placode/primodium, pfb: pectoral fin bud, sw: swim bladder, tb: tail bud, tbr: three brain

regionalization, and tl: tail lift from yolk. The ranks of missing data are marked by ?.

FIGURE 2 Phylogenetic relationships of the 31 fish

The phylogenetic tree of the 31 fish examined in this study. The asterisk marks the anadromous

fish, while all the others are freshwater fish. The numbers aside the branches indicate the

divergent times (Mya). In some representative branches, the numbers of heterochronic shifts

detected under acctran and deltran optimizations are shown in boxes.

FIGURE 3 Distribution of ranks of events in the developmental sequence

The boxplot shows the statistical distribution (minimum, first quartile, median, third quartile,

maximum and outliers) of normalized ranks for individual developmental events obtained from

the extant in-group 30 fish data (a) and reconstructed ancestral developmental sequences by

acctran (b) and deltran (c) optimizations. In all of the panels, the developmental events are

horizontally aligned from left to right according to average ranks in the extant fish sequences. In

the ancestral sequences (b, c), the average sequence is reversed between first somite (fs) and tail

bud (tb), between heart beats (hb) and olfactory vesicle/pit/placode (olf), and between blood

circulation (bc) and otolithes (oto). An additional inversion is observed between swim bladder

(sb) and caudal fin rays (cfr) in the deltran optimization (c).

FIGURE 4 Rank changeability of individual developmental events

The variation of the ranks is shown as the average value of pairwise rank distances, which are

calculated from all the pairs of ancestral developmental sequences reconstructed under acctran

(left) and deltran (right) optimizations. The events are arranged along the standard ontogenic

time frame defined by the average developmental sequence in extant fish (Figure 2) from top to

bottom. *significant differences (P<0.05) by Mann-Whitney U-test when comparing the values

of Kupffer's vesicle (kv) and three brain regionalization (tbr) and those of lens formation (le)

and tail lift from yolk (tl).

FIGURE 5 Sequence orders of event pairs in extant developmental sequences

The event sequence matrix represents all the pairwise combinations of developmental events.

The number shows the percentage of the sequences in which the row event occurs later than the

column event, and was calculated from the dataset of extant 30 fish excluding the missing event

data. The individual cells are differently heatmap color-coded depending on the percentage.

FIGURE 6 Distribution of heterochronic shifts in the fish phylogeny

(a,b) The relationship between the phylogenetic branch length and the number of herterochronic

shifts detected from the extant and ancestral developmental sequences (black circle) and

theoretically estimated by simulation (open circle) under acctran (a) and deltran (b)

optimizations. In the simulation, the branch length and the number of shifts are highly

correlated (Spearman's rank correlation coefficients; 0.9995 (acctran), 0.9995 (deltran)). (c) The

coefficient of variance for the number of heterochronic shifts in each branch. The black and

open circles show the experimental and simulated values, respectively. The vertical bars

indicate 95% confident intervals for the simulated value. The analysis was conducted with three

different branch categories: all branches, and the branches shorter than 50 and 20 million years

(Mys). (d) The number of branches with no heterochronic shifts calculated from experimental

(black circle) and simulation data (open circle) in three different branch length categories.

Vertical bars indicate 95% confident intervals of the simulated value. (e, f) The number of

heterochronic shifts detected in the external (e) and internal (f) branches calculated from

experimental (black circle) and simulation data (open circle). The branches are categorized into

three groups according to their lengths. Vertical bars indicate 95% confident intervals of the

simulated value.

Taxon	Common name	Reference	es	op	fs	tb	kv	tbr	ot	le	tl	hb	olf	mff	oto	bc	pfb	ep	h	mo	sb	cfr
Amia calva	Bowfins	Ballard, 1986	1	2	2	2	?	3	4	5	5	5	5	6	8	7	7	7	10	8	9	9
Alosa sapidissima	American shad	Shardo, 1995	1	4	2	5	?	3	5	5	6	7	5	6	7	7	7	8	10	9	?	11
Catostomus commersoni	White sucker	Long & Ballard, 1976	1	3	2	3	4	3	4	5	5	6	7	6	7	7	8	8	8	10	11	9
Danio rerio	Zebrafish	Kimmel et al., 1995	1	4	3	2	4	5	5	6	5	8	7	7	7	9	9	8	11	12	10	12
Cyprinus carpio	Common carp	Verma, 1970	1	3	4	?	5	2	5	6	6	7	7	11	8	8	9	8	10	13	12	14
Carassius auratus	Minnows	Tsai et al., 2013	1	3	3	2	4	4	5	5	5	6	7	5	?	?	6	6	8	8	9	?
Barbodes gonionotus	Silver barb	Basak et al., 2014	1	3	2	2	?	3	7	4	5	6	?	6	?	5	6	11	9	10	12	8
Heteropneustes fossilis	Stinging catfish	Puvaneswari et al, 2009	1	2	2	2	4	3	6	4	4	5	5	4	?	7	9	8	7	10	?	8
Heterobranchus bidorsalis	African catfish	Olaniyi & Omitogue, 2014	1	2	2	2	3	7	4	8	4	5	8	?	4	8	?	5	6	9	?	10
Oncorhynchus mykiss	Rainbow trout	Ballard, 1973	1	3	2	?	2	3	7	5	6	6	4	7	9	7	7	8	12	10	?	11
Salmo salar	Atlantic salmon	Pelluet, 1944, Gorodilov, 1996	1	3	2	6	2	?	4	5	?	7	8	8	9	8	9	10	12	6	13	11
Galaxias maculatus	Common galaxias	Benzie, 1968	?	1	2	1	2	1	3	2	5	2	3	7	8	?	6	4	10	10	?	9
Gadus morhua	Atlantic cod	Hall et al., 2004	1	2	1	3	3	3	4	6	?	7	5	4	5	8	8	9	10	12	11	11
Gobius niger	Black goby	Ballard, 1969	1	2	2	3	3	6	5	4	4	8	?	5	7	9	8	8	10	11	9	12
Leucopsarion petersii	Ice goby	Arakawa et al., 1999	1	3	3	2	3	4	5	6	5	7	?	?	9	8	10	8	13	11	12	?
Gasterosteus aculeatus	Three-spined stickleback	Swarup, 1958	1	3	4	?	5	2	5	5	?	6	6	?	6	8	9	7	11	10	12	12
Stizostedion vitreum	Walleye	McElman & Balon, 1979	1	2	3	2	3	4	4	5	6	7	4	6	8	9	11	10	13	14	15	12
Channa striatus	Striped snakehead	Marimuthu & Haniffa, 2007	1	2	2	2	4	3	3	4	4	5	5	4	6	5	7	?	6	8	7	9
Anabas testudineus	Climbing gouramies	Zalina et al., 2012	1	3	4	?	2	4	4	5	?	6	?	7	7	7	?	10	8	9	9	11
Amphilophus xiloaensis	Cichlids	Kratochwil et al., 2015	1	2	2	2	?	5	2	4	3	5	?	?	?	7	8	8	6	9	11	10
Cichlasoma dimerus	South American cichlids	Meijide & Guerrero, 2000	1	4	3	2	?	5	5	5	6	5	?	8	6	6	9	8	7	10	?	10
Oreochromis niloticus	Nile tilapia	Fujimura & Okada, 2007	1	3	2	3	?	4	5	6	5	6	6	7	8	8	8	9	10	10	12	11
Labeotropheus trewavasae	Scrapermouth mbuna	Balon, 1977	1	2	4	3	?	2	3	5	5	6	11	9	5	7	9	8	10	12	14	13
Haplochromis piceatus	Victoria cichlids	de Jong et al., 2009	1	2	2	4	?	4	3	4	5	5	4	6	6	6	7	7	8	9	6	9
Melanotaenia splendida	Eastern rainbow fish	Humphrey et al., 2003	1	2	4	3	3	?	6	5	6	7	?	11	8	7	9	5	12	11	10	13
Adinia xenica	Diamond killifish	Cunningham & Balon, 1985	1	2	3	?	3	2	3	5	?	6	8	9	4	7	8	9	13	12	11	10
Fundulus heteroclitus	Mummichog	Armstrong & Swope Child, 1965	1	2	4	?	3	3	5	5	7	6	5	?	8	7	9	10	12	12	12	11
Xiphophorus maculatus	Southern platyfish	Tavolga & Rugh	1	2	2	4	?	3	3	5	4	5	6	7	8	6	5	7	11	10	?	9
Austrofundulus myersi	Rivulines	Wourms, 1998	1	4	3	2	2	5	6	7	10	7	?	11	9	8	9	10	14	13	11	12
Oryzias latipes	Japanese ricefish	Iwamatsu & Hirata, 1984	1	3	4	?	2	5	4	6	9	7	13	12	8	8	10	11	17	15	14	16
Oryzias javanicus	Javanese ricefish	Iwamatsu, 2004	1	3	4	?	2	5	5	6	9	7	7	?	8	8	9	10	14	13	11	12

723 **FIGURE 1**

FIGURE 2

a

b

c

748   **FIGURE 3**

761   **FIGURE 4**

	es	op	fs	tb	kv	tbr	ot	le	tl	hb	olf	oto	mff	bc	pfb	ep	h	mo	sb
op	0%																		
fs	0%	33%																	
tb	0%	27%	41%																
kv	0%	27%	27%	7%															
tbr	0%	11%	21%	15%	35%														
ot	0%	0%	3%	14%	5%	18%													
le	0%	0%	0%	5%	0%	7%	20%												
tl	0%	0%	0%	0%	0%	13%	16%	24%											
hb	0%	0%	0%	0%	0%	4%	13%	3%	24%										
olf	0%	0%	0%	0%	0%	0%	9%	14%	39%	32%									
oto	0%	0%	0%	0%	0%	5%	9%	5%	0%	36%	41%								
mff	0%	0%	0%	0%	0%	4%	0%	12%	14%	23%	20%	37%							
bc	0%	0%	0%	0%	0%	0%	4%	0%	13%	4%	15%	40%	32%						
pfb	0%	0%	0%	0%	0%	0%	4%	0%	4%	0%	19%	27%	13%	8%					
ep	0%	0%	0%	0%	0%	4%	3%	3%	8%	3%	19%	29%	20%	19%	37%				
h	0%	0%	0%	0%	0%	4%	0%	3%	0%	0%	9%	9%	0%	7%	14%	17%			
mo	0%	0%	0%	0%	0%	0%	0%	0%	0%	3%	5%	5%	4%	4%	4%	10%	37%		
sb	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	6%	0%	0%	5%	9%	39%	48%	
cfr	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%	4%	4%	39%	39%	43%

775 **FIGURE 5**

788   **FIGURE 6**