Transgenerational cues about local mate competition affect offspring sex

ratios in the spider mite *Tetranychus urticae*

Authors: Alison B. Duncan[1*†], Cassandra Marinosci[1†], Céline Devaux[1], Sophie

Lefèvre[1], Sara Magalhães[2], Joanne Griffin[1], Adeline Valente[1], Ophélie Ronce[1],

Isabelle Olivieri[1].

1. Institut des Sciences de l'Évolution, Université de Montpellier, CNRS, IRD, EPHE,

CC065, Place Eugène Bataillon, 34095 Montpellier Cedex 05, France

2. cE3c: Centre for Ecology, Evolution, and Environmental Changes, Faculty of

Sciences, University of Lisbon, Edifício C2, 38 piso, 1749-016 Lisboa, Portugal

*Correspondence: alison.duncan@umontpellier.fr.

[†]Equal authorship contribution

KEYWORDS: local mate competition, sex-ratio, sib-mating, haplodiploid

<u>Abstract</u>

In structured populations, competition for mates between closely related males,

termed Local Mate Competition (LMC), is expected to select for female-biased

offspring sex ratios. However, the cues underlying sex allocation decisions remain

poorly studied. Here, we test for several cues in the spider mite *Tetranychus urticae*,

a species that was previously found to adjust the sex ratio of its offspring in response

to the number of females in an isolated population, i.e. a patch. We here investigate

whether the offspring sex ratio of *T. urticae* females changes in response to 1) the

current number of females in the same patch, 2) the number of females in the

patches of their mothers and 3) their relatedness to their mate. Single females on

patches produced similar sex ratios to those of groups of 15 females; their mothers

had been in identical conditions of panmixia. The offspring sex ratios of females

mated with their brother did not differ from those of females mated with an unrelated

male. Females however produced a more female-biased offspring sex ratio if their

mothers were alone on a patch compared to 15 other females. Thus, maternal

environment is used as a cue for the sex allocation of daughters. We discuss the

conditions under which the maternal environment may be a reliable predictor of LMC

experienced by grand-sons.

INTRODUCTION

In structured populations, dispersal of one sex can cause competition to be more intense between kin of the non-dispersing sex than between kin of the dispersing one. This asymmetry in competition can cause a bias in the offspring sex ratio to reduce competitive interactions between brothers or sisters. If females are the dispersing sex, related males can compete for mates, a phenomenon termed Local Mate Competition (LMC). Female-biased offspring sex ratios are expected to evolve to reduce this competition between males, both through a decrease in the number of male competitors and an increase in the number of mates per son (Hamilton, 1967; Maynard Smith, 1978; Taylor, 1981; Charnov, 1982; Frank, 1985; Herre, 1985). In haplodiploid species, in addition to LMC, a female-biased sex ratio can also arise because mothers that mate with a brother are more related to their daughters than they are to their sons (Greeff, 1996; Reece *et al.*, 2004).

Population structure can vary in time, and consequently so might the intensity of LMC and sib-mating. In such a situation, selection should favour females that facultatively (plastically) adjust their sex allocation in response to cues in the environment that reflect the intensity of LMC their sons will experience as well as whether they have mated with kin (Frank, 1985; Herre, 1985; Nagelkerke & Sabelis, 1996; Macke *et al.*, 2011a). For example, females could adjust their offspring sex ratios in response to the number of females laying eggs (foundresses) in the same patch or to the level of relatedness of their mate. Consistent with expectations that females respond to current cues in the environment, more female biased offspring sex ratios are observed in patches formed by a single foundress as compared to patches founded by multiple foundresses, across a wide range of species (Shuker *et al.*, 2005; West *et al.*, 2005; Sato & Saito, 2007; Wang *et al.*, 2015). Several studies,

however, found that females do not adjust their offspring sex ratio in response to

different numbers of foundresses on a patch (Innocent *et al.*, 2007; Abe *et al.*, 2014;

Lievens *et al.*, 2016). Moreover, there is currently no empirical support for the

prediction that sib-mating results in more female biased offspring sex ratios than

random mating (Reece *et al.*, 2004; Burton-Chellew *et al.*, 2008).

Alternatively, LMC and inbreeding may change in a predictable fashion over

generations. This scenario is predicted by haystack models of LMC in which sub-

divided populations may persist over multiple generations, and relatedness between

females sharing a patch may increase across generations (Nagelkerke & Sabelis,

1996). This increased relatedness can select for even more female biased offspring

sex ratios due to more intense LMC (Frank, 1985; Gardner *et al.*, 2009). If change in

the sub-divided population structure is predictable through time, transfer of

information from mothers to daughters could allow for a better adjustment of sex

allocation.

Discrepancies among studies investigating sex allocation may be caused by the

underlying cues being unknown and thus overlooked when designing experiments. In

principle, these cues could stem from both the current or past environment, that is

factors experienced while a female is laying her eggs, or those encountered as a

juvenile, or even in previous generations. For example, *Nasonia vitripennis* females

change their sex allocation in response to immediate cues in the environment,

including the current number of foundresses, female odours and the presence of

eggs in the environment (Shuker *et al.*, 2005, 2007). In contrast, it is the environment

that house wrens experience as juveniles, not the current environment, that affects

the sex ratio of their own offspring: they produce a more female-biased offspring sex

ratio when raised in a smaller brood (Bowers *et al.*, 2017). Also, female mice

exposed to stress in utero produce more daughters (Edwards et al., 2016).

Here we investigate the impact of immediate and longer-term cues on sex

allocation in the spider mite *Tetranychus urticae.* An earlier study showed that spider

mites evolving under constant, high LMC lost the ability to facultatively adjust their

offspring sex ratio, whilst those evolving under weaker LMC retained the ability for

facultative sex allocation in response to the number of foundresses (Macke *et al.*,

2011a). In that study, sex ratio was measured after mites had spent two generations

on patches containing the same number of foundresses. Thus, both mothers and

their daughters experienced cues indicating the same intensity of LMC for their sons.

Therefore, the set-up did not allow disentangling the current vs past cue(s) used by

females that retained the ability for facultative sex allocation. Other studies have

shown that maternal cues impact a number of traits in spider mites including sex

allocation, dispersal distance, fecundity and diapause (Oku *et al.*, 2003; Magalhães

et al., 2011; Bitume *et al.*, 2014; Marinosci *et al.*, 2015). *T. urticae* females also

respond to current cues and produce more female-biased offspring sex ratios when

there are fewer females sharing a patch (Wrensch & Young, 1978; Roeder, 1992)

and they are able to recognise kin (Tien *et al.*, 2011; Bitume *et al.*, 2014).

We aim at uncovering the interplay between immediate and past cues in the

environment in determining sex ratios in *T. urticae*. We investigate whether females

alter their sex allocation in response to the number of females in the same patch (i.e.

current indicators of LMC that their sons will experience) or the number of females in

the patch of their mothers (i.e. past LMC conditions). Because sib-mating is

negatively correlated with the number of foundresses on a patch, and a certainty in

our high LMC treatments, we evaluate the effect of sib-mating on sex allocation in a

separate experiment.

MATERIAL AND METHODS

Biological model

The two-spotted spider mite, *T. urticae* Koch (Acari: Tetranychidae)*,* is a generalist

herbivore (Migeon & Dorkeld, n.d.; Helle & Sabelis, 1985) with a short generation

time (less than 15 days at 20°C on a suitable host; (Helle & Sabelis, 1985; Riahi *et*

al., 2013)). It has an arrhenotokous haplodiploid life cycle in which females produce

haploid sons parthenogenetically and diploid daughters from fertilised eggs. After the

egg stage, all individuals pass through a larval and two nymphal (protonymph and

deutonymph) stages, with short quiescent stages between each, before reaching

maturity. Mature females are approximately twice as big as, and rounder than,

mature males. The sex ratios measured in this study are tertiary sex ratios defined as

the proportion of adult males on a patch; offspring sex ratio is a trait of the mother.

Study population

This study uses the same base population as Macke et al. 2011a. This population

was collected from a cucumber greenhouse in Pijnacker, The Netherlands, in May

1994. Mites were transferred to a climate chamber at the University of Amsterdam

where they were maintained on the same host plant. In September 2007, after ~240

months (~480 generations), approximately 5000 mites were sampled from the

Amsterdam population to create two populations on cucumber plants at the

University of Montpellier. Mite populations were maintained under the same

conditions in a single climate chamber at 25°C±2°C, with a photoperiod of 14L:10D.

A new, more genetically variable, base population was created in February 2011 by

mixing and transferring approximately 200 adult females from the two previous

populations. This new base population was maintained on 4 - 6 cucumber plants

contained in a single plastic box (505mm length x 310mm width x 380mm height)

with a hole in the lid covered in a fine mesh to avoid condensation. Plants were

watered every week and rotten plants removed and replaced with young, mite-free

plants; plants were placed beside infested ones to facilitate mite dispersal.

To start the experiment (Generation 0), 12 independent samples from the base

population consisting of 40 adult females each were created; this ensures that

females seeding the experiment were of the same age. Each sample was placed on

a whole cucumber leaf resting on water-saturated cotton wool in a plastic box

(255mm x 185mm x 77mm) to prevent both leaf dehydration and mite dispersal.

Females were allowed to lay eggs and, 14 days later, their mature adult daughters

were sampled to start the experiment '*sex allocation in response to LMC across*

generations'. In the *'sib mating'* experiment, slightly older daughters were used as the

experiment was started 20 and 21 days after females started laying eggs in

generation 0. In all experiments, mites were maintained in the same plastic boxes on

leaves or leaf fragments placed on saturated cotton wool in climate chambers at 25 ±

2ºC with a 16L:8D cycle. Prior to starting and during the experiments, we confirmed

that our *T. urticae* populations were *Wolbachia*-free (see Supplementary Materials,

Appendix 1 for details).

Sex allocation in response to LMC across generations

This experiment investigated the sex allocation of females over two generations. In

Generation 1 (the 'maternal generation'), we measured the offspring sex ratios in

response to high (1 foundress) versus low (15 foundresses) LMC experienced by

laying females (these are the daughters of females from Generation 0). Each patch

from Generation 0 contributed equal numbers of females, arbitrarily sampled, to the

high and low LMC treatments. In Generation 2 (the 'daughter generation'), we

investigated the sex allocation under high LMC of daughters from females of

Generation 1, which had experienced either high or low LMC. Thus, in Generation 2

we tested whether LMC experienced by mothers influenced the offspring sex ratios of

their daughters (see Figure S1 for an outline of the experimental design).

Generation 1 (maternal generation):

In the low LMC treatment, a total of 360 females, in groups of 15 females, were

placed on 60-cm^2 cucumber leaf patches. We had 6 boxes, each containing 4 low

LMC patches. In the high LMC treatment, 240 females were individually placed on a

4-cm^2 leaf patch, and spread across 5 boxes, each containing 48 patches. Female

density was the same for both low and high LMC conditions (0.25 females $\times$ cm^{-2}).

Each box was divided into 4 quarters, each quarter containing 12 patches of single

females in high LMC boxes and 1 patch of 15 females in low LMC boxes. This was

done to get the same numbers of replicates (quarters) for each treatment with similar

numbers of females laying eggs (12 for low LMC and 15 for high LMC). Note that we

did not control for female mortality, i.e. there may have been fewer females

contributing to the next generation in each treatment. In both LMC treatments,

females laid eggs, and 14 days later we counted the number of emerging adults on

each patch, and determined their sex, using a stereo light microscope. Only

measurements from a subset of 3 high LMC boxes and 2 low LMC boxes were

included in the statistical analyses due to experimenter error.

Generation 2 (daughter generation):

Independently of LMC levels, 12 females were randomly collected from each quarter

of each box from Generation 1 and individually placed under high LMC on a 4 cm^2

leaf patch (same density as in Generation 1). Due to differential fecundity, some

females from Generation 1 may have contributed more offspring to Generation 2 than

did others. All patches were randomly spread across 6 boxes (48 patches per box).

Note that some females in Generation 2 were sampled from patches that were not

used in the analysis of sex ratio in Generation 1 due to experimenter error. Females

were allowed to lay eggs and their offspring sex ratio was measured 14 days after

transfer. In total, we measured the offspring sex ratio and number of emerging adults

on 114 patches from 20 quarters in boxes testing the effect of low maternal LMC (5 -

6 females per quarter), and 116 patches, from 14 quarters, in boxes testing the effect

of high maternal LMC (3 - 12 females per quarter). Some patches were lost due to

female mortality before they laid eggs or to leaf patches rotting.

Sex allocation adjustment in response to sib mating

This experiment investigated whether females mated with their brothers produced a

more female biased offspring sex ratio than did females mated with unrelated males.

In total, 96 females sampled from Generation 0 were individually placed on 4 cm^2

patches spread across 2 boxes (48 patches per box). This experiment was done in

temporal blocks, over two days. Females were allowed to lay eggs and 9 days later

quiescent daughters (2 from each patch, later allocated to each of the two

treatments) were individually placed on 4 cm^2 patches to ensure all mites assigned to

our mating treatments were virgin. The same was done for emerged or quiescent

sons. Mating pairs were formed three days later, when all quiescent females were

adult. Females were assigned to one of two mating treatments: (1) the sib-mating

treatment, in which one haphazardly sampled brother was added to their patch or (2)

the unrelated mating treatment, in which a male from another patch was added to

their patch. Males were removed after three days, females were allowed to lay eggs

and 14 days later the sex ratio and number of emerging adults was measured. In

total, we formed 24 patches with unrelated mates and 28 patches with sib-mating.

Statistical analyses

For all analyses, except sex ratio of females from generation 1 (see below), we

analysed variation in the number of sons (offspring sex ratio) and the total number of

offspring that emerged as adults for females using generalised linear mixed models,

assuming a binomial distribution and a Poisson distribution, respectively. These

models were performed in SAS using the GLIMMIX model procedure with the

lsmeans statement to account for unbalanced designs (SAS script provided in

Appendix 2, Supplementary Materials).

Sex allocation in response to LMC across two generations

At generation 1, the sex ratio was measured for single females for the high LMC

treatment and for groups of 15 females for the low LMC treatment. To analyse

variation in the number of sons across groups of females of similar sizes, we used a

two-tailed permutation test, randomly sampling groups of 12 and 15 females

(corresponding to the experimental design) under the null hypothesis of no effect of

LMC. We generated 10,000 empirical values of the z statistics and could thus obtain

the p-value for the effect of LMC on sex ratio. The observed statistic was generated

using the number of sons produced in quarters of boxes by groups of 12 females for

the high LMC treatment and by groups of 15 females for the low LMC treatment.

At Generation 1, we analysed variation in mean individual offspring production

in each quarter as the number of offspring reaching adulthood in 14 days (this will

correspond to ~4 days of laying eggs) accounting for LMC treatment as a fixed effect,

and the box containing each quarter as a random effect nested within LMC

treatment. At Generation 2, we tested the effect of the LMC conditions experienced

by the mother at generation 1 on male production and offspring production. We

accounted for the LMC treatment of the mothers as a fixed effect, and quarter from

which each female originated as a random factor nested within LMC treatment.

Sex allocation adjustment in response to sib mating

In the sib-mating experiment, we accounted for the mating treatments of the mothers

as a fixed effect and the patch of each mother, and temporal block as random

factors. Note, block could not be included in the analysis for offspring production

because the model did not converge.

RESULTS

In all experiments, offspring production was not affected by the intensity of LMC in

current (mean offspring per female $\pm$ 1 SE, 19.01 $\pm$ 1.630 offspring for females in

high LMC and 20.16 $\pm$ 0.920 for females in low LMC; $F_{1,3} = 0.13$, p-value = 0.7397) or

past generations (13.05 $\pm$ 0.860 for high LMC and 14.79 $\pm$ 0.744 for low LMC; $F_{1,32} =$

2.37, p-value = 0.1335), or by the relatedness between mates (14.54 $\pm$ 1.088

offspring for females mated with brothers and 16.42 $\pm$ 0.934 for females mated with

unrelated males; $F_{1,21} = 2.95$, p-value = 0.1005). The absence of an effect on

offspring production ensures that the estimates of sex-ratio have the same precision

between LMC and sib-mating treatments.

Sex allocation in response to LMC across two generations

At Generation 1, females did not modify their offspring sex ratio according to the

number of other females in the same patch (p-value = 0.938; Figure 1A). In contrast,

at Generation 2, the maternal environment affected the sex allocation of their

daughters, as females produced a less female biased offspring sex ratio if their

mothers had experienced low, compared to high, LMC ($F_{1,32}$ = 10.98, p-value =

0.002, Figure 1B).

Sex allocation adjustment in response to sib-mating

Mating with a brother or an unrelated male did not change sex allocation ($F_{1,21}$ =

0.06, p-value = 0.82; Fig .1C). This means the effect of high LMC over two

generations on sex allocation are not due to sib-mating.

Figure 1. Proportion of male offspring for (A) females experiencing high (1 female) or

low (15 females) LMC, (B) females whose mothers experienced high or low LMC and

(C) females mated with their brother or an unrelated male. Figures show empirical

means ± 2 standard errors (confidence interval).

DISCUSSION

We found that sex allocation in the spider mite *T. urticae* does not depend on the

number of females on the same patch or on the relatedness between mates. Instead,

we found that females alter their offspring sex ratio according to the number of

females on their mother's patch, showing that the maternal environment may be

important for sex allocation. Additionally, none of the treatments -current or maternal

levels of LMC, or mating with a sibling- impacted the number of offspring that were

laid over a ~ 4day period and emerged as adults.

292   Female *T. urticae* did not produce a more female-biased sex ratio when they

mated with their brother compared to an unrelated male. As spider mites do

recognise kin (Tien *et al.*, 2011; Bitume *et al.*, 2013), this calls for an explanation.

One possibility may be that there is sexual conflict over the optimal offspring sex ratio

(Shuker *et al.*, 2009; Macke *et al.*, 2014). In haplodiploids, males only pass their

genes to the next generation via daughters, thus it is always beneficial to produce a

more female-biased offspring sex ratio for males, whereas for females the optimal

offspring sex ratio depends on levels of LMC (Shuker *et al.*, 2009; Macke *et al.*, 2014;

Shuker & Cook, 2014). Since *T. urticae* males can manipulate females to produce

more female-biased offspring sex ratios (Macke *et al.*, 2014) females may be

reluctant to alter their sex allocation in response to any cue provided by their mate.

Another possible explanation is that females do not alter their sex allocation in

response to sib mating due to selection to reduce inbreeding (Greeff, 1996), from

which they suffer (Tien *et al.*, 2014; Yoshioka & Yano, 2014). Furthermore, *T. urticae*

actively avoids mating with kin as females prefer to mate with unrelated males over

brothers (Tien *et al.*, 2011) and males prefer to mate with unfamiliar females

(Yoshioka & Yano, 2014).

As found in a number of other species (Reece *et al.*, 2005; Innocent *et al.*,

2007; Abe *et al.*, 2014) we did not find that female *T. urticae* alter their sex allocation

in response to the number of other females in the same patch (but see Wrensch &

Young, 1978; Roeder, 1992). For example, in *Nasonia vitripennis* other cues,

specifically egg presence and the relative number of eggs a female contributes to a

patch, had a stronger influence on sex allocation than the number of foundresses

(Shuker & West, 2004; Burton-Chellew *et al.*, 2008). It could be, for our population

under our experimental conditions, that the number of females in the current patch

does not provide reliable information about the LMC level that the offspring will

experience. This may be because (a) females often move away from the patch or (b)

females have high mortality during the egg laying period. In spider mites, the second

hypothesis is possible if females are on low-quality plants (Magalhaes *et al.*, 2007;

Marinosci *et al.*, 2015).

It may be that females respond to density and not the absolute number of

females. In other studies that found an effect of foundress number on LMC, density

was not controlled, which we did in our experiment (Wrensch & Young, 1978;

Roeder, 1992). Alternatively, the absolute number of females on a patch could have

been too low to elicit a response. Spider mites use mechanical and chemosensory

processes via web and faeces to detect conspecifics (Royalty *et al.*, 1992; Clotuche

et al., 2014). *T. urticae* can also use volatiles released by infected plants to detect

conspecifics, but at densities much higher than in our experiment, and on whole

plants, and not leaves as here (Pallini *et al.*, 1997). These cues may be not available

in our experiment with low densities of females.

In Generation 2 densities were also identical to those of Generation 1 and

between treatments, and here we did detect a difference in sex allocation. Adult

females whose mothers experienced low LMC produced less female biased offspring sex ratios. In both Macke et al, 2011 and our experiment sex allocation is more female biased when females experienced at least two generations of LMC. In both experiments sex allocation could be in response to the number of foundresses in the mother's patch, or the number, or relatedness, of individuals in the juvenile environment.

There could be a true maternal effect in sex allocation, meaning that the mother's experience of her environment influences her offspring's phenotype (Mousseau & Fox, 1998). Maternal effects have been shown to have a role determining many offspring traits including offspring size and juvenile survival (Pick *et al.*, 2016), resistance to parasites (Little *et al.*, 2003) and competitive ability (Bentz *et al.*, 2016). In *T. urticae* maternal effects have been shown to partly determine offspring phenotypes for dispersal distance (Bitume *et al.*, 2014), diapause incidence (Oku *et al.*, 2003) and offspring life-history traits (Magalhães *et al.*, 2011; Marinosci *et al.*, 2015). To the best of our knowledge, maternal effects have not been previously shown to play a role in determining offspring sex allocation. An exception may be studies in which stress during the embryonic period influences the offspring sex ratio; one in mice, which found that females produce a more female bias offspring sex ratio if they experience stress *in utero* (Edwards *et al.*, 2016), and another showing that paternal incubation temperature influences offspring sex ratio in jacky dragons (Warner *et al.*, 2013). Whether these can be considered as maternal (or paternal) effects is debatable, as the developing individual is itself experiencing the stress. If our results are due to a maternal effect the same one might be responsible for the results observed in the selection experiment by Macke *et al.* (2011a). Indeed, in that

experiment, mothers and daughters experienced the same predictable LMC level, so

sex ratio adjustment may be explained by the maternal environment.

If female sex allocation in *T. urticae* is in part influenced by maternal cues for

LMC this could come about via the manipulation of a daughter's egg production. Egg

size is thought to be one way by which female *T. urticae* can plastically adjust their

sex allocation, as fertilisation probability increases with egg size, bigger eggs being

more likely to develop into females (Macke *et al.*, 2011b). Furthermore, *T. urticae*

evolving under high LMC produce larger eggs and a more female biased sex ratio

(Macke *et al.*, 2012). Female *T. urticae* possess a single ovary within which eggs

develop and are fertilised (Feiertag-Koppen & Pijnacker, 1982). Female larvae that

have recently emerged from their eggs possess cells in their ovaries undergoing

meiosis (Feiertag-Koppen & Pijnacker, 1982; Mothes-Wagner, 1984); the conditions

or resources of the mothers could influence the development and size of their

daughters' eggs, which could translate into a given sex ratio.

Another possibility is that the juvenile environment acts as a possible proximal

cause of sex allocation, which itself depends on maternal LMC. When mothers

experience high LMC, and are alone on a patch, females will develop with their

siblings whereas under low LMC, juveniles will share their patch with both related

and unrelated individuals. Furthermore, although we controlled for density across

treatments, under low LMC there is a higher absolute number of juveniles than there

is under high LMC. Offspring sex ratio has been linked to the environment a mother

experiences during the juvenile period in wrens and voles; when mothers

experienced juvenile resource limitation they produced less male biased offspring

sex ratios (Helle *et al.*, 2012; Bowers *et al.*, 2017). Nevertheless, if females respond

to cues experienced whilst juvenile and before they disperse to find their own patch,

the result remains that factors due to maternal LMC influence a daughter's sex
allocation.

Why then, should individuals use maternal cues, or cues available from the
juvenile environment, over those present in the immediate environment to alter the
sex ratio of their offspring? It might be that our observed response is maladaptive.
Indeed, maternal effects do not necessarily increase offspring fitness (Sheriff & Love,
2013; Barks & Laird, 2015). However, it might adaptive if levels of LMC change in a
predictable fashion across generations (for example, increasing or decreasing LMC)
for a female to integrate information provided by mothers (Mousseau & Fox, 1998). It
has been hypothesised that *T. urticae* life-history follows predictable population
groupings across multiple generations on the same host plant (Nagelkerke & Sabelis,
1996). If this is the case, maternal cues may provide reliable information about the
current state of a population, especially if relevant cues are not available in the patch.
Mitchell (1972) shows that patches in experimental populations can be relatively
small (1 cm^2) compared to the size of a leaf, and the patch sizes we used. Patch size
and population structure remain to be measured in natural populations.

We can conclude from our experiment and others that multiple cues act
together to influence sex allocation decisions, with the contribution of each
depending on context. Maternal cues (maternal LMC or juvenile environment) might
provide a more integrative measure of population structure, which females use for
their sex allocation decisions at low densities. Clearly, more studies are needed to
assess the relative value of cues informing on population structure and on the
precise scale at which competition for mates actually occurs.

ACKNOWLEDGEMENTS

We would like to thank David Carbonell for growing and maintaining plants for this

experiment. This project was funded by a joint grant from the Agence Nationale de la

Recherche and the Fundação para a Ciência e a Tecnologia to IO and SM (FCT-

ANR/BIA-EVF/0013/2012).

REFERENCES

Abe, J., Kamimura, Y. & West, S.A. 2014. Inexplicably female-biased sex ratios in Melittobia wasps. *Evolution* **68–9**: 2709–2717.

Barks, P.M. & Laird, R.A. 2015. Senescence in duckweed: Age-related declines in survival, reproduction and offspring quality. *Funct. Ecol.* **29**: 540–548.

Bentz, A.B., Becker, D.J. & Navara, K.J. 2016. Evolutionary implications of interspecific variation in a maternal effect: a meta-analysis of yolk testosterone response to competition. *R. Soc. Open Sci.* **3**: 160499.

Bitume, E. V., Bonte, D., Ronce, O., Olivieri, I. & Nieberding, C.M. 2014. Dispersal distance is influenced by parental and grand-parental density. *Proc. R. Soc. B* **281**: 20141061–20141061.

Bitume, E. V, Bonte, D., Ronce, O., Bach, F., Flaven, E., Olivieri, I., *et al.* 2013. Density and genetic relatedness increase dispersal distance in a subsocial organism. *Ecol. Lett.* **16**: 430–7.

Bowers, E.K., Thompson, C.F. & Sakaluk, S.K. 2017. Maternal natal environment and breeding territory predict the condition and sex ratio of offspring. *Evol. Biol.* **44**: 11–20.

Burton-Chellew, M.N., Koevoets, T., Grillenberger, B.K., Sykes, E.M., Underwood, S.L., Bijlsma, K., *et al.* 2008. Facultative sex ratio adjustment in natural populations of wasps: cues of local mate competition and the precision of adaptation. *Am. Nat.* **172**.

Charnov, E.L. 1982. *The theory of sex allocation.* Princeton University Press.

Clotuche, G., Yano, S., Akino, T. & Amano, H. 2014. Chemical investigation of aggregation behaviour in the two-spotted spider mite Tetranychus urticae. *Exp. Appl. Acarol.* **63**: 377–387.

Edwards, A.M., Cameron, E.Z., Pereira, J.C., Wapstra, E., Horton, S.R. & Thomasson, K. 2016. Gestational experience alters sex allocation in the subsequent generation. *R. Soc. Open Sci.* **3**: 160210.

Feiertag-Koppen, C.C.M. & Pijnacker, L.P. 1982. Development of the female germ cells and process of internal fertilization in the two-spotted spider mite Tetranychus urticae Koch (Acariformes: Tetranychidae). *Int. J. Insect Morphol. Embryol.* **11**: 271–284.

Frank, S.A. 1985. Hierarchical selection theory and sex ratios. II. On applying the theory, and a test with fig wasps. *Evolution* **39**: 949–964.

Gardner, A., Arce, A. & Alpedrinha, J. 2009. Budding dispersal and the sex ratio. *J. Evol. Biol.* **22**: 1036–1045.

Greeff, J.M. 1996. Alternative mating strategies, partial sex ratios in haplodiploid species sibmating and split. *J. Evol. Biol.* **9**: 855–869.

Hamilton, W.D. 1967. Extraordinary sex ratios. *Science* **156**: 477–488.

Helle, H., Koskela, E. & Mappes, T. 2012. Life in varying environments: Experimental evidence for delayed effects of juvenile environment on adult life history. *J. Anim. Ecol.* **81**: 573–582.

Helle, M. & Sabelis, M.W. 1985. *Spider mites: Their biology, natural enemies and control*. Elsevier.

Herre, E.A. 1985. Sex ratio adjustment in fig wasps. *Science* **228**: 896–898.

Innocent, T.M., Savage, J., West, S.A. & Reece, S.E. 2007. Lethal combat and sex ratio evolution in a parasitoid wasp. *Behav. Ecol.* **18**: 709–715.

Lievens, E.J.P., Henriques, G.J.B., Michalakis, Y. & Lenormand, T. 2016. Maladaptive sex ratio adjustment in the invasive brine shrimp Artemia

462 franciscana. *Curr. Biol.* **26**: 1–5.

463 Little, T.J., O'Connor, B., Colegrave, N., Watt, K. & Read, A.F. 2003. Maternal
464 transfer of strain-specific immunity in an invertebrate. *Curr. Biol.* **13**: 489–492.

465 Macke, E., Magalhaes, S., Bach, F. & Olivieri, I. 2011a. Experimental evolution of
466 reduced sex ratio adjustment under local mate competition. *Science* **334**: 1127–
467 1129.

468 Macke, E., Magalhães, S., Bach, F. & Olivieri, I. 2012. Sex-ratio adjustment in
469 response to local mate competition is achieved through an alteration of egg size
470 in a haplodiploid spider mite. *Proc. R. Soc. B* **279**: 4634–42.

471 Macke, E., Magalhães, S., Khan, H.D.-T., Luciano, A., Frantz, A., Facon, B., *et al.*
472 2011b. Sex allocation in haplodiploids is mediated by egg size: evidence in the
473 spider mite Tetranychus urticae Koch. *Proc. R. Soc. B* **278**: 1054–63.

474 Macke, E., Olivieri, I. & Magalhaes, S. 2014. Local mate competition mediates sexual
475 conflict over sex ratio in a haplodiploid spider mite. *Curr. Biol.* **24**: 2850–2854.

476 Magalhães, S., Blanchet, E., Egas, M. & Olivieri, I. 2011. Environmental effects on
477 the detection of adaptation. *J. Evol. Biol.* **24**: 2653–62.

478 Magalhaes, S., Fayard, J., Janssen, A., Carbonell, D., Olivieri, I. & Magalhães, S.
479 2007. Adaptation in a spider mite population after long-term evolution on a single
480 host plant. *J. Evol. Biol.* **20**: 2016–2027.

481 Marinosci, C., Magalhaes, S., Macke, E., Navajas, M., Carbonell, D., Devaux, C., *et*
482 *al.* 2015. Effects of host plant on life-history traits in the polyphagous spider mite
483 Tetranychus urticae. *Ecol. Evol.* **5**: 3151–3158.

484 Maynard Smith, J. 1978. *The evolution of sex*. Cambridge University Press,
485 Cambridge.

486 Migeon, A. & Dorkeld, F. Spider mites web: a comprehensive database for the
487 Tetranychidae. http://www1.montpellier.inra.fr/CBGP/spmweb/

488 Mothes-Wagner, U. 1984. Post-embryonic development of the two-spotted spider
489 mite Tetranychus urticae with special reference to the differentiation of the
490 female reproductive system. *Int. J. Invertebr. Reprod. Dev.* **7**: 203–216.

491 Mousseau, T.A. & Fox, C.W. 1998. The adaptive significance of maternal effects.
492 *Trends Ecol. Evol.* **13**: 403–407.

493 Nagelkerke, C.J. & Sabelis, M.W. 1996. Hierarchical levels of spatial structure and
494 their consequences for the evolution of sex allocation in mites and other
495 arthropods. *Am. Nat.* **148**: 16–39.

496 Oku, K., Yano, S. & Takafuji, A. 2003. Different maternal effects on diapause
497 induction of tetranychid mites, Tertranychus urticae and T. kanzawai (Acari:
498 Tetranychidae). *Appl. Entomol. Zool.* **38**: 267–270.

499 Pallini, A., Janssen, A. & Sabelis, M.W. 1997. Odour-mediated responses of
500 phytophagous mites to conspecific and heterospecific competitors. *Oecologia*
501 **110**: 179–185.

502 Pick, J.L., Ebneter, C., Hutter, P. & Tschirren, B. 2016. Disentangling genetic and
503 prenatal maternal effects on offspring size and survival. *Am. Nat.* **188**: 628–639.

504 Reece, S.E., Shuker, D.M., Pen, I., Duncan, A.B., Choudhary, A., Batchelor, C.M., *et*
505 *al.* 2004. Kin discrimination and sex ratios in a parasitoid wasp. *J. Evol. Biol.* **17**:
506 208–216.

507 Reece, S.E., Wherry, R.N. & Bloor, J.M.G. 2005. Sex allocation and interactions
508 between relatives in the bean beetle, Callosobruchus maculatus. *Behav.*
509 *Processes* **70**: 282–288.

510 Riahi, E., Shishehbor, P., Nemati, A.R. & Saeidi, Z. 2013. Temperature effects on
511 development and life table parameters of Tetranychus urticae (Acari:

Tetranychidae). *J. Agric. Sci. Technol.* **15**: 661–672.

Roeder, C. 1992. Sex ratio response of the two-spotted spider mite. *Can. J. Zool.* **70**: 1965–1967.

Royalty, R.N., Phelan, P.L. & Hall, F.R. 1992. Arrestment of male twospotted spider mite. *J. Chem. Ecol.* **18**: 137–153.

Sato, Y. & Saito, Y. 2007. Can the extremely female-biased sex ratio of the social spider mites be explained by Hamilton's local mate competition model? *Ecol. Entomol.* **32**: 597–602.

Sheriff, M.J. & Love, O.P. 2013. Determining the adaptive potential of maternal stress. *Ecol. Lett.* **16**: 271–280.

Shuker, D.M. & Cook, N. 2014. Evolution: conflict by the sexes, for the sexes. *Curr. Biol.* **24**: R1135–R1137.

Shuker, D.M., Moynihan, A.M. & Ross, L. 2009. Sexual conflict, sex allocation and the genetic system. *Biol. Lett.* **5**: 682–685.

Shuker, D.M., Pen, I., Duncan, A.B., Reece, S.E. & West, S.A. 2005. Sex ratios under asymmetrical local mate competition: Theory and a test with parasitoid wasps. *Am. Nat.* **166**: 301–316.

Shuker, D.M., Reece, S.E., Lee, a, Graham, a, Duncan, A.B. & West, S.A. 2007. Information use in space and time: sex allocation behaviour in the parasitoid wasp Nasonia vitripennis. *Anim. Behav.* **73**: 971–977.

Shuker, D.M. & West, S.A. 2004. Information constraints and the precision of adaptation: sex ratio manipulation in wasps. *Proc. Natl. Acad. Sci. U. S. A.* **101**: 10363–7.

Taylor, P. 1981. Intra-sex and inter-sex sibling interactions. *Nature* **291**: 64–66.

Tien, N., Massourakis, G., Sabelis, M. & Egas, M. 2011. Mate choice promotes inbreeding avoidance in the two-spotted spider mite. *Exp. Appl. Acarol.* **54**: 119–124.

Tien, N.S.H., Sabelis, M.W. & Egas, M. 2014. Inbreeding depression and purging in a haplodiploid: gender-related effects. *Heredity* **114**: 327–332.

Wang, R., Sun, B., He, J. & Dunn, D.W. 2015. Non-quantitative adjustment of offspring sex ratios in pollinating fig wasps. *Sci. Rep.* **5**: 13057.

Warner, D.A., Uller, T. & Shine, R. 2013. Transgenerational sex determination: The embryonic environment experienced by a male affects offspring sex ratio. *Sci. Rep.* **3**: 1–4.

West, S.A., Shuker, D.M. & Sheldon, B.C. 2005. Sex-ratio adjustment when relatives interact: a test of constraints on adaptation. *Evolution* **59**: 1211–28.

Wrensch, D.L. & Young, S.S.Y. 1978. Effects of density and host quality on rate of development, survivorship and sex ratio in the Carmine spider mite. *Environ. Entomol.* **7**: 499–501.

Yoshioka, T. & Yano, S. 2014. Do Tetranychus urticae males avoid mating with familiar females? *J. Exp. Biol.* **217**: 2297–2300.