Title: Scaling of G1 duration with population doubling time by a cyclin in *Saccharomyces*

cerevisiae

Authors: Heidi M. Blank, Michelle Callahan, Ioannis P.E. Pistikopoulos, Aggeliki O. Polymenis[1] and

Michael Polymenis

Affiliations: Department of Biochemistry and Biophysics, Texas A&M University, College Station, TX

77843, U.S.A.

[1]Present address: Department of Physics and Astronomy, University of California, Los

Angeles, CA 90095

Correspondence: blank@tamu.edu (H.M.B); polymenis@tamu.edu (M.P.)

Running Title: Scaling G1 with doubling time

ABSTRACT

A quantitative relation that defines the population doubling time (T_d) as a function of the time

eukaryotic cells spend in specific cell cycle phases would be instrumental for estimating rates of cell

proliferation and for evaluating introduced perturbations. Here, we show that in human cells the length

of the G1 phase (T_{G1}) regressed on T_d with a slope of ≈0.75, while in the yeast *Saccharomyces cerevisiae*

the slope was smaller, at ≈0.60. Levels of the yeast G1 cyclin Cln3p were positively associated with the

growth rate over a broad range through translational control mediated by a short uORF in the *CLN3*

transcript. Cln3p was also necessary for the scaling between T_{G1} and T_d. In contrast, yeast lacking the

Whi5p transcriptional repressor maintained the scaling between T_{G1} and T_d. These data reveal

fundamental scaling relations between the duration of eukaryotic cell cycle phases and rates of cell

proliferation, point to the necessary role of Cln3p in these relations in yeast and provide a mechanistic

basis linking Cln3p levels to growth rate and the scaling of G1 with doubling time.

Keywords: cell size / growth law / START / chemostat

INTRODUCTION

Recurring shapes and patterns in nature are sometimes described with mathematical relationships. As a

result, these natural processes can be predicted and understood better. Regarding patterns of

eukaryotic cell division, one could ask: How are the lengths of eukaryotic cell cycle phases related to

each other and to the total doubling time of the population? Can such relations be described

mathematically, in the form of a scaling formula? If so, what are the molecular mechanisms that govern

the scaling? A scaling relation that describes eukaryotic cell division would be an important advance. For

example, it could serve as a point of reference against which the effects of genetic or other

perturbations can be evaluated.

The 'textbook' view in the coordination of growth and division in the eukaryotic cell cycle (e.g., see

Fig. 10-26 in (MORGAN 2007)) is that expansion of the G1 phase of the eukaryotic cell cycle accounts for

most of the lengthening of the cell cycle in slower proliferating cells, in budding yeast (JOHNSTON *et al.*

1977; BRAUER *et al.* 2008) or humans (BASERGA 1985; FISHER 2016). However, there is no report in the

literature of a quantitative relation that defines the doubling time (T_d) as a function of the time yeast or

human cells spend in the G1 phase (T_{G1}). Here, we collected data for yeast and human cell populations

to derive, for the first time in the field, scaling relations between T_{G1} and T_d. These scaling relations also

allowed us to critically evaluate the role of cell cycle regulators in yeast cells growing at different rates.

Two key regulators of the length of the G1 phase in yeast are the Cln3p and Whi5p proteins. The G1

cyclin Cln3p promotes initiation of DNA replication (CROSS 1988; NASH *et al.* 1988). In contrast, the

transcriptional repressor Whi5p acts analogously to the retinoblastoma gene product in animals, to

inhibit the G1/S transition (COSTANZO *et al.* 2004; DE BRUIN *et al.* 2004; PALUMBO *et al.* 2016). It has been

reported that while synthesis of Cln3p parallels cell size, the synthesis of Whi5p is independent of cell

size (SCHMOLLER *et al.* 2015), arguing that dilution of Whi5p as cells get bigger in G1 governs the length of

the G1 phase (SCHMOLLER AND SKOTHEIM 2015; SCHMOLLER *et al.* 2015). We obtained the first

measurements of Cln3p and Whi5p levels as a function of growth rate in steady-state cultures. The

levels of Cln3p varied over a broad range as a function of growth rate, due to a uORF affecting

translation of *CLN3*. Our data also show that Whi5p is fully dispensable for the normal scaling relation

between T_d and T_{G1}, and provide instead the functional and molecular basis for the necessary role of

Cln3p in this process.

MATERIALS AND METHODS

Strains. Unless stated otherwise, *S. cerevisiae* wild-type, *cln3Δ* and *whi5Δ strains* were in the BY4741

background (NCBI Taxon 559292; *MATa, his3Δ1, leu2Δ0, ura3Δ0, met15Δ0*) and they have been

described previously (SOMA *et al.* 2014). For protein surveillance, we constructed an otherwise wild type

strain that carried epitope-tagged *WHI5* and *CLN3* alleles at their endogenous chromosome locations.

First, a commercially available *WHI5-TAP::HIS3* strain (BY4741 otherwise; GE Healthcare) was

backcrossed three times into the W303 background (NCBI Taxon 580240; *MATα leu2-3,112 trp1-1 can1-*

100 ura3-1 ade2-1 his3-11,15). Then, it was crossed with an otherwise wild-type strain carrying a *CLN3-*

13MYC allele (W303 background) described elsewhere (THORBURN *et al.* 2013), and kindly provided by

Dr. A. Amon (MIT and HHMI). The resulting diploid was sporulated and dissected, to obtain *MATa*

haploid segregants carrying both the epitope-tagged *WHI5* and *CLN3* alleles (strains HB94/97; *MATa*

CLN3-13MYC::TRP⁺ WHI5-TAP::HIS⁺ leu2 ura3 met15), which were used in the experiments shown in

Figure 4. We verified expression of Whi5p-TAP and Cln3p-(Myc)₁₃ in this strain (see Source Data,

Supplementary File 1), and their absence in *whi5Δ* or *cln3Δ* strains, respectively. We also generated a

derivative of this strain, which lacks the uORF in the 5'-leader of the *CLN3* mRNA. To this end, we used

plasmid A-315T-pMT10 we had described previously (POLYMENIS AND SCHMIDT 1997), as a template in a

PCR reaction with forward (5'-CAAGAACTACCATTCGACAGG-3') and reverse primers (5'-

CGTACAGAAAGCGTATCAAA-3') to generate a product that carries in the 5'-leader of *CLN3* the *URA3-*

marked A-315T mutation that inactivates the uORF. We then used this PCR product to transform strain

HB94 (*WHI5-TAP, CLN3-13MYC*). Genomic DNA of transformants was sequenced to verify the presence

of the A-315T mutation. Confirmed A-315T mutants were then backcrossed with wild type (W303) to

segregate away possible secondary mutations at other loci. The resulting heterozygote was sporulated

and dissected to isolate a *WHI5-TAP, A-315T-CLN3-13MYC* segregant (HB104), which was used in the

experiments shown in Figure 4.

Measurements of G1 length. The values we show in Tables S1 and S2 were obtained from studies

reporting on the doubling time (T_d) of asynchronous cell populations and the fraction of cells that are in

the G1 phase, reflecting the *relative* duration of the G1 phase. To estimate the *absolute* length of the G1

phase, T_{G1}, we multiplied the relative G1 length by T_d (Tables S1, S2). This simple equation is the

appropriate one to use with chemostat data when as many cells are removed as are being produced in

the culture (HOFFMAN 1949). Subtracting T_{G1} from T_d yields the duration of the rest of the cell cycle

phases (T_{nonG1}). For non-chemostat data, we recognize that other more elaborate equations could be

used, especially for the asymmetric patterns of division of budding yeast (HARTWELL AND UNGER 1977;

JOHNSTON *et al.* 1980). However, for simplicity and ease of comparison across systems, we uniformly

applied the simple equation mentioned above. Furthermore, inaccuracies in the absolute values of T_{G1}

may only affect the intercept of the linear relation between T_d and T_{G1}, but not the slope that describes

the fundamental scaling between T_d and T_{G1}, or any of our conclusions. Indeed, when we plotted in

exactly the same manner as in Figure 1 only the chemostat values in Table S1 (94 data points), the slope

of LnT_d vs. LnT_{G1} was 0.6107. For the remaining, non-chemostat values (55 data points), the slope of LnT_d

vs. LnT_{G1} was 0.6074.

Overall, the obtained variables represent population averages. They do not resolve

intergenerational differences in cell cycle progression of the same cells in successive cell cycles. It is also

assumed that cell death contributes negligibly to the doubling time of the population. This assumption is

reasonable in microbial culture systems because young cells vastly outnumber older ones approaching

senescence. However, it may be of concern in mammalian culture systems. Hence, our data with human

cells should be interpreted with caution, for at least two reasons: First, the fraction of growing cells in

the culture may be significantly lower than one. Second, although we are looking at trends that

seemingly hold across a multitude of different cells, the data were overwhelmingly derived from cancer

cell lines, which in many cases have altered cell cycles. In the context of this study, however, population

averages hold significant advantages: First, they are easily obtained; Second, they are ubiquitously used

and reported in the literature; Third, they allow straightforward comparisons between different

systems, for example between yeast and human cells (see Figure 1).

For yeast, the data we collected (Table S1) were from wild type strains from various backgrounds,

except in the few cases where they carried temperature-sensitive alleles, such as *cdc* mutations

(JAGADISH AND CARTER 1977), to estimate the length of the G1 phase upon transfer to the non-permissive

temperature. The methods used to calculate the fraction of G1 cells included: measurements of the DNA

content of the cells by flow cytometry (SLATER *et al.* 1977; JOHNSTON *et al.* 1980; GUO *et al.* 2004; BRAUER

et al. 2008; HENRY *et al.* 2010); budding (TYSON *et al.* 1979; RIVIN AND FANGMAN 1980); sensitivity to cell

cycle arrest before DNA replication by pheromone (HARTWELL AND UNGER 1977; JAGADISH AND CARTER

1977), or *cdc* (JAGADISH AND CARTER 1977) mutations. In this study, to obtain the fraction of G1 cells (e.g.,

see Figures 3, 4), we used DNA content measurements by flow cytometry, as described previously

(HOOSE *et al.* 2012; HOOSE *et al.* 2013). For human cells, earlier studies employed ^{3}H-thymidine pulses or

division waves after thymidine block (BASERGA 1985). The doubling times of the NCI-60 human cancer

cell lines we included in Table S2 are known (ROSS *et al.* 2000; SCHERF *et al.* 2000), but there was no

quantitative cell cycle data for most of the cell lines. However, images of DNA content profiles for the

NCI-60 panel, albeit with no quantification, have been published (GARNER AND EASTMAN 2011). We

requested and obtained high-resolution files of these images from Dr. Alan Eastman (Geisel School of

Medicine - Dartmouth College). From the entire DNA content histogram, to quantify the fraction of cells

in G1, we used imaging software to measure the area on the left side of the G1 peak (from peak to

valley) and multiplied this area by two, as has been described previously (JOHNSTON *et al.* 1980). This

approach avoids complications from heavy right side tails due to S phase cells and yields an acceptable

estimate of the relative G1 length as a fraction of total cell cycle time. We then combined these values

with all others available from human cells, cancerous and normal (Sɪsκεɴ ᴀɴᴅ Kɪɴοsɪᴛᴀ 1961; Dᴇғᴇɴᴅɪ ᴀɴᴅ

Mᴀɴsοɴ 1963; Lᴇɴɴᴀʀᴛᴢ ᴀɴᴅ Mᴀᴜʀᴇʀ 1964; Aοκɪ ᴀɴᴅ Mοοʀᴇ 1970; Bᴀsᴇʀɢᴀ 1985; Kᴜᴍᴇɪ *et al.* 1989; Bʀοɴs

et al. 1992; Lᴜᴄɪᴀɴɪ *et al.* 2001; Hᴀʜɴ *et al.* 2009), to compile a dataset of 96 values for G1 length (T_{G1})

and doubling time (T_d) for human cells, shown in Table S2.

Chemostat cultures. The experiments were done using a New Brunswick BioFlo (BF-110) reactor with a

working volume of 880 mL, run at room temperature, as described earlier(Hᴇɴʀʏ *et al.* 2010). In each

experiment and at each dilution rate the reactor was sampled several times to measure protein levels by

immunoblots, the DNA content with flow cytometry, and the cell size and cell density of the culture

using a Beckman Z2 channelyzer (Hᴇɴʀʏ *et al.* 2010; Hοοsᴇ *et al.* 2012), as indicated. We measured the

cell density at every sampling, to ensure that we never reached 'wash-out' conditions at the high

dilution rates. In every experiment the cell density remained >1E+07 cells/ml, and did not vary more

than 3-fold between the lowest and highest dilution rates.

Protein surveillance. Protein were resolved onto 4-12% Tris-Glycine gels (Thermo Scientific, Cat#:

XP04125BOX). Cln3p-$(Myc)_{13}$ was detected with an anti-Myc antibody (Abcam, Cat #: ab13836). All other

procedures for TAP-tagged protein detection, extract preparation for immunoblots and their analysis

has been described elsewhere (Bʟᴀɴκ *et al.* 2017).

Statistical analysis. All data were analyzed and displayed with R language packages. To build the linear

models we described using the values for the yeast (Tables S1) and human (Tables S2) datasets, we first

examined if the assumptions for building simple, linear parametric models were satisfied. The diagnostic

residual plots evaluating whether the errors were independent of each other, normally distributed

around a mean of zero and equal variance, are shown in Figure S2. In both the yeast and human

datasets, the existence of a few outlier points appeared to violate the necessary assumptions (Figure S2;

$p<0.05$ for assessment of the assumptions using the global test on 4 degrees-of-freedom, with the

'gvlma' function of the *gvlma* R language package (PENA AND SLATE 2006)). Hence, we opted for non-

parametric, robust linear regression models based on Siegel repeated medians (Table 1).

RESULTS

Association between T_{G1} and T_d

We compiled the available values for T_d and T_{G1} from the literature for budding yeast (Table S1) and

human (Table S2) cells (see Materials and Methods). To model T_{G1} against T_d, it is necessary to first

examine the distribution of T_{G1} values. We found that T_{G1} values were not normally distributed for yeast

(p = 4.434E-14, Shapiro-Wilk test, using the 'shapiro.test' function of the *stats* R package) or human cells

(p = 1.039E-07, Shapiro-Wilk test). Instead, T_{G1} values fit better a log-normal distribution. For example,

for yeast T_{G1} values, the Anderson-Darling statistic was the lowest for the log-normal distribution

(0.347), compared to other distributions (Weibull: 1.693; gamma: 1.521; exponential: 2.462), calculated

with the 'ad.test' function of the *nortest* R language package. As expected for log-normal distributions,

log-transformed values of T_{G1} were normally distributed for yeast (Figure S1A-D, p = 0.1871, Shapiro-

Wilk test) and human cells (Figure S1E-H, p = 0.3099, Shapiro-Wilk test). The apparent log-normal

distribution of T_{G1} values is consistent with a multiplicative process of many, positive, independent

random variables that determine G1 length (KOCH 1966). In cell proliferation, log-normality has been

proposed to reflect exponential patterns of growth in mass. Despite fluctuations in the growth rate

constant, the growth of the overwhelming majority of cellular components is influenced similarly,

leading to log-normality (KOCH AND SCHAECHTER 1962; KOCH 1966). In budding yeast and other cell types

there is evidence for exponential patterns of protein synthesis (ELLIOTT AND MCLAUGHLIN 1978; DI TALIA *et*

al. 2007) and increase in mass in the cell cycle (BRYAN *et al.* 2012; SON *et al.* 2012). Our data are

accommodated by such considerations and, for display purposes, we log-transformed the relevant

values (e.g., see Figure 1).

To test for association between T_{G1} and T_d we used the distribution-free Spearman's and Kendall's

tests for independence based on ranks. We opted for these non-parametric, distribution-free tests

because of the existence of outliers even in log-transformed T_{G1} values (e.g., see Figure S1). The high

values (> 0.75) of the rank correlation coefficients (τ for Kendall's and r for Spearman's; see Table 1)

show a strong positive association between T_{G1} and T_d, for both yeast (Figure 1A) and human (Figure 1C)

cells. Interestingly, however, the duration of the non-G1 phases (T_{nonG1}) of the cell cycle were also

positively correlated with T_d (Figures 1B, D; and Table 1) in both organisms, albeit less so in yeast

(τ=0.398; Figure 1B) than in human cells (τ=0.579; Figure 1D). Overall, these data document the tight

association between the duration of the G1 phase and population doubling time. Additionally, they

clearly show that growth requirements for cell division are also registered later in the cell cycle, in

agreement with observations from other groups (ANASTASIA *et al.* 2012; FERREZUELO *et al.* 2012; DOWLING

et al. 2014; SOIFER AND BARKAI 2014; CERULUS *et al.* 2016; MAYHEW *et al.* 2017).

Lack of association between T_{G1}, or T_d, and cell size

The size of yeast cells was not significantly associated with T_d (p-value= 0.171; Figure 2A) or T_{G1} (p-value=

0.2449; Figure 2B). The lack of association between cell size and T_d or T_{G1} may seem surprising in the

context of some earlier reports in yeast (TYSON *et al.* 1979), and the growth law in bacteria where an

exponential relation between cell size and doubling time (SCHAECHTER *et al.* 1958) accommodates

overlapping DNA replication cycles (SI *et al.* 2017). Based on those models, one would have expected a

negative association between size and growth rate, with cells getting smaller with larger T_d values. If

anything, in some nutrients yeast cells may become larger as they divide slower (note that some of the

largest cell sizes were recorded for long T_d values in Figure 2; see also Figure 4D, under leucine

limitation, where contrary to expectations cells appeared to increase in size as they divide slower

instead of getting smaller). Yet previous studies reported a negative association between growth rate

and cell size (TYSON *et al.* 1979). For example, we had also reported that cells proliferating slower in

media with poor carbon sources were smaller than cells proliferating faster in glucose-replete medium

(SOMA *et al.* 2014). However, all these examples were from batch cultures, where growth rate cannot be

adjusted independently of a specific nutrient. In contrast, in all examples from steady-state cultures,

when growth rate was adjusted independently of nutrient composition in yeast, there was no clear

pattern between T_d and cell size (GUO *et al.* 2004; BRAUER *et al.* 2008) (see also Figures 3, 4). Importantly,

in all these previous chemostat studies, cell size measurements were internally calibrated, and the lack

of any pattern of correlation between cell size and growth-rate cannot be attributed to experimental

variabilities of different studies incorporated in our meta-analysis of the literature in Table 1. Likewise,

from a genome-wide survey of single gene deletions (HOOSE *et al.* 2012), there was no pattern of

correlation between cell size and the relative duration of the G1 phase. Hence, although cell size can be

modulated by changes in nutrient composition in yeast, it is more likely that these are nutrient-specific

effects, not causally linked to growth rate changes. This interpretation is also consistent with models in

bacterial systems, which view cell size as a spandrel and the reported cell size effects of growth rate as

secondary to other causes (AMIR 2017).

A scaling relation between T_{G1} and T_d

To estimate a predictive and quantitative relation between T_{G1} and T_d we derived non-parametric,

robust linear regression models using the Siegel repeated median estimates (SIEGEL 1982), (Figure 1 and

Table 1; see also Materials and Methods). The intercepts of the linear T_{G1} vs. T_d plots reflect the

apparent minimum duration of the S+G2+M phases in yeast (1.4 h; Table 1) and human (7.6 h; Table 1)

cells. The slope of the regression of T_{G1} on T_d appeared to differ between the yeast and human datasets

(Figure 1 and Table 1). Applying the non-parametric Sen-Adichie test for parallelism, using the function

'sen.adichie' of the *NSM3* R package, confirmed that the difference in the slopes of the regression lines

of T_{G1} on T_d between yeast and human cells was statistically significant (V statistic = 7.324, p-value =

0.007). The particular slopes (≈0.6 for yeast and ≈0.75 for human cells) represent the exponents in the

linear relations of the corresponding log-transformed T_{G1} and T_d values. Such values are often

incorporated in scaling relations between the measured variables (CHAN AND MARSHALL 2010). For

example, a slope of ≈0.75 of the regression of T_d on T_{G1} brings to mind the famous Kleiber's law that an

animal's metabolic rate scales to the ¾ power of the animal's mass (CALDER 1981). The evidence we

presented above is not sufficient to make a case for any mechanistic connection with Kleiber's

parameters and the decision of human cells to initiate a new cell cycle. Instead, the quantitative

relations we identified linking T_{G1} with T_d are significant because they provide a framework to interpret

experimental perturbations in cell cycle progression and cell proliferation, as we describe next for yeast

cells.

Loss of Cln3p, but not of Whi5p, alters the relation between T_{G1} and T_d

To understand how the relation between the length of the G1 phase and doubling time is established in

budding yeast, we next examined the role of the Cln3p and Whi5p proteins, which regulate the G1/S

transition in this organism. It has been proposed that dilution of Whi5p as cells get bigger in G1 is the

key event controlling the timing of the G1/S transition (SCHMOLLER AND SKOTHEIM 2015; SCHMOLLER *et al.*

2015). Since in cells proliferating at different rates there is not a significant correlation with cell size

(Figure 2, Table S1), it is not clear how the inhibitor dilution model would apply to the conditions we

examine in this study. To test the role of Cln3p and Whi5p in the relation between T_{G1} and T_d, we

examined the cell cycle profile of *cln3Δ* or *whi5Δ* cells in continuous, steady-state chemostat cultures. To

our knowledge, the kinetics of cell cycle progression in cells lacking Cln3p or Whi5p have not been

examined previously in chemostat cultures. We measured T_{G1} in *cln3Δ* or *whi5Δ* cells from the DNA

content of the cultures under glucose (0.08% $^w/_v$) limitation and at different dilution rates (0.068 h^{-1} to

0.348 h^{-1}; corresponding to T_d values between 10 h and 2 h, respectively). As expected, cells lacking

Whi5p were very small ($\approx$18 fL), and their size did not change significantly as a function of T_d (Figure 3C).

Furthermore, the intercept of the linear fit between the log-transformed T_d and T_{G1} values of *whi5Δ* cells

was higher than the intercept of the linear fit of these parameters in wild-type cells (1.17 vs. 0.92; see

Table 1), consistent with the shortened G1 phase of *whi5Δ* cells. Remarkably, however, the slope of the

linear relation between T_d and T_{G1} was similar to what we observed in wild-type cells (Figures 1A, 3A,

and Table 1; Sen-Adichie V statistic = 1.775, p-value = 0.183). Hence, although in absolute terms the G1

phase is shortened at all growth rates tested, *whi5Δ* cells nonetheless maintain the proportional

changes in T_{G1} and T_d upon nutrient limitation. Dilution of Whi5p in cells dividing in a given growth state

may account for the size dependency of the G1/S transition in that physiological state (SCHMOLLER *et al.*

2015). However, our data show that in different physiological states, and despite their shortened G1

phase and small size, *whi5Δ* cells remain fully responsive to different environments, displaying the

expected proportional changes between T_d and T_{G1}. In contrast, cells lacking Cln3p had an abnormal

behavior. T_{G1} did not even have a straightforward linear relation with T_d in cultures of *cln3Δ* cells. At

shorter division times, from T_d between 2 h and 6 h, the T_d and T_{G1} values of *cln3Δ* cultures were related

linearly (Figure 3B; slope = 0.7685), but this relation breaks down completely at low dilution rates, with a

T_d greater than $\approx$7-8 h (Figure 3B). At those long doubling times, the size of *cln3Δ* cells increased even

more than their already large cell size (Figures 3D). These data suggest that Cln3p, but not Whi5p, is

necessary for imposing the proper scaling relation between T_{G1} and T_d in wild type cells.

Measurements of Cln3p and Whi5p steady-state levels at different growth rates

Given the necessary role of Cln3p in establishing the physiological relation between T_d and T_{G1} (Figure 3),

we sought to measure the levels of Cln3p and Whi5p as a function of growth rate. There are no reports

of the steady-state levels of Cln3p or Whi5p in cell populations proliferating at different rates in

chemostats. To measure Whi5p and Cln3p levels from the same cells, we generated a strain that carries

WHI5-TAP and *CLN3-13MYC* alleles, providing the only source of these gene products in the cells,

expressed from their endogenous chromosomal locations (Figure 4A, see Materials and Methods). The

expressed proteins were epitope-tagged, but otherwise un-mutated, wild type Whi5p-TAP and Cln3p-

$(Myc)_{13}$. These cells were then cultured in continuous, steady-state chemostat cultures under glucose

(0.08% $^w/_v$) or leucine (0.0015% $^w/_v$) limitation. Although the *CLN3-13MYC* allele provides the means for

reliable detection of otherwise wild type Cln3p, it is known to be hypermorphic, stabilizing the Cln3p

protein somewhat and shortening the G1 phase of the cell cycle (THORBURN *et al.* 2013). Indeed, the

intercept of the linear fit between the log-transformed T_d and T_{G1} values of this strain (HB94; *WHI5-TAP*,

CLN3-13MYC) was higher than the intercept of the linear fit of these parameters in wild-type cells (0.97

vs. 0.92; see Table 1), consistent with a shortened G1 phase. Furthermore, the slope of the linear

relation between the log-transformed values of T_d and T_{G1} was increased (0.68 vs. 0.6; see Table 1),

arguing that even moderate stabilization of the Cln3p protein alters the relation between T_d and T_{G1}.

Nonetheless, these cells still displayed a strong, linear, positive association between T_d and T_{G1} (τ=0.77;

r=0.92; see Table 1). Hence, we proceeded to quantify the levels of both Whi5p-TAP and $Cln3p-(Myc)_{13}$,

from separate chemostat experiments under glucose or leucine limitation, each run at ≥5 different

dilution rates (Figure 4; see Materials and Methods). Surprisingly, the levels of Whi5p-TAP were reduced

in slower proliferating cells (Figure 4A, B). Previously, it was reported that Whi5p abundance increases

≈3-fold in cells growing in poorer carbon sources (LIU *et al.* 2015). There are several variables that were

different between that study and ours, which could account for the disagreement in the findings: First,

different epitope-tagged alleles were used (*WHI5-tdTomato* vs. *WHI5-TAP*). Second, different detection

methods were applied (fluorescence live cell imaging vs. immunoblots). Third, and most significantly,

nutrient-specific effects could not be separated from growth rate-specific ones in (LIU *et al.* 2015).

We observed a much more significant and disproportionate reduction of Cln3p-(Myc)$_{13}$ levels in

slower-proliferating cells (Figure 4A, B). Normalized against the total cellular protein content, the fastest

proliferating populations had >10-fold higher levels of Cln3p-(Myc)$_{13}$ compared to the slowest

proliferating cells (Figure 4A, B). Furthermore, because these estimates rely on the hypermorphic *CLN3-*

13MYC allele that produces a more stabilized Cln3p protein (THORBURN *et al.* 2013), the dynamic range of

Cln3p levels as a function of growth rate is likely even greater.

A uORF in *CLN3* adjusts the levels of Cln3p at different growth rates

What is the mechanism that underpins the growth rate-dependent control of Cln3p abundance? We had

predicted that a uORF in *CLN3* may inhibit disproportionately its translational efficiency in poor media

(POLYMENIS AND SCHMIDT 1997). However, such predictions have not been accompanied with

measurements of Cln3p levels. A kinetic model of protein synthesis (LODISH 1974) forecasts that

removing the uORF would de-repress synthesis of Cln3p in low growth rates when the ribosome content

of the cell is low. In contrast, removing the *CLN3* uORF would have minimal effects in cells that

proliferate fast, when the ribosome content is high. To test this model, we introduced an A-315T

substitution that mutates the start codon of the uORF in *CLN3* without affecting *CLN3* mRNA levels

(POLYMENIS AND SCHMIDT 1997), in the strain that expresses otherwise wild type Whi5p-TAP and Cln3p-

(Myc)$_{13}$ (see Materials and Methods). Indeed, in *CLN3* uORF mutant cells proliferating slower (T$_d$ > 4h),

the dynamic range of Cln3p levels as a function of growth rate was narrower and indistinguishable from

the range of Whi5p levels (Figure 4A, B and Source Data). A hallmark phenotypic readout of gain-of-

function *CLN3* alleles is a reduction in cell size (NASH *et al.* 1988). Consistent with a de-repression of

Cln3p synthesis, not only cells that lack the *CLN3* uORF were smaller than their wild type counterparts

were, but also this effect was growth rate-dependent (see Figure 4C, D) in glucose and especially in

leucine-limited cells, with the latter displaying pronounced enlargement that was suppressed by

removing the *CLN3* uORF. These data argue that translational control contributes to the

disproportionate reduction of Cln3p levels as a function of growth rate. Overall, higher Cln3p levels, by

stabilizing the protein or by de-repressing Cln3p synthesis via translational control, alter the slope of the

linear relation between T_d and T_{G1} (see Table 1). Conversely, loss of Cln3p severely perturbs the linear

relation altogether (Figure 3). These results underscore the critical role of the G1 cyclin Cln3p in the

physiological coupling between growth and division.

DISCUSSION

The scaling relations between G1 length and population time in yeast and human cells we report are

important for several reasons. First, if the duration of G1 is estimated, they allow predictions of

proliferation rates, which could be useful in diverse settings, such as in tissues at an organismal level.

Second, they serve as benchmarks against which the effects of genetic or other perturbations can be

evaluated, as we demonstrated for Whi5p and Cln3p, two cell cycle regulators in yeast. Third, scaling

relations of cellular physiology may ultimately point to general, physical mechanisms that organize life at

the cellular level.

How do the results we report here impinge on the Whi5p dilution model (SCHMOLLER *et al.* 2015),

which was reported to account for the coupling between growth and division in budding yeast? As we

noted above, our data dealt with growth rate-dependent changes in the cell cycle, which was not

addressed in (SCHMOLLER *et al.* 2015). Hence, the two studies are not directly comparable. A critical

aspect of the Whi5p dilution model, however, was that while Whi5p levels were disproportionately

lower than expected from cell growth, Cln3p levels were roughly constant and proportional to the

increase in size from birth to START (SCHMOLLER *et al.* 2015). Given the strong and disproportionate

dependency of Cln3p levels on growth rate that we report here and had predicted earlier (POLYMENIS AND

SCHMIDT 1997), could the uORF-mediated translational control of *CLN3* affect Cln3p synthesis in the G1

phase from birth to START? We think this is unlikely because the uORF-mediated translational control

we described operates when the concentration of active ribosomes in the cell changes (LODISH 1974), for

example in poor vs. rich nutrients. To our knowledge, there is no report of cell cycle-dependent changes

in ribosome content.

Other mechanisms may contribute to significant changes in Cln3p abundance in G1, which could

have implications for the Whi5p dilution model. There are conflicting reports in the literature about

whether Cln3p 'cycles' in the cell cycle. The initial report claiming that Cln3p levels are constant in the

cell cycle did not interrogate the early G1 phase (Tyers *et al.* 1993). In that report, although early G1,

small (25 fL), elutriated daughter cells were collected, Cln3p levels were not measured until much later

in G1 (at 35 fL, when by 40 fL 25% of the cells were already budded; see Fig. 4 in (Tyers *et al.* 1993)).

Based on that result, it had been assumed for decades that Cln3p levels were constant in the cell cycle.

Cln3p abundance cannot be properly addressed in the single-cell microscopy methods of (Schmoller *et*

al. 2015), because a mutated *CLN3* allele had to be used, producing a stabilized Cln3p protein so that it

can accumulate at high enough, but non-physiological levels to be visible with microscopy. Recently,

however, two independent studies by the Kellogg (Zapata *et al.* 2014) and Amon (Thorburn *et al.* 2013)

labs, assayed elutriated synchronous cells carrying epitope-tagged, but otherwise wild type *CLN3* alleles.

Both studies showed that Cln3p levels oscillate >10-fold in G1. Cln3p is absent in early G1 cells, while it

rises dramatically before START. Hence, although a 2-fold dilution of Whi5p is observed in G1 (Schmoller

et al. 2015), the changes in Cln3p levels are likely more pronounced (Thorburn *et al.* 2013; Zapata *et al.*

2014), through transcriptional (Zapata *et al.* 2014), or other mechanisms. It is important to stress,

however, those changes of Cln3p levels in G1 do not affect the fundamental aspect of the inhibitor

dilution model, namely that Whi5p levels are reduced by cell growth. Instead, we may be dealing with a

more complex, 'mixed' model of inhibitor dilution *and* activator accumulation.

ACKNOWLEDGEMENTS

Author Contributions: Methodology, H.M.B., M.P.; Formal Analysis, H.M.B., M.P.; Investigation, H.M.B.,

M.C., I.P.E.P., A.O.P., M.P.; Resources, M.P.; Data Curation, H.M.B., M.P.; Writing – Original Draft, M.P.,

Writing – Review and Editing, H.M.B., M.P.; Visualization, H.M.B., M.P.; Supervision, H.M.B., M.P.,

Funding Acquisition, M.P.

This work was supported from the NIH (grant GM123139 to M.P.). M.C. was supported by a National

Science Foundation REU award (DBI-1358941). The authors declare no conflicts of interest.

LITERATURE CITED

Amir, A., 2017 Is cell size a spandrel? Elife 6.

Anastasia, S. D., D. L. Nguyen, V. Thai, M. Meloy, T. MacDonough *et al.*, 2012 A link between mitotic

entry and membrane growth suggests a novel model for cell size control. J Cell Biol 197**:** 89-104.

Aoki, Y., and G. E. Moore, 1970 Comparative study of mitotic stages of cells derived from human

peripheral blood. Exp Cell Res 59**:** 259-266.

Baserga, R., 1985 *The Biology of Cell Reproduction*.

Blank, H. M., R. Perez, C. He, N. Maitra, R. Metz *et al.*, 2017 Translational control of lipogenic enzymes in

the cell cycle of synchronous, growing yeast cells. EMBO J 36**:** 487-502.

Brauer, M. J., C. Huttenhower, E. M. Airoldi, R. Rosenstein, J. C. Matese *et al.*, 2008 Coordination of

growth rate, cell cycle, stress response, and metabolic activity in yeast. Mol Biol Cell 19**:** 352-

367.

Brons, P. P., J. M. Raemaekers, M. J. Bogman, P. E. van Erp, J. B. Boezeman *et al.*, 1992 Cell cycle kinetics

in malignant lymphoma studied with in vivo iododeoxyuridine administration, nuclear Ki-67

staining, and flow cytometry. Blood 80**:** 2336-2343.

Brunner, N., D. Bronzert, L. L. Vindelov, K. Rygaard, M. Spang-Thomsen *et al.*, 1989 Effect on growth and

cell cycle kinetics of estradiol and tamoxifen on MCF-7 human breast cancer cells grown in vitro

and in nude mice. Cancer Res 49**:** 1515-1520.

Bryan, A. K., A. Engler, A. Gulati and S. R. Manalis, 2012 Continuous and long-term volume

measurements with a commercial Coulter counter. PLoS One 7**:** e29866.

Calder, W. A., 3rd, 1981 Scaling of physiological processes in homeothermic animals. Annu Rev Physiol

43**:** 301-322.

Carter, B. L., and M. N. Jagadish, 1978 Control of cell division in the yeast Saccharomyces cerevisiae

cultured at different growth rates. Exp Cell Res 112**:** 373-383.

Cerulus, B., A. M. New, K. Pougach and K. J. Verstrepen, 2016 Noise and Epigenetic Inheritance of Single-

Cell Division Times Influence Population Fitness. Curr Biol 26**:** 1138-1147.

Chan, Y. H., and W. F. Marshall, 2010 Scaling properties of cell and organelle size. Organogenesis 6**:** 88-

96.

Costanzo, M., J. L. Nishikawa, X. Tang, J. S. Millman, O. Schub *et al.*, 2004 CDK activity antagonizes Whi5,

an inhibitor of G1/S transcription in yeast. Cell 117**:** 899-913.

Cross, F. R., 1988 DAF1, a mutant gene affecting size control, pheromone arrest, and cell cycle kinetics of

Saccharomyces cerevisiae. Mol Cell Biol 8**:** 4675-4684.

de Bruin, R. A., W. H. McDonald, T. I. Kalashnikova, J. Yates, 3rd and C. Wittenberg, 2004 Cln3 activates

G1-specific transcription via phosphorylation of the SBF bound repressor Whi5. Cell 117**:** 887-

898.

Defendi, V., and L. A. Manson, 1963 Analysis of the life-cycle in mammalian cells. Nature 198**:** 359-361.

Di Talia, S., J. M. Skotheim, J. M. Bean, E. D. Siggia and F. R. Cross, 2007 The effects of molecular noise

and size control on variability in the budding yeast cell cycle. Nature 448**:** 947-951.

Dowling, M. R., A. Kan, S. Heinzel, J. H. Zhou, J. M. Marchingo *et al.*, 2014 Stretched cell cycle model for

proliferating lymphocytes. Proc Natl Acad Sci U S A 111**:** 6377-6382.

Elliott, S. G., and C. S. McLaughlin, 1978 Rate of macromolecular synthesis through the cell cycle of the

yeast Saccharomyces cerevisiae. Proc Natl Acad Sci U S A 75**:** 4384-4388.

Ferrezuelo, F., N. Colomina, A. Palmisano, E. Gari, C. Gallego *et al.*, 2012 The critical size is set at a single-

cell level by growth rate to attain homeostasis and adaptation. Nat Commun 3**:** 1012.

Fisher, R. P., 2016 Getting to S: CDK functions and targets on the path to cell-cycle commitment.

F1000Research 5**:** 2374.

Garner, K. M., and A. Eastman, 2011 Variations in Mre11/Rad50/Nbs1 status and DNA damage-induced

S-phase arrest in the cell lines of the NCI60 panel. BMC Cancer 11**:** 206:201-213.

Guo, J., B. A. Bryan and M. Polymenis, 2004 Nutrient-specific effects in the coordination of cell growth

with cell division in continuous cultures of Saccharomyces cerevisiae. Arch Microbiol 182**:** 326-

330.

Hahn, A. T., J. T. Jones and T. Meyer, 2009 Quantitative analysis of cell cycle phase durations and PC12

differentiation using fluorescent biosensors. Cell Cycle 8**:** 1044-1052.

Hartwell, L. H., and M. W. Unger, 1977 Unequal division in Saccharomyces cerevisiae and its implications

for the control of cell division. J Cell Biol 75**:** 422-435.

Henry, K. A., H. M. Blank, S. A. Hoose and M. Polymenis, 2010 The unfolded protein response is not

necessary for the G1/S transition, but it is required for chromosome maintenance in

Saccharomyces cerevisiae. PLoS One 5**:** e12732.

Hoffman, J. G., 1949 Theory of the mitotic index and its application to tissue growth measurement. Bull

Math Biophys 11**:** 139-144.

Hoose, S. A., J. A. Rawlings, M. M. Kelly, M. C. Leitch, Q. O. Ababneh *et al.*, 2012 A systematic analysis of

cell cycle regulators in yeast reveals that most factors act independently of cell size to control

initiation of division. PLoS Genet 8**:** e1002590.

Hoose, S. A., J. T. Trinh, M. C. Leitch, M. M. Kelly, R. F. McCormick *et al.*, 2013 Saccharomyces cerevisiae

deletion strains with complex DNA content profiles. FEMS Microbiol Lett 345**:** 72-76.

Jagadish, M. N., and B. L. Carter, 1977 Genetic control of cell division in yeast cultured at different

growth rates. Nature 269**:** 145-147.

Johnston, G., R. Singer, S. Sharrow and M. Slater, 1980 Cell division in the yeast Saccharomyces

cerevisiae growing at different rates. Microbiology 118**:** 479-484.

Johnston, G. C., J. R. Pringle and L. H. Hartwell, 1977 Coordination of growth with cell division in the

yeast Saccharomyces cerevisiae. Exp Cell Res 105**:** 79-98.

Koch, A. L., 1966 The logarithm in biology. 1. Mechanisms generating the log-normal distribution exactly.

J Theor Biol 12**:** 276-290.

Koch, A. L., and M. Schaechter, 1962 A model for statistics of the cell division process. J Gen Microbiol

29**:** 435-454.

Kumei, Y., T. Nakajima, A. Sato, N. Kamata and S. Enomoto, 1989 Reduction of G1 phase duration and

enhancement of c-myc gene expression in HeLa cells at hypergravity. Journal of Cell Science 93**:**

221-226.

Lennartz, K. J., and W. Maurer, 1964 Autoradiographische Bestimmung der Dauer der DNS-Verdopplung

und der Generationszeit beim Ehrlich-Ascitestumor der Maus durch Doppelmarkierung mit 14C-

und 3H-Thymidin. Zeitschrift für Zellforschung und Mikroskopische Anatomie 63**:** 478-495.

Liu, X., X. Wang, X. Yang, S. Liu, L. Jiang *et al.*, 2015 Reliable cell cycle commitment in budding yeast is

ensured by signal integration. Elife 4.

Lodish, H. F., 1974 Model for the regulation of mRNA translation applied to haemoglobin synthesis.

Nature 251**:** 385-388.

Luciani, A. M., A. Rosi, P. Matarrese, G. Arancia, L. Guidoni *et al.*, 2001 Changes in cell volume and

internal sodium concentration in HeLa cells during exponential growth and following lonidamine

treatment. Eur J Cell Biol 80**:** 187-195.

Mayhew, M. B., E. S. Iversen and A. J. Hartemink, 2017 Characterization of dependencies between

growth and division in budding yeast. J R Soc Interface 14.

Morgan, D. O., 2007 Control of Cell Proliferation and Growth, pp. 221 in *The Cell Cycle*. New Science

Press Ltd, London, UK.

Nash, R., G. Tokiwa, S. Anand, K. Erickson and A. B. Futcher, 1988 The WHI1+ gene of Saccharomyces

cerevisiae tethers cell division to cell size and is a cyclin homolog. EMBO J 7**:** 4335-4346.

Palumbo, P., M. Vanoni, V. Cusimano, S. Busti, F. Marano *et al.*, 2016 Whi5 phosphorylation embedded

in the G1/S network dynamically controls critical cell size and cell fate. 7**:** 11372.

Pena, E. A., and E. H. Slate, 2006 Global Validation of Linear Model Assumptions. J Am Stat Assoc 101**:**

341.

Polymenis, M., and E. V. Schmidt, 1997 Coupling of cell division to cell growth by translational control of

the G1 cyclin CLN3 in yeast. Genes Dev 11**:** 2522-2531.

Rivin, C. J., and W. L. Fangman, 1980 Cell cycle phase expansion in nitrogen-limited cultures of

Saccharomyces cerevisiae. J Cell Biol 85**:** 96-107.

Ross, D. T., U. Scherf, M. B. Eisen, C. M. Perou, C. Rees *et al.*, 2000 Systematic variation in gene

expression patterns in human cancer cell lines. Nat Genet 24**:** 227-235.

Schaechter, M., O. Maaloe and N. O. Kjeldgaard, 1958 Dependency on medium and temperature of cell

size and chemical composition during balanced grown of Salmonella typhimurium. J Gen

Microbiol 19**:** 592-606.

Scherf, U., D. T. Ross, M. Waltham, L. H. Smith, J. K. Lee *et al.*, 2000 A gene expression database for the

molecular pharmacology of cancer. Nat Genet 24**:** 236-244.

Schmoller, K. M., and J. M. Skotheim, 2015 The Biosynthetic Basis of Cell Size Control. Trends Cell Biol

25**:** 793-802.

Schmoller, K. M., J. J. Turner, M. Koivomagi and J. M. Skotheim, 2015 Dilution of the cell cycle inhibitor

Whi5 controls budding-yeast cell size. Nature 526**:** 268-272.

Si, F., D. Li, S. E. Cox, J. T. Sauls, O. Azizi *et al.*, 2017 Invariance of Initiation Mass and Predictability of Cell

Size in Escherichia coli. Curr Biol 27**:** 1278-1287.

Siegel, A. F., 1982 Robust regression using repeated medians. Biometrika 69**:** 242-244.

Sisken, J. E., and R. Kinosita, 1961 TIMING OF DNA SYNTHESIS IN THE MITOTIC CYCLE IN VITRO. J Biophys

Biochem Cytol 9**:** 509-518.

Slater, M. L., S. O. Sharrow and J. J. Gart, 1977 Cell cycle of Saccharomycescerevisiae in populations

growing at different rates. Proc Natl Acad Sci U S A 74**:** 3850-3854.

Soifer, I., and N. Barkai, 2014 Systematic identification of cell size regulators in budding yeast. Molecular

systems biology 10.

Soma, S., K. Yang, M. I. Morales and M. Polymenis, 2014 Multiple metabolic requirements for size

homeostasis and initiation of division in Saccharomyces cerevisiae. Microb Cell 1**:** 256-266.

Son, S., A. Tzur, Y. Weng, P. Jorgensen, J. Kim *et al.*, 2012 Direct observation of mammalian cell growth

and size regulation. Nature methods 9**:** 910-912.

Thorburn, R. R., C. Gonzalez, G. A. Brar, S. Christen, T. M. Carlile *et al.*, 2013 Aneuploid yeast strains

exhibit defects in cell growth and passage through START. Mol Biol Cell 24**:** 1274-1289.

Tyers, M., G. Tokiwa and B. Futcher, 1993 Comparison of the Saccharomyces cerevisiae G1 cyclins: Cln3

may be an upstream activator of Cln1, Cln2 and other cyclins. Embo j 12: 1955-1968.

Tyson, C. B., P. G. Lord and A. E. Wheals, 1979 Dependency of size of Saccharomyces cerevisiae cells on

growth rate. J Bacteriol 138**:** 92-98.

Zapata, J., N. Dephoure, T. Macdonough, Y. Yu, E. J. Parnell *et al.*, 2014 PP2ARts1 is a master regulator of

pathways that control cell size. J Cell Biol 204**:** 359-376.

FIGURE LEGENDS

FIGURE 1. Linking the length of the G1 phase with population doubling time. Scatter plots of T_{G1} (**A, C**)

or T_{nonG1} (**B, D**) values on the x-axis, against T_d values (y-axis). All plots used the natural logarithms of the

values for yeast (**A, B**) and human (**C, D**) cells from Tables S1 and S2, respectively. The Kendall's (τ) rank

correlation coefficient is shown in each case. In red are regression lines of the Siegel repeated medians.

The slope and the associated 95% confidence intervals of the linear model are shown in each case.

Additional statistical parameters associated with these plots are shown in Table 1.

FIGURE 2. Cell size does not correlate with T_d or T_{G1} in yeast. Scatter plots of cell size values (x-axis)

against T_d (**A**) or T_{G1} (**B**) values (y-axis) from the data shown in Table S1. In red are regression lines of the

Siegel repeated medians. The Kendall's (τ) and Spearman's (r) rank correlation coefficients are shown in

each case.

FIGURE 3. Cln3p, but not Whi5p, imposes the proper relation between T_{G1} and T_d. Scatter plots of T_{G1}

values on the x-axis, against T_d values (y-axis). All plots used the natural logarithms of the values for

whi5Δ (**A**) or *cln3Δ* (**B**) cells, sampled from chemostat cultures several times at each dilution rate, as

indicated. In red are regression lines of the Siegel repeated medians. For *whi5Δ* cells, the slope of the

linear model is shown (additional statistical parameters are in Table 1). For *cln3Δ* cells, the linear fit is

shown in dashed form and did not include Ln(T_d) values >2.0, because the relation between T_d and T_{G1}

breaks down at those proliferation rates for those cells. Scatter plots of the relation of cell size and T_d in

cells lacking Whip5 (**C**) or Cln3p (**D**), with cell size values (x-axis) plotted against T_d (y-axis) from the same

cultures described in (**A**) and (**B**).

539 **FIGURE 4. The levels of the G1 cyclin Cln3p vary over a broad range as a function of growth rate, due**

540 **to a uORF affecting translation of *CLN3*.** Scatter plots of the relative abundance (y-axis) of Cln3p-(Myc)$_{13}$

541 and Whi5p-TAP in otherwise wild type *CLN3-13MYC*, *WHI5-TAP* cells (**A**), or *CLN3* uORF (*A-315T-CLN3*)

542 mutant cells (**B**), against T$_d$ (x-axis). Each data point in the scatter plots is the average of immunoblot

543 signal intensities run in duplicate, and detected with antibodies against the Myc or TAP epitopes (see

544 Materials and Methods). All the raw immunoblots used to quantify protein levels are shown in the

545 Source Data (Supplementary File 1). Before averaging, each individual signal intensity value was

546 normalized against loading in the corresponding immunoblot lane (visualized with Ponceau staining, see

547 Supplementary File 1 and Materials and Methods). For each Cln3p-(Myc)$_{13}$ or Whi5p-TAP relative unit

548 (r.u.) shown in the scatterplots (**A, B**), the normalized, averaged intensities were scaled by the lowest

549 value (set to 1) for each protein in the given chemostat experiment run at different dilution rates.

550 Scatter plots of the relation of cell size (x-axis) and T$_d$ (y-axis) in the indicated strains, cultured under

551 glucose (**C**) or leucine (**D**) limitation, from the same cultures described in (**A**) and (**B**).

552   **TABLE 1. Statistical parameters of linear relations**

Variables		Coefficients[a]		Confidence Intervals (level = 0.95)[b]				τ^c	r^d
Y	X	Intercept	Slope	Intercept		Slope			
$(T_d)_{yeast}$	$(T_{G1})_{yeast}$	1.429	1.091	1.3817	1.5049	1.0598	1.1133	0.815	0.938
$(T_d)_{yeast}$	$(T_{nonG1})_{yeast}$	0.36	1.75	0.1900	0.8712	1.6244	2.0278	0.398	0.532
$Ln(T_d)_{yeast}$	$Ln(T_{G1})_{yeast}$	0.9168	0.5955	0.9047	0.9418	0.5815	0.6136	0.815	0.938
$Ln(T_d)_{yeast}$	$Ln(T_{nonG1})_{yeast}$	0.8682	0.8745	0.7764	0.9128	0.7827	0.9659	0.398	0.532
$(T_d)_{human}$	$(T_{G1})_{human}$	7.558	1.279	6.8401	8.5971	1.2433	1.3339	0.775	0.921
$(T_d)_{human}$	$(T_{nonG1})_{human}$	6.838	1.738	3.6212	7.3905	1.6919	2.0059	0.579	0.762
$Ln(T_d)_{human}$	$Ln(T_{G1})_{human}$	1.2202	0.7656	1.1770	1.3663	0.7211	0.7843	0.775	0.921
$Ln(T_d)_{human}$	$Ln(T_{nonG1})_{human}$	1.3608	0.7816	1.1374	1.4674	0.7487	0.8762	0.579	0.762
$Ln(T_d)_{whi5\Delta}$[e]	$Ln(T_{G1})_{whi5\Delta}$[e]	1.1654	0.6166	1.1613	1.1910	0.6083	0.6362	0.928	0.987
$Ln(T_d)_{HB94/7}$[f]	$Ln(T_{G1})_{HB94/7}$	0.9691	0.6848	0.9031	0.9843	0.6588	0.7461	0.765	0.918
$Ln(T_d)_{HB104}$[g]	$Ln(T_{G1})_{HB104}$	0.9913	0.7048	0.9902	1.0074	0.6930	0.7120	0.934	0.989

[a]From the linear fit of Siegel repeated medians, calculated with the 'mblm' function of the *mblm* R

package.

[b]Calculated with the 'confint.mblm' function of the *mblm* R package.

[c]Kendall's (τ) rank correlation coefficient, calculated with the 'cor.test' function of the *stats* R package.

[d]Spearman's (r) rank correlation coefficient, calculated with the 'cor.test' function of the *stats* R

package.

[e]The values for *whi5Δ* cells were from the experiment shown in Figure 3A.

[f]The values for HB94/7 cells (*CLN3-13MYC, WHI5-TAP*) were from the experiments in Figure 4A.

[g]The values for HB104 cells (*A-315T-CLN3-13MYC, WHI5-TAP*) were from the experiments in Figure 4B.

SUPPLEMENTARY INFORMATION

SUPPLEMENTARY FIGURE LEGENDS

FIGURE S1. Goodness-of-fit plots for log-normal distribution of yeast (A-D), and human (E-H), T_{G1}

values, from Tables S1, and S2, respectively. A, E. Histogram of experimental T_{G1} values, with the

theoretical log-normal density plot shown in red. **B, F.** Plot of the cumulative distribution function (CDF)

of experimental T_{G1} values (shown as open symbols), along with the theoretical CDF of the log-normal

distribution (shown in red). **C, G.** Quantile-Quantile (Q-Q) plot, comparing the experimental quantiles of

T_{G1} values (shown as open symbols) against the theoretical quantiles of a log-normal distribution (shown

as the diagonal line). Q-Q plots emphasize poor fit at the tails of the distribution. **D, H.** Probability-

Probability (P–P) plot of the CDF of experimental T_{G1} values plotted against the theoretical CDF of the

log-normal distribution (shown as the diagonal line). P-P plots emphasize poor fit at the center of the

distribution. The plots were generated with the *fitdistrplus* R package.

FIGURE S2. Diagnostic plots of simple linear regression models for T_d and T_{G1} values of yeast (A-D), and

human (E-H), cells. The log-transformed T_d and T_{G1} values shown in Tables S1 and S2 were used to build

simple linear regression models with the 'lm' function of the *stats* R package. **A, E.** Plot of residuals (y-

axis) against fitted values (x-axis). Numbered data points indicate problematic outliers. **B, F**. Normal Q-Q

plot of the ordered standardized residuals (y-axis), against the expected order statistics from a standard

normal distribution (x-axis). Points away from the straight dashed line indicate departures from

normality. **C, G**. Scale-Location plot of the square root of the absolute residuals (y-axis) against fitted

values (x-axis), in order to diminish skewness. **D, H**. Plot of residuals (y-axis) against leverages (x-axis).

The residual-leverage plot shows contours (shown with dashed red lines) of equal Cook's distance (by

default 0.5 and 1). Plots in **B, C, D, F, G, H** use standardized residuals which have identical variance

(under the hypothesis of normality). The solid red lines in **A, C, D, E, G, H** indicate the locally weighted

scatterplot smoothing.

Table S1. *S. cerevisiae* **cell cycle kinetics in different media**

Culture medium	Cell size (fL)	T_d (h)	T_{G1} (h)	Reference
YP+Dextrose	34.3	1.3	0.3	(TYSON *et al.* 1979)
YP+Fructose	36.6	1.3	0.3	(TYSON *et al.* 1979)
YP+Sucrose	38.7	1.3	0.3	(TYSON *et al.* 1979)
YP+Dextrose		1.4	0.3	(SLATER *et al.* 1977)
MM2+Fructose	34	1.5	0.4	(TYSON *et al.* 1979)
MM2+Dextrose	25.9	1.6	0.5	(TYSON *et al.* 1979)
YP+Raffinose		1.7	0.4	This study
YP+Dextrose		1.7	0.5	This study
YP+Raffinose	18	1.8	0.6	(TYSON *et al.* 1979)
MM+Dextrose+Ammonia		1.8	0.9	(RIVIN AND FANGMAN 1980)
MM+Dextrose+Glutamine		1.8	1.0	(RIVIN AND FANGMAN 1980)
MM1+Dextrose	17.8	2.0	0.7	(TYSON *et al.* 1979)
MM1+Maltose	25.5	2.0	0.7	(TYSON *et al.* 1979)
MM1+Fructose	22.8	2.1	0.8	(TYSON *et al.* 1979)
YP+Maltose	21.2	2.1	0.9	(TYSON *et al.* 1979)
SF		2.2	0.7	(SLATER *et al.* 1977)
YP+Glycerol		2.2	0.7	This study
YP+Ethanol		2.2	0.8	This study
Glucose-limited chemostat	38.8	2.3	0.5	(BRAUER *et al.* 2008)
Leucine-limited chemostat	27.6	2.3	0.7	(BRAUER *et al.* 2008)
Uracil-limited chemostat	41	2.3	0.7	(BRAUER *et al.* 2008)
Phosphorous-limited chemostat	46.3	2.3	0.8	(BRAUER *et al.* 2008)
YP+Sorbitol	15.9	2.3	0.8	(TYSON *et al.* 1979)
YP+Galactose		2.3	0.8	This study
Nitrogen-limited chemostat	30.6	2.3	1	(BRAUER *et al.* 2008)
Sulfur-limited chemostat	29.9	2.3	1	(BRAUER *et al.* 2008)
Glucose-limited chemostat	18.9	2.3	1	(GUO *et al.* 2004)
Nitrogen-limited chemostat	30	2.3	1	(GUO *et al.* 2004)
Glucose-limited chemostat		2.3	1.5	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.3	1.5	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.3	1.6	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.3	1.6	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.3	2.1	(HENRY *et al.* 2010)
YP+Gluconate	25.8	2.4	0.6	(TYSON *et al.* 1979)
MM+Dextrose+Glutamate		2.4	0.7	(JOHNSTON *et al.* 1980)
MM+Dextrose+Ammonia		2.6	0.9	(JOHNSTON *et al.* 1980)
YP+Galactose	21.5	2.6	1.1	(TYSON *et al.* 1979)
Glucose-limited chemostat		2.7	0.3	(JAGADISH AND CARTER 1977)
Glucose-limited chemostat		2.7	0.4	(JAGADISH AND CARTER 1977)
YP+Glycerol	24.1	2.7	1.2	(TYSON *et al.* 1979)
Uracil-limited chemostat	37.3	2.8	0.7	(BRAUER *et al.* 2008)
Glucose-limited chemostat	33.6	2.8	1	(BRAUER *et al.* 2008)
YP+Dextrose		2.8	1.0	(JAGADISH AND CARTER 1977)

Nitrogen-limited chemostat	23.9	2.8	1.1	(BRAUER *et al.* 2008)
Nitrogen-limited chemostat	29.9	2.8	1.1	(GUO *et al.* 2004)
Leucine-limited chemostat	36.6	2.8	1.2	(BRAUER *et al.* 2008)
Phosphorous-limited chemostat	37.3	2.8	1.2	(BRAUER *et al.* 2008)
Glucose-limited chemostat	19	2.8	1.3	(GUO *et al.* 2004)
Sulfur-limited chemostat	26.1	2.8	1.5	(BRAUER *et al.* 2008)
Glucose-limited chemostat		2.8	1.8	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.8	1.8	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.8	1.8	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.8	1.9	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.8	2.1	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		2.8	2.2	(HENRY *et al.* 2010)
Glucose-limited chemostat		2.9	0.7	(CARTER AND JAGADISH 1978)
YP+Dextrose		3	0.8	(CARTER AND JAGADISH 1978)
YP+Mannitol	20	3.2	1.0	(TYSON *et al.* 1979)
YP+Trehalose	17.8	3.2	1.0	(TYSON *et al.* 1979)
MM3+Acetate	21.6	3.2	1.1	(TYSON *et al.* 1979)
Glucose-limited chemostat		3.2	1.2	(CARTER AND JAGADISH 1978)
MM+Dextrose+Proline		3.3	1.4	(JAGADISH AND CARTER 1977)
MM+Dextrose+Ammonia+CHX		3.3	1.6	(HARTWELL AND UNGER 1977)
SEY		3.3	1.7	(SLATER *et al.* 1977)
MM+Dextrose+Methionine		3.3	1.9	(RIVIN AND FANGMAN 1980)
Glucose-limited chemostat		3.5	1.1	(JAGADISH AND CARTER 1977)
Nitrogen-limited chemostat	29.3	3.5	1.3	(GUO *et al.* 2004)
Leucine-limited chemostat	35.1	3.5	1.5	(BRAUER *et al.* 2008)
Uracil-limited chemostat	41	3.5	1.5	(BRAUER *et al.* 2008)
Phosphorous-limited chemostat	32.1	3.5	1.7	(BRAUER *et al.* 2008)
Glucose-limited chemostat	17.9	3.5	1.8	(GUO *et al.* 2004)
Glucose-limited chemostat	32.1	3.5	1.9	(BRAUER *et al.* 2008)
Sulfur-limited chemostat	26.9	3.5	2	(BRAUER *et al.* 2008)
Nitrogen-limited chemostat	24	3.5	2.1	(BRAUER *et al.* 2008)
Nitrogen-limited chemostat		3.5	2.1	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		3.5	2.3	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		3.5	2.5	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		3.5	2.8	(HENRY *et al.* 2010)
Glucose-limited chemostat		3.5	2.9	(HENRY *et al.* 2010)
Nitrogen-limited chemostat		3.5	2.9	(HENRY *et al.* 2010)
MM+Dextrose+Proline		3.8	1.6	(CARTER AND JAGADISH 1978)
Glucose-limited chemostat		3.9	1.4	(JAGADISH AND CARTER 1977)
MM+Dextrose+Leucine		4	1.8	(JOHNSTON *et al.* 1980)
Nitrogen-limited chemostat		4	3.1	(HENRY *et al.* 2010)
Glucose-limited chemostat		4.2	1.8	(CARTER AND JAGADISH 1978)
Glucose-limited chemostat		4.3	2.1	(CARTER AND JAGADISH 1978)
SE		4.3	2.4	(SLATER *et al.* 1977)
MM+Dextrose+Threonine		4.3	2.6	(RIVIN AND FANGMAN 1980)
EMM+Galactose	22.9	4.5	2.2	(TYSON *et al.* 1979)

Uracil-limited chemostat	41.8	4.6	2	(BRAUER et al. 2008)
Nitrogen-limited chemostat	29.6	4.6	2	(GUO et al. 2004)
Phosphorous-limited chemostat	31.3	4.6	2.3	(BRAUER et al. 2008)
Glucose-limited chemostat		4.6	2.3	(JAGADISH AND CARTER 1977)
Glucose-limited chemostat	29.9	4.6	2.6	(BRAUER et al. 2008)
Glucose-limited chemostat	19	4.6	2.6	(GUO et al. 2004)
Leucine-limited chemostat	38.1	4.6	2.9	(BRAUER et al. 2008)
Nitrogen-limited chemostat		4.6	3.1	(HENRY et al. 2010)
Sulfur-limited chemostat	27.6	4.6	3.2	(BRAUER et al. 2008)
Nitrogen-limited chemostat	23.9	4.6	3.5	(BRAUER et al. 2008)
Nitrogen-limited chemostat		4.6	3.5	(HENRY et al. 2010)
Nitrogen-limited chemostat		4.6	3.6	(HENRY et al. 2010)
Nitrogen-limited chemostat		4.6	3.8	(HENRY et al. 2010)
Nitrogen-limited chemostat		4.6	3.9	(HENRY et al. 2010)
Glucose-limited chemostat		4.6	4	(HENRY et al. 2010)
MM1+Glycerol	16.9	4.9	3.1	(TYSON et al. 1979)
YP+Glycerol		5.2	3.0	(CARTER AND JAGADISH 1978)
YP+Glycerol		5.2	3.2	(JAGADISH AND CARTER 1977)
MM+Dextrose+Ammonia+CHX		5.4	3.6	(HARTWELL AND UNGER 1977)
MM+Dextrose+Tryptophan		5.6	3.8	(JOHNSTON et al. 1980)
Glucose-limited chemostat		6.2	2.6	(JAGADISH AND CARTER 1977)
Glucose-limited chemostat		6.2	3.9	(JAGADISH AND CARTER 1977)
MM4+Acetate	19.5	6.4	3.4	(TYSON et al. 1979)
MM+Dextrose+Proline		6.4	4.5	(JOHNSTON et al. 1980)
Glucose-limited chemostat		6.6	4.5	(CARTER AND JAGADISH 1978)
MM+Dextrose+Proline		6.7	4.7	(RIVIN AND FANGMAN 1980)
Nitrogen-limited chemostat	33.2	6.9	3	(GUO et al. 2004)
Glucose-limited chemostat	18.9	6.9	4.3	(GUO et al. 2004)
Glucose-limited chemostat	36.6	6.9	4.4	(BRAUER et al. 2008)
Nitrogen-limited chemostat		6.9	4.8	(HENRY et al. 2010)
Uracil-limited chemostat	47.8	6.9	5	(BRAUER et al. 2008)
Phosphorous-limited chemostat	40.3	6.9	5.1	(BRAUER et al. 2008)
Nitrogen-limited chemostat		6.9	5.1	(HENRY et al. 2010)
Leucine-limited chemostat	38.1	6.9	5.3	(BRAUER et al. 2008)
Sulfur-limited chemostat	27.6	6.9	5.3	(BRAUER et al. 2008)
Nitrogen-limited chemostat	29.9	6.9	5.5	(BRAUER et al. 2008)
Nitrogen-limited chemostat		6.9	5.8	(HENRY et al. 2010)
Nitrogen-limited chemostat		6.9	6.1	(HENRY et al. 2010)
Glucose-limited chemostat		6.9	6.6	(HENRY et al. 2010)
Nitrogen-limited chemostat		6.9	6.6	(HENRY et al. 2010)
Glucose-limited chemostat		7	4.6	(CARTER AND JAGADISH 1978)
MM3+Citrate	16.9	7.3	3.6	(TYSON et al. 1979)
MM+Ethanol+Ammonia		7.5	4.7	(CARTER AND JAGADISH 1978)
EMM+Acetate+Pthalate	18.9	7.6	4.2	(TYSON et al. 1979)
MM+Dextrose+Threonine		7.6	4.6	(JOHNSTON et al. 1980)
MM+Dextrose+Ammonia+CHX		8.2	6.5	(HARTWELL AND UNGER 1977)

MM+Ethanol+Ammonia		8.3	5.9	(JAGADISH AND CARTER 1977)
MM+Acetate		8.5	6.1	(JAGADISH AND CARTER 1977)
Glucose-limited chemostat		8.6	6.5	(CARTER AND JAGADISH 1978)
Glucose-limited chemostat		9.1	5.9	(JAGADISH AND CARTER 1977)
Glucose-limited chemostat		11.7	8.4	(CARTER AND JAGADISH 1978)
MM+Dextrose+Ammonia+CHX		11.9	10.6	(HARTWELL AND UNGER 1977)
Glucose-limited chemostat	42.5	13.9	11.3	(BRAUER *et al.* 2008)
Leucine-limited chemostat	43.3	13.9	11.5	(BRAUER *et al.* 2008)
Phosphorous-limited chemostat	40.4	13.9	11.7	(BRAUER *et al.* 2008)
Sulfur-limited chemostat	31.3	13.9	11.9	(BRAUER *et al.* 2008)
Nitrogen-limited chemostat	32.8	13.9	12	(BRAUER *et al.* 2008)
Uracil-limited chemostat	47.8	13.9	12	(BRAUER *et al.* 2008)
Glucose-limited chemostat		17.9	12.5	(CARTER AND JAGADISH 1978)
Glucose-limited chemostat	34.3	21.6	15.6	(JAGADISH AND CARTER 1977)

590   **Table S2. Human cell cycle kinetics**

Cell type	T_d (h)	T_{G1} (h)	Reference
acute myeloblastic leukemia cells	49	24	(BASERGA 1985)
amnion cells	18.2	7.3	(SISKEN AND KINOSITA 1961)
amnion cells	19.5	9.5	(BASERGA 1985)
amnion cells	21.4	10.3	(SISKEN AND KINOSITA 1961)
ascites carcinoma cells	113	50	(BASERGA 1985)
basal cell carcinoma cells	67	36	(BASERGA 1985)
bone marrow cells	18	2	(BASERGA 1985)
BR:BT-549; NCI60 panel	53.9	30.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
BR:HS 578T; NCI60 panel	53.8	30.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
BR:MCF7; NCI60 panel	25.4	14.7	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
BR:MDA-MB-231; NCI60 panel	41.9	23.7	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
BR:T-47D; NCI60 panel	45.5	30.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:SF-268; NCI60 panel	33.1	17.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:SF-295; NCI60 panel	29.5	17.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:SF-539; NCI60 panel	35.4	23.59	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:SNB-19; NCI60 panel	34.6	21.3	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:SNB-75; NCI60 panel	62.8	40.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CNS:U251; NCI60 panel	23.8	9.8	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:COLO 205; NCI60 panel	23.8	14.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:HCC-2998; NCI60 panel	31.5	19.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:HCT-116; NCI60 panel	17.4	8.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:HCT-15; NCI60 panel	20.6	10.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:HT29; NCI60 panel	19.5	13	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:KM12; NCI60 panel	23.7	11.4	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
CO:SW-620; NCI60 panel	20.4	13.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study

colon epithelial cells	25	9	(BASERGA 1985)
epidermoid carcinoma cells	24	9	(BASERGA 1985)
Fibroblasts	18	5.5	(DEFENDI AND MANSON 1963)
HeLa	16.2	7.6	(KUMEI et al. 1989)
HeLa	19	12.1	(LUCIANI et al. 2001)
HeLa	19.4	11	(HAHN et al. 2009)
HeLa	20.4	14.9	(LUCIANI et al. 2001)
HeLa	21	8	(BASERGA 1985)
HeLa	25	13	(LENNARTZ AND MAURER 1964)
HeLa	26.6	19.7	(LUCIANI et al. 2001)
HeLa	27.6	21.8	(LUCIANI et al. 2001)
HeLa	28	13.5	(DEFENDI AND MANSON 1963)
HeLa	28	14	(LENNARTZ AND MAURER 1964)
high grade non-Hodgkins lymphoma cells	57	36	(BRONS et al. 1992)
Hodgkins lymphoma cells	62	37	(BRONS et al. 1992)
intermediate grade non-Hodgkins lymphoma cells	69	47	(BRONS et al. 1992)
KB	31.5	6.5	(BASERGA 1985)
Ki-67 reactive hyperplasia lymphoma cells	22	14	(BRONS et al. 1992)
LC:A549/ATCC; NCI60 panel	22.9	13.8	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:EKVX; NCI60 panel	43.6	29.2	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:HOP-92; NCI60 panel	79.5	46. 2	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:NCI-H226; NCI60 panel	61	41.2	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:NCI-H23; NCI60 panel	33.4	15.4	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:NCI-H322M; NCI60 panel	35.3	22.8	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:NCI-H460; NCI60 panel	17.8	12.6	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LC:NCI-H522; NCI60 panel	38.2	25.3	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LE:CCRF-CEM; NCI60 panel	26.7	12.1	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LE:HL-60(TB) ; NCI60 panel	28.6	16.2	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LE:K-562; NCI60 panel	19.6	10.1	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LE:MOLT-4; NCI60 panel	27.9	15.9	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study
LE:RPMI-8226; NCI60 panel	33.5	12.4	(SCHERF et al. 2000; GARNER AND EASTMAN 2011), and this study

LE:SR; NCI60 panel	28.7	14.0	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
low grade non-Hodgkins lymphoma cells	107	80	(BRONS *et al.* 1992)
Lymphocytes	30.4	14.2	(AOKI AND MOORE 1970)
Lymphocytes	33.1	15.1	(AOKI AND MOORE 1970)
Lymphocytes	38.4	21.2	(AOKI AND MOORE 1970)
Lymphocytes	43.5	25.9	(AOKI AND MOORE 1970)
Lymphocytes	72.5	49.4	(AOKI AND MOORE 1970)
Lymphocytes	76.5	55	(AOKI AND MOORE 1970)
MCF-7 breast cancer cells	24	14.4	(BRUNNER *et al.* 1989)
ME:LOX IMVI; NCI60 panel	20.5	11.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:M14; NCI60 panel	26.3	15.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:MALME-3M; NCI60 panel	46.2	30.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:MDA-MB-435; NCI60 panel	25.8	10.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:SK-MEL-2; NCI60 panel	45.5	25	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:SK-MEL-28; NCI60 panel	35.1	24.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:SK-MEL-5; NCI60 panel	25.2	17.7	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:UACC-257; NCI60 panel	38.5	28.3	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
ME:UACC-62; NCI60 panel	31.3	21.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
melanoma cells	46	20	(BASERGA 1985)
NCI-H292	18.5	6.8	(HAHN *et al.* 2009)
OV:IGROV1; NCI60 panel	31	21.6	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:NCI/ADR-RES; NCI60 panel	34	23.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:OVCAR-3; NCI60 panel	34.7	19.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:OVCAR-4; NCI60 panel	41.4	19.0	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:OVCAR-5; NCI60 panel	48.8	28.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:OVCAR-8; NCI60 panel	26.1	13.3	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
OV:SK-OV-3; NCI60 panel	48.7	39.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
PR:DU-145; NCI60 panel	32.3	18.2	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study

PR:PC-3; NCI60 panel	27.1	15.4	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:786-0; NCI60 panel	22.4	13.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:A498; NCI60 panel	66.8	41.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:ACHN; NCI60 panel	27.5	14.9	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:CAKI-1; NCI60 panel	39	25.0	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:RXF 393; NCI60 panel	62.9	31.1	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:SN12C; NCI60 panel	29.5	15.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:TK-10; NCI60 panel	51.3	29.4	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
RE:UO-31; NCI60 panel	41.7	26.5	(SCHERF *et al.* 2000; GARNER AND EASTMAN 2011), and this study
rectum epithelial cells	48	33	(BASERGA 1985)
stomach epithelial cells	24	9	(BASERGA 1985)
H. sapiens diploid fibroblasts	18	6.6	(BASERGA 1985)

An additional file (**Supplementary File 1**) is attached and contains all raw immunoblot images generated

and used in this study.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE S1

FIGURE S2

Western blot controls for detection of Cln3p-13Myc and Whi5p-TAP.

A. Western blot of proteins extracted from the strains indicated. The blot was first probed with the PAB antibody to show the detection of Whi5p-TAP in the appropriate lanes at the right size. B. Same blot as in A but probed additionally with the anti-Myc antibody followed by anti-rabbit HRP antibody for detection. This method allowed us to detect both Cln3p-13Myc and Whi5p-TAP on the same blot using the same exposure. C and D. Additional earlier control western blot performed with an earlier lot of the anti-Myc antibody. C. Detection of Whi5p-TAP probed with the PAP antibody. D. Different blot of the same samples run in C, probed with the anti-Myc antibody. Note the presence of dark non-specific bands detected with this lot of the anti-Myc antibody.

Western blots for quantification of Cln3p and Whi5p at different doubling times (T_D) from chemostat experiments.

Cell samples collected and frozen at the indicated doubling times, controlled by media in/out flow rate.

Chemostat

Proteins extracted and run on gel in duplicate, followed by transfer to a membrane and staining with Ponceau S (column 1).

Blot probed first with PAP antibody (not shown) and then with anti-Myc antibody and anti-Rabbit HRP secondary antibody for detection and quantification of Whi5p-TAP and Cln3p-(Myc)$_{13}$ (columns 3 and 4).

Column 1 Ponceau S (total protein)	**Column 2** Marker overlay (for size only)	**Column 3** Raw signal	**Column 4** Signal (cropped with background subtract for quantification)

Glucose limiting, Chemostat 1, Set 1 (Strain HB94)

Glucose limiting, Chemostat 1, Set 2 (Strain HB94)

Glucose limiting, Chemostat 2, Set 1 (Strain HB97)

Glucose limiting, Chemostat 2, Set 2 (Strain HB97)

Leucine limiting (Strain HB97)

Glucose limiting (Strain HB104-A315T) set1 wb2

Glucose limiting (Strain HB104-A315T set2 wb1

Leucine limiting (Strain HB104-A315T set1 wb1

* indicates a non-specific band that was detected with an earlier lot of the anti-MYC antibody.