1

Murine polyomavirus microRNAs promote viuria during the acute phase of infection

James M. Burke[1], Clovis R. Bass[1], Emin T. Ulug[1], Christopher S. Sullivan[1,*]

*Correspondence:

Chris_sullivan@austin.utexas.edu

[1]The University of Texas at Austin, Center for Systems and Synthetic Biology, Center for

Infectious Disease and Dept. Molecular Biosciences, 1 University Station A5000, Austin TX

78712-0162

Running title: Polyomavirus miRNAs promote acute viuria

Keywords: Polyomavirus, miRNA, viral shedding, viral persistence

Abstract word count: 183

Manuscript word count: 4,103

Abstract

Polyomaviruses (PyVs) can cause serious disease in immunosuppressed hosts. Several

pathogenic PyVs encode microRNAs (miRNAs), small RNAs that regulate gene expression

via RNA silencing. Despite recent advances in understanding the activities of PyV miRNAs,

the biological functions of PyV miRNAs during *in vivo* infections are mostly unknown.

Studies presented here use murine polyomavirus (MuPyV) as a model to assess the roles of

the PyV miRNAs in a natural host. This analysis reveals that a MuPyV mutant that is unable

to express miRNAs has enhanced viral DNA loads in select tissues at late times after

infection, indicating that during infection of a natural host, PyV miRNAs function to reduce

viral replication during the persistent phase of infection. Additionally, MuPyV miRNAs

promote viuria during the acute phase of infection as evidenced by a defect in shedding

during infection with the miRNA mutant virus. The viuria defect of the miRNA mutant virus

could be rescued by infecting Rag2-/- mice. These findings implicate miRNA activity in both

the persistent and acute phases of infection and suggest a role for MuPyV miRNA in evading

the adaptive immune response.

Importance

MicroRNAs are expressed by diverse viruses, but for only a few is there any understanding

of their *in vivo* function. PyVs can cause serious disease in immunocompromised hosts.

Therefore, increased knowledge of how these viruses interact with the immune response is

of possible clinical relevance. Here we show a novel activity for a viral miRNA in promoting

virus shedding. This work indicates that in addition to any role for the PyV miRNA in long-

term persistence, that it also has biological activity during the acute phase. As this mutant

phenotype is alleviated by infection of mice lacking an effective adaptive immune response,

our work also connects the *in vivo* activity of a PyV miRNA to the immune response. Given

that PyV-associated disease is associated with alterations in the immune response, our

findings may help to better understand how the balance between PyV and the immune

response becomes altered in pathogenic states.

Introduction

The polyomavirus (PyV) family is comprised of a large number of viruses that predominantly infect vertebrates (1). The founding member of the PyVs, murine polyomavirus (MuPyV), has been the most tractable *in vivo* model for studying the PyV life cycle and has played an important role in understanding PyV-mediated transformation and the antiviral response (2, 3, 4). MuPyV is thought to be transmitted to newborn mice via a respiratory route (5, 6, 7). Primary replication, which occurs 1-6 days post-infection in nonciliated epithelial clara cells of the lungs (7), is followed by dissemination and replication in secondary tissues 7-12 days post infection (6, 8). Viral replication has been observed in various secondary organs, including kidneys, salivary glands, spleen, lymph nodes, heart, liver, skin, lungs, and mammary glands (8, 9, 10). Virus replication peaks around 12 days after infection, concurrent with a rise in viral shedding in the urine and saliva (6). A rise in anti-MuPyV antibody titers occurs 7-15 days p.i. and precedes clearance of detectable virus in most organs by 22-30 days (8).

A reduction in viral loads marks the transition from the acute phase to the persistent phase, which is generally defined by the continued presence of viral DNA in tissues (3). Viral DNA is maintained in select tissues, such as bone marrow, spleen, kidney, salivary glands, and mammary glands (8, 9). The viral DNA in these tissues is thought to represent a persistent reservoir of 'smoldering' viral replication in semi-permissive cells (4), although a latent state has yet to be ruled out. A change in the microenvironment (e.g. tissue damage), immune status (e.g. pregnancy or immunosuppression), or hormone levels (e.g. pregnancy, stress), can 'reactivate' or increase replication of MuPyV from persistent reservoirs (11, 12). This results in episodic shedding of

MuPyV in urine and saliva, which contaminates the surrounding environment with infectious virus

(6). Similar observations have been made for human PyVs, whereby increased viuria during

pregnancy coincides with a rise in anti-PyV neutralizing antibodies (13, 14, 15).

Many polyomaviruses – including MuPyV, RacPyV, SV40, JCPyV, MCPyV, and BKPyV –

encode microRNAs (miRNAs) (16, 17, 18, 19, 20), small regulatory RNAs (~22 nt) that repress

gene expression via RNA silencing (21). The PyV miRNAs are encoded in the late orientation,

opposite to the T antigen mRNAs. Consequently, miRNAs are perfectly complementary to the T

antigen transcripts, permitting Argonaute 2 (Ago2)-mediated cleavage of these mRNAs. This

results in a reduction in T antigen protein levels late in infection, a conserved function of PyV

miRNAs (16, 17, 18, 19, 20, 22). Despite established roles of T antigen in promoting viral

replication, miRNA-null laboratory strains of SV40 and MuPyV replicate at rates similar to their

wild type counterparts under standard cell culture conditions (16, 17). However, miRNAs

expressed by non-rearranged BKPyV strains, which more closely resemble circulating BKPyV in

humans and express lower levels of the early transcripts, are able to downregulate T antigen levels

to sufficiently reduce viral replication in primary renal proximal tubule epithelial cells (23).

Consistent with these results, MCPyV miRNA promotes long-term persistence by inhibiting DNA

replication in a cell culture model (24), and the SV40 miRNAs reduce persistent SV40 DNA loads

in Syrian Golden hamster tissues (25). These studies suggest that the PyV miRNAs reduce virus

replication by down-regulating T antigen expression. However, the effects of miRNA expression

on virus replication in a natural host have yet to be established.

In addition to regulating viral replication, independent studies implicate a role for the PyV

miRNAs in inhibiting the host immune response. Downregulation of T-antigen by the SV40

miRNAs decreases CD8 T-cell-mediated lysis in cell culture (16). miRNA-mediated

downregulation of the host ULBP3 stress-induced ligand has been suggested to reduce killing of

BKV- and JCV-infected cells by natural killer cells (26). MuPyV miRNAs reduce Smad2-

mediated apoptosis during infection (27). These studies support a model whereby the PyV

miRNAs modulate viral and host transcripts in order to extend the life of infected cells by reducing

innate and adaptive immune responses.

Despite advances in understanding the expression and activities of PyV miRNAs, their functions

during the viral lifecycle in a natural host are poorly understood. In this study, we use MuPyV as

a model to investigate the role of PyV miRNAs on viral dissemination, replication, and shedding

during the acute and persistent phases of infection in mice. This work reveals activities of the

MuPyV miRNA in both the persistent and acute phases of infection and suggests a link between

miRNA function and the adaptive immune response.

Results

Sequencing of the MuPyV miRNAs

MuPyV miRNAs were initially identified via secondary structure predictions and confirmed indirectly (17). This approach could only estimate the 5' and 3' termini of the derivative miRNAs. It also did not resolve whether MuPyV contains additional miRNA loci. To better characterize the MuPyV miRNAs, we conducted high throughput sequencing of small RNA from NIH3T3 cells infected with the MuPyV PTA strain (Figure 1). These results map with precision the termini of the MuPyV miRNAs and confirm that only the single, previously identified pre-miRNA locus, gives rise to MuPyV-encoded miRNA derivatives.

MuPyV miRNAs correlate with reduced viral DNA loads during the persistent phase of infection.

PTA-dll013 is a deletion mutant of the MuPyV-PTA strain that does not express miRNAs (17). Previous studies reported that adult C57BL/6 mice infected with PTA-dll013 maintained comparable or slightly higher (up to 5-fold) vDNA loads as compared to PTA in spleen and kidney during the acute phase of infection (up to 34 days p.i.) (17). To confirm these results and determine whether the miRNA function at later times post-infection, we examined MuPyV DNA loads by qPCR in spleens, kidneys, and livers of adult C57BL/6 mice infected with 10^5 IU of PTA or PTA-dl1013 via IP injection during the acute (1 and 4 weeks p.i.) and persistent (10, 16, and 28 weeks p.i.) phases of infection (Figure 2).

High virus yields (2×10^5 -5×10^6 genome equivalents/ug DNA) were observed in spleen and

kidney after one and four weeks of infection with PTA, consistent with robust virus replication

during the acute phase of replication (Figure 2). Lower levels were observed in liver during these

times (Figure 2). Virus loads diminished to $\leq 10^3$ genome equivalents/ug DNA at later times after

infection (16 and 28 weeks p.i.), consistent with a low level of virus replication during the

persistent phase of replication. These levels were only slightly above the limit of detection of our

PCR assay, even when increased input DNA was assayed.

In comparison to PTA, PTA-dl1013 levels were 4.1-fold lower in spleen, but not kidney, at one

week p.i. (Figure 2, closed symbols). No significant differences in PTA and PTA-dl1013 DNA

levels were detected at 4 weeks or 10 weeks after infection. However, at 16 weeks, PTA-dl1013

genome levels were approximately 10-fold higher than PTA in spleen and kidney. By 28 weeks,

only low levels of both PTA and PTA-dl1013 genome levels were detected and no significant

differences were observed. These results confirm that the miRNAs are not required for infection

and replication in tissues during the acute phase of infection, and indicate that miRNA expression

correlates with reduced genomic DNA in certain tissues during the persistent phase of infection.

The results summarized in Figure 2 reveal a substantial variation in viral copies between different

mice. To minimize variations due to inter-host differences, we compared levels of PTA and PTA-

dl1013 in individual animals. C57BL/6 mice were co-infected with equivalent amounts of PTA

and PTA-dl1013 by IP injection. Tissues were then harvested during the acute (2 and 4 weeks) and

persistent (10 and 16 weeks) phases of infection, and genome levels were determined by strain-

specific qPCR (Supplemental Figure 1A-C). Similar to individually infected mice, we observed

high levels of both PTA and PTA-dl1013 in tissues during the acute phase of infection, with

MuPyV levels decreasing during the persistent phase of infection (Figure 3). We note that PTA

was not detected in the spleen or kidney by 16 weeks p.i., while PTA-dl1013 was present at 10^3 -

10^4 copies/ug DNA (Figure 3). To compensate for variation in different animals, we calculated the

ratio of PTA to PTA-dl1013 genomes in individual mice. This analysis reveals that during the

acute phase of infection (2 and 4 weeks p.i.), the levels of PTA and PTA-dl1013 were comparable

in spleen and kidney (Figure 4). At later times, PTA-dl1013 copy numbers were 10-fold higher

than PTA in the spleen and kidney at weeks 10 and 16 and 100-fold higher in the kidney at week

16 (Figure 4). We did not observe substantial differences in the liver, but did observe 7-fold higher

PTA-dl1013 in the bladder at week 4 (Figure 4). These results are consistent with the elevated

levels of PTA-dl1013 detected in the spleen and kidney at 16 weeks following infections with the

individual virus strains (Figure 2) and suggest that MuPyV miRNA expression correlates with

reduced viral loads in select tissues during the persistent phase of infection.

MuPyV miRNAs promote viuria during the acute phase of infection.

Early epidemiological studies of MuPyV revealed that high titers of infectious virus are excreted

in the urine of mice during the acute phase of infection and then episodically shed during the

persistent phase of infection (6). The periodic shedding of MuPyV and many other PyVs in the

urine is considered a primary route of transmission (6,28). Therefore, we investigated whether the

MuPyV miRNAs affects the magnitude and/or frequency of MuPyV shedding in the urine during

the acute and persistent phases of infection. To test this, we collected urine specimens periodically

post-infection and quantitated viral genomes. To assess whether PTA-dl1013 could maintain long-

term persistence, we treated mice with cyclosporine at day 474 post-infection in order to

immunosuppress the mice, which is correlated with increased PyV replication (4).

In the majority of mice individually infected with PTA or PTA-dl1013, we observed vDNA

shedding of both strains during the acute phase of infection (13-34 days p.i.) (Supplemental Figure

2), indicating that the MuPyV miRNAs are not required for establishing viuria. Both strains were

episodically shed throughout the persistent phase (34-87 day p.i.), which demonstrates that the

MuPyV miRNAs are not required for persistent viuria (Supplemental Figure 2). The magnitude

and timing of virus shedding by individual mice infected varied substantially (Supplemental Figure

2), and thus no statistical differences between PTA and PTA-dl1013 levels could be determined.

To circumvent the differences in shedding due to variations between individual mice, we tested

whether expression of miRNA could correlate with viral shedding in mice co-infected with both

PTA and PTA-dl1013. Adult C57BL/6 female mice were co-infected with equal amounts of the

PTA and PTA-dl1013, and PTA and PTA-dl1013 DNA was quantitated during the acute and

persistent phases of infection (Figure 5A). We did not detect substantial quantities of either PTA

or PTA-dl1013 genomes in the urine of ~40% of the mice at any time point during the acute phase

of infection (Supplemental file). The remaining "high-shedder" mice expressed readily detectable

MuPyV DNA in the urine. MuPyV in these samples was detected at high levels beginning by 10

days p.i. and peaked between 13-27 days, after which the number of shedding mice and the

magnitude of shedding began to decrease. Thereafter, we observed episodic shedding events

throughout the persistent phase of infection (34-87 days p.i.). These findings reveal that while

some mice rarely or never shed detectable virus under our assay conditions, ~60% that are

competent for acute and persistent viuria.

The median PTA levels were significantly higher (5- to 100- fold) than PTA-dl1013 during the

majority of the acute phase of infection (10, 13. 15, 17, 20 and 27 days p.i.) in co-infected mice

(Figure 5A). To minimize the variance in individual mice, we determined the ratio of PTA:PTA-

dl1013 DNA in each individual mouse. This data confirmed that PTA was shed at significantly

higher levels on average (~10-fold) than PTA-dl1013 during days 10-17 of the acute phase of

infection in individual mice co-infected mice (Figure 5B). We did not observe statistically

significant differences in the magnitude of viral shedding at the majority of times tested during the

persistent phase of infection (Figure 5A, 5B, Supplemental file). Although this may be due to high

variance in the timing and the magnitude of shedding, as well as the lower number of mice tested,

this suggests that the MuPyV miRNAs do not promote viuria during the persistent phase of

infection. In fact, we observed eighty-four PTA-dl1013 shedding events that were above the limit

of detection from days 34 through 87, compared to thirty-three for PTA (Figure 5A). This implies

that the MuPyV miRNAs may limit detectable shedding during persistent phase of infection,

consistent with the reduced PTA levels in tissues (Figures 2-4). At late times post-infection (467-

492), we observed PTA and PTA-dl1013 shedding events in a small number of mice. Thus, the

MuPyV miRNAs are not required to maintain long-term persistent infections. These data further

indicate that the MuPyV miRNAs promote viral shedding during the acute phase of infection, but

limit the number of MuPyV shedding event during the persistent phase of infection.

The shedding defect of miRNA-null MuPyV is mitigated in Rag2-/- mice.

The host immune response plays an important role in controlling MuPyV infection (2-4), and two lines of evidence suggest that PyV miRNAs may function to reduce the adaptive immune response. First, miRNA-mediated down-regulation of T-antigens decreases susceptibility to cytotoxic T-cell-mediated lysis to SV40-infected cultured cells (16). Second, miRNA-null strains of SV40 and JCV have only been isolated from severely immunocompromised hosts, suggesting that an active immune response provides selective pressure to maintain miRNA expression (29). Since the absence of the MuPyV miRNA results in decreased virus shedding during the acute phase of infection, we tested whether the defect in acute viral shedding of PTA-dl1013 is mitigated in C57BL/6 *Rag2*-/- mice, which lack mature T- and B- cells. C57BL/6 *Rag2*-/- mice were co-infected with PTA and PTA-dl1013, and the shedding of DNA in urine was monitored. Unlike wild type mice, where ~40% of animals did not shed detectable vDNA during the acute phase of infection, all infected C57BL/6 *Rag2*-/- mice shed detectable levels of MuPyV by 10 days post-infection (Figure 6 and Supplemental file). Levels of both PTA and PTA-dl1013 DNA in urine were found to continually increase in the co-infected C57BL/6 *Rag2*-/- mice up to 20 days p.i., whereas MuPyV levels leveled off and began to decrease by 20 days p.i. (Figure 6). These observations are consistent with the host adaptive immune response being a major factor that controls MuPyV shedding. Importantly, unlike in wild type C57BL/6 mice, C57BL/6 *Rag2*-/- mice shed PTA-dl1013 DNA at levels similar to PTA during the acute phase of infection (10-13 days p.i.) (Figure 6). These data indicate that the MuPyV miRNAs promote viuria during the acute phase of infection in a manner that is dependent on the host having an intact adaptive immune response.

Discussion

The natural *in vivo* functions of PyV miRNAs are poorly understood. Results presented here reveal that, consistent with previous studies in cell culture models and a non-natural host (23-25), the PyV miRNAs function to reduce viral loads during the persistent infection of a natural host. In addition, we uncover novel effects of the MuPyV miRNAs on viuria during the persistent and acute phases of infection that provide new insights into PyV miRNA biology.

Similar to previous studies (17), we did not observe large differences in the gross levels of PTA (wild type) and PTA-dl1013 (miRNA-null) strains of MuPyV in tissues during acute phase of infection (weeks 1, 2 and 4 p.i.). This further confirms that the MuPyV miRNAs are not required for acute viral infection and replication *in vivo* under laboratory conditions. However, at weeks 10 and 16 p.i., we observed higher levels of PTA-dl1013 DNA, relative to PTA, in kidney and spleen of mice following single- or co-infection with PTA and PTA-dl1013 (Figures 2-4). Similar findings were observed in tissues of Syrian golden hamsters infected with miRNA-null strains of the simian virus SV40 (25). These results are consistent with the notion that miRNA-mediated downregulation of T antigen reduces virus replication in specific tissues/cell-types at late times after infection to promote viral persistence (23,24).

Our assessment of MuPyV viuria revealed that the MuPyV miRNAs are not required to establish or maintain long-term viral persistence (Figure 5 and Supplemental Figure 2). However, we did observe more PTA-dl1013 than PTA shedding events during the persistent phase of infection (34-87 days p.i.). This is consistent with increased PTA-dl1013 levels in tissues and further indicates

that the MuPyV miRNAs may limit viral replication events during the persistent phase of infection.

Though the function of this activity remains unclear, it may be important to prevent priming of the

host immune response in order to transmit to naïve hosts more efficiently. Alternatively, this

activity could prevent viral-associated pathologies to the host, which would presumably decrease

viral fitness. Future experiments will address the relevant role of miRNA-mediated reduction of

viral loads in urine and tissues during the persistent phase of infection.

During the acute phase of infection (days 10-20) in wild type C57BL/6 mice, only ~60% of mice

shed substantial levels of MuPyV DNA (Figure 5 and Supplemental Figure 2). This is in contrast

to infections in immunodeficient C57BL/6 *Rag2*-/- mice, in which 100% of mice shed substantial

levels of MuPyV DNA. This suggests that the adaptive immune response is a major factor

controlling viuria. Analysis of wild-type C57BL/6 mice co-infected with PTA and PTA-dl1013

revealed that PTA-dl1013 DNA was shed at significantly lower levels than PTA (Figure 5), which

indicates the MuPyV miRNAs promote viral shedding at early times post-infection. Importantly,

the defect in PTA-dl1013 was largely mitigated during infection of immunodeficient C57BL/6

Rag2-/- mice (Figure 6), suggesting that the MuPyV miRNAs help evade a component of the host

adaptive immune response to promote viuria. We note that PTA-dl1013 shedding was not fully

rescued by infecting C57BL/6 *Rag*2-/- mice, possibly indicating that the MuPyV miRNAs have

additional activities [for example, reduction of smad2-mediated apoptosis (27)] in promoting

viuria independent of the adaptive immune response. These combined data demonstrate that the

MuPyV miRNAs promote viral shedding by at least indirectly interacting with the adaptive

immune response. This activity could be important in the wild as a means to increase transmission

of the virus to susceptible hosts during the acute phase of infection.

The exact mechanism(s) by which MuPyV miRNA expression enhances virus shedding in immunocompetent mice remains unclear. Because we observed miRNA-dependent promotion of shedding in co-infected immunocompetent mice, this indicates that MuPyV miRNAs are likely functioning via a cell autonomous mechanism to repress the effects of the adaptive immune response. This is consistent with previous findings that the SV40 miRNAs can reduce CD8 T-cell-mediated lysis. However, it is unclear why the miRNAs promote viuria only during a narrow window of time post-infection. One possibility is that the MuPyV miRNAs increase dissemination to the target cells required for subsequent viuria, possibly by reducing the CTL response to promote infection/replication in lymphocytes early during the acute phase of infection. In support of this, at one week p.i., but not at 2 and 4 weeks p.i., we observed ~4-fold higher PTA DNA levels specifically in the spleen (Figures 2-4), indicating the MuPyV miRNAs may promote infection of lymphocytes early during infection. Alternatively, the cell-mediated adaptive immune response to MuPyV may change over time, making the miRNAs less effective at promoting viuria. While the CD8+ T cell response to MuPyV is predominately to T-antigen, the CD8+ T cell response has also been shown to be directed toward a VP2–derived nonamer peptide (30). Thus, as the infection progresses, miRNA-mediated down-regulate of T-antigens may become less effective at muting the CTL response as the host T-cell response develops more so against the late VP proteins. Future studies are required to more precisely define the relevant components of the immune response important for hindering the miRNA mutant virus.

In summary, this study demonstrates functionality of MuPyV miRNAs during both the persistent and acute phases of infection and at least indirectly implicates their function in altering some

317   component of the adaptive immune response. Given the clinical relevance of understanding the

318   interplay between polyomaviruses and the immune response, this work provides a useful,

319   quantitative, and non-invasive system for future efforts at understanding PyV miRNA effects on

320   adaptive immunity.

Materials and Methods

Cell cultures and virus strains

Primary baby mouse kidney (BMK) cell cultures were prepared from 2-4 week old female C57BL/6 mice (Jackson Laboratories). Kidneys were excised, washed in PBS, and incubated in DMEM containing 0.25% trypsin overnight at 4°C. Kidneys were homogenized, incubated in trypsin at 37°C with agitation, and suspended in DMEM+10% FBS (Cellgro). Cells were washed by centrifugation and plated at a density of 10^7 cells per 100 mm dish. NIH3T3 cells were maintained in DMEM+10% FBS.

The PTA and PTA-dl1013 virus strains have been described previously (17). Virus was propagated by infecting primary BMK cells (80% confluent) with PTA or PTA-dl1013 at an MOI of 0.05. Virus stocks were prepared after 10 days by scraping the cells into a portion of the culture medium. The cell extracts were then subjected to three freeze/thaw cycles, clarified by centrifugation, and stored at -80°C.

Virus titers were determined by immunofluorescence. 80-90% confluent NIH3T3 cells in 12- or 24-well plates were infected with dilutions of the PTA and PTA-dl1013 virus stocks for one hour. Infected cells were maintained in DMEM+2% FBS for 40 hours and then fixed and permeabilized with 4% paraformaldehyde and 1% NP-40. After blocking with 10% goat serum overnight, cells were incubated with rabbit-anti-PyV antibody (a gift from Dr. Richard Consigi) for 30 min. After three PBS washes, cells were stained with Cy3-goat-anti-rabbit IgG (Abcam) for 10 min. and

344 washed. The numbers of total and stained cells per field were quantified by using an inverted

345 fluorescent microscope (Leica), and infectious titers (IU/ml) were determined.

Animal infections and tissue analysis

Female C57BL/6 (Jackson Laboratories) and C57BL/6 *Rag2*-/- mice were inoculated with 1 x 10^5

IU of either PTA or PTA-dl1013 via intraperitoneal (IP) injection between 4 and 5 weeks of age.

For co-infections, mice were inoculated (IP) with 5 x 10^4 IU of PTA and 5 x 10^4 IU of PTA-

dl1013. Cylcospoin was administered as described in (31). Tissues were harvested at the indicated

times and flash frozen in liquid nitrogen. Homogenates were prepared by grinding with a mortar

and pestle chilled with liquid nitrogen. DNA was extracted from 25 mg of tissue by using the

DNeasy Blood and Tissue kit (Qiagen). Recovered DNA was quantitated by the Nanodrop

procedure and diluted to 100 ng/ul or 10 ng/ul for qPCR analysis.

Urine analysis

358 To collect urine, mice were placed on wire-bottomed cages for 2-4 hours. Urine was collected on

359 Parafilm, and 50 ul aliquots (or the total if less than 50 ul) were used for DNA extraction with the

360 QIAamp viral RNA mini kit (Qiagen) following the manufacturer's protocol to remove PCR

361 inhibitors. DNA was eluted in 60 ul of TE buffer. MuPyV copy numbers were normalized to that

362 in 2 ul of urine.

Quantitative PCR

Concentrated pBluescript-sk+PTA and pBluescript-sk+PTA-dl1013 vectors were obtained via

Maxi Prep (Invitrogen). Plasmids were prepared at 85.5 ng/ul (10^{10} copies of PTA or PTA-dl1013

per ul) and then serial diluted. For tissue analysis from individually infected mice, 20 ng of DNA

(for samples taken 1, 4, 10 weeks p.i.) or 200 ng of DNA (10 and 28 weeks p.i.) in 2 ul of TE

buffer was added to an 8 ul reaction mixture containing 5 ul of 2X gene expression master mix

(Applied Biosystems), 0.75 uM of MuPyV sense primer (CGCACATACTGCTGGAAGAAGA),

1.0 uM MuPyV antisense primer (TCTTGGTCGCTTTCTGGATACAG), and 100 nM of MuPyV

TaqMan MGB (Applied Biosystems) probe (FAM-ATCCTTGTGTTGCTGAGCCCGATGA-

NFQ) as described (32). For specific detection of PTA and PTA-dl1013 in urine, 2 ul of DNA (of

the 60 ul extracted unless otherwise indicated in Supplemental file) was added to an 8 ul reaction

mixture containing 5 ul 2X gene expression master mix (Applied Biosystems), 0.75 uM of

PTA/PTA-dl1013 sense primer (GATGAGCTGGGGTACTTGT), 0.75 uM of PTA/PTA-dl1013

antisense primer (TGTATCCAGAAAGCGACCAAG), and 100 nM of either the PTA-specific

TaqMan MGB (Applied Biosystems) probe (FAM-TAGGATGTCCAAATACAGATCCTC-

NFQ) or PTA-dl1013-specific TaqMan MGB (Applied Biosystems) probe (FAM-

CTCCGGTTCCATTGGCATGT-NFQ). The PTA and PTAdl1013 standards (10^7/ul -10^1/ul) were

confirmed by using the universal MuPyV primers/probe, which detects both PTA and PTA-

dl1013, to ensure that the copy number of each standard was equal. Assays were performed using

384-well format plates in a ViiA™ 7 Real-Time PCR System (Applied Biosystems). The limit of

detection was 10 copies.

Statistical analyses

One sample T-test (two-tailed) and Mann-Whitney U-test (two-tailed) were performed using real

statistics resource package for excel (http://www.real-statistics.com/).

References

1. Johne R, Buck CB, Allander T, Atwood WJ, Garcea RL, Imperiale MJ, Major EO, Ramqvist

T, Norkin LC. 2011. Taxonomical developments in the family Polyomaviridae. Arch Virol

156: 1627–1634.

2. Benjamin TL. 2001. Polyoma virus: old findings and new challenges. Virology 289: 167-173.

3. Ramqvist T, Dalianis T. 2009. Murine polyomavirus tumour specific transplantation antigens

and viral persistence in relation to the immune response, and tumour development. Semin

Cancer Biol 19: 236–243.

4. Swanson PA, Lukacher AE, Szomolanyi-Tsuda E. 2009. Immunity to polyomavirus infection:

The polyomavirus–mouse model. Semin Cancer Biol 19: 244-251.

5. Rowe WP, Hartley JW, Brodsky I, Huebner RJ, Law LW. 1958. Observations on the spread

of mouse polyoma virus infection. Nature 182:1617-16179.

6. Rowe W P. 1961. The epidemiology of mouse polyoma virus infection. Bacteriol Rev 25:18-

31.

7. Gottlieb K, Villarreal LP. 2000. The Distribution and Kinetics of Polyomavirus in Lungs of

Intranasally Infected Newborn Mice. Virology 266: 52-65.

8. Dubensky TW, Villarreal LP. 1984. The primary site of replication alters the eventual site of

persistent infection by polyomavirus in mice. J Virol 50: 541-546.

9. Wirth JJ, Amalfitano A, Gross R, Oldstone MB, Fluck MM. 1992. Organ- and age-specific

replication of polyomavirus in mice. J Virol 66:3278-86.

10. Berke Z, Dalianis T. 1993. Persistence of polyomavirus in mice infected as adults differs from

that observed in mice infected as newborns. J Virol 67: 4369-4371.

11. Atencio IA, Shadan FF, Zhou XJ, Vaziri ND, Villarreal LP. 1993. Adult mouse kidneys

become permissive to acute polyomavirus infection and reactivate persistent infections in

response to cellular damage and regeneration. J Virol 67: 1424–1432.

12. McCance DJ, Mims CA. 1979. Reactivation of polyoma virus in kidneys of persistently

infected mice during pregnancy. Infect Immun 25: 998–1002.

13. Coleman DV, Gardner SD, Mulholland C, Fridiksdottir V, Porter AA, Lilford R, Valdimarsson

H. 1983. Human polyomavirus in pregnancy. A model for the study of defense mechanisms to

virus reactivation. Clin Exp Immunol 53: 289-296.

14. Eash S, Manley K, Gasparovic M, Querbes W, Atwood WJ. 2006. The human polyomaviruses.

Cell Mol Life Sci 63:865-876.

15. Gibson PE, Field AM, Gardner SD, Coleman DV. 1981. Occurrence of IgM antibodies against

BK and JC polyomaviruses during pregnancy. J Clin Pathol 34: 674-679.

16. Sullivan CS, Grundhoff AT, Tevethia S, Pipas JM, Ganem D. 2005. SV40-encoded

microRNAs regulate viral gene expression and reduce susceptibility to cytotoxic T cells.

Nature 435: 682-686.

17. Sullivan CS, Sung CK, Pack CD, Grundhoff AT, Lukacher AE, Benjamin T, Ganem D. 2009.

Murine polyomavirus encodes a microRNA that cleaves early RNA transcripts but is not

essential for experimental infection. Virology 387: 157-167.

18. Seo GJ, Chen CJ, Sullivan CS. 2009. Merkel cell polyomavirus encodes a microRNA with the

ability to autoregulate viral gene expression. Virology 383: 183-187.

19. Seo GJ, Fink LH, O'Hara B, Atwood WJ, Sullivan CS. 2008. Evolutionarily conserved

function of a viral microRNA. J Virol 82: 9823-9828.

20. Chen CJ, Cox JE, Azarm K, Wylie KN, Woolard KD, Pesavento PA, Sullivan CS. 2015.

Identification of a Polyomavirus microRNA Highly Expressed in Tumors. Virology 476: 43–

53.

21. Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell 116: 281-

297.

22. Chen CJ, Cox JE, Kincaid RP, Martinez A, Sullivan CS. 2013. Divergent MicroRNA

Targetomes of Closely Related Circulating Strains of a Polyomavirus. J Virol 87:11135-11147.

23. Broekema NM, Imperiale MJ. 2013. miRNA regulation of BK polyomavirus replication during

early infection. Proc Natl Acad Sci U S A 110: 8200-8205.

24. Theiss JM, Günther T, Alawi M, Neumann F, Tessmer U, Fischer N, Grundhoff A. 2015. A

Comprehensive Analysis of Replicating Merkel Cell Polyomavirus Genomes Delineates the

Viral Transcription Program and Suggests a Role for mcv-miR-M1 in Episomal Persistence.

PLoS Pathog 11:e1004974.

25. Zhang S, Sroller V, Zanwar P, Chen CJ, Halvorson SJ, Ajami NJ, Hecksel CW, Swain JL,

Wong C, Sullivan CS, Butel JS. 2014. Viral MicroRNA Effects on Pathogenesis of

Polyomavirus SV40 Infections in Syrian Golden Hamsters. PLoS Pathog 10: e1003912.

26. Bauman Y, Nachmani D, Vitenshtein A, Tsukerman P, Drayman N, Stern-Ginossar N, Lankry

D, Gruda R, Mandelboim O. 2011. An identical miRNA of the human JC and BK polyoma

viruses targets the stress-induced ligand ULBP3 to escape immune elimination. Cell Host

Microbe 9:93-102.

27. Sung SK, Yim H, Andrews E, Benjamin TL. 2014. A Mouse Polyomavirus-encoded

microRNA Targets the Cellular Apoptosis Pathway through Smad2 Inhibition. Virology 468-

470: 57-62.

28. Gosert R, Rinaldo CH, Funk GA, Egli A, Ramos E, Drachenberg CB, Hirsch HH. 2008.

Polyomavirus BK with rearranged noncoding control region emerge in vivo in renal transplant

patients and increase viral replication and cytopathology. J Exp Med 205:841–852.

29. Chen CJ, Burke JM, Kincaid RP, Azarm KD, Mireles N, Butel JS, Sullivan CS. 2014. Naturally

Arising Strains of Polyomaviruses with Severely Attenuated MicroRNA Expression. J Virol

88:12683-12693.

30. Swanson PA, Pack CD, Hadley A, Wang C-R, Stroynowski I, Jensen PE, Lukacher AE. 2008.

An MHC class Ib–restricted CD8 T cell response confers antiviral immunity. J Exp Med

205:1647–1657.

31. Gattazzo F, Molon S, Morbidoni V, Braghetta P, Blaauw B, Urciuolo A, Bonaldo P. 2014.

Cyclosporin A Promotes in vivo Myogenic Response in Collagen VI-Deficient Myopathic

Mice. Front Aging Neurosci 244.

32. Kemball CC, Lee ED, Vezys V, Pearson TC, Larsen CP, Lukacher AE. 2005. Late priming

and variability of epitope-specific CD8+ T cell responses during a persistent virus infection. J

Immunol.174:7950–60.

Acknowledgements

The work described in this article was supported by a Burroughs Wellcome Investigators in

Pathogenesis Award to CSS and a grant from the Cancer Prevention and Research Institute of

Texas [RP140842]. The authors would like to thank Dr. Lauren Ehrlich and Dr. Edward

Marcotte for reagents, and Dr. Luis Villarreal for insights regarding viral persistence.

Figure Legends

Figure 1. Identification of MuPyV miRNAs by small RNA sequencing.

NIH3T3 cells were infected with MuPyV-PTA at an MOI of 0.1. A small RNA library was

prepared 7 days p.i. Dots on the graph represent the read counts of the 5' end of small RNA reads

mapping to the MuPyV-PTA genome (NCBI accession number: U27812), positioned on the x-

axis. The arrow below the graph represents the pre-miRNA hairpin position and direction. The

MuPyV-PTA reference sequence encoding the miRNAs is indicated. Sequences of the three most

abundant small RNAs mapping to the 5-prime (bold/red) and 3-prime (bold/blue) arms of pre-

miRNA hairpin are shown aligned to the pre-miRNA. Numbers represent the read counts.

Figure 2. Quantitation of PTA and PTA-dl1013 during the acute and persistent phases of

infection.

Adult C57BL/6 female mice were inoculated with $1\text{X}10^5$ IU of either PTA (white circles) or PTA-

dl1013 (gray circles) via IP injection, and genome levels in spleen and kidney were quantified at

times post infection by using a TaqMan probe and primers that recognize both genomes. The limit

of detection for the qPCR reactions was 10 copies. The input DNA amounts were 20 ng (at 1, 4,

and 10 weeks p.i.), 40 ng (at 16 weeks), or 200 ng (at 28 weeks). Genome copy numbers were

normalized per 1ug. Dots represent individual animals. The black bar represents the average

genome copy number. The gray bars represent the limit of detection (L.O.D,) of the assay after

normalization. Samples that were below the limit of detection are graphed at the limit of detection.

P-values were calculated using the Mann-Whitney U test.

Figure 3. Quantitation of PTA and PTA-dl1013 in tissues of co-infected mice.

qPCR analysis of PTA and PTA-dl1013 genome levels in the spleen, kidney, liver, and bladder of

adult C57BL/6 mice inoculated with 5×10^4 IUs of PTA and 5×10^4 IUs of PTA-dl1013 via IP

injection. Tissues were harvested at indicated time points. DNA was recovered and subject to

qPCR analysis using PTA- and PTA-dl1013- specific probes. The limit of detection (L.O.D.) was

10 copies. Samples in which PTA or PTA-dl1013 were below the limit of detection were graphed

at the limit of detection. The quantities were normalized to 1-ug of total DNA. P-values were

determined using Mann-Whitney U test.

Figure 4. Ratio of PTA and PTA-dl1013 in tissues of co-infected mice.

The ratio of PTA and PTA-dl1013 genome copies from (A) in the indicated organs of individual

mice co-infected with PTA and PTA-dl1013. Values below the limit of detection were set at the

limit of detection. The dots represent Log_{10}(PTA/PTA-dl1013). P-values were calculated using

one-sample t-test.

Figure 5. Quantitation of PTA and PTA-dl1013 DNA shedding in the urine.

Mice were co-infected with 5×10^4 IU of PTA and 5×10^4 IU of PTA-dl1013 via IP injection, urine

samples were collected at the indicated times, and strain-specific qPCR was used to quantify wt

and mutant genomes. The number of mice tested at each time ranged from 18-30 mice during days

1-27 p.i., 10-20 mice during days 29-55 p.i., and 4 mice during days 62-492 p.i. Panel A presents

a dot plot of genome equivalents of PTA (blue circles) and PTA-dl1013 (yellow triangles) in 2 ul

of urine from mice that shed viral genome copies equal to or greater that the limit of detection of

the qPCR assay. Mice that did not shed above the limit of detection or time points in which

genomes were not detected were not included in this analysis. Lines represent the median values

of PTA (solid blue) and PTA-dl1013) (dotted yellow). Significant differences (P ≤ 0.05) between

PTA and PTA-dl1013 genome levels are marked by asterisks. Panel B compares the ratio of wt

and mutant genomes in individual mice co-infected with PTA and PTA-dl1013. When only one

genome could be detected (either PTA or PTA-dl1013), the other genome was set at the limit of

detection in order to calculate the PTA/PTA-dl1013 ratio. The line represents the average

log10(PTA/PTA-dl1013) ± SEM. P values were calculated by using a one sample t-test (asterisks

indicate P ≤ 0.05).

Figure 6. Analysis of PTA and PTA-dl1013 shedding in C57BLC57bl/6 *Rag2*-/- mice.

Nine female C57BL/6-RAG2-/- mice were co-infected with 5 x10^4 IU of PTA and 5 x10^4 IU of

PTA-dl1013 via IP injection. Urine was collected during the acute phase of infection, and genomic

DNA was quantified by strain-specific qPCR analysis. Wild type C57BL/6 mice co-infected with

PTA (blue circles/dashed line) and PTA-dl1013 virus (yellow triangles/dashed line) are from

Figure 6. PTA genomes (blue circles/solid line) and PTA-dl1013 virus (yellow triangles/solid

line) from C57BL/6-Rag2-/- mice are presented. Plots represent the average (±SEM, n=8-9) at the

indicated times.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6