Symbiont switching and alternative resource acquisition strategies drive

mutualism breakdown

Gijsbert DA Werner*[1,2], Johannes HC Cornelissen[3], William K Cornwell[4], Nadejda A

Soudzilovskaia[5], Jens Kattge[6,7], Stuart A West[1], E Toby Kiers[3]

*gijsbert.werner@zoo.ox.ac.uk

1: Department of Zoology, University of Oxford, OX1 3PS, Oxford UK

2: Balliol College, University of Oxford, OX1 3BJ, Oxford, UK

3: Department of Ecological Sciences, Vrije Universiteit Amsterdam, Amsterdam, The

Netherlands

4: School of Biological, Earth and Environmental Sciences, University of New South Wales,

Sydney, Australia

5: Conservation Biology Department, Institute of Environmental Sciences, CML, Leiden

University

6: Max Planck Institute for Biogeochemistry, Jena, Germany

7: German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig,

Germany

Abstract

Cooperative interactions among species, termed mutualisms, have played a crucial role in the

evolution of life on Earth. However, despite key potential benefits to partners, there are many

cases where two species cease to cooperate, and mutualisms break down. What factors drive

the evolutionary breakdown of mutualism? We examined the pathways towards breakdowns

of the mutualism between plants and arbuscular mycorrhizal (AM) fungi. Using a

comparative approach, we identify ~25 independent cases of complete mutualism breakdown

across global seed plants. We found that breakdown of cooperation was only stable when

host plants either: (i) partner with other root symbionts or (ii) evolve alternative resource

acquisition strategies. Our results suggest that key mutualistic services are only permanently

lost if hosts evolve alternative symbioses or adaptations.

Significance Statement

Cooperative interactions among species – mutualisms – are major sources of evolutionary innovation. However, despite their importance, two species that formerly cooperated sometimes cease their partnership. Why do mutualisms breakdown? We asked this question in the partnership between arbuscular mycorrhizal (AM) fungi and their plant hosts, one of the most ancient mutualisms. We analyse two potential trajectories towards evolutionary breakdown of their cooperation, symbiont switching and mutualism abandonment. We find evidence that plants stop interacting with AM fungi when they switch to other microbial mutualists or when they evolve alternative strategies to extract nutrients from the environment. Our results show vital cooperative interactions can be lost - but only if successful alternatives evolve.

Introduction

Mutualisms, cooperative partnerships among different species, have shaped much of Earth's biodiversity, allowing organisms to outsource crucial functions like nutrition, cleaning, transport and defence (1, 2). Both theoretical and empirical work has provided us with a good understanding of the mechanisms, such as co-transmission and sanctions, that stabilise mutualism and maintain cooperation among species (3–5). Because of these mechanisms, beneficial interactions can be maintained over millions of years, and in some cases give rise to extreme mutualistic dependence (6, 7).

Despite reciprocal benefits, mutualisms do not always persist, and conflict among partners can remain. Theoretical and experimental work suggests that even when mutual benefits occur, fitness interests of both partners are generally not perfectly aligned, potentially selecting for cheaters and exploiters of mutualism (5, 8–11). This finding is further reinforced by the observation that in many mutualisms, there are mechanisms to evaluate partner quality and reward cooperation or sanction non-cooperative cheats (12–14). Furthermore, over ecological time, short-term breakdowns of cooperation in response to shifting environmental conditions, have been observed in many mutualisms, including plant rhizobial and mycorrhizal mutualisms, coral symbioses, protection and pollination mutualisms (15–19). Together, these observations raise the question in which conditions we should expect cooperation among species to fail, and partners in previously successful mutualisms to cease cooperating.

Even mutualisms that have become highly dependent over millions of years of co-evolution,
have broken down in some occasions. This is the case, for example, when free-living fungi
evolved from a previously lichenised lifestyle, or when parasitic moths evolved from
pollinating ancestors (20–22). Yet, while we have a good understanding of why mutualistic
cooperation is favoured, we lack a general understanding of the drivers of these evolutionary
breakdowns of mutualisms. A number of non-exclusive reasons for the breakdown of
mutualisms have been proposed (23, 24). Benefits provided by a mutualistic partner can
become redundant through the evolution of alternative adaptations. In these cases, one of the
partners switches from relying on another species to acquiring a function autonomously. For
example, the evolution of large amounts of small-diameter pollen enabled the reversion back
to an autonomous, wind-pollinated lifestyle in some angiosperms (25). A second trajectory
occurs when one side of the interaction is replaced with a new mutualist species. While
partner switching by definition leads to the evolution of a new partnership, the ancestral
interaction is lost and thus a previously functional mutualism breaks down. This is illustrated
in cases where plant species stop cooperating with birds and switch to insect pollination (26).
83
Our aim was to study the ancient and ubiquitous mutualism between plants and arbuscular
mycorrhizal (AM) fungi to understand pathways towards mutualism breakdown. We focus on
the plant-AM mutualism for three reasons. First, AM fungi (Glomeromycota) are among the
most important terrestrial mutualists. AM fungi form extensive hyphal networks in the soil
(up to 100 m cm^{-3} soil), providing plants with a key solution to the problem of extracting
immobile nutrients, especially phosphorus (27). The partnership is crucial for plant growth,
providing hosts with primarily phosphorus, but also nitrogen, water and trace elements (28).
Second, even though the large majority of plants can be successfully colonised by AM fungi,
10-20% of plant species across a number of divergent clades do not interact with any AM
fungi (27, 29). These repeated losses of the interaction, separated by millions of years of
evolution, enable us to test general patterns and explanatory factors driving cooperation loss
in a comparative framework. Third, the tools and databases allowing for broad comparative
analyses are becoming available for plants, including a comprehensive phylogeny of seed
plants (30), and large scale databases of plant traits including their association with AM fungi
and other root symbionts (31–33).
99
In our analysis of the plant – AM mutualism, we take a plant-centric perspective. We are
interested in cases where plants completely cease to interact with *all* AM fungi and where

this lack of interaction persists over evolutionary time: *i.e.* where the loss of the interaction is
not followed by host plant extinction. Thus, we do not study when plant-AM cooperation
dissolves in the short-term due to ecological conditions, such as under high nutrient
conditions. Rather, our aim is to first quantify stable losses of cooperation, and then test the
importance of two types of evolutionary breakdown: partner switching and mutualism
abandonment. By partner switching, we mean a situation where a host plant that ancestrally
interacted with AM fungi, switched to interacting with a novel root symbiont with similar
function and ceased interacting with AM fungi. We analyse switches to other mycorrhizal
fungi, as well as to N_2-fixing symbioses with rhizobial and *Frankia* bacteria (28, 34). We
refer to mutualism abandonment, when plants have evolved an alternative strategy to acquire
resource in a non-symbiotic way, for instance carnivory or cluster roots (35, 36).

113
114

Results

Evolutionary reconstruction of the plant-AM fungal mutualism

Our first aim was to quantify the evolutionary stability of the plant-AM mutualism,
determining the number of losses of plant-AM interactions across the plant phylogeny. We
compiled a global database of plant mycorrhizal fungal status across the seed plants
(angiosperms and gymnosperms). We scored the reported interactions of plants with AM
fungi in 3,736 plant species present in the most recent and comprehensive phylogeny of
gymnosperms and angiosperms (30). We then established patterns of AM loss and gain using
a Hidden Rate Model (HRM) approach to ancestral state reconstructions (37). This technique
permits variation in the speed of binary character evolution so we can detect changes in rates
of evolution, such as shifts in the evolutionary stability of plant-AM associations.

Table 1: AICc-values and weights for all HRM models

Number of rate classes	Number of parameters	AICc	Δ-AICc	AICc-weight
1	2	3231.25	770.0	0.0%
2	8	2597.78	136.5	0.0%
3	**14**	**2461.25**	**0**	**74.8%**
4	20	2463.44	2.2	25.1%
5	26	2473.79	12.5	0.1%

Table of the five different HRMs explored to analyse our AM fungal association data (see Methods). We used AICc-weights (corrected Akaike information criterion) to determine the model with the best fit (bold).

Our reconstructions revealed that the evolution of AM interactions across seed plants was
best characterised by heterogeneity in speed of evolution: the best evolutionary model
contains three different rate classes of evolution (Table 1). Specifically, we find strong
evidence for the existence of an evolutionary class where AM interactions are strongly
favoured (which we termed Stable AM), a class where an absence of AM interactions is
strongly favoured (Stable Non-AM), and a class where AM interactions are evolutionarily
labile (SI Figure 1).

Mapping these different evolutionary states back onto the phylogeny (SI Figure 2), we found
that: (i) Association with AM fungi was likely the ancestral state of seed plants (99.6%
likelihood); (ii) Stable AM fungal associations have been widely retained throughout the seed
plants for over 350 million years, and represent the large majority of all historical and
contemporary plant species and families (Table 2); (iii) some plant lineages evolve to either
an evolutionarily labile state or a state where AM fungi are disfavoured (SI Figure 1, Table 2).
Specifically, (iv) there have been an estimated ~25 evolutionary losses of the AM mutualism
throughout the history of seed plants, found across 69 families (median over 100 bootstrap
phylogenies 25.4, SD: 7.73). Which evolutionary trajectories are most important in
explaining these breakdowns of cooperation among plants and AM fungi?

Table 2: Number of contemporary species and families in three AM Classes

	Species numbers		Family numbers	
Stability Class	Best tree	Median	Best tree	Median
Stable AM	2,616	2613 (SD 178)	172	171 (SD 12)
Labile	829	833 (SD 180)	77	79 (SD 13)
Stable Non-AM	291	288 (SD 21)	8	7 (SD 3.7)

Numbers of contemporary species and families per AM stability class.
Median value and SD across 100 bootstrap phylogenies are indicated.

Symbiont switching and mutualism abandonment drive breakdown
We tested the hypotheses that AM loss is driven by shifts to other symbionts (partner
switching) or by alternative adaptations for resource acquisition (abandonment). We
generated a database of other major root symbionts with functional roles (providing
phosphorus and nitrogen) similar to AM fungi. Specifically, based on a previously published
database, we included presence or absence of a potential to interact with symbiotic N_2-fixing
bacteria (both rhizobial and *Frankia* bacteria) for all our host plant species (34, 38). We also
included reported interactions with other mycorrhizal fungi (*i.e.* non-AM fungi that live in

symbiotic association with plant roots). This included ectomycorrhizal (EM), ericoid

mycorrhizal (ER), orchid mycorrhizal (ORM) and arbutoid mycorrhizal (ARB) fungi. All

AM fungi belong to the division *Glomeromycota*, while other mycorrhizal fungi are only

distantly related, belonging to a wide range of divisions, mainly *Basidiomycota* (ECM, ARB

and ORM, some ER) and *Ascomycota* (some ECM and ORM and most ER). Some plant

species interact with multiple types of mycorrhizal fungi (28, 39).

We scored our species for the reported presence of alternative resource acquisition strategies.

These included parasitism as a plant strategy (both plants parasitising other plants and full

mycoheterotrophs, *i.e.* plants parasitising mycorrhizal fungi) (40, 41), carnivory (35) and

cluster roots (36) (Figure 1). These strategies have in common that they represent alternative

solutions to the problem of acquiring scarce mineral resources: they acquire resources by

seizing them from other organisms (plant parasitism), through direct predation (carnivorous

plants), or through investing in a unique root architecture characterised by a high density of

finely-branched roots and root hairs, known as cluster roots (Figure 1). To study congruence

between losses of AM interactions and alternative strategies, we again performed ancestral

state reconstructions, to study the origins of: (i) other symbionts (*i.e.* non-AM mycorrhizal

fungal symbionts or symbiotic N_2-fixation), which were present in 820 of our 3,736 plant

species; and (ii) alternative resource acquisition strategies, present in 109 plant species.

We found a high degree of congruence between the different origins of AM losses and of

various AM-alternatives (SI Table 1, SI Figures 3-9). To study this quantitatively, we

compared models of dependent *vs.* independent evolution (42), analysing the relationship

between AM loss and presence of alternative partners or alternative resources acquisition

strategies. We studied a binary variable coding for presence of any AM alternative and found

that a dependent model of evolution vastly outperformed an independent model (Δ-AICc

428.90, AICc-weight 99.9%). This means that over evolutionary time, AM loss (shift from

the bottom left plane to the top right in the transitions matrix, Figure 2) is strongly associated

with the presence of another mycorrhizal fungal partner, or alternative resource strategy.

Thus, partner switching and mutualism abandonment are important in enabling evolutionary

breakdown of the ancestral plant-AM fungal mutualism throughout the seed plants.

More specifically, from the inferred transition matrix and associated ancestral state
reconstruction (Figure 2), we conclude that: (i) The AM mutualism is generally highly stable
- transition rates towards the AM state (green) are about ten times as high as losses (from
green to yellow) (ii) AM fungal loss is only stable when an alternative is present (orange to
red) and not without (green to yellow). (iii) While evolutionary stability is high when plants
associate with either AM fungal symbionts (green state) or an alternative symbiont or
acquisition strategy (red), having neither (yellow) is evolutionarily unstable. For instance, all
the origins of this type (*e.g.* in the Brassicales) have occurred relatively recently in
evolutionary terms (within the last 30 million years). (iv) Similarly, it is evolutionarily less
stable to have both AM and an alternative simultaneously (orange).

Our reconstructions show that both the evolutionary scenario of initial AM loss followed by
alternative strategy evolution and the reverse order are possible. Initial acquisition of an AM-
alternative (move from green to orange state), in some cases may have resulted in released
selection to maintain the AM interaction, allowing for its subsequent evolutionary breakdown
(orange to red). In other cases, the AM interaction was lost first (yellow state), for instance
through symbiosis gene loss (43), and survival of host plants was subsequently favoured
when rapidly evolving an AM alternative state. Thus, our analysis indicates there is no single
dominant trajectory in the transition from an AM plant to a stable non-AM plant, but that
both routes can occur.

Sensitivity Analyses

To verify the robustness of our results, we considered the sensitivity of our main conclusions
to two forms of uncertainty, (1) phylogenetic uncertainty and (2) uncertainty in the
underlying AM data. We analysed phylogenetic uncertainty by replicating our initial AM
fungal reconstruction analysis over 100 bootstrap phylogenies, and found highly similar
relative loss rates of the plant AM-interaction throughout 100% of our bootstrap replicates
(SI Figure 10) and highly similar ancestral state reconstructions (SI Figure 11). We also
found that across the 100 bootstrap phylogenies a dependent model of evolution always
outperformed an independent model (mean Δ-AICc 390.52). This further confirms the deep
evolutionary link between AM loss and the evolution of other symbionts and resource
acquisition strategies regardless of the details of the phylogenies used (SI Figure 12).

A second main source of uncertainty is in the AM status of plants. This is because AM fungi
are notoriously difficult to score: it is easy to misidentify other fungi as AM fungi (false
positive) or to miss AM hyphae (false negative). To address this, we implemented a re-
simulation approach which takes into account the number of independent reports of AM
status, and allows us to test separate false positive and false negative rates for these
underlying reports. We found that even if one in four of the AM reports in our database is
incorrect (*e.g.* a saprotrophic fungus) while simultaneously 25% of our AM absence reports
in fact were mycorrhizal, we still draw highly similar conclusions (SI Figure 13, SI Table 2).
Therefore, even if we assume the underlying mycorrhizal data are of very poor quality, we
recover qualitatively highly similar patterns. Thus, overall, we conclude that all our main
conclusions are robust to substantial phylogenetic and data uncertainty.

As a final analysis, we compared our results with a recent comparative analysis of plant-
mycorrhizal symbioses (29). This analysis used an alternative scoring approach that divided
plant species in four categories: AM plants, non-mycorrhizal plants (NM-plants), ECM plants,
and plants that are commonly found in either AM or NM states (AMNM-plants), and found
that transitions from AM towards NM states primarily go through the AMNM state. We re-
confirmed this result, in that we find in our best HRM-model of plant-AM interactions that
plants transition through the labile state to the stable non-AM state, where the loss of plant-
AM mutualism becomes evolutionarily entrenched (SI Figure 1). We also find that the
species-level percentage of observations with AM-presence has a median value of 100% (SE:
0.89%; mean 83.4%) for species inferred to be in the stable AM class and 0% in the stable
non-AM class (SE: 0.73%; mean 1.58%), while in the evolutionarily labile class this is 16.7%
(SE: 1.57%; mean 22.0%; SI Figure 14). This indicates that the labile state inferred under our
deep evolutionary model effectively recovers the notion of an AMNM presupposed by
Maherali *et al*. While their analysis allows for direct inclusion of AMNM and ECM states,
with our approach of binary coding the presence or absence of AM and other mycorrhizal
interactions we can answer different questions: (i) It allows us to infer the *variation* in loss
rate of the AM mutualism across seed plant evolutionary history (which is only possible in
the HRM-framework for binary traits (37)) (ii) Rather than *a priori* defining an intermediate
state, it allows us to verify if an evolutionarily labile state is actually inferred in our best
model. (iii) It allows us to study the dependent evolution of AM and other mycorrhizal
interactions as separate traits. This is especially important because, while rare, dual
colonisation of plants by two types simultaneously is possible and could represent an

important evolutionary intermediary state, as confirmed by our analysis (Figure 2). (4) It

allows us to include in our analysis not just ECM fungi, but also other root symbionts such as

symbiotic N_2-fixation, ericoid (ERM) and orchid (ORM) mycorrhizal fungi, which turned out

to be drivers of major evolutionary losses of the plant-AM mutualism (Figure 2).

Discussion

Our analyses revealed that the ancient and ubiquitous plant–AM fungal mutualism has

broken down in ~25 cases across the seed plants. We found that stable and persistent

mutualism breakdown is driven both by acquisitions of other root symbionts (partner

switching) and by the evolution of alternative non-symbiotic resource acquisition strategies

(mutualism abandonment).

These results in turn raise the question of what underlying ecological factors favour

transitions to these alternative solutions, and the mechanisms that enable them.

Mechanistically, an important step is likely the loss of key genes in the 'symbiotic toolkit'

encoding crucial root mutualism effectors (43, 44). This must either be followed or preceded

by molecular evolution in the genes encoding alternative symbioses or resource acquisition

traits. Ecologically, these alternatives can potentially be favoured by a range of ultimate

factors, such as environmental change, habitat shifts (for instance to high-nutrient soils),

migration, invasion or partner abundance (22, 24, 45–49), although discriminating these over

deep evolutionary time is challenging. One hypothesis is that switching from the AM nutrient

uptake strategy to rarer alternative strategies has enabled plants to compete in a range of

(micro)habitats. Evolution of carnivory in temperate swamps (35), cluster-roots in extremely

phosphorus-impoverished soils (36), cold-resistant ectomycorrhizal interactions in lower

temperature habitats (50) and ericoid mycorrhizal fungi in resource-poor heath lands (51) has

helped host plants to thrive in environments where the more common AM interaction is a less

successful solution to obtain nutrients.

We emphasise that our estimate of ~25 breakdowns represents a conservative lower bound,

since we study plants that stopped interacting with *all* AM fungal species and subsequently

persisted over evolutionary times. The number of breakdowns of plant mutualism with

specific AM fungal species or lineages while cooperation with other AM continued, is likely

to be considerably higher. More generally, studies that analyse breakdowns of symbiotic

interactions with entire taxa of organisms - such as among corals and any photosynthetic

dinoflagellates (52, 53) – will underestimate the number of breakdowns with specific
symbiont species.

We find that dual symbioses – simultaneously being able to interact with two types of root
symbionts - is unlikely to be evolutionarily stable (Figure 2). This is a different pattern
compared to what is documented in insect endosymbioses, which often acquire secondary
partners while retaining the ancestral mutualism (54, 55). In insects, maintenance of two
endosymbionts could be favoured by different microbial partners subsequently specialising
on different mutualistic functions (56, 57). In root symbioses, nutritional benefits provided by
AM fungi and by alternative root symbionts may often be too similar to outweigh the costs of
maintaining them both. For instance, while AM fungi are thought to provide primarily
phosphorus, they also contribute nitrogen to their hosts (58). This could help explain why
plants only rarely associate with both ECM fungi and AM fungi simultaneously (131 species
in our dataset). However, our reconstructions suggest that such dual symbioses can be a
transitory state on the path towards a complete switch and breakdown of the original
mutualism (Figure 2), as was previously hypothesised (59). Other root symbionts may
provide more complementary benefits to their plant hosts, which could select for maintenance
of dual symbioses. For instance, AM fungi and N_2-fixing rhizobial bacteria are often thought
to provide complementary benefits to their legume hosts (60), although a meta-analysis did
not generally find synergistic effects on host growth (61).

If breakdown of the AM fungal mutualism is driven by acquisition of other root symbionts or
alternative resources strategies, how can we explain plants that have neither AM fungi nor an
alternative (yellow in Figure 2)? Recently, a member of the Brassicaceae, a family generally
lacking mycorrhizal symbionts, was found to engage in a specific and beneficial interaction
with fungi from the order Helotiales (Ascomycota), which provides soil nutrients (phosphate)
to their hosts (62). While we do not know how widespread this phenomenon yet is, it raises
the intriguing possibility that some of our species without AM fungi have in fact evolved
interactions with yet unknown beneficial root symbionts functionally similar to mycorrhizal
fungi. This would further strengthen the relationship we observed among AM loss and
switches to alternative symbionts. Another, non-mutually exclusive possibility is that plants
abandoning the AM fungal mutualism - without evolving alternatives - are likely to go
extinct after an evolutionarily short period of time, or rapidly re-establish the mutualism. In
line with this, all cases of AM breakdown not coupled to an alternative (yellow state; Figure

2), have evolved fairly recently (<30 MYA), compared to many much older losses associated

with symbiont switching or alternative strategies (e.g. the switch to ECM fungi in Pines,

more than 200 MYA).

An alternative potential reason for mutualism breakdown is when cheaters, low quality

partners or parasites, arise in one of the partner lineages (23, 24). This can drive the

interaction from mutual benefit to parasitism, and cause the other partner to abandon the

interaction (9). Theory and empirical work suggests that hosts are particularly vulnerable to

cheating when partners are acquired directly from the environment, like AM fungi (4, 5, 63–

65). However, in bacteria, phylogenetic work has shown that while transitions towards

cooperative states are common, loss of mutualist status is rare for bacterial symbionts (66,

67). When these losses occur, bacteria are more likely to revert to a free-living state than to

become parasites (66, 67). In our case, such a reversion to a free-living state would

correspond to a plant evolving an abiotic adaptation to replace AM fungi, such as cluster

roots. While most of our ~25 losses can be explained in terms of symbiont switches or

alternative resource strategies (Figure 2), some of the switches to other root symbionts or

resource strategies we observed could initially have been driven by the fitness cost of

parasitic AM fungi.

Our analyses show that cooperation among plants and AM fungi has generally persisted in a

highly stable state for over 350 million years. This illustrates the importance of mutualistic

services provided by AM fungi for most host plant species. Yet, even ancient and versatile

mutualists like AM fungi can be completely and permanently lost in the right circumstances:

we estimate this happened ~25 times. In general, mutualistic partnerships allow organisms to

outsource crucial functions to other species, thereby obtaining these services more efficiently

(5). Our results highlight how a key mutualistic service like nutrient acquisition is only

permanently lost if hosts evolve either symbiotic or abiotic alternatives to obtain these

functions.

Methods

More detailed Extended Methods can be found in the online Supporting Information.

Mycorrhizal status database

We compiled our database of reported plant mycorrhizal status by obtaining data from both primary

literature and publicly accessible databases. Our full data source list, as well as our scoring criteria

can be found in the Extended Methods (Supporting Information). Our analysed database contained

data for a total of 3,736 spermatophyte species (3,530 angiosperms, 206 gymnosperms, 61 orders, 230

families and 1,629 genera) that overlapped with the phylogeny used in our analysis (30), is available

online (Supporting Data 1).

Reconstruction of the evolution of AM interactions

We used a Hidden Markov Model approach called 'Hidden Rate Models' (HRMs) which allows for

heterogeneity in the loss and gain rates of a binary trait across a phylogeny (37). We used the R-

package *corHMM* (37)(version 1.18) in R 3.2.3 to analyse our mycorrhizal data and explored HRMs

with one to five rate classes, using AICc-weights to select the best HRM among this family of

candidate models (Table 1). We used the marginal method to perform ancestral state reconstructions

and employed Yang's method to compute the root state (68). We *a posteriori* labelled the three rate

classes under the best model 'Stable AM', 'Labile' and 'Stable Non-AM' (SI Figure 1).

Database Alternative Resource Acquisition Strategies

We generated a second database for all our 3,376 analysed species and scored each species for the

presence or absence of three main resource strategies, which each represent an alternative way of

extracting minerals from the environment: carnivory (35), parasitism (40, 41) and cluster roots (36).

Based on our previously generated database of plant species associating with symbiotic nitrogen-

fixing bacteria (34, 38), we also assigned all analysed species a binary symbiotic nitrogen-fixation

status. We describe our full data sources and scoring procedures in the Extended Methods.

Correlated evolution of AM interactions and AM-alternatives

We generated HRM-models (37) of both non-AM mycorrhizal fungi and adaptations for resource

acquisition (SI Table 1), plotted them onto our AM ancestral state reconstruction and visually

identified the origins of these AM-alternatives (SI Figure 3-9). We then tested the potential for

correlated evolution among AM fungi, other mycorrhizal fungi and resource acquisition adaptations.

Using AIC-criteria, we compared models of dependent and independent evolution (42, 69) among the

binary variables AM and AM-alternatives. We utilised the Maximum Likelihood implementation of

the Discrete-module in *BayesTraits V2*, and constrained the ancestral node of the phylogeny to have

AM fungi but none of the alternatives, as that is what our previous analyses had revealed (SI Figures

3-9).

Sensitivity analysis to phylogenetic and data uncertainty

We studied the robustness of our main conclusions to two main sources of uncertainty: phylogenetic

uncertainty and uncertainty in the underlying mycorrhizal data. We reran our key models (three rate

class HRM and correlated evolution models in *BayesTraits*) across hundred bootstrap phylogenies

(30) (SI Figure 11 and 12). To test for effects of data uncertainty, we used a resimulation approach

that for each species takes into the number of observations of a given mycorrhizal state and simulates

different error rates for underlying mycorrhizal observations (SI Figure 13, SI Table 2). We detail our

full approach to the sensitivity analyses in the Extended Methods.

Data Availability

Our full dataset, including number of reports of various mycorrhizal states across databases

(see Extended Methods), our resulting assignment of AM, ECM, ORM, ER, ARB and symbiotic N_2-

fixation states, and our assignment of alternative resources acquisition strategies (carnivory,

parasitism, mycoheterotrophy) is available online (Supporting Data 1).

References

1.   Bronstein JL (2015) *Mutualism* (Oxford University Press, Oxford, UK).

2.   Weber MG, Agrawal A a. (2014) Defense mutualisms enhance plant diversification. *Proc Natl Acad Sci*
111(46):16442–16447.

3.   West SA, Griffin AS, Gardner A (2007) Evolutionary Explanations for Cooperation. *Curr Biol* 17(16):R661–R672.

4.   Sachs JL, Mueller UG, Wilcox TP, Bull JJ (2004) The evolution of cooperation. *Q Rev Biol* 79(2):135–160.

5.   Leigh EG (2010) The evolution of mutualism. *J Evol Biol* 23(12):2507–2528.

6.   Bennett GM, Moran NA (2015) Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *Proc*
Natl Acad Sci 112(33):10169–10176.

7.   Neiman M, Lively CM, Meirmans S (2017) Why Sex? A Pluralist Approach Revisited. *Trends Ecol Evol*.
doi:10.1016/j.tree.2017.05.004.

8.   Herre EA, West SA (1997) Conflict of interest in a mutualism: documenting the elusive fig wasp-seed trade-off.
Proc R Soc London Ser B, Biol Sci 264:1501–1507.

9.   Ghoul M, Griffin AS, West S a (2014) TOWARD AN EVOLUTIONARY DEFINITION OF CHEATING.
Evolution (N Y) 68(2):318–331.

10.   Bao, Addicott (1998) Cheating in mutualism: defection of Yucca baccata against its yucca moths. *Ecol Lett*
1(3):155–159.

11.   Sachs JL, Ehinger MO, Simms EL (2010) Origins of cheating and loss of symbiosis in wild Bradyrhizobium. *J Evol*
Biol 23(5):1075–1089.

12.   Kiers ET, et al. (2011) Reciprocal Rewards Stabilize Cooperation in the Mycorrhizal Symbiosis. *Science (80-)*
333(6044):880–882.

13.   Kiers ET, Rousseau RA, West SA, Denison RF (2003) Host sanctions and the legume–rhizobium mutualism.
Nature 425(6953):78–81.

14.   Wang R-W, Dunn DW, Sun BF (2014) Discriminative host sanctions in a fig–wasp mutualism. *Ecology*
95(5):1384–1393.

15.   Husband R, Herre EA, Young JPW (2002) Temporal variation in the arbuscular mycorrhizal communities
colonising seedlings in a tropical forest. *FEMS Microbiol Ecol* 42(1):131–6.

16. Baker DM, Freeman CJ, Wong JCY, Fogel ML, Knowlton N (2018) Climate change promotes parasitism in a coral
symbiosis. *ISME J*. doi:10.1038/s41396-018-0046-8.

17. Batterman SA, et al. (2013) Key role of symbiotic dinitrogen fixation in tropical forest secondary succession.
Nature 502(7470):224–227.

18. Ohm JR, Miller TEX (2014) Balancing anti-herbivore benefits and anti-pollinator costs of defensive mutualists.
Ecology 95(10):2924–2935.

19. MENZEL F, KRIESELL H, WITTE V (2014) Parabiotic ants: the costs and benefits of symbiosis. *Ecol Entomol*
39(4):436–444.

20. Lutzoni F, Pagel M, Reeb V (2001) Major fungal lineages are derived from lichen symbiotic ancestors. *Nature*
411(6840):937–940.

21. Pellmyr O, Leebens-Mack J (2000) Reversal of Mutualism as a Mechanism for Adaptive Radiation in Yucca Moths.
Am Nat 156(S4):S62–S76.

22. Kawakita A, Mochizuki K, Kato M (2015) Reversal of mutualism in a leafflower-leafflower moth association: The
possible driving role of a third-party partner. *Biol J Linn Soc* 116(3):507–518.

23. Sachs J, Simms E (2006) Pathways to mutualism breakdown. *Trends Ecol Evol* 21(10):585–592.

24. Kiers ET, Palmer TM, Ives AR, Bruno JF, Bronstein JL (2010) Mutualisms in a changing world: an evolutionary
perspective. *Ecol Lett* 13(12):1459–1474.

25. Culley TM, Weller SG, Sakai AK (2002) The evolution of wind pollination in angiosperms. *Trends Ecol Evol*
17(8):361–369.

26. Whittall JB, Hodges S a (2007) Pollinator shifts drive increasingly long nectar spurs in columbine flowers. *Nature*
447(7145):706–709.

27. Parniske M (2008) Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nat Rev Microbiol* 6(10):763–
775.

28. van der Heijden MGA, Martin FM, Selosse M-A, Sanders IR (2015) Mycorrhizal ecology and evolution: the past,
the present, and the future. *New Phytol* 205(4):1406–1423.

29. Maherali H, Oberle B, Stevens PF, Cornwell WK, McGlinn DJ (2016) Mutualism Persistence and Abandonment
during the Evolution of the Mycorrhizal Symbiosis. *Am Nat* 188(5):E113–E125.

30. Zanne AE, et al. (2014) Three keys to the radiation of angiosperms into freezing environments. *Nature*
506(7486):89–92.

31. Kattge J, et al. (2011) TRY - a global database of plant traits. *Glob Chang Biol* 17(9):2905–2935.

32. Wang B, Qiu Y-L (2006) Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza*
16(5):299–363.

33. Akhmetzhanova AA, et al. (2012) A rediscovered treasure: mycorrhizal intensity database for 3000 vascular plant
species across the former Soviet Union. *Ecology* 93(3):689–690.

34. Werner GDA, Cornwell WK, Sprent JI, Kattge J, Kiers ET (2014) A single evolutionary innovation drives the deep
evolution of symbiotic N2-fixation in angiosperms. *Nat Commun* 5:4087.

35. Ellison AM, Gotelli NJ (2001) Evolutionary ecology of carnivorous plants. *Trends Ecol Evol* 16(11):623–629.

36. Neumann G, Martinoia E (2002) Cluster roots – an underground adaptation for survival in extreme environments.
Trends Plant Sci 7(4):162–167.

37. Beaulieu JM, O'Meara BC, Donoghue MJ (2013) Identifying hidden rate changes in the evolution of a binary
morphological character: the evolution of plant habit in campanulid angiosperms. *Syst Biol* 62(5):725–37.

38. Werner GDA, Cornwell WK, Cornelissen JHC, Kiers ET (2015) Evolutionary signals of symbiotic persistence in
the legume–rhizobia mutualism. *Proc Natl Acad Sci* 112(33):10262–10269.

39. Martin F, Kohler A, Murat C, Veneault-Fourrey C, Hibbett DS (2016) Unearthing the roots of ectomycorrhizal
symbioses. *Nat Rev Microbiol*. doi:10.1038/nrmicro.2016.149.

40.   Merckx VS (2013) *Mycoheterotrophy* ed Merckx V (Springer New York, New York, NY) doi:10.1007/978-1-4614-
5209-6.

41.   Těšitel J (2016) Functional biology of parasitic plants□: a review. *Plant Ecol Evol* 149(1):5–20.

42.   Pagel M (1994) Detecting Correlated Evolution on Phylogenies: A General Method for the Comparative Analysis
of Discrete Characters. *Proc R Soc B Biol Sci* 255(1342):37–45.

43.   Delaux P-M, et al. (2014) Comparative Phylogenomics Uncovers the Impact of Symbiotic Associations on Host
Genome Evolution. *PLoS Genet* 10(7):e1004487.

44.   Bravo A, York T, Pumplin N, Mueller LA, Harrison MJ (2016) Genes conserved for arbuscular mycorrhizal
symbiosis identified through phylogenomics. *Nat Plants* 2(2):15208.

45.   Oliver KM, Degnan PH, Burke GR, Moran N a (2010) Facultative symbionts in aphids and the horizontal transfer
of ecologically important traits. *Annu Rev Entomol* 55:247–266.

46.   Weese DJ, Heath KD, Dentinger BTM, Lau J a. (2015) Long-term nitrogen addition causes the evolution of less-
cooperative mutualists. *Evolution (N Y)* 69(3):631–642.

47.   Gutiérrez-Valencia J, Chomicki G, Renner SS (2017) Recurrent breakdowns of mutualisms with ants in the
neotropical ant-plant genus Cecropia (Urticaceae). *Mol Phylogenet Evol* 111:196–205.

48.   Chomicki G, Renner SS (2017) Partner abundance controls mutualism stability and the pace of morphological
change over geologic time. *Proc Natl Acad Sci* 114(15):3951–3956.

49.   Sheldrake M, et al. (2017) A phosphorus threshold for mycoheterotrophic plants in tropical forests. *Proc R Soc B*
Biol Sci 284(1848):20162093.

50.   Treseder KK, et al. (2014) Evolutionary histories of soil fungi are reflected in their large-scale biogeography. *Ecol*
Lett 17(9):1086–1093.

51.   READ D (1996) The Structure and Function of the Ericoid Mycorrhizal Root. *Ann Bot* 77(4):365–374.

52.   Kitahara M V., Cairns SD, Stolarski J, Blair D, Miller DJ (2010) A comprehensive phylogenetic analysis of the
scleractinia (Cnidaria, Anthozoa) based on mitochondrial CO1 sequence data. *PLoS One* 5(7).
doi:10.1371/journal.pone.0011490.

53.   Barbeitos MS, Romano SL, Lasker HR (2010) Repeated loss of coloniality and symbiosis in scleractinian corals.
Proc Natl Acad Sci 107(26):11877–11882.

54.   Hudson LN, et al. (2016) The database of the PREDICTS (Projecting Responses of Ecological Diversity In
Changing Terrestrial Systems) project. *Ecol Evol* (September 2016):145–188.

55.   Bennett GM, McCutcheon JP, MacDonald BR, Romanovicz D, Moran NA (2014) Differential genome evolution
between companion symbionts in an insect-bacterial symbiosis. *MBio* 5(5):e01697-14.

56.   Douglas AE (2016) How multi-partner endosymbioses function. *Nat Rev Microbiol* 14(12):731–743.

57.   Szabó G, et al. (2017) Convergent patterns in the evolution of mealybug symbioses involving different
intrabacterial symbionts. *ISME J* 11(3):715–726.

58.   Govindarajulu M, et al. (2005) Nitrogen transfer in the arbuscular mycorrhizal symbiosis. *Nature* 435(7043):819–
23.

59.   Brundrett MC, Tedersoo L (2018) Evolutionary history of mycorrhizal symbioses and global host plant diversity.
New Phytol. doi:10.1111/nph.14976.

60.   Larimer AL, Clay K, Bever JD (2014) Synergism and context dependency of interactions between arbuscular
mycorrhizal fungi and rhizobia with a prairie legume. *Ecology* 95(4):1045–1054.

61.   Larimer AL, Bever JD, Clay K (2010) The interactive effects of plant microbial symbionts: a review and meta-
analysis. *Symbiosis* 51(2):139–148.

62.   Almario J, et al. (2017) Root-associated fungal microbiota of nonmycorrhizal Arabis alpina and its contribution to
plant phosphorus nutrition. *Proc Natl Acad Sci* 114(44):E9403–E9412.

63.   Porter SS, Simms EL (2014) Selection for cheating across disparate environments in the legume-rhizobium

mutualism. *Ecol Lett* 17(9):1121–1129.

64. Salem H, Onchuru TO, Bauer E, Kaltenpoth M (2015) Symbiont transmission entails the risk of parasite infection. *Biol Lett* 11(12):20150840.

65. Gordon BR, et al. (2016) Decoupled genomic elements and the evolution of partner quality in nitrogen-fixing rhizobia. *Ecol Evol* 6(5):1317–1327.

66. Sachs JL, Skophammer RG, Regus JU (2011) Evolutionary transitions in bacterial symbiosis. *Proc Natl Acad Sci* 108(Supplement_2):10800–10807.

67. Sachs JL, Skophammer RG, Bansal N, Stajich JE (2013) Evolutionary origins and diversification of proteobacterial mutualists. *Proc R Soc B Biol Sci* 281(1775):20132146–20132146.

68. Yang Z (2006) *Computational Molecular Evolution* (Oxford University Press, Oxford, UK).

69. Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401(October 1999):877–884.

70. Armstrong A (2015) Ecophysiology: Root-cluster control. *Nat Plants* 1(2):15009.

71. Beyrle HF, Smith SE, Franco CMM, Peterson RL (1995) Colonization of Orchis morio protocorms by a mycorrhizal fungus: effects of nitrogen nutrition and glyphosate in modifying the responses. *Can J Bot* 73(8):1128–1140.

Acknowledgements

We thank SURFsara (www.surf-sara.nl) for support in using the Lisa Computing Cluster. The AM and AM-alternative illustrations in Figures 1 and 2 are based on figures in the public domain (CC0) with the exception of the illustrations for ericoid (courtesy Dr. David Midgley, CC-BY-SA) and arbuscular mycorrhizae (courtesy dr. Yoshihiro Kobae), root nodules (courtesy Dr. Euan James) and the illustrations for cluster roots (70) and orchid mycorrhizae (71) which were reprinted with permission from the respective publishers. The study has been supported by the TRY initiative on plant traits (http://www.trydb.org). GDAW was funded by a Royal Society Newton International Fellowship and a Junior Research Fellowship at Balliol College Oxford. ETK was funded by Netherlands Organisation for Scientific Research Grants 836.10.001 and 864.10.005 and European Research Council ERC Grant Agreement 335542.

Figure Legends

Figure 1: We explored the evolution of various alternatives to interacting with AM fungi, the ancestral state of seed plants. Examples of six important alternatives we considered are depicted, with columns indicating two potential pathways towards evolutionary breakdown of the plant-AM mutualism. In both pathways, the ancestral mutualism with AM fungi breaks down. Coloured borders match coloured bars in Figure 2, indicating distribution of these traits across global seed plants.

Figure 2: Transition rates and ancestral state reconstruction of the dependent evolutionary model for plant AM status and AM-alternatives. The four potential evolutionary states are represented by colours in the transition matrix and on the seed plant phylogeny. Transition rates are expressed as number of transitions per 100 million years per lineage. The ancestral state is AM presence with no AM-alternatives (green), red indicates the switch to one of the AM alternatives (i.e. another mycorrhizal fungus, symbiotic N_2-fixation,

563   parasitism, carnivory or cluster roots). From inside to outside, coloured bands around the phylogeny

564   indicate the presence or absence of (i) AM interactions, (ii) other root symbionts and (iii) alternative

565   resource acquisition strategies. Key clades that have lost AM fungal interactions are indicated with

566   schematic images of their evolved alternatives. Grey and white concentric circles indicate 50 million years.

567   An expanded version with fully legible species labels for all 3,736 species is available online.

Symbiont Switching

Mutualism Abandonment

Ancestral Mutualism

Arbuscular mycorrhizal (AM) fungi

Arbuscular Mycorrhizal Fungi Alternatives

Ectomycorrhizal (ECM) fungi

Carnivory

Orchid mycorrhizal (ORM) fungi

Cluster Roots

Bacterial symbiotic N_2-fixation

Parasitic Plants

Legend Bands

AM Interaction (Inner Band)
- ▮ AM Present

Other symbionts (Middle Band)
- ▮ Ectomycorrhizal
- ▮ Orchid
- ▮ Ericoid / Arbutoid
- ▮ Symbiotic N₂-fixation

Alternative strategies (Outer Band)
- ▮ Carnivorous Plants
- ▮ Cluster Roots
- ▮ Parasitic Plants

Proteaceae

Cyperaceae

Myrtaceae

Orchids

Fagales

Pines

Detariae

Orobanchaceae

Lentibulariaceae

Fabaceae

Santales

Drosera

Monotropoidea

Ericaceae

No AM No alternatives

11.90   0.04

1.54   0.50

AM No alternatives   **No AM Alternatives**

0.84   0.63

0.22   1.52

AM & Alternatives