Symbiont switching and alternative resource acquisition strategies drive

mutualism breakdown

Gijsbert DA Werner*[1,2], William K Cornwell[3], Johannes HC Cornelissen[4], Nadejda A

Soudzilovskaia[5], Jens Kattge[6,7], Stuart A West[1], E Toby Kiers[4]

*gijsbert.werner@zoo.ox.ac.uk

1: Department of Zoology, University of Oxford, OX2 6GG, Oxford UK

2: Balliol College, University of Oxford, OX1 3BS, Oxford, UK

3: School of Biological, Earth and Environmental Sciences, University of New South Wales,

Sydney, Australia

4: Department of Ecological Sciences, Vrije Universiteit Amsterdam, Amsterdam, The

Netherlands

5: Conservation Biology Department, Institute of Environmental Sciences, CML, Leiden

University

6: Max Planck Institute for Biogeochemistry, Jena, Germany

7: German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig,

Germany

Abstract

Cooperative interactions among species, termed mutualisms, have played a crucial role in the

evolution of life on Earth. However, despite key potential benefits to partners, there are many

examples in which mutualisms break down, and two species cease to cooperate. What factors

drive these breakdowns? We examined the pathways towards the evolutionary breakdown of

the mutualism between plants and arbuscular mycorrhizal (AM) fungi. Using a comparative

approach, we identify ~25 independent cases of mutualism breakdown across global seed

plants. We found that breakdown of cooperation was only stable when host plants either: (i)

partner with other root symbionts or (ii) evolve alternative resource acquisition strategies.

Our results suggest that key mutualistic services are only permanently lost if hosts evolve

alternative symbioses or adaptations.

Significance Statement

Cooperative interactions among species – mutualisms – are major sources of evolutionary

innovation. However, despite their importance, two species that formerly cooperated

sometimes cease their partnership. Why do mutualisms breakdown? We asked this question

in the partnership between Arbuscular mycorrhizal (AM) fungi and their plant hosts, one of

the most ancient mutualisms. We analyse two potential trajectories towards breakdown of

their cooperation, symbiont swichting and mutulism abandonment. We find evidence that

plants stop interacting with AM fungi when they switch to other microbial mutualists or

when they evolve alternative strategies to extract nutrients from the environment. Our results

show vital cooperative interactions can be lost - but only if successful alternatives evolve.

Introduction

Mutualisms, cooperative partnerships among different species, have shaped much of the

Earth's biodiversity. Chemosynthetic symbionts enable animal colonisation of extreme deep

sea habitats (1), nutrient-providing mutualists allow hosts to thrive in resource-poor

conditions (2, 3) and plant outsourcing of defence functions to insects has driven host plant

diversification (4). The existence and success of these partnerships pose the question why

individuals should help members of another species? Theoretical and empirical studies have

provided us with a good understanding of the mechanisms, such as co-transmission and

sanctions, that stabilise mutualism and maintain cooperation (5–7). These factors allow stable

mutualisms to be maintained over millions of years, in some cases giving rise to extreme

mutualistic dependence (8–11).

However, in some cases, cooperation fails and species that used to be successful mutualists,

cease to cooperate (12, 13). For instance, free-living fungi have evolved from symbiotic

lichen ancestors (14). The shipworm *Kuphus polythalamia* abandoned its ancestral

cellulolytic symbionts, replacing them with chemoautotrophic bacteria that allowed it to

switch from a wood-boring lifestyle to living in marine sediments (15). Parasitic moths

evolved from mutualistic pollinating ancestors, leading to the collapse of an ancestral

pollination partnership (16–18). Why do mutualisms dissolve? In contrast to our relatively

good understanding of why mutualistic cooperation is favoured, we lack a general

understanding of why mutualisms breakdown.

In general, a mutualistic interaction is only expected to persist over evolutionary time if its

benefits outweigh the costs. This cost-to-benefit ratio can shift for a number of non-mutually

exclusive reasons, leading to different potential trajectories towards the evolutionary

breakdown of mutualism (12, 13). For instance, the benefits provided by the mutualistic

partner can become redundant through the evolution of alternative adaptations supplying the

previously mutualistically provided function (mutualism abandonment). In these cases, one

of the partners switches from relying on another species to acquiring a function

autonomously. For example, the evolution of large amounts of small-diameter pollen enabled

the reversion back to an autonomous, wind-pollinated lifestyle in some angiosperms (19). A

second trajectory occurs when one side of the interaction is replaced with a new mutualist

species (partner switching). While partner switching by definition leads to the evolution of a

new partnership, the ancestral interaction is lost and thus a previously functional mutualism

breaks down. This is illustrated in cases where plant species stop cooperating with birds and

switch to insect pollination (20). Breakdowns have been documented in a range of

mutualisms, but we do not have a good understanding of the driving factors.

Our aim was to study the ancient and ubiquitous mutualism between plants and arbuscular

mycorrhizal (AM) fungi to understand trajectories towards mutualism breakdown. We focus

on the plant-AM mutualism for three reasons. First, AM fungi (Glomeromycota) are among

the most important terrestrial mutualists (21). AM fungi form extensive hyphal networks in

the soil (up to 100 m cm^{-3} soil), providing plants with a key solution to the problem of

extracting immobile nutrients, especially phosphorus (21). The partnership is crucial for plant

growth, providing hosts with primarily phosphorus, but also nitrogen, water and trace

elements (22, 23). Second, even though the large majority of plants can be successfully

colonised by AM fungi, 10-20% of species across a number of divergent clades do not

interact with AM fungi (21, 24). These repeated losses of the interaction, separated by

millions of years of evolution, enable us to test general patterns and explanatory factors

driving cooperation loss in a comparative framework. Third, the tools and databases allowing

for broad comparative analyses are becoming available for plants, including a comprehensive

phylogeny of seed plants (25), and large scale databases of plant traits including their

association with AM fungi and other root symbionts (26–28).

In our analysis of the plant – AM mutualism, we take a plant-centric perspective. We are

interested in cases where plants cease to interact with AM fungi and where this lack of

interaction persists over evolutionary time: *i.e.* where the loss of the interaction is not followed by host plant extinction. Our aim is to first quantify how common stable losses of cooperation are, and then test the relative importance of two evolutionary pathways to mutualism breakdown: (i) replacement of the AM fungal mutualist with a novel symbiont with a similar function in providing soil nutrients (partner switching) and (ii) evolution of alternative resource acquisition strategies by the plant (mutualism abandonment). By partner switching, we mean a situation where a host plant that ancestrally interacted with AM fungi, switched to interacting with other root symbionts and ceased interacting with AM fungi. We analyse both switches to other mycorrhizal fungi as well as to N_2-fixing symbioses with rhizobial and *Frankia* bacteria (23, 29). We refer to mutualism abandonment, when plants have evolved an alternative strategy to acquire resource in a non-symbiotic way, for instance carnivory or cluster roots (30, 31).

Results

Evolutionary reconstruction of the plant-AM fungal mutualism

Our first aim was to quantify the evolutionary stability of the plant-AM mutualism, determining the number of losses of plant-AM interactions across the plant phylogeny. We compiled a global database of plant mycorrhizal fungal status across the seed plants (angiosperms and gymnosperms). We scored the reported interactions of plants with AM fungi in 3,736 plant species present in the most recent and comprehensive phylogeny of gymnosperms and angiosperms (25). We then established patterns of AM loss and gain using a Hidden Rate Model (HRM) approach to ancestral state reconstructions (32). This technique permits variation in the speed of binary character evolution so we can detect changes in rates of evolution, such as shifts in the evolutionary stability of plant-AM associations.

Table 1: AICc-values and weights for all HRM models

Number of rate classes	Number of parameters	AICc	Δ-AICc	AICc-weight
1	2	3231.25	770.0	0.0%
2	8	2597.78	136.5	0.0%
3	**14**	**2461.25**	**0**	**74.8%**
4	20	2463.44	2.2	25.1%
5	26	2473.79	12.5	0.1%

Table of the five different HRMs explored to analyse our AM fungal association data (see Methods). We used AICc-weights to determine the model with the best fit (bold).

Our reconstructions revealed that the evolution of AM interactions across seed plants was
best characterised by heterogeneity in speed of evolution: the best evolutionary model
contains three different rate classes of evolution (Table 1). Specifically, we find strong
evidence for the existence of an evolutionary class where AM interactions are strongly
favoured (which we termed Stable AM), a class where an absence of AM interactions is
strongly favoured (Stable Non-AM), and a class where AM interactions are evolutionarily
labile (SI Figure 1).

Mapping these different evolutionary states back onto the phylogeny (SI Figure 2), we found
that: (i) Association with AM fungi was likely the ancestral state of seed plants (99.6%
likelihood); (ii) Stable AM fungal associations have been widely retained throughout the seed
plants for over 350 million years, and represent the large majority of all historical and
contemporary plant species and families (Table 2); (iii) some plant lineages evolve to either
an evolutionarily labile state or a state where AM are disfavoured (SI Figure 1, Table 2).
Specifically, (iv) there have been an estimated 25 evolutionary losses of the AM mutualism
throughout the history of seed plants, found across 69 families (median over 100 bootstrap
phylogenies 25.4, SD: 7.73). Which evolutionary trajectories are most important in
explaining these breakdowns of cooperation among plants and AM fungi?

Table 2: Number of contemporary species and families in three AM Classes

Stability Class	Species numbers			Family numbers		
	Best tree	Median	SD	Best tree	Median	SD
Stable AM	2,616	2613	178	172	171	12.4
Labile	829	833	180	77	79	12.6
Stable Non-AM	291	288	21	8	7	3.7

Columns indicate the numbers of contemporary species and families per AM stability class. Median value and standard deviations across 100 bootstrap phylogenies are indicated. Supporting Data 1 provides the precise likelihood of the three stability classes for each species.

Symbiont switching and mutualism abandonment drive breakdown
We tested the hypotheses that AM loss is driven by shifts to other symbionts (partner
switching) or by alternative adaptations for resource acquisition (abandonment). We
generated a database of other major root symbionts with functional roles (providing
phosphorus and nitrogen) similar to AM fungi. Specifically, based on a previously published
database, we included presence or absence of a potential to interact with symbiotic N_2-fixing
bacteria (both rhizobial and *Frankia* bacteria) for all our host plant species (29, 33). We also
included reported interactions with other mycorrhizal fungi (*i.e.* fungi that live in symbiotic

association with plant roots but are not AM fungi). This included ectomycorrhizal (EM),

ericoid mycorrhizal (ER), orchid mycorrhizal (ORM) and arbutoid mycorrhizal (ARB) fungi.

All AM fungi belong to the division *Glomeromycota*, while other mycorrhizal fungi are only

distantly related, belonging to a wide range of divisions, mainly *Basidiomycota* (ECM, ARB

and ORM, some ER) and *Ascomycota* (some ECM and ORM and most ER). Some plant

species interact with multiple types of mycorrhizal fungi (23, 34).

We scored our species for the reported presence of alternative resource acquisition strategies.

These included parasitism as a plant strategy (both plants parasitising other plants and full

mycoheterotrophs, *i.e.* plants parasitising mycorrhizal fungi) (35, 36), carnivory (30) and

cluster roots (31) (Figure 1). These strategies have in common that they represent alternative

solutions to the problem of acquiring scarce mineral resources: they acquire resources by

seizing them from other organisms (plant parasitism), through direct predation (carnivorous

plants), or through investing in a unique root architecture characterised by a high density of

finely-branched roots and root hairs, known as cluster roots (Figure 1). To study congruence

between losses of AM interactions and alternative strategies, we again performed ancestral

state reconstructions, to study the origins of: (i) other symbionts (*i.e.* non-AM mycorrhizal

fungal symbionts or symbiotic N_2-fixation), which were present in 820 of our 3,736 plant

species; and (ii) alternative resource acquisition strategies, present in 109 plant species

(Figure 1).

We found a high degree of congruence between the different origins of AM losses and of

various AM-alternatives (SI Table 1, SI Figures 3-9). To study this quantitatively, we

compared models of dependent *vs* independent evolution (37), analysing the relationship

between AM loss and presence of alternative partners or alternative resources acquisition

strategies. We studied a binary variable coding for presence of any AM alternative and found

that a dependent model of evolution vastly outperformed an independent model (Δ-AICc

428.90, AICc-weight 99.9%). This means that over evolutionary time, AM loss (shift from

the bottom left plane to the top right in the transitions matrix, Figure 2) is strongly associated

with the presence of another mycorrhizal fungal partner, or alternative resource strategy.

Thus, both partner switching and mutualism abandonment are important in enabling

evolutionary breakdown of the ancestral plant-AM fungal mutualism throughout the seed

plants.

More specifically, from the inferred transition matrix and associated ancestral state

reconstruction (Figure 2), we conclude that: (i) The AM mutualism is generally highly stable:

transition rates towards the AM state (green) are about ten times as high as losses (from green

to yellow) (ii) AM fungal loss is only stable when an alternative is present (orange to red) and

not without (green to yellow). (iii) While evolutionary stability is high when plants associate

with either AM fungal symbionts (green state) or an alternative symbiont or acquisition

strategy (red), having neither (yellow) is evolutionarily unstable. For instance, all the origins

of this type (*e.g.* in the Brassicales) have occurred relatively recently in evolutionary terms

(within the last 30 million years). (iv) Similarly, it is evolutionarily less stable to have both

simultaneously (orange, *e.g.* AM fungi and alternatives together).

Our reconstructions show that both the evolutionary scenario of initial AM loss followed by

alternative strategy evolution and the reverse order are possible. Initial acquisition of an AM-

alternative (move from green to orange state), in some cases may have resulted in released

selection to maintain the AM interaction, allowing for its subsequent evolutionary breakdown

(orange to red). In other cases, the AM interaction was lost first (yellow state), for instance

through symbiosis gene loss (38), and survival of host plants was subsequently favoured

when rapidly evolving an AM alternative state. Thus, our analysis indicates that there is no

single dominant trajectory in the transition from an AM plant to a stable non-AM plant, but

that both routes can occur.

Sensitivity Analyses

To verify the robustness of our results, we considered the sensitivity of our main conclusions

to two forms of uncertainty, (1) phylogenetic uncertainty and (2) uncertainty in the

underlying AM data. We analysed phylogenetic uncertainty by replicating our initial AM

fungal reconstruction analysis over 100 bootstrap phylogenies, and found highly similar

relative loss rates of the plant AM-interaction throughout 100% of our bootstrap replicates

(SI Figure 10) and highly similar ancestral state reconstructions (SI Figure 11). Studying the

sensitivity of our correlated evolution models of AM fungi and AM-alternatives (Figure 2),

we found that across the 100 bootstrap phylogenies the dependent model always

outperformed the independent model (mean Δ-AICc 390.52). This further confirms the deep

evolutionary link between AM loss and the evolution of other symbionts and resource
acquisition strategies regardless of the details of the phylogenies used (SI Figure 12).
222
A second main source of uncertainty is in the AM status of plants. This is because AM fungi
are notoriously difficult to score: it is easy to misidentify other fungi as AM fungi (false
positive) or to miss AM hyphae (false negative). To address this, we implemented a re-
simulation approach which takes into account the number of independent reports of AM
status, and allows us to test separate false positive and false negative rates for these
underlying reports. We found that even if one in four of the AM reports in our database is
incorrect (*e.g.* a saprotrophic fungus) while simultaneously 25% of our AM absence reports
in fact were mycorrhizal, we still draw highly similar conclusions (SI Figure 13, SI Table 2).
Therefore, even if we assume the underlying mycorrhizal data are of poor quality, we recover
qualitatively highly similar patterns. Thus, overall, we conclude that all our main conclusions
are robust to substantial phylogenetic and data uncertainty.
234
As a final analysis, we compared our results with a recent comparative analysis of plant-
mycorrhizal symbioses (24). This analysis used an alternative scoring approach that divided
plant species in four categories: AM plants, non-mycorrhizal plants (NM-plants), ECM plants,
and plants that are commonly found in either AM or NM states (AMNM-plants) (24), and
found that transitions from AM towards NM states primarily go through the AMNM state.
We re-confirmed this result, in that we find in our best HRM-model of plant-AM interactions
that plants transition through the labile state to the stable non-AM state, where the loss of
plant-AM mutualism becomes evolutionarily entrenched (SI Figure 1). We also find that the
species-level percentage of observations with AM-presence has a median value of 100% (SE:
0.89%; mean 83.4%) for species inferred under our HRM model as being in the stable AM
class and 0% in the stable non-AM class (SE: 0.73%; mean 1.58%), while in the
evolutionarily labile class this is 16.7% (SE: 1.57%; mean 22.0%; SI Figure 14). This
indicates that the labile state inferred under our deep evolutionary model effectively recovers
the notion of an AMNM presupposed by Maherali *et al*. While their analysis allows for direct
inclusion of AMNM and ECM states, with our approach of binary coding the presence or
absence of AM and other mycorrhizal interactions we can answer different questions: (i) It
allows us to infer the *variation* in loss rate of the AM symbiosis across seed plant
evolutionary history (which is only possible in the HRM-framework for binary traits (32)) (ii)
Rather than *a priori* defining an intermediate state, it allows us to verify if an evolutionarily

labile state is actually inferred in our best model. (iii) It allows us to study the dependent
evolution of AM and other mycorrhizal interactions as separate traits. This is especially
important because, while rare, dual colonisation of plants by two types simultaneously is
possible and could represent an important evolutionary intermediary state, as confirmed by
our analysis (Figure 2). (4) It allows us to include in our analysis not just ECM fungi, but also
other root symbionts such as symbiotic N_2-fixation, ericoid (ERM) and orchid (ORM)
mycorrhizal fungi, which turned out to be drivers of major evolutionary losses of the plant-
AM mutualism (Figures 1 & 2).

Discussion

Our analyses revealed that the ancient and ubiquitous plant–AM fungal mutualism has
broken down in ~25 cases across the seed plants. We found that mutualism breakdown is
driven both by acquisitions of other root symbionts (partner switching) and by the evolution
of alternative non-symbiotic resource acquisition strategies (mutualism abandonment)
(Figure 2). This in turn raises the question of what underlying ecological factors favour
transitions to these alternative solutions, and the mechanisms that enable them.
Mechanistically, an important step is likely the loss of key genes in the 'symbiotic toolkit'
encoding crucial root mutualism effectors (38, 39). This must either be followed or preceded
by molecular evolution in the genes encoding alternative symbioses or resource acquisition
traits. Ecologically, these alternatives can potentially be favoured by a range of ultimate
factors, such as environmental change, habitat shifts (for instance to high-nutrient soils),
migration, invasion or partner abundance (13, 18, 40–44), although discriminating these over
deep evolutionary time is challenging. One hypothesis is that switching from the AM nutrient
uptake strategy to rarer alternative strategies has enabled plants to compete in a range of
(micro)habitats. Evolution of carnivory in temperate swamps (30), cluster-roots in extremely
phosphorus-impoverished soils (31), cold-resistant ectomycorrhizal interactions in lower
temperature habitats (45) and ericoid mycorrhizal fungi in resource-poor heath lands (46) has
helped host plants to thrive in environments where the more common AM interaction is a less
successful solution to obtain nutrients.

We find that interacting with two types of root symbionts is unlikely to be evolutionarily
stable (Figure 2). This is a different pattern from what has been documented in many insect
endosymbioses, which often acquire secondary partners while retaining the ancestral
mutualism (2, 47, 48). In insects, maintenance of two endosymbionts could be favoured by

different microbial partners subsequently specialising on different mutualistic functions (2,
49). This can be driven by symbionts losing key functions due to extreme bottlenecks and
genetic drift associated with obligate vertical transmission of symbionts across insect
generations (8). In contrast, mycorrhizal fungi are horizontally transmitted and do not
experience the same selection for genome reduction (11, 50). In mycorrhizal symbioses,
nutritional benefits provided by AM fungi and by alternative root symbionts may be too
similar to outweigh the costs of maintaining them both. For instance, symbiotic N_2-fixation,
which is generally found in plants that also interact with AM fungi, appears to be lost at a
high rate (29). Our reconstruction suggests that simultaneous interaction with two types of
root symbionts can be a transitory state on the path towards a complete switch and
breakdown of the original mutualism (Figure 2).
299
If breakdown of the AM fungal mutualism is driven by acquisition of other root symbionts or
resources strategies, how can we explain plants that have neither AM nor an alternative
(yellow in Figure 2)? Recently, a member of the Brassicaceae, a family generally lacking
mycorrhizal symbionts, was found to engage in a specific and beneficial interaction with
fungi from the order Helotiales (Ascomycota), which provides soil nutrients (phosphate) to
their hosts (51). While we do not know how widespread this phenomenon yet is, it raises the
intriguing possibility that some of our species without AM fungi have in fact evolved
interactions with yet unknown beneficial root symbionts functionally similar to mycorrhizal
fungi. This would further strengthen the relationship we observed among AM loss and
switches to alternative symbionts. Another, non-mutually exclusive, possibility is that plants
abandoning the AM fungal mutualism without evolving alternatives are likely to go extinct
after an evolutionarily short period of time or rapidly re-establish the mutualism. In line with
this, all cases of AM breakdown not coupled to an alternative (yellow state; Figure 2), have
evolved fairly recently (<30 MYA), compared to many much older losses associated with
symbiont switching or alternative strategies (e.g. the switch to ECM fungi in Pines, more than
200 MYA).
316
An alternative potential pathway to mutualism breakdown is when cheaters, low quality
partners or parasites, arise in one of the partner lineages (12, 13). This can drive the
interaction from mutual benefit to parasitism and cause the other partner to abandon the
interaction (52). Theory and empirical work suggests that hosts are particularly vulnerable to
cheating when partners are acquired directly from the environment, like AM fungi (6, 7, 53–

55). However, phylogenetic analysis suggested that a biotroph fungal lifestyle – the lifestyle
of AM fungi – is likely an evolutionarily stable state, unlike antagonistic (necrotrophic) and
neutral (endophytic) lifestyles (56, 57). Similarly, in bacteria, phylogenetic work has shown
that while transitions towards cooperative states are common, loss of mutualist status is rare
for bacterial symbionts (58, 59). When these losses occur, bacteria are more likely to revert to
a free-living state than to become parasites (58, 59). In our case, such a reversion to a free-
living state would correspond to a plant evolving an abiotic adaptation to replace AM fungi,
such as cluster roots. While most of our ~25 independent losses can be explained in terms of
symbiont switches or alternative resource strategies (Figure 2), some of the switches to other
root symbionts or resource strategies we observed could initially have been driven by the
fitness cost of parasitic AM fungi. Testing this hypothesis over deep evolutionary time
requires data on the fitness effects of individual AM fungal strains and on their coexistence
with host plants, across different habitats in a large number of species.

In conclusion, our analyses show that cooperation among plants and AM fungi is
extraordinarily persistent over evolutionary time. In most cases, it has been in a highly stable
state for over 350 million years. This illustrates the importance of mutualistic services
provided by AM fungi for most host plant species. In general, mutualistic partnerships allow
organisms to outsource crucial functions to other species, thereby obtaining these services
more efficiently (7). Our results highlight how a key mutualistic service like nutrient
acquisition is only permanently lost if hosts evolve either symbiotic or abiotic alternatives to
obtain these functions.

Methods
More detailed Extended Methods can be found in the online Supporting Information.

Mycorrhizal status database
We compiled our database of reported plant mycorrhizal status by obtaining data from both primary
literature and publicly accessible databases. Our full data source list, as well as our scoring criteria
can be found in the Extended Methods (Supporting Information). Our analysed database contained
data for a total of 3,736 spermatophyte species (3,530 angiosperms, 206 gymnosperms, 61 orders, 230
families and 1,629 genera) that overlapped with the phylogeny used in our analysis (25).

Reconstruction of the evolution of AM interactions

We used a Hidden Markov Model approach called 'Hidden Rate Models' (HRMs) which allows for
heterogeneity in the loss and gain rates of a binary trait across a phylogeny (32). We used the R-
package *corHMM* (32)(version 1.18) in R 3.2.3 to analyse our mycorrhizal data and explored HRMs
with one to five rate classes, using AICc-weights to select the best HRM among this family of
candidate models (Table 1). We used the marginal method to perform ancestral state reconstructions
and employed Yang's method to compute the root state (60). We *a posteriori* labelled the three rate
classes under the best model 'Stable AM', 'Labile' and 'Stable Non-AM' (SI Figure 1).
363
Database Alternative Resource Acquisition Strategies
We generated a second database for all our 3,376 analysed species and scored each species for the
presence or absence of three main resource strategies, which each represent an alternative way of
extracting minerals from the environment: carnivory (30), parasitism (35, 36) and cluster roots (31).
Based on our previously generated database of plant species associating with symbiotic nitrogen-
fixing bacteria (29, 33), we also assigned all analysed species a binary symbiotic nitrogen-fixation
status. We describe our full data sources and scoring procedures in the Extended Methods.
371
Correlated evolution of AM interactions and AM-alternatives
We generated HRM-models (32) of both non-AM mycorrhizal fungi and adaptations for resource
acquisition (SI Table 1), plotted them onto our AM ancestral state reconstruction and visually
identified the origins of these AM-alternatives (SI Figure 3-9). We then tested the potential for
correlated evolution among AM fungi, other mycorrhizal fungi and resource acquisition adaptations.
Using AIC-criteria, we compared models of dependent and independent evolution (37, 61) among the
binary variables AM and AM-alternatives. We utilised the Maximum Likelihood implementation of
the Discrete-module in *BayesTraits V2*, and constrained the ancestral node of the phylogeny to have
AM fungi but none of the alternatives, as that is what our previous analyses had revealed (SI Figures
3-9).
382
Sensitivity analysis to phylogenetic and data uncertainty
We studied the robustness of our main conclusions to two main sources of uncertainty: phylogenetic
uncertainty and uncertainty in the underlying mycorrhizal data. We reran our key models (three rate
class HRM and correlated evolution models in *BayesTraits*) across hundred bootstrap phylogenies
(25) (SI Figure 11 and 12). To test for effects of data uncertainty, we used a resimulation approach
that takes into the number of observations per species of a given mycorrhizal state and simulates
different error rates for underlying mycorrhizal observations (SI Figure 13, SI Table 2). We detail our
full approach to the sensitivity analyses in the Extended Methods.
391
References

1. Kleiner M, et al. (2012) Metaproteomics of a gutless marine worm and its symbiotic microbial community reveal
unusual pathways for carbon and energy use. *Proc Natl Acad Sci* 109(19):E1173–E1182.

2. Douglas AE (2016) How multi-partner endosymbioses function. *Nat Rev Microbiol* 14(12):731–743.

3. Brock DA, Douglas TE, Queller DC, Strassmann JE (2011) Primitive agriculture in a social amoeba. *Nature*
469(7330):393–396.

4. Weber MG, Agrawal A a. (2014) Defense mutualisms enhance plant diversification. *Proc Natl Acad Sci*
111(46):16442–16447.

5. West SA, Griffin AS, Gardner A (2007) Evolutionary Explanations for Cooperation. *Curr Biol* 17(16):R661–R672.

6. Sachs JL, Mueller UG, Wilcox TP, Bull JJ (2004) The evolution of cooperation. *Q Rev Biol* 79(2):135–160.

7. Leigh EG (2010) The evolution of mutualism. *J Evol Biol* 23(12):2507–2528.

8. Bennett GM, Moran NA (2015) Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *Proc*
Natl Acad Sci 112(33):10169–10176.

9. Currie CR, et al. (2003) Ancient tripartite coevolution in the attine ant-microbe symbiosis. *Science* 299(5605):386–
8.

10. Mueller UG, et al. (2011) Evolution of cold-tolerant fungal symbionts permits winter fungiculture by leafcutter ants
at the northern frontier of a tropical ant-fungus symbiosis. *Proc Natl Acad Sci U S A* 108(10):4053–6.

11. Fisher RM, Henry LM, Cornwallis CK, Kiers ET, West SA (2017) The evolution of host-symbiont dependence. *Nat*
Commun 8:15973.

12. Sachs J, Simms E (2006) Pathways to mutualism breakdown. *Trends Ecol Evol* 21(10):585–592.

13. Kiers ET, Palmer TM, Ives AR, Bruno JF, Bronstein JL (2010) Mutualisms in a changing world: an evolutionary
perspective. *Ecol Lett* 13(12):1459–1474.

14. Lutzoni F, Pagel M, Reeb V (2001) Major fungal lineages are derived from lichen symbiotic ancestors. *Nature*
411(6840):937–940.

15. Distel DL, et al. (2017) Discovery of chemoautotrophic symbiosis in the giant shipworm *Kuphus polythalamia*
(Bivalvia: Teredinidae) extends wooden-steps theory. *Proc Natl Acad Sci* 114(18):E3652–E3658.

16. Pellmyr O, Leebens-Mack J, Huth CJ (1996) Non-mutualistic yucca moths and their evolutionary consequences.
Nature 380(6570):155–6.

17. Pellmyr O, Leebens-Mack J (2000) Reversal of Mutualism as a Mechanism for Adaptive Radiation in Yucca Moths.
Am Nat 156(S4):S62–S76.

18. Kawakita A, Mochizuki K, Kato M (2015) Reversal of mutualism in a leafflower-leafflower moth association: The
possible driving role of a third-party partner. *Biol J Linn Soc* 116(3):507–518.

19. Culley TM, Weller SG, Sakai AK (2002) The evolution of wind pollination in angiosperms. *Trends Ecol Evol*
17(8):361–369.

20. Whittall JB, Hodges S a (2007) Pollinator shifts drive increasingly long nectar spurs in columbine flowers. *Nature*
447(7145):706–709.

21. Parniske M (2008) Arbuscular mycorrhiza: the mother of plant root endosymbioses. *Nat Rev Microbiol* 6(10):763–
775.

22. Hoeksema JD, et al. (2010) A meta-analysis of context-dependency in plant response to inoculation with
mycorrhizal fungi. *Ecol Lett* 13(3):394–407.

23. van der Heijden MGA, Martin FM, Selosse M-A, Sanders IR (2015) Mycorrhizal ecology and evolution: the past,
the present, and the future. *New Phytol* 205(4):1406–1423.

24. Maherali H, Oberle B, Stevens PF, Cornwell WK, McGlinn DJ (2016) Mutualism Persistence and Abandonment
during the Evolution of the Mycorrhizal Symbiosis. *Am Nat* 188(5):E113–E125.

25. Zanne AE, et al. (2014) Three keys to the radiation of angiosperms into freezing environments. *Nature*
506(7486):89–92.

26.   Kattge J, et al. (2011) TRY - a global database of plant traits. *Glob Chang Biol* 17(9):2905–2935.

27.   Wang B, Qiu Y-L (2006) Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza*
440   16(5):299–363.

441   28.   Akhmetzhanova AA, et al. (2012) A rediscovered treasure: mycorrhizal intensity database for 3000 vascular plant
442   species across the former Soviet Union. *Ecology* 93(3):689–690.

443   29.   Werner GDA, Cornwell WK, Sprent JI, Kattge J, Kiers ET (2014) A single evolutionary innovation drives the deep
444   evolution of symbiotic N2-fixation in angiosperms. *Nat Commun* 5:4087.

445   30.   Ellison AM, Gotelli NJ (2003) Evolutionary ecology of carnivorous plants. *Trends Ecol Evol* 16(11):623–629.

446   31.   Neumann G, Martinoia E (2002) Cluster roots – an underground adaptation for survival in extreme environments.
447   *Trends Plant Sci* 7(4):162–167.

448   32.   Beaulieu JM, O'Meara BC, Donoghue MJ (2013) Identifying hidden rate changes in the evolution of a binary
449   morphological character: the evolution of plant habit in campanulid angiosperms. *Syst Biol* 62(5):725–37.

450   33.   Werner GDA, Cornwell WK, Cornelissen JHC, Kiers ET (2015) Evolutionary signals of symbiotic persistence in
451   the legume–rhizobia mutualism. *Proc Natl Acad Sci* 112(33):10262–10269.

452   34.   Martin F, Kohler A, Murat C, Veneault-Fourrey C, Hibbett DS (2016) Unearthing the roots of ectomycorrhizal
453   symbioses. *Nat Rev Microbiol*. doi:10.1038/nrmicro.2016.149.

454   35.   Merckx VS (2013) *Mycoheterotrophy* ed Merckx V (Springer New York, New York, NY) doi:10.1007/978-1-4614-
455   5209-6.

456   36.   Těšitel J (2016) Functional biology of parasitic plants⸱: a review. *Plant Ecol Evol* 149(1):5–20.

457   37.   Pagel M (1994) Detecting Correlated Evolution on Phylogenies: A General Method for the Comparative Analysis
458   of Discrete Characters. *Proc R Soc B Biol Sci* 255(1342):37–45.

459   38.   Delaux P-M, et al. (2014) Comparative Phylogenomics Uncovers the Impact of Symbiotic Associations on Host
460   Genome Evolution. *PLoS Genet* 10(7):e1004487.

461   39.   Bravo A, York T, Pumplin N, Mueller LA, Harrison MJ (2016) Genes conserved for arbuscular mycorrhizal
462   symbiosis identified through phylogenomics. *Nat Plants* 2(2):15208.

463   40.   Oliver KM, Degnan PH, Burke GR, Moran N a (2010) Facultative symbionts in aphids and the horizontal transfer
464   of ecologically important traits. *Annu Rev Entomol* 55:247–266.

465   41.   Weese DJ, Heath KD, Dentinger BTM, Lau J a. (2015) Long-term nitrogen addition causes the evolution of less-
466   cooperative mutualists. *Evolution (N Y)* 69(3):631–642.

467   42.   Gutiérrez-Valencia J, Chomicki G, Renner SS (2017) Recurrent breakdowns of mutualisms with ants in the
468   neotropical ant-plant genus Cecropia (Urticaceae). *Mol Phylogenet Evol* 111:196–205.

469   43.   Chomicki G, Renner SS (2017) Partner abundance controls mutualism stability and the pace of morphological
470   change over geologic time. *Proc Natl Acad Sci* 114(15):3951–3956.

471   44.   Sheldrake M, et al. (2017) A phosphorus threshold for mycoheterotrophic plants in tropical forests. *Proc R Soc B*
472   *Biol Sci* 284(1848):20162093.

473   45.   Treseder KK, et al. (2014) Evolutionary histories of soil fungi are reflected in their large-scale biogeography. *Ecol*
474   *Lett* 17(9):1086–1093.

475   46.   Read DJ (1996) The Structure and Function of the Ericoid Mycorrhizal Root. *Ann Bot* 77(4):365–374.

476   47.   Husnik F, et al. (2013) Horizontal gene transfer from diverse bacteria to an insect genome enables a tripartite nested
477   mealybug symbiosis. *Cell* 153(7):1567–78.

478   48.   Bennett GM, McCutcheon JP, MacDonald BR, Romanovicz D, Moran NA (2014) Differential genome evolution
479   between companion symbionts in an insect-bacterial symbiosis. *MBio* 5(5):e01697-14.

480   49.   Szabó G, et al. (2017) Convergent patterns in the evolution of mealybug symbioses involving different
481   intrabacterial symbionts. *ISME J* 11(3):715–726.

50.   Tisserant E, et al. (2013) Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant

symbiosis. *Proc Natl Acad Sci U S A* 110(50):20117–22.

51.   Almario J, et al. (2017) Root-associated fungal microbiota of nonmycorrhizal Arabis alpina and its contribution to
plant phosphorus nutrition. *Proc Natl Acad Sci* 114(44):E9403–E9412.

52.   Ghoul M, Griffin AS, West S a (2014) Toward and evolutinary definition of cheating. *Evolution (N Y)* 68(2):318–
331.

53.   Porter SS, Simms EL (2014) Selection for cheating across disparate environments in the legume-rhizobium
mutualism. *Ecol Lett* 17(9):1121–1129.

54.   Salem H, Onchuru TO, Bauer E, Kaltenpoth M (2015) Symbiont transmission entails the risk of parasite infection.
Biol Lett 11(12):20150840.

55.   Gordon BR, et al. (2016) Decoupled genomic elements and the evolution of partner quality in nitrogen-fixing
rhizobia. *Ecol Evol* 6(5):1317–1327.

56.   García-Guzmán G, Heil M (2014) Life histories of hosts and pathogens predict patterns in tropical fungal plant
diseases. *New Phytol* 201(4):1106–1120.

57.   Delaye L, García-Guzmán G, Heil M (2013) Endophytes versus biotrophic and necrotrophic pathogens—are fungal
lifestyles evolutionarily stable traits? *Fungal Divers* 60(1):125–135.

58.   Sachs JL, Skophammer RG, Regus JU (2011) Evolutionary transitions in bacterial symbiosis. *Proc Natl Acad Sci*
108(Supplement_2):10800–10807.

59.   Sachs JL, Skophammer RG, Bansal N, Stajich JE (2013) Evolutionary origins and diversification of proteobacterial
mutualists. *Proc R Soc B Biol Sci* 281(1775):20132146–20132146.

60.   Yang Z (2006) *Computational Molecular Evolution* (Oxford University Press, Oxford, UK).

61.   Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401(October 1999):877–884.

62.   Armstrong A (2015) Ecophysiology: Root-cluster control. *Nat Plants* 1(2):15009.

63.   Smith SE, Read DJ (2002) Orchid mycorrhizas. *Mycorrhizal Symbiosis* (Elsevier), pp 349–375.

Acknowledgements

The study has been supported by the TRY initiative on plant traits (http://www.trydb.org).
GDAW was funded by a Royal Society Newton International Fellowship and a Junior Research
Fellowship at Balliol College Oxford. ETK was funded by Netherlands Organisation for Scientific
Research Grants 836.10.001 and 864.10.005 and European Research Council ERC Grant Agreement
335542. We thank SURFsara (www.surf-sara.nl) for support in using the Lisa Computing Cluster and
Floortje Bouwkamp for providing the botanical illustrations in Fig. 1. The AM-alternative illustrations
in Figures 1 and 2 are based on figures in the public domain (CC0) with the exception of the
illustrations for ericoid mycorrhizae (courtesy Dr. David Midgley, CC-BY-SA) and the illustrations
for cluster roots (62) and orchid mycorrhizae (63) which were reprinted with permission from the
respective publishers.

Figure Legends

Figure 1: We explored the evolution of various alternatives to interacting with AM fungi, the ancestral
state of seed plants. Examples of the six most important alternatives we considered. The two columns
indicate two potential trajectories towards evolutionary breakdown of the plant-AM mutualism: switches

to other symbiotic partners (left) and abandonment of mutualism through the evolution of alternative

resource acquisition strategies (right). In both trajectories, the ancestral mutualism with AM fungi breaks

down. Coloured borders in this figure are the same as the coloured bars around the phylogeny in Figure 2

indicating where across global seed plants these traits are present.

Figure 2: Transition rates and ancestral state reconstruction of the dependent model of the evolution of

plant AM status and AM-alternatives. The four potential symbiotic states of a plant species are represented

by the transition matrix and by the same colours in the internal branches of the seed plant phylogeny.

Transition rates are expressed as number of transitions per 100 million years per lineage. The ancestral

state is AM presence with no AM-alternatives (green), red indicates the switch to one of the AM

alternatives (i.e. another mycorrhizal fungus, symbiotic N_2-fixation, parasitism, carnivory or cluster roots).

Coloured bands around the phylogeny (from inside to outside) indicate the presence or absence of (i) AM

interactions, (ii) other mycorrhizal fungi and (iii) alternative resource acquisition strategies. Key clades

that have lost AM fungal interactions and made an evolutionary switch to one of the AM alternatives are

indicated with schematic images of that alternative. Grey and white concentric circles within the

phylogeny indicate 50 million years.

Symbiont Switching

Mutualism Abandonment

Ancestral Mutualism

Arbuscular Mycorrhizal Fungi Alternatives

Arbuscular mycorrhizal (AM) fungi

Ectomycorrhizal (ECM) fungi

Carnivory

Orchid mycorrhizal (ORM) fungi

Cluster Roots

Bacterial symbiotic N_2-fixation

Parasitic Plants

Legend Bands

AM Interaction (Inner Band)
- ⬛ AM Present

Other symbionts (Middle Band)
- 🟪 Ectomycorrhizal
- 🟦 Orchid
- 🟩 Ericoid / Arbutoid
- 🟦 Symbiotic N₂-fixation

Alternative strategies (Outer Band)
- 🟥 Carnivorous Plants
- 🟧 Cluster Roots
- 🟩 Parasitic Plants

Proteaceae
Cyperaceae
Myrtaceae
Orchids
Fagales
Pines
Detariae
Orobanchaceae
Lentibulariaceae
Fabaceae
Santales
Drosera
Monotropoidea
Ericaceae

No AM No alternatives → *AM No alternatives*: **11.90**
AM No alternatives → *No AM No alternatives*: **1.54**
No AM No alternatives → *No AM Alternatives*: **0.04**
No AM Alternatives → *No AM No alternatives*: **0.50**
AM & Alternatives → *AM No alternatives*: **0.84**
AM No alternatives → *AM & Alternatives*: **0.22**
No AM Alternatives → *AM & Alternatives*: **0.63**
AM & Alternatives → *No AM Alternatives*: **1.52**