# Universal temporal rate of DNA replication origin firing: A balance between origin activation and passivation

**Jean-Michel Arbona**[1], **Arach Goldar**[2], **Olivier Hyrien**[3], **Alain Arneodo**[4], **Benjamin Audit**[1]

**\*For correspondence:**
benjamin.audit@ens-lyon.fr (BA)

[1]Univ Lyon, Ens de Lyon, Univ Claude Bernard Lyon 1, CNRS, Laboratoire de Physique, F-69342 Lyon, France.; [2]Ibitec-S, CEA, Gif-sur-Yvette, France.; [3]Institut de biologie de l'Ecole normale supérieure (IBENS), Ecole normale supérieure, CNRS, INSERM, PSL Research University, 75005 Paris, France.; [4]LOMA, Univ de Bordeaux, CNRS, UMR 5798, 351 Cours de le Libération, F-33405 Talence, France.

**Abstract**  The time-dependent rate $I(t)$ of origin firing per length of unreplicated DNA presents a universal bell shape in eukaryotes that has been interpreted as the result of a complex time-evolving interaction between origins and limiting firing factors. Here we show that a normal diffusion of replication fork components towards localized potential replication origins (*p-oris*) can more simply account for the $I(t)$ universal bell shape, as a consequence of a competition between the origin firing time and the time needed to replicate DNA separating two neighboring *p-oris*. We predict the $I(t)$ maximal value to be the product of the replication fork speed with the squared *p-ori* density. We show that this relation is robustly observed in simulations and in experimental data for several eukaryotes. Our work underlines that fork-component recycling and potential origins localization are sufficient spatial ingredients to explain the universality of DNA replication kinetics.

## Introduction

Eukaryotic DNA replication is a stochastic process (*Hyrien et al., 2013*; *Hawkins et al., 2013*; *Hyrien, 2016*). Prior to entering the S(ynthesis)-phase of the cell cycle, a number of DNA loci called potential origins (*p-oris*) are *licensed* for DNA replication initiation (*Machida et al., 2005*; *Hyrien et al., 2013*; *Hawkins et al., 2013*). During S-phase, in response to the presence of origin *firing* factors, pairs of replication *forks* performing bi-directional DNA synthesis will start from a subset of the *p-oris*, the active replication origins for that cell cycle (*Machida et al., 2005*; *Hyrien et al., 2013*; *Hawkins et al., 2013*). Note that the inactivation of *p-oris* by the passing of a replication fork called origin *passivation*, forbids origin firing in already replicated regions (*de Moura et al., 2010*; *Hyrien and Goldar, 2010*; *Yang et al., 2010*).

   The time-dependent rate of origin firing per length of unreplicated DNA, $I(t)$, is a fundamental parameter of DNA replication kinetics. $I(t)$ curves present a universal bell shape in eukaryotes (*Goldar et al., 2009*), increasing toward a maximum after mid-S-phase and decreasing to zero at the end of S-phase. An increasing $I(t)$ results in a tight dispersion of replication ending times, which provides a solution to the random completion problem (*Hyrien et al., 2003*; *Bechhoefer and Marshall, 2007*; *Yang and Bechhoefer, 2008*).

Models of replication in *Xenopus* embryo (*Goldar et al., 2008*; *Gauthier and Bechhoefer, 2009*) proposed that the initial $I(t)$ increase reflects the progressive import during S-phase of a limiting origin firing factor and its recycling after release upon forks merge. The $I(t)$ increase was also reproduced in a simulation of human genome replication timing that used a constant number of firing factors having an increasing reactivity through S-phase (*Gindin et al., 2014*). In these 3 models, an additional mechanism was required to explain the final $I(t)$ decrease by either a subdiffusive motion of the firing factor (*Gauthier and Bechhoefer, 2009*), a dependency of firing factors' affinity for *p-oris* on replication fork density (*Goldar et al., 2008*), or an inhomogeneous firing probability profile (*Gindin et al., 2014*). Here we show that when taking into account that *p-oris* are distributed at a finite number of localized sites then it is possible to reproduce the universal bell shape of the $I(t)$ curves without any additional hypotheses than recycling of fork factor components. $I(t)$ increases following an increase of fork mergers, each merger releasing a firing factor that was trapped on DNA. Then $I(t)$ decreases due to a competition between the time $t_c$ to fire an origin and the time $t_r$ to replicate DNA separating two neighboring *p-ori*. We will show that when $t_c$ becomes smaller than $t_r$, *p-ori* density over unreplicated DNA decreases, and so does $I(t)$. Modeling random localization of active origins in *Xenopus* embryo by assuming that every site is a (weak) *p-ori*, previous work implicitly assumed $t_r$ to be close to zero (*Goldar et al., 2008*; *Gauthier and Bechhoefer, 2009*) forbidding the observation of a decreasing $I(t)$. Licensing of a limited number of sites as *p-ori* thus appears to be a critical property contributing to the observed canceling of $I(t)$ at the end of S-phase in all studied eukaryotes.

## Results

### Emergence of a bell-shape $I(t)$

In our modeling of replication kinetics, a bimolecular reaction between a firing factor and a *p-ori* results in an origin firing event; then the diffusing element is trapped and travels with the replication forks until two converging forks merge (termination, Fig. 1 (a)). Under the assumption of a well-mixed system, for every time step $dt$, we consider each interaction between the $N_{FD}(t)$ free diffusing firing factors and the $N_{p\text{-}ori}(t)$ *p-oris* as potentially leading to a firing with a probability $k_{on}dt$. The resulting simulated firing rate per length of unreplicated DNA is then:

$$I_S(t) = \frac{N_{fired}(t, t+dt)}{L_{unrepDNA}(t)dt} \, , \qquad (1)$$

where $N_{fired}(t, t+dt)$ is the number of *p-oris* fired between times $t$ and $t+dt$, and $L_{unrepDNA}(t)$ is the length of unreplicated DNA a time $t$. Then we propagate the forks along the chromosome with a constant speed $v$, and if two forks meet, a free firing factor is released. Finally we simulate the chromosomes as 1D chains where prior to entering S-phase, the *p-oris* are precisely localized. For *Xenopus* embryo, the *p-ori* positions are randomly sampled, so that each simulated S-phase corresponds to a different positioning of the *p-oris*. We compare results obtained with periodic or uniform *p-ori* distributions. For *S. cerevisiae*, the *p-ori* positions, identical for each simulation, are taken from the OriDB database (*Siow et al., 2012*). As previously simulated in human (*Löb et al., 2016*), we model the entry in S-phase using an exponentially relaxed loading of the firing factors with a time scale shorter than the S-phase duration $T_{phase}$ (3 mins for *Xenopus* embryo, where $T_{phase} \sim 30$ mins, and 10 mins for *S. cerevisiae*, where $T_{phase} \sim 60$ mins). After the short loading time, the total number of firing factors $N_D^T$ is constant. As shown in Fig. 1 (b) (see also Fig. 2), the universal bell shape of the $I(t)$ curves (*Goldar et al., 2009*) spontaneously emerges from our model when going from weak to strong interaction, and decreasing the number of firing factors below the number of *p-oris*. The details of the firing factor loading dynamics do not affect the emergence of a bell shaped $I(t)$, even though it can modulate its precise shape, especially early in S-phase.

In a simple bimolecular context, the rate of origin firing is $i(t) = k_{on} N_{p\text{-}ori}(t) N_{FD}(t)$. The firing rate by element of unreplicated DNA is then given by

$$I(t) = k_{on} N_{FD}(t) \rho_{p\text{-}ori}(t) \, , \qquad (2)$$

**Figure 1.** (a) Sketch of the different steps of our modeling of replication initiation and propagation. (b) $I_S(t)$ (Eq. (1)) obtained from numerical simulations of one chromosome of length 3000 kb, with a fork speed $v = 0.6$ kb/min. The firing factors are loaded with a characteristic time of 3 mins. From blue to green to red the interaction is increased and the number of firing factors is decreased: blue ($k_{on} = 5 \times 10^{-5}$ min$^{-1}$, $N_D^T = 1000$, $\rho_0 = 0.3$ kb$^{-1}$), green ($k_{on} = 6 \times 10^{-4}$ min$^{-1}$, $N_D^T = 250$, $\rho_0 = 0.5$ kb$^{-1}$), red ($k_{on} = 6 \times 10^{-3}$ min$^{-1}$, $N_D^T = 165$, $\rho_0 = 0.28$ kb$^{-1}$)). (c) Corresponding normalized densities of *p-oris* (solid lines), and corresponding normalized numbers of free diffusing firing factors (dashed line): blue ($N_{FD}^* = 3360$), green ($N_{FD}^* = 280$), red ($N_{FD}^* = 28$); the light blue horizontal dashed line corresponds to the critical threshold value $N_{FD}(t) = N_{FD}^*$. (d) Corresponding number of passivated origins over the number of activated origins (solid lines). Corresponding histograms of replication time (dashed lines).

where $\rho_{p\text{-}ori}(t) = N_{p\text{-}ori}(t)/L_{unrepDNA}(t)$. In the case of a strong interaction and a limited number of firing factors, all the diffusing factors react rapidly after loading and $N_{FD}(t)$ is small (Fig. 1 (c), dashed curves). Then follows a stationary phase where as long as the number of *p-oris* is high (Fig. 1 (c), solid curves), once a diffusing factor is released by the encounter of two forks, it reacts rapidly, and so $N_{FD}(t)$ stays small. Then, when the rate of fork mergers increases due to the fact that there are as many active forks but a smaller length of unreplicated DNA, the number of free firing factors increases up to $N_D^T$ at the end of S-phase. As a consequence, the contribution of $N_{FD}(t)$ to $I(t)$ in Eq. (2) can only account for a monotonous increase during the S phase. For $I(t)$ to reach a maximum $I_{max}$ before the end of S-phase, we thus need that $\rho_{p\text{-}ori}(t)$ decreases in the late S-phase. This happens if the time to fire a *p-ori* is shorter than the time to replicate a typical distance between two neighboring *p-oris*. The characteristic time to fire a *p-ori* is $t_c = 1/k_{on}N_{FD}(t)$. The mean time for a fork to replicate DNA between two neighboring *p-oris* is $t_r = d(t)/v$, where $d(t)$ is the mean distance between unreplicated *p-oris* at time $t$. So the density of origins is constant as long as:

$$\frac{d(t)}{v} < \frac{1}{k_{on}N_{FD}(t)} \,, \tag{3}$$

or

$$N_{FD}(t) < N_{FD}^* = \frac{v}{k_{on}d(t)} \,. \tag{4}$$

Thus, at the beginning of the S-phase, $N_{FD}(t)$ is small, $\rho_{p\text{-}ori}(t)$ is constant (Fig. 1 (c), solid curves) and so $I_S(t)$ stays small. When $N_{FD}(t)$ starts increasing, as long as Eq. (4) stays valid, $I_S(t)$ keeps

increasing. When $N_{FD}(t)$ becomes too large and exceeds $N_{FD}^*$, then Eq. (4) is violated and the number of *p-oris* decreases at a higher rate than the length of unreplicated DNA, and $\rho_{p\text{-}ori}(t)$ decreases and goes to zero (Fig. 1 (c), red solid curve). As $N_{FD}(t)$ tends to $N_D^T$, $I_S(t)$ goes to zero, and its global behavior is a bell shape (Fig. 1 (b), red).

Let us note that if we decrease the interaction strength ($k_{on}$), then the critical $N_{FD}^*$ will increase beyond $N_D^T$ (Fig. 1 (c), dashed blue and green curves). $I_S(t)$ then monotonously increase to reach a plateau (Fig. 1 (b), green), or if we decrease further $k_{on}$, $I_S(t)$ present a very slow increasing behavior during the S-phase (Fig. 1 (b), blue). Now if we come back to strong interactions and increase the number of firing factors, almost all the *p-oris* are fired immediately and $I_S(t)$ drops to zero after firing the last *p-ori*.

Another way to look at the density of *p-oris* is to compute the ratio of the number of passivated origins by the number of activated origins (Fig. 1 (d)). After the initial loading of firing factors, this ratio is higher than one. For weak and moderate interactions (Fig. 1 (d), blue and green solid curves, respectively) this ratio stays bigger than one during all the S-phase, where $I_S(t)$ was shown to be monotonously increasing (Fig. 1 (b)). For a strong interaction (Fig. 1 (b), red solid curve), this ratio reaches a maximum and then decreases below one, at a time corresponding to the maximum observed in $I_S(t)$ (Fig. 1 (d), red solid curve). Hence, the maximum of $I(t)$ corresponds to a switch of the balance between origin passivation and activation, the latter becoming predominant in late S-phase. We have seen that up to this maximum $\rho_{p\text{-}ori}(t) \approx cte \approx \rho_0$, so $I_S(t) \approx k_{on}\rho_0 N_F(t)$. When $N_{FD}(t)$ reaches $N_{FD}^*$, then $I_S(t)$ reaches its maximum value:

$$I_{max} = k_{on}\rho_0 N_{FD}^* \approx \frac{\rho_0 v}{d(t)} \approx v\rho_0^2 \,, \tag{5}$$

where we have used the approximation $d(t) \approx d(0) = 1/\rho_0$ (which is exact for periodically distributed *p-oris*). $I_{max}$ can thus be predicted from two measurable parameters, providing a direct test of the model.

## Comparison with different eukaryotes

*Xenopus* embryo. Given the huge size of *Xenopus* embryo chromosomes, to make the simulations more easily tractable, we rescaled the size $L$ of the chromosomes, $k_{on}$ and $N_D^T$ to keep the duration of S-phase $T_{phase} \approx L/2vN_D^T$ and $I(t)$ (Eq. (2)) unchanged ($L \rightarrow \alpha L$, $N_D^T \rightarrow \alpha N_D^T$, $k_{on} \rightarrow k_{on}/\alpha$). In Fig. 2 (a) are reported the results of our simulations for a chromosome length $L = 3000$ kb. We see that a good agreement is obtained with experimental data (*Goldar et al., 2009*) when using either a uniform distribution of *p-oris* with a density $\rho_0 = 0.70$ kb$^{-1}$ and a number of firing factors $N_D^T = 187$, or a periodic distribution with $\rho_0 = 0.28$ kb$^{-1}$ and $N_D^T = 165$. A higher density of *p-oris* was needed for uniformly distributed *p-oris* where $d(t)$ (slightly) increases with time, than for periodically distributed *p-oris* where $d(t)$ fluctuates around a constant value $1/\rho_0$. The uniform distribution, which is the most natural to simulate *Xenopus* embryo replication, gives a density of activated origins of 0.17 kb$^{-1}$ in good agreement with DNA combing data analysis (*Herrick et al., 2002*) but twice lower than estimated from real time replication imaging of surface-immobilized DNA in a soluble *Xenopus* egg extract system (*Loveland et al., 2012*).

*S. cerevisiae*. To test the robustness of our minimal model with respect to the distribution of *p-oris*, we simulated the replication in *S. cerevisiae*, whose *p-oris* are known to be well positioned as reported in OriDB (*Siow et al., 2012*). 829 *p-oris* were experimentally identified and classified into three categories: confirmed origins (410), likely origins (216), and dubious origins (203). When comparing the results obtained with our model to the experimental $I(t)$ data (*Goldar et al., 2009*) (Fig. 2 (b)), we see that to obtain a good agreement we need to consider not only the confirmed origins but also the likely and the dubious origins. However in regard to the uncertainty in the value of the replication fork velocity and the possible experimental contribution of the *p-oris* in the rDNA part of chromosome 12 (not taken into account in our modeling), this conclusion needs to be confirmed in future experiments. It is to be noted that even if 829 *p-oris* are needed, on

**Figure 2.** (a) *Xenopus* embryo: Simulated $I_S(t)$ (Eq. (1)) for a chromosome of length $L = 3000$ kb and a uniform distribution of *p-oris* (blue: $v = 0.6$ kb/min, $k_{on} = 3{\times}10^{-3}$ min$^{-1}$, $N_D^T = 187$, $\rho_0 = 0.70$ kb$^{-1}$) or a periodic distribution of *p-oris* (red: $v = 0.6$ kb/min, $k_{on} = 6{\times}10^{-3}$ min$^{-1}$, $N_D^T = 165$, $\rho_0 = 0.28$ kb$^{-1}$); (red squares) 3D simulations with the same parameter values as for periodic *p-ori* distribution; (black) experimental $I(t)$: raw data obtained from *Goldar et al.* (*2009*) were binned in groups of 4 data points; the mean value and standard error of the mean of each bin were represented. (b) *S. cerevisiae*: Simulated $I_S(t)$ for the 16 chromosomes with the following parameter values: $v = 1.5$ kb/min, $N_D^T = 143$, $k_{on} = 3.6{\times}10^{-3}$ min$^{-1}$, when considering only confirmed origins (light blue), confirmed and likely origins (yellow) and confirmed, likely and dubious origins (purple); the horizontal dashed lines mark the corresponding predictions for $I_{max}$ (Eq. (5)); (purple squares) 3D simulations with the same parameter values considering confirmed, likely and dubious origins; (black) experimental $I(t)$ from *Goldar et al.* (*2009*). (c) *Eukaryotic organisms:* $I_{max}$ as a function of $v\rho_0^2$; (squares and bullets) simulations performed for regularly spaced origins (blue) and uniformly distributed origins (green) with two sets of parameter values: $L = 3000$ kb, $v = 0.6$ kb/min, $k_{on} = 1.2{\times}10^{-2}$ min$^{-1}$ and $N_D^T = 12$ (dashed line) or 165 (solid line); (black diamonds) experimental data points for *Xenopus* embryo, *S. cerevisiae*, *S. cerevisae* grown in Hydroxyurea (HU), *S. pombe*, *D. melanogaster*, human (see text and Table 1).

average only 352 origins have fired by the end of S-phase. For *S. cerevisiae* with well positioned
*p-oris*, we have checked the robustness of our results with respect to a stochastic number of firing
factors $N_D^T$ from cell to cell (Poisson distribution, *Iyer-Biswas et al.* (*2009*)). We confirmed the $I(t)$
bell shape with a robust duration of the S-phase of $58.6 \pm 4.3$ min as compared to $58.5 \pm 3.3$ min
obtained previously with a constant number of firing factors. Interestingly, in an experiment where
$T_{phase}$ was lengthened from 1 h to 16 h by adding hydroxyurea (HU) in yeast growth media, the
pattern of activation of replication origins was shown to be conserved (*Alvino et al., 2007*). HU
slows down the DNA synthesis to a rate of $\sim 50$ bp min$^{-1}$ corresponding to a 30 fold decrease
of the fork speed (*Sogo et al., 2002*). In our model with a constant number of firing factors,
$T_{phase} \sim 1/vN_D^T$: a two fold increase of the number $N_D^T$ of firing factors is sufficient to account
for the 16 fold increase of $T_{phase}$, which is thus mainly explained by the HU induced slowdown of
the replication forks. In a model where the increase of $I(t)$ results from the import of replication
factors, the import rate would need to be reduced by the presence of HU in proportion with
the lengthening of S-phase in order to maintain the pattern of origin activations. Extracting $I(t)$
from experimental replication data for cells grown in absence (HU$^-$) or presence (HU$^+$) (*Alvino*
*et al., 2007*), we estimated $I_{max}^{HU-} \sim 6.0$ Mb$^{-1}$min$^{-1}$ and $I_{max}^{HU+} \sim 0.24$ Mb$^{-1}$min$^{-1}$ for HU$^-$ and HU$^+$ cells,
respectively. The ratio $I_{max}^{HU-}/I_{max}^{HU+} \simeq 24.8 \sim v^{HU-}/v^{HU+}$ is quite consistent with the prediction of the

**Table 1.** Experimental data for various eukaryotic organisms with genome length $L$ ($Mb$), replication fork velocity $v$ (kb/min), number of $p$-oris ($N_{p\text{-}ori}(t{=}0)$), $\rho_0 = N_{p\text{-}ori}(t{=}0)/L$ (kb$^{-1}$) and $I_{max}$ (Mb$^{-1}$min$^{-1}$). All $I_{max}$ data are from *Goldar et al.* (*2009*), except for *S. cerevisiae* grown in presence or absence of hydroxyurea (HU) which were computed from the replication profile of *Alvino et al.* (*2007*). For *S. cerevisiae* and *S. pombe*, confirmed, likely, and dubious origins were taken into account. For *D. melanogaster*, $N_{p\text{-}ori}(t{=}0)$ was obtained from the same Kc cell type as the one used to estimate $I_{max}$. For *Xenopus* embryo, we used the experimental density of activated origins to estimate $N_{p\text{-}ori}(t{=}0)$ which is probably lower than the true number of $p$-oris. For human, we averaged the number of origins experimentally identified in K562 (62971) and in MCF7 (94195) cell lines.

| | $L$ | $v$ | $N_{p\text{-}ori}$ | $\rho_0$ | $I_{max}$ | Ref. |
|---|---|---|---|---|---|---|
| *S. cerevisiae* | 12.5 | 1.60 | 829 | 0.066 | 6.0 | *Sekedat et al.* (*2010*); *Siow et al.* (*2012*) |
| *S. cerevisiae* in presence of HU | 12.5 | 0.05 | 829 | 0.066 | 0.24 | *Alvino et al.* (*2007*). Same $N_{p\text{-}ori}$ and $\rho_0$ as *S. cerevisiae* in normal growth condition. |
| *S. pombe* | 12.5 | 2.80 | 741 | 0.059 | 10.0 | *Siow et al.* (*2012*); *Kaykov and Nurse* (*2015*) |
| *D. melanogaster* | 143.6 | 0.63 | 6184 | 0.043 | 0.5 | *Ananiev et al.* (*1977*); *Cayrou et al.* (*2011*) |
| *Xenopus* embryo | 2233.0 | 0.52 | 744333 | 0.333 | 70.0 | *Loveland et al.* (*2012*) |
| human | 6469.0 | 1.46 | 78000 | 0.012 | 0.3 | *Conti et al.* (*2007*); *Martin et al.* (*2011*) |

scaling law (Eq. (5)) for a constant density of *p-oris*.

*D. melanogaster* and human. We gathered from the literature experimental estimates of $I_{max}$, $\rho_0$
and $v$ for different eukaryotic organisms (Table 1). As shown in Fig. 2 (c), when plotting $I_{max}$ vs $v\rho_0^2$,
all the experimental data points remarkably follow the diagonal trend indicating the validity of the
scaling law (Eq. (5)) for all considered eukaryotes. We performed two series of simulations for fixed
values of parameters $k_o$, $N_D^T$ and $v$ and decreasing values of $\rho_0$ with both periodic distribution (blue)
and uniform (green) distributions of *p-oris* (Fig. 2 (c)). The first set of parameters was chosen to cover
high $I_{max}$ values similar the one observed for *Xenopus* embryo (bullets, solid lines). When decreasing
$\rho_0$, the number of firing factors becomes too large and $I(t)$ does no more present a maximum.
We thus decreased the value of $N_D^T$ keeping all other parameters constant (boxes, dashed line) to
explore smaller values of $I_{max}$ in the range of those observed for human and *D. melanogaster*. We
can observe that experimental data points' deviation from Eq. (5) is smaller than the deviation due
to specific *p-oris* distributions.

## Discussion

To summarize, we have shown that within the framework of 1D nucleation and growth models of
DNA replication kinetics (*Herrick et al., 2002*; *Jun and Bechhoefer, 2005*), the sufficient conditions
to obtain a universal bell shaped $I(t)$ as observed in eukaryotes are a strong bimolecular reaction
between localized *p-oris* and limiting origin firing factors that travel with replication forks and are
released at termination. Under these conditions, the density of *p-oris* naturally decreases by the
end of the S-phase and so does $I_S(t)$. Previous models in *Xenopus* embryo (*Goldar et al., 2008*;
*Gauthier and Bechhoefer, 2009*) assumed that all sites contained a *p-ori* implying that the time $t_r$ to
replicate DNA between two neighboring *p-oris* was close to zero. This clarifies why they needed
some additional mechanisms to explain the final decrease of the firing rate. Moreover our model
predicts that the maximum value for $I(t)$ is intimately related to the density of *p-oris* and the fork
speed (Eq. (5)), and we have shown that without free parameter, this relationship holds for 5 species
up to a 300 fold difference of $I_{max}$ and $v\rho_0^2$ (Table 1, Fig. 2 (c)).
In contrast with models where replication kinetics is explained by properties specific to each
*p-oris* (*Bechhoefer and Rhind, 2012*), our model assumes that all *p-oris* are governed by the same
rule of initiation resulting from physicochemically realistic particulars of their interaction with
limiting replication firing factors. To confirm this simple physical basis of our modeling, we used
molecular dynamics rules as previously developed for *S. cerevisiae* (*Arbona et al., 2017*) to simulate
S-phase dynamics of chromosomes confined in a spherical nucleus. We added firing factors that
are free to diffuse in the covolume left by the chain and that can bind to proximal *p-oris* to initiate

replication, move along the chromosomes with the replication forks and be released when two fork merges. As shown in Fig. 2 (a, b) for *Xenopus* embryo and *S. cerevisiae*, results confirmed the physical relevance of our minimal modeling and the validity of its predictions when the 3D diffusion of the firing factors is explicitly taken into account. This opens new perspectives for understanding correlations between firing events along chromosomes that could result in part from the spatial transport of firing factors. For example in *S. cerevisiae* (*Knott et al., 2012*) and in *S. pombe* (*Kaykov and Nurse, 2015*), a higher firing rate has been reported near origins that have just fired (but see *Yang et al.* (*2010*)). In mammals, megabase chromosomal regions of synchronous firing were first observed long ago (*Huberman and Riggs, 1968*; *Hyrien, 2016*). Recently, profiling of replication fork directionality obtained by Okazaki fragment sequencing have suggested that early firing origins located at the border of Topologically Associating Domains (TADs) trigger a cascade of secondary initiation events propagating through the TAD (*Petryk et al., 2016*). Early and late replicating domains were associated with nuclear compartments of open and closed chromatin (*Ryba et al., 2010*; *Boulos et al., 2015*; *Goldar et al., 2016*; *Hyrien, 2016*). In human, replication timing U-domains (0.1-3 Mb) were shown to correlate with chromosome structural domains (*Baker et al., 2012*; *Moindrot et al., 2012*; *Pope et al., 2014*) and chromatin loops (*Boulos et al., 2013*, *2014*).

Understanding to which extent spatio-temporal correlations of the replication program can be explained by the diffusion of firing factors in the tertiary chromatin structure specific to each eukaryotic organism is a challenging issue for future work.

We thank F. Argoul for helpful discussions. This work was supported by Institut National du Cancer (PLBIO16-302), Fondation pour la Recherche Médicale (DEI20151234404) and Agence National de la Recherche (ANR-15-CE12-0011-01). BA acknowledges support from Science and Technology Commission of Shanghai Municipality (15520711500) and Joint Research Institute for Science and Society (JoRISS). We gratefully acknowledge support from the PSMN (Pôle Scientifique de Modélisation Numérique) of the ENS de Lyon for the computing resources. We thank BioSyL Federation and Ecofect LabEx (ANR-11-LABX-0048) for inspiring scientific events.

## References

**Alvino GM**, Collingwood D, Murphy JM, Delrow J, Brewer BJ, Raghuraman MK. Replication in Hydroxyurea: It's a Matter of Time. Molecular and Cellular Biology. 2007 Sep; 27(18):6396–6406. http://mcb.asm.org/content/27/18/6396, doi: 10.1128/MCB.00719-07.

**Ananiev EV**, Polukarova LG, Yurov YB. Replication of chromosomal DNA in diploid *Drosophila melanogaster* cells cultured in vitro. Chromosoma. 1977 Feb; 59:259–272.

**Arbona JM**, Herbert S, Fabre E, Zimmer C. Inferring the physical properties of yeast chromatin through Bayesian analysis of whole nucleus simulations. Genome Biol. 2017; 18(1):81.

**Baker A**, Audit B, Chen CL, Moindrot B, Leleu A, Guilbaud G, Rappailles A, Vaillant C, Goldar A, Mongelard F, d'Aubenton Carafa Y, Hyrien O, Thermes C, Arneodo A. Replication fork polarity gradients revealed by megabase-sized U-shaped replication timing domains in human cell lines. PLoS Comput Biol. 2012; 8(4):e1002443. http://dx.doi.org/10.1371/journal.pcbi.1002443, doi: 10.1371/journal.pcbi.1002443.

**Bechhoefer J**, Rhind N. Replication timing and its emergence from stochastic processes. Trends Genet. 2012; 28(8):374–381. doi: 10.1016/j.tig.2012.03.011.

**Bechhoefer J**, Marshall B. How Xenopus laevis replicates DNA reliably even though its origins of replication are located and initiated stochastically. Phys Rev Lett. 2007 Mar; 98:098105. doi: 10.1103/PhysRevLett.98.098105.

**Boulos RE**, Arneodo A, Jensen P, Audit B. Revealing Long-Range Interconnected Hubs in Human Chromatin Interaction Data Using Graph Theory. Phys Rev Lett. 2013 Sep; 111(11):118102. https://link.aps.org/doi/10.1103/PhysRevLett.111.118102, doi: 10.1103/PhysRevLett.111.118102.

**Boulos RE**, Drillon G, Argoul F, Arneodo A, Audit B. Structural organization of human replication timing domains. FEBS Lett. 2015 Oct; 589(20 Pt A):2944–2957. http://dx.doi.org/10.1016/j.febslet.2015.04.015, doi: 10.1016/j.febslet.2015.04.015.

**Boulos RE**, Julienne H, Baker A, Chen CL, Petryk N, Kahli M, Yves d'Aubenton-Carafa, Goldar A, Jensen P, Hyrien O, Thermes C, Arneodo A, Audit B. From the chromatin interaction network to the organization of the human genome into replication N/U-domains. New J Phys. 2014; 16(11):115014. http://stacks.iop.org/1367-2630/16/i=11/a=115014, doi: 10.1088/1367-2630/16/11/115014.

**Cayrou C**, Coulombe P, Vigneron A, Stanojcic S, Ganier O, Peiffer I, Rivals E, Puy A, Laurent-Chabalier S, Desprat R, Méchali M. Genome-scale analysis of metazoan replication origins reveals their organization in specific but flexible sites defined by conserved features. Genome Res. 2011 Sep; 21(9):1438–1449. http://dx.doi.org/10.1101/gr.121830.111, doi: 10.1101/gr.121830.111.

**Conti C**, Sacca B, Herrick J, Lalou C, Pommier Y, Bensimon A. Replication fork velocities at adjacent replication origins are coordinately modified during DNA replication in human cells. Mol Biol Cell. 2007 Aug; 18(8):3059–3067. http://dx.doi.org/10.1091/mbc.E06-08-0689, doi: 10.1091/mbc.E06-08-0689.

**Gauthier MG**, Bechhoefer J. Control of DNA replication by anomalous reaction-diffusion kinetics. Phys Rev Lett. 2009 Apr; 102:158104. doi: 10.1103/PhysRevLett.102.158104.

**Gindin Y**, Valenzuela MS, Aladjem MI, Meltzer PS, Bilke S. A chromatin structure-based model accurately predicts DNA replication timing in human cells. Mol Syst Biol. 2014; 10:722.

**Goldar A**, Arneodo A, Audit B, Argoul F, Rappailles A, Guilbaud G, Petryk N, Kahli M, Hyrien O. Deciphering DNA replication dynamics in eukaryotic cell populations in relation with their averaged chromatin conformations. Sci Rep. 2016 Mar; 6:22469. http://www.nature.com/srep/2016/160303/srep22469/full/srep22469.html, doi: 10.1038/srep22469.

**Goldar A**, Labit H, Marheineke K, Hyrien O. A dynamic stochastic model for DNA replication initiation in early embryos. PLoS One. 2008; 3(8):e2919. http://dx.doi.org/10.1371/journal.pone.0002919, doi: 10.1371/journal.pone.0002919.

**Goldar A**, Marsolier-Kergoat MC, Hyrien O. Universal temporal profile of replication origin activation in eukaryotes. PLoS One. 2009; 4(6):e5899. http://dx.doi.org/10.1371/journal.pone.0005899, doi: 10.1371/journal.pone.0005899.

**Hawkins M**, Retkute R, Müller CA, Saner N, Tanaka TU, de Moura APS, Nieduszynski CA. High-resolution replication profiles define the stochastic nature of genome replication initiation and termination. Cell Rep. 2013 Nov; 5:1132–1141. doi: 10.1016/j.celrep.2013.10.014.

**Herrick J**, Jun S, Bechhoefer J, Bensimon A. Kinetic model of DNA replication in eukaryotic organisms. J Mol Biol. 2002 Jul; 320:741–750.

**Huberman JA**, Riggs AD. On the mechanism of DNA replication in mammalian chromosomes. J Mol Biol. 1968 Mar; 32(2):327–341. http://www.sciencedirect.com/science/article/pii/0022283668900132, doi: 10.1016/0022-2836(68)90013-2.

**Hyrien O**. Up and Down the Slope: Replication Timing and Fork Directionality Gradients in Eukaryotic Genomes. In: Kaplan DL, editor. *The Initiation of DNA Replication in Eukaryotes* Switzerland: Springer International Publishing; 2016.p. 65–85.

**Hyrien O**, Goldar A. Mathematical modelling of eukaryotic DNA replication. Chromosome Res. 2010; 18(1):147–161.

**Hyrien O**, Marheineke K, Goldar A. Paradoxes of eukaryotic DNA replication: MCM proteins and the random completion problem. BioEssays. 2003 Feb; 25(2):116–125. http://onlinelibrary.wiley.com/doi/10.1002/bies.10208/abstract, doi: 10.1002/bies.10208.

**Hyrien O**, Rappailles A, Guilbaud G, Baker A, Chen CL, Goldar A, Petryk N, Kahli M, Ma E, d'Aubenton Carafa Y, Audit B, Thermes C, Arneodo A. From Simple Bacterial and Archaeal Replicons to Replication N/U-Domains. J Mol Biol. 2013 Nov; 425(23):4673–4689. http://www.sciencedirect.com/science/article/pii/S0022283613006050, doi: 10.1016/j.jmb.2013.09.021.

**Iyer-Biswas S**, Hayot F, Jayaprakash C. Stochasticity of gene products from transcriptional pulsing. Phys Rev E. 2009 Mar; 79(3):031911. https://link.aps.org/doi/10.1103/PhysRevE.79.031911, doi: 10.1103/PhysRevE.79.031911.

**Jun S**, Bechhoefer J. Nucleation and growth in one dimension. II. Application to DNA replication kinetics. Phys Rev E. 2005; 71:011909. doi: 10.1103/PhysRevE.71.011909.

**Kaykov A**, Nurse P. The spatial and temporal organization of origin firing during the S-phase of fission yeast.
Genome Res. 2015 Mar; 25:391–401. doi: 10.1101/gr.180372.114.

**Knott SRV**, Peace JM, Ostrow AZ, Gan Y, Rex AE, Viggiani CJ, Tavaré S, Aparicio OM. Forkhead transcription
factors establish origin timing and long-range clustering in *S. cerevisiae*. Cell. 2012 Jan; 148:99–111. doi:
10.1016/j.cell.2011.12.012.

**Löb D**, Lengert N, Chagin VO, Reinhart M, Casas-Delucchi CS, Cardoso MC, Drossel B. 3D replicon distributions
arise from stochastic initiation and domino-like DNA replication progression. Nat Commun. 2016 Apr; 7:11207.
doi: 10.1038/ncomms11207.

**Loveland AB**, Habuchi S, Walter JC, van Oijen AM. A general approach to break the concentration barrier in
single-molecule imaging. Nat Methods. 2012 Oct; 9:987–992. doi: 10.1038/nmeth.2174.

**Machida YJ**, Hamlin JL, Dutta A. Right place, right time, and only once: replication initiation in metazoans. Cell.
2005 Oct; 123(1):13–24. http://dx.doi.org/10.1016/j.cell.2005.09.019, doi: 10.1016/j.cell.2005.09.019.

**Martin MM**, Ryan M, Kim R, Zakas AL, Fu H, Lin CM, Reinhold WC, Davis SR, Bilke S, Liu H, Doroshow JH, Reimers
MA, Valenzuela MS, Pommier Y, Meltzer PS, Aladjem MI. Genome-wide depletion of replication initiation
events in highly transcribed regions. Genome Res. 2011 Nov; 21:1822–1832. doi: 10.1101/gr.124644.111.

**Moindrot B**, Audit B, Klous P, Baker A, Thermes C, de Laat W, Bouvet P, Mongelard F, Arneodo A.
3D chromatin conformation correlates with replication timing and is conserved in resting cells. Nu-
cleic Acids Res. 2012 Oct; 40(19):9470–9481. https://academic.oup.com/nar/article/40/19/9470/2414837/
3D-chromatin-conformation-correlates-with, doi: 10.1093/nar/gks736.

**de Moura AP**, Retkute R, Hawkins M, Nieduszynski CA. Mathematical modelling of whole chromosome replica-
tion. Nucleic Acids Res. 2010; 38(17):5623–5633.

**Petryk N**, Kahli M, d'Aubenton Carafa Y, Jaszczyszyn Y, Shen Y, Silvain M, Thermes C, Chen CL, Hyrien O.
Replication landscape of the human genome. Nat Commun. 2016 Jan; 7:10208. http://www.nature.com/
ncomms/2016/160111/ncomms10208/full/ncomms10208.html, doi: 10.1038/ncomms10208.

**Pope BD**, Ryba T, Dileep V, Yue F, Wu W, Denas O, Vera DL, Wang Y, Hansen RS, Canfield TK, Thurman RE,
Cheng Y, Gülsoy G, Dennis JH, Snyder MP, Stamatoyannopoulos JA, Taylor J, Hardison RC, Kahveci T, Ren B,
et al. Topologically-associating domains are stable units of replication-timing regulation. Nature. 2014 Nov;
515(7527):402–405. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4251741/, doi: 10.1038/nature13986.

**Ryba T**, Hiratani I, Lu J, Itoh M, Kulik M, Zhang J, Schulz TC, Robins AJ, Dalton S, Gilbert DM. Evolutionarily
conserved replication timing profiles predict long-range chromatin interactions and distinguish closely
related cell types. Genome Res. 2010 Jun; 20(6):761–770. http://dx.doi.org/10.1101/gr.099655.109, doi:
10.1101/gr.099655.109.

**Sekedat MD**, Fenyö D, Rogers RS, Tackett AJ, Aitchison JD, Chait BT. GINS motion reveals replication fork progres-
sion is remarkably uniform throughout the yeast genome. Mol Syst Biol. 2010; 6:353. doi: 10.1038/msb.2010.8.

**Siow CC**, Nieduszynska SR, Müller CA, Nieduszynski CA. OriDB, the DNA replication origin database updated
and extended. Nucleic Acids Res. 2012 Jan; 40:D682–D686. doi: 10.1093/nar/gkr1091.

**Sogo JM**, Lopes M, Foiani M. Fork reversal and ssDNA accumulation at stalled replication forks owing to
checkpoint defects. Science. 2002; 297(5581):599–602. doi: 10.1126/science.1074023.

**Yang SCH**, Bechhoefer J. How Xenopus laevis embryos replicate reliably: investigating the random-completion
problem. Phys Rev E. 2008; 78(4):041917.

**Yang SCH**, Rhind N, Bechhoefer J. Modeling genome-wide replication kinetics reveals a mechanism for regulation
of replication timing. Mol Syst Biol. 2010; 6(1):404.