1

# Is phylogenetic diversity a surrogate for functional diversity across clades and space?

Florent Mazel[1,2,3,*], Matthew W. Pennell[3,4], Marc Cadotte[5,6,$], Sandra Diaz[7,$], Giulio Valentino

Dalla Riva[8,$], Richard Grenyer[9,$], Fabien Leprieur[10], Arne O. Mooers[1,$], David Mouillot[10,11,$],

Caroline M. Tucker[12,$], and William D. Pearse[13]

[1] Department of Biological Sciences, Simon Fraser University, Burnaby, BC, Canada

[2] Department of Botany, University of British Columbia, Vancouver, BC V6T 1Z4, Canada;

[3] Biodiversity Research Centre, University of British Columbia, Vancouver, BC V6T 1Z4, Canada;

[4] Department of Zoology, University of British Columbia, Vancouver, BC V6T 1Z4, Canada;

[5] Biological Sciences, University of Toronto-Scarborough, Scarborough M1C 1A4, Canada

[6] Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

[7] Instituto Multidisciplinario de Biología Vegetal. FECFyN - Universidad Nacional de Córdoba.

Casilla de Correo 495, 5000 Córdoba. Argentina

[8] Department of Statistics, University of British Columbia, Vancouver, BC V6T 1Z4, Canada;

[9] School of Geography and the Environment, University of Oxford, Oxford OX1 3QY, U.K.

[10] Marine Biodiversity, Exploitation, and Conservation (MARBEC), UMR 9190, Université

Montpellier, Montpellier, France

[11] Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook

University, Townsville, QLD, Australia

[12] Department of Biology, University of North Carolina-Chapel Hill

[13] Ecology Center and Department of Biology, Utah State University, Logan, Utah

* Corresponding Author; flo.mazel@gmail.com

$ Authors listed alphabetically

Keywords: conservation, EDGE, evolutionary distinct, trait diversity, evolutionary diversity

**Abstract**

In the face of limited funding and widespread threats to biodiversity, conserving the widest possible variety of biological traits (functional diversity, FD) is a reasonable prioritization objective. Because species traits are often similar among closely related species (phylogenetic signal), many researchers have advocated for a "phylogenetic gambit": maximizing phylogenetic diversity (PD) should indirectly capture FD. To our knowledge, this gambit has not been subject to a focused empirical test. Here we use data from >15,000 vertebrate species to empirically test it. We delineate >10,000 species pools and test whether prioritizing the most phylogenetically diverse set of species results in more or less FD relative to a random choice. We find that, across species pools, maximizing PD results in an average gain of 18% of FD relative to a random choice, suggesting that PD is a sound conservation prioritization strategy. However, this averaged gain hides important variability:  for 10% of the species pools, maximizing PD can capture less FD than an averaged random scheme because of recent trait divergence and/or very strong trait conservatism.  In addition, within a species pool, many random sets of species actually yield more FD than the PD-maximized selection – on average 36% of the time per pool. If the traits we used are representative of traits we wish to conserve, our results suggest that conservation initiatives focusing on PD will, on average, capture more FD than a random strategy, but this gain will not systematically yield more FD than random and thus can be considered risky.

We are in the midst of a period of heightened biological extinction, with rates several orders of

magnitude higher than background rates estimated from the fossil record [1–3]. In addition to

having potentially widespread consequences for the functioning of ecosystems and the

provisioning of valuable ecosystem services, this situation poses a huge moral challenge [4–8].

And, to the extent that resources for conservation actions remain limited, agonizing choices as

to which species most warrant attention become necessary [9,10]. To keep humanity's options

open, and our common legacy as rich as possible, it is widely argued that we should seek to

maximize the biological diversity of forms and functions [6–12] . The biological diversity of

forms and functions can be measured as functional diversity [FD] (see methods). However, in

practice, it is challenging to prioritize species on the basis of FD: we have imperfect knowledge

about which, and how many traits and functions are important in a given context, how these

traits and functions vary among species and across space, and how the importance of traits may

change in the future [13]. Many researchers have therefore advocated for a "phylogenetic

gambit"; that is, if species traits reflect their shared evolutionary history, then the pattern of

that evolutionary history –their phylogeny– should serve as a useful stand-in for unmeasured

and unmeasurable traits [9,14,15]. The phylogenetic gambit implies that maximizing

phylogenetic diversity (PD), i.e. the breadth of evolutionary history, will ensure that a wide

variety of forms and functions are present within a species set [14–17].

Following this logic, PD has formed the basis of global conservation schemes, notably the EDGE

program [18], has been used by restoration biologists [19], and has been widely embraced by

researchers across the biodiversity sciences [20–23]. Despite this enthusiasm, the critical

question of whether maximizing PD will actually capture more FD than prioritization schemes

that ignore phylogeny has, to our knowledge, never been empirically tested [16]. While it may

seem obvious that sampling species across the tree of life will capture high amounts of FD, a

recent theoretical study demonstrated that PD could be a poor surrogate for FD and, in some

scenarios, prioritizing species on the basis of PD could actually lead to capture *less* FD than if

species were simply selected at random [16].

We clarify what our goals are in testing the utility of PD to capture FD.  First, we take as given

that maximizing PD is not the overarching goal *per se* of PD-maximization schemes, but rather

that a PD maximization strategy is valued for its ability to capture more FD compared to a

strategy that ignores phylogeny. Second, asking whether PD maximization captures more FD

than a random choice is fundamentally distinct (and a lower bar) from asking whether

maximizing PD also *maximizes* FD [e.g. 15,19–21,23,24]. Finally, it is important to note that we

are selecting *species* sets to maximize PD or FD within a region. While this is a simplification, as

conservation actions often aim to select sets of *areas* (e.g. in reserve design), the only global

phylogenetically-informed conservation initiative is species-centered (EDGE; Isaac et al. 2007).

More fundamentally, the framework we use here allows us to directly test the fundamental

phylogenetic gambit at the heart of all PD-based conservation [16]. Critically, the question we

raise has been shown to be distinct from asking whether traits have phylogenetic signal

(whether closely related species tend to share similar sets of traits), since  PD can be a poor

surrogate for FD *even if* traits exhibit phylogenetic signal [16].

This points to the need for empirical tests of whether —within a given species pool— sets of

species selected to maximize PD actually contain more FD than sets of species selected without

regard to evolutionary relatedness. We evaluate the PD~FD relationship for different species

pools (taxonomic families and geographical assemblages, *i.e.*, sets of species co-occurring at a

given scale) using a large global dataset including trait, phylogenetic, and geographic range data

for 4,616 species of mammals, 9,993 species of birds, and 1,5036 species of tropical fish.

Specifically, we measure FD as functional richness (see methods) and compute, for any given

species pool, an estimate of surrogacy ($S_{PD\_FD}$, [26,27], Figure 1). $S_{PD\_FD}$ represents the amount

of FD sampled by the set of species chosen to maximize PD, relative to the FD sampled by

optimal set of species selected to maximize FD directly, with both components controlled for

the expected FD from a random species set of the same size. $S_{PD\_FD}$ will be positive if the

averaged PD-maximized set contains more FD than the averaged random set, and negative if

not. $S_{PD\_FD}$ will equal 100% if the PD-maximization strategy is optimal (*i.e.* to maximize FD). We

integrate $S_{PD\_FD}$ for each species pool across all deciles of species richness (Eqn. 1) but because

they are many sets of species that can maximize PD or than can be chosen at random, we

computed $S_{PD\_FD}$ based on the averaged FD over 1000 PD-maximized sets and 1000 random sets

[16].

We find that selecting the most phylogenetically diverse sets of species within a given

taxonomic family or within a given geographical location (large grid-cells across the globe)

captures, on average, 18% more FD than that of randomly chosen species (i.e. $S_{PD\_FD}$ = 18%, SD

+/- 6.5% across pools, see Figure 1). Although the surrogacy is generally positive, there was

variation across species pools. For example, the surrogacy of PD varies widely from a minimum

of -85% to a maximum of 92%, meaning that selecting the most phylogenetically diverse sets of

taxa can capture either 85% less (or 92% more) FD than that of randomly chosen taxa (Fig. 2-3

and Fig. S1-2). However, in 88% of the species pools, choosing sets of species according to PD

captured more FD than would be expected at random (*i.e.*, surrogacy values > 0 in 88% of the

cases, see Fig. 2-3). This suggest that, on average, maximizing PD is a sound strategy to capture

FD.

However, even if in the majority cases maximizing PD does, on average, better than an

averaged random selection, this does not capture the *reliability* of its performance. The PD-

maximization and the random selection strategies exhibit variation: simply by chance, random

selection of species can capture very high (or, conversely, very low) FD, and the same may be

true (to a previously unstudied degree) for PD. The extent of this variation is important: if it is

less than the average difference, PD-maximization is a reliable strategy as it will always yield

more FD, but if it does not, then PD-maximization could be unreliable for individual

conservation interventions. To contrast these two situations, we measured the fraction of times

that, within each species pool, the PD-maximization strategy yielded more FD than random

selection (see methods). PD-based selection was the best choice in 64% of cases (SD across

species pool=9%, see Supplementary Table 1 and Fig. S3), making it the better strategy but not

a perfectly reliable one. Thus, while the PD-maximization strategy has a consistent positive

effect (i.e. the average PD-maximization strategy yields more FD than the average random

strategy), its effect is weak (i.e. the PD-maximization strategy still yields less FD than the

random strategy in 36% of the trials within a species pool).

We next explored the drivers of surrogacies values across species pools. Surrogacy of PD

appears to weaken as the species pool richness increases (on average, Spearman Rho between

absolute surrogacies and species richness = -.15), most clearly seen in the tropics and in

species-rich families such as the Muridae (rats mice and allies) and Columbidae (pigeons and

allies) (Fig. 2-3). This is likely because our measure of FD (see Methods) rapidly saturates as the

number of selected species increases and species from these large pools harbor high functional

redundancy, such that a random prioritization scheme performs relatively well, or at least no

worse than other strategies (Fig. S4). In contrast, FD can be greatly increased by prioritization of

species using PD from species poor assemblages or clades [see also 28]. This is particularly the

case in spatial assemblages containing multiple taxonomic orders, which are both

phylogenetically and ecologically divergent from one another. Interestingly, the PD-FD

relationship was not consistent across *taxonomic* scale: we found that, in contrast to patterns

at the family level, for certain mammalian and avian orders (which are older than the families

described above), using PD to select species is much worse for capturing FD than choosing

species at random (see, for example, the Afrosoricidae, Chiroptera, and Charadriiformes in Fig.

S5).

We explored whether it is possible to explain this variability within- and between-

datasets, and in particular, why for some assemblages/clades, a PD-prioritization strategy fails

to capture more FD than random choice. It is often implicitly assumed that phylogenetic signal

(i.e. the degree to which closely related species tend to harbor similar sets of traits) can be used

to evaluate the effectiveness of PD as a surrogate for FD [5,15–17]. Surprisingly perhaps, the

value of PD as a surrogate for FD was only weakly correlated with the phylogenetic signal of the

underlying traits (Fig. S6-7, on average Spearman Rho = 0.17). Similarly, tree imbalance, which

is known to affect surrogacy in simulations [16], did not explain surrogacy in these empirical

data (Fig. S6-7).

For mammals, regions where PD did worse than random were located in the Sahara,

south western Patagonia, southern Africa including parts of Madagascar, and New Guinea

(Figure 2). These latter two in particular are of concern, since they are global conservation

priorities on account of species endemism and habitat loss. We suggest two historical reasons

for such idiosyncratic poor performance of PD. First, there is a tendency for a large carnivore

species, either a top predator (*e.g.*, cheetahs in the Sahara or foxes in Patagonia) or a large

scavenger (*e.g.*, the hyena in South Africa) to co-occur with a close relative with distinct traits in

these areas (*e.g.*, a desert cat with the cheetah or the aardwolf with the hyena, see Fig. S8).

Only one of these closely-related species will tend to be selected under prioritization schemes

that maximize PD, thus reducing the volume of the convex hull on average when the

functionally distinct one is not selected (the large predator or scavenger). This seems also to

drive the low surrogacy of PD in Charadriiformes (especially *Larus* and *Sterna*; see Figure S8).

Second, lineages in which traits evolve very slowly will contribute little to FD, even over long

periods of time (branch lengths) that contribute greatly to PD. For example, in New Guinea

many co-occurring bats with similar traits diverged long ago, such that they are always selected

in the PD maximizing set, but do not add much to the convex hull, resulting in a poor surrogacy

of PD for FD. Such strong ecological niche conservatism is common in mammals [29], e.g. in the

Geomyidae: two basal branches of the Geomyidae tree harbor very similar traits (species

descending from these branches are actually grouped in the same genus *Thomomys)* while

being distantly related in the phylogenies we used (Fig. S8).  As such, they will be selected in all

PD maximizing sets, but will not contribute greatly to FD.

In summary, while in specific cases maximizing PD actually captures less FD than a

random set, in the majority of cases PD performs well (at least, better than random) as a

surrogate (in 88% of the species pool sets the mean surrogacy value ≥0). This represents an

important and necessary test of the motivations of conservation planning activities that

incorporate PD. However, we simplistically and implicitly assume that chosen species will either

be saved or will go extinct and we have not linked our various scenarios to any particular policy

position or conservation objective other than maximizing FD within a phylogenetic clade or

region [16]. In reality, conservation decisions reflect the interplay of social, economic, political,

and scientific priorities, and do not necessarily result in the saving of target species (and

therefore of their associated FD or PD). However, our approach allows us to test a long-

standing idea in biodiversity science, and identify two critical ways in which PD-based

conservation can become disconnected from FD. We have not made any assumptions regarding

the macroevolutionary history of these traits (for example, assuming some particular model of

trait evolution) or how phenotypes are distributed across geographical and ecological gradients,

and therefore hope that our results generalize beyond the species we study here.

The spatial scale of our analysis reflects the scale of available data appropriate for

making a general statement about the surrogacy value of PD for FD. The scale of conservation

activities can vary, from the global scale of the hotspots approach to local protected areas

within a single country. Unfortunately, the connection between these scales remains unclear.

For example, if the motivation for protecting FD is to maintain community-driven ecosystem

functions and services [5,6,30], the value of a regional or global scale focus may be

questionable [31]. This worry has motivated studies that focus on local scales [6]. This is an

important area for further research; the patterns linking PD and FD in the regional pool may still

be important, since this is the species pool from which local sites are assembled and

maintained.

The motivator of our test of the surrogacy value of PD for FD is the fact that ecologically-

relevant trait data is in short supply, especially for rare and data-deficient species. Indeed, if it

were not for this relative paucity of data, we could simply prioritize species based on their

unique contribution to FD directly [e.g. 36]. Although there have been massive and well-funded

efforts to collect and curate trait data from across the Tree of Life [33–35], we are still far from

having comprehensive coverage. Furthermore, despite recent progress [e.g. 34,35], it is still not

fully understood which traits are most relevant for responses to environmental change, or that

contribute most to certain ecosystem functions and services, and how these vary among

systems. Our analysis suffers from a similar data limitation. We chose these traits because they

are frequently collected in ecological studies, not because we know they are ecologically

important. Our assumption is that their phylogenetic distribution is typical of those traits that

are most desirable for the purpose of conservation and that our primary results are therefore

widely applicable. We urge others to expand our simple test to other clades and traits in order

to test the generality of our findings.

**Conclusion**

Prioritizing the most phylogenetically diverse set of taxa in a region or clade will result in an

average gain of 18% functional diversity relative to applying the same conservation effort

without considering phylogeny, but this gain will decrease as species richness increases. This

suggests that PD is a reasonable conservation prioritization strategy, especially in species-poor

clades or regions, or in the absence of meaningful data on functional traits. However, we note

two important drawbacks of this strategy. First, in cases of either recent trait divergence or,

alternatively, very strong trait conservatism, a PD prioritization scheme can capture less FD

than a random scheme. Second, we found that while this strategy, on average, captures FD

well, it is also somewhat unreliable, and 36% of the time will not capture more FD than random

choice. Critically, and in opposition to what has previously been implicitly assumed [15,16], we

find weak empirical evidence that the presence of phylogenetic signal in traits predicts whether

PD-based conservation will prioritize FD. Assuming that the traits we have used are

representative of a broader array of ecologically relevant traits, our results provide a baseline

for how well we should expect PD-based prioritization to perform when ecologically important

traits are unknown, unmeasured, or unmeasurable.  By clearly outlining the cases and parts of

the world in which phylogenetically-based conservation prioritization can (and cannot)

effectively act as a surrogate for functional diversity, we hope to inform and improve

conservation interventions globally.

239
240
241

**Figure 1 – A conceptual approach for evaluating if PD a good surrogate for FD.** To evaluate if
PD is a good surrogate of FD, we measure to what extent a species prioritization strategy that
maximize PD captures FD relative to an optimal and a random strategy. To do so, we compare
FD accumulation curves (i.e. FD computed for increasing proportion of the species pool
considered) across these three different sampling strategies: the random sampling (i.e.
rarefaction curve, averaged over 1000 sets), the maxPD (surrogacy, averaged over 1000 sets)
sampling (i.e. the sets that maximize PD) and the maxFD (optimal) sampling (i.e. sets that
maximize FD, see legends).  Then, we measure the surrogacy of PD for FD ($SD_{PD\text{-}FD}$) as the area
between the random and the maxPD curve ('A', see legend) divided by the area between the
random and the maxFD curve ('A+B', see legend). If $SD_{PD\text{-}FD}$ is positive, PD is a good surrogate
for FD (the maximum value being 1 where PD is an optimal surrogate) while when $SD_{PD\text{-}FD}$ is
negative preserving species based on PD is worse than preserving them at random.

**Figure 2 – PD is a good surrogate of FD across space.** The figure presents the distribution and correlates of $S_{PD-FD}$ for mammals (panels A-C), birds (panels D-F) and tropical fishes (G-I) separately across space. For each of the three groups, the $SD_{PD-FD}$ frequency distribution is presented in top panels (B, E and H) along with its mean (vertical line) and the color code that is common to all panels, with blue indicating positive $S_{PD-FD}$ (maximizing PD captures more FD than random). $SD_{PD-FD}$ geographical distribution is presented in middle panels (A, D, G). Relationships between $SD_{PD-FD}$ and species pool richness are presented in panels C, F and I. In each grid cell, $SD_{PD-FD}$ values are based on the mean over 1000 repetitions of random and PDmax set draw (there is only one maxFD set).

**Figure 3 – PD is a good surrogate of FD across clades.** The figure presents the distribution and
correlates of $SD_{PD-FD}$ for mammals (panels A-C) and birds (panels D-F) across families.  For each
of the two groups, the $SD_{PD-FD}$ frequency distribution is presented (B and E) along with its mean
(vertical line). The colour code that is common to all panels. $SD_{PD-FD}$ phylogenetic distribution is
presented in panels A and D. Relationships between $SD_{PD-FD}$ and species pool richness are
presented in panels C, F and I. For each taxonomic family, $SD_{PD-FD}$ values are based on the mean
over 1000 repetitions of random and maxPD set draw (there is only one maxFD set).

**Methods**

We use two classes of data to address the question of whether choosing sets of species according to PD captures the underlying trait diversity (as measured with FD) well. First, we used taxonomic groups (clades) of species as our unit of analysis ('species pool' hereafter) and, second, we investigated broad assemblages found across the globe. The former is more explicitly evolutionary, ensuring that our results are not driven by well-established relationships across large taxonomic groups (*e.g.*, monotremes are distinct from placental mammals) and the latter is likely more relevant to actual conservation practice.

**1. Data**

We use distribution data to delineate geographical assemblage species pool and taxonomy to delineate clade-based species pools (namely families and orders).

**Distribution data –**  For mammals, we used the distribution maps provided by the Mammal Red List Assessment (http://www.iucnredlist.org/) for 4,616 species. For birds, full (breeding and wintering) and breeding ranges distribution maps were extracted from BirdLife (http://www.birdlife.org/) for 9,993 species. The best resolution at which these maps should be used is still under discussion in the literature, so we decided to use the 40 000km$^2$ resolution (200x200km gird cell at the equator) that is commonly used at global scale [38,39]. The total number of grid cells was 3,646. Domestic and aquatic mammals were excluded from the analysis. In order to make sure our results were not driven by the important trait difference between volant and non volant mammals, we repeated our results excluding bats. For birds we repeated our analysis using the full ranges (*i.e.*, summer and winter ranges). Finally, we evaluated the robustness of our result to the spatial resolution considered by repeating our analysis at a resolution of 100x100km (number of grid cells was 13,330) for birds and mammals; we present these results in the supplementary materials, as they are qualitatively identical to those conducted at 200x200km (fig. S1). For fishes, we used a database of 1536 species, for which we had distribution data, phylogenetic and functional data. Distribution data were extracted from a global-scale distribution database

[40]. Species composition was then extracted from grid cells of 5°x5°, corresponding to

approximately 555x555 km at the equator [41]. This grain size of the grid was chosen

because it represents a good compromise between the desired resolution and the

geographical density of information.

**Phylogenies –** In order to prioritize species to maximize PD, phylogenies of each species pool

are needed. We used the first 100 published calibrated ultrametric trees of Jetz et al.[42] for

birds and Faurby and Svenning [43] for mammals. By repeating our analyses across a posterior

distribution of phylogenetic hypotheses, we control and account for phylogenetic uncertainty.

For tropical reef fishes, we built a phylogeny for 18 families (i.e. Labridae, Scaridae,

Pomacentridae, Chaetodontidae, Acanthuridae, Haemulidae, Balistidae, Carangidae,

Serranidae, Lutjanidae, Sparidae, Caesionidae, Holocentridae, Mullidae, Muraenidae,

Tetraodontidae, Lethrinidae and Siganidae) by pruning a dated molecular phylogenetic tree for

7,822 extant fish species [44]. These families were selected as the most representative tropical

reef fish families, that is, they are abundant and speciose on tropical reefs. We grafted missing

species on the pruned phylogenetic tree (circa 50% among the 1536 studied species) based on

published phylogenies for these families, supplemented by taxonomic information from fish

identification guides and FishBase (www.fishbase.org) [44,45]. We recorded, for each of these

trees, a measure of imbalance (as measured by **β** [46]) and 'tipiness' (as measured by Gamma

[47]). For both mammals and birds, we chose to group species in families and orders. We used

these groupings when calculating the purely phylogenetic, clade-based analyses (to address

question 1), but not within the spatial, assemblage-based analyses (question 2). For the

taxonomic analysis of mammal families we removed two families (Dipodidae and Echimyidae)

because of their very poor phylogenetic resolution (i.e. polytomies for an important number of

species).

**Traits –** For birds and mammals, four traits (diet, (log transformed) body mass, activity cycle,

and foraging height) were extracted from Elton Traits1.0 [35]. These traits are generally

assumed to appropriately represent Eltonian niche dimensions within an assemblage or clade

of mammals or birds [35,48,49]. For fishes, we used a previously published database [12]. We used 6 categorical traits: size, mobility, period of activity, schooling, vertical position in the water column, and diet (for a full description of the dataset, see Mouillot et al. [12]). These traits have already been used to investigate community assembly rules [50] and to seek vulnerable fish functions [11]. For each clade and assemblage, we used the raw trait (only body mass was log-transformed and rescaled by the clade/assemblage range of body masses) values to compute distance between species using Gower distance [19] and use PCoA to summarize the trait space in few dimensions. We retained the numbers of PCoA axes necessary to represent 70% of the total initial variability (using a 80% threshold did not quantitatively change our conclusions, see Fig. S9). We also recorded phylogenetic signal for each PCoA axis using Blomberg's K [51].

## 2. Approach

Our aim was to evaluate, across a wide range of clades and regions, the ability of PD-informed prioritization scheme to capture FD in comparison with two other prioritization schemes: selecting species to directly maximize FD ('maxFD' hereafter) and selecting species randomly (Figure 1). Our premise was that we often do not know or have not measured the traits that are most relevant for ecosystem function and services such that maximizing FD is not generally feasible. By focusing on a subset of traits and assuming that they are representative of ecologically relevant traits, we were able to get an estimate of how well PD does compared to the best we could possibly do. We used performance relative to choosing on the basis of FD as an upper-limit to the performance of PD as a surrogate for FD, and used random species selection as a lower benchmark.

**Random prioritization scheme –** For each pool (i.e. each clade and each geographical assemblage) and each number of selected species (10, 20, 30, 40, 50, 60, 70, 80, 90, and 100% of the total pool), 1000 random sets of species were produced, from which the average FD was recorded.

**Prioritization scheme maximizing PD (maxPD) –** While there are many, overlapping metrics for measuring the evolutionary history encompassed by a set of species [52,53], the most common is the sum of all branch lengths (often in units of time) connecting a set of species to a common root[14], called Phylogenetic Diversity (PD). This is the metric whose maximization has most commonly been proposed as a conservation prioritization metric [14,54,55], and as a measure of phylogenetic 'richness' it most naturally maps onto our chosen FD metric [52]. We used the greedy algorithm proposed by Bordewich et al. [56] to find our maxPD set of species $S$. For a given tree there are likely multiple, and possibly very many, sets of species with the same PD as $S$. As a consequence, we produced, for each pool, each number of selected species, and each alternative phylogenetic trees, 10 maxPD sets of species. We then averaged the FD of these sets across our 100 phylogenetic tree, so that each value is an average of 1000 sets (10 sets for each of the 100 trees).

**Prioritization scheme maximizing FD (maxFD) –** Functional diversity was estimated using a functional richness index (FRic; [57–59]). The FRic index relies on a multidimensional Euclidean space, where the axes are traits (or factorial axes from a principal coordinates analysis (PCoA) computed using these traits) along which species are placed according to their trait values. This index measures the volume of trait space occupied by a given species assemblage by calculating the convex hull volume [59], defined by the species at the vertices of the functional space, that encompasses the entire trait space filled by all species in this assemblage. In a single dimension, this simply equals the range of values [59]. This broadly used metric in ecology is set monotonic with species richness, a property generally assumed desirable in conservation whereby the addition of a new species can never decrease the metric's value [60]. FD measures the total amount of variation in trait values, making it conceptually comparable to PD [52]. We used the FRic index instead of the FD index based on a functional dendrogram (Petchey & Gaston, 2006) since recent studies showed that the FD index may lead to biased assessments of functional diversity and inaccurate ecological conclusions [61]. The most straightforward way to obtain the maximal FD for $n$ species is to compute FD for all possible combinations of $n$ species and simply record the greatest value (the brute force approach). However, this is not feasible in

practice as the numbers of combinations of selected species was too high (e.g., $10^{71}$ possible

sets for all mammal assemblages). To rapidly and efficiently find the set of species that aim to

maximize FD, we developed a novel (at least in ecology) greedy algorithm. In brief, our

approach iteratively (starting with two species) select the species that is the furthest from the

centroid of the already selected set. To avoid selecting two species that are far from the

centroid but close to each other, we penalized the distance to the centroid by the distance to

the closest neighbour in the already selected set.  Here we present in details the greedy

algorithm we used to find the set of species that maximize FD:

Step 1. Select the two species with the highest trait distance

Step 2. Compute the centroid of these two selected species

Step 3. Compute distances between species not in the set and this 'set centroid'.

Step 4. Penalize these distance by adding the following factor f (Eq. 4)

400                          $$f = K \times e^{L \times minD}$$                          (eq. 4)

with K and L being penalizing factors and minD the distance between a given candidate

species and the nearest species already in the selected set.

Step 5. Select the species that maximized the penalized distance

Step 6. Go back to step one with this new set of species until the desired number of

species is reached.

To avoid arbitrarily setting the penalizing parameters, we tested 1000 pairs of parameters

drawn from a truncated normal distribution (mean=1, sd=.5) and retained the parameter pairs

that yielded the maximal FD.

In tests of subsets of the data for which finding the true maxFD was feasible, we found our

approach to adequately approximate the true maxFD and to produce a very good

approximation of the true degree of PD's surrogacy for FD (fig. S10).

**Measuring performance and surrogacy of prioritization schemes.**

We use a common approach [26,27] to quantify the extent to which a given surrogate (here,

the maxPD choice) reaches a certain objective (here, maximize FD). Species from a given pool

(i.e., for each dataset (clade and assemblages) independently,) were prioritized and selected

according to (1) the objective, i.e. maximize FD, producing the 'optimal curve' (maxFD curve in

Figure 1), (2) the surrogate i.e. maximize PD, producing the 'surrogate curve' (maxPD curve in

Figure 1) and (3) at random (random curve in Figure 1), i.e. producing the 'random curve'

(Figure 1). To compute a 'surrogacy' estimate of PD ($S_{PD-FD}$), we compare the position of the

surrogate curve (1) to the random curve (2) relative to the optimal curve (2) (Figure 1 and Eq. 1)

across the deciles of species richness of the pool (given as an interval 0-1):

$$S_{PD-FD} = \int_0^1 \frac{FD_{maxPD}-FD_{random}}{FD_{maxPF}-FD_{random}} \qquad\qquad \text{(Equation 1)}$$

This surrogacy metric is at 100% when the surrogate perfectly meets the objective (i.e., the

maxFD and maxPD curves are identical and the max PD set is the maxFD set), 0% when the

surrogate is not better than randomly chosen sets of species (i.e., the random and maxPD

curves are identical) and is negative if the surrogate choice is worse than random (i.e., the

maxPD curve is below the random curve). Correlates of $S_{PD-FD}$ were evaluated using Spearman

correlations.

Apart from focusing on average tendencies, we quantified the variability of the FD yielded by

the PD—maximized selection strategy and the random selection strategy within each species

pools. To do so, we compute, for each species pool and for each % of selected species

independently, the number of cases where $FD_{random}>FD_{maxPD}$ across the 1000 random *1000

maxPD sets combinations (i.e. $10^6$ comparisons). We then averaged theses number across % of

selected species and report statistics across datasets (Supp. Table 1).

1.  Pimm, S.L., Jenkins, C.N., Abell, R., Brooks, T.M., Gittleman, J.L., Joppa, L.N., Raven, P.H., Roberts, C.M., and Sexton, J.O. (2014). The biodiversity of species and their rates of extinction, distribution, and protection. Science (80-. ). *344*.

2.  Hull, P.M., Darroch, S.A.F., and Erwin, D.H. (2015). Rarity in mass extinctions and the future of ecosystems. Nature *528*, 345–51.

3.  Ceballos, G., Ehrlich, P.R., and Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. Proc. Natl. Acad. Sci. U. S. A. *114*, E6089–E6096.

4.  Tilman, D., Wedin, D., and Knops, J. (1996). Productivity and Sustainability Influenced by Biodiversity in Grassland Ecosystems. Nature *379*, 718–720.

5.  Srivastava, D.S., Cadotte, M.W., MacDonald, A.A.M., Marushia, R.G., and Mirotchnick, N. (2012). Phylogenetic diversity and the functioning of ecosystems. Ecol. Lett. *15*, 637–48.

6.  Cadotte, M.W., Carscadden, K., and Mirotchnick, N. (2011). Beyond species: functional diversity and the maintenance of ecological processes and services. J. Appl. Ecol. *48*, 1079–1087.

7.  Isbell, F., Gonzalez, A., Loreau, M., Cowles, J., Díaz, S., Hector, A., Mace, G.M., Wardle, D.A., O'Connor, M.I., Duffy, J.E., *et al.* (2017). Linking the influence and dependence of people on biodiversity across scales. Nature *546*, 65–72.

8.  Pascual, U., Balvanera, P., Diaz, S., Pataki, G., Roth, E., Stenseke, M., Watson, R.T., Wickson, F., and Yagi, N. (2017). Valuing nature's contributions to people: the IPBES approach. Curr. Opin. Environ. Sustain. *26–27*, 7–16.

463 9. Vane-Wright, R.I., Humphries, C.J., and Williams, P.H. (1991). What to protect?—

464   Systematics and the agony of choice. Biol. Conserv. *55*, 235–254.

465 10. Bottrill, M.C., Joseph, L.N., Carwardine, J., Bode, M., Cook, C., Game, E.T., Grantham, H.,

466   Kark, S., Linke, S., McDonald-Madden, E., *et al.* (2008). Is conservation triage just smart

467   decision making? Trends Ecol. Evol. *23*, 649–654.

468 11. Mouillot, D., Bellwood, D.R., Baraloto, C., Chave, J., Galzin, R., Harmelin-vivien, M.,

469   Kulbicki, M., Lavergne, S., Lavorel, S., Mouquet, N., *et al.* (2013). Rare species support

470   vulnerable functions in high-diversity ecosystems. PLOS Biol. *11*.

471 12. Mouillot, D., Villeger, S., Parravicini, V., Kulbicki, M., Arias-Gonzalez, J.E., Bender, M.,

472   Chabanet, P., Floeter, S.R., Friedlander, A., Vigliola, L., *et al.* (2014). Functional over-

473   redundancy and high functional vulnerability in global fish faunas on tropical reefs. Proc.

474   Natl. Acad. Sci. *111*, 13757–13762.

475 13. Cadotte, M.W., Jonathan Davies, T., and Peres-Neto, P.R. (2017). Why phylogenies do not

476   always predict ecological differences. Ecol. Monogr.

477 14. Faith, D.P. (1992). Conservation evaluation and phylogenetic diversity. Biol. Conserv. *61*,

478   1–10.

479 15. Winter, M., Devictor, V., and Schweiger, O. (2013). Phylogenetic diversity and nature

480   conservation: where are we? Trends Ecol. Evol.

481 16. Mazel, F., Mooers, A.O., Riva, G.V.D., and Pennell, M.W. (2017). Conserving Phylogenetic

482   Diversity Can Be a Poor Strategy for Conserving Functional Diversity. Syst. Biol. *In press*.

483 17. Kelly, S., Grenyer, R., and Scotland, R.W. (2014). Phylogenetic trees do not reliably

484   predict feature diversity. Divers. Distrib. *20*, 600–612.

485    18.    Isaac, N.J.B., Turvey, S.T., Collen, B., Waterman, C., and Baillie, J.E.M. (2007). Mammals

486           on the EDGE: conservation priorities based on threat and phylogeny. PLoS One *2*, e296.

487    19.    Barak, R.S., Hipp, A.L., Cavender-Bares, J., Pearse, W.D., Hotchkiss, S.C., Lynch, E.A.,

488           Callaway, J.C., Calcote, R., and Larkin, D.J. (2016). Taking the Long View: Integrating

489           Recorded, Archeological, Paleoecological, and Evolutionary Data into Ecological

490           Restoration. Int. J. Plant Sci. *177*, 90–102.

491    20.    Pollock, L.J., Thuiller, W., and Jetz, W. (2017). Large conservation gains possible for global

492           biodiversity facets. Nature *546*, 141–144.

493    21.    Brum, F.T., Graham, C.H., Costa, G.C., Hedges, S.B., Penone, C., Radeloff, V.C., Rondinini,

494           C., Loyola, R., and Davidson, A.D. (2017). Global priorities for conservation across

495           multiple dimensions of mammalian diversity. Proc. Natl. Acad. Sci. U. S. A., 201706461.

496    22.    Mazel, F., Guilhaumon, F., Mouquet, N., Devictor, V., Gravel, D., Renaud, J., Marcus

497           Vinicius Cianciaruso, R.L., Diniz-Filho, J.A.F., Mouillot, D., and Thuiller, W. (2014).

498           Multifaceted diversity–area relationships reveal global hotspots of mammalian species,

499           trait and lineage diversity. Glob. Ecol. Biogeogr. *23*, 836–847.

500    23.    Thuiller, W., Maiorano, L., Mazel, F., Guilhaumon, F., Ficetola, G.F., Lavergne, S., Renaud,

501           J., Roquet, C., and Mouillot, D. (2015). Conserving the functional and phylogenetic trees

502           of life of European tetrapods. Philos. Trans. R. Soc. Lond. B. Biol. Sci. *370*, 20140005.

503    24.    Devictor, V., Mouillot, D., Meynard, C., Jiguet, F., Thuiller, W., and Mouquet, N. (2010).

504           Spatial mismatch and congruence between taxonomic, phylogenetic and functional

505           diversity: the need for integrative conservation strategies in a changing world. Ecol. Lett.

506           *13*, 1030–1040.

25.   Albouy, C., Delattre, V.L., Mérigot, B., Meynard, C.N., and Leprieur, F. (2017).

Multifaceted biodiversity hotspots of marine mammals for conservation priorities.

Divers. Distrib. *23*, 615–626.

26.   Ferrier, S., and Watson, G. (1997). An evaluation of the effectiveness of environmental

surrogates and modelling techniques in predicting the distribution of biological diversity.

Armidale NSW Environ. Aust.

27.   Rodrigues, A.S.L., and Brooks, T.M. (2007). Shortcuts for Biodiversity Conservation

Planning: The Effectiveness of Surrogates. Annu. Rev. Ecol. Evol. Syst. *38*, 713–737.

28.   Rodrigues, A.S.L., Brooks, T.M., and Gaston, K.J. (2005). Integrating phylogenetic diversity

in the selection of priority areas for conservation: does it make a difference? In

Phylogeny and Conservation, A. Purvis, J. L. Gittleman, and T. M. Brooks, eds.

(Cambridge, UK: Cambridge Univ. Press), pp. 101–119.

29.   Mazel, F., Wüest, R.O., Gueguen, M., Renaud, J., Ficetola, G.F., Lavergne, S., and Thuiller,

W. (2017). The Geography of Ecological Niche Evolution in Mammals. Curr. Biol. *27*,

1369–1374.

30.   Díaz, S., Lavorel, S., de Bello, F., Quétier, F., Grigulis, K., and Robson, T.M. (2007).

Incorporating plant functional diversity effects in ecosystem service assessments. Proc.

Natl. Acad. Sci. U. S. A. *104*, 20684–9.

31.   Srivastava, D.S., and Vellend, M. (2005). Biodiversity-Ecosytem Function Research: Is It

Relevant to Conservation? Annu. Rev. Ecol. Evol. Syst. *36*, 267–294.

32.   Cornwell, W.K., Westoby, M., Falster, D.S., FitzJohn, R.G., O'Meara, B.C., Pennell, M.W.,

McGlinn, D.J., Eastman, J.M., Moles, A.T., Reich, P.B., *et al.* (2014). Functional

distinctiveness of major plant lineages. J. Ecol. *102*, 345–356.

33.    Kattge, J., Diaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Bönisch, G., Garnier, E., Westoby,

M., Reich, P.B., Wright, I.J., *et al.* (2011). TRY--a global database of plant traits. Glob.

Chang. Biol. *17*, 2905–2935.

34.    Jones, K.E., Bielby, J., Cardillo, M., Fritz, S.A., O'Dell, J., Orme, C.D.L., Safi, K., Sechrest, W.,

Boakes, E.H., Carbone, C., *et al.* (2009). PanTHERIA: a species-level database of life

history, ecology, and geography of extant and recently extinct mammals. Ecology *90*,

2648–2648.

35.    Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M.M., and Jetz, W.

(2014). EltonTraits 1.0: Species-level foraging attributes of the world's birds and

mammals. Ecology *95*, 2027–2027.

36.    Lavorel, S., and Garnier, E. (2002). Predicting changes in community composition and

ecosystem functioning from plant traits: revisiting the Holy Grail. Funct. Ecol. *16*, 545–

556.

37.    Díaz, S., Purvis, A., Cornelissen, J.H.C., Mace, G.M., Donoghue, M.J., Ewers, R.M.,

Jordano, P., and Pearse, W.D. (2013). Functional traits, the phylogeny of function, and

ecosystem service vulnerability. Ecol. Evol. *3*, 2958–75.

38.    Hurlbert, A.H., and Jetz, W. (2007). Species richness, hotspots, and the scale dependence

of range maps in ecology and conservation. Proc. Natl. Acad. Sci. U. S. A. *104*, 13384–9.

39.    Holt, B.G., Lessard, J.P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., Fabre,

P.H., Graham, C.H., Graves, G.R., Jønsson, K.A., *et al.* (2013). An Update of Wallace's

Zoogeographic Regions of the World. Science (80-. ). *339*, 74–78.

40.  Parravicini, V., Kulbicki, M., Bellwood, D.R., Friedlander, A.M., Arias-Gonzalez, J.E.,

Chabanet, P., Floeter, S.R., Myers, R., Vigliola, L., D?Agata, S., *et al.* (2013). Global

patterns and predictors of tropical reef fish species richness. Ecography (Cop.). *36*, 1254–

1262.

41.  Leprieur, F., Descombes, P., Gaboriau, T., Cowman, P.F., Parravicini, V., Kulbicki, M.,

Melián, C.J., de Santana, C.N., Heine, C., Mouillot, D., *et al.* (2016). Plate tectonics drive

tropical reef biodiversity dynamics. Nat. Commun. *7*, 11461.

42.  Jetz, W., Thomas, G., Joy, J., and Mooers, A.Ø. (2012). The global diversity of birds in

space and time. Nature *491*, 444–448.

43.  Faurby, S., and Svenning, J.-C. (2015). A species-level phylogeny of all extant and late

Quaternary extinct mammals using a novel heuristic-hierarchical Bayesian approach.

Mol. Phylogenet. Evol. *84*, 14–26.

44.  Leprieur, F., Descombes, P., Gaboriau, T., Cowman, P.F., Parravicini, V., Kulbicki, M.,

Melián, C.J., de Santana, C.N., Heine, C., Mouillot, D., *et al.* (2016). Plate tectonics drive

tropical reef biodiversity dynamics. Nat. Commun. *7*, 11461.

45.  Ottimofiore, E., Albouy, C., Leprieur, F., Descombes, P., Kulbicki, M., Mouillot, D.,

Parravicini, V., and Pellissier, L. (2017). Responses of coral reef fishes to past climate

changes are related to life-history traits. Ecol. Evol. *7*, 1996–2005.

46.  Blum, M., François, O., and Steel, M. (2006). Which Random Processes Describe the Tree

of Life? A Large-Scale Study of Phylogenetic Tree Imbalance. Syst. Biol. *55*, 685–691.

47.  Pybus, O., and Harvey, P. (2000). Testing macro–evolutionary models using incomplete

molecular phylogenies. Proc. R. Soc. London, Ser. B *267*, 2267–2272.

48.   Safi, K., Cianciaruso, M. V, Loyola, R.D., Brito, D., Armour-Marshall, K., and Diniz-Filho,

J.A.F. (2011). Understanding global patterns of mammalian functional and phylogenetic

diversity. Philos. Trans. R. Soc. London, Ser. B *366*, 2536–2544.

49.   Belmaker, J., and Jetz, W. (2015). Relative roles of ecological and energetic constraints,

diversification rates and region history on global species richness gradients. Ecol. Lett. *18*,

563–571.

50.   Guillemot, N., Kulbicki, M., Chabanet, P., Vigliola, L., Possingham, H., Garnier, E.,

Goldberg, D., Hooper, D., Jackson, S., and Navas, M. (2011). Functional Redundancy

Patterns Reveal Non-Random Assembly Rules in a Species-Rich Marine Assemblage. PLoS

One *6*, e26735.

51.   Blomberg, S.P., Garland, T.J.R., and Ives, A. (2003). Testing for phylogenetic signal in

compartive data : behavioral traits are more labile. Evolution (N. Y). *57*, 717–745.

52.   Tucker, C.M., Cadotte, M.W., Carvalho, S.B., Davies, T.J., Ferrier, S., Fritz, S.A., Grenyer,

R., Helmus, M.R., Jin, L.S., Mooers, A.Ø., *et al.* (2017). A guide to phylogenetic metrics for

conservation, community ecology and macroecology. Biol. Rev. Camb. Philos. Soc.

53.   Winter, M., Devictor, V., and Schweiger, O. (2013). Phylogenetic diversity and nature

conservation: where are we? Trends Ecol. Evol. *28*, 199–204.

54.   Forest, F., Grenyer, R., Rouget, M., Davies, T.J., Cowling, R.M., Faith, D.P., Balmford, A.,

Manning, J.C., Procheş, S., van der Bank, M., *et al.* (2007). Preserving the evolutionary

potential of floras in biodiversity hotspots. Nature *445*, 757–760.

55.   Rodrigues, A.S.L., and Gaston, K.J. (2002). Maximising phylogenetic diversity in the

selection of networks of conservation areas. Biol. Conserv. *105*, 103–111.

56.  Bordewich, M., Rodrigo, A., Semple, C., and Collins, T. (2008). Selecting Taxa to Save or

Sequence: Desirable Criteria and a Greedy Solution. Syst. Biol. *57*, 825–834.

57.  Villéger, S., Mason, N., and Mouillot, D. (2008). New multidimensional functional

diversity indices for a multifaceted framework in functional ecology. Ecology *89*, 2290–

2301.

58.  Pavoine, S., and Bonsall, M.B. (2011). Measuring biodiversity to explain community

assembly: a unified approach. Biol. Rev. *86*, 792–812.

59.  Cornwell, W.K., Schwilk, D.W., Ackerly, D.D., and Schwilk, L.D.W. (2006). A trait-based

test for habitat filtering: convex hull volume. Ecology *87*, 1465–1471.

60.  Ricotta, C. (2005). A note on functional diversity measures. Basic Appl. Ecol. *6*, 479–486.

61.  Villéger, S., Maire, E., and Leprieur, F. (2017). On the risks of using dendrograms to

measure functional diversity and multidimensional spaces to measure phylogenetic

diversity: a comment on Sobral *et al* . (2016). Ecol. Lett. *20*, 554–557.