High accuracy haplotype-derived allele frequencies from ultra-low coverage pool-seq samples

Susanne Tilk[1], Alan Bergland[1,2], Aaron Goodman[1], Paul Schmidt[3], Dmitri Petrov[1], Sharon Greenblum[1]

[1]Department of Biology, Stanford University, Stanford CA 94305
[2]Department of Biology, University of Virginia, Charlottesville VA 22904
[3]Department of Biology, University of Pennsylvania, Philadelphia PA 19104

Abstract

Evolve-and-resequence experiments leverage next-generation sequencing technology to track allele frequency dynamics of populations as they evolve. While previous work has shown that adaptive alleles can be detected by comparing frequency trajectories from many replicate populations, this power comes at the expense of high-coverage (>100x) sequencing of many pooled samples, which can be cost-prohibitive. Here we show that accurate estimates of allele frequencies can be achieved with very shallow sequencing depths (<5x) via inference of known founder haplotypes in small genomic windows. This technique can be used to efficiently estimate frequencies for any number of alleles in any model system. Using both experimentally-pooled and simulated samples of *Drosophila melanogaster,* we show that haplotype inference can improve allele frequency accuracy by orders of magnitude, and that high accuracy is maintained after up to 200 generations of recombination, even in the presence of missing data or incomplete founder knowledge. By reducing sequencing costs without sacrificing accuracy, our method enables analysis of samples from more time-points and replicates, increasing the statistical power to detect adaptive alleles.

Introduction

A major barrier to understanding the genetic basis of rapid adaptation has been the lack of robust experimental frameworks for assaying allele frequency dynamics. Recently, evolve and resequence (E+R) experiments [1], which leverage next-generation sequencing technology to track real-time genome-wide allele frequency changes during evolution, have become a powerful step forward in studying adaptation [2]. In most E+R studies, replicate populations are evolved over tens to hundreds of generations in an artificial or natural selection regime and allele frequency measurements from multiple time-points are used to identify genomic targets of selection. To date, E+R approaches have already been successfully applied in a variety of model systems, including RNA molecules, viruses, *Escherichia coli*, *Saccharomyces cerevisiae*, and *Drosophila melanogaster* [3–7]. The ability to concurrently observe both trait and genome-wide allele frequency changes across multiple systems offers the potential to answer long-standing

questions in molecular evolution. Careful analysis of the patterns and magnitude of allele frequency change may reveal the extent of the genome that is under selection, how interacting alleles contribute to adaptive traits, and the speed of adaptation in different evolutionary regimes.

Crucially, however, the power to address such questions depends on the replication, time-resolution, and accuracy of allele frequency trajectories, with tradeoffs between these often incurred due to high sequencing costs. Recommended E+R schemes with even minimal power to detect selection involve sampling tens to hundreds of individuals from at least three replicate populations over a minimum of ten generations [8,9]. Since individual-based, genome-wide DNA sequencing at sufficient coverages is enormously cost-prohibitive, most E+R studies rely instead on pooled sequencing [10–13] of all individuals sampled from a given time-point and replicate. While this approach sacrifices information about individual genotypes and linkage, pooled sequencing has been shown to provide a reliable measure of population-level allele frequencies [14,15]. Still, forward-in-time simulations suggest that each pooled sample must be sequenced at a minimum of 50x coverage to detect strong selection and even higher coverage to detect weak selection [9]. Given that optimized experimental designs often involve >100 samples, total costs for *Drosophila melanogaster* E+R experiments that achieve reasonable detection power can reach well above $25,000 at current sequencing costs. Thus achieving sufficient accuracy remains a major limiting factor in capitalizing on the promise of E+R.

The short timescales for which E+R is most appropriate may, however, facilitate ways to reduce sequencing costs without sacrificing experimental power. First, there is a growing body of evidence that in sexual populations, the bulk of short-term adaptation, especially in fairly small populations, relies on standing genetic variation rather than new mutations [5,12]. Many E+R schemes involve experimental populations derived from a fixed number of inbred founder lines [6,16,17], so the identity, starting frequency, and haplotype structure of all segregating variants are often either already well-known or can easily be obtained by sequencing each founder line. Tracking only the frequencies of these validated variants can still provide enough power to detect selection, while reducing the depth of sequencing usually required to differentiate new mutations from sequencing error.

Second, at short timescales haplotype structure can be leveraged to provide more accurate allele frequency estimates. In the time frame of most E+R experiments, recombination does not fully break apart haplotype blocks and disrupt linkage, and thus genomes in an evolving population are each expected to be a mosaic of founder haplotypes. In this scenario, recently developed haplotype inference tools [18–23] can integrate information from sequencing reads across multiple nearby sites to efficiently infer the relative frequency of each founder haplotype within small genomic windows. These haplotype frequencies can then be used as weights to calculate pooled allele frequencies for local segregating variants. With this approach, the accuracy of an allele frequency estimate depends less on the number of mapped reads at the individual site, and instead relies on the discriminatory power of all mapped reads in the surrounding genomic window when inferring haplotype frequencies. Haplotype inference methods

such as HARP [22] have been shown to accurately predict haplotype frequencies at coverages as low as 25x, and simulations of pool-seq data from a small genomic region at fixed read depth indicate that the use of haplotype frequency information increases the power to detect selection [24]. However, these tools have not yet been used to infer allele frequencies from real pooled samples in an E+R framework, nor has a thorough analysis been performed to fully examine how empirical depth of pooled coverage scales with the accuracy of haplotype-derived allele frequency estimates at individual SNPs genome-wide, across many parameters relevant for E+R.

Here, we focus on defining the accuracy of haplotype-derived allele frequencies (HAFs) in order to provide new optimal design recommendations for haplotype-informed E+R given realistic financial constraints, real-life experimental noise, and real-life levels of missing data. Since haplotype inference will be affected by 1) read depths throughout genomic windows, 2) recombination events, and 3) incomplete founder information, we begin by leveraging both simulated and experimental data to assess how the accuracy of HAFs scales with each of these parameters. To do so, we introduce a new metric, 'effective coverage', that associates the error from HAF estimates to the expected sampling error of pooled sequences at various read depths. We find that haplotype inference can significantly increase the accuracy of allele frequency estimations across a range of genomic window sizes, multiple generations of recombination, and with incomplete information about founders. Although we primarily focus on simulated and experimental data from *Drosophila melanogaster*, we later describe how our results can be extended to other model organisms as well. We conclude our findings by offering recommendations about the most powerful way to integrate haplotype inference into E+R experimental schemes, paving the way for deeper insight into the genomic underpinnings of adaptation.

Results

Similar to many E+R studies that approximate population-wide allele frequencies at a given time-point by randomly sampling and pooling ~100 individuals [6,17,25], we pooled two biological replicates of 99 *Drosophila melanogaster* individuals, and performed high-coverage sequencing of each replicate. In our pooled samples however, each individual was derived from a different previously sequenced isogenic founder line.

All reads were mapped to the *D. melanogaster* reference genome, and HAFs along with raw AFs, were calculated at each of the 283,437 known segregating bi-allelic sites on chromosome 2L (for simplicity, the remainder of the analysis focuses just on this chromosome). Raw AFs were calculated by evaluating the fraction of mapped reads containing the alternate allele. To calculate HAFs, founder haplotypes were first constructed from founder genotype calls at the same 283k sites. Haplotype frequency estimation was performed with HARP, a haplotype inference tool that uses both sequence identity and base quality scores to probabilistically assign pool-seq reads to founder haplotypes, and then obtain maximum likelihood estimates of haplotype frequency in discrete chromosomal windows. After inferring the frequency of each founder haplotype in sliding windows across the chromosome, we calculated HAFs at

each SNP site by evaluating the average weighted sum of local founder haplotypes containing the alternate allele.

To determine the accuracy of HAFs and raw AFs, estimated allele frequencies were compared to 'true' allele frequencies derived from the known composition of founder haplotypes that incorporated estimates of uneven pooling (see Supp. Fig. 1 and Supp. Text). Chromosome-wide accuracy of HAFs and raw AFs was quantified using a new metric, *effective coverage,* which represents the theoretical coverage at which the expected binomial variance from read sampling equals the average error from observed allele frequency estimates. While expected error from extreme allele frequencies is lower than that of intermediate allele frequencies under a binomial model, by taking the ratio of the expected error and the observed error for the same set of true allele frequencies, effective coverage values do not depend on the underlying allele frequency spectrum (see *Methods* for full description). Note that while this metric specifically focuses on the variance from sampling of pooled sequences, in practice, the ability of both HAFs and raw AFs to accurately reflect true population-level allele frequencies will also depend on variance from sampling of individuals from the population for pooling. This independent source of error has however been well-treated elsewhere [26,27] and will not be impacted by haplotype inference.

In the following sections, we explore how the accuracy of HAFs differs from raw AFs, and how it scales with empirical coverage, inference window size, different founder sets, incomplete founder information, and the number of generations of recombination.

Imputation of missing genotype calls reduces bias in haplotype frequency assignment

Ambiguous or missing founder genotype calls are common due to residual heterozygosity in inbred lines and uneven sampling during initial founder sequencing. On average, at each segregating site in our set of 99 founders, genotype calls for 4 founders were unresolved. Initial rounds of haplotype inference produced a clear negative correlation between the number of missing calls per founder and the haplotype frequencies estimated for that founder (Supp. Fig. 2). However, we found that imputation of missing founder genotypes prior to haplotype inference both significantly reduced the skewed haplotype frequency assignment and produced more accurate HAFs overall. Thus, we include imputation as a key step in our HAF calculation pipeline for the rest of the analysis (see Supp. Text).

Haplotype inference significantly increases the accuracy of allele frequency estimations

The accuracy of HAFs depends on the power to estimate haplotype frequencies, which in turn is affected by the coverage of mapped reads throughout the genomic window used for haplotype inference. In order to compare the accuracy of HAFs to raw AFs and test how each scales with empirical coverage, reads from the two biological replicates originally sequenced at ~140x were down-sampled to chromosome-wide empirical

coverages of 1x to 100x, and then used to calculate the effective coverage of allele
frequencies for each replicate (Fig. 1a). Haplotype inference was initially performed
using 100kb sliding windows, and accuracy here is assessed at the 27k sites with
known genotype information for every founder. As expected, effective coverage of both
HAFs and raw AFs is similar between the two biological replicates, and increases with
greater chromosome-wide empirical coverage. Yet for all empirical coverages tested,
HAFs have strikingly higher effective coverages than raw AFs. This substantial gain in
accuracy from haplotype inference was most prominent at lower empirical coverages,
with a 40-fold increase in accuracy at 10x empirical coverage, from 10x to effectively
400x. At higher empirical coverages, haplotype inference appears to produce
diminishing returns and effective coverage begins to plateau.
12
We next tested the effect of using smaller (10kb; Fig. 1b) or larger (1000kb; Fig. 1d)
windows for haplotype inference at empirical coverages up to 10x. We find that for both
pooled replicates, larger window sizes provide the most accurate HAFs, since more
reads are available to infer haplotype frequencies in each window. Specifically, 1000kb
HAFs derived from empirical coverages of 1x and 5x reached effective coverages of
>400x and >900x, respectively.
19
We also confirmed that the same results would be achieved by simulated samples with
known sources of error. To do so, we simulated pooled synthetic reads with a standard
Illumina sequencing error rate of 0.002 [28] and corresponding base quality scores [22] from
the same proportions of the 99 founder lines included in the first biological replicate, and
calculated effective coverage with the same empirical coverages and window sizes as
above. Effective coverages for these simulated samples closely mirror effective
coverages obtained from matched experimental samples (Fig. 1b-d). Slight differences
at higher empirical coverages and larger window sizes are most likely caused by
compounded experimental error from DNA extractions, PCR reactions and sequencing,
as well as ambiguity in the 'true' genotypes estimated for individually sequenced lines.
30
Finally, we confirmed that the increased accuracy due to haplotype inference applies
equally to pooled samples derived from other founder sets. We simulated pooled
samples from an entirely different founder set composed of lines from the DGRP [29] and
found that the relationship between HAFs and raw AFs in all ranges of empirical
coverages and window sizes tested is qualitatively the same between the two different
founder sets (Supp. Fig. 3).
37
38 Together, these results suggest that HAFs derived from both simulated samples and
39 multiple biologically distinct samples sequenced at low empirical coverages can be
40 orders of magnitude more accurate than raw AFs. In the following analyses we focus
41 on simulated data from the 99 original founder lines in order to precisely and reliably
42 test how recombination and founder ambiguity affect HAFs in realistic E+R scenarios.
43
44
45
46

Recombination and selection minimally affect accuracy over short time scales

Since recombination will unlink neighboring alleles resulting in smaller haplotype blocks,
we hypothesized that as an experiment proceeds, the accuracy of HAFs calculated with
a given inference window size will decrease. To test this prediction, we used forqs to
perform forward-in-time simulation of neutral recombination for 200 generations in a
population of 1000 individuals using a *D. melanogaster* recombination map [30] , and
tracked the breakpoints and haplotype of origin for all recombined segments at every
generation. At various timepoints, we randomly selected 198 recombined chromosomes
(ie. 99 diploid individuals), and constructed the full sequences of these 'sampled'
chromosomes from corresponding segments of the 99 individually sequenced founder
haplotypes. Reads were simulated from the pooled set of 198 chromosomes after
assigning 'true' genotypes to all missing calls that were then hidden during downstream
haplotype inference calculations (see methods for full description). The accuracy of
HAFs calculated from these simulated reads was assessed using 1000kb, 100kb, and
10kb inference windows at all 283k segregating sites. This is contrast to the analysis
above that only computed HAF accuracy at sites with complete founder information, and
allowed us to realistically incorporate missing information found in experimental
samples.

We first confirmed that, with all window sizes tested, the accuracy of HAFs decreases after more generations of recombination (Fig. 2). Additionally, there is a tradeoff between window sizes (1000kb vs 100kb) after 50 generations, as the accuracy gained from extra information in larger windows is outweighed by the incorrect assumption of complete haplotype blocks. However, even after 50 generations of recombination, HAFs calculated with 100kb windows and an empirical coverage of 5x achieve an effective coverage of >100x, and the same empirical coverage can achieve effective coverages >50x at any timepoint tested. Conversely, 10kb windows may only be useful after >200 generations of recombination, longer than most E+R experiments to date, and therefore the rest of our analysis focuses on using 100kb and 1000kb windows.

In order to ensure that these results also extend to different evolutionary dynamics and are not specific any particular set of chromosomal breakpoints, we repeated this simulation 10 additional times, starting with the same founder population (Supp. Fig. 4). During the first few generations of recombination, evolutionary dynamics and sampling do result in notable variance in the accuracy of HAFs, although this variance rapidly diminishes as recombination proceeds. We also confirmed that our estimates of effective coverage are robust to non-neutral dynamics in the presence of selected sites with varying selective strength (Supp. Fig. 5).

Figure 2. Effect of recombination on the accuracy of HAFs. Effective coverage values of HAFs on chromosome 2L after 0 to 200 generations of recombination in simulated pooled samples of founders (n=99) using 1000kb (dark red), 100kb (red) and 10kb (pink) windows.

HAF accuracy is impacted by haplotype inference with incomplete founder sets

Although recombination is a major factor affecting the accuracy of allele frequency estimates, ambiguity in the set of founders represented in the pool can also influence accuracy. As evolution proceeds during E+R experiments, whole founder haplotypes can be lost from the population. Similarly, contamination may create scenarios in which unintended haplotypes become part of the pool. Loss or gain of founder haplotypes over time has the potential to affect the accuracy of HAFs. We tested the accuracy of HAFs in the first scenario of haplotype loss by allowing haplotype frequency assignment to an extended set of founders, in addition to those included in the original population. Conversely, we tested the accuracy of HAFs in the second scenario by constraining haplotype frequency assignment to only 50% of the original founders. In both cases, HAF accuracy was evaluated in multiple window sizes after 5-20 generations of recombination.

We find that the first scenario (founder haplotype loss) has a notable effect on effective coverage (Fig. 3), reducing accuracy by ~40% and ~50% when 25% and 50% of the

founders were lost from the population, respectively, after 15 generations of
recombination at 10x coverage. The second scenario (haplotype contamination)
resulted in a 50% decrease of effective coverage after 15 generations of recombination
when the population was contaminated by additional haplotypes, increasing the number
of haplotypes in the population by 25% compared to the original founder set of 99
haplotypes. Effective coverage was further decreased by 63% when the number of
haplotypes in the population was increased by 50%, and by 77% when the number of
haplotypes was doubled. However, even after founder losses and gains of 50%,
effective coverage remained >100x and >50x, respectively, in all E+R scenarios tested
here.
11

12
13

Figure 3. Measuring HAF accuracy after loss or gain of founder haplotypes.
Effective coverage values of HAFs after intermediate amounts of recombination (5 gens, left; 15 gens, middle; 20 gens, right) using 1000kb (top panel), or 100kb (bottom panel) windows. Effective coverage values were calculated after founder haplotype loss, where fewer founders were represented in the pool than the number of potential haplotype frequency assignments (red lines), and founder haplotype gain, where more founders were represented in the pool than the number of potential haplotype frequency assignments (blue lines).

Estimating effective coverage with other model organisms

Finally, we explored how the utility of HAFs may extend to any genomic region, for any founder set with known SNPs, and in any model organism with a known recombination rate. To do so, we evaluated effective coverage separately within each 1000kb inference window of chromosome 2L, for all generations and empirical coverages tested in the analyses above. As recombination rate and SNP density varies across the chromosome, we developed a more fine-scale map of effective coverage, parameterized by the expected number of recombination events per sampled chromosome within each window (a function of the number of generations and the local recombination rate), and the expected number of reads mapped to SNP sites within each window (a function of empirical coverage and the local SNP density) (Supp. Fig. 6). With this map, chromosome-wide effective coverage values reported here can then be translated to any founder set or model organism by comparing SNP densities and recombination rates. For example, *Drosophila melanogaster* chromosome 2L has an average recombination rate of 2.39 cM/Mb [30] and our founder set contains an average of 124 SNPs per Mb per founder line. Chromosome I in *Caenorhabditis elegans*, however, has a recombination rate that is 1.4x that of Drosophila (3.33 cM/Mb [31]) and a commonly used reference panel of 249 *C. elegans* strains [32] contains 0.6x as many SNPs per line (74 SNPs/Mb). Thus, to achieve an effective coverage of 150x after 45 generations of *C. elegans* recombination (with the expected number of breakpoints equivalent to ~60 D. melanogaster generations), one should aim to sequence pooled samples at an empirical coverage of >10x (with the expected number of informative reads per line equivalent to *D. melanogaster* coverage of ~6x) .

Discussion

E+R experiments have become a powerful tool to assay the underpinnings of rapid adaptation by tracking allele frequency trajectories within populations over time. Previous studies have shown that the greatest power to detect adaptive variants comes from an optimized experimental design that tracks allele frequencies in multiple replicate populations, samples each replicate population at multiple timepoints, and maximizes the coverage of each pooled sample. Incorporating all of these factors into an E+R framework, however, can present significant financial challenges. Here, we offer a way to mitigate these high sequencing costs without sacrificing statistical power.

Our framework uses haplotype inference to increase the accuracy of pooled allele frequency estimates at low coverages. Since the accuracy of haplotype-derived allele

frequencies relies on the total discriminatory power of reads across a genomic window, rather than coverage at a single site, this approach allows us to sequence less but still maintain high accuracy in allele frequency estimations. Namely, our method achieves the same accuracy expected from sequencing each sample at 100x (as recommended in order to reliably detect strong selection), while only requiring empirical coverage of 1x or less, bringing total sequencing costs from >$25,000 down to less than $200.

There are, however, limitations to this approach. First, as presented, this framework requires the founder population to be derived entirely from fully-inbred lines. As a result, the population dynamics of loci under selection may differ slightly from trajectories in natural populations due to the genetic diversity lost in the inbreeding process (i.e. natural haplotypes, homozygous lethal mutations, and rare variants), as well as higher levels of linkage disequilibrium. Reconstituting an outbred population using inbred lines, however, can be an effective way to mitigate effects of the inbreeding process, and has been experimentally shown to have negligible bias and effect on adaptive dynamics [33]. Alternatively, the use of inbred lines may not be necessary with more sophisticated founder sequencing approaches that incorporate phasing. Newer long-read technologies may make this an achievable reality for a number of systems in the near future.

Second, this approach requires a reliable and comprehensive account of the variants present in each founder line. Since previous studies recommend upwards of 100 founders, sequencing each individual founder line to a sufficiently high depth (in our work, we have found sequencing coverages >10x to be sufficient) may present a high upfront cost. However, this cost represents a one-time investment, which can be applied toward all future experiments using the same set of founders. Furthermore, a number of consortiums already maintain publically available stocks of large numbers of *Drosophila* lines with full, high-quality genome sequences [29,34]. We anticipate that these resources will continue to rapidly expand, facilitating experiments with even greater haplotype diversity at minimal costs.

In addition, this approach is limited to studying short-term adaptation on the scale of tens of generations. In fact, an assumption of our method is that within an inference window, recombination breakpoints minimally affect the ability to accurately call haplotype frequencies. For a given window size however, this assumption becomes less valid as recombination proceeds, and haplotypes blocks decay. Conversely, using increasingly smaller windows reduces the information used for haplotype inference, to the point at which HAFs are no longer more accurate than raw AFs. Though our results here demonstrate that recombination will limit the ability to detect adaptation on timescales of more than 200 generations, the short-term adaptive dynamics which E+R is best suited for fall well within this range. Furthermore, it is at these short timescales, when large numbers of replicate populations are critical for reliably detecting selection, that the cost savings associated with haplotype inference methods will be most beneficial.

Finally, this approach relies on tracking the trajectories of known bi-allelic polymorphisms derived from the founder population, and thus, de novo mutations will not be assayed in this framework. Nonetheless our approach should sufficiently capture the salient features of short-term adaptive dynamics, as there is a growing body of experimental evidence suggesting that selection acts primarily on standing genetic variation in sexual organisms, and that de novo beneficial mutations do not play a large role in rapid adaptation [5,35–37]. Additionally, by tracking only known well-validated polymorphisms, the approach is largely robust to error from small non-SNP chromosomal variants such as indels.

Despite the above limitations, collectively our results show that integrating haplotype inference into future E+R experiments is the most cost-effective way to achieve accuracy in allele frequency estimates, which will directly improve the ability to detect genome-wide signatures of adaptation. Consequently, we offer specific recommendations for future E+R experimental schemes that take advantage of this approach. First, each founder line should be initially sequenced to a sufficient depth that minimizes any missing genotypes. Preliminary analysis reveals that the loss of even 1 called founder genotype per site results in notable drops in accuracy. If missing genotype calls do exist in founder lines, imputing sites prior to haplotype inference can mitigate some of this error. When calculating haplotype frequencies, we find that using large inference window sizes (1000kb) and providing information for the most comprehensive set of founders maximizes the accuracy of allele frequencies and effective coverages attained for each relevant E+R scenario tested here. Together, these guidelines and the analysis above form a framework for achieving effective coverages of close to 100x with empirical coverages as low as 1x even after 50 generations of recombination, reducing sequencing costs by 100-fold. Ultimately, these cost savings, which can be extended to experiments with a variety of model organisms, will facilitate E+R frameworks that can incorporate large numbers of replicate populations. These improvements will be crucial to the statistical power to distinguish between beneficial and neutral alleles [5], and ultimately the future of E+R as a sustainable and feasible experimental tool.

References

1.   Long, A., Liti, G., Luptak, A. & Tenaillon, O. Elucidating the molecular architecture of adaptation via evolve and resequence experiments. *Nat Rev Genetics* **16,** 567–82 (2015).

2.   Burke, MK. How does adaptation sweep through the genome? Insights from long-term selection experiments. *Proceedings of the Royal Society of …* (2012). at <http://rspb.royalsocietypublishing.org/content/early/2012/07/17/rspb.2012.0799.short>

3.   Pitt, JN & Ferré-D'Amaré, AR. Rapid construction of empirical RNA fitness landscapes. *Science* (2010). at <http://science.sciencemag.org/content/330/6002/376.short>

4.   Barrick, JE, Yu, DS, Yoon, SH, Jeong, H & Oh, TK. Genome evolution and adaptation in a long-term experiment with Escherichia coli. *Nature* (2009). at

<http://search.proquest.com/openview/6675f97b6f9c1df27431787caadadc8e/1?pq-origsite=gscholar&cbl=40569>

5.   Burke, MK, Liti, G & Long, AD. Standing genetic variation drives repeatable experimental evolution in outcrossing populations of Saccharomyces cerevisiae. *Molecular biology and evolution* (2014). doi:10.1093/molbev/msu256

6.   OROZCO-terWENGEL, P, Kapun, M & Molecular …, N.-V. Adaptation of Drosophila to a novel laboratory environment reveals temporally heterogeneous trajectories of selected alleles. *Molecular …* (2012). doi:10.1111/j.1365-294X.2012.05673.x

7.   Wichman, HA, Badgett, MR & Scott, LA. Different trajectories of parallel evolution during viral adaptation. … (1999). at <http://science.sciencemag.org/content/285/5426/422.short>

8.   Kofler, R & Schlötterer, C. A guide for the design of evolve and resequencing studies. *Molecular biology and evolution* (2013). at <https://academic.oup.com/mbe/article-abstract/31/2/474/1000475>

9.   Schlötterer, C, Kofler, R, Versace, E & Tobler, R. Combining experimental evolution with next-generation sequencing: a powerful tool to study adaptation from standing genetic variation. *Heredity* (2015). at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4815507/>

10.   Burke, MK, Dunham, JP, Shahrestani, P & Thornton, KR. Genome-wide analysis of a long-term evolution experiment with Drosophila. *Nature* (2010). at <http://search.proquest.com/openview/33de3b0d4f540e9392f9b5faff2a3368/1?pq-origsite=gscholar&cbl=40569>

11.   Illingworth, C. J., Parts, L., Schiffels, S., Liti, G. & Mustonen, V. Quantifying selection acting on a complex trait using allele frequency time series data. *Molecular biology and evolution* **29,** 1187–1197 (2011).

12.   Graves, J. L. *et al.* Genomics of Parallel Experimental Evolution in Drosophila. *Mol. Biol. Evol.* **34,** 831–842 (2017).

13.   Barghi, N., Tobler, R., Nolte, V. & Schlötterer, C. Drosophila simulans: A Species with Improved Resolution in Evolve and Resequence Studies. *G3 (Bethesda)* **7,** 2337–2343 (2017).

14.   Zhu, Y, Bergland, AO, González, J & one, P.-D. Empirical validation of pooled whole genome population re-sequencing in Drosophila melanogaster. *PloS one* (2012). at <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0041901>

15.   Fracassetti, M., Griffin, P. & Willi, Y. Validation of Pooled Whole-Genome Re-Sequencing in Arabidopsis lyrata. *Plos One* **10,** e0140462 (2015).

16.   Turner, T. L. & Miller, P. M. Investigating natural variation in Drosophila courtship song by the evolve and resequence approach. *Genetics* **191,** 633–642 (2012).

17.   Jha, AR, Miles, CM, Lippert, NR & Brown, CD. Whole-genome resequencing of experimental populations reveals polygenic basis of egg-size variation in Drosophila melanogaster. *… biology and evolution* (2015). at <http://mbe.oxfordjournals.org/content/32/10/2616.short>

18.   Cao, CC & Sun, X. Accurate estimation of haplotype frequency from pooled sequencing data and cost-effective identification of rare haplotype carriers by overlapping pool sequencing. *Bioinformatics* (2014). at <https://academic.oup.com/bioinformatics/article-abstract/31/4/515/2748165>

19.　　Long, Q. *et al.* PoolHap: inferring haplotype frequencies from pooled samples by
next generation sequencing. *PLoS ONE* **6,** e15292 (2011).
20.　　Pirinen, M. Estimating population haplotype frequencies from pooled SNP data
using incomplete database information. *Bioinformatics* (2009).
doi:10.1093/bioinformatics/btp584
21.　　Jajamovich, G. H., Iliadis, A., Anastassiou, D. & Wang, X. Maximum-parsimony
haplotype frequencies inference based on a joint constrained sparse representation of
pooled DNA. *BMC Bioinformatics* **14,** 270 (2013).
22.　　Kessner, D., Turner, T. & Novembre, J. Maximum Likelihood Estimation of
Frequencies of Known Haplotypes from Pooled Sequence Data. *Molecular Biology and*
Evolution **30,** (2013).
23.　　Franssen, S. U., Barton, N. H. & Schlötterer, C. Reconstruction of Haplotype-
Blocks Selected during Experimental Evolution. *Mol. Biol. Evol.* **34,** 174–184 (2017).
24.　　Lynch, M, Bost, D, Wilson, S & biology and …, M.-T. Population-genetic
inference from pooled-sequencing data. *Genome biology and …* (2014). at
<https://academic.oup.com/gbe/article-abstract/6/5/1210/604081>
25.　　Turner, TL, Stewart, AD, Fields, AT & Rice, WR. Population-based
resequencing of experimentally evolved populations reveals the genetic basis of body
size variation in Drosophila melanogaster. *PLoS …* (2011).
doi:10.1371/journal.pgen.1001336
26.　　Feder, A., Petrov, D. & Bergland, A. LDx: Estimation of Linkage Disequilibrium
from High-Throughput Pooled Resequencing Data. *PLoS ONE* **7,** (2012).
27.　　Kolaczkowski, B, Kern, AD, Holloway, AK & Genetics, B.-D. Genomic
Differentiation Between Temperate and Tropical Australian Populations of Drosophila
melanogaster. *Genetics* (2011). at <http://www.genetics.org/content/187/1/245.full-
text.pdf+html>
28.　　Schirmer, M., D'Amore, R., Ijaz, U. Z., Hall, N. & Quince, C. Illumina error
profiles: resolving fine-scale variation in metagenomic sequencing data. *BMC*
bioinformatics **17,** 125 (2016).
29.　　Huang, W, Massouras, A, Inoue, Y & Peiffer, J. Natural variation in genome
architecture among 205 Drosophila melanogaster Genetic Reference Panel lines.
Genome … (2014). at <http://genome.cshlp.org/content/24/7/1193.short>
30.　　Comeron, J. M., Ratnappan, R. & Bailin, S. The many landscapes of
recombination in Drosophila melanogaster. *PLoS genetics* **8,** e1002905 (2012).
31.　　Rockman, MV & Kruglyak, L. Recombinational landscape and population
genomics of Caenorhabditis elegans. *PLoS genetics* (2009).
doi:10.1371/journal.pgen.1000419
32.　　Zdraljevic, S, Roberts, JP & acids , A.-E. CeNDR, the Caenorhabditis elegans
natural diversity resource. *Nucleic acids …* (2017). at
<https://academic.oup.com/nar/article-abstract/45/D1/D650/2770657>
33.　　Nouhaud, P., Tobler, R., Nolte, V. & Schlötterer, C. Ancestral population
reconstitution from isofemale lines as a tool for experimental evolution. *Ecol Evol* **6,**
7169–7175 (2016).
34.　　Lack, JB, Lange, JD & biology …, T.-A. A Thousand Fly Genomes: An
Expanded Drosophila Genome Nexus. *Molecular biology …* (2016).
doi:10.1093/molbev/msw195

35. Teotónio, H, Chelo, IM, Bradić, M, Rose, MR & Long, AD. Experimental evolution reveals natural selection on standing genetic variation. *Nature genetics* (2009). at <https://www.nature.com/articles/ng.289>

36. Pettersson, ME & Honaker, CF. Standing genetic variation as a major contributor to adaptation in the Virginia chicken lines selection experiment. *Genome …* (2015). at <https://genomebiology.biomedcentral.com/articles/10.1186/s13059-015-0785-z>

37. Turchin, MC, Chiang, C. & Palmer, CD. Evidence of widespread selection on standing variation in Europe at height-associated SNPs. *Nature …* (2012). at <https://www.nature.com/articles/ng.2368>

38. Behrman, EL, Howick, VM & B, K.-M. Rapid seasonal evolution in innate immunity of wild Drosophila melanogaster. *… . R. Soc. B* (2018). doi:10.1098/rspb.2017.2599

39. Kessner, D. & Novembre, J. forqs: forward-in-time simulation of recombination, quantitative traits and selection. *Bioinform Oxf Engl* **30,** 576–7 (2013).

40. Money, D, Gardner, K, Genes, M.-Z. & Genomes . LinkImpute: Fast and accurate genotype imputation for nonmodel organisms. *G3: Genes* (2015). doi:10.1534/g3.115.021667

41. Yang, Y *et al.* A new genotype imputation method with tolerance to high missing rate and rare variants. *PloS one* (2014). doi:10.1371/journal.pone.0101025

42. Browning, BL & of Genetics, B.-S. A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *The American Journal of Human Genetics* (2009). at <http://www.sciencedirect.com/science/article/pii/S0002929709000123>

Methods

Establishment and sequencing of founder set

As described in (Behrman et al., 2018) [38] , 207 iso-female lines were established from wild individuals sampled from Maine and Pennsylvania in spring and autumn, and inbred for ~20 generations of full-sib mating to produce viable, fertile inbred lines. 30-50 individuals from each line were pooled for DNA extraction. Whole flies were homogenized with lysis buffer and 1mm beads and DNA was precipitated from the homogenate before resuspension in TE buffer. Libraries were prepared with a modified Nextera protocol developed by (Baym, et al 2015). All samples were indexed with Illumina's TruSeq Dual Index Sequencing Primer Kit (PE-121-1003) and pooled equimolarly into 3 sets of ~70 samples each. Each set of pooled DNA libraries were purified using Ampure XP and size-selected to 450-500 bp with a SizeSelect E-Gel. After an additional 5 rounds of PCR, DNA libraries were purified using Ampure XP beads, quantified, and diluted to the appropriate concentration before sequencing on the HiSeq 3000. All sequences were deposited in SRA (BioProject PRJNA383555). Adapter sequences were trimmed (Trimmomatic v0.33) and overlapping reads were merged (PEAR v0.9.6), then reads were mapped (bwa v 0.7.9) to the *D.melanogaster* reference genome (v5.39) using default parameters. PCR duplicates were removed using PicardTools (v1.12). Base quality score recalibration, indel realignment, and novel SNP discovery were carried out using GATK's Unified Genotyper (version 3.4-46), on all sequenced inbred lines together with a set of 143

previously sequenced *Drosophila melanogaster* lines derived from multiple Europe and North America populations (see Table S2 for SRA accessions). Only SNPs segregating in the 99 inbred strains pooled for re-sequencing were used to simulate reads and estimate haplotype frequencies.

Generating Experimentally Pooled Samples
One male each was selected from each of 99 inbred strains, and all individuals were pooled for re-sequencing. A second biological replicates was constructed from 99 additional individuals. DNA isolation was performed as described above. 3 separate libraries were prepared from each of the two biological replicates using different library prep methods: [1] according to protocols described in Nextera DNA Library Prep Reference Guide (15027987 v01); [2] a modified Nextera protocol (as described above); [3] a Covaris shearing protocol. Final results from the 3 library prep methods were similar. All libraries were size-selected and PCR amplified using two replicate PCR reactions and a high volume of template DNA to prevent PCR-jackpotting. DNA was purified, quantified, and diluted before sequencing on the HiSeq 3000. Raw, 150bp pair-end reads were trimmed for adapter sequences with Skewer (version 0.1.127). Read merging, mapping, and PCR duplicate removal was performed as above.

Generating Simulated Pooled Samples
150-bp paired end pre-aligned reads were simulated from a table of alternate founder genotypes and the *D. melanogaster* reference genome with simreads, a software tool included with the HARP package. All reads were simulated with an error rate of 0.2%. No read trimming or PCR duplicate removal was done. All SNP tables used to generate reads underwent imputation before read simulation.

Haplotype Frequency Estimation
All haplotype likelihoods and frequencies were estimated with HARP. Haplotype frequencies were evaluated in 1000kb, 100kb or 10kb window width sizes, with 100kb, 10kb, or 1kb window step sizes, respectively. SNP tables used to assign haplotype frequencies were re-imputed separately from any SNP table used to simulate reads. For reference, inferring haplotype frequencies for 99 founder lines at 283k segregating sites on chromosome 2L in 1000kb windows took 8 minutes and required 450Mb RAM for samples sequenced at 5x empirical coverage and took 15 minutes and required 860Mb RAM for samples sequenced at 10x. Using 100kb windows took 9.5 minutes / 70Mb and 17.5 minutes / 132Mb for 5x and 10x samples, respectively.

HAF Estimations
Allele frequencies calculated using haplotype inference (HAFs) were estimated as the sum of founder haplotypes containing the alternate allele, each weighted by their average inferred haplotype frequency in all haplotype inference windows overlapping the site. Founder haplotypes with missing calls were given a fractional alternate allele count equal to the mean of called founders with alternate alleles.

Accuracy Estimations Using Effective Coverage
Effective coverage was used as a metric to assess accuracy of all HAFs and raw AFs. Effective coverage was calculated by equating the total theoretical binomial variance (*BV*) of the true frequencies given an average coverage C [where BV= $\sum \frac{pp(1-pp)}{cC}$] to the sum of the squared error (*SSE*) of estimated allele frequencies. Solving for C with $C = \frac{\sum pp(1-p)}{\text{SSE}}$ yields the theoretical coverage at which binomial sampling of reads would be expected to contain the observed amount of error from estimated frequencies.

HAFs with an alternate founder set

SNPs were derived from 205 strains initially isolated from Raleigh, NC that were independently sequenced as part of freeze 2 of the *Drosophila* Genetic Reference Panel (DGRP) [29] . Genotype data was downloaded directly from http://dgrp2.gnets.ncsu.edu and read simulation, haplotype inference and effective coverage calculations were carried out as above.

Recombination

Forward-in-time simulations of recombination were performed with the software tool forqs [39] using a *D. melanogaster* recombination map [30] . As a conservative metric, here we refer to the female *D. mel* recombination rate. Since male *D mel*. do not undergo recombination, our estimates of the number of recombination events per generation are higher than that expected in real populations and our estimates of effective coverage serve as a lower bound on effective coverage expected at the same number of generations in real populations. Recombination breakpoints were simulated for up to 200 generations of the same evolutionary trajectory in 10 replicate populations with a constant population size of 1000 individuals. When selected sites were added, each replicate contained the same parameters for selection but included different selected sites. 198 recombination breakpoints were used to construct 'sampled' sets of chromosome pairs from corresponding segments of the 99 individually sequenced founder haplotypes. Read simulations were performed as above with recombined sets of SNPs. True allele frequencies were calculated as above from the same set of SNPs that were used to generate simulated reads.

Founder Ambiguity

To simulate reads with founder ambiguity, SNP calls from additional North American lines were added to calls from the 99 seasonal inbred lines pooled for sequencing. Additional sets of lines were then added to obtain SNP calls from 124, 149 and 209 lines respectively. SNP tables were also made by removing 25 and 50 lines from the original set of 99 lines. Each SNP table contained the same number of SNPs and was imputed as above. Reads were simulated from each new SNP table and further downstream analysis was conducted as described above.

Code Accessibility

Scripts to calculate HAFs are available at https://github.com/petrov-lab/HAFpipe-line . At minimum, the pipeline requires a) called biallelic variants from sequenced founder lines, and b) mapped reads from one or more pool-seq samples, and uses harp for haplotype inference.

Statement on Data and Reagent Availability

Sequence data from seasonal strains is available at SRA (BioProject PRJNA383555) and genotype data is available at *https://github.com/petrov-lab/HAFpipe-line/blob/master/99.clean.SNP.HARP.segregating*. Strains are available upon request. Code used to generate the simulated data is provided at https://github.com/petrov-lab/HAFpipe-line/tree/master/simulations .

Supplemental Text

Incorporating uneven pooling of individuals produces more realistic estimates of true allele frequencies

Our ability to measure the accuracy of HAFs and raw AFs depends on our ability to determine the true contribution of each pooled individual. Since uneven pooling is a source of error known to affect pool-seq samples[12], we estimated the relative contribution of DNA from each individual by calculating the average genome-wide allele frequency at sites private to each founder. While each founder could be detected in the pool, we found substantial variation in their relative representation (Supp. Fig. 2). 'True' frequencies for the experimental pooled sample were thus calculated by weighting founders known to contain the alternate allele by their estimated representation in the pool. We assessed whether these 'true' allele frequencies were better recapitulated by experimental reads than 'true' allele frequencies calculated without incorporating uneven pooling at all fully genotyped sites (both private and common). We found that the effective coverage using unevenly pooled weighted values (126x) was higher than the effective coverage assuming evenly pooled individuals (120x). We used these same estimates of uneven pooling to simulate reads in uneven proportions from different haplotypes for the synthetic sample as well.

Imputing missing founder genotypes increases the accuracy of HAFs

While missing information can be accommodated by many haplotype inference tools (i.e. an N in place of a missing call), it is unclear how missing calls affect inference accuracy, and what the best practices should be when missing calls are present in the reference founder set.

We first examined whether haplotype frequencies estimated for founders with many missing calls or few missing calls systematically deviated from an expected haplotype frequency of 0.101 (1/99). We found that across individual inference windows, there was a clear negative correlation between the number of missing calls per founder, and the haplotype frequencies estimated for that founder (Supp. Fig. 1). To determine whether the observed skewed haplotype frequencies were directly associated with the presence of missing sites, we tested whether imputing genotype calls for missing sites would reduce bias in haplotype frequency assignment. While a number of sophisticated methods for imputing rare SNPs do exist [40–42], here we used a simple approach. To perform imputation, at each site we first calculated the allele frequency among called founder genotypes and used this value as a probability for assigning genotypes to missing calls. We found that imputation significantly reduced the skewed haplotype frequency distribution by 4-6-fold for all empirical coverages and window sizes tested. We expect that imputation with more advanced tools would achieve even better results.

We next examined how imputation of haplotype frequencies can impact the overall accuracy of HAFs. We found that imputation increased the accuracy of HAFs in all

1 window sizes, but was most effective in 1000kb windows where accuracy increased by
2 more than two-fold at higher empirical coverages (Supp. Table 2). We also confirmed
3 that haplotype inference using imputed calls produced more accurate HAFs than using
4 a subset of sites with no missing calls. Thus, we include imputation as a key step in our
5 analysis pipeline.

Supplemental Figures and Tables

7
8 **Supplemental Figure 1.** Contribution of DNA from each pooled individual in experimental
9 replicate 1, estimated by average genome-wide allele frequency across all singleton sites. The
10 dashed line represents theoretical expectation for evenly pooled individuals. Points are colored
11 by number of singletons sites per founder.

1
2 **Supplemental Figure 2**. Median haplotype frequency across all windows on chromosome 2L
3 for each founder (n=99), calculated with different window sizes and empirical coverages.
4 Haplotype frequencies calculated before imputation (filled circles) and after imputation (open
5 circles) are plotted as a function of the log of the total number of ambiguous genotypes (aka "N-
6 count"). Best fit lines for each dataset were calculated with standard linear regression.
7
8
9

1
2 **Supplemental Figure 3.** Effective coverage values of HAFs using 1000kb (dark red), 100kb
3 (red) and 10kb (pink) inference windows, and raw AFs (black) within simulated pooled samples
4 using a DGRP founder set (n=205) on chromosome 2L.
5

1
2 **Supplemental Figure 4.** Effective coverage values of HAFs calculated with 1000kb inference
3 windows in 10 replicate populations after 0 to 200 generations of neutral recombination.
4
5

Supplemental Figure 5. Effective coverage of simulated samples from populations under a) the standard neutral selection regime, compared to populations under different selection regimes including b) 4 randomly distributed strongly selected sites contributing additively to a quantitative trait, c) same as panel b but with 50% reduced selection strength, and d) same as panel b but with twice as many selected sites. In each panel, effective coverage was calculated for 10 simulated replicate populations sampled at generations 5 (red boxplot) and 15 (yellow boxplot) and sequenced at an empirical coverage of 5x. Effective coverages from the same replicate are connected by a gray line.

Supplemental Figure 6. Effective coverage as a function of the expected number of breakpoints per sampled chromosome and the expected number of informative reads per founder line, in each haplotype inference window. While all simulation parameters and results reported here were derived from a panel of 99 *D. melanogaster* lines, "generations" and "empirical coverage" are easily translated to other founder panels and model systems with known recombination rates and SNP densities. ˙*Comeron et al., 2012;* ˙˙*Rockman et al., 2009;* *http://dgrp2.gnets.ncsu.edu/ ; **https://www.elegansvariation.org

Window Size	Coverage	Ambiguous	Imputed	Subsetted
10kb	1	9.7085	10.9811	10.5440
	5	34.0194	39.8801	37.3980
	10	58.4309	70.9542	63.7056
100kb	1	51.3650	60.5870	51.3393
	5	164.6956	215.1379	141.2237
	10	240.8699	345.4896	189.6102
1000kb	1	202.5682	329.7044	172.0310
	5	321.9890	701.5851	240.6798
	10	346.1369	809.1494	248.5619

Supplemental Table 1. Effective coverage values of HAFs obtained by different methods for treating missing founder genotype calls. Column 1 ('Ambiguous') refers to effective coverages obtained by performing haplotype inference with missing genotypes denoted by 'N'. Column 2 ('Imputed') refers to effective coverages obtained by performing haplotype inference with genotypes assigned to missing calls by randomly selecting the reference or alternate allele with a probability determined by the ratio of *called* ref and alt alleles at the site. Column 3 ('Subsetted') refers to effective coverages obtained by performing haplotype inference using only sites with full genotype information for every founder.

Population	Sample	SRA Accession
Birmingham_AL	21_36	SRR1820252
Birmingham_AL	21_39	SRR1820253,SRR1820254,SRR1820255,SRR1820256,SRR1820257,SRR1820258,SRR1820259,SRR1820260,SRR1820261
BullocksHarbor_BerryIslands	40_10	SRR1820282,SRR1820283,SRR1820284,
BullocksHarbor_BerryIslands	40_23	SRR1820288
CO	CO1	SRR189041
CO	CO10N	SRR189044
CO	CO13N	SRR189045
CO	CO14	SRR189046
CO	CO15N	SRR189047
CO	CO2	SRR189048
CO	CO4N	SRR189050
CO	CO8N	SRR189053
CO	CO9N	SRR306633
Cockburn_SanSalvador	42_23	SRR1820294
Cockburn_SanSalvador	42_20	SRR1820318,SRR1820319,SRR1820320,SRR1820321
dgrp	DGRP_100	SRR833244
dgrp	DGRP_319	SRR834508
dgrp	DGRP_31	SRR834509
dgrp	DGRP_32	SRR834512
dgrp	DGRP_348	SRR834514
dgrp	DGRP_385	SRR834518
dgrp	DGRP_390	SRR834519
dgrp	DGRP_397	SRR834522
dgrp	DGRP_189	SRR834523
dgrp	DGRP_223	SRR834527
dgrp	DGRP_382	SRR834552
dgrp	DGRP_361	SRR834553
dgrp	DGRP_48	SRR835034
dgrp	DGRP_505	SRR835039
dgrp	DGRP_528	SRR835043
dgrp	DGRP_530	SRR835044
dgrp	DGRP_551	SRR835047
dgrp	DGRP_843	SRR835059
dgrp	DGRP_849	SRR835060
dgrp	DGRP_850	SRR835061
dgrp	DGRP_913	SRR835077
dgrp	DGRP_634	SRR835086
dgrp	DGRP_627	SRR835097
dgrp	DGRP_630	SRR835098
dgrp	DGRP_360	SRR835260
dgrp	DGRP_748	SRR835336
dgrp	DGRP_821	SRR835347
dgrp	DGRP_584	SRR835942
dgrp	DGRP_900	SRR846989
dpgp3	ZI157	SRR202089
dpgp3	ZI188	SRR202123
dpgp3	ZI184	SRR203068
dpgp3	ZI181	SRR203069
dpgp3	ZI103	SRR203226
dpgp3	ZI129	SRR203232
dpgp3	ZI179	SRR203233
dpgp3	ZI173	SRR203330
dpgp3	ZI126	SRR203336
dpgp3	ZI182	SRR203473
dpgp3	ZI138	SRR203474
dpgp3	ZI178	SRR203496
dpgp3	ZI10	SRR203502
dpgp3	ZI136	SRR204007
dpgp3	ZI190	SRR204018
dpgp3	ZI170	SRR210787
dpgp3	ZI176	SRR210794
dpgp3	ZI165	SRR248129
dpgp3	ZI117	SRR248130
dpgp3	ZI152	SRR326790
dpgp3	ZI177	SRR326796
dpgp3	ZI172	SRR326800
dpgp3	ZI134N	SRR326933
dpgp3	ZI164	SRR346921
dpgp3	ZI161	SRR353759
dpgp3	ZI104	SRR654551
dpgp3	ZI167	SRR654553

Population	Sample	SRA Accession
dpgp3	ZI114N	SRR654662
dpgp3	ZI118N	SRR654664
dpgp3	ZI185	SRR654669
France	FR14	SRR189088
France	FR151	SRR189089
France	FR180	SRR189090
France	FR207	SRR189091
France	FR217	SRR189092
France	FR229	SRR189093
France	FR310	SRR189094
France	FR361	SRR189095
France	FR70	SRR189096
Freeport_GrandBahamasWest	33_11	SRR1820273,SRR1820274
Freeport_GrandBahamasWest	33_16	SRR1820275,SRR1820276,SRR1820277,SRR1820278,SRR1820279
GeorgeTown_Exumas	36_12	SRR1820280
GeorgeTown_Exumas	36_9	SRR1820281
GU	GU10	SRR189114
GU	GU2	SRR189117
GU	GU6	SRR189120
GU	GU7	SRR189121
GU	GU9	SRR189122
Ithica	I01	SRR1663542
Ithica	I02	SRR1663543
Ithica	I03	SRR1663544
Ithica	I04	SRR1663545
Ithica	I06	SRR1663546
Ithica	I07	SRR1663547
Ithica	I13	SRR1663548
Ithica	I16	SRR1663549
Ithica	I17	SRR1663550
Ithica	I22	SRR1663551
Ithica	I23	SRR1663552
Ithica	I24	SRR1663553
Ithica	I26	SRR1663554
Ithica	I29	SRR1663555
Ithica	I31	SRR1663556
Ithica	I33	SRR1663557
Ithica	I34	SRR1663558
Ithica	I35	SRR1663559
Ithica	I38	SRR1663560
Mayaguana_Mayaguana	43_18	SRR1820302
Mayaguana_Mayaguana	43_19	SRR1820303
Meridian_MS	24_2	SRR1820262,SRR1820263,SRR1820264,SRR1820265,SRR1820266,SRR1820267
Meridian_MS	24_9	SRR1820309,SRR1820310,SRR1820311,SRR1820312,SRR1820313,SRR1820314,SRR1820315,SRR1820316,SRR1820317
NL	Netherlands01	SRR1663561
NL	Netherlands02	SRR1663562
NL	Netherlands03	SRR1663563
NL	Netherlands04	SRR1663564
NL	Netherlands07	SRR1663565
NL	Netherlands10	SRR1663566
NL	Netherlands11	SRR1663567
NL	Netherlands13	SRR1663568
NL	Netherlands14	SRR1663569
NL	Netherlands15	SRR1663570
NL	Netherlands16	SRR1663571
NL	Netherlands17	SRR1663572
NL	Netherlands18	SRR1663573
NL	Netherlands19	SRR1663574
NL	Netherlands22	SRR1663575
NL	Netherlands23	SRR1663576
NL	Netherlands25	SRR1663577
NL	Netherlands29	SRR1663578
NL	Netherlands30	SRR1663579
PortAuPrince_Haiti	H_25	SRR1820304,SRR1820305,SRR1820306,SRR1820307,SRR1820308
PortAuPrince_Haiti	H_29	SRR1820322,SRR1820323,SRR1820324,SRR1820325
Sebastian_FL	28_8	SRR1820268,SRR1820269,SRR1820270,SRR1820271,SRR1820272
Selba_AL	20_17	SRR1820244,SRR1820245,SRR1820246,SRR1820247,SRR1820248,SRR1820249
Selba_AL	20_28	SRR1820250,SRR1820251
TampaBay_FL	4_12	SRR1820289,SRR1820290,SRR1820291,SRR1820292,SRR1820293
TampaBay_FL	4_27	SRR1820295,SRR1820296,SRR1820297,SRR1820298,SRR1820299,SRR1820300,SRR1820301,SRR1845494,SRR1845495,SRR1845496
Thomasville_GA	13_29	SRR1820240,SRR1820241
Thomasville_GA	13_34	SRR1820242,SRR1820243

1
2 **Supplemental Table 2.** SRA accessions for sequenced global *D. melanogaster lines* used for
3 joint variant calling.