Title

Pyrimidine starvation activates a bistable phenotypic switch leading to ribosome provisioning and
rapid exit from stationary phase

Authors:

P. Remigi[1], G.C. Ferguson[2], S. De Monte[3] and P.B. Rainey[1,4,5]

[1] New Zealand Institute for Advanced Study, Massey University, Auckland 0745, New Zealand

[2] Institute of Natural and Mathematical Sciences, Massey University, Auckland 0745, New Zealand

[3] Institut de Biologie de l'Ecole Normale Superieure, UMR CNRS 8197 INSERM 1024, F-75005 Paris,
France

[4] Department of Microbial Population Biology, Max Planck Institute for Evolutionary Biology, Plön
24306, Germany

[5] Ecole Superieure de Physique et de Chimie Industrielles de la Ville de Paris (ESPCI Paris Tech), CNRS
UMR 8231, PSL Research University, 75231 Paris, France

Corresponding authors:

Philippe Remigi (p.remigi@massey.ac.nz) and Paul B. Rainey (rainey@evolbio.mpg.de)

Abstract:

Observations of bacteria at the single-cell level have revealed many instances of phenotypic

heterogeneity within otherwise clonal populations, but the selective causes, molecular bases and

broader ecological relevance remain poorly understood. In an earlier experiment in which the

bacterium *Pseudomonas fluorescens* SBW25 was propagated under a selective regime that mimicked

the host immune response, a genotype evolved that stochastically switched between capsulation

states. The genetic cause was a mutation in *carB* that decreased the pyrimidine pool (and growth

rate) lowering the activation threshold of a pre-existing but hitherto unrecognised phenotypic

switch. Genetic components surrounding bifurcation of UTP flux towards DNA/RNA or UDP-glucose

(a precursor of colanic acid forming the capsules) were implicated as key components. Extending

these molecular analyses – and based on a combination of genetics, transcriptomics, biochemistry

and mathematical modelling – we show that pyrimidine limitation triggers an increase in ribosome

biosynthesis and that switching is caused by competition between ribosomes and CsrA/RsmA

proteins for the mRNA transcript of a feed-forward regulator of colanic acid biosynthesis. We

additionally show that in the ancestral bacterium the switch is part of a programme that determines

stochastic entry into the semi-quiescent capsulated state, ensures that such cells are provisioned

with excess ribosomes, and enables provisioned cells to exit rapidly from stationary phase under

permissive conditions.

Introduction:

Phenotypic variation between isogenic cells growing in homogeneous environments can have adaptive consequences, allowing populations to survive unpredictable environmental changes or promoting interactions between different cell types (ACKERMANN 2015; VAN BOXTEL et al. 2017). Natural selection, by means of genetic mutations affecting the integration of stochastic noise within signaling pathways, can fine-tune epigenetic switches (ROTEM et al. 2010; FRIDMAN et al. 2014; NEW et al. 2014; RICHARD AND YVERT 2014; VAN DEN BERGH et al. 2016; BODI et al. 2017) but the molecular details underpinning the evolution of phenotypic heterogeneity remain poorly understood.

An opportunity to study the genetic bases of the evolution of phenotypic heterogeneity arose from a selection experiment where the capacity to switch between different colony phenotypes evolved *de novo* in the bacterium *Pseudomonas fluorescens* SBW25 (BEAUMONT et al. 2009). In this experiment, bacteria were passaged through consecutive cycles comprised of single-cell bottlenecks and negative frequency-dependent selection, a regime mimicking essential features of the adaptive immune system in animals. A strain emerged (1B4) that forms distinct opaque or translucent colonies on agar plates. At the single-cell level, this behaviour reflects an epigenetic switch characterized by the bistable production of an extracellular capsule, resulting in the coexistence of two sub-populations of capsulated (Cap+) or non-capsulated (Cap-) cells. Capsules are made of a colanic acid-like polymer, whose production arises from the activity of the *wcaJ-wzb* locus and requires the precursor UDP-glucose. A mutation in *carB*, a gene involved in *de novo* pyrimidine biosynthesis, is responsible for heterogeneous capsule production via a decrease in intracellular pyrimidine pools (BEAUMONT et al. 2009; GALLIE et al. 2015). Additionally, the switch between capsulated and uncapsulated cells was shown to function in the ancestral gentoype devoid of the *carB* mutation – the *carB* mutation having altered the threshold at which the switch is activated – and underpins the stochastic entry of cells into a semi-quiescent state.

Here we extend earlier work and present a detailed mechanistic model that explains how the *carB* mutation impacts on capsulation heterogeneity. Central to the model is evidence that ribosome biosynthesis is up-regulated upon pyrimidine limitation and that this favors translation of a positively auto-regulated activator of capsular exopolysaccharide biosynthesis that is otherwise inhibited by CsrA/Rsm proteins. The switch comprises part of a programme that facilitates stochastic entry into a semi-quiescent state and rapid exit from this state upon realisation of permissive conditions through modulation of ribosome levels.

Results

Capsulation is not induced by UDP-glucose depletion

The paradoxical observation that pyrimidine limitation induced capsulation, a phenotypic trait

depleting the pyrimidine pool even more through the consumption of UDP-glucose, caused

attention to focus on the link between capsulation, growth and pyrimidine metabolism (GALLIE *et al.*

2015). In particular, the fact that capsulation may represent a semi-quiescent state whose initiation

is triggered by fluctuations in the concentration of an intermediate of the pyrimdine biosynthetic

pathway suggested that this metabolite may act both as a signal for capsulation and as a cell cycle

checkpoint. UDP-glucose appeared a prime candidate for this function, since it is a widely conserved

signal regulating cell size and bacterial growth (VADIA AND LEVIN 2015). A transposon insertion in *galU*

(encoding the enzyme required for UDP-glucose biosynthesis) prevented capsulation (GALLIE *et al.*

2015), but a role for UDP-glucose as a building block for colanic acid or as a signal for its activation

remained undetermined. Introduction of a translational reporter of capsulation in the *galU* mutant

(with undetectable UDP-glucose levels; GALLIE *et al.* 2015) revealed that the proportion of cells

expressing this reporter was lower in the mutant compared to 1B4 (Supplementary Figure 1). This is

in contrast to the predicted response if a reduction in the concentration of UDP-glucose was the

signal inducing capsulation. This key result caused us to revisit the mechanistic links between

pyrimidine limitation, growth rate and the heterogeneous production of colanic acid capsules.

Ribosomes are over-produced in *carB* mutants

We re-analysed transcriptomic data (published in GALLIE *et al.* 2015) to identify signalling pathways

that may display different levels of activity as a result of the *carB* mutation. KEGG-enrichment

analyses showed over-representation of ribosomal components among the genes that are expressed

at least two-fold more in the capsulated sub-population of 1B4 (1B4 Cap+) compared to its (non-

capsulated) ancestor 1A4 (Supplementary Table 1). By extracting raw expression values from the

available RNAseq datasets, we observed that the average expression levels of ribosomal protein

genes increase in 1B4 (either Cap- or Cap+) compared to 1A4 or SBW25 (Fig. 1a). Such a finding is

surprising since ribosome production is usually proportional to growth rate (DENNIS *et al.* 2004; PAUL

et al. 2004) and was expected to be reduced in the slower growing strain 1B4 (Supplementary Figure

2)**.**

In bacteria, correct adjustment of ribosomal production to nutrient availability is essential to

maximise growth rate. Bacteria do so by modulating transcriptional activity at the promoters of

ribosomal RNA (*rrn*) operons (DENNIS *et al.* 2004; PAUL *et al.* 2004). The production of ribosomal

proteins is then adjusted to the level of available rRNA (KEENER AND NOMURA 1996). The over-

expression of ribosomal protein genes in 1B4 may therefore reflect transcriptional up-regulation at

rrn promoters. Using a chromosomally-integrated reporter, an increase in P*rrnB*-GFP transcriptional

activity was detected in *carB* mutants compared to immediate ancestral types (Fig. 1b). This

difference was most obvious when cultures reached OD~2, a density at which 1B4 cultures undergo

a noticeable increase in capsulation (Supplementary Figure 3). Supplementing growth media with

2mM uracil – a treatment known to suppress capsulation (GALLIE *et al.* 2015) – restored wild-type

P*rrnB*-GFP expression in strains carrying the mutant *carB* allele (Fig. 1b). Together with the higher

total RNA concentration measured in *carB* mutants (Fig. 1c), these results show that pyrimidine

limitation triggers the production of an enhanced pool of active ribosomes.

High ribosome levels are required for capsulation

The counterintuitive effect of the mutation in *carB* on ribosome levels suggested a causal connection

between ribosome concentration and capsulation. In support of this hypothesis, a previous

transposon-mutagenesis screen found that insertions in ribosome- or translation-associated genes

(*prfC, rluB, rluC, glu/gly* tRNA) decreased or abolished capsulation in 1B4 (GALLIE *et al.* 2015). We set

out to manipulate ribosome concentration in 1B4 in order to test directly if ribosome abundance has

an impact on capsulation. *P. fluorescens* SBW25 harbours 5 *rrn* copies (*rrnA-E*) and, in bacteria, the

deletion of a single operon can often be compensated for by the over-expression of those remaining

(CONDON *et al.* 1993; BOLLENBACH *et al.* 2009; GYORFY *et al.* 2015). Capsulation was quantified in single

or double *rrn* deletion mutants carrying the chromosomally-inserted translational reporter

P*pflu3655*-GFP (Gallie *et al.*, 2015) by measuring the proportion of GFP positive cells in cultures at

the onset of stationary phase (OD~2). Whereas single mutants were not significantly affected in their

capsulation levels, double mutants produced fewer capsulated cells (Fig. 2a and Supplementary

Figure 4). A reduction in RNA concentration (Supplementary Figure 5) was observed in the *rrnAE*

double mutant, but not in *rrnAC* that showed a milder capsulation phenotype. Together, these

results show that capsulation is induced by increased ribosome dosage, which is itself a consequence

of the *carB* mutation.

A ribosome-Rsm competition model for the control of capsulation

Next, we asked how ribosome abundance may influence the heterogeneous production of

capsulated cells in an isogenic population. Results from a previous transposon mutagenesis screen

indicated that the Gac/Rsm two-component signalling pathway is required for the production of

colanic acid-like capsules (GALLIE *et al.* 2015) . The Gac/Rsm signalling pathway controls important

ecological traits in many Gram-negative bacteria (VAKULSKAS *et al.* 2015; VALENTINI *et al.* 2017). Its

activity is mediated through post-transcriptional regulators of the Csr/Rsm family that can prevent

translation of mRNA targets by binding sites adjacent to or overlapping ribosome-binding sites (RBS)

(LAPOUGE *et al.* 2008; VAKULSKAS *et al.* 2015). The phenotypic effect of the Gac/Rsm pathway was

investigated by creating deletion mutants for the response regulator *gacA* and the two Csr/Rsm

homologs present in SBW25, namely *rsmA* and *rsmE.* Capsulation was completely abolished in the

gacA deletion strain, confirming the transposon-mutagenesis results (Fig. 2b). Deletion of *rsmA* or

rsmE increased the production of capsulated cells, consistent with their typical inhibitory role in

Gac/Rsm signalling pathways.

We postulated that variations in the relative concentration of free ribosomes and RsmA/E may

determine translational output of a key positive regulator of capsule biosynthesis (Fig. 2c). While

searching for such a regulator, attention turned to *PFLU3655*. The first gene of a putative operon

(*PFLU3655-3657*) localized just upstream of the colanic acid biosynthetic operon, *PFLU3655* is

annotated as a hypothetical protein carrying a two-component response regulator C-terminal

domain (PFAM PF00486). *PFLU3655* is among the most highly up-regulated genes in 1B4 Cap+ cells

and transposon insertions in its promoter were shown to abolish capsulation in 1B4 (GALLIE *et al.*

2015). A non-polar deletion of *PFLU3655* in 1B4 abolished capsule formation, while

complementation of the mutant with an IPTG-inducible copy of *PFLU3655* on a low copy number

plasmid (pME6032) restored capsulation (Fig. 3a). Over-expression of *PFLU3655* in SBW25 or 1B4

was sufficient to induce high capsulation levels in these strains, showing that PFLU3655 is a key

positive regulator of colanic acid biosynthesis, the expression of which is necessary and sufficient for

capsulation. Moreover, expression of PFLU3655 is also necessary and sufficient to induce the

expression of its own promoter, as measured with the chromosomally-encoded P*pflu3655*-GFP

translational fusion (Fig. 3b). This result shows that PFLU3655 expression can generate a positive

feedback loop, a motif that can sustain bistable gene expression (VEENING *et al.* 2008; NORMAN *et al.*

2015).

Ribosome-Rsm interaction at the PFLU3655 promoter

Existence of a positive fedback loop does not warrant bistability, which often requires the additional

presence of an ultrasensitive switch to convert small input deviations (typically, molecular noise)

into large output differences (FERRELL AND HA 2014). When signalling components are present in large

numbers, ultrasensitive responses and threshold effects can arise through molecular titration where

a molecule (RNA or protein) is sequestered and inhibited by another protein (BUCHLER AND LOUIS

2008; BUCHLER AND CROSS 2009; FERRELL AND HA 2014).

The possibility that titration of a target mRNA by RsmA/E may trigger ultrasensitivity was considered.

Interestingly, two putative Rsm binding sites are located in the promoter and 5' region of the coding

sequence of *PFLU3655* (Fig. 4a), indicating that PFLU3655 mRNA could be a direct target of RsmA/E.

The P*pflu3655*-GFP capsulation reporter was used to test this hypothesis. This translational reporter

contains ~500 nucleotides upstream of PFLU3655 start codon and the first 39 coding nucleotides

fused to GFP in frame; the two putative RsmA/E binding sites are conserved in this synthetic

construct. Using site-directed mutagenesis, nucleotides located in the putative Rsm binding sites

were substituted and effects on GFP production in the P*pflu3655*-GFP reporter determined (Fig.

4a,b). Both G-8A and A33T point mutations increased GFP production, albeit to a different extent,

consistent with the expected effects arising from reduction in binding of an inhibitor. Altering the

putative RBS (GG-7AC) completely abolished GFP production (Fig. 4b), which is similarly to be

expected given the need for translation. Re-introducing these point mutations at the original locus in

1B4 showed that the G-8A mutation, but not the A33T, increased the proportion of capsulated cells

(Fig. 4c). This difference mirrors the extent of presumed translational mis-regulation by the two

point mutations and suggests that the increase in PFLU3655 translation mediated by the A33T

mutation is not sufficient to increase the likelihood to jump-start the positive feedback loop,

contrary to G-8A.

Together, these data support the model proposed earlier (Fig. 2c). If this model is correct, one would

expect other RsmA/E targets to be over-expressed in *carB* mutants. To this end, a list of genes that

were differentially expressed in a SBW25 *gacS* mutant (CHENG *et al.* 2013) was extracted and their

expression levels were compared using the RNAseq dataset. On average, genes that were up-

regulated in the *gacS* mutant were expressed at lower levels in 1B4 (both Cap- and Cap+) than in

SBW25 or 1A4 (Supplementary Figure 6). Genes that were down-regulated in *gacS* showed a slight

bias towards higher expression in 1B4 Cap+ but this difference was not statistically significant. These

results are consistent with the opposing effects of *gacS* inactivation (leading to constitutive

activation of RsmA/E) and *carB*-dependent increase in ribosome concentration on RsmA/E targets.

A qualitative mathematical model of the genetic network controlling bistable capsule production explains experimental results

We produced a mathematical model based on our experimental observations in order to describe the qualitative behaviour of the cells, notably the bistability of the internal state, and to predict how changes in parameters are likely to affect the fraction of capsulated cells.

The abundance of PFLU3655, represented by the variable x, is described by the following ordinary differential equation:

$$\frac{dx}{dt} = P(x) - dx$$

where $P(x)$ is the production rate and the constant d measures the rate of degradation/dilution (or decay) per molecule. The existence of a positive feedback loop in PFLU3655 production and the plausible assumption that RsmA/E and ribosome compete to bind *PFLU3655* mRNA led us to choose an ultrasensitive curve for the production term $P(x)$ (Fig. 5a; see Supplementary text for full details on the model), analogous to other cases of molecular titration (BUCHLER AND LOUIS 2008; BUCHLER AND CROSS 2009; ZHANG *et al.* 2013). For fixed concentrations of RsmA/E and ribosomes, the shape of this curve is characterized by a nonlinear increase of production with x. For low x, even though *PFLU3655* is transcribed proportionally to x, sequestration of its mRNA by RsmA/E maintains translation, thus the concentration of the protein, at low levels. When the number of mRNA molecules exceeds the buffering capacity of the RsmA/E pool, then translation increases massively, but it eventually saturates for high values of x, due to the competition for the ribosome pool.

The equilibrium solutions of the system x_E are found when the production equalizes the decay ($P(x_E) = d\, x_E$). When the decay term is sufficiently small (dashed purple line in Fig. 5a), the system is bistable, with two stable (corresponding to the OFF ($x_E=0$) and the ON ($x_E=x^{on}$) states) and one unstable ($x_E=x^S$) equilibria. The unstable solution separates the space of concentrations x in two basins of attraction of the stable equilibria. Stochastic fluctuations in the components of the system can activate or inactivate capsulation by letting x change the basin of attraction and subsequently approach the corresponding equilibrium. The smaller the basin of attraction of the OFF equilibrium, (smaller x^S) the higher the probability that random fluctuations cause the system to switch to the ON state, and vice-versa. Thus, the position of x^S defines the proportion of cells found in the capsulated ON state.

Qualitative changes in the behaviour of the system can be explored by a graphical analysis of the equilibria. An increase in ribosome concentration leads to an increase in the maximal production rate $P(x)$, which can be sufficient to generate bistability (Fig. 5b, blue line). Starting from a bistable

situation (blue line), the probability of switching ON can be enhanced through an increase of the

basal production rate of PFLU3655 (Fig. 5c**,** light blue line). These observations qualitatively match

experimental results previously described (Fig. 3), indicating that this simple model encapsulates

fundamental aspects of the switching mechanism.

Consequences of ribosome accumulation on growth resumption in capsulated cells

A corollary of our model is that, if ribosome levels fluctuate between isogenic cells, cells with highest

ribosome concentrations should be more likely to produce capsules. Indeed, RNAseq data indicate

that ribosomal protein genes are expressed at higher levels in 1B4 cap+ compared to 1B4 cap- (Fig.

1a). When the quality and/or quantity of nutrients rises abruptly, differences in ribosome

abundance can be significant for bacterial fitness (MORI *et al.* 2017). Reaching higher ribosome

concentration – required for maximal growth rate after nutrient up-shift – is time-consuming and

means a time-delay between environment change and future growth (EHRENBERG *et al.* 2013). Cells

with higher ribosome concentrations before nutrient up-shift might be considered as provisioned for

rapid acclimation to the new conditions. If it is true that (i) capsulated cells have more ribosomes

and (ii) ribosomes promote growth resumption after nutrient up-shift, then capsulated cells should

have a fitness advantage in those conditions. To test this prediction, 1B4 cells grown to late

exponential phase (OD_{600nm}~1) and cell suspensions enriched in Cap- or Cap+ cells were used to

inoculate fresh cultures. The initial growth rate after nutrient up-shift in batch cultures was

positively correlated with the proportion of capsulated cells (Fig. 6a). Time-lapse microscopy on solid

agar pads confirmed that colonies founded by GFP+ (capsulated) cells grew approximately 10%

faster than those originating from GFP- (non-capsulated) cells (Fig. 6b).

Capsulation in SBW25

The identification of mechanisms promoting the production of colanic acid capsules in evolved

P. fluorescens genotypes raises questions as to the role and conditions for expression of colanic acid

in ancestral SBW25. In bacteria, extracellular capsules are important for bacterial pathogenicity

(ROBERTS 1996) and are associated with broader environmental versatility (RENDUELES *et al.* 2017).

Recent work has also demonstrated a role for colanic acid-like capsules in the positioning of the cells

at the surface of bacterial colonies, providing them with a better access to oxygen and a fitness

advantage (KIM *et al.* 2014; KIM *et al.* 2016).

In order to test if capsulation could occur in wild-type SBW25 colonies, we spot-inoculated bacterial

suspensions on agar plates and incubated the plates at 28°C for several days. Mucoid swellings were

observed from 5 days post-inoculation and increased over time (Fig. 7a). These swellings were not

observed with a colanic acid mutant and their appearance was delayed when 2mM uracil was added

to agar plates (Fig. 7b). In SBW25, cells sampled from mucoid regions showed a high proportion of

capsulated cells, approximately 60% of which expressed the Ppflu3655-GFP reporter (Fig 7c).

However, when streaked on new agar plates, no phenotypic difference was observed between cells

coming from these mucoid areas and regular SBW25 colonies (data not shown), indicating that

mucoidy was not dependent on *de novo* mutation. These results are in accord with predictions and

indicate that capsulation in starved SBW25 colonies is also a consequence of bistable colanic acid

production resulting from nutrient limitation.

Given previous results with 1B4, the possibility arose that capsulated SBW25 cells might also benefit

from a growth advantage upon exposure to rich medium. Cells were collected from 7-day old

colonies and enriched in capsulated or non-capsulated cells by gentle centrifugation. When

transferred to fresh batch cultures, cell suspensions enriched in capsulated cells showed a faster

initial growth rate (Supplementary Figure 7) that resulted in a fitness advantage during head-to-head

competition (Fig. 7d). Moreover, time-lapse microscopy revealed that micro-colonies founded by

GFP+ cells display an initial growth rate significantly higher than colonies arising from GFP- cells (Fig

7e).

To test if capsulation status in SBW25 is also associated with higher ribosome content, we measured

expression of the ribosomal protein gene *rpsL* in capsulated and non-capsulated cells originating

from old SBW25 colonies by RT-qPCR. We found an expression ratio of 1.67 (+/- 0.35 s.d., n=5, two-

tailed t-test: p=0.012) in cap+ vs. cap- cells, suggesting that SBW25 capsulated cells contain more

ribosomes than their non-capsulated counter-parts.

Overall, the results from experiments carried out in SBW25 are consistent with the capsulation

model proposed for 1B4. During 'long-term' (*i.e.*, 7 days) starvation, nutrient limitation (possibly flux

through the pyrimidine pathway and likely associated with spatial heterogeneity within colonies)

establishes a sub-population of cells with increased ribosome content. This population then has an

enhanced chance of flipping to the capsulated state where growth remains slow and cells enter a

semi-quiescent state. This same sub-population then stands primed for rapid growth upon nutrient

upshift.

Discussion:

Studies of adaptive phenotypes derived from selection experiments are by nature multifaceted, but two parallel lines of inquiry are of particular importance. The first concerns the nature of the adaptive phenotype, including its selective and molecular causes. The second concerns the ecological significance of the traits affected by adaptive evolution, prior to the occurrence of the adaptive mutation(s). A satisfactory answer to the first requires understanding of the latter.

Consider initial discovery of the 1B4 switching genotype (BEAUMONT et al. 2009): a remarkable behaviour that owes its existence to reduction of function of carbamoyl phosphate synthase (GALLIE et al. 2015). In and of itself discovery of the causal switch-generating mutation shed no light on the adpative phenotype. Understanding began to emerge upon recognition that the carB mutation had altered the activation threshold of a pre-existing switch.

Here we have substantially extended understanding of both the mechanistic bases of phenoytpic switching in the derived 1B4 genotype, and also the ecological significance of the underlying traits in the ancestral genotype. The former is specific to the selection experiment, whereas the latter has likely relevance of a more general nature. Indeed, as is often the case, studies of phenotypes derived from in vitro selection experiments can shed light on hitherto unrecognised aspects of bacterial physiology (HINDRE et al. 2012; RAINEY et al. 2017; and references therein).

With attention on the mechanistic basis of the 1B4 switch, experiments examining function of the switch in a galU mutant showed that the previous conclusions concerning the central role of UTP or related molecules required revision. While pyrimidine biosynthesis and the UTP decision point are important components of the pathways leading to capsulation, the mechanism of switching resides elsewhere. Data presented here have led to formulation of a compelling new model of the switch. Central to the proposed model (Fig. 2c) is titration of PFLU3655 mRNA by RsmA/E that has the potential to generate an ultrasensitive response. The increase in ribosome production resulting from the carB mutation brings the system closer to its activation threshold, where small fluctuations in ribosome and/or in RsmA/E activities initiate a positive feedback loop leading to capsulation. Given the prevalence of inhibitory interactions in signaling networks, pathways with these features offer a common route for the evolution of or evolutionary tuning of bistable switches through relatively minor changes to the basal level (or interaction affinity) of threshold-defining components (BUCHLER AND CROSS 2009; ROTEM et al. 2010).

Data supporting this revised model comes largely from experiments described here, but also incorporates data presented in Gallie et al. (2015) that at the time were inexplicable. Most notably

data indicating a role for ribosomes, which stemmed from transcriptome data, but also from analysis

of mutants defective in switching obtained from the original suppressor study. Here numerous non-

switching types carried transposon insertions in genes implicated in ribosome or translation-

associated activities. Additional mutants that remained unincorporated included that Gac pathway

and its connection to RsmA/E and the newly recognised regulator of colanic-acid production

(PFLU3655).

Understanding the switch requires understsanding of its role in the ancestral genotype. Previous

work showed that switching to the capsulated type was accompanied by a reduction in growth rate.

This work also showed that the probability of switching to the capsulated state was more likely in

starved cells (particularly in cells starved of pyrimidines). This was understood as a mechanism that

allowed cells entering starvation conditions to hedge their bets in the face of uncertainty

surrounding the future state of the environment (GALLIE *et al.* 2015). Discovery (from this work) that

capsulated cells, despite slow growth, are replete in ribosomes was perplexing, but caused attention

to focus on exit from the semi-quiescent state. Just as cells entering a slow growing phase stand to

be out-competed by conspecific types that remain in the active growth phase if the environment

unexpectedly returns to one conducive for growth, cells exiting from a slow growth state stand to be

out-competed by types that are already actively growing unless they can rapdily resume "life in the

fast lane".

Although the subject of little attention in the microbiological world (but see (MORI *et al.* 2017) and

below), the importance of provisioning future generations with resources sufficient to aid

estabishment is a central part of life-history evolution theory (STEARNS 1992). It is not difficult to

conceive that cells entering a slow or non-growing state, such as persistors, or the capsulated cells of

SBW25, will through evolutionary time, experience selection for mechanisms that facilitate rapid re-

entry to active growth. Our data here is strongly indicative of such an evolutionary response.

Nutrient starvation leads to stochastic activation of capsulation and entry into a semi-quiescent

growth phase. At the same time such capsulated cells contain above average concentrations of

ribosomes. Upon encountering conditions conducive for growth capsulated cells rapidly switch to

the non-capsulated state (GALLIE *et al.* 2015), but remarkably, out-compete non-capsulated cells. A

phenomenom of ribosome over-capacity that bears similarity to what we have interpreted as

'ribosome provisioning', was described previously in fast growing bacteria (KOCH 1971): production

of excess ribosomes during exponential growth, while being deleterious in the short term, can

provide an advantage during nutrient up-shift (MORI *et al.* 2017). In environments where resource

availability fluctuates, it appears that the control of ribosome biosynthesis might be subjected to a

trade-off between maximising growth rate before starvation and growth resumption upon nutrient

up-shift (MORI *et al.* 2017). Our findings add a single-cell perspective showing that cell-to-cell

heterogeneity in ribosome activity during stationary phase may contribute to maximisation of long-

term geometric mean fitness.

Material and methods:

Bacterial strains and growth conditions

Bacterial strains used in this study are listed in Supplementary Table 2. Unless otherwise stated,

Pseudomonas fluorescens strains were cultivated in King's Medium B (KB; KING *et al.* 1954)) at 28°C

with orbital shaking. *Escherichia coli* DH5α *λpir* was used for clonings and was grown on Lysogeny

Broth at 37°C. Bacteria were plated on their respective growth medium containing 1.5% agar.

Antibiotics were used at the following concentrations: ampicillin (50-100 µg mL^{-1}), gentamicin (10 µg

mL^{-1}), tetracycline (10 µg mL^{-1}), kanamycin (25 or 50 µg mL^{-1} for *E. coli* or *P. fluorescens*, respectively)

and nitrofuorantoin (100 µg mL^{-1}). Uracil (Sigma-Aldrich) was added to culture medium at 2mM final

concentration when indicated. For competition experiments, 5-bromo-4-chloro-3-indolyl-β -d-

galactopyranoside (X-gal) was used at a concentration of 60 mg L^{-1} in agar plates.

For capsulation assays, pre-cultures were inoculated from pre-calibrated dilutions of frozen glycerol

aliquots in order to reach an optical density of 0.3-0.5 after overnight culture.

For colony assays with SBW25, 5µl of cell suspensions were spot-inoculated on KB agar plates and

incubated for 7 days at 28°C. Cells from the center of these colonies were resuspended in PBS or

Ringer's solution for growth and competition assays and time-lapse microscopy, or in RNAlater

solution (Invitrogen) for RT-qPCR.

Molecular techniques

Oligonucleotides and plasmids used in this study are listed in Supplementary Tables 3 and 4,

respectively. Standard molecular biology techniques were used for DNA manipulations (SAMBROOK *et*

al. 1989). DNA fragments used to generate promoter fusions and gene deletion constructs were

prepared by splicing by overhang extension polymerase chain reaction (SOE-PCR; HO *et al.* 1989). All

DNA fragments generated by SOE-PCR were first cloned into the pGEM-T easy vector (Promega) and

their fidelity was verified by Sanger sequencing (Macrogen, Seoul, South-Korea). Plasmids were

introduced into *P. fluorescens* by tri-parental conjugations with the helper plasmid pRK2013,

carrying the *tra* and *mob* genes required for conjugation. Tn7-based plasmids were mobilized into

recipient strains with the additional helper plasmid pUX-BF13 (BAO *et al.* 1991).

To generate deletion mutants (*rrn* operons, *gacA, rsmA, rsmE* and *pflu3655*), regions flanking the

genes or operons of interest were amplified from SBW25 genomic DNA and assembled by SOE-PCR.

Deletion cassettes were inserted into the pUIC3 plasmid (RAINEY 1999) as SpeI fragments and

mutants were obtained following the two-step allelic exchange protocol described previously (ZHANG

AND RAINEY 2007). Deletion mutants were checked by PCR. To check *rrn* copy number after *rrn*

deletions, quantitative PCR were performed using a protocol described previously (FARR *et al.* 2017).

For complementation and over-expression studies, *PFLU3655* was amplified with oPR206/oPR207

and cloned into pME6032 (HEEB *et al.* 2002) as *Eco*RI/*Xho*I restriction fragment, downstream of the

P*tac* promoter.

To generate the Tn7-P*rrnB*-GFP reporter, a ~600bp fragment upstream of the *rrnB* operon (*PFLUr7-*

11) was amplified from SBW25 genomic DNA (oPR148/oPR151) and fused by SOE-PCR to *gfp-mut3.1*

sequence containing the T0 terminator previously amplified with oPR152/FluomarkerP2 from the

miniTn7(Gm)-P*rrnB1-gfp-a* plasmid (Lambertsen *et al.* 2003). The resulting fragment was cloned into

pUC18R6K-miniTn7T-Gm (CHOI *et al.* 2005) as a *Spe*I restriction fragment.

Site-directed mutagenesis of putative RsmA/E binding sites in pGEMTeasy-P*pflu3655*-GFP plasmid

was performed using the Quick Change mutagenesis kit (Stratagene) according to manufacturer's

instructions. Mutagenized fragments were then cloned into pUC18R6K-miniTn7T-Gm (CHOI *et al.*

2005) as a *Spe*I restriction fragment. In order to re-engineer point mutations in RsmA/E into

Pseudomonas fluorescens genome, a 1.5kb fragment spanning equal length on each site of the

target sites was amplified with oPR37/oPR223 and cloned into pGEM-T easy vector. Site-directed

mutagenesis was performed on this plasmid as described above, and the resulting DNA fragments

were cloned in pUIC3 as *Spe*I restriction fragments and introduced into *P. fluorescens* genome by

double recombination.

RNA extractions and RT-qPCR

For quantification of RNA concentration in bacterial cultures, cells were harvested from 1ml of

cultures at OD 0.5-1 and resuspended in 200ul of RNAlater solution (Invitrogen). For total RNA

quantification, we followed the method described by You *et al.* (2013), except that, before

processing, cells cultures were resuspended in RNAlater (Invitrogen) instead of being fast frozen on

dry ice. To normalise total RNA concentrations, the relationship between cell density and OD_{600nm}

was established for each strain by counting cells with a hemocytometer in 5 independent cultures of

similar OD to those used for RNA extractions.

Reverse-transcription quantitative PCR were performed as described previously (FARR *et al.* 2017),

using *gyrA* as an internal control. Oligonucleotide primers used for RT-qPCR are listed in

Supplementary Table 3.

Capsulation and gene expression assays

For capsulation test, cells were grown from standardized glycerol aliquots stored at -80°C. Aliquots

were diluted in KB and pre-cultures were grown overnight in order to reach an OD ~0.3-0.5 in the

morning. For gene expression studies, overnight pre-cultures were grown to saturation. In both

cases, pre-cultures were diluted to OD_{600nm}=0.05 in KB and incubated at 28°C. IPTG was added to a

final concentration of 0.1-1mM when indicated. Samples were taken at different time points for flow

cytometry and OD measurements. GFP fluorescence in bacterial populations was measured with a

BD FACS Canto II flow cytometer. Cell suspensions were diluted to a density of ~ 10^5 cells ml^{-1} in

filter-sterilised PBS and at least 20,000 cells were analysed by flow cytometry. Cellular debris were

filtered using side- and forward-scatter channels. GFP fluorescence was detected with a 488 nm

laser with 530/30 bandwidth filter. Laser intensity was set to 600V, except for P*rrnB*-GFP analyses

where intensity was lowered to 300V. Flow cytometry data files were analysed in R using the

flowCore package (ELLIS *et al.* 2017).

Growth curves in microplate reader

Overnight precultures were adjusted to OD_{600nm}=0.05 in fresh KB and used to start growth curves in a

Synergy 2 microplate reader (Biotek) in 96-well plates. Cultures were incubated for at least 24h at

28C with constant shaking and OD_{600nm} was read every 5 minutes.

To measure growth rates of cultures enriched in capsulated or non-capsulated cells, cells were

harvested from late-exponential phase (OD_{600nm}~1; 1B4) or 7-day old colonies (SBW25) and

centrifuged (1 min, 3000 rpm). Supernatants and pellets were collected, representing sub-

populations enriched in capsulated and non-capsulated cells, respectively. Cell suspension density

was adjusted to OD_{600nm}=0.05 in KB to start growth curves. Initial growth rates were calculated by

performing a linear regression on the logarithm of the measure OD values during the first 2h of

growth. Slopes of these linear regressions are reported as growth rate values.

Microscopy

Indian ink staining of capsulated cells was performed as described previously (GALLIE *et al.* 2015). For

time-lapse microscopy, bacteria were harvested from late exponential phase (1B4, OD~1-2) or from

7 day-old colonies (SBW25), diluted 1:1000 in KB and 2μl of the resulting suspension was

immediately transferred on a gel pad (1% agar KB) located on a glass slide within an adhesive frame

(GeneFrame, Thermo-Fisher). When dry, a cross-section of the pad was removed with a razor blade

in order to allow gaz exchanges to occur, the preparation was sealed with a glass cover-slip and

transferred to the microscope incubation chamber pre-heated at 28°C. Time lapse microscopy

experiments was performed with an Olympus BX61 upright microscope equipped with an F-View II

monochrome camera, a motorized stage and a temperature-controlled chamber set at 28°C. Devices

were operated by the Olympus Cell^P or CellSens softwares. Phase-contrast images were acquired

every 10 minutes with an oil-immersion 100x objective. One GFP image was taken before starting

the experiment in order to determine the capsulation status of each cell.

Phase contrast images were segmented with Fiji (SCHINDELIN *et al.* 2012) and individual colony areas

were extracted. For each colony analysed, the capsulation status of founding cell was scored

manually. The logarithm of the growth rate of individual 1B4 colonies was then fitted using a linear

regression. For SBW25, segmented linear regressions were found to better fit the data and the slope

of the first line was reported.

RNAseq analyses

RNAseq data were published previously (GALLIE *et al.* 2015). KEGG orthology terms for SBW25

genome were downloaded from the KEGG Orthology database (www.genome.jp; accessed in April

2016). KEGG enrichment statistics were computed with a hypergeometric test and was performed

separately for up- and down-regulated genes. Data from the transcriptome analysis of gacS mutant

were retrieved from supplementary table 3 in Cheng *et al.* (2013).

Competition experiments

SBW25 or SBW25-*lacZ* cell suspensions were spot-inoculated on separate KB agar plates and grown

for 7 days. Cells from the central ring were harvested, resuspended in Ringer's solution and gently

centrifuged to enrich suspensions in capsulated or non-capsulated cells. Capsulated SBW25 cells

were mixed with non-capsulated SBW-*lacZ* cells, and vice-versa. Mixed suspensions were diluted

100x in fresh KB and grown at 28°C in shaken tubes for 4h. Appropriate dilutions of the cultures

were plated on KB + X-gal plates at 0, 2h and 4h post-inoculation in order to measure the ratio of

white-blue colonies.

Acknowledgements:

The authors thank H. Hendrickson and P. Lind for help with microscopy and flow cytometry,

respectively, XX. Zhang for the generous gift of pUIC3-Δ*gacA* deletion plasmid, and E. Denisenko for

assistance with KEGG enrichment analyses. This work was supported in part by the Marsden Fund

Council and a James Cook Research Fellowship from government funding administered by the Royal

Society of New Zealand.

 Supplementary text: description of the mathematical model

The protein PFLU3655 is a key regulator for capsulation; we assume that the amount of colanic acid

(and fluorescence from the P*pflu3655-GFP* reporter) produced by the cell is proportional to

PFLU3655 concentration, described by the variable *x*. The rate of change of *x* will be the result of

production, at rate P(x), and of degradation and dilution, whose rates can be for simplicity assumed

proportional to x. The time change of x will thus be described by the following ordinary differential

equation:

$$\frac{dx}{dt} = P(x) - dx$$

where the constant d measures the rate of degradation/dilution (or decay). The equilibrium

solutions of the system x_E are such that the production equalizes the decay, that is when P(x_E)= d x_E.

We can constrain the shape of the production P(x) term based on the experimental results.

PFLU3655 generates a positive feedback loop, meaning that P(x) is a positive function and the

production of the corresponding mRNA will be increasing with x (the more PFLU3655 protein, the

higher the production of mRNA). If there was no post-trancriptional regulation, the amount of mRNA

would determine the amount of protein produced, and P(x) would follow a Michealis-Menten

kinetics whose saturation value would be set by the concentration R of ribosomes (the higher the

concentration of ribosomes, the higher the maximal amount of protein that can be produced). In this

case, the system would be monostable and cells would be either on or off (x_E=0 or x_E=x^{on}), depending

on whether the baseline production is smaller or larger than the decay, respectively.

When protein production is regulated post-transcriptionally, however, the competition of RsmA/E

and ribosomes (whose concentrations are assumed to be constant) for binding to the mRNA can

produce an ultrasensitive dependence of P(x) on x (Fig. 5a; black line). For small x, the produced

PFLU3655 mRNA will be almost entirely sequestered by RsmA/E and translation will be prevented.

Above a threshold concentration x_T, the titration by RsmA/E will be no longer sufficient to prevent

PFLU3655 proteins being produced, so that the production will be positive (but maybe insufficient to

overcome the decay term). If the affinity of RsmA/E is much stronger than that of ribosomes, the

shape of P(x) will be very nonlinear, with the production term being effectively zero for x<x_T. Finally,

for high x we expect production to saturate because of the finite amount of available ribosomes.

It is well known that ultrasensitivity can produce bistability (MUKHERJI *et al.* 2011; ZHANG *et al.* 2013;

FERRELL AND HA 2014). This is graphically illustrated in Fig. 5a: when the decay term (the slope of the

dashed red line) is sufficiently small, there will be three equilibrium solutions. Two of these solutions

are stable, corresponding to the OFF (x_E=0) and the ON (x_E=x^{on}) states, and one unstable (x_E=x^S). The

unstable solution separates the space of concentrations x in two domains: for all $x<x^S$, the

concentration will decrease and eventually approach the OFF state, for larger concentrations it will

converge to the ON state. The position of x^S, thus the extension of the basins of attraction of the two

stable equilibria, determines what is the probability that random fluctuations in the protein

concentration or in its production will lead the system to change equilibrium. For instance, the

probability to switch ON will be higher the smaller the value of x^S.

To examine what qualitative changes in the state of the system are expected following a variation in

its parameters, let us first imagine that for the WT strain in exponential phase of growth the decay

line lays just above the production curve (Fig. 5a; solid red line): cells will be almost exclusively in the

OFF state, even though rare large fluctuations in the protein concentration may lead them to be

'trapped' for some time in the region around the ON state. In the *carB* mutants, an increase in

ribosome concentration leads to an increase in the maximal production rate such that the two

curves cross in three points (Fig. 5b). There are now two alternative stable states, and the population

will contain a sizeable fraction of capsulated cells.

Changes in the parameters of the system can lead both to the transition from mono- to bistability

and to a change in the extent of the basins of attraction of the two stable equilibria, thus of the

population composition. When the baseline level of protein production is increased, as for instance

when the gene *PFLU3655* is inserted on a plasmid (Fig. 3b), the curve P(x) is displaced towards

higher values (Fig. 5c). This causes both the threshold concentration (thus the proportion of ON

cells) and the protein concentration (thus cell fluorescence) to increase.

References:

Ackermann, M., 2015 A functional perspective on phenotypic heterogeneity in microorganisms. Nat Rev Microbiol 13**:** 497-508.

Bao, Y., D. P. Lies, H. Fu and G. P. Roberts, 1991 An improved Tn7-based system for the single-copy insertion of cloned genes into chromosomes of gram-negative bacteria. Gene 109**:** 167-168.

Beaumont, H. J. E., J. Gallie, C. Kost, G. C. Ferguson and P. B. Rainey, 2009 Experimental evolution of bet hedging. Nature 462**:** 90-U97.

Bodi, Z., Z. Farkas, D. Nevozhay, D. Kalapis, V. Lazar *et al.*, 2017 Phenotypic heterogeneity promotes adaptive evolution. PLoS Biol 15**:** e2000644.

Bollenbach, T., S. Quan, R. Chait and R. Kishony, 2009 Nonoptimal microbial response to antibiotics underlies suppressive drug interactions. Cell 139**:** 707-718.

Buchler, N. E., and F. R. Cross, 2009 Protein sequestration generates a flexible ultrasensitive response in a genetic network. Mol Syst Biol 5**:** 272.

Buchler, N. E., and M. Louis, 2008 Molecular titration and ultrasensitivity in regulatory networks. J Mol Biol 384**:** 1106-1119.

Cheng, X., I. de Bruijn, M. van der Voort, J. E. Loper and J. M. Raaijmakers, 2013 The Gac regulon of Pseudomonas fluorescens SBW25. Environ Microbiol Rep 5**:** 608-619.

Choi, K. H., J. B. Gaynor, K. G. White, C. Lopez, C. M. Bosio *et al.*, 2005 A Tn7-based broad-range bacterial cloning and expression system. Nat Methods 2**:** 443-448.

Condon, C., S. French, C. Squires and C. L. Squires, 1993 Depletion of functional ribosomal RNA operons in Escherichia coli causes increased expression of the remaining intact copies. EMBO J 12**:** 4305-4315.

Dennis, P. P., M. Ehrenberg and H. Bremer, 2004 Control of rRNA synthesis in Escherichia coli: a systems biology approach. Microbiol Mol Biol Rev 68**:** 639-668.

Ehrenberg, M., H. Bremer and P. P. Dennis, 2013 Medium-dependent control of the bacterial growth rate. Biochimie 95**:** 643-658.

Ellis, B., P. Haaland, F. Hahne, N. Le Meur, N. Gopalakrishnan *et al.*, 2017 flowCore: Basic structures for flow cytometry data, R package version 1.44.0.

Farr, A. D., P. Remigi and P. B. Rainey, 2017 Adaptive evolution by spontaneous domain fusion and protein relocalization. Nat Ecol Evol 1**:** 1562-1568.

Ferrell, J. E., Jr., and S. H. Ha, 2014 Ultrasensitivity part II: multisite phosphorylation, stoichiometric inhibitors, and positive feedback. Trends Biochem Sci 39**:** 556-569.

Fridman, O., A. Goldberg, I. Ronin, N. Shoresh and N. Q. Balaban, 2014 Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. Nature 513**:** 418-421.

Gallie, J., E. Libby, F. Bertels, P. Remigi, C. B. Jendresen *et al.*, 2015 Bistability in a metabolic network underpins the de novo evolution of colony switching in Pseudomonas fluorescens. PLoS Biol 13**:** e1002109.

Gyorfy, Z., G. Draskovits, V. Vernyik, F. F. Blattner, T. Gaal *et al.*, 2015 Engineered ribosomal RNA operon copy-number variants of E. coli reveal the evolutionary trade-offs shaping rRNA operon number. Nucleic Acids Res 43**:** 1783-1794.

Heeb, S., C. Blumer and D. Haas, 2002 Regulatory RNA as mediator in GacA/RsmA-dependent global control of exoproduct formation in Pseudomonas fluorescens CHA0. J Bacteriol 184**:** 1046-1056.

Hindre, T., C. Knibbe, G. Beslon and D. Schneider, 2012 New insights into bacterial adaptation through in vivo and in silico experimental evolution. Nat Rev Microbiol 10**:** 352-365.

Ho, S. N., H. D. Hunt, R. M. Horton, J. K. Pullen and L. R. Pease, 1989 Site-directed mutagenesis by overlap extension using the polymerase chain reaction. Gene 77**:** 51-59.

Keener, J., and M. Nomura, 1996 Regulation of ribosome synthesis, pp. 1417-1431 in *Escherichia coli and Salmonella: cellular and molecular biology*, edited by F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low *et al.* American Society for Microbiology, Washington, D.C.

Kim, W., S. B. Levy and K. R. Foster, 2016 Rapid radiation in bacteria leads to a division of labour. Nat
Commun 7: 10508.
Kim, W., F. Racimo, J. Schluter, S. B. Levy and K. R. Foster, 2014 Importance of positioning for
microbial evolution. Proc Natl Acad Sci U S A 111: E1639-1647.
King, E. O., M. K. Ward and D. E. Raney, 1954 Two simple media for the demonstration of pyocyanin
and fluorescin. J Lab Clin Med 44: 301-307.
Koch, A. L., 1971 The adaptive responses of Escherichia coli to a feast and famine existence. Adv
Microb Physiol 6: 147-217.
Lapouge, K., M. Schubert, F. H. Allain and D. Haas, 2008 Gac/Rsm signal transduction pathway of
gamma-proteobacteria: from RNA recognition to regulation of social behaviour. Mol
Microbiol 67: 241-253.
Mori, M., S. Schink, D. W. Erickson, U. Gerland and T. Hwa, 2017 Quantifying the benefit of a
proteome reserve in fluctuating environments. Nat Commun 8: 1225.
Mukherji, S., M. S. Ebert, G. X. Zheng, J. S. Tsang, P. A. Sharp et al., 2011 MicroRNAs can generate
thresholds in target gene expression. Nat Genet 43: 854-859.
New, A. M., B. Cerulus, S. K. Govers, G. Perez-Samper, B. Zhu et al., 2014 Different levels of
catabolite repression optimize growth in stable and variable environments. PLoS Biol 12:
e1001764.
Norman, T. M., N. D. Lord, J. Paulsson and R. Losick, 2015 Stochastic Switching of Cell Fate in
Microbes. Annu Rev Microbiol 69: 381-403.
Paul, B. J., W. Ross, T. Gaal and R. L. Gourse, 2004 rRNA transcription in Escherichia coli. Annu Rev
Genet 38: 749-770.
Rainey, P. B., 1999 Adaptation of Pseudomonas fluorescens to the plant rhizosphere. Environ
Microbiol 1: 243-257.
Rainey, P. B., P. Remigi, A. D. Farr and P. A. Lind, 2017 Darwin was right: where now for experimental
evolution? Curr Opin Genet Dev 47: 102-109.
Rendueles, O., M. Garcia-Garcera, B. Neron, M. Touchon and E. P. C. Rocha, 2017 Abundance and co-
occurrence of extracellular capsules increase environmental breadth: Implications for the
emergence of pathogens. PLoS Pathog 13: e1006525.
Richard, M., and G. Yvert, 2014 How does evolution tune biological noise? Front Genet 5: 374.
Roberts, I. S., 1996 The biochemistry and genetics of capsular polysaccharide production in bacteria.
Annu Rev Microbiol 50: 285-315.
Rotem, E., A. Loinger, I. Ronin, I. Levin-Reisman, C. Gabay et al., 2010 Regulation of phenotypic
variability by a threshold-based mechanism underlies bacterial persistence. Proc Natl Acad
Sci U S A 107: 12541-12546.
Sambrook, J., E. F. Fritsch and T. Maniatis, 1989 Molecular cloning: a laboratory manual. Cold Spring
Harbor Laboratory, Cold Spring Harbor, N.Y.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair et al., 2012 Fiji: an open-source
platform for biological-image analysis. Nat Methods 9: 676-682.
Stearns, S. C., 1992 The evolution of life histories. Oxford University Press, Oxford ; New York.
Vadia, S., and P. A. Levin, 2015 Growth rate and cell size: a re-examination of the growth law. Curr
Opin Microbiol 24: 96-103.
Vakulskas, C. A., A. H. Potts, P. Babitzke, B. M. Ahmer and T. Romeo, 2015 Regulation of bacterial
virulence by Csr (Rsm) systems. Microbiol Mol Biol Rev 79: 193-224.
Valentini, M., D. Gonzalez, D. A. Mavridou and A. Filloux, 2017 Lifestyle transitions and adaptive
pathogenesis of Pseudomonas aeruginosa. Curr Opin Microbiol 41: 15-20.
van Boxtel, C., J. H. van Heerden, N. Nordholt, P. Schmidt and F. J. Bruggeman, 2017 Taking chances
and making mistakes: non-genetic phenotypic heterogeneity and its consequences for
surviving in dynamic environments. J R Soc Interface 14.

652 Van den Bergh, B., J. E. Michiels, T. Wenseleers, E. M. Windels, P. V. Boer *et al.*, 2016 Frequency of
653    antibiotic application drives rapid evolutionary adaptation of Escherichia coli persistence.
654    Nat Microbiol 1**:** 16020.
655 Veening, J. W., W. K. Smits and O. P. Kuipers, 2008 Bistability, epigenetics, and bet-hedging in
656    bacteria. Annu Rev Microbiol 62**:** 193-210.
657 Zhang, Q., S. Bhattacharya and M. E. Andersen, 2013 Ultrasensitive response motifs: basic amplifiers
658    in molecular signalling networks. Open Biol 3**:** 130031.
659 Zhang, X. X., and P. B. Rainey, 2007 Construction and validation of a neutrally-marked strain of
660    Pseudomonas fluorescens SBW25. J Microbiol Methods 71**:** 78-81.

Figure 1: Increased ribosome production in *carB* mutants
a. Transcriptional induction of ribosomal protein genes in SBW25, 1A4, 1B4 Cap- and 1B4 Cap+ cells. Absolute expression levels of ribosomal protein genes (KEGG pathway '0310-Ribosomes', n=26) were extracted from a previous RNA-seq dataset (Gallie *et al.* 2015) and normalised to SBW25. Letter groups indicate statistical significance; $P < 0.05$, Kruskall-Wallis test with Dunn's post-hoc correction.
b. Expression kinetics of the P*rrnB*-GFP transcriptional reporter. Fluorescence in individual cells was measured by flow cytometry. Mean fluorescence values of bacterial populations are reported. Means ± standard deviation are shown, n=4. Data are representative of 3 independent experiments.
c. Total RNA concentration in bacterial cells during exponential phase (OD=0.5-0.6) normalized per cell count. Values were normalized to SBW25 control within each experiment. Means ± standard deviation are shown, n=6. ** $P<0.01$, Student's t-test.

Figure 2: Genetic bases of capsulation

a. Capsulation in *rrn* deletion mutants. The Tn7-P*pflu3655*-GFP reporter was introduced in 1B4 bacteria ("WT") and its derived *rrn* mutants. Capsulation was measured by quantifying the proportion of GFP positive cells by flow cytometry at the onset of stationary phase (OD~2). Means ± standard error of the mean are shown, n≥8. Data were pooled from 4 independent experiments.* $P < 0.05$, *** $P < 0.001$, Dunn's test with Benjamini-Hochberg post-hoc correction, comparison to WT.

b. Capsulation in *gac/rsm* mutants. Means ± standard error of the mean are shown, n=9. Data are pooled from 3 independent experiments. *** $P < 0.001$, Dunnett's multiple comparison test on log-transformed data.

c. A model for capsulation in 1B4. See text for details.

Figure 3: *PFLU3655* expression is necessary and sufficient for capsulation
a. Capsulation in SBW25, 1B4Δ*pflu3655* and 1B4 strains carrying the pME6032-*PFLU3655* plasmid or empty vector (EV) after induction with 1mM IPTG. Phase contrast microscopy images of bacterial suspensions counter-stained with indian ink. White halo around cells indicate capsulation. Scale bar = 10 μm.
b. PFLU3655 establishes a positive feedback loop. GFP fluorescence from the P*pflu3655*-GFP reporter in SBW25 (left), 1B4Δ*pflu3655* (middle) or 1B4 (right) cells carrying the pME6032-*PFLU3655* plasmid or empty vector. *PFLU3655* expression was induced with IPTG at indicated concentration and fluorescence was measured by flow cytometry. Smoothed fluorescence distributions are shown. Data are representative of 3 independent experiments.

Figure 4: *PFLU3655* is a direct target of RsmA/E

a. Schematic diagram of *PFLU3655* region. Two putative RsmA/E binding sites (orange squares) are located in the promoter and 5' region of the gene. Numbers indicate nucleotide position relative to start codon (not to scale). Sequences of putative RsmA/E binding sites are shown, the putative ribosome-binding site (RBS) is underlined. Below: point mutations introduced in the different sequences by site-directed mutagenesis.

b. Expression of the P*pflu3655*-GFP reporter carrying the different point mutations in the 1B4 background. GFP fluorescence was measured by flow cytometry, smoothed distributions are shown. Data are representative of 3 independent experiments.

c. Mutations in putative RsmA/E binding sites affect capsulation. Individual point mutations were re-introduced into SBW25 and 1B4 carrying the wild-type P*pflu3655*-GFP reporter and the proportion of GFP positive cells in late exponential phase (OD~1-2) was measured by flow-cytometry.

Figure 5: A mathematical model of capsulation allows to qualitatively explain switching behaviour

Graphical analysis of the effect of parameters on switching behaviour. Growth rate (a), ribosome concentration (b) and basal transcription rate of *PFLU3655* (c) can all affect the position of the equilibrium points.

Production curves P(x) are represented in black for wild-type SBW25 (a,b), in blue for *carB* mutants (b,c), and light blue for *carB* mutants with increased basal transcription of *PFLU3655* (c). Decay curves are drawn in purple. Stable or unstable equilibria are located at the intersections between production and decay curves.

Figure 6: Capsulation and growth in 1B4

a. Initial growth rate after nutrient upshift depends on the proportion of capsulated cells in 1B4 populations. Data points pooled from 2 independent experiments. n=36, r^2=0.91. Shaded area indicates 95% confidence interval.

b. Growth rate of micro-colonies founded by cap- (GFP-) or cap+ (GFP+) cells measured by time-lapse microscopy. Growth rate was measured as the coefficient of the linear fit of the logarithm of colony area over time during the first 2h of growth. n=97 for GFP-; n=94 for GFP+. *** $P < 0.001$, Wilcoxon test.

Figure 7: Capsulation and growth in SBW25

a-b. SBW25 colony grown on KB agar plate (a) or KB agar plate supplemented with 2mM uracil (b) for 7 days. Scale bar = 2 mm.

c. SBW25 cells carrying the Tn7-P*pflu3655*-GFP reporter and sampled from a 7-days old colony were counter-strained with indian ink to detect the presence of colanic-acid capsules. GFP images are overlaid to the phase contrast image. Scale bar = 10 μm.

d. Competitive fitness difference between SBW25 Cap- and Cap+ cells. Boxplot of he differences in Malthusian parameters between cultures enriched in Cap+ vs. Cap- cells are shown, n=12. * $P<0.05$, *** $P<0.001$, comparison to 0 with Student's t-test.

e. Growth rate of microcolonies founded by GFP- or GFP+ cells measured by time-lapse microscopy. Growth rate was measured as the first coefficient from the segmented fit of the logarithm of colony area over time during the first 2h of growth. n=65. *** $P < 0.001$, Wilcoxon test.

Supplementary Figure 1: Capsulation in the *galU* mutant
The Tn7-P*pflu3655*-GFP reporter was introduced in 1B4 and *galU* mutant.
Capsulation was measured by quantifying the proportion of GFP positive cells
by flow cytometry at the onset of stationary phase. Means ± standard
deviation are shown. * $P < 0.05$, Student's t-test, n = 12.

Supplementary Figure 2: Growth curves of *carB* mutants

Growth kinetics of SBW25, SBW25 *carB*, 1A4 and 1B4 strains in KB medium. Overnight cultures were diluted and used to inoculate 200μl KB cultures in a 96-well plate. OD_{600nm} was read in a microplate reader every 5min during incubation at 28°C with continuous shaking. Lines represent mean from 3 technical replicates.

<u>Supplementary figure 3: Capsulation kinetics in 1B4</u>
The Tn7-P*pflu3655*-GFP reporter was introduced in 1B4. OD_{600nm} (inset) and
the size of GFP positive subpopulation (main panel) were monitored over
>15h. Capsulation was measured by quantifying the proportion of GFP
positive cells by flow cytometry. Means ± standard deviation are shown, n=3.
Data are representative of 2 independent experiments.

Supplementary Figure 4: Capsulation frequency in double *rrn* mutants
The Tn7-Pp*flu3655*-GFP reporter was introduced in 1B1 bacteria ("WT") and
its derived *rrn* mutants. Capsulation was measured by quantifying the
proportion of GFP positive cells by flow cytometry at the onset of stationary
phase (OD~2). Color indicate the number of *rrn* operons per genotype. Means
and standard error of the mean are shown, n≥9. Data are pooled from 3
independent experiments. * *P* < 0.05 compared to WT, Dunn's test with
Benjamini-Hochberg post-hoc correction.

<u>Supplementary Figure 5: RNA quantification in *rrn* mutants</u>
Total RNA concentration in bacterial cells during exponential phase
(OD=0.5-0.6) normalized per cell count. Values were normalized to SBW25
control within each experiment. Means ± standard deviation are shown, n=6.
* *P*<0.05, Student's t-test. n.s. = not signifcant.

Supplementary figure 6: Expression of *gacS*-regulated genes
Genes that are up-regulated (left) or down-regulated (right) more than 4
times in a *gacS* mutant compared to WT SBW25 were recovered from Cheng
et al. 2013 (dataset from supplementary table 3). The distribution of induction
or repression values (after normalization by the base expression in SBW25) in
the different RNAseq datasets is shown for each set of genes. n=125 (a) or 165
(b). Letter groups indicate statistical significance at $P < 0.05$, Kruskall-Wallis
test with Dunn's post-hoc correction.

<u>Supplementary Figure 7: Growth rate of SBW25 cultures enriched in Cap- or Cap+ cells</u>

Cells from 7-day old colonies were resuspended in KB medium and suspensions were enriched in Cap- (pellet) or Cap+ (supernatant) cells by gentle centrifugation. The growth of these suspensions in a 96-well plate was measured with a microplate reader and the slopes from the fit of logarithm of density increase over time during the first 2h of growth are shown. n=8. ** $P <$ 0.01, Wilcoxon test.