RESEARCH ARTICLE

Adult plant resistance in maize to northern leaf spot is a feature of partial loss-of-

function alleles of *Hm1*

Sandeep R. Marla[1,#a], Kevin Chu[1,#b], Satya Chintamanani[1,#c], Dilbag Multani[2], Antje

Klempien[1,#d], Alyssa DeLeon[1,#e], Kim Bong-suk[1], Larry D. Dunkle[1], Brian P. Dilkes[3],

Gurmukh S. Johal[1*]

[1]Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN
47907
[2]DowDuPont, Johnston, IA 50131
[3]Department of Biochemistry, Purdue University, West Lafayette, IN 47907
12
[#a]Current address: Department of Agronomy, Kansas State University, Manhattan, KS
66506
[#b]Current address: Donald Danforth Plant Science Center, St. Louis, MO 63132
[#c]Current address: Syngenta, Slater, IA 50244
[#d]Current address: KeyGene USA, Rockville, MD 20850
[#e]Current address: DowDuPont, Johnston, IA 50131
19
20   [*]**Corresponding author**

21   E-mail: gjohal@purdue.edu

23   **Short title:** A causal link between weak resistance and APR in maize

24   **One sentence summary:** Characterization of adult plant resistance in the maize-CCR1

25   pathosystem reveals a causal link between weak resistance and APR.

ABSTRACT

Adult plant resistance (APR) is an enigmatic phenomenon in which resistance genes are ineffective in protecting seedlings from disease but confer robust resistance at maturity. Maize has multiple cases in which genes confer APR to northern leaf spot, a lethal disease caused by *Cochliobolus carbonum* race 1 (CCR1). The first identified case of APR in maize is encoded by a hypomorphic allele, $Hm1^A$, at the *hm1* locus. In contrast, wild type alleles of *hm1* provide complete protection at all developmental stages and in every part of the maize plant. *Hm1* encodes an NADPH-dependent reductase, which inactivates HC-toxin, a key virulence effector of CCR1. Cloning and characterization of $Hm1^A$ ruled out differential transcription or translation for its APR phenotype and identified an amino acid substitution that reduced HC-toxin reductase (HCTR) activity. The possibility of a causal relationship between the weak nature of $Hm1^A$ and its APR phenotype was confirmed by the generation of two new APR alleles of *Hm1* by mutagenesis. The HCTRs encoded by these new APR alleles had undergone relatively conservative missense changes that partially reduced their enzymatic activity similar to $HM1^A$. No difference in accumulation of HCTR was observed between adult and juvenile plants, suggesting that the susceptibility of seedlings derives from a greater need for HCTR activity, not reduced accumulation of the gene product. Conditions and treatments that altered the photosynthetic output of the host had a dramatic effect on resistance imparted by the APR alleles, demonstrating a link between the energetic or metabolic status of the host and disease resistance affected by HC-toxin catabolism by the APR alleles of HCTR.

AUTHOR SUMMARY

Adult plant resistance (APR) is a phenomenon in which disease resistance genes are able to confer resistance at the adult stages of the plant but somehow fail to do so at the seedling stages. Despite the widespread occurrence of APR in various plant diseases, the mechanism underlying this trait remains obscure. It is not due to the differential transcription of these genes, and here we show that it is also not due to the differential translation or activity of the APR alleles of the maize *hm1* gene at different stages of development. Using a combination of molecular genetics, biochemistry and physiology, we present multiple lines of evidence that demonstrate that APR is a feature or symptom of weak forms of resistance. While the mature parts of the plant are metabolically robust enough to manifest resistance, seedling tissues are not, leaving them vulnerable to disease. Growth conditions that compromise the photosynthetic output of the plant further deteriorate the ability of the seedlings to protect themselves from pathogens.

INTRODUCTION

Plant responses to pathogens are dynamic, and they involve a number of inducible mechanisms that are tightly regulated both in space and time (1). They are called into action only at the time and site of infection. The tight regulation of innate immunity is due to disease resistance (DR) genes that plants inherit from their parents and which often segregate with the trait of resistance (1–3). A vast majority of these DR genes function in every part of the plant and at every stage of development. However, many exceptions exist where resistance is manifested in a tissue- or developmental stage-specific manner. In most instances of developmentally regulated resistance, plants are susceptible at the seedling stage but become increasingly resistant toward maturity. The term commonly

used to define such developmentally regulated resistance is adult plant resistance (APR),

although other terms such as age-associated resistance, ontogenic resistance, mature plant

resistance, or flowering-induced resistance have also been used in the literature to

describe the same phenomenon (4–8).

Adult plant resistance (APR) often manifests gradually with the advancement of

plant age, but a few cases have been reported where the onset is abrupt, happening

sharply at a certain stage of development (9–12). An example of the latter kind is the

wheat *Lr34* gene-mediated resistance, in which the onset against the leaf rust pathogen,

Puccinia triticina, is largely confined to the uppermost leaf (flag leaf) (13). In contrast, in

the rice-*Xanthomonas oryzae* pv. *oryzae* pathosystem, resistance conferred by the *Xa21*

gene is almost negligible during the first three weeks of age but then increases steadily

each week, reaching full efficacy at maturity (14,15). Similarly, the *Yr36*-conferred

resistance in wheat to *Puccinia striiformis* (16) and the *Hm2*-conferred resistance in

maize to *Cochliobolus carbonum* race 1 (CCR1) increase gradually with plant age (12).

In efforts to understand the mechanistic basis of APR, several genes conferring

this form of resistance were isolated in different pathosystems. Some of these genes

include *Cf-9B* from tomato conferring resistance to leaf mold (17), *Mi-1* from tomato

conferring resistance to aphids (18), *Xa21* from rice conferring resistance to leaf blight

(14), *Lr67* and *Lr34* from wheat conferring resistance to leaf rust (13,19), *Yr36* from

wheat conferring resistance to stripe rust (16), and *Hm2* from maize conferring resistance

to leaf blight (12). Two of these genes, *Cf-9* and *Mi-1*, clearly follow the gene-for-gene

(GFG) paradigm in conferring resistance, while four others, *Lr67, Lr34, Yr36* and *Hm2,*

do not, suggesting that any disease resistance gene has the potential to confer an APR

phenotype.

What makes a gene behave in an APR manner? This question still eludes us, even

though a number of APR genes, including those described in the preceding paragraph,

have been cloned and characterized. One logical expectation was that the phenotype of

APR genes may derive from their differential expression at different stages of plant

development and that the level of gene expression would match their phenotypic efficacy

closely. However, this has been ruled out with the majority of the APR genes, as their

transcript levels do not reflect changes in their resistance phenotype (12,13,15,17,20).

Other possibilities that may affect the APR behavior of these genes are differential

translation, differential post-translational modifications, and developmental changes in

plant physiology and metabolism.

To gain insight into the mechanistic basis of APR in maize, we have been

studying the northern leaf spot (NLS) disease of maize (*Zea mays*) caused by *C.*

carbonum race 1. A classic APR syndrome is described in this pathosystem where alleles

at two homeologous loci can confer resistance in a developmentally programmed fashion

(9). These duplicate genes, *Hm1* and *Hm2*, encode NADPH-dependent HC-toxin

reductases (HCTR), which utilize NADPH as a cofactor to reduce an essential ketone

function in HC-toxin (HCT), the key disease causing effector of CCR1, and abolish its

activity (21–23). There is one prominent difference between the HCTRs encoded by *hm1*

and *hm2*: whereas the HCTR encoded by wild type (WT) *Hm1* contains 356 amino acids,

the HCTR encoded by the functional *Hm2* allele is truncated and lacks the last 52 amino

acids compared to HM1 (12). This truncated allele is the only functional allele that has

been identified at *hm2*, and it confers APR against CCR1 when *hm1* is null. *Hm2* is

expressed throughout the age of the plant (12), ruling out developmentally regulated

transcript accumulation as the mechanism of APR. Like *Hm2*, an allele of *hm1* conferring

APR has also been described (9). Designated $Hm1^A$, this APR allele is recessive to the

WT *Hm1* allele and dominant to the *hm1* null allele (9).

To explore why and how the $Hm1^A$ allele leads to an APR phenotype, we have

cloned and characterized it in detail. Comparison of the sequence of $Hm1^A$ with those of

the WT haplotypes from a number of resistant inbreds and accessions revealed a single

amino acid substitution in the $HM1^A$ peptide that is unique to its APR behavior. $HM1^A$

transcripts accumulated to similar levels throughout plant growth and development, as

did the translational product of the gene. However, the HCTR activity in $Hm1^A$ plants

was intermediate between WT (*Hm1Hm*1) and null mutant (*hm1hm1hm2hm2*) plants.

This, along with the truncated nature of the APR allele at *hm2*, prompted us to consider if

the hypomorphic *Hm1* allele in $Hm1^A$ was the reason for its APR phenotype. This

hypothesis was addressed by mutagenesis, generating two new APR mutants of the B73

maize inbred, which is homozygous for the WT allele at *hm1* and the null allele at *hm2*

(*Hm1Hm1hm2hm2*). Both new APR alleles were found to contain single amino acid

substitutions in HM1-B73 and reduced HCTR activity. Thus, APR is a symptom of

partial loss-of-function mutations in *Hm1* that result in seedling susceptibility.

RESULTS

Detailed genetics of APR-conferring $Hm1^A$ as an allele of *hm1*

The APR trait attributed to $Hm1^A$ was first noticed in the inbred P8, developed at Purdue

University in the early 1960s (9). The genetic evidence linking the APR of P8 with an

allele of *hm1* (*Hm1^A*) made use of two segregating populations, a testcross and an F$_2$

population, generated by crossing P8 (*Hm1^{A}Hm1^{A}hm2hm2*) with the resistant inbred

WF9 (*Hm1Hm1Hm2Hm2*). The susceptible inbred for the testcross was Pr, which is

homozygous for null mutations at both the *hm1* and *hm2* loci. There were at least two

concerns with this study. First, it used a relatively small number of progenies, comprising

about 90 plants each for both the F$_2$ and testcross populations. Second, the resistant

inbred WF9 also contained an APR allele at the *hm2* locus, leaving room for error in

extrapolation from these data.

These concerns necessitated that we revisit these findings, to clone and

characterize *Hm1^A*. We acquired P8 from the Germplasm Resources Information

Network (GRIN). To confirm that this source of P8 harbored the *Hm1^A* allele reported by

Nelson and Ullstrup (1964), we conducted a thorough analysis of the genetics of P8

resistance to CCR1. We first crossed P8 twice with Pr (*hm1hm1hm2hm2*) to produce a

BC$_1$F$_1$ testcross population. Of 384 BC$_1$F$_1$ plants inoculated with CCR1, 186 plants were

susceptible at both the seedling and adult stage while 198 plants were susceptible as

seedlings, but later emerging leaves were fully resistant, consistent with the APR

phenotype of P8. The recessive null *hm1* allele of Pr (designated as *hm1^{Pr}*) contains a

256-bp *Drone* transposon insertion in exon 4 (24). All 186 plants susceptible at maturity

were homozygous for *hm1^{Pr}*, whereas all 198 plants that were initially susceptible and

then displayed APR were heterozygous for *hm1^{Pr}*. This 1:1 ratio of susceptible *vs.* APR

plants (X^2 - 0.375, $P > 0.05$, 1 d.f.) indicated that a single gene at or near the *hm1* locus

controlled the APR behavior of P8.

Next we crossed P8 to Pr1, a near isogenic line (NIL) of Pr in which the mutant

hm1 allele was replaced by a WT *Hm1* (25). The resulting $Hm1^A Hm1$ F_1 hybrid was

testcrossed to Pr, the *hm1hm2* null stock. The inheritance of $Hm1^{Pr1}$ vs. $Hm1^A$ in this

population was tracked with a PCR-based marker that differentiated between those two

alleles. Of the 540 F_1 test cross progeny, 276 were susceptible as seedlings and later

exhibited APR, while the remaining 264 were completely resistant to CCR1 regardless of

age. All 264 completely resistant plants had inherited the WT *Hm1* allele from Pr1, while

the 274 plants that exhibited APR had inherited the $Hm1^A$ allele from P8. Chi-squared

tests supported the 1:1 expected inheritance of monogenic inheritance (X^2-0.266667, *P* >

0.05, 1 d.f.). No recombinants between the genotypes at the *hm1* locus and the expression

of CCR1 susceptibility were found in either population (924 opportunities for crossover).

This confirmed that the source of P8 we obtained recapitulated the phenomenon

described in 1964 (9) and that the APR of P8 is likely conferred by the $Hm1^A$ allele.

To incorporate $Hm1^A$ into a uniform background for detailed phenotypic

comparisons, we introgressed this APR allele into the B73 inbred by crossing P8

($Hm1^A Hm1^A$ *hm2hm2*) to B73 (*Hm1Hm1hm2hm2*). As $Hm1^A$ is recessive to WT *Hm1*, we

utilized sequence polymorphism between $Hm1^A$ and $Hm1^{B73}$ to construct a PCR-based

marker. After seven crosses to B73 with selection for the $Hm1^A$ genotype, BC_7F_2 progeny

were generated by self-pollinating a heterozygous plant. This BC_7F_2 population

segregated in a 3:1 ratio for complete resistance and APR, again consistent with $Hm1^A$

being responsible for APR of P8. Homozygous $Hm1^A$ plants from this population were

selected and maintained as an $Hm1^A$ near-isogenic line in B73.

Phenotypic manifestation of adult plant resistance in maize to CCR1

To develop a comprehensive account of the onset of APR by $Hm1^A$, we also introgressed the null $hm1^{Pr}$ allele into the B73 background over seven generations, and crossed with $Hm1^A$ B73 NIL to generate plants heterozygous for $Hm1^A$. Both homozygous ($Hm1^AHm1^A$) and heterozygous ($Hm1^Ahm1^{Pr}$) $Hm1^A$ plants were inoculated with CCR1 at weekly intervals, starting at 1 week-after-planting (wap) and culminating at 10 wap. Their infection phenotypes were measured using a 1-10 disease rating scale (12) and compared with those of B73 and a B73 NIL containing the null $hm1$ allele ($hm1^{Pr}$ B73 NIL). A rating of 10 on this scale indicated highly susceptible plants, while a rating of 1 indicated complete resistance.

The susceptible $hm1^{Pr}$ B73 NILs scored 10 on the disease rating scale regardless of age, and the resistant controls (B73 inbred), which produced small chlorotic flecks in response to CCR1 infection, scored 1 throughout development. Plants containing $Hm1^A$ exhibited very little resistance at the seedling stage, but severity scores decreased with age (Fig 1A and 1C). At the age of week-1, $Hm1^A$ seedlings were consistently rated 8 or higher. This disease rating dropped to 5 or less by week-5. At week-10, $Hm1^A$ plants resembled the resistant controls, receiving a rating of 1 (Fig 1B and 1C). The level of resistance conferred by $Hm1^A$ correlated with the age of the whole plant at the time of inoculation and not the age of the inoculated leaf. Inoculating each leaf of $Hm1^AHm1^A$ and $hm1hm1$ plants at week-5 of plant growth confirmed this observation. All the leaves of $Hm1^A$ plants were equally resistant regardless of their age, and all the leaves of $hm1hm1$ plants were equally susceptible (data not shown).

Similar to the APR conferred by the *Hm2* gene (12), the resistance conferred by

Hm1^A was dosage dependent. Plants homozygous for *Hm1^A* were slightly more resistant

to CCR1 at almost all stages of development compared to plants heterozygous for *Hm1^A*

and the null allele (*Hm1^A hm1^{Pr}*) indicating that *Hm1^A* is haploinsufficient (Fig 1C). The

dosage effect was more pronounced at week-5 and declined after week-7 as the plants

matured and became completely resistant.

Molecular characterization of the *Hm1^A* allele

Atypical behavior of a disease resistance gene can sometimes result from complex

structural changes at the locus, such as an increase in the copy number of the gene or a

part of the gene (26,27). To address if such a genetic mechanism also led to the *Hm1^A*

APR, we conducted a Southern blot analysis with P8 DNA digested with a variety of

restriction enzymes. Consistent with the genetic data, a single *BamHI* restriction fragment

hybridized to *Hm1*-specific probes on these blots, indicating that *Hm1^A* was a single copy

gene in the P8 inbred and that the entire gene was present on a 13 kb restriction fragment

(Fig 2A). To clone the *Hm1^A* gene, a lambda library was constructed from the *Bam*H1-

digested P8 DNA restriction fragments migrating on a gel as 12 to 15 kb fragments. We

identified and sequenced a clone containing the 13 kb *hm1*-encoding fragment. Sequence

analysis indicated that our clone contained the entire coding region of the *Hm1* gene, as

well as 3.8 kb of the promoter region.

To determine the structural changes in *Hm1^A,* its sequence was compared with

that of the B73 reference sequence. Significant changes were encountered in the promoter

regions of *Hm1^A* and *Hm1^{B73}*. Except for a few indels and SNPs, the first -200 bp from

the translation start site of the promoter region are similar in *Hm1^A* and B73 (Fig S1). The

next -1.5 kb region upstream, however, is completely different between the two alleles,

though this does not seem to be due to the insertion of a transposable element.

Interestingly, the promoter region of $Hm1^A$ is identical to that of $hm1^{Pr}$, the null $hm1$

allele from the susceptible inbred Pr. To examine if any other resistant lines containing a

wild type $Hm1$ allele also had a promoter region identical to that of $Hm1^A$, we used a

primer pair designed from the $Hm1^A$ promoter region to PCR amplify DNA from a

number of resistant inbreds. Two inbreds, Pr1 and Va35, were found whose $Hm1$ WT

alleles have the promoter regions identical to that of $Hm1^A$ (Fig S1). Taken together,

these results indicate that the promoter polymorphism between $Hm1^A$ and $Hm1^{B73}$

predicted neither resistance nor susceptibility and thus may be inconsequential to the

APR phenotype of $Hm1^A$.

The coding region of $Hm1^A$ also differed from that of $Hm1^{B73}$, containing nine

SNPs. Although four of these SNPs were silent or synonymous, five led to amino acid

substitutions in the predicted $HM1^A$ peptide (Fig 2B). Relative to the B73 HM1

reference, these substitutions were: a Serine to Tyrosine change at residue 99 (S99Y), an

Aspartic acid to Tyrosine change at residue 110 (D110Y), a Leucine to Histidine change

at residue 116 (L116H), a Serine to Asparagine change at residue 191 (S191N), and a

Leucine to Proline change at residue 240 (L240P) (Fig 2B).

The L116H substitution is the likely causative polymorphism in the $Hm1^A$ allele

As $Hm1$ is one of the most polymorphic genes in maize (28), we decided to examine the

peptide sequence of various resistance alleles to potentially pinpoint the amino acid

change(s) responsible for the APR behavior of $Hm1^A$. We first amplified and evaluated

the HM1 sequences of Pr1 and Va35, the two resistant inbreds that share their promoters

with $Hm1^A$, and compared them with the sequences of both HM1^A and HM1^{B73}. HM1^{Pr1}

was found to differ by five amino acids from HM1^{B73}, with two of these polymorphisms,

S99Y and L240P, also being present in HM1^A (Fig S2). These same two changes were

also found in HM1^{Va35}, which differed from HM1^{B73} by six amino acids. Another

resistant $Hm1$ allele that differed from B73 by six amino acids was in the inbred W22,

but none of those changes matched those of HM1^A. However, the predicted HM1 of the

landrace Enano from Bolivia (28) shared with HM1^A the two polymorphisms D110Y and

S191N. And most importantly, the HM1 of the landrace Pira from Colombia (28) shared

four of the five amino acid changes between HM1^A and HM1^{B73}. These are S99Y,

D110Y, S191N, and L240P, thereby leaving only the L116H polymorphism unique to

HM1^A.

To examine the functional status of the $Hm1$ allele of Pira, we acquired this

landrace from GRIN and inoculated it with CCR1. It was found to be completely resistant

to CCR1, even at the seedling stage. This demonstrated that despite having four of the

five amino acid changes of HM1^A, the $Hm1^{Pira}$ allele is fully functional and not APR.

These results highlight the importance of the L116H substitution in defining the

phenotype of $Hm1^A$. Consistent with this hypothesis, the Leucine at 116 is highly

conserved not only in all the homoeologs and orthologs of the $Hm1$ gene across the grass

lineage, but also in the maize dihydroflavonol 4-reductase (DFR), an NADPH-dependent

enzyme of the anthocyanin pathway predicted to be a progenitor of HM1 (Fig S2). All

these findings suggest that the HM1^A L116H substitution is unique to $Hm1^A$ and may

underlie its APR behavior to CCR1 in maize by somehow negatively impacting HCTR

activity.

HM1 transcript accumulation is not developmentally regulated in *Hm1^A*

To examine if the transcriptional activity of *Hm1^A* undergoes any change during plant

development, reverse transcription (RT)-PCR was conducted on RNA extracted from

CCR1-inoculated *Hm1^A* plants of diverse ages. Using a semi-quantitative form of this

assay, no dramatic changes could be observed in the level of the *Hm1^A* transcript between

the seedling and mature-plant stages (Fig 3A). Likewise, quantitative real time PCR

(qRT-PCR) measurements of transcript abundance of *Hm1^A* plants inoculated with CCR1

at different ages did not detect any rise in HM1 expression as the susceptible plants

became resistant over time (Fig 3B). These results ruled out the differential transcription

of the *Hm1^A* allele as the basis for its APR phenotype.

The level and activity of *Hm1^A*-encoded HCTR stays the same during plant

development

To address if the differential translational activity of *Hm1^A* had any impact on its APR

behavior, we first conducted western analysis to examine the level and stability of the

HM1^A protein. While an antibody raised against the entire HM1^A peptide lacked

specificity, a multiple antigenic peptide (MAP) (29) antibody generated against a 13 aa

peptide corresponding to residues 312 to 324 of the HM1 peptide worked well and

reacted to a single product on Western blots generated from *Hm1^A* homozygous or

heterozygous plants (Fig 3C). No change in the level of the HM1^A protein could be

detected over time, indicating that the APR phenotype of *Hm1^A* is not due to differential

translation either.

We next addressed if the activity of HCTR encoded by *Hm1^A* had any role in its

APR behavior. To do this, an LC-MS/MS-based *in vitro* HCTR activity assay that

quantified the reduction of HC-toxin by crude protein extracts was developed. The *in*

vitro measurements were normalized to total protein content, allowing us to estimate the

level of the functional HCTR in plant tissues. To examine the level of HCTR over time,

proteins were extracted from CCR1-inoculated leaves of 3- and 7-week-old plants of

Hm1^A and control stocks, and their HCTR activity was measured in replicated samples.

Two trends were noted as shown in Fig 3D. First, the HCTR activity encoded by HM1^A

was lower than by the WT allele but not null as that of *hm1^{Pr}*. At both stages of

development, the HCTR activity of HM1^A was about 3-fold lower than that of HM1.

Second, the level of active HCTR differed little if any in 3- or 7-week *Hm1^A* plants (Fig

3D). Likewise, the HCTR activity of the WT allele also did not differ between week-3

and week-7-old plants (Fig 3D). Two conclusions can be drawn from these results. First,

Hm1^A encodes an HCTR that is relatively weaker than the enzyme encoded by the WT

allele. Second, the level of the active HCTR stays constant over development and does

not account for the APR phenotype of *Hm1^A*.

Partial loss-of-function mutations confer adult plant resistance in the maize-CCR1

pathosystem

What aspect of the *Hm1^A* gene structure or function restricts it to be an APR gene, i.e.,

conferring resistance only at the mature-plant stage but not the seedling stage? Having

ruled out differential transcription or translation as possible mechanisms, we paid

attention to an attribute of *Hm1^A* that differentiates it from both the WT and null mutant

alleles of *hm1* - the relatively weak nature of the HCTR activity encoded by *Hm1^A*. This

partial enzymatic activity of HM1^A mirrored exactly the phenotypic strength of resistance

conferred by this APR allele, which is recessive to that of WT *Hm1* but dominant to that

of null *hm1*. Given that the APR allele at the *hm2* locus also confers partial resistance to

CCR1 (12), we pondered if this could be a requirement for a resistance gene to have an

APR phenotype.

If this hypothesis that a *Hm1* APR allele owes its phenotype to being a weak or

partial loss-of-function allele is correct, we should be able to confirm it by generating

new APR alleles from the WT *Hm1* allele by mutagenesis. To address this possibility, we

first tried a random mutagenesis screen to generate new alleles of *Hm1*, in large part

because of the lethal nature of CCR1 infection on field-grown plants lacking functional

Hm1. About 1,000 M_2 families of B73 were generated by treating pollen with the

mutagen ethyl methanesulfonate (EMS). Twenty-four plants per M_2 family were planted

in a field and inoculated with CCR1 at the seedling stage. One M_2 family was identified

in which CCR1-susceptible plants segregated in a recessive fashion. These plants

remained susceptible throughout their growth, suggesting they were the result of a null

mutation. Sequence analysis of the *hm1* allele from this mutant (named *hm1-2*) confirmed

its null status and revealed a single G to A transition at the junction of exon3/intron3 as

the cause of mutation (Fig S3). Since this change is expected to abolish the splicing of

intron 3, it would result in a truncated protein lacking all the amino acids encoded by

exons 4 and 5 (Fig S3). It is unlikely that such a grossly truncated protein would have any

HCTR activity, and as shown in Fig 6, *hm1-2* exhibited very little enzymatic activity.

We next conducted a targeted mutagenesis screen to generate a series of mutant

alleles of *Hm1*. To accomplish this, EMS-mutagenized *Hm1^B73* pollen was applied to ears

of completely susceptible Pr plants in a greenhouse (Fig 4). Approximately 4,500 M_1

seeds obtained from this cross were planted in the field and inoculated with CCR1 at

week-2. Seven plants were identified as CCR1 susceptible at this seedling stage. When

inoculated again at week-5, five of them were still fully susceptible, suggesting they were

null mutants. The other two plants however exhibited APR as they developed different

levels of resistance (Fig S4). Sequencing the *Hm1* gene (Fig S5) from all seven mutants

revealed that they all carried GC to AT transitions in the coding region of *Hm1*. The two

APR-exhibiting alleles (designated *Hm1-3* and *Hm1-4*) had missense mutations resulting

in single amino acid substitutions, T90M in *Hm1-3* and V210M in *Hm1-4*, in the HM1

peptide (Table 1). Of the five null mutants, three (named *hm1-6* to *hm1-8*) had nonsense

mutations, one a C82Y substitution (*hm1-5*), and one a splice-site mutation (*hm1-9*) at the

junction of intron 4/exon 5 that also produced a pre-mature stop codon (Table 1). The

new APR alleles were introgressed back into B73 for seven generations using CAPs

markers. Comparison of the resistance phenotype of the two new APR alleles with $Hm1^A$

revealed that all three APR alleles differ markedly from each other in this trait. *Hm1-3*

confers the highest level of resistance at all stages of development, followed by $Hm1^A$

and *Hm1-4* (Fig 5). This screen thus provided us with a series of APR alleles at the *hm1*

locus.

Table 1. The nature of molecular changes in the mutant alleles of *Hm1* generated by

mutagenesis and their respective disease/resistance phenotypes to infection by CCR1 at

maturity.

Allele No.	Disease Response	Mutation
Hm1-3	APR	T90M
Hm1-4	APR	V210M
hm1-5	Null	C82Y
hm1-6	Null	Nonsense
hm1-7	Null	Nonsense
hm1-8	Null	Nonsense
hm1-9	Null	Nonsense

Like *Hm1^A*, the new APR alleles encode HCTRs with intermediate activity

To evaluate if the HCTR activity encoded by *Hm1-3* and *Hm1-4* was also partially

compromised like that of *Hm1^A*, we used the aforementioned LC-MS/MS based activity

assay on samples derived from these two mutants as well as their positive and negative

controls. During weeks-3 and 7 (when APR plants are susceptible and resistant,

respectively), crude protein was extracted from the leaf tissue following inoculation with

CCR1. The HCTR activity of extracts from APR plants was found to be significantly

reduced when compared with B73 at both week-3 and week-7, indicating that HM1-3 and

HM1-4 proteins display partially compromised HCTR activity during both susceptible

and resistant plant ages like HM1^A (Fig 6). Furthermore, and consistent with HM1^A (Fig

3C), the levels of their HCTR did not change significantly with age (Fig 6),

demonstrating that the APR encoded by these new alleles was also expressed without a

concomitant increase in HCTR levels in mature plants.

Differences in the disease/resistance ratings of the new APR alleles predicted

corresponding differences in their HCTR activities. This indeed was found to be true. The

disease severity of APR plants at 3 weeks of age was found to be linearly correlated with

HCTR activity (Fig 5 and 6).   The APR allele with the highest degree of HCTR activity

was HM1-3, followed by HM1^A, and HM1-4 being the weakest (Fig 6). This variation in

enzymatic activity is consistent with the gradient of CCR1 resistance displayed by *Hm1-*

3, Hm1^A, and *Hm1-4* plants from strongest to weakest (Fig 5). At maturity, however,

plants carrying any of these weak alleles of *Hm1* were all indistinguishable from WT

B73. This was not the case with plants carrying only the null allele; they remained

uniformly susceptible to CCR1 infection even at maturity.

Modulation of photosynthesis output alters susceptibility to CCR1 in *Hm1^A*

seedlings

If the HCTR levels of the APR alleles remain largely uniform throughout plant

development, why then are weak alleles unable to confer protection at the seedling stage?

Some anecdotal observations that we have made about plants with APR alleles suggested

that the availability of fixed carbon for energy production played a role in determining

the ability of these weak alleles to suppress disease. The APR mutants always exhibited

greater disease susceptibility and prolonged sensitivity in winter greenhouses as

compared to the field. In the winter greenhouse, those plants closest to supplemental

lights were more resistant than plants growing distant from light fixtures. Third, the

resistance phenotype of APR alleles was compromised in the dominant *oil-yellow1-*

N1989 allele that has a chlorophyll deficiency (30).

400   We grew the *Hm1^A* plants at extended and reduced photoperiods to test the

hypothesis that energy availability from fixed carbon could determine disease

susceptibility in APR mutants. We grew *Hm1^A* B73 NIL homozygotes in a growth

chamber with a light regimen of 12h light (L) and 12h dark (D) for 2 weeks. Following

inoculation with CCR1 and overnight incubation, half of the seedlings were shifted to a

growth chamber adjusted at 18h L and 6h D. $Hm1^A$ seedlings grown in 12:12 L:D

photoperiod were susceptible to CCR1 when examined at 72 hours post-inoculation (hpi)

(Fig 7A) and showed no ability to suppress expanding lesions at 96 hpi (Fig 7B).

However, the $Hm1^A$ plants that were shifted to 18:6 L:D developed a resistant reaction

instead (Fig 7C and D). Thus, the seedling susceptibility of $Hm1^A$ conferred by low

HCTR activity could be overcome by providing a longer period of photosynthetically

active radiation.

We reasoned that if greater photosynthate availability provides enhanced

resistance sufficient to permit the weak $Hm1$ alleles to confer seedling resistance,

disruption of energy balance should negate their ability to confer any resistance. To test

this, we treated $Hm1^A$ and $Hm1$-3 homozygotes with extended darkness or with the

herbicide (3-(3,4-dichlorophenyl)-1,1-dimethylurea) (DCMU), which disrupts electron

transfer during the light reactions of photosynthesis. We inoculated two-week-old plants

with CCR1 and grew them in 14:10 L:D or 4:20 L:D. Extending the dark period of the

diurnal cycle resulted in an increase in disease severity after 7 days of growth for $Hm1$-3

plants (Fig 8A). If extended darkness renders plants susceptible to CCR1 due to a lack of

photosynthesis, then disruption of photosynthesis by herbicide treatment should effect the

same result. To test this, plants grown at 14:10 L:D were inoculated with CCR1 and

grown for 24 h. At 24 hpi, plants were divided into two groups with one receiving a

solution of DCMU applied to the leaf whorl and then grown for 6 days under the 14:10

L:D cycles. Observation of plants 7 dpi and 6 days after the DCMU treatment

demonstrated that a single DCMU application rendered both $Hm1^A$ and $Hm1$-3

homozygotes completely susceptible to CCR1 (Fig 8b).

Together, these two experiments demonstrate that light, and perhaps the energy

status of the plant, were key determinants of resistance to CCR1, and provide a direct link

between plant primary metabolism and physiology and disease resistance.

DISCUSSION

This study reveals one fundamental aspect of adult plant resistance (APR) in maize to

CCR1. APR alleles at the *hm1* locus are weak determinants of resistance that fail to

protect plants at the seedling stage but are sufficient to confer complete protection to

CCR1 at maturity. This conclusion is supported by multiple lines of evidence derived

from a combination of genetic, molecular, and biochemical experimentation. Genetic

analysis demonstrated that all APR alleles of *hm1* confer partial resistance that exhibits

haploinsufficiency (gene-dosage sensitivity) during most stages of plant development.

This contrasts with resistance conferred by the wild type (WT) alleles of *hm1* that are

completely dominant and protect every part of the plant regardless of age or maturity.

Plants with null alleles of *hm1,* on the other hand, are susceptible to CCR1 at all stages of

development. CCR1 infection typically results in plant lethality for these alleles, and the

ubiquitous nature of this pathogen makes them difficult to propagate in the field. The

APR alleles of *hm1* are recessive to the WT alleles (e.g., $Hm1^{B73}$) but dominant to null

alleles of *hm1* (e.g., $hm1^{Pr}$).

Consistent with the idea that APR is a symptom of weak or partial loss-of-

function alleles, we were able to generate two new APR alleles from the WT $Hm1^{B73}$

allele by mutagenesis with EMS. Five completely susceptible mutants were also

recovered in this mutant screen, which presumably encoded null mutations. In keeping

with these predictions, molecular analysis of these null alleles showed that four of the

five null mutants were the result of nonsense mutations that truncated their predicted

peptides by introducing premature stop codons. The fifth null mutant, which was caused

by a missense mutation, changed a highly conserved cysteine residue (C82Y) that is

perhaps critical for protein function. In sharp contrast, both novel APR alleles underwent

relatively conservative mutational changes: T90M in *Hm1-3* and V210M in *Hm1-4*. Even

Hm1^A, which differs from the WT *Hm1^{B73}* allele by five amino acids, seems to owe its

APR phenotype to a single L116H change. HCTR activity was encoded by all of the APR

alleles, indicating that none of these mutations completely eliminates the function of the

enzyme. Their HCTR activities were compromised, however, being intermediate to that

of the fully functional WT allele (which confers completely dominant protection) and the

recessive null *hm1* alleles, which impart no resistance to CCR1. These results indicate

that at some level HCTR activity is unable to deter the pathogen from colonizing maize

plants at the seedling stage but that level of activity is sufficient to prevent CCR1 from

colonizing at maturity.

A cause-and-effect relationship between APR and partial-loss-of-function alleles

of *hm1* is further substantiated by the correlation between the strength of the resistance

reaction conferred by an APR allele and its HCTR activity. The level of HCTR activity

matched perfectly with the strength of CCR1 resistance conditioned by the three APR

alleles. These results demonstrate that alleles of *hm1* with partial loss-of-function

mutations encode HCTR with a compromised activity and that the weaker activity results

in later onset of disease resistance. The resistance of seedlings encoding WT *Hm1*

demonstrates that efficient toxin deactivation is sufficient for maize seedlings to resist

CCR1 infection and, therefore, they express all of the required machinery for defense.

Likewise, mature plants lacking *hm1* function are completely susceptible, demonstrating

that HCTR is absolutely required for CCR1 infection, and mature maize plants are not

protected from toxin-mediated disease spread. These interpretations depend on the *in*

vitro assay correctly reflecting *in vivo* activity. Our *in vitro* HCTR activity assay did not

detect the *in vivo* activity of the enzyme but instead the level of the functional protein

present at a given time point. It is possible that *in vivo* activity did not correspond to the

in vitro activity identified by this method.

A seemingly mechanistic relationship between partial resistance and APR is also

evident in many other pathosystems where such genes have been cloned and studied in

detail. One example is that of *Cf-9B*, which mediates incomplete resistance to *C. fulvum*

in a developmentally specified fashion (17). Its paralog *Cf9*, which encodes a receptor

like protein, confers complete protection in all plant tissues at every stage of development

(31). Another example is that of *Xa21*, a receptor-like kinase that confers weak resistance

to Xanthomonas leaf blight in rice (14,15). The maize *Hm2* APR allele provides another

example. The weak CCR1 resistance provided by this allele is conferred by a truncated

HCTR (12).

In wheat, APR genes are rather common and have been used widely to protect

this crop from all forms of the disease caused by three different species of rust pathogens

(reviewed in Ellis et al., 2014). Even though APR genes confer little or no protection in

wheat seedlings, the broad-spectrum and durable nature of resistance provided by such

genes in adult plants have many breeders proclaim that breeding for rust resistance

should deploy only APR genes (32). Three of these wheat APR genes have been cloned

recently and, interestingly, they all appear to confer resistance by different mechanisms.

One of them, *Yr36*, a mediator of resistance to yellow rust, encodes a kinase with an

unusual domain (16), while *Lr34* and *Lr67*, both of which mediate APR to both rust and

powdery mildew pathogens, encode an ABC transporter and a hexose transporter,

respectively (13,19). Exactly how these genes confer APR remains unresolved, but one

thread that unifies them is their ability to confer only weak or partial resistance (32).

Overexpression of *Lr34*, one of the best studied APR genes, however, did enable it to

confer seedling resistance in durum wheat (33). Furthermore, the efficacy of this

transgene in conferring seedling resistance improved even further under extended

daylight conditions (34). These results echo what we have discovered with the APR

alleles in maize and suggest that the connection between weak resistance and APR is not

unique to the maize-CCR1 pathosystem but perhaps is a general feature of most disease

resistance genes that are weak and provide only partial protection.

A second major finding is that APR is not the result of the enhanced level or

activity of proteins encoded by APR alleles at the mature-plant stage. Rather, it must be

the result of a change in seedlings vs mature plants that affects differential resistance. It

was previously shown in a number of cases that the differential transcriptional activity of

an APR gene did not account for its APR phenotype (12,13,15,17,20). Here we extend

this to the level and HCTR activity of the accumulated HM1 proteins, which remained

stable across development. At the onset of APR, resistance manifests uniformly in all

parts of the plant, including the youngest leaves that are still unfurled, indicating that the

APR-inducing factor is not accumulated over a long period of time in aging tissues, but

rather is available in every part of the plant regardless of the age of the organ and

determined solely by the plant maturity.

Considering that the HCTR activity is present at equivalent levels in APR mutant

extracts regardless of plant stage, why then are seedlings susceptible? Though the studies

presented here do not resolve this question, the biochemical mechanism by which *hm1*

confers resistance to CCR1 suggests a plausible scenario. Although this resistance is

conferred by *hm1*-encoded HCTR, the HC-toxin (HCT) inactivation reaction requires the

reducing power of NADPH as a co-substrate. The direct involvement of NADPH in HC-

toxin reduction suggests this molecule could be very critical in regulating resistance in

the maize-CCR1 pathosystem. Supporting this hypothesis are our results showing that

light and photosynthetic activity have a great impact on resistance mediated by APR

alleles, either boosting them to confer seedling resistance or limiting them to prevent

APR.

Based on these results, it could well be the availability of NADPH that determines

the difference in resistance between seedling and mature stages in the *hm1* APR mutants.

NADPH is produced during the light reactions of photosynthesis, the C4 malate shuttle,

and sugar oxidation, along with other energy carriers such as ATP. Maize seedlings not

only have a limited photosynthetic capacity to assimilate carbon (C), but also strong sinks

to consume these assimilates (35). As a result, seedling leaves become C-deficient at

night and that may negatively impact the availability of NADPH and ATP. Since

NADPH is required for HCTR activity, its depletion at night may negatively impact the

activity of hypomorphic mutants of HCTR, thereby leaving HCT active to induce

susceptibility to CCR1. Bolstering this hypothesis is the observation that the *Hm1-3* and

Hm1-4 mutations occur at residues predicted to be critical for the binding of NADPH to

HCTR (36). The WT HCTR has likely evolved to require lower NADPH levels for

optimal activity, buffering any impact from the likely diurnal dip in its cofactor at night

and thereby allowing sufficient HCT inactivation. This scenario also explains why plants

with the APR genes become more resistant as they mature; the increased output of

photosynthates may outstrip the sink requirements, allowing excess photosynthates to be

stored as starch during the day and then used at night to fuel NADPH production.

Although several other aspects of plant bioenergetics are expected to support the

resistance phenotype of the APR genes in most pathosystems, NADPH appears to be the

most critical in energizing APR in the maize-CCR1 pathosystem. This, of course, is due

to the direct involvement of this molecule in the resistance mechanism mediated by

HCTR, and is supported by the fact that maize plants carrying the WT *Hm1* gene are

completely resistant to CCR1 at all stages of development, including as seedlings. This

study thus provides direct evidence linking, for the first time, primary host metabolism to

the realm of disease resistance in plants.

An intriguing implication of this study concerns the metabolic cost of resistance

in plants. This topic is not only of fundamental interest to plant pathologists and

entomologists but also has huge agricultural relevance (37–39). Our study demonstrates

that, compared to strong resistance, the weak form of resistance has a much higher

metabolic cost for the host. As shown in the case of APR, this cost can be so high that the

seedlings are not robust enough metabolically to express such resistance effectively. This

argument also extends to the quantitative form of resistance that is often relatively weak

and easily affected by the environment (40,41). An additional complication is that the

vulnerability of seedlings to diseases increases even further by conditions that

compromise photosynthesis. This phenomenon is analogous to what has been well

established in the animal world that malnutrition compromises the immunity of infants

much more than that of adults (42,43).

METHODS

Plant materials

The inbred P8 and landraces Pira and Enano were obtained from Germplasm Resources

Information Network (GRIN) of the U.S. National Plant Germplasm System. The CCR1-

susceptible maize inbred Pr, and the CCR1-resistant inbreds B73, Va35, W22, and Pr1 (a

near-isogenic line of Pr) were previously available in our research program. To determine

whether $Hm1^A$ is an allele of $Hm1$, P8 was crossed with Pr and the F_1 hybrid was

backcrossed to Pr to generate a BC_1F_1 population. Additionally, P8 was crossed with Pr1

and the resulting F_1 hybrid was testcrossed to the $hm1$ null stock Pr. Near-isogenic lines

of B73 displaying APR to CCR1 infection were generated by backcrossing $hm1$ APR

alleles with the B73 inbred, to determine the behavior of the APR alleles in a uniform

genetic background.

Pathogen growth and inoculation

The protocol for culturing CCR1 pathogen on carrot juice agar medium was the same as

previously described (23). One-hundred µl of 10^5 spores/ml of CCR1 conidial suspension

was used for leaf whorl inoculations. To study the phenotypic manifestation of APR by

the $Hm1^A$ allele, both homozygous ($Hm1^AHm1^A$ introgressed into B73) and heterozygous

($Hm1^Ahm1^{Pr}$ also in B73) plants were planted in isolation at the Purdue ACRE farm and

inoculated with 100 µl of 10^5 spores/mL of CCR1 spore suspension. Wild type B73

encoding $Hm1^{B73}$ and the susceptible $hm1^{Pr}$ B73 NIL plants were used as resistant and

susceptible controls, respectively. A fresh set of five rows of ~40 plants per row was

inoculated every week, and disease severity rating was determined 5 days post-

inoculation (dpi) as described previously (12). To determine if $Hm1^A$ is an allele of $Hm1$,

genetic crosses were made at the ACRE farm and the resulting segregating progeny was

evaluated under field conditions again at the ACRE farm.

Amplification of $Hm1^A$ genomic DNA

Four primer pairs were designed to amplify $Hm1^A$ based on its sequence homology with

$Hm1^{B73}$. The promoter region was amplified using a primer pair based on the promoter of

$hm1$ from Pr. Touchdown PCR (44) was carried out with 10 consecutive cycles of

denaturation at 94ºC for 30 sec, annealing at 63ºC for 30 sec with a decrease in 0.5ºC per

cycle to a "touchdown" of 58ºC, and extension at 72ºC for 30 sec; followed by 35 cycles

of 94ºC for 30 sec, 58ºC for 30 sec, and 72ºC for 45 sec. Three separate PCR reactions

were carried out for every primer so that any errors initiated by either the GoTaq DNA

Polymerase (Promega, Madison, WI, USA) or by sequencing could be ruled out. The

PCR products were cleaned by running them through an agarose column, BigDye

sequencing reactions were conducted, and the products precipitated with sodium acetate

and ethanol before final resuspension in 20 µl of double-distilled water (ddH$_2$O). These

samples were submitted to the Purdue Genomics Facility for low throughput sequencing.

Forward and reverse complementary sequences for each primer were compared using the

ClustalW2 multiple alignment program. In order to assemble the $Hm1^A$ sequence without

sequencing errors, only sequences with at least three perfect reads for each primer

sequence were considered.

Cloning of *Hm1^A* cDNA

P8 (*Hm1^A Hm1^A*) seeds were planted in 500M MetroMix and grown in Conviron growth chambers for two weeks. One-hundred μl of 10^5 spores/mL CCR1 spore suspension was used for whorl inoculation, and plants were covered with a hood overnight to maintain humidity required for spore germination and penetration into the leaf tissue. At 24 h post-inoculation (hpi), affected leaf tissue was collected from the plants and snap-frozen in liquid nitrogen. RNA was extracted with a Qiagen RNeasy extraction kit (Qiagen, Germantown, MD), and cDNA was synthesized by RT-PCR using random hexamer mix (New England BioLabs, Ipswich, MA).

Generating near-isogenic lines of B73 manifesting APR and susceptibility to CCR1

The P8 maize inbred line was crossed with the maize reference B73 inbred, and the resulting F_1 hybrid was backcrossed to B73. To introgress *Hm1^A* into the B73 inbred, the resulting BC_1F_1 progeny was backcrossed to B73 for six generations. Since the promotor region of *Hm1^{B73}* differed from that of *Hm1^A*, PCR-based markers designed from the promotor region were used for introgressing *Hm1^A* into B73 (primer sequences are available in Table S1). After the BC_7 generation, *Hm1^A* containing plants (*Hm1Hm1^A*) were self-pollinated to generate homozygous *Hm1^A* B73 NIL plants. Homozygous *Hm1^A* B73 NIL plants were identified with PCR-based markers and were self-pollinated to generate seed. Similar to *Hm1^A*, the two novel APR alleles *Hm1-3* and *-4* generated through EMS mutagenesis were introgressed into the B73 inbred for seven generations using a Cleaved Amplified Polymorphic sequences (CAPs) assay (primer sequences in Table S1). The restriction enzyme *NlaIII* (New England BioLabs, Ipswich, MA) was used to differentiate the *Hm1^{B73}* allele from the two novel APR alleles. Similar to the

novel APR alleles, the novel null allele *hm1-2* identified in the EMS-mutagenized B73

M_2 family screen was backcrossed for five generations into B73 using PCR-based

markers and self-pollinated to obtain a homozygous *hm1-2* NIL in B73. Marker-assisted

backcrossing using PCR-based genotyping was conducted on plants grown at the Purdue

Agronomy Center for Research and Education (ACRE) farm during the summer and in

the Purdue University Botany and Plant Pathology greenhouses during the winter season.

Transcriptional activity of *Hm1^A*

Hm1^A plants were inoculated with CCR1 spore suspension as described above at weekly

intervals from the seedling stage to maturity (week-1 through week-8). Total RNA was

isolated from CCR1-infected leaf tissue as described by Eggermont et al. (1996) and

treated with RNase-free DNase I to eliminate genomic DNA using the TURBO DNA-

free Kit (Ambion, Austin, TX). One µg of treated RNA was reverse-transcribed to cDNA

in a total volume of 25 µl using the iScript^{TM} cDNA Synthesis kit from Bio-Rad

(Hercules, CA). RT-PCR was conducted using gene specific primers with the maize actin

transcript as a control (see Table S1 for primer information). RT-PCR was conducted

under the following conditions: denaturation at 94°C for 1 min, annealing at 60°C for 1

min, extension at 72°C for 1 min, and a terminal extension steps for 10 min. 30 and 28

cycles of PCR were conducted to amplify *Hm1^A* and the control actin gene, respectively.

Amplified PCR products were separated on a 0.8% agarose gel to visualize the

expression of the *Hm1^A* transcript. Three replicates for each time point were used for this

experiment.

654   Additionally, qRT-PCR was conducted on cDNA from *Hm1^A* plants inoculated

with CCR1 at week-1, -3, -5, and -7 using gene specific primers. For relative

quantification, Molybdenum co-factor biosynthesis protein (MOL, GRMZM2G067176)

was used as a reference gene (46). All primer combinations had an efficiency of 90-

100%. Individual qRT-PCR reactions contained 5 µl of SYBR® Select Master Mix

(Applied Biosystems, Foster City, CA), 2 µl of cDNA template (20x dilution), and the

appropriate amount of forward and reverse primers plus water. A three-step qRT-PCR

amplification (40 cycles of 95°C for 5 s followed by 61°C for 20 s and 72°C for 30 s)

was performed using the Mx3000P qPCR system (Stratagene–Agilent Technologies,

Santa Clara, CA). Semi-quantitative RT-PCR was conducted using gene-specific primers

for $Hm1^A$ and the reference gene *Actin* (primer sequence in Table S5). Three replicates

for each time point were used for this experiment.

666   Translational activity of $Hm1^A$

For Western analysis, samples were collected at weekly intervals from week 2 through

week 8 after planting. Since $Hm1^A$ is infection inducible, plants were inoculated with

CCR1 for 24 h before collecting leaf samples. Protein was extracted using an alkaline

lysis protocol and quantified using the Bradford method. Equal amounts of the protein

(10 ug) was loaded onto SDS-PAGE gels and western blots were developed using a MAP

antibody raised against a synthesized 13 amino acid peptide (RPARDRLGELGFK)

corresponding to amino acids 312 to 324 of the HM1 peptide.

For the HCTR activity assay, $Hm1^{B73}$, $hm1^{Pr}$, and $Hm1^A$ plants grown in the field

were inoculated with 200 μl of 10^5 spores/mL CCR1 spore suspension into the leaf whorl

at weeks-3 and -7. Four biological replicates of three inoculated plants were sampled 24

hpi and stored at -80°C until used. Total plant protein was extracted using a protocol

adapted from Hayashi et al. (2005) and desalted using a Sephadex G-50 Fine column (GE

Healthcare, Chicago, IL). After determining protein concentration with a Bradford assay,

13.55 μg of protein was used to start reactions containing 25 mM Tris-HCl (pH 7.0), 160

mM NADPH, and 55 μM HC-toxin. The assays were run at 30°C in the dark for 45 min

and then stopped by the addition of 1.25 ml cold acetone. After centrifugation at 15,000 x

g for 15 min at 4°C, 10 μL of the supernatant was injected onto an Atlantis T3 column

(2.1 x 150 mm, 3 μm, 100 Å, Waters) maintained at room temperature and analyzed

using an Agilent 1200 series LC instrument coupled to an Agilent 6460 triple quadrupole

mass spectrometer (Agilent Technologies, Santa Clara, CA) at the Bindley Bioscience

Center in Purdue Discovery Park.

The solvent system contained solvents A (0.1% formic acid in ddH$_2$O) and B

(0.1% acetonitrile). The column was eluted with 85% A and 15% B (0 to 1 min),

followed by a linear gradient from 1 to 16 min to 40% A and 60% B, and a hold from 16

to 16.5 min at 40% A and 60% B. The column solvent was then reduced from 60% B to

15% B (16.5 to 17 min) and kept isocratic at 15% B from 17 to 22 min with a flow rate of

0.3 ml/min. HC-toxin (Sigma-Aldrich, St. Louis, MO) and its reduced form eluted from

the column at 8.5–11.5 min under these conditions. During the analysis, the column

effluent was directed to the MS/MS, with the Jetstream ESI set to positive mode with

nozzle and capillary voltages at 1000 – 4000 V. The nebulizer pressure was set at 35 psi,

the nitrogen drying gas was set at 325°C with a flow rate of 8 L/min, and the sheath gas

was held at 250°C at a flow rate of 7 L/min. Fragmentation was achieved with 70 V for

both analytes. Multiple reaction monitoring (MRM) was used to selectively detect HC-

toxin and its reduced form. The first quadrupole was set to transition between the [M-H]$^+$

of the analytes, whereas the last quadrupole monitored *m/z* 411 and 409 for reduced and

normal HC-toxin respectively. Each transition was monitored with a dwell time of 150

ms and collision energy of 15 V, with ultrapure nitrogen used as the collision gas. Mass

selection was achieved using the following ions: 439.3 for reduced HC-toxin and 437.3

for HC-toxin. Data were collected and analyzed via the MassHunter Workstation (version

B.06.00, Agilent Technologies, Santa Clara, CA), and peak areas were determined by

integration. Similar to *Hm1^A*, the HCTR activity of the new APR alleles generated by

targeted EMS mutagenesis (*Hm1-3* and *Hm1-4*) along with resistant (*Hm1*) and

susceptible (*hm1-2*) controls were also evaluated by LC-MS/MS.

Generating novel APR manifesting alleles of *Hm1* by EMS mutagenesis

The B73 (*Hm1Hm1hm2hm2*) maize inbred, which exhibits complete resistance to CCR1

at all stages of plant development (23), was the pollen parent for the targeted EMS

mutagenesis screen. The CCR1-susceptible maize inbred Pr (*hm1hm1hm2hm2*) (9,24),

which exhibited complete susceptibility to CCR1 at all stages of plant development, was

used as the female parent. This experiment was conducted in a greenhouse facility, as the

Pr plants do not survive in the field due to high levels of disease pressure.

To conduct pollen EMS mutagenesis, EMS stock solution was prepared by adding

1 ml of EMS (Sigma-Aldrich, St. Louis, MO) to 99 ml of paraffin oil (Sigma-Aldrich, St.

Louis, MO). Tassels of the Pr plants were removed before starting the experiment. On the

day of conducting pollen mutagenesis, EMS working solution was prepared by mixing 1

ml of EMS stock solution with 14 ml of paraffin oil. This working solution of EMS was

mixed gently for one hour to uniformly disperse the EMS in paraffin oil. B73 pollen was

collected in tassel bags, measured and transferred to a 50-ml Nalgene bottle. For every 1

ml of pollen collected, 10 ml of EMS working solution was added. The EMS-treated

pollen was placed on ice and mixed gently every 5 min for 45 min. About two to three

drops of EMS-treated B73 pollen was then applied to the silks of Pr ears. Ears from these

Pr plants were harvested 45 days after pollination. The M_1 seeds (~4500) obtained from

this genetic cross were planted at the Purdue ACRE farm. At both week-2 and week-5,

plants were whorl-inoculated with 100 µl of 10^5 spores/mL of CCR1 conidial suspension

and screened for their disease response one week post-inoculation.

Amplification of $Hm1^{B73}$ allele from heterozygous CCR1-susceptible mutants

Based on sequence polymorphisms between the wild type *Hm1* from B73, $Hm1^{B73}$ and

the null *hm1* allele from Pr, $hm1^{Pr}$, four primer pairs amplifying -560-bp of the promoter

region from the translation start site and the entire coding region of *Hm1* were designed

to preferentially amplify the WT $Hm1^{B73}$ from heterozygous M_1 plants (Fig S5), which

were obtained by crossing Pr plants with EMS-treated B73 pollen. Four overlapping

primer combinations (primer sequences in Table S1) were used to preferentially amplify

$Hm1^{B73}$ over the $hm1^{Pr}$ allele. Amplified PCR fragments were processed as described

above for $Hm1^{A}$ amplification and submitted to the Purdue Genomics Facility for low-

throughput sequencing.

Differential photoperiod treatments of $Hm1^{A}$ plants

$Hm1^{A}$ B73 NIL plants were grown in Conviron growth chambers providing a 12:12 L:D

photoperiod. Two-week-old $Hm1^{A}$ plants were inoculated with 100 µl of 10^5 spores/mL

of CCR1 spore suspension into the leaf whorl. CCR1-inoculated plants were incubated

overnight in a humidity chamber at 80% relative humidity. These plants were then

subjected to 12:12 L:D or 18:6 L:D photoperiods. The response reaction to CCR1

infection was evaluated every 24 h for a 96 h period. Digital photographs of lesion

progression were taken using a Canon EOS Digital Rebel XSi camera.

Additional extended darkness and DCMU treatment experiments were performed

in growth chambers on plants homozygous $Hm1^A$ and $Hm1-3$ in the B73 genetic

background. Plants were grown in a growth chamber under 14:10 L:D for two weeks. We

inoculated these plants with CCR1 and subjected them to two different light regimes,

14:10 L:D or 4:20 L:D. On a subset of CCR1 inoculated plants transferred to 14:10 L:D,

the herbicide (3-(3,4-dichlorophenyl)-1,1-dimethylurea) (DCMU) at a concentration of

100 μM was applied to the leaf whorl 24 hpi. Disease severity of these plants was

determined at 7 dpi.

ACKNOWLEDGEMENTS

This work was partially supported by GSJ's Hatch project (IND011280), by the IOS-NSF

grant 0547132 to G.S.J., and by the National Science Foundation Plant Genome Research

Program grant 1444503 to B.P.D. and G.S.J.

AUTHOR CONTRIBUTIONS

DSM, SC, SM and GSJ designed and performed linkage analysis studies. DSM did the

Southern blots and also constructed and screened the lambda library to clone $Hm1^A$. SM

conducted sequence alignment analysis with $Hm1$ orthologs and different maize inbreds.

SC performed $Hm1^A$ transcriptome analysis using semi-quantitative RT-PCR and AK

conducted qRT-PCR. KC and AK designed and performed HCTR assays. SC, SM, and

GJ designed and conducted EMS mutagenesis. SM, KC, and GJ screened EMS-generated

M_1 plants for their disease response to CCR1. SM, AD, BK, BD and GJ designed and

performed experiments to look at the effect of photosynthesis on APR. LD initially

provided P8, the inbred possessing *Hm1*A, and worked with GJ to characterize its APR.

SM, SC, KC, BD, and GJ wrote the paper. All authors reviewed the manuscript and LD

provided the detailed editorial changes.

REFERENCES

1. Johal GS, Gray J, Gruis D, Briggs SP. Convergent insights into mechanisms
determining disease and resistance response in plant–fungal interactions. Can J Bot.
1995 Dec 31;73(S1):468–74.

2. Jones JDG, Dangl JL. The plant immune system. Nature. 2006 Nov
16;444(7117):323–9.

3. Jones JDG, Vance RE, Dangl JL. Intracellular innate immune surveillance devices in
plants and animals. Science. 2016 Dec 2;354(6316):aaf6395-1– 8.

4. Dyck PL, Samborski, D. J., Anderson, R. G. Inheritance of adult-plant leaf rust
resistance derived from the common wheat varieties Exchange and Frontana. Can J
Genet Cytol. 1966 Dec 1;8(4):665–71.

5. Kus JV, Zaton K, Sarkar R, Cameron RK. Age-related resistance in Arabidopsis is a
developmentally regulated defense response to *Pseudomonas syringae*. Plant Cell.
2002 Feb;14(2):479–90.

6. Panter SN, Jones DA. Age-related resistance to plant pathogens. In: Advances in
Botanical Research [Internet]. Academic Press; 2002 [cited 2018 Mar 13]. p. 251–
80.

7. Whalen MC. Host defence in a developmental context. Mol Plant Pathol. 2005 May
1;6(3):347–60.

8. Develey-Rivière M-P, Galiana E. Resistance to pathogens and host developmental
stage: a multifaceted relationship within the plant kingdom. New Phytol. 2007 Aug
1;175(3):405–16.

9. Nelson OE, Ullstrup AJ. Resistance to leaf spot in maize: genetic control of
resistance to Race I of *Helminthosporium carbonum* Ull. J Hered. 1964 Sep
1;55(5):195–9.

10. Jones IT, Hayes JD. The effect of sowing date on adult plant resistance to *Erysiphe*
graminis f.sp. avenae in oats. Ann Appl Biol. 1971 May 1;68(1):31–9.

11. Abedon BG, Tracy WF. Corngrass 1 of maize (*Zea mays* L.) delays development of
adult plant resistance to common rust (*Puccinia sorghi* Schw.) and European corn
borer (*Ostrinia nubilalis* Hubner). J Hered. 1996 May 1;87(3):219–23.

12. Chintamanani S, Multani DS, Ruess H, Johal GS. Distinct mechanisms govern the
dosage-dependent and developmentally regulated resistance conferred by the maize
Hm2 gene. Mol Plant Microbe Interact. 2008;21(1):79–86.

13. Krattinger SG, Lagudah ES, Spielmeyer W, Singh RP, Huerta-Espino J, McFadden
H, et al. A putative ABC transporter confers durable resistance to multiple fungal
pathogens in wheat. Science. 2009 Feb 19;323(5919):1360–1363.

14. Song W-Y, Wang G-L, Chen L-L, Kim H-S, Pi L-Y, Holsten T, et al. A receptor
kinase-like protein encoded by the rice disease resistance gene, *Xa21*. Science. 1995
Dec 15;270(5243):1804–6.

15. Century KS, Lagman RA, Adkisson M, Morlan J, Tobias R, Schwartz K, et al.
Developmental control of Xa21-mediated disease resistance in rice. Plant J. 1999
Oct 1;20(2):231–6.

16. Fu D, Uauy C, Distelfeld A, Blechl A, Epstein L, Chen X, et al. A kinase-START
gene confers temperature-dependent resistance to wheat stripe rust. Science. 2009
Mar 6;323(5919):1357–60.

17. Panter SN, Hammond-Kosack KE, Harrison K, Jones JDG, Jones DA.
Developmental control of promoter activity Is not responsible for mature onset of
Cf-9B-mediated resistance to leaf mold in tomato. Mol Plant Microbe Interact. 2002
Nov 1;15(11):1099–107.

18. Milligan SB, Bodeau J, Yaghoobi J, Kaloshian I, Zabel P, Williamson VM. The root
knot nematode resistance gene *Mi* from tomato is a member of the leucine zipper,
nucleotide binding, leucine-rich repeat family of plant genes. Plant Cell. 1998 Aug
1;10(8):1307–19.

19. Moore JW, Herrera-Foessel S, Lan C, Schnippenkoetter W, Ayliffe M, Huerta-
Espino J, et al. A recently evolved hexose transporter variant confers resistance to
multiple pathogens in wheat. Nat Genet. 2015 Dec;47(12):1494–8.

20. McDowell JM, Williams SG, Funderburg NT, Eulgem T, Dangl JL. Genetic analysis
of developmentally regulated resistance to downy mildew (*Hyaloperonospora*
parasitica) in *Arabidopsis thaliana*. Mol Plant Microbe Interact. 2005 Nov
1;18(11):1226–34.

21. Kim S-D, Knoche HW, Dunkle LD. Essentiality of the ketone function for toxicity of
the host-selective toxin produced by *Helminthosporium carbonum*. Physiol Mol
Plant Pathol. 1987 May 1;30(3):433–40.

22. Meeley RB, Johal GS, Briggs SP, Walton JD. A biochemical phenotype for a disease
resistance gene of maize. Plant Cell. 1992 Jan 1;4(1):71–7.

23. Johal GS, Briggs SP. Reductase activity encoded by the HM1 disease resistance gene
in maize. Science. 1992 Nov 6;258(5084):985–7.

24. Multani DS, Meeley RB, Paterson AH, Gray J, Briggs SP, Johal GS. Plant–pathogen
microevolution: Molecular basis for the origin of a fungal disease in maize. Proc
Natl Acad Sci U S A. 1998 Feb 17;95(4):1686–91.

25. Ullstrup AJ. Further studies on a species of Helminthosporium parasitizing Corn.
Phytopathology. 1944;34(2):214–22.

26. Piffanelli P, Ramsay L, Benabdelmouna A, D'Hont A, Jørgensen JH, Hollricher K, et
al. A barley cultivation-associated polymorphism conveys resistance to powdery
mildew. Nature. 2004 Aug 19;430(7002):887.

27. Cook DE, Lee TG, Guo X, Melito S, Wang K, Bayless AM, et al. Copy number
variation of multiple genes at *Rhg1* mediates nematode resistance in soybean.
Science. 2012 Nov 30;338(6111):1206–9.

28. Zhang L, Peek AS, Dunams D, Gaut BS. Population genetics of duplicated disease-
defense genes, *hm1* and *hm2*, in maize (*Zea mays* ssp. *mays* L.) and its wild ancestor
(*Zea mays* ssp. *parviglumis*). Genetics. 2002 Oct 1;162(2):851–60.

29. Tam JP, Zavala F. Multiple antigen peptide. A novel approach to increase detection
sensitivity of synthetic peptides in solid-phase immunoassays. J Immunol Methods.
1989 Nov 13;124(1):53–61.

30. Sawers RJH, Viney J, Farmer PR, Bussey RR, Olsefski G, Anufrikova K, et al. The
maize *Oil Yellow1* gene encodes the I subunit of magnesium chelatase. Plant Mol
Biol. 2006 Jan 1;60(1):95–106.

31. Parniske M, Hammond-Kosack KE, Golstein C, Thomas CM, Jones DA, Harrison K,
et al. Novel disease resistance specificities result from sequence exchange between
tandemly repeated genes at the Cf-4/9 locus of tomato. Cell. 1997 Dec
12;91(6):821–32.

32. Ellis JG, Lagudah ES, Spielmeyer W, Dodds PN. The past, present and future of
breeding rust resistant wheat. Front Plant Sci. 2014 Nov 24;5:1–13.

33. Risk JM, Selter LL, Krattinger SG, Viccars LA, Richardson TM, Buesing G, et al.
Functional variability of the *Lr34* durable resistance gene in transgenic wheat. Plant
Biotechnol J. 2012 May 1;10(4):477–87.

34. Rinaldo A, Gilbert B, Boni R, Krattinger SG, Singh D, Park RF, et al. The *Lr34* adult
plant rust resistance gene provides seedling resistance in durum wheat without
senescence. Plant Biotechnol J. 2017 Jul;15(7):894–905.

35. Kalt-Torres W, Kerr PS, Usuda H, Huber SC. Diurnal changes in maize leaf
photosynthesis 1. Plant Physiol. 1987 Feb;83(2):283–8.

36. Dehury B, Patra MC, Maharana J, Sahu J, Sen P, Modi MK, et al. Structure-based
computational study of two disease resistance gene homologues (*Hm1* and *Hm2*) in
maize (*Zea mays* L.) with implications in plant-pathogen interactions. PLOS ONE.
2014 May 21;9(5):e97852.

37. Huot B, Yao J, Montgomery BL, He SY. Growth–defense tradeoffs in plants: A
balancing act to optimize fitness. Mol Plant. 2014 Aug 1;7(8):1267–87.

38. Karasov TL, Chae E, Herman JJ, Bergelson J. Mechanisms to mitigate the trade-off
between growth and defense. Plant Cell. 2017 Apr 1;29(4):666–80.

39. Guo Q, Major IT, Howe GA. Resolution of growth–defense conflict: mechanistic
insights from jasmonate signaling. Curr Opin Plant Biol. 2018 Aug 1;44:72–81.

40. Poland JA, Balint-Kurti PJ, Wisser RJ, Pratt RC, Nelson RJ. Shades of gray: the
world of quantitative disease resistance. Trends Plant Sci. 2009 Jan 1;14(1):21–9.

41. French E, Kim B-S, Iyer-Pascuzzi AS. Mechanisms of quantitative disease resistance
in plants. Semin Cell Dev Biol. 2016 Aug 1;56:201–8.

42. Katona P, Katona-Apte J. The interaction between nutrition and infection. Clin Infect
Dis. 2008 May 15;46(10):1582–8.

43. Walson JL, Berkley JA. The impact of malnutrition on childhood infections. Curr
Opin Infect Dis [Internet]. 2018 Mar 22 [cited 2018 Apr 3];Publish Ahead of Print.
Available from: https://insights.ovid.com/pubmed?pmid=29570495

44. Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS. "Touchdown" PCR to
circumvent spurious priming during gene amplification. Nucleic Acids Res. 1991
Jul 25;19(14):4008.

45. Eggermont K, Goderis IJ, Broekaert WF. High-throughput RNA extraction from
plant samples based on homogenisation by reciprocal shaking in the presence of a
mixture of sand and glass beads. Plant Mol Biol Report. 1996 Sep 1;14(3):273–9.

46. Hartwig T, Chuck GS, Fujioka S, Klempien A, Weizbauer R, Potluri DPV, et al.
Brassinosteroid control of sex determination in maize. Proc Natl Acad Sci. 2011
Dec 6;108(49):19814–9.

47. Hayashi K, Yoshida K, Matsui Y. A histone H3 methyltransferase controls epigenetic
events required for meiotic prophase. Nature. 2005 Nov 17;438(7066):374–8.

FIGURE LEGENDS

Fig 1. Developmental onset of the adult plant resistance phenotype of *Hm1^A*.

(A) A seedling *Hm1^A* leaf exhibiting susceptibility to *Cochliobolus carbonum* race 1

(CCR1) at the 2-week age. **(B)** A 9-week old *Hm1^A* leaf completely resistant to CCR1.

(C) The disease/resistance phenotype of *Hm1^A* plants homozygous and heterozygous

(*Hm1^A hm1^{Pr}*) for the APR allele to CCR1 at weekly intervals from week-1 through week-

10. Ratings were established by controls *Hm1^{B73} Hm1^{B73}* (rated 1 and resistant

throughout) and *hm1^{Pr} hm1^{Pr}* (rated 10 and susceptible throughout). All *hm1* alleles were

in the B73 genetic background. Error bars represent standard error calculated using R

statistical package.

Fig 2. Molecular characteristics of *Hm1^A*.

(A) Southern blot analysis of DNA of inbreds P8 (*Hm1^A Hm1^A*) and Pr1 (*Hm1^{Pr1} Hm1^{Pr1}*)

demonstrating that *Hm1^A* is a single copy gene. Sample genotypes (inbreds P8 or Pr1) are

indicated below the restriction endonuclease used for DNA digestions (BamHI, EcoRI, or

HindIII) and M corresponds to the DNA marker lane. **(B)** Schematic representation of the

gene structure of *Hm1^A* comprised by five exons (gray boxes) and four introns, identical

to *Hm1^{B73}*. The locations and the nature of five amino acids that differ between HM1^A

and HM1^{B73} are indicated by red lines. The locations of the start and termination codons

are also indicated.

Fig 3. Transcriptional and translational activities of *Hm1^A* during the seedling and

mature stages.

(A) Reverse transcription (RT)-PCR assay showing no change in $Hm1^A$ accumulation in

leaves from week-1 through week-8 after planting. The *actin* gene was used as a control.

(B) Quantitative real time PCR (qRT-PCR) measurements of the expression of $Hm1^A$ also

demonstrates no change in $Hm1^A$ accumulation across the time period when APR is

established. **(C)** Western blots showing that the level or stability of the $HM1^A$ protein do

not change over time during plant development. Equal amounts of protein were loaded

following quantification with the Bradford method. **(D)** *In vitro* HC-toxin reductase

(HCTR) assays showing that the relative enzymatic activity encoded by $Hm1^A$ is less than

$Hm1^{B73}$ but higher than $hm1^{Pr}$, the null allele. The specific activity of HCTR varies

between alleles but not over time between weeks 3 and 7 in any genotype. The HCTR

assay was based on the determination via LC-MS/MS of the amount of HC-toxin reduced

by leaf protein extracts from the leaves of all genotypes. Different letters indicate

significant differences between genotypes ($p_{adj} < 0.05$).

Fig 4. Design of the targeted EMS mutagenesis screen to generate new mutant

alleles of *Hm1*.

Pollen collected from the fully resistant inbred B73 ($Hm1^{B73}Hm1^{B73}$) was treated with

ethyl methanesulfonate (EMS) and used to pollinate ears of the fully susceptible inbred

Pr ($hm1^{Pr}hm1^{Pr}$) in a greenhouse. The resultant M1 seeds ($Hm1^{B73}/hm1^{Pr}$) were planted in

the field, inoculated with CCR1, and screened for disease resistance at both the seedling

stage and at maturity to identify rare susceptible mutants, designated as $Hm1^{B73*}/hm1^{Pr}$.

M1 mutants that were susceptible at the seedling stage that became resistant with the

progression of age were considered APR. Out of about 4,500 M1 plants screened, 7

susceptible mutants were found and two became resistant at maturity.

Fig 5. Relative strength of the three APR alleles of *hm1* in conferring protection

against CCR1.

Like $Hm1^A$, both new APR alleles (*Hm1-3 and Hm1-4*) were introgressed into B73 for six

generations for comparison of their resistance phenotypes. Plants homozygous for the

$Hm1^{B73}$ and *hm1-2* alleles were fully resistant and susceptible, respectively. Disease

resistance was evaluated three times, at week-2, week-5 and week-9 after planting, and a

scale of 1 (completely resistant) to 10 (completely susceptible) was used to rate the

interaction phenotypes. Letters represent whether differences among each age group were

significant ($p_{adj} < 0.05$). The relative order of strength observed was $Hm1^{B73} > Hm1\text{-}3 >$

$Hm1^A > Hm1\text{-}4 > hm1\text{-}2$.

Fig 6. *In vitro* enzymatic activities of HCTRs encoded by the new APR alleles of

***hm1*.**

Protein extracts from the leaf tissue of near-isogenic lines of the APR alleles *Hm1-3* and

Hm1-4 in the B73 background were used to conduct *in vitro* HCTR assays. The fully

resistant ($Hm1^{B73}$) and susceptible (*hm1-2*) alleles of *hm1* were used as controls. HCTR

activities, measured at age week-3 and week-7, relied on to determining the amount of

HC-toxin reduced via LC-MS/MS. Letters represent whether differences among each age

group were significant ($p_{adj} < 0.05$).

Fig 7. Resistance of *Hm1^A* seedlings to CCR1 in increased by extended photoperiod.

Two-week-old homozygous *Hm1^A* seedlings were inoculated with CCR1 and incubated

under two different photoperiods of 12 h daylight (12 h L:12 h D) and 18 h daylight (18 h

L:6 h D). *Hm1^A* seedlings grown under 12 h daylight were susceptible to CCR1 at 72 hpi

(A) and 96 hpi **(B)**. *Hm1^A* seedlings incubated under the extended photoperiod of 18 h

light exhibited notably enhanced resistance at both 72 hpi **(C)** and 96 hpi **(D)**.

Fig 8. Decreased photoperiod and photosynthesis inhibition by DCMU enhanced the

susceptibility of APR genotypes to CCR1.

(A) Two-week-old homozygous *Hm1^A* and *Hm1-3* B73 NIL plants were inoculated with

CCR1 and incubated with a shortened photoperiod of 4:20 L:D or longer 14:10 L:D

photoperiod. Plants grown under a decreased photoperiod were completely susceptible to

CCR1 while control plants were relatively less susceptible. **(B)** *Hm1^A* and *Hm1-3* B73

NIL plants were grown for two-weeks in the longer photoperiod conditions (14:10 L:D)

and half of the plants were sprayed with DCMU, a photosynthesis inhibiting herbicide.

Application of DCMU rendered both *Hm1^A* and *Hm1-3* plants highly susceptible to

CCR1 compared to control plants. Pictures were taken 6 days after inoculation.

TABLES

Table 1. The nature of molecular changes in the mutant alleles of *Hm1* generated by

mutagenesis and their respective disease/resistance phenotypes to infection by CCR1 at

maturity.

A *Hm1^A* seedling leaf **B** *Hm1^A* mature leaf **C**

$Hm1^A hm1^{Pr}$
$Hm1^A Hm1^A$
$hm1^{Pr} hm1^{Pr}$
$Hm1^{B73} Hm1^{B73}$

Figure 1. Developmental onset of the adult plant resistance phenotype of *Hm1^A*.

(A) A seedling *Hm1^A* leaf exhibiting susceptibility to *Cochliobolus carbonum* race 1 (CCR1) at the 2-week age. **(B)** A 9-week old *Hm1^A* leaf completely resistant to CCR1. **(C)** The disease/resistance phenotype of *Hm1^A* plants homozygous and heterozygous (*Hm1^A hm1^{Pr}*) for the APR allele to CCR1 at weekly intervals from week-1 through week-10. Ratings were established by controls *Hm1^{B73} Hm1^{B73}* (rated 1 and resistant throughout) and *hm1^{Pr} hm1^{Pr}* (rated 10 and susceptible throughout). All *hm1* alleles were in the B73 genetic background. Error bars represent standard error calculated using R statistical package.

Figure 2. Molecular characteristics of *Hm1ᴬ*.

(A) Southern blot analysis of DNA of inbreds P8 (*Hm1ᴬHm1ᴬ*) and Pr1 (*Hm1ᴾʳ¹Hm1ᴾʳ¹*) demonstrating that *Hm1ᴬ* is a single copy gene. Sample genotypes (inbreds P8 or Pr1) are indicated below the restriction endonuclease used for DNA digestions (BamHI, EcoRI, or HindIII) and M corresponds to the the DNA marker lane. **(B)** Schematic representation of the gene structure of *Hm1ᴬ* comprised by five exons (grey boxes) and four introns, identical to *Hm1ᴮ⁷³*. The locations and the nature of five amino acids that differ between HM1ᴬ and HM1ᴮ⁷³ are indicated by red lines. The locations of the start and termination codons are also indicated.

Figure 3. Transcriptional and translational activities of *Hm1^A* during the seedling and mature stages.

(A) Reverse transcription (RT)-PCR assay showing no change in *Hm1^A* accumulation in leaves from week-1 through week-8 after planting. The *actin* gene was used as a control. **(B)** Quantitative real time PCR (qRT-PCR) measurements of the expression of *Hm1^A* also demonstrates no change in *Hm1^A* accumulation across the time period when APR is established. **(C)** Western blots showing that the level or stability of the HM1^A protein do not change over time during plant development. Equal amounts of protein were loaded following quantification with the Bradford method. **(D)** *In vitro* HC-toxin reductase (HCTR) assays showing that the relative enzymatic activity encoded by *Hm1^A* is less than *Hm1^B73* but higher than *hm1^Pr*, the null allele. The specific activity of HCTR varies between alleles but not over time between weeks 3 and 7 in any genotype. The HCTR assay was based on the determination via LC-MS/MS of the amount of HC-toxin reduced by leaf protein extracts from the leaves of all genotypes. Different letters indicate significant differences between genotypes ($p_{adj} < 0.05$).

Figure 4. Design of the targeted EMS mutagenesis screen to generate new mutant alleles of *Hm1*.

Pollen collected from the fully resistant inbred B73 (*Hm1^{B73}Hm1^{B73}*) was treated with ethyl methanesulfonate (EMS) and used to pollinate ears of the fully susceptible inbred Pr (*hm1^{Pr}hm1^{Pr}*) in a greenhouse. The resultant M1 seeds (*Hm1^{B73}/hm1^{Pr}*) were planted in the field, inoculated with CCR1, and screened for disease resistance at both the seedling stage and at maturity to identify rare susceptible mutants, designated as *Hm1^{B73*}/hm1^{Pr}*. M_1 mutants that were susceptible at the seedling stage that became resistant with the progression of age were considered APR. Out of about 4,500 M_1 plants screened, 7 susceptible mutants were found and two became resistant at maturity.

Figure 5. Relative strength of the three APR alleles of *hm1* in conferring protection against CCR1.

Like *Hm1^A*, both new APR alleles (*Hm1-3 and Hm1-4*) were introgressed into B73 for six generations for comparison of their resistance phenotypes. Plants homozygous for the *Hm1*^B73 and *hm1-2* alleles were fully resistant and susceptible, respectively. Disease resistance evaluations were done three times, at week-2, week-5 and week-9 after planting, and a scale of 1 (completely resistant) to 10 (completely susceptible) was used to rate the interaction phenotypes. Letters represent whether differences among each age group were significant ($p_{adj} < 0.05$). The relative order of strength observed was *Hm1*^B73 > *Hm1-3* > *Hm1^A* > *Hm1-4* > *hm1-2*.

Figure 6. *In vitro* enzymatic activities of HCTRs encoded by the new APR alleles of *hm1*.

Protein extracts from the leaf tissue of near-isogenic lines of the APR alleles *Hm1-3* and *Hm1-4* in the B73 background were used to conduct *in vitro* HCTR assays. The fully resistant (*Hm1B73*) and susceptible (*hm1-2*) alleles of *hm1* were used as controls. HCTR activities, measured at age week-3 and week-7, relied on to determining the amount of HC-toxin reduced via LC-MS/MS. Letters represent whether differences among each age group were significant ($p_{adj} < 0.05$).

Fig. 7

A 72 hpi   **B** 96 hpi   **C** 72 hpi   **D** 96 hpi

12 h L:12 h D   **18 h L:6 h D**

Figure 7. Resistance of *Hm1^A* seedlings to CCR1 in increased by extended photoperiod.

Two-week-old homozygous *Hm1^A* seedlings were inoculated with CCR1 and incubated under two different photoperiods of 12 h daylight (12 h L:12 h D) and 18 h daylight (18 h L:6 h D). *Hm1^A* seedlings grown under 12 h daylight were susceptible to CCR1 at 72 hpi **(A)** and 96 hpi **(B)**. *Hm1^A* seedlings incubated under the extended photoperiod of 18 h light exhibited notably enhanced resistance at both 72 hpi **(C)** and 96 hpi **(D)**.

Figure 8. Decreased photoperiod and photosynthesis inhibition by DCMU enhanced the susceptibility of APR genotypes to CCR1.

(**A**) Two-week-old homozygous *Hm1ᴬ* and *Hm1-3* B73 NIL plants were inoculated with CCR1 and incubated with a shortened photoperiod of 4:20 L:D or longer 14:10 L:D photoperiod. Plants grown under a decreased photoperiod were completely susceptible to CCR1 while control plants were relatively less susceptible. (**B**) *Hm1ᴬ* and *Hm1-3* B73 NIL plants were grown for two-weeks in the longer photoperiod conditions (14:10 L:D) and half of the plants were sprayed with DCMU, a photosynthesis inhibiting herbicide. Application of DCMU rendered both *Hm1ᴬ* and *Hm1-3* plants highly susceptible to CCR1 compared to control plants. Pictures were taken 6 days after inoculation.