Insect wings and body wall evolved from ancient leg segments

Heather S. Bruce* and Nipam H. Patel
Department of Molecular and Cell Biology, University of California, Berkeley, CA
*Correspondence: hbruce@berkeley.edu

Researchers have long debated the origin of insect wings. One theory proposes that the proximal portion of the ancestral crustacean leg became incorporated into the body[1], which moved the leg's epipod (multi-functional lobe, e.g. gill) dorsally, up onto the back to form insect wings[2]. Another theory proposes that the dorsal insect body wall co-opted crustacean epipod genes to form wings[3]. Alternatively, wings may be derived from both leg and body wall (dual origin)[4]. To determine whether wings can be traced to ancestral, pre-insect structures, or arose by co-option, comparisons are necessary between insects and other arthropods more representative of the ancestral state, where the hypothesized proximal leg region is not fused to the body wall. To do so, we examined the function of five leg gap genes in the crustacean *Parhyale hawaiensis* and compared this to previous functional data from insects. Here we show, using CRISPR-Cas9 mutagenesis, that leg segment deletion phenotypes of all five leg gap genes in *Parhyale* align to those of insects only by including the hypothesized fused ancestral proximal leg region. We also argue that possession of eight leg segments is the ancestral state for crustaceans. Thus, *Parhyale* incorporated one leg segment into the body, which now bears the tergal plate, while insects incorporated two leg segments into the body, the most proximal one bearing the wing. We propose a model wherein much of the body wall of insects, including the entire wing, is derived from these two ancestral proximal leg segments, giving the appearance of a "dual origin" [4-9]. This model explains many observations in favor of either the body wall origin, proximal leg origin, or dual origin of insect wings.

Arthropod appendages are key to their spectacular success, but their incredible diversity has complicated comparisons between distantly related species. The origin of the most debated appendage, insect wings, pivots on the alignment of leg segments, because wings may be derived from an epipod (e.g. gill or plate, Fig. 1b)[5] of ancestral leg segments that fused to the body[2], or alternatively, may represent a co-option of the epipod-patterning pathway by the insect body wall[3], or a combination of both[4,6-10]. To answer this, functional comparisons are necessary between insects and arthropods more representative of the ancestral state, where the hypothesized proximal leg region is not fused to the body wall.

Towards this aim, we examined five leg gap genes, *Distalless* (*Dll*), *Sp6-9*, *dachshund* (*dac*), *extradenticle* (*exd*), and *homothorax* (*hth*), in an amphipod crustacean, *Parhyale hawaiensis*. While we have documented their expression at several developmental stages (Fig. S1), our comparative analysis does not rely solely on these expression patterns, given that expression is not always a reliable indication of function, and expression is often temporally dynamic[11]. Instead, we have systematically knocked out these genes in *Parhyale* using CRISPR-Cas9 mutagenesis and compared this to our understanding of their function in *Drosophila* and other insects (Figs. 2, S2).

Insects have six leg segments, while *Parhyale* has seven (Fig. 1). In insects, *Dll* is required for the development of leg segments 2 – 6[12-15] . In *Parhyale*, the canonical *Dll* gene, *Dll*-e[16-18], is required for the development of leg segments 3 – 7 (Fig. 2b). In insects, *Sp6-9*[19] is

required for the development of leg segments 1 – 6 [20], and in addition in *Drosophila*, loss of *Sp6-*
9 (i.e. D-Sp1[19]) occasionally transforms the leg towards wing and lateral body wall identity[20]. In
Parhyale, *Sp6-9*[19,21] is required for the development of leg segments 2 – 7 (Fig. 2c), and in some
legs, segment 2 is occasionally transformed towards a leg segment 1 identity (Fig S3). In
Drosophila, *dac* is required in the trochanter through proximal tarsus (leg segments 2 – 4, and
first tarsus)[21,22]. *Parhyale* has two *dac* paralogs. *dac1* does not seem to be expressed in the legs
or have a morphologically visible knockout phenotype. *dac2* is required to pattern leg segments
3 – 5 (Fig. 2d). *Exd* and *hth* are expressed in the body wall and proximal leg segments of
insects[23-26] and *Parhyale*[27] (Fig S1). They form heterodimers and therefore have similar
phenotypes [23-26]. In insects, *exd* or *hth* knockout results in deletions/fusions of the coxa through
proximal tibia (leg segments 1 – 3, and proximal tibia)[24-28]. In *Parhyale*, *exd* or *hth* knockout
results in deletions/fusions of the coxa through proximal carpus (leg segments 1 – 4, and
proximal carpus; Figs. 2e, f). In both insects[23,24,28] and *Parhyale*, the remaining distal leg
segments are sometimes transformed towards a generalized thoracic leg identity (compare Fig. 2
e, f and Fig S4). In both insects[23-26] and *Parhyale* (Fig. S4), *exd* or *hth* knockout results in
deletions/fusions of body segments.
In summary, the expression and function of *Dll*, *Sp6-9*, *dac*, *exd*, and *hth* in *Parhyale* are
shifted distally by one segment relative to insects. This shift is accounted for if insects fused an
ancestral proximal leg segment to the body wall (Fig. 2g). Thus, there is a one-to-one homology
between insect and *Parhyale* legs, displaced by one segment, such that the insect coxa is
homologous to the crustacean basis, the insect femur is the crustacean ischium, and so on for all
leg segments. This also means that at least part of the insect body wall is homologous to the
crustacean coxa.
While our CRISPR-Cas9 data is agnostic regarding the origin of the insect wing, we
noted that *Parhyale* has what appears to be an epipod, the tergal plate, emerging dorsal to the
coxa (Fig. 3). Since epipods are characteristic of leg segments[29], this suggested to us that
Parhyale, like most groups of crustaceans, might retain an additional proximal leg segment, the
ancestral precoxa (Fig. 3a). We therefore examined carefully dissected *Parhyale* using confocal
and brightfield microscopy and identified what appears to be a precoxa (Fig. 3). In fact, this
structure has been previously described in amphipods[30], but it was assumed to be body wall.
However, this structure meets the criteria for a true leg segment: it protrudes from the body wall;
it forms a true, muscled joint; and it extends musculature to another leg segment (Figs. 3 and
S5)[29,31,32]. Importantly, the tergal plate emerges not from the body wall, but from this precoxa
(Fig. 3e). Molecular evidence of a precoxa in *Parhyale* is provided by Clark-Hachtel in the
accompanying manuscript. They show that the tergal plate is indeed an epipod just like the coxal
plate, and that *nubbin*, a marker of arthropod leg segments, is expressed in a distinct stripe above
the *Parhyale* tergal plate, suggesting there is indeed a precoxa leg segment here. Thus, much of
what appears to be lateral body wall in *Parhyale* is in fact proximal leg.
Since insects evolved from crustaceans, if the insect coxa is homologous to the
crustacean basis, then one would expect to find two leg segments incorporated into the insect
body wall, each equipped with an epipod (Fig. 4). As predicted, two leg-like segments can be
observed proximal to the coxa in basal hexapods[1] including collembolans[33], as well as in the
embryos of many insects[9,34,35]. In insect embryos, these two leg-like segments flatten out before
hatching to form the lateral body wall[1,9,33-37] (Fig 1c). Furthermore, insects indeed appear to have
two epipods proximal to the insect coxa. When "wing" genes are depleted in insects via RNAi,
two distinct regions are affected: the wing, but also the protruding plate adjacent to the leg (Fig.

1c)[38-41]. Based on these data, we hypothesize that insects have incorporated the ancestral precoxa
and crustacean coxa into the body wall, with the precoxa epipod later forming the wing and the
crustacean coxa epipod later forming the plate.
The results presented here may settle the long-standing debate regarding the origin of
insect wings as derived from (a) the epipod of the leg, (b) the body wall, or, more recently, (c)
from both (dual-origin[4,6-10]). Our model accounts for all observations in favor of either of these
hypotheses, including the dorsal position of insect wings relative to their legs, the loss of
ancestral leg segments in insects, the two-segmented morphology of the insect subcoxa in both
embryos and adults, the complex musculature for flight, and the shared gene expression between
wings and epipods. The realization that crustaceans have a precoxa accounts for the apparent
"dual origin" of insect wings: much of what appears to be insect body wall is in fact the remnant
of two ancestral leg segments.

105
106
107   Fig. 1. Crustacean and insect legs. (a) Adult *Parhyale*, with third thoracic leg (T3) outlined. (b)
108   Cartoon of *Parhyale* T3. The coxal plate extends over the leg. (c) Adult *Oncopeltus*, with T2
109   outlined. Inset shows magnified proximal leg, with body wall plate extending over the leg. (d)
110   Cartoon of *Oncopeltus* T2 leg.

Fig. 2. Knockout phenotypes of leg gap genes. (a-f) *Parhyale* CRISPR-Cas9 phenotypes in
dissected third thoracic legs (T3). Graded cyan in f indicates deletion/fusion of proximal leg
segment 5. (g) Leg gap gene function in *Parhyale* and insects aligns only if insects incorporated
the red leg segment into the body wall (0). Color bars correspond to remaining leg segments
following knockout, transparent bars indicate deleted leg segments. Open bar in *dac* indicates
slight extension of *dac* function into tarsus 1 of insects. Coxal plate (Cp), gill (G), tergal plate
(Tp). Scale bar 50um.

120
Fig. 3. Evidence for a precoxa in *Parhyale*. (a) Phylogeny based on Oakley 2012, precoxa
references in supplements. (b) Confocal image of *Parhyale* hatchling, autofluorescent cuticle in
cyan. T5 and T6 tergal plates (dotted outlines). (c) Confocal image of *Parhyale* hatchling,
autofluorescent cuticle in cyan, muscle phalloidin stain in red. Compare the blocks of simple,
anterior-posterior muscles of the body to the orthogonal, complexly arranged muscles of the leg
segments. Outline of tergal plates (dotted lines) relative to orthogonal muscle. (d) Brightfield
image of right half of adult *Parhyale,* sagittal dissection, internal tissues removed, lateral view.
Wire used to position sample (w). The same orthogonal muscles in b are visible as striations that
continue above the wire. The precoxa forms a joint with the coxa[30] (arrow). The dorsal limit of
the precoxa is unclear, but the most conservative estimate is to begin at the joint (arrow) and
follow the leg up to where it meets the adjacent leg, denoted by (<). Comparing (<) and (→), it
can be seen that the precoxa protrudes quite a bit from the body wall. However, the precoxa
appears to continue farther up the body wall (compare orthogonal muscle striations). (e)
Posterior-lateral view of right T6, looking edge-on at tergal plate. The tergal plate (dotted
outline) emerges from the precoxa (contiguous pink between ←, >, and ---), just as the coxal
plate (dashed line) emerges from the coxa. In c, d, coxa is red (coxal plate not shaded), gills
(teal) partially cut for visibility, basis orange, precoxa pink. Note that all three plates (tergal,
coxal, and basal) form contiguous cuticle with their leg segment, i.e. there is no distinguishing
suture.

140
141
Fig 4. Gene function alignment (a) and proposed leg segment homologies (colors) between an
ancestral crustacean, *Parhyale*, and insects (b). Ancestral precoxa epipod (ep), *Parhyale* tergal
plate (Tp), and insect wing are homologous (pink). Ancestral coxa epipod, *Parhyale* coxal plate
(Cp) and gill (G), and insect plate (see Fig. 1c) are homologous (red). *Parhyale* basal plate (Bp).
Insect numbering based on crustaceans.

147
148
Fig S1. Expression of leg gap genes in whole embryos and dissected third thoracic legs (T3). (a –
d): *Dll*-e. (e – h): *Sp6-9*. (i – l): dac2. (m – p): *exd*. (q – t): *hth*. Embryonic expression data for
Dll-e[16-18], *Sp6-9*[19], and *exd* and *hth*[27] have been previously characterized, but not at the level of
individual leg segments. (d) *Dll*-e is expressed in leg segments 3 – 7; in the interior of the tergal
plate (Tp), coxal plate (Cp), and gill (G), where it may be playing a sensory role, similar to the
expression of Dll that patterns sensory hairs in the Drosophila wing margin[13]; and marks the
bristle (^) of leg segment 2. This bristle is deleted in *Dll*-e KO (compare Fig. 2a, b). (h) *Sp6-9* is
expressed in leg segments 2 – 7. (l) dac2 is expressed in leg segments 3 – 5. Expression in
segment 5 may be stronger at other time points. (p) *exd* is expressed in the body wall through leg
segment 5, and perhaps a little in 6. *Exd* is not expressed in the gill (not visible here). (t) *hth* is
expressed in the body wall through leg segment 3. *Hth* is not expressed in the gill. Note that both
insects and *Parhyale* share a peculiar disparity between *hth* expression and function, wherein *hth*
knockout deletes one more leg segment than would be predicted by the *hth* expression domain.

162
163
Fig. S2. T7 endonuclease assay to confirm CRISPR-Cas9 mutagenesis. For each gene, one or
two wild type (WT) hatchlings were assayed, and one, two, or three KO hatchlings were assayed.
T7 endonuclease was either added (+) or not added (–) to the heteroduplex mixture. In brief, a
~1kb region flanking the CRISPR-Cas9 target site by at least 300bp to either side was amplified
by PCR from either WT or KO hatchlings. The purified PCR products were denatured, then
slowly cooled to allow WT DNA and mutant DNA with indels to anneal, resulting in a "bubble"
of unpaired DNA (heteroduplex) at the target site. T7 endonuclease was added to the (+)
samples, incubated, and run on a 1.5% agarose gel. KO animals are mosaic, so if the target site
was cut, the indels will cause heteroduplexes when annealed with either a WT strand, or a
different indel. When a single deletion is present, each half of the cut heteroduplex adds up to
approximately 1kb (see *Sp6-9* KO 1 and 2). Some deletions are large enough to be seen without
the T7 endonuclease assay (see *Dll*-e KO), and some hatchlings had multiple deletions which
produced multiple bands when cut with T7 (see *exd* KO 1, *hth* KO 2, dac2 KO).

Gene	sgRNA	total injected	# dead	death %	# hatch w/phenotype	% phenotype of hatched
Dll-e	1+2	151	45	30%	57	54%
exd	1+2	206	90	44%	86	74%
exd	1	204	102	50%	84	82%
exd	2	173	36	21%	85	62%
hth	1+2	124	71	57%	32	60%
hth	1	131	30	23%	36	36%
hth	2	99	62	63%	22	59%
dac2	1+2	80	28	35%	41	79%
dac2	1	84	31	37%	9	17%
dac2	2	88	18	20%	16	23%
Sp6-9	1+2	165	88	53%	51	66%
Sp6-9	1	54	22	41%	9	28%
Sp6-9	2	37	3	8%	15	44%

177
178 Table 1. CRISPR-Cas9 injection numbers. Two sgRNAs per gene were made, and either one or
179 both were injected as indicated. Both guides for each gene gave the same phenotype. # dead is
180 the number of embryos that did not survive to hatching. For each gene, sgRNA 1 and 2 produced
181 the same phenotypes.

182
183
Fig S3. *Sp6-9* knockout sometimes causes a homeotic transformation of orange leg segment 2
towards a red leg segment 1 identity in jumping legs (thoracic legs T6 – 8). In WT jumping legs
(a, c), orange leg segment 2 is very large and wide, due to the epipod on this segment (compare
to skinny orange leg segment of WT T3 leg, inset in c). In WT T6 legs (a), the red coxal plate is
bilobed, while in the WT T8 legs (c), the coxal plate is small and oval. In T6 *Sp6-9* KO (b), the
epipod of orange leg segment 2 is bilobed, indicating a transformation towards red leg segment
1. In T8 *Sp6-9* KO (d), the large epipod of orange leg segment 2 has been reduced to the size and
shape of the coxal plate, indicating a transformation towards red leg segment 1. Note that the
tergal plates are unaffected (d, pink, Tp), which is similar to *Drosophila Sp6-9* knockouts, where
the wings are unaffected[20]. The bilobed shape of the transformed T6 basal plate demonstrates
that these are transformations towards a coxal plate rather than tergal plate, because the tergal
plates are never bilobed. Therefore, these represent a homeotic transformation of one leg
segment into another. This argues that the transformation of *Drosophila* leg to wing following
loss of *Sp6-9* is also a transformation of one leg segment into another, and thus that insect wings
are appendicular. Scale bar 50um.

199
200
Fig. S5. *Parhyale* precoxa forms a true, muscled joint and extends musculature to another leg
segment. Confocal images. (a) Phalloidin stain of muscle in right half of *Parhyale* hatchling.
Contrast simple, anterior-posterior body muscles to orthogonal, complexly arranged leg muscles.
No muscles cross the coxa-basis joint, as noted by Boxshall 1998. Note that all three plates
(tergal, coxal, and basal) form contiguous cuticle with their leg segment, i.e. there is no
distinguishing suture between leg segment and epipod. (b) Optical section showing superficial
muscles of right half. Confocal colors are partially desaturated: cuticle in grey-blue, muscle in
grey-pink. The precoxa forms two articulations with the coxa: an anterior, bifurcated, load-
bearing hinge articulation (arrowhead), and a posterior gliding articulation ($\rightarrow$) (see also Fig. 3e).
Both articulations were previously identified by Hessler 1982. Coxa is red (coxal plate not
shaded, to focus on joints), basis is orange, precoxa is magenta pink. Adjacent legs meet on their
ventral sides at (<) and on their dorsal sides at (*). Outline of tergal plate (dashed line) relative to
muscle and joints shows that tergal plate emerges from precoxa. Muscles in green insert on the
precoxa-coxa joint, indicating that this is a true joint, and not merely a point of flexure in the
exoskeleton (annulation)[29,31,32]. The shorter, anterior muscle originates in the protruding precoxa
to insert on the rim of the next leg segment, the coxa. This muscle is therefore an intrinsic
muscle, a hallmark of a true leg segment[29,31,32]. (c) Confocal of dissected left half, medial view.
Coxal plate and basis partially cut. The precoxa forms a joint with two articulations with the
coxa: an anterior, bifurcated, load-bearing hinge articulation (arrowhead), and a posterior gliding
articulation (arrow). Orthogonal muscles visible as striations on T4 precoxa. (d) Close-up of left
T4, medial-anterior view, showing bifurcated hinge articulation.

METHODS

BIOINFORMATICS

      Partial or complete sequences for *Parhyale Dll*, *Sp6-9*, *Exd*, and *Hth* have been previously identified. These were >99% identical at the nucleotide level to sequences in the *Parhyale* assembled transcriptome. In order to confirm their orthology, identify potential *Parhyale* paralogs and identify *Parhyale dac*, we ran reciprocal best Blast hit searches. For each gene, orthologs from several arthropods and vertebrates were downloaded from NCBI and EMBL and aligned against the *Parhyale* transcriptome[42] using standalone NCBI blastp. The *Parhyale* hits with the lowest E-values were used to run a blastp against the NCBI database, restricted to Arthropoda. We confirmed that the original set of orthologs from several arthropods were the best hits to our *Parhyale* candidates (i.e. were each other's reciprocal best Blast hits). These reciprocal best Blast hits are listed in the tables below, and were deposited in Genbank under Accession Numbers MG457799 - MG457804.

      No *Parhyale* buttonhead/Sp5 was recovered in the assembled transcriptome. Buttonhead/Sp5 was also not found in the genome of the related amphipod Hyalella azteca. The assembled transcriptome only recovered fragments of *Parhyale* Sp1-4, so the previously sequenced *Parhyale* Sp1-4 (CBH30980.1) was used for the table below (asterisk).

      Parhyale has three Dll paralogs, which appear to be an amphipod-specific duplication, because a related amphipod, Hyalella azteca, also has these same three Dll paralogs. The three *Parhyale* Dll paralogs had the lowest E-values to all Dll orthologs examined, but which of the three *Parhyale* Dll paralogs had the lowest E-value was variable, as expected for a clade-specific duplication.

      The coding region for *Parhyale* exd and hth in the assembled transcriptome are longer than those previously identified. Exd is 204 amino acids longer, and hth is 166 amino acids longer. This explains the higher-than-expected E-values between the *Parhyale* exd and hth sequences identified previously and the *Parhyale* exd and hth sequences used in this study.

Extradenticle		
Query id	Subject id	E-value
Daphnia_pulex exd EFX62563.1	*Parhyale* exd MG457802	8.00E-177
Drosophila exd AAF48555.1	*Parhyale* exd MG457802	7.00E-173
Hyalella exd XP_018011298.1	*Parhyale* exd MG457802	2.00E-166
Parhyale exd CAO98909.1	*Parhyale* exd MG457802	6.00E-126
Tribolium exd NP_001034501.1	*Parhyale* exd MG457802	1.00E-173
Homo Pbx1 NP_002576.1	*Parhyale* exd MG457802	3.00E-166

Homothorax		
Query id	Subject id	E-value
Daphnia hth EFX75948.1	*Parhyale* hth MG457803	0
Drosophila hth NP_476578.3	*Parhyale* hth MG457803	6.00E-179
Homo Meis2 AAH07202.1	*Parhyale* hth MG457803	1.00E-148

Hyalella hth XP_018016731.1	*Parhyale* hth MG457803	0
Parhyale hth CAO98908.1	*Parhyale* hth MG457803	0
Tribolium hth NP_001034489.1	*Parhyale* hth MG457803	0

253
254

Sp6-9, Sp1-4, buttonhead/Sp5		
Query id	Subject id	E-value
Drosophila btd NP_511100.1	*Parhyale* Sp6-9 MG457804	4.00E-47
Drosophila Sp1-4 NM_142975.3	* *Parhyale* Sp1-4 CBH30980.1	5.00E-62
Drosophila Sp6-9 NP_727360.1	*Parhyale* Sp6-9 MG457804	6.00E-109
Homo Sp4 NP_003103.2	* *Parhyale* Sp1-4 CBH30980.1	2.00E-66
Homo Sp5 NP_001003845.1	*Parhyale* Sp6-9 MG457804	7.00E-62
Homo Sp8 NP_874359.2	*Parhyale* Sp6-9 MG457804	3.00E-105
Hyalella Sp1-4 XP_018012207.1	* *Parhyale* Sp1-4 CBH30980.1	0
Hyalella Sp6-9 XP_018014881.1	*Parhyale* Sp6-9 MG457804	0
Parhyale Sp1-4 CBH30980.1	* *Parhyale* Sp1-4 CBH30980.1	0
Parhyale Sp6-9 CBH30981.1	*Parhyale* Sp6-9 MG457804	0
Tribolium btd NP_001107792.1	*Parhyale* Sp6-9 MG457804	7.00E-59
Tribolium Sp1-4 XP_015833716.1	*Parhyale* Sp6-9 MG457804	3.00E-62
Tribolium Sp6-9 XP_008198341.1	*Parhyale* Sp6-9 MG457804	6.00E-159

255
256

Distalless		
Query id	Subject id	E-value
Drosophila Dll ACL83212.1	PhDllL2	2.00E-54
Drosophila Dll ACL83212.1	PhDllL1	2.00E-48
Drosophila Dll ACL83212.1	PhDlle MG457801	4.00E-42
Homo DLX-2 AAB40902.1	PhDlle MG457801	3.00E-35
Homo DLX-2 AAB40902.1	PhDllL2	6.00E-35
Homo DLX-2 AAB40902.1	PhDllL1	3.00E-34
Hyalella DLX-2 XP_018023955.1	PhDlle MG457801	0
Hyalella DLX-2 XP_018023955.1	PhDllL1	1.00E-49
Hyalella DLX-2 XP_018023955.1	PhDllL2	3.00E-45
Hyalella DLX-6 XP_018023956.1	PhDllL2	4.00E-102
Hyalella DLX-6 XP_018023956.1	PhDllL1	1.00E-51
Hyalella DLX-6 XP_018023956.1	PhDlle MG457801	1.00E-40
Hyalella unchar. protein XP_018023484.1	PhDllL1	8.00E-83
Hyalella unchar. protein XP_018023484.1	PhDllL2	0.89
Parhyale Dll-e ACT78885.1	PhDlle MG457801	0

Parhyale Dll-e ACT78885.1	PhDllL1	7.00E-48
Parhyale Dll-e ACT78885.1	PhDllL2	1.00E-44
Tribolium Dll AAG39634.1	PhDllL1	7.00E-48
Tribolium Dll AAG39634.1	PhDllL2	1.00E-46
Tribolium Dll AAG39634.1	PhDlle MG457801	5.00E-39

257
258

Dachshund		
Query id	Subject id	E-value
Daphnia pulex dac EFX90187.1	*Parhyale* Dac1 MG457799	3.00E-67
Drosophila dac AAF53538.3	*Parhyale* Dac2 MG457800	2.00E-64
Homo dach2 Q96NX9	*Parhyale* Dac1 MG457799	4.00E-52
Hyalella Dac1 XP_018011787.1	*Parhyale* Dac1 MG457799	7.00E-109
Hyalella Dac1 XP_018011787.1	*Parhyale* Dac2 MG457800	2.00E-55
Hyalella Dac2 XP_018011801.1	*Parhyale* Dac2 MG457800	0
Hyalella Dac2 XP_018011801.1	*Parhyale* Dac1 MG457799	1.00E-59
Tribolium dac1 XP_015834662.1	*Parhyale* Dac2 MG457800	6.00E-72

259
260
261
262 IN SITU PRIMER SEQUENCES
263

Primer name	product size	seq
hth FORWARD	941	GTTATGGGCTCCGTACCTGA
hth REVERSE	941	GCCAGCTGTTTCTTCTGGTC
exd FORWARD	734	AGCGAGTCCTCAACAAAGGA
exd REVERSE	734	AGGAGGCGTGTGCTATTCTG
Dll FORWARD	725	TGGGTCCAGTTCAACCTCTC
Dll REVERSE	725	GACATCGTCCTCCAAAGCAT
dac 1 FORWARD	638	GGAGAGCAGAGGGGACTTTT
dac 1 REVERSE	638	CCACTTCACGACCTCCTCAT
dac 2 FORWARD	699	CTTCAACCCCCTCCAGTACA
dac 2 REVERSE	699	TGTCTGTCGTCGTCTTCCTG
Sp6-9 FORWARD	789	CAAATGGCTCGCATGTATTG
Sp6-9 REVERSE	789	CAGTGCGTTCAAACTTCCAA

264
265
CLONING AND RNA PROBE SYNTHESIS
Total RNA was extracted from a large pool of *Parhyale* embryos at multiple stages of
embryogenesis, from Stages 12 to 26 using Trizol. cDNA was generated using Superscript III.

Primers were generated with Primer3 (http://bioinfo.ut.ee/primer3-0.4.0), with a preferred
product size of 700bp, and did not include the DNA binding domain. Inserts were amplified with
Platinum Taq (ThermoFisher 10966026), ligated into pGem T-Easy vectors (ProMega A1360),
and transformed into E coli. The resulting plasmids were cleaned with a QiaPrep mini-prep kit
(Qiagen A1360), and sequenced to verify the correct insert and determine sense and anti-sense
promoters. In situ templates were generated by PCR from these plasmids using M13F/R primers
and purified with Qiagen PCR Purification kit (Qiagen 28104). The resulting PCR products were
used to make DIG-labeled RNA probes (Roche 11175025910) using either T7 or Sp6 RNA
polymerase. RNA probes were precipitated with LiCl, resuspended in water, and run on an
agarose gel to check that probes were the correct size, and concentration was determined using a
Nanodrop 10000. Probes were used at 1-5ng/uL concentration.
280
IN SITU PROTOCOL
Embryo collection, fixation, and dissection as previously described[43]. In situ performed as
previously described[44]. In brief, embryos were fixed in 4% paraformaldehyde (PFA) in artificial
seawater for 45 minutes, dehydrated to methanol, and stored overnight at -20C to discourage
embryos from floating in later hybridization solution (Hyb) step. Embryos were rehydrated to
1xPBS with 0.1% Tween 20 (PTw), post-fixed for 30 minutes in 9:1 PTw:PFA, and washed in
PTw. Embryos were incubated in Hyb at 55C for at least 36 hours. Embryos were blocked with
5% normal goat serum and 1x Roche blocking reagent (Roche 11096176001) in PTw for 30
minutes. Sheep anti-DIG-AP antibody (Roche 11093274910) was added at 1:2000 and incubated
for 2 hours at room temperature. Embryos were developed in BM Purple (Roche 11442074001)
for a few hours to overnight. After embryos were sufficiently developed, they were dehydrated to
methanol to remove any pink background, then rehydrated to PTw. Embryos were then moved to
1:1 PBS:glycerol with 0.1mg/mL DAPI, then 70% glycerol in PBS.
294
CRISPR-CAS9 GUIDE RNA GENERATION, INJECTION, AND IMAGING
Guide RNAs were generated using ZiFit[45,46] as previously described[47]. sgRNAs were ordered
from Synthego. Injection mixes had a final concentration of 333ng/uL Cas9 protein, 150ng/uL
sgRNA (for both single and double guide injection mixes), and 0.05% phenol red for
visualization during injection, all suspended in water. One- or two-cell embryos were injected
with approximately 40 – 60 picoliters of sgRNA mixture as previously described[47]. Resulting
knockout hatchlings were fixed in 4% paraformaldehyde in artificial seawater at 4C for 1 – 2
days, then moved to 70% glycerol in 1xPBS. Dissected hatchling limbs were visualized with
Zeiss 700 and 780 confocal microscopes using the autofluorescence in the DAPI channel. Z-
stacks were assembled with Volocity. Hatchling images were desaturated, levels adjusted, and
false-colored using Overlay with Adobe Photoshop CS6.
306
T7 ENDONUCLEASE I ASSAY
Genomic primers were designed using Primer3, and flanked the target site by at least 400bp to
either side. DNA isolation and subsequent PCR amplification of the region of interest was
modified from previously described protocols[48]. Genomic DNA was amplified directly from
fixed hatchlings in 70% glycerol using ExTaq (Takara RR001A). The resulting PCR products
were purified with the Qiaquick PCR purification kit (Qiagen 28104). Heteroduplexes were
annealed and digested by T7 endonuclease I according to NEB protocols (NEB M0302L). The

digested products were run out on a 1.5% agarose gel. Genomic primers used for the T7
endonuclease I assay are listed below.
316
317 GENOMIC DNA PRIMERS
318

Primer name	product size	seq
exd left	907	CTTGAGATTCGTTCAGGTGCA
exd right	907	TTCTCCCCAGTTCCTTGCAA
hth left	943	TGTTCGTGTACCCGCAGAT
hth right	943	TCGGGCATACTAGAAGGCAG
Sp6-9 left	935	GCCCAGCTACTAACGATTTTCA
Sp6-9 right	935	GATCCGCTTCCTGACAGTTG
Dll-e left	922	GGAATGGTGAAGGAAGAGCG
Dll-e right	922	TCAGCAGTGCAGACTCATGT
dac2 left	983	CACGCGACACTCATACACAG
dac2 right	983	GATGCTCCTCCCACCGAATA

333
334
335 PRECOXA PHYLOGENY REFERENCES
336 Branchiura[49,50]. Mystacocarida[51,52]. Ostracoda[51,53-55]. Copepoda[29,51,56,57]. Cirripedia[55].
337 Decapoda[51,58-61]. Leptostraca[55]. Stomatopod[49,62]. Amphipoda[30]. Notostraca[55]. Spinicaudata[63].
338 Remipedia[33,51]. Collembola[33,64,65]. Insecta[1,29,34,35,37,66].
339
340
341
342 AUTHOR CONTRIBUTIONS
343 H.S.B. and N.H.P. conceived of the experiments. H.S.B. performed all experiments, conceived
344 of model, and wrote the manuscript. N.H.P. edited and revised the manuscript.

References

1. Snodgrass, R. E. Morphology and mechanism of the insect thorax. *Smithsonian Miscellaneous Collections* **80,** (1927).

2. Kukalová-Peck, J. Origin of the insect wing and wing articulation from the arthropodan leg. *Can. J. Zool.* **61,** 1618–1669 (1983).

3. Snodgrass, R. E. *Principles of insect morphology.* (McGraw-Hill Book Company, Inc, 1935).

4. Clark-Hachtel, C. M. & Tomoyasu, Y. Exploring the origin of insect wings from an evo-devo perspective. *Curr Opin Insect Sci* **13,** 77–85 (2016).

5. Averof, M. & Cohen, S. M. Evolutionary origin of insect wings from ancestral gills. *Nature* **385,** 627–630 (1997).

6. Clark-Hachtel, C. M. & Tomoyasu, Y. Two sets of wing homologs in the crustacean, Parhyale hawaiensis. *bioRxiv.* (2017). doi:10.1101/236281

7. Requena, D. *et al.* Origins and Specification of the Drosophila Wing. *Current Biology* **27,** 3826–3836.e5 (2017).

8. Prokop, J. *et al.* Paleozoic Nymphal Wing Pads Support Dual Model of Insect Wing Origins. *Curr. Biol.* **27,** 263–269 (2017).

9. Mashimo, Y. & Machida, R. Embryological evidence substantiates the subcoxal theory on the origin of pleuron in insects. *Sci Rep* **7,** 1–9 (2017).

10. Niwa, N. *et al.* Evolutionary origin of the insect wing via integration of two developmental modules. *Evol. Dev.* **12,** 168–176 (2010).

11. Estella, C. A dynamic network of morphogens and transcription factors patterns the fly leg. *Curr. Top. Dev. Biol.* **98,** 173–198 (2012).

12. Estella, C., Rieckhof, G., Calleja, M. & Morata, G. The role of buttonhead and Sp1 in the development of the ventral imaginal discs of Drosophila. *Development* **130,** 5929–5941 (2003).

13. Campbell, G. & Tomlinson, A. The roles of the homeobox genes aristaless and Distal-less in patterning the legs and wings of Drosophila. *Development* **125,** 4483–4493 (1998).

14. Angelini, D. R. & Kaufman, T. C. Functional analyses in the hemipteran Oncopeltus fasciatus reveal conserved and derived aspects of appendage patterning in insects. *Developmental Biology* **271,** 306–321 (2004).

15. Beermann, A. *et al.* The Short antennae gene of Tribolium is required for limb development and encodes the orthologue of the Drosophila Distal-less protein. *Development* **128,** 287–297 (2001).

16. Serano, J. M. *et al.* Comprehensive analysis of Hox gene expression in the amphipod crustacean Parhyale hawaiensis. *Developmental Biology* **409,** 297–309 (2015).

17. Browne, W. E., Price, A. L., Gerberding, M. & Patel, N. H. Stages of embryonic development in the amphipod crustacean, Parhyale hawaiensis. *genesis* **42,** 124–149 (2005).

18. Liubicich, D. M. *et al.* Knockdown of Parhyale Ultrabithorax recapitulates evolutionary changes in crustacean appendage morphology. *Proc. Natl. Acad. Sci. U.S.A.* **106,** 13892–13896 (2009).

19. Schaeper, N. D., Prpic, N.-M. & Wimmer, E. A. A clustered set of three Sp-family
genes is ancestral in the Metazoa: evidence from sequence analysis, protein domain
structure, developmental expression patterns and chromosomal location. *BMC Evol.*
Biol. **10,** 88 (2010).
20. Estella, C. & Mann, R. S. Non-Redundant Selector and Growth-Promoting
Functions of Two Sister Genes, buttonhead and Sp1, in Drosophila Leg
Development. *PLoS Genet* **6,** e1001001 (2010).
21. Mardon, G., Solomon, N. M. & Rubin, G. M. dachshund encodes a nuclear protein
required for normal eye and leg development in Drosophila. *Development* **120,**
3473–3486 (1994).
22. Tavsanli, B. C. *et al.* Structure–function analysis of the Drosophila retinal
determination protein Dachshund. *Developmental Biology* **272,** 231–247 (2004).
23. Mito, T. *et al.* Divergent and conserved roles of extradenticle in body segmentation
and appendage formation, respectively, in the cricket Gryllus bimaculatus.
Developmental Biology **313,** 67–79 (2008).
24. Ronco, M. *et al.* Antenna and all gnathal appendages are similarly transformed by
homothorax knock-down in the cricket Gryllus bimaculatus. *Developmental Biology*
313, 80–92 (2008).
25. Rauskolb, C., Smith, K. M., Peifer, M. & Wieschaus, E. extradenticle determines
segmental identities throughout Drosophila development. *Development* **121,** 3663–
3673 (1995).
26. Wu, J. & Cohen, S. M. Proximodistal axis formation in the Drosophila leg:
subdivision into proximal and distal domains by Homothorax and Distal-less.
Development **126,** 109–117 (1999).
27. Prpic, N.-M. & Telford, M. J. Expression of homothorax and extradenticle mRNA in
the legs of the crustacean Parhyale hawaiensis: evidence for a reversal of gene
expression regulation in the pancrustacean lineage. *Dev Genes Evol* **218,** 333–339
(2008).
28. Casares, F. & Mann, R. S. The ground state of the ventral appendage in Drosophila.
Science **293,** 1477–1480 (2001).
29. Boxshall, G. A. The evolution of arthropod limbs. *Biol. Rev.* **79,** 253–300 (2004).
30. Hessler, R. R. The Structural Morphology of Walking Mechanisms in
Eumalacostracan Crustaceans. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **296,** 245–298 (1982).
31. Boxshall, G. *Arthropod Limbs and their Development. Arthropod Biology and*
Evolution 241–267 (2013). doi:10.1007/978-3-662-45798-6_11
32. Shultz, J. W. Morphology of locomotor appendages in Arachnida: evolutionary
trends and phylogenetic implications. *Zool J Linn Soc* **97:1-56,** (1989).
33. Bäcker, H., Fanenbruck, M. & Wägele, J. W. A forgotten homology supporting the
monophyly of Tracheata: The subcoxa of insects and myriapods re-visited.
Zoologischer Anzeiger - A Journal of Comparative Zoology **247,** 185–207 (2008).
34. Kobayashi, Y. Formation of Subcoxae-1 and 2 in Insect Embryos: The Subcoxal
Theory Revisited. *Proc Arthropod Embryol Soc Jpn* **48,** 33–38 (2017).
35. Kobayashi, Y., Niikura, K., Oosawa, Y. & Takami, Y. Embryonic development of
Carabus insulicola (Insecta, Coleoptera, Carabidae) with special reference to

435		external morphology and tangible evidence for the subcoxal theory. *J. Morphol.*
436		**274,** 1323–1352 (2013).
437	36.	Hansen, H. J. *Studies on the Arthropoda III. On the Comparative Morphology of the*
438		*Appendages in the Arthropoda. B. Crustacea (Supplement), Insecta, Myriapoda*
439		*and ....* (Copenhagen: Gyldendalske Boghandel, 1930).
440	37.	Deuve, T. The epipleural field in hexapods. *Annales De La Societe Entomologique*
441		*De France* **37,** 195–231 (2001).
442	38.	Clark-Hachtel, C. M., Linz, D. M. & Tomoyasu, Y. Insights into insect wing origin
443		provided by functional analysis of vestigial in the red flour beetle, Tribolium
444		castaneum. *Proc. Natl. Acad. Sci. U.S.A.* **110,** 16951–16956 (2013).
445	39.	Ohde, T., Yaginuma, T. & Niimi, T. Insect morphological diversification through
446		the modification of wing serial homologs. *Science* **340,** 495–498 (2013).
447	40.	Medved, V. *et al.* Origin and diversification of wings: Insights from a neopteran
448		insect. *Proc. Natl. Acad. Sci. U.S.A.* **112,** 15946–15951 (2015).
449	41.	Wang, D. *et al.* spalt is functionally conserved in Locusta and Drosophila to promote
450		wing growth. *Sci Rep* **7,** 1–9 (2017).
451	42.	Kao, D. *et al.* The genome of Parhyale hawaiensis: a model for animal development,
452		regeneration, immunity and ligno-cellulose digestion. *Elife* 1–76 (2016).
453	43.	Rehm, E. J., Hannibal, R. L., Chaw, R. C., Vargas-Vila, M. A. & Patel, N. H.
454		Fixation and Dissection of Parhyale hawaiensis Embryos. *Cold Spring Harbor*
455		*Protocols* **4,** (2009).
456	44.	Rehm, E. J., Hannibal, R. L., Chaw, R. C., Vargas-Vila, M. A. & Patel, N. H. In Situ
457		Hybridization of Labeled RNA Probes to Fixed Parhyale hawaiensis Embryos. *Cold*
458		*Spring Harbor Protocols* **4,** (2009).
459	45.	Sander, J. D. *et al.* ZiFiT (Zinc Finger Targeter): an updated zinc finger engineering
460		tool. *Nucleic Acids Res.* **38,** W462–W468 (2010).
461	46.	Sander, J. D., Zaback, P., Joung, J. K., Voytas, D. F. & Dobbs, D. Zinc Finger
462		Targeter (ZiFiT): an engineered zinc finger/target site design tool. *Nucleic Acids*
463		*Res.* **35,** W599–W605 (2007).
464	47.	Martin, A. *et al.* CRISPR/Cas9 Mutagenesis Reveals Versatile Roles of Hox Genes
465		in Crustacean Limb Specification and Evolution. *Curr. Biol.* **26,** 14–26 (2016).
466	48.	Gloor, G. B., Nassif, N. A., Johnson-Schlitz, D. M., Preston, C. R. & Engels, W. R.
467		Targeted gene replacement in Drosophila via P element-induced gap repair. *Science*
468		**253,** 1110–1117 (1991).
469	49.	Schram, F. R. *Crustacea.* (Oxford University Press, USA, 1986).
470	50.	Tanzola, R. D. & Villegas-Ojeda, M. A. Argulus ventanensis sp. n.(Crustacea,
471		Branchiura) parasite of Hypsiboas pulchellus tadpoles (Anura, Hylidae). *Pan Am J A*
472		*S* **12,** 218–226 (2017).
473	51.	Boxshall, G. Comparative limb morphology in major crustacean groups: the coxa-
474		basis joint in postmandibular limbs. *Arthropod Relationships* (1997).
475	52.	Stachowitsch, M. & Proidl, S. *The Invertebrates.* (University of Texas Press, 1992).
476	53.	Horne, D. J. Homology and homoeomorphy in ostracod limbs. *Hydrobiologia* **538,**
477		55–80 (2005).
478	54.	Cohen, A. C., Martin, J. W. & Kornicker, L. S. Homology of Holocene ostracode
479		biramous appendages with those of other crustaceans: the protopod, epipod, exopod
480		and endopod. *Lethaia* **31:251-265,** (1998).

55.   Hansen, H. J. *Studies on Arthropoda. II. At the Expense of the Rask-Ørsted Fund.*
(Copenhagen: Gyldendalske Boghandel, 1925).
56.   Karanovic, T. Two new genera and three new species of subterranean cyclopoids
(Crustacea, Copepoda) from New Zealand, with redescription of Goniocyclops
silvestris Harding, 1958. *Contributions to Zoology* **74, 3/4,** 223–254 (2005).
57.   Fiers, F. & Jocque, M. Leaf litter copepods from a cloud forest mountain top in
Honduras (Copepoda: Cyclopidae, Canthocamptidae). *Zootaxa* **3630,** 270–290
(2013).
58.   Borradaile, L. A. XXVII.— Notes upon Crustacean limbs. *Journal of Natural*
History Series 11 **17,** 193–213 (1926).
59.   Snodgrass, R. E. *A textbook of arthropod anatomy.* (Ithaca, N.Y., Comstock Pub.
Associates, 1952).
60.   Hong, S. Y. Development of epipods and gills in some pagurids and brachyurans.
Journal of Natural History Series 11 **22,** 1005–1040 (1988).
61.   Fukuda, Y. Existence of Precoxa in the Thoracic Appendages of Decapod
Crustaceansand its Functions. *Proc. Japan. Soc. Sys. Zool.* **52,** 47–64 (1994).
62.   Watling, L., Hof, C. H. J. & Schram, F. R. The place of the Hoplocarida in the
Malacostracan Pantheon. *Journal of Crustacean Biology* **20,** 1–11 (2000).
63.   Ferrari, F. D. & Grygier, M. J. Comparative morphology among trunk limbs of
Caenestheriella gifuensis and Leptestheria kawachiensis (Crustacea: Branchiopoda:
Spinicaudata). *Zool J Linn Soc* **139,** 547–564 (2003).
64.   Sun, X. & Wu, D. Review of Chinese Oligaphorurini (Collembola, Onychiuridae)
with descriptions of two new Palaearctic species. *Zookeys* **192,** 15–26 (2012).
65.   Fjellberg, A. *The Collembola of Fennoscandia and Denmark, Part II:*
Entomobryomorpha and Symphypleona. (BRILL, 2007).
66.   Coulcher, J. F., Edgecombe, G. D. & Telford, M. J. Molecular developmental
evidence for a subcoxal origin of pleurites in insects and identity of the subcoxa in
the gnathal appendages. *Sci Rep* **5,** 1–8 (2015).
509