Title: Insect wings and body wall evolved from ancient leg segments

One Sentence Summary: CRISPR-Cas9 knockout of leg gap genes in a crustacean reveals that insect wings are not novel structures, they evolved from crustacean leg segments

Authors: Heather S. Bruce* and Nipam H. Patel

Affiliations: Department of Molecular and Cell Biology, University of California, Berkeley, CA

*Correspondence to: hbruce@berkeley.edu

Abstract: The origin of insect wings has long been debated. Central to this debate is whether wings evolved from an epipod (outgrowth, e.g., a gill) on ancestral crustacean leg segments, or represent a novel outgrowth from the dorsal body wall that co-opted some of the genes used to pattern the epipods. To determine whether wings can be traced to ancestral, pre-insect structures, or arose by co-option, comparisons are necessary between insects and arthropods more representative of the ancestral state, where the hypothesized proximal leg region is not fused to the body wall. To do so, we examined the function of five leg patterning genes in the crustacean Parhyale hawaiensis and compared this to previous functional data from insects. By comparing gene knockout phenotypes of leg patterning genes in a crustacean with those of insects, we show that two ancestral crustacean leg segments were incorporated into the insect body, moving the leg's epipod dorsally, up onto the back to form insect wings. Thus, our data shows that much of the body wall of insects, including the entire wing, is derived from these two ancestral proximal leg segments. This model explains all observations in favor of either the body wall origin or proximal leg origin of insect wings. Thus, our results show that insect wings are not novel structures, but instead evolved from existing, ancestral structures.

Main Text: The origin of insect wings has fascinated researchers for over 130 years. One theory proposes that the proximal portion of the ancestral crustacean leg became incorporated into the body (*1*), which moved the leg's epipod (lobe-shaped outgrowth, e.g. gill) dorsally, up

onto the back to form insect wings (*2*). Another theory proposes that the wing is a novel

outgrowth from the dorsal body wall that co-opted some of the genes used to pattern the epipods

of leg segments (*3*). Alternatively, wings may be derived from a combination of leg and body

wall (dual origin, (*4*)). To determine whether wings can be traced to ancestral, pre-insect

structures, or arose by co-option, comparisons are necessary between insects and other

arthropods more representative of the ancestral state, where the hypothesized proximal leg region

is not fused to the body wall.

Towards this aim, we examined five leg gap genes, *Distalless* (*Dll*), *Sp6-9*, *dachshund*

(*dac*), *extradenticle* (*exd*), and *homothorax* (*hth*), in an amphipod crustacean, *Parhyale*

hawaiensis. While we have documented their expression at several developmental stages (Fig.

S1), our comparative analysis does not rely solely on these expression patterns, given that

expression is not always a reliable indication of function, and expression is often temporally

dynamic (*5*). Instead, we have systematically knocked out these genes in *Parhyale* using

CRISPR- Cas9 mutagenesis and compared this to our understanding of their function in

Drosophila and other insects (Figs. 2, S2).

Insects have six leg segments, while *Parhyale* has seven (Fig. 1). In insects, *Dll* is

required for the development of leg segments 2 – 6 (*6-9*) . In *Parhyale*, the canonical *Dll* gene,

Dll-e (*10-12*), is required for the development of leg segments 3 – 7 (Fig. 2b). In insects, *Sp6-9*

(*13*) is required for the development of leg segments 1 – 6 (*14*), and in addition in *Drosophila*,

loss of *Sp6-9* (i.e. D-Sp1, (*13*)) occasionally transforms the leg towards wing and lateral body

wall identity (*14*). In *Parhyale*, *Sp6-9* (*13*, *15*) is required for the development of leg segments 2

– 7 (Fig. 2c), and in some legs, segment 2 is occasionally transformed towards a leg segment 1

identity (Fig S3). In *Drosophila*, *dac* is required in the trochanter through proximal tarsus (leg

segments 2 – 4, and first tarsus, (*15*, *16*). *Parhyale* has two *dac* paralogs. *dac1* does not seem to

be expressed in the legs or have a morphologically visible knockout phenotype. *dac2* is required

to pattern leg segments 3 – 5 (Fig. 2d). *exd* and *hth* are expressed in the body wall and proximal

leg segments of insects (*17-20*) and *Parhyale* (*21*) (Fig S1). They form heterodimers and

therefore have similar phenotypes (*17-20*). In insects, *exd* or *hth* knockout results in

deletions/fusions of the coxa through proximal tibia (leg segments 1 – 3, and proximal tibia, *17-*

20). In *Parhyale*, *exd* or *hth* knockout results in deletions/fusions of the coxa through proximal

carpus (leg segments 1 – 4, and proximal carpus; Figs. 2e, f). In both insects (*17, 18, 22*) and

Parhyale, the remaining distal leg segments are sometimes transformed towards a generalized

thoracic leg identity (compare Fig. 2 e, f and Fig S4). In both insects (*17-20*) and *Parhyale* (Fig.

S4), *exd* or *hth* knockout results in deletions/fusions of body segments.

In summary, the expression and function of *Dll*, *Sp6-9*, *dac*, *exd*, and *hth* in *Parhyale* are

shifted distally by one segment relative to insects. This shift is accounted for if insects fused an

ancestral proximal leg segment into the body wall (Fig. 2g). Thus, there is a one-to-one

homology between insect and *Parhyale* legs, displaced by one segment, such that the insect coxa

is homologous to the crustacean basis (*23*), the insect femur is the crustacean ischium, and so on

for all leg segments. This also means that part of the insect body wall is homologous to the

crustacean coxa.

Clark-Hachtel (accompanying manuscript) show that the plates on the *Parhyale* basis,

coxa, and lateral body wall are epipods. The body wall epipod is notable, because epipods are

characteristic of leg segments (*23*). This suggested to us that part of the *Parhyale* body wall

might actually be an additional leg segment. In fact, most groups of crustaceans have an

additional proximal leg segment, the precoxa (Fig. 3a). To determine whether *Parhyale* retains

the precoxa, we examined dissected *Parhyale* using confocal and brightfield microscopy. We

identified a precoxal structure that meets the criteria for a true leg segment: it protrudes

conspicuously from the body wall; it forms a true, muscled joint; and it extends musculature to

another leg segment (Figs. 3 and S5, (*23-25*)). Importantly, the plate does not emerge from the

body wall, but from the precoxa (Fig. 3e), like the plates of the coxa and basis. Thus, much of

what appears to be lateral body wall in *Parhyale* is in fact proximal leg.

If the insect coxa is homologous to the crustacean basis, what happened to the leg

segments corresponding to the ancestral crustacean precoxa and crustacean coxa in insects? If

these two leg segments became incorporated into the body wall, then one would expect to find

two leg segments and two epipods dorsal to the insect coxa (Fig. 4a). As predicted, two leg-like

segments can be observed proximal to the coxa in basal hexapods (*1*) including collembolans

(*26*), as well as in the embryos of many insects (*27-29*), where these two leg-like segments

flatten out before hatching to form the lateral body wall (*1, 26-31*). Insects also appear to have

two epipods dorsal to the insect coxa, because when "wing" genes are depleted in insects via

RNAi, two distinct outgrowths are affected, the wing and the plate adjacent to the leg (Fig. 1c,

(*32-35*)).

Based on these data, insects incorporated two ancestral leg segments, the precoxa and

crustacean coxa, into the body wall (Fig. 4a). Thus, like *Parhyale*, much of what appears to be

lateral body wall in insects is in fact proximal leg. Clark-Hachtel's interpretation of the dual

origin theory proposes that these two leg segments and their two epipods fused to form the wing.

While we agree that both leg segments may contribute wing muscle, we propose that only the

more dorsal precoxa epipod formed the insect wing, while the more ventral crustacean coxa

epipod formed the insect plate (Fig. 4b).

Our results may settle the long-standing debate regarding the origin of insect wings as

derived from either an epipod of the leg or from body wall. Our model accounts for all

observations in favor of either of these hypotheses, including the dorsal position of insect wings

relative to their legs, the loss of ancestral leg segments in insects, the two-segmented

morphology of the insect subcoxa in both embryos and adults, the complex musculature for

flight, and the shared gene expression between wings and epipods. Our model also explains the

apparent "dual origin" of insect wings from both body wall and leg epipod: much of what

appears to be insect body wall is in fact the remnant of two ancestral leg segments and their

epipods.

In fact, a number leg-associated outgrowths in arthropods could be explained by this

model, in addition to insect wings. The Daphnia carapace(*36*) is the epipod of the precoxa(*37*);

the *Oncopeltus* small plate outgrowth (Fig. 1c) is the epipod of the crustacean coxa; and the

thoracic stylus of jumping bristletails (Fig. 4, st) is the epipod of the crustacean basis(*38*, *39*).

This also explains many insect abdominal appendages, like gills(*40*), gin traps(*33*), prolegs(*41*),

and sepsid fly appendages(*42*), which are often proposed as de novo structures(*43-45*). However,

most insects form abdominal appendages as embryos(*40*, *46*), some even with an epipod nub, but

these fuse to the body wall before hatching to form the sternites(*28*, *39*). The existence of insect

abdominal appendages is supported by a re-analysis of the expression of *Sp6-9* and its paralog,

buttonhead, in insect embryos in a previous study (*13*). According to the leg segment homology

model presented here (Fig. 4), the paired dots of *btd* expression on each abdominal segment of

insect embryos demonstrates that these appendages are comprised of three leg segments: the

precoxa (pink), crustacean coxa (red), and insect coxa (orange). These abdominal appendages are

truncated, lacking all distal appendages from the trochanter (yellow) down, because *Dll* and *dac*,

which mark the trochanter and more distal leg segments, are not expressed in the insect

abdomen. Thus, rather than de novo co-options, abdominal appendages were always there,

persisting in a truncated, highly modified state, and de-repressed in various lineages to form

apparently novel structures. This provides a model for how insect wings can be both homologous

to the epipod of the crustacean precoxa, and yet not be continuously present in the fossil record:

epipod fields may persist in a truncated state, perhaps only visible as a nub in the embryo. We

therefore propose cryptic persistence via truncation as a general mechanism for the origin of

apparently novel structures that appear to be derived from serial homologs, rather than the

current model of extensive gene co-option.

References and Notes:

1.   R. E. Snodgrass, Morphology and mechanism of the insect thorax. *Smithsonian*
Miscellaneous Collections. **80** (1927).

2.   J. Kukalová-Peck, Origin of the insect wing and wing articulation from the arthropodan
leg. *Can. J. Zool.* **61**, 1618–1669 (1983).

3.   R. E. Snodgrass, *Principles of insect morphology*. (McGraw-Hill Book Company, Inc,
NEW YORK, First edition second impression. 1935).

4.   C. M. Clark-Hachtel, Y. Tomoyasu, Exploring the origin of insect wings from an evo-
devo perspective. *Curr Opin Insect Sci*. **13**, 77–85 (2016).

5.   C. Estella, A dynamic network of morphogens and transcription factors patterns the fly
leg. *Curr. Top. Dev. Biol.* **98**, 173–198 (2012).

6.   C. Estella, G. Rieckhof, M. Calleja, G. Morata, The role of buttonhead and Sp1 in the
development of the ventral imaginal discs of Drosophila. *Development*. **130**, 5929–5941
(2003).

7.   G. Campbell, A. Tomlinson, The roles of the homeobox genes aristaless and Distal-less in
patterning the legs and wings of Drosophila. *Development*. **125**, 4483–4493 (1998).

8.   D. R. Angelini, T. C. Kaufman, Functional analyses in the hemipteran Oncopeltus
fasciatus reveal conserved and derived aspects of appendage patterning in insects.
Developmental Biology. **271**, 306–321 (2004).

9.   A. Beermann *et al.*, The Short antennae gene of Tribolium is required for limb
development and encodes the orthologue of the Drosophila Distal-less protein.
Development. **128**, 287–297 (2001).

10.   J. M. Serano *et al.*, Comprehensive analysis of Hox gene expression in the amphipod
crustacean Parhyale hawaiensis. *Developmental Biology*. **409**, 297–309 (2015).

11.   W. E. Browne, A. L. Price, M. Gerberding, N. H. Patel, Stages of embryonic development
in the amphipod crustacean, Parhyale hawaiensis. *genesis*. **42**, 124–149 (2005).

12.   D. M. Liubicich *et al.*, Knockdown of Parhyale Ultrabithorax recapitulates evolutionary
changes in crustacean appendage morphology. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 13892–
13896 (2009).

13.   N. D. Schaeper, N.-M. Prpic, E. A. Wimmer, A clustered set of three Sp-family genes is
ancestral in the Metazoa: evidence from sequence analysis, protein domain structure,
developmental expression patterns and chromosomal location. *BMC Evol. Biol.* **10**, 88
(2010).

14.   C. Estella, R. S. Mann, Non-Redundant Selector and Growth-Promoting Functions of Two
162   Sister Genes, buttonhead and Sp1, in Drosophila Leg Development. *PLoS Genet*. **6**,
163   e1001001 (2010).

164   15.   G. Mardon, N. M. Solomon, G. M. Rubin, dachshund encodes a nuclear protein required
165   for normal eye and leg development in Drosophila. *Development*. **120**, 3473–3486 (1994).

166   16.   B. C. Tavsanli *et al.*, Structure–function analysis of the Drosophila retinal determination
167   protein Dachshund. *Developmental Biology*. **272**, 231–247 (2004).

168   17.   T. Mito *et al.*, Divergent and conserved roles of extradenticle in body segmentation and
169   appendage formation, respectively, in the cricket Gryllus bimaculatus. *Developmental*
170   *Biology*. **313**, 67–79 (2008).

171   18.   M. Ronco *et al.*, Antenna and all gnathal appendages are similarly transformed by
172   homothorax knock-down in the cricket Gryllus bimaculatus. *Developmental Biology*. **313**,
173   80–92 (2008).

174   19.   C. Rauskolb, K. M. Smith, M. Peifer, E. Wieschaus, extradenticle determines segmental
175   identities throughout Drosophila development. *Development*. **121**, 3663–3673 (1995).

176   20.   J. Wu, S. M. Cohen, Proximodistal axis formation in the Drosophila leg: subdivision into
177   proximal and distal domains by Homothorax and Distal-less. *Development*. **126**, 109–117
178   (1999).

179   21.   N.-M. Prpic, M. J. Telford, Expression of homothorax and extradenticle mRNA in the legs
180   of the crustacean Parhyale hawaiensis: evidence for a reversal of gene expression
181   regulation in the pancrustacean lineage. *Dev Genes Evol*. **218**, 333–339 (2008).

182   22.   F. Casares, R. S. Mann, The ground state of the ventral appendage in Drosophila. *Science*.
183   **293**, 1477–1480 (2001).

184   23.   G. A. Boxshall, The evolution of arthropod limbs. *Biol. Rev.* **79**, 253–300 (2004).

185   24.   G. Boxshall, *Arthropod Limbs and their Development* (2013).

186   25.   J. W. Shultz, Morphology of locomotor appendages in Arachnida: evolutionary trends and
187   phylogenetic implications. *Zool J Linn Soc*. **97:1-56** (1989).

188   26.   H. Bäcker, M. Fanenbruck, J. W. Wägele, A forgotten homology supporting the
189   monophyly of Tracheata: The subcoxa of insects and myriapods re-visited. *Zoologischer*
190   *Anzeiger - A Journal of Comparative Zoology*. **247**, 185–207 (2008).

191   27.   Y. Kobayashi, Formation of Subcoxae-1 and 2 in Insect Embryos: The Subcoxal Theory
192   Revisited. *Proc Arthropod Embryol Soc Jpn*. **48**, 33–38 (2017).

28. Y. Kobayashi, K. Niikura, Y. Oosawa, Y. Takami, Embryonic development of Carabus
insulicola (Insecta, Coleoptera, Carabidae) with special reference to external morphology
and tangible evidence for the subcoxal theory. *J. Morphol.* **274**, 1323–1352 (2013).

29. Y. Mashimo, R. Machida, Embryological evidence substantiates the subcoxal theory on
the origin of pleuron in insects. *Sci Rep.* **7**, 1–9 (2017).

30. H. J. Hansen, *Studies on the Arthropoda III. On the Comparative Morphology of the*
Appendages in the Arthropoda. B. Crustacea (Supplement), Insecta, Myriapoda and …
(Copenhagen: Gyldendalske Boghandel, 1930).

31. T. Deuve, The epipleural field in hexapods. *Annales De La Societe Entomologique De*
France. **37**, 195–231 (2001).

32. C. M. Clark-Hachtel, D. M. Linz, Y. Tomoyasu, Insights into insect wing origin provided
by functional analysis of vestigial in the red flour beetle, Tribolium castaneum. *Proc. Natl.*
Acad. Sci. U.S.A. **110**, 16951–16956 (2013).

33. T. Ohde, T. Yaginuma, T. Niimi, Insect morphological diversification through the
modification of wing serial homologs. *Science*. **340**, 495–498 (2013).

34. V. Medved *et al.*, Origin and diversification of wings: Insights from a neopteran insect.
Proc. Natl. Acad. Sci. U.S.A. **112**, 15946–15951 (2015).

35. D. Wang *et al.*, spalt is functionally conserved in Locusta and Drosophila to promote wing
growth. *Sci Rep.* **7**, 1–9 (2017).

36. Y. Shiga *et al.*, Repeated co-option of a conserved gene regulatory module underpins the
evolution of the crustacean carapace, insect wings and other flat outgrowths,
doi:10.1101/160010.

37. H. J. Hansen, *Studies on Arthropoda. II. At the Expense of the Rask-Ørsted Fund*
(Copenhagen: Gyldendalske Boghandel, 1925).

38. N. Niwa *et al.*, Evolutionary origin of the insect wing via integration of two
developmental modules. *Evol. Dev.* **12**, 168–176 (2010).

39. J. Bitsch, The controversial origin of the abdominal appendage-like processes in immature
insects: Are they true segmental appendages or secondary outgrowths? (Arthropoda
hexapoda). *J. Morphol.* **273**, 919–931 (2012).

40. S. Komatsu, Y. Kobayashi, Embryonic development of a whirligig beetle, Dineutus
mellyi, with special reference to external morphology (insecta: Coleoptera, Gyrinidae). *J.*
Morphol. **273**, 541–560 (2012).

41. Y. Suzuki, D. C. Squires, L. M. Riddiford, Developmental Biology. *Developmental*
Biology. **326**, 60–67 (2009).

42.   J. H. Bowsher, H. F. Nijhout, Partial co-option of the appendage patterning pathway in the
228   development of abdominal appendages in the sepsid fly Themira biloba. *Dev Genes Evol.*
229   **219**, 577–587 (2010).

230   43.   H. Hoch *et al.*, Non-sexual abdominal appendages in adult insects challenge a 300 million
231   year old bauplan. *Curr. Biol.* **24**, R16–7 (2014).

232   44.   D. R. Angelini, T. C. Kaufman, Comparative developmental genetics and the evolution of
233   arthropod body plans. *Annu. Rev. Genet.* **39**, 95–119 (2005).

234   45.   A. P. Moczek, On the origins of novelty in development and evolution. *Bioessays.* **30**,
235   432–447 (2008).

236   46.   R. Matsuda, *Morphology and Evolution of the Insect Abdomen* (Elsevier, 1976).

237   47.   R. R. Hessler, The Structural Morphology of Walking Mechanisms in Eumalacostracan
238   Crustaceans. *Philosophical Transactions of the Royal Society B: Biological Sciences.* **296**,
239   245–298 (1982).

Acknowledgments: We thank Courtney Clark-Hachtel and Yoshinori Tomoyasu for sharing

their preprint. **Funding:** This work is supported by the National Science Foundation (NSF) (to

NHP), NSF Graduate Research Fellowship (to HSB). **Author contributions:** H.S.B. and N.H.P.

conceived of the experiments. H.S.B. performed the experiments, conceived of model, and wrote

the manuscript. N.H.P. edited and revised the manuscript. **Competing interests**: Authors declare

no competing interests. **Data and materials availability:** All data is available in the main text or

the supplementary materials.

Supplementary Materials:

Materials and Methods

Figures S1-S5

Table S1

References 37-43

Fig. 1. Crustacean and insect legs. (a) Adult *Parhyale*, with third thoracic leg (T3) outlined. (b)

Cartoon of *Parhyale* T3. The coxal plate extends over the leg. (c) Adult *Oncopeltus*, with T2

outlined. Inset shows magnified proximal leg, with body wall plate extending over the leg. (d)

Cartoon of *Oncopeltus* T2 leg.

Fig. 2. Knockout phenotypes of leg gap genes. (a-f) *Parhyale* CRISPR-Cas9 phenotypes in
dissected third thoracic legs (T3). Graded cyan in f indicates deletion/fusion of proximal leg
segment 5. (g) Leg gap gene function in *Parhyale* and insects aligns only if insects incorporated
the red leg segment into the body wall (0). Color bars correspond to remaining leg segments
following knockout, transparent bars indicate deleted leg segments. Open bar in *dac* indicates
slight extension of *dac* function into tarsus 1 of insects. Coxal plate (Cp), gill (G), tergal plate
(Tp). Scale bar 50um.

Fig. 3. Evidence for a precoxa in *Parhyale*. (a) Phylogeny based on Oakley 2012, precoxa

references in supplements. (b) Confocal image of *Parhyale* hatchling, autofluorescent cuticle in

blue. T5, T6 tergal plates (dotted outlines). (c) Confocal image of *Parhyale* hatchling,

autofluorescent cuticle in blue, muscle phalloidin stain in red. Compare blocks of simple,

anterior-posterior muscles of the body to orthogonal, complexly arranged muscles of the leg

segments. Note overlap between tergal plate (dotted lines) and orthogonal leg muscle. (d)

Brightfield image of right half of *Parhyale,* sagittal dissection, internal tissues removed, lateral

view. Wire used to position sample (w). The same orthogonal muscles in b are visible as

striations extending above the wire. The precoxa forms a joint with the coxa (*47*) (arrow). The

dorsal limit of the precoxa is unclear: a conservative estimate is to begin at the joint (arrow) and

follow the leg up to where it meets the adjacent leg, denoted by (<); however, the orthogonal

muscle striations continue farther up (pink outline). Either way, the precoxa protrudes quite a bit

from the body wall. (e) Posterior-lateral view of right T6, looking edge-on at tergal plate. The

tergal plate (dotted outline) emerges from the precoxa (contiguous pink between ←, >, and ---),

just as the coxal plate (dashed line) emerges from the coxa. In c, d, coxa is red (coxal plate not

shaded), gills (teal) partially cut for visibility, basis orange, precoxa pink. Note that all three

plates (tergal, coxal, and basal) form contiguous cuticle with their leg segment, i.e. there is no

distinguishing suture. Scale bar 100um.

Fig. 4. Proposed leg segment homologies (colors) between insects, *Parhyale*, and an ancestral

crustacean (a) based on gene function alignment (b). Ancestral precoxa epipod (pink ep),

Parhyale tergal plate (Tp), and insect wing are homologous (pink). Ancestral coxa epipod,

Parhyale coxal plate (Cp) and gill (G), and insect plate (see Fig. 1c) are homologous (red).

Parhyale basal plate (Bp). Insect numbering based on crustaceans.