1

# Expression of the Eight GABA$_A$ Receptor α Subunits in the Developing Zebrafish Central Nervous System

Bryan Monesson-Olson[1,2], Jon J. McClain[2], Abigail E. Case[2], Hanna E. Dorman[2], Daniel R.

Turkewitz[3], Aaron B. Steiner[3] and Gerald B. Downes[1,2]*

[1]Neuroscience and Behavior Graduate Program

[2]Biology Department, University of Massachusetts, Amherst, MA 01003, USA

[3]Department of Biology and Health Sciences, Pace University, Pleasantville, NY 10570, USA

*Corresponding author:

Gerald B. Downes

Email: gbdownes@bio.umass.edu

# ABSTRACT

GABA is a robust regulator of both developing and mature neural networks. It exerts many of its effects through $GABA_A$ receptors, which are heteropentamers assembled from a large array of subunits encoded by distinct genes. In mammals, there are 19 different $GABA_A$ subunit types, which are divided into the α, β, γ, δ, ε, π, θ and ρ subfamilies. The immense diversity of $GABA_A$ receptors is not fully understood. However, it is known that specific isoforms, with their distinct biophysical properties and expression profiles, tune responses to GABA. Although larval zebrafish are well-established as a model system for neural circuit analysis, little is known about $GABA_A$ receptors diversity and expression in this system. Here, using database analysis, we show that the zebrafish genome contains at least 23 subunits. All but the mammalian θ and ε subunits have at least one zebrafish ortholog, while five mammalian $GABA_A$ subunits have two zebrafish orthologs. Zebrafish contain one subunit, β4, which does not have a clear mammalian ortholog. Similar to mammalian $GABA_A$ receptors, the zebrafish α subfamily is the largest and most diverse of the subfamilies. In zebrafish there are eight α subunits, and RNA *in situ* hybridization across early zebrafish development revealed that they demonstrate distinct patterns of expression in the brain, spinal cord, and retina. Some subunits were very broadly distributed, whereas others were restricted to small populations of cells. Subunit-specific expression patterns were similar between zebrafish, frogs and rodents, which suggests that the roles of different $GABA_A$ receptor isoforms are largely conserved among vertebrates. This study provides a platform to examine isoform specific roles of $GABA_A$ receptors within zebrafish neural circuits and it highlights the potential of this system to better understand the remarkable heterogeneity of $GABA_A$ receptors.

# INTRODUCTION

Neural networks throughout the central nervous system rely upon a diversity of

neurotransmitter systems for both their initial formation and mature function. GABA is the major

inhibitory neurotransmitter throughout most of the mature nervous system [1]. It exerts its effects

through its receptors, which are divided into two classes. GABA$_A$ receptors are ligand-gated

chloride channels responsible for most of the rapid effects of GABA, while GABA$_B$ receptors are

heterotrimeric G-protein coupled receptors. In mammalian systems, GABA$_A$ receptors

demonstrate incredible subtype diversity and are targets for classes of clinically important drugs,

such as benzodiazepines and barbituates [2, 3]. GABA$_A$ receptors are heteropentamers

composed of various combinations of 19 different subunits: α1-6, β1-3, γ1-3, δ, ε, π, θ and ρ1-3

(previously referred to as GABA$_C$ receptors) [4, 5]. Each subunit is encoded by a discrete gene

that is spatially and developmentally regulated to generate distinct, but often overlapping,

expression patterns [6-9]. Alternative splicing and RNA editing of some subunits further

increases the number of subtypes available [10]. Although this extensive receptor heterogeneity

is not fully understood, some subunits confer distinct biophysical and pharmacological

properties, interact with specific cytoplasmic proteins, and localize to specific subcellular

domains [11-13]. Ultimately, this receptor diversity provides a capacity to tailor responses to

GABA within neural circuits.

The α subunits play a prominent role in GABA$_A$ receptor function. They are thought to be

essential components of 'classic' GABA$_A$ receptors, which exclude the pharmacologically

distinct receptors composed from ρ subunits [3]. Exogenous expression studies and analysis of

native receptors indicate that each receptor contains two α subunits which, along with β

subunits, form the GABA binding site. α subunits can also dictate isoform-selective

pharmacology. For example, receptors containing α1, α2, α3 and α5 subunits confer

benzodiazepine-sensitivity, whereas receptors that contain α4 and α6 subunits do not bind to

clinically used benzodiazepines [2]. In addition, α subunits regulate subcellular localization.

GABA receptors that contain α1, α2, and α3 demonstrate a synaptic localization and mediate

transient or phasic inhibition, while receptors that contain α4, α5 and α6 are principally

extrasynaptic and mediate tonic inhibition [14, 15]. Through both transient and tonic inhibition,

these receptors mediate the majority of inhibition in neural circuits throughout the vertebrate

brain.

75            Developing zebrafish have several features that make them well-suited to study the

formation and function of neural circuits [16, 17]. First, zebrafish embryos and larvae develop

external to the mother and are optically transparent. These features make the central nervous

system easily accessible throughout development and particularly amenable to optical

physiological approaches, such as optogenetics or calcium imaging. Second, the central

nervous system contains fewer cells compared to mammalian preparations, yet many cell types

and mechanisms are conserved among vertebrates. Lastly, larval zebrafish develop rapidly,

therefore many sensory and motor systems are present and functional within five days post-

fertilization. Combined, these features have helped establish developing zebrafish as a

prominent system to examine neural circuit formation and function.

85            GABA$_A$ receptors are expressed in larval zebrafish and essential for normal nervous

system function. For example, bath application of pentylenetetrazole, a convulsant and

noncompetitive GABA$_A$ receptor antagonist, induces hyperactive behavior that can serve as a

model of seizures [18]. Similarly, insecticides known to target GABA$_A$ receptors generate

hyperactive behavior in larvae [19]. Pharmacological blockade of GABA$_A$ receptors has also

been shown to regulate the activity of specific cells, including retinal bipolar cells [20], optic

tectum neurons [21], Mitral cells in the olfactory bulb [22], spinal cord CoPA interneurons and

motoneurons [23], and the Mauthner cells, a pair of well-studied reticulospinal neurons found in

the amphibian and teleost hindbrain [24]. In these studies, the GABA$_A$ receptor isoforms were

not identified. However, patch-clamp recordings of the Mauthner cells showed three distinct

GABA$_A$ kinetic profiles, which were proposed to be caused by different receptor isoforms [25].

There is limited information about the heterogeneity of GABA$_A$ receptors in zebrafish.

The most extensive study to date identified 23 GABA$_A$ receptor subunits and examined their

expression broadly in adult zebrafish brain via RT-qPCR [26]. Despite playing prominent roles in

developing animals, little is known about the expression of GABA$_A$ receptor subunits in

zebrafish embryos and larvae. α1 has been shown to be expressed at 50 and 72 hours post-

fertilization (hpf), γ2 at 50 hpf, and α2a (then named α2) from ~14 – 96 hpf. [27-29].

Unpublished α6a expression data has also been deposited in the ZFIN database [30]. The

expression patterns of other GABA$_A$ receptor subunits has not been reported.

In this study we performed phylogenetic analysis, which indicates that zebrafish contain

at least 23 different GABA$_A$ receptor subunits. Although we observed some differences between

the zebrafish and mammalian GABA$_A$ receptor subunit gene families, zebrafish contain

orthologs for most of the GABA$_A$ receptor subunits found in mammals. To examine where and

when these GABA$_A$ receptors are expressed in the zebrafish nervous system, we performed

whole-mount RNA *in situ* hybridization. We present the embryonic and larval expression of the

eight identified α subunit-encoding genes: *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, *gabra5*,

*gabra6a* and *gabra6b*. These data show that each subunit has a distinct expression pattern,

broadly similar to the reported expression of amphibian and rodent orthologs. Combined, these

results suggest that GABA$_A$ isoform-specific roles are conserved among vertebrates, and they

establish a foundation to use the zebrafish system to better understand how GABA$_A$ receptor

heterogeneity mediates neural circuit formation and function.

# MATERIALS AND METHODS

## Phylogenetic analysis

Homologous gene queries were performed using the National Center for Biotechnology Information tBLASTn search tool. Protein sequences for each of the 19 mouse GABA$_A$ receptor subunits were used to search the zebrafish nucleotide collection. Matching zebrafish sequences were evaluated to determine if they were splice variants from the same gene or generated from different genes. When splice variants were identified, only the longest variant was used for subsequent analysis. 19 mouse and 23 zebrafish protein sequences were used to assemble a phylogenetic tree. The genes and Genbank accession numbers for the mouse sequences are: *Gabra1*- NP_034380, *Gabra2*- NP_032092, *Gabra3*- NP_032093, *Gabra4*- NP_034381, *Gabra5*- NP_795916*, Gabra6*- NP_00109311, *Gabrb1*- NP_032095, *Gabrb2*- NP_032096, *Gabrb3*- NP_032097, *Gabrd*- NP_032098, *Gabre*- NP_059065, *Gabrg1*- NP_034382, Gabrg2- NP_032099, *Gabrg3*- NP_032100, *Gaprp*- NP_666129, *Gabrq*- NP_065234, *Gabrr1*- NP_032101, *Gabrr2*- NP_032102, and *Gabrr3*- NP_001074659. The Genbank accession numbers for the zebrafish sequences are: *gabra1*- NM_001077326, *gabra2a*- XM_009307207, gabra2b- XM_017359049, *gabra3*- XM_009295708, *gabra4*- NM_001017822, *gabra5*- XM_001339475, *gabra6a*- XM_005173112, *gabra6b*- XM_002667357, *gabrb1*- XM_002664133, *gabrb2*- NM_001024387, *gabrb3*- XM_005166079, *gabrb4*- XM_005173874, *gabrd*- XM_695007, *gabrg1*- XM009307208, *gabrg2*- NM_001256250, *gabrg3*- XM_009302568, *gabrp*- XM_005173293, *gabrr1*- NM_001025553, *gabrr2a*- NM_001045376, *gabrr2b*- XM_009294512, *gabrr3a*- NM_001128760, *gabrr3b*- XM_009297450, *gabrz*- XM_005156247. To construct the tree, a ClustalW alignment of amino acid sequences was performed using the Geneious software package [31]. An unrooted PHYML consensus tree was then generated using 100 bootstrap replicates [32].

## Fish maintenance and breeding

Zebrafish were raised and maintained using established husbandry procedures.

Embryos were kept at 28.5 °C in E3 media and staged according to morphological criteria [33].

All experiments were performed using Tuebingen (Tu) or tub longfin (TLF) wild type embryos.

From 0-24 hpf, embryos were grown in 0.01% Methylene Blue in E3 medium. At 24 hpf the

solution was changed to 0.0045% Phenylthiourea (PTU) in E3 to inhibit pigment formation. This

solution was changed every 24 hours until the fish were sacrificed. All animal protocols were

approved by the University of Massachusetts Institutional Animal Care and Use Committees

(IACUC).

## Whole-mount in situ hybridization

Antisense digoxigenin probes were generated against *gabra1*, *gabra2a*, *gabra2b*,

*gabra3*, *gabra4*, *gabra5*, *gabra6a* and *gabra6b* (S1 Table). *In situ* probe synthesis utilized the

digoxigenin RNA labeling kit with SP6 or T7 RNA polymerases (Roche Diagnostics, Mannheim,

Germany). Whole-mount, colorimetric *in situ* hybridization was performed using established

protocols [34] and examined using a compound microscope (Zeiss, Thornwood, NY) attached to

a digital camera (Zeiss, Thornwood, NY). Cross sections were generated by imbedding *in situ*

hybridization stained embryos in 1.5% agar, 5% sucrose. The blocks were submerged in a 30%

sucrose solution overnight then cut into 20mm thick sections using a cryostat (Leica, Buffalo

Grove, IL). Representative larvae at 24, 48, and 96 hpf were mounted in either 100% glycerol or

dehydrated through a methanol series, equilibrated in a 2:1 benzylbenzoate/benzylalchol

solution and mounted in a 10:1 Canada balsam/methyl saliylate misture. Images were captured

using a Zeiss Discovery v12 stereomicroscope with a 1.5x objective or a Zeiss Axioskop

Microscope with a 10x objective and an AxioCam MRc camera. All images were processed

using Adobe Photoshop, in which multiple focal planes were merged to produce single

representative images. The identify of neuroanatomical structures was determined using

zebrafish brain atlases [35-37].

## Reverse-Transcriptase PCR

RT-PCR was used to analyze whether the *gabra3* and *gabra3*-like sequence were

portions of the same transcript. Primers designed against *gabra3*-like (Primer 1 5'

GGACGGCGGATGATGAGAAA-3') and *gabra3* (Primer 2 5'-CACGACCGTCCTGACTA-3',

Primer 3 5'-GTGGAGTAGATGTGGTGGGC-3') were used to amplify cDNA from wild-type

zebrafish larvae. RNA was extracted from 48 hpf zebrafish larvae using the RNAeasy kit

(Qiagen, Venlo, Netherlands) and reverse transcribed using the Accuscript RT-PCR system

(Stratagene). The PCR products were sequenced to confirm that they are portions from the

same transcript.

# RESULTS

## The zebrafish GABA$_A$ receptor subunit gene family is similar in size and diversity to the mammalian GABA$_A$ receptor gene family.

19 GABA$_A$ receptor subunit genes have been identified in humans, mice, and rats [4, 5].

To establish the number and organization of zebrafish GABA$_A$ receptor subunits, we used

mouse GABA$_A$ receptor subunit amino acid sequences to query zebrafish genome databases.

We identified 23 zebrafish GABA$_A$ receptor subunits, each encoded by a distinct gene.

Frequently splice variants were observed. In these cases, the principle splice isoform, as

indicated by the databases, was selected for analysis. 12 of the 19 mouse GABA$_A$ receptor

subunits were found to have a single ortholog in zebrafish (Fig 1). Amino acid identities between

mouse and zebrafish orthologs ranged from 53.1 to 86.3%. Five additional mouse subunits, α2,

α6, ρ2, ρ3, and π, each exhibit similarity with two subunits in zebrafish: *gabra2a* and *gabra2b*,

*gabra6a* and *gabra6b*, *gabrr2a* and *gabrr2b*, *gabrr3a* and *gabrr3b*, *gabrp* and *gabrz*,

respectively. The relatively high percentage of amino acid identity of these zebrafish subunits

with each other suggests that the zebrafish paralogs are duplicated genes. Duplicated genes

are often observed in zebrafish, and they are thought to be due to a whole genome duplication

within the teleost lineage [38, 39]. It was reported previously that there is a *gabra3* and an α3-

like gene. While the α3-like gene has been localized to chromosome 21, the genomic location of

the α3-like gene is not known. Our database analysis suggested that these sequences could be

non-overlapping portions of the same gene, therefore RT-PCR was performed using primers

targeting α3-like and *gabra3* sequences (S1 Fig). Transcripts were amplified that spanned the

region between *gabra3* and α3-like sequences, indicating that these are different portions of the

same gene and that there is only one α3 encoding gene, *gabra3*.

**Fig 1. Phylogenetic analysis shows that the zebrafish GABA$_A$ subunit gene family is**

**similar in size, diversity, and organization to the mouse GABA$_A$ subunit gene family.**

Amino acid sequence alignments were used to generate a consensus tree using 100 bootstrap

replicates. The genes that encode the GABA$_A$ subunits are shown at the tip of each branch and

bootstrap proportions are shown at the branch points. The zebrafish and mouse GABA$_A$ subunit

sequences showed high amino acid identity and grouped into the α, β, γ, δ, π, and ρ

subfamilies. Most mouse GABA$_A$ subunits have a single zebrafish ortholog, while six mouse

GABA$_A$ subunits have two zebrafish orthologs, likely due to gene duplication.

There are two mouse GABA$_A$ receptor subunits for which zebrafish orthologous are less

clear or have yet to be identified, the θ and ε subunits. The θ subunit, encoded by the gene

*Gabrq*, exhibits only 31.3% amino acid identity with zebrafish β4, encoded by *gabrb4*, which is

the greatest percentage identity of this subunit with any zebrafish sequence. There are three β

subunits in mice compared to the four β subunits in zebrafish so it is possible that, despite the

low amino acid sequence identity, zebrafish β4 is orthologous to the mouse θ subunit. No clear

zebrafish ortholog has yet to be identified for the mouse ε subunit, which is encoded by the

*Gabre* gene.

## α subunits demonstrate distinct expression patterns across early zebrafish development

Determining the temporal and spatial expression of GABA$_A$ subunits is instrumental to

identify isoform-specific roles in zebrafish neural circuit development and function. Therefore,

given the large size of the α subunit subfamily and their prominent role in GABA$_A$ receptor

function, we examined the expression of the eight α subunit encoding genes: *gabra1*, *gabra2a*,

*gabra2b*, *gabra3*, *gabra4*, *gabra6a* and *gabra6b*. Using whole-mount RNA *in situ* hybridization,

we determined their expression patterns at 24, 48, and 96 hpf. These time points span much of

larval zebrafish locomotor network development [40].

Expression of *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4* and *gabra5* was detected at 24

hpf (Fig 2). Neither *gabra6a* or *gabra6b* were detected at this developmental stage. *gabra2a*

and *gabra2b* were detected broadly and did not appear to be spatially restricted (Fig 2C-F). To

distinguish whether these gene transcripts were widespread or background due to the probe, a

second probe was generated for each gene (S1 Table). These probes yielded widespread

staining, very similar to the initial probes used, which suggests that *gabra2a* and *gabra2b* are

widely distributed. Consistent with these findings, a previous study reported that *gabra2a*

exhibits a diffuse, broad pattern of expression in larval zebrafish [28].

**Fig 2. Expression of GABA$_A$ α subunits at 24 hpf.** *gabra1* (A, B), *gabra2a* (C, D), *gabra2b*

(E, F), *gabra3* (G, H), *gabra4* (I, J), and *gabra5* (K, L) were detected at this time point. Whole

mount lateral (A, C, E, G, I, K) views are shown along with dorsal views of the head (B, D, F, H,

J, L).The scale bar (A) is 0.1 mm. The brackets in G, I, and K indicates the regions shown at

higher magnification in the corresponding insets. Abbreviations: drc, dorsal rostral cluster; hbc,

hindbrain cluster; op, olfactory placode; SC, spinal cord; t, tegmentum; vcc, ventral caudal

cluster; vrc, ventral rostral cluster.

*gabra1*, *gabra3*, *gabra4*, and *gabra5* were each expressed in discrete cells in the brain

or spinal cord at 24 hpf. *gabra1* was detected in the olfactory placodes, the ventral rostral

cluster, the ventral caudal cluster and small clusters of cells within each rhombomere of the

hindbrain (Fig 2A, B). Although expressed in discrete cells, *gabra3* was widely expressed, and

found within the tegmentum, hindbrain, and ventral and intermediate domains in the spinal cord

(Fig 2G, H). *gabra4* was expressed in a diffuse manner throughout the brain but specifically

within a select population of ventral spinal cord cells (Fig 2I, J). The location and distribution of

these cells suggests that *gabra4* is expressed selectively within Kolmer-Agduhr (KA) neurons.

Lastly, *gabra5* is prominently expressed in small groups of cells in each rhombomere of the

hindbrain and in an intermediate domain of the spinal cord (Fig 2K, L). The location and

distribution of these spinal cord cells suggests that they are CoPA neurons.

The expression of all GABA$_A$ receptor subunit genes, except for *gabra6a*, was observed

at 48 hpf (Fig 3). Similar to the pattern observed at 24 hpf, the *gabra2* paralogs were again

detected broadly, with little spatial restriction (Fig 3C-F). Similar to its expression at 24hpf,

*gabra1* is prominently expressed in the olfactory bulbs, the pallium and discrete clusters of cells

in medial portions of the medulla (Fig 3A, B). *gabra3* was observed in relatively small clusters of

cells in the pallium, thalamus, and the medulla (Fig 3G, H). *gabra4* transcripts were identified in

distinct cells in the subpallium and lateral potions of the medulla. *gabra5* transcripts were

detected most prominently in the medulla. The location and expression this subunit suggests it

exhibits robust expression in the Mauthner cells, therefore α5 containing receptors are likely

some of the GABA$_A$ isoforms identified previously via electrophysiology [25]. *gabra6b* is

expressed prominently in the olfactory bulbs.

**Fig 3. Expression of the GABA$_A$ α subunits in the brain at 48hpf.** *gabra1* (A, B), *gabra2a* (C,

D), *gabra2b* (E, F) *gabra3* (G, H) *gabra4* ( I, J) *gabra5* (K, L) and *gabra6b* (M, N) were detected

at this time point. Lateral (A, C, E, G, I, K, M) and dorsal (B, D, F, H, J, L, N) views are shown.

The scale bar (A) is 0.1 mm. Brackets in A, G, I, and K indicate the region shown in cross

sections within the insets. Abbreviations: cb, cerebellum; di, diencephalon; hth, hypothalamus;

M, Mauthner cell; mo, medulla oblongata; ob, olfactory bulb; p, pallium; sp, subpallium; t,

tegmentum; th, thalamus.

Although *gabra1*, *gabra3*, *gabra4*, and *gabra5* transcripts were all detected in the

medulla, they do not appear to be in the same cells. Instead, an intriguing medial to lateral

organization was observed. *gabra1* is expressed strongly in the most medial cells, *gabra3* and

*gabra5* transcripts were found in an intermediate domain, and *gabra4* was observed laterally

(compare insets in Fig A, G, I, K).

Transcripts for all α subunits are detected in larval zebrafish at 96 hpf (Fig 4). As with the

earlier developmental time points, *gabra2a* and *gabra2b* transcripts were detected broadly,

although *gabra2b* appears less diffuse and more discrete compared to at 24 and 48 hpf (Fig 4C-

F). *gabra1*, *gabra3*, and *gabra6b* are also expressed more broadly compared to earlier stages.

In contrast, *gabra4*, *gabra5* and *gabra6a* transcripts were detected in smaller groups of cells.

*gabra4* is expressed in a stripped pattern in the outer nuclear layer of the retina, the posterior

tuberculum area, which is a portion of the diencephalon, and the tectum, cerebellum, and

medulla (Fig I, J). *gabra5* is expressed in discrete cells in the pallium, hypothalamus, cerebellum

and medulla (Fig K, L). Within the medulla, *gabra5* expression in the Mauthner Cells is robust,

as it is at earlier developmental stages.

**Fig 4. Expression of the GABA$_A$ α subunits in the brain at 96hpf.** Transcripts encoding all α

subunits were detected at this time point. Lateral (A, C, E, G, I, K, M, O) and dorsal (B, D, F, H,

J, L, N, P) are shown The scale bar (A) is 0.1 mm. Abbreviations: cb, cerebellum; gcl, ganglion

cell layer; hth, hypothalamus; inl, inner nuclear layer;M, mauthner cell; mo, medulla oblongata;

ob, olfactory bulb; opn, optic nerve; p, pallium; pcl, photoreceptive cell layer; po, preoptic region;

poc, post optic commissure; pta, posterior tubercular area; sp, subpallium; t, tegmentum; te,

telencephalon; th, thalamus; to, tectum opticum;

Transcripts for several α subunits are expressed in the retina at 96 hpf, however most

are restricted to one or two cell layers (Fig 4). *gabra1* and *gabra6b* are both expressed in the

ganglion cell and outer nuclear layers, *gabra2a* and *gabra2b* both seem to be expressed

broadly, and *gabra3* and *gabra4* transcripts both are restricted to the outer nuclear layer.

*gabra6a* is expressed prominently in the photoreceptor cell layer, and does not demonstrate

robust expression outside of the retina.

# DISCUSSION

In this study, we showed that the GABA$_A$ receptor gene family exhibits a size and diversity very similar to those found in mammals. We identified sequences for 23 GABA$_A$ receptor encoding genes, and phylogenetic analysis indicates that isoform are conserved among vertebrates for the majority of subunits. We determined the expression of the eight zebrafish α subunit encoding genes, the largest and most diverse gene family, and observed that they are expressed in distinct, often overlapping expression patterns. Taken together, these data argue that larval zebrafish can serve as a useful model to investigate the functional roles of GABA$_A$ receptor subtypes within developing neural circuits.

A previous study identified 23 GABA$_A$ receptor genes in zebrafish. Our data largely confirm their results, except ρ3b, which is encoded by *gabrr3b*, was not reported. A α3 and an α3-like gene were also described, but our analysis indicates that zebrafish likely contain only one α3 gene. Given the amount of zebrafish genome data that is available, it seems unlikely that additional GABA$_A$ receptor subunit encoding genes exist. However, similar to mammalian systems, sequence data indicates that several zebrafish GABA$_A$ receptor encoding genes are alternatively spliced, which further enhances the already extensive diversity of receptor isoforms.

The evolution of the GABA$_A$ receptor gene family has been examined by using sequence data from a wide variety of species, including humans, rodents, canary, chicken, frogs and pufferfish, tunicates, *C. elegans* and *drosophila* [41, 42]. Our results are in line with these studies in finding that mammalian θ and ε subunits are more distinct compared to other subunits, such that their existence outside of the mammalian lineage is unclear [43]. Even among mammalian species, for example comparing humans to rodents, these two subunits are very diverse, which suggests they are evolving at a much faster rate than other subunits [44].

The roles of θ and ε subunits may be unique to mammals or, alternatively, fulfilled in zebrafish

and other species using other subtypes. Experiments to investigate the functional ability of

zebrafish subtypes to substitute for θ and ε in rodent systems could distinguish between these

possibilities.

The α subunits demonstrate a wide range of expression patterns across early zebrafish

development (Table 1). On one end of the spectrum are *gabra2a* and *gabra2b*, which appeared

to be expressed very broadly at each of the three time points examined. At the other end of the

spectrum is *gabra6a*, which we detected only at 96 hpf and mostly in the photoreceptor cell

layer of the retina. The other subunits fall between these extremes, with expression in discrete

cells in various brain and spinal cord regions. When GABA$_A$ receptor α subtypes are expressed

in the same region, they often occupy different domains. For example, *gabra1*, *gabra3*, *gabra4*,

and *gabra5* are all expressed in the medulla, but are organized in medial to lateral stripes. The

zebrafish medulla has a structural and functional organization in which neurons of shared

neurotransmitter, phenotype, age, morphology and functional properties are arranged into

stripes [45, 46]. Another example of different GABA$_A$ α subtypes occupying different domains

within the same region is found in the spinal cord. *gabra4* transcripts were found in ventral cells,

likely KA neurons, *gabra5*-expressing cells, likely CoPA neurons, were observed more dorsally,

and *gabra3* transcripts were detected more broadly in both ventral and more dorsal domains. It

is not yet clear how the expression of different GABA$_A$ subtypes correspond to the

organizational stripes of the hindbrain or cell types in the spinal cord, but it seems likely that

some subtypes demonstrate cell-type specific expression and confer distinct responses to

GABA. Conclusive cell-type identification will require colabeling experiments that examine

GABA$_A$ subunit expression along with specific markers or techniques that determine neuronal

identity by revealing cell morphology [47].

**359** **Table 1. Expression of GABA$_A$ receptor α subunits at 24, 48, and 96 hours post fertilization.**

| | | gabra1 | gabra2a | gabra2b | gabra3 | gabra4 | gabra5 | gabra6a | gabra6b |
|---|---|---|---|---|---|---|---|---|---|
| 24 hpf | dorsal rostral cluster | | • | • | | | | | |
| | hindbrain cluster | • | • | • | • | • | • | | |
| | olfactory placode | • | | | | | | | |
| | spinal cord | | | | • | • | • | | |
| | tegmentum | | | | • | | | | |
| | ventral caudal cluster | • | • | • | • | • | | | |
| | ventral rostral cluster | • | • | • | | | | | |
| 48 hpf | cerebellum | | • | • | | | | | • |
| | diencephalon | • | | | | | | | |
| | hypothalamus | | | | | | • | | |
| | Mauthner cell | | | | | | • | | |
| | medulla oblongata | • | | | • | • | | | |
| | olfactory bulb | • | | | | | | | • |
| | pallium | | | | • | • | • | | |
| | subpallium | | | | • | | | | |
| | tegmentum | | | | • | | | | |
| | thalamus | | | | • | | | | |
| 96 hpf | cerebellum | • | • | • | | • | • | • | • |
| | hypothalamus | • | | • | | | • | | |
| | Mauthner cell | | | | | | • | | |
| | medulla oblongata | • | | • | • | • | • | | |
| | olfactory bulb | | | | | | | | • |
| | optic nerve | • | | | | | | | |
| | Pallium | | | • | • | • | • | | |
| | post-optic commissure | | | • | | | | | |
| | pre-optic region | | | • | | | | | |
| | tectum opticum | • | | | | • | | | |
| | Tegmentum | • | | • | • | | | | |
| | thalamus | | | | • | | | | |
| | Retina- | | | | | | | | |
| | inner nuclear layer | • | | • | • | • | | | • |
| | ganglion cell layer | • | | • | | | | | • |
| | photoreceptive cell layer | | | | | | | • | |

**Table 1**Shaded boxes indicate broad, diffuse expression was observed and dots indicate

punctate expression was detected.

The expression patterns of GABA$_A$ α subunits are well-conserved across species. The

expression of five α subunits has been described in *Xenopus laevis* during development [48].

Consistent with our observations in zebrafish, α2 was found to be expressed broadly and early

in development. In fact, α2 was found to be deposited in eggs as a maternal message. α6

showed the latest onset of expression during development and demonstrated robust

photoreceptor expression. Likewise the expression patterns of α1, α3, and α5 are similar

between frogs and zebrafish. These similarities in expression extend to mammalian systems.

For example, in rat, similar to zebrafish α2, transcripts were detected relatively early in

development and were widespread, and α6 is expressed much later in development and

exhibits restricted expression [7]. One notable difference is that in mammals α6 transcripts were

detected in the cerebellum and retinal expression has not been reported, whereas in zebrafish

α6a exhibits robust retinal expression and α6b is more widespread. It is unclear how much

these differences in expression between zebrafish and mammals can be attributed to the

various development stages and tissues selected for investigation versus authentic differences

in patterns of expression.

Developing zebrafish are a powerful system for neural circuit analysis and, more

recently, they have been cultivated as a model of epilepsy [17, 49-53]. Although GABA$_A$

receptors are robust regulators of many neural circuits and play a central role in epilepsy

syndromes, GABA$_A$ receptors have not been well characterized in zebrafish. By reporting the

diversity of zebrafish GABA$_A$ receptor subunits and describing the expression patterns of the α

subfamily during early development, we have laid a foundation to leverage the strengths of the

zebrafish system to investigate isoform specific roles of GABA$_A$ receptors and further develop

zebrafish as a model of epilepsy. In mammalian systems, genetic inactivation of individual α

subunits has revealed surprisingly subtle defects compared to what would be predicted from

pharmacological blockade. Compensatory upregulation of other subunits is thought to mask

some of the effects of inactivating a single subunit [54, 55]. The ability easily mutate multiple

genes in zebrafish may help overcome such compensatory changes in expression to shed new

light into the subtype specific roles of $GABA_A$ receptors.

## ACKNOWLEDGEMENTS

392 The authors thank members of the Downes lab, Kelly Anne McKeown at Westfield State

393 University, and Josef Trapani at Amherst College for helpful advice and discussion.

## REFERENCES

1.      Bowery NG, Smart TG. GABA and glycine as neurotransmitters: a brief history. Br J
Pharmacol. 2006;147 Suppl 1:S109-19. Epub 2006/01/13. doi: 10.1038/sj.bjp.0706443. PubMed
PMID: 16402094; PubMed Central PMCID: PMCPMC1760744.

2.      Mohler H. GABA(A) receptor diversity and pharmacology. Cell Tissue Res.
2006;326(2):505-16. Epub 2006/08/29. doi: 10.1007/s00441-006-0284-3. PubMed PMID:
16937111.

3.      Sigel E, Steinmann ME. Structure, function, and modulation of GABA(A) receptors. J
Biol Chem. 2012;287(48):40224-31. Epub 2012/10/06. doi: 10.1074/jbc.R112.386664. PubMed
PMID: 23038269; PubMed Central PMCID: PMCPMC3504738.

4.      Chua HC, Chebib M. GABAA Receptors and the Diversity in their Structure and
Pharmacology. Adv Pharmacol. 2017;79:1-34. Epub 2017/05/23. doi:
10.1016/bs.apha.2017.03.003. PubMed PMID: 28528665.

5.      Olsen RW, Sieghart W. International Union of Pharmacology. LXX. Subtypes of gamma-
aminobutyric acid(A) receptors: classification on the basis of subunit composition,
pharmacology, and function. Update. Pharmacol Rev. 2008;60(3):243-60. Epub 2008/09/16.
doi: 10.1124/pr.108.00505. PubMed PMID: 18790874; PubMed Central PMCID:
PMCPMC2847512.

6.      Laurie DJ, Seeburg PH, Wisden W. The distribution of 13 GABAA receptor subunit
mRNAs in the rat brain. II. Olfactory bulb and cerebellum. J Neurosci. 1992;12(3):1063-76.
Epub 1992/03/01. PubMed PMID: 1312132.

7.       Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABAA receptor subunit
mRNAs in the rat brain. III. Embryonic and postnatal development. J Neurosci.
1992;12(11):4151-72. PubMed PMID: 1331359.

8.       Wisden W, Laurie DJ, Monyer H, Seeburg PH. The distribution of 13 GABAA receptor
subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. J Neurosci.
1992;12(3):1040-62. Epub 1992/03/01. PubMed PMID: 1312131.

9.       Delgado-Lezama R, Loeza-Alcocer E, Andres C, Aguilar J, Guertin PA, Felix R.
Extrasynaptic GABA(A) receptors in the brainstem and spinal cord: structure and function. Curr
Pharm Des. 2013;19(24):4485-97. Epub 2013/01/31. PubMed PMID: 23360278.

10.      Simon J, Wakimoto H, Fujita N, Lalande M, Barnard EA. Analysis of the set of GABA(A)
receptor genes in the human genome. J Biol Chem. 2004;279(40):41422-35. Epub 2004/07/20.
doi: 10.1074/jbc.M401354200. PubMed PMID: 15258161.

11.      Olsen RW, Sieghart W. GABA A receptors: subtypes provide diversity of function and
pharmacology. Neuropharmacology. 2009;56(1):141-8. Epub 2008/09/02. doi:
10.1016/j.neuropharm.2008.07.045. PubMed PMID: 18760291; PubMed Central PMCID:
PMCPMC3525320.

12.      Tretter V, Moss SJ. GABA(A) Receptor Dynamics and Constructing GABAergic
Synapses. Front Mol Neurosci. 2008;1:7. Epub 2008/10/24. doi: 10.3389/neuro.02.007.2008.
PubMed PMID: 18946540; PubMed Central PMCID: PMCPMC2526003.

13.      Fritschy JM, Panzanelli P. GABAA receptors and plasticity of inhibitory
neurotransmission in the central nervous system. Eur J Neurosci. 2014;39(11):1845-65. Epub
2014/03/19. doi: 10.1111/ejn.12534. PubMed PMID: 24628861.

14.      Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of
GABA(A) receptors. Nat Rev Neurosci. 2005;6(3):215-29. Epub 2005/03/02. doi:
10.1038/nrn1625. PubMed PMID: 15738957.

15.      Egawa K, Fukuda A. Pathophysiological power of improper tonic GABA(A)
conductances in mature and immature models. Front Neural Circuits. 2013;7:170. Epub
2013/10/30. doi: 10.3389/fncir.2013.00170. PubMed PMID: 24167475; PubMed Central PMCID:
PMCPMC3807051.

16.      Wolman M, Granato M. Behavioral genetics in larval zebrafish: learning from the young.
Dev Neurobiol. 2012;72(3):366-72. Epub 2012/02/14. doi: 10.1002/dneu.20872. PubMed PMID:
22328273.

17.      McLean DL, Fetcho JR. Movement, technology and discovery in the zebrafish. Curr Opin
Neurobiol. 2011;21(1):110-5. Epub 2010/10/26. doi: 10.1016/j.conb.2010.09.011. PubMed
PMID: 20970321; PubMed Central PMCID: PMCPMC3057273.

18.      Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in
zebrafish behavior, neural activity and c-fos expression. Neuroscience. 2005;131(3):759-68.
Epub 2005/02/26. doi: 10.1016/j.neuroscience.2004.11.031. PubMed PMID: 15730879.

19.      Stehr CM, Linbo TL, Incardona JP, Scholz NL. The developmental neurotoxicity of
fipronil: notochord degeneration and locomotor defects in zebrafish embryos and larvae. Toxicol
Sci. 2006;92(1):270-8. Epub 2006/04/14. doi: 10.1093/toxsci/kfj185. PubMed PMID: 16611622.

20.      Connaughton VP, Nelson R, Bender AM. Electrophysiological evidence of GABAA and
GABAC receptors on zebrafish retinal bipolar cells. Vis Neurosci. 2008;25(2):139-53. Epub
2008/04/30. doi: 10.1017/S0952523808080322. PubMed PMID: 18442437; PubMed Central
PMCID: PMCPMC4272819.

21.    Del Bene F, Wyart C, Robles E, Tran A, Looger L, Scott EK, et al. Filtering of visual
information in the tectum by an identified neural circuit. Science. 2010;330(6004):669-73. Epub
2010/10/30. doi: 10.1126/science.1192949. PubMed PMID: 21030657; PubMed Central PMCID:
PMCPMC3243732.

22.    Mack-Bucher JA, Li J, Friedrich RW. Early functional development of interneurons in the
zebrafish olfactory bulb. Eur J Neurosci. 2007;25(2):460-70. Epub 2007/02/08. doi:
10.1111/j.1460-9568.2006.05290.x. PubMed PMID: 17284187.

23.    Hubbard JM, Bohm UL, Prendergast A, Tseng PB, Newman M, Stokes C, et al.
Intraspinal Sensory Neurons Provide Powerful Inhibition to Motor Circuits Ensuring Postural
Control during Locomotion. Curr Biol. 2016;26(21):2841-53. Epub 2016/10/11. doi:
10.1016/j.cub.2016.08.026. PubMed PMID: 27720623.

24.    Triller A, Rostaing P, Korn H, Legendre P. Morphofunctional evidence for mature
synaptic contacts on the Mauthner cell of 52-hour-old zebrafish larvae. Neuroscience.
1997;80(1):133-45. Epub 1997/09/01. PubMed PMID: 9252227.

25.    Roy B, Ali DW. Multiple types of GABAA responses identified from zebrafish Mauthner
cells. Neuroreport. 2014;25(15):1232-6. Epub 2014/08/28. doi:
10.1097/WNR.0000000000000258. PubMed PMID: 25162782.

26.    Cocco A, Ronnberg AM, Jin Z, Andre GI, Vossen LE, Bhandage AK, et al.
Characterization of the gamma-aminobutyric acid signaling system in the zebrafish (Danio rerio
Hamilton) central nervous system by reverse transcription-quantitative polymerase chain
reaction. Neuroscience. 2017;343:300-21. Epub 2016/07/28. doi:
10.1016/j.neuroscience.2016.07.018. PubMed PMID: 27453477.

27.    Hortopan GA, Dinday MT, Baraban SC. Spontaneous seizures and altered gene
expression in GABA signaling pathways in a mind bomb mutant zebrafish. J Neurosci.
2010;30(41):13718-28. Epub 2010/10/15. doi: 10.1523/JNEUROSCI.1887-10.2010. PubMed
PMID: 20943912; PubMed Central PMCID: PMCPMC2962868.

28.    Gonzalez-Nunez V. Role of gabra2, GABAA receptor alpha-2 subunit, in CNS
development. Biochem Biophys Reports. 2015;3:11.

29.    Baxendale S, Holdsworth CJ, Meza Santoscoy PL, Harrison MR, Fox J, Parkin CA, et al.
Identification of compounds with anti-convulsant properties in a zebrafish model of epileptic
seizures. Dis Model Mech. 2012;5(6):773-84. Epub 2012/06/26. doi: 10.1242/dmm.010090.
PubMed PMID: 22730455; PubMed Central PMCID: PMCPMC3484860.

30.    Thisse B, Thisse, C. . Fast Release Clones: A High Throughput Expression Analysis.
ZFIN Direct Data Submission. 2004.

31.    Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. Nucleic Acids Res. 1994;22(22):4673-80. Epub 1994/11/11.
PubMed PMID: 7984417; PubMed Central PMCID: PMCPMC308517.

32.    Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. Syst Biol. 2010;59(3):307-21. Epub 2010/06/09. doi: 10.1093/sysbio/syq010.
PubMed PMID: 20525638.

33.    Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic
development of the zebrafish. Dev Dyn. 1995;203(3):253-310. PubMed PMID: 8589427.

34.    Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount zebrafish
embryos. Nat Protoc. 2008;3(1):59-69. PubMed PMID: 18193022.

35.    Mueller T, Wullimann MF. Atlas of early zebrafish brain development : a tool for
molecular neurogenetics. Second edition. ed. xiii, 244 pages p.

36.    Folgueira M, Bayley P, Navratilova P, Becker TS, Wilson SW, Clarke JD.
Morphogenesis underlying the development of the everted teleost telencephalon. Neural Dev.
2012;7:32. Epub 2012/09/20. doi: 10.1186/1749-8104-7-32. PubMed PMID: 22989074; PubMed
Central PMCID: PMCPMC3520737.

37.    Sabaliauskas NA, Foutz CA, Mest JR, Budgeon LR, Sidor AT, Gershenson JA, et al.
High-throughput zebrafish histology. Methods. 2006;39(3):246-54. Epub 2006/07/28. doi:
10.1016/j.ymeth.2006.03.001. PubMed PMID: 16870470.

38.    Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, et al. Zebrafish hox clusters
and vertebrate genome evolution. Science. 1998;282(5394):1711-4. Epub 1998/11/30. PubMed
PMID: 9831563.

39.    Postlethwait JH, Yan YL, Gates MA, Horne S, Amores A, Brownlie A, et al. Vertebrate
genome evolution and the zebrafish gene map. Nat Genet. 1998;18(4):345-9. Epub 1998/04/16.
doi: 10.1038/ng0498-345. PubMed PMID: 9537416.

40.    Saint-Amant L, Drapeau P. Time course of the development of motor behaviors in the
zebrafish embryo. J Neurobiol. 1998;37(4):622-32. PubMed PMID: 9858263.

41.    Tsang SY, Ng SK, Xu Z, Xue H. The evolution of GABAA receptor-like genes. Mol Biol
Evol. 2007;24(2):599-610. Epub 2006/12/01. doi: 10.1093/molbev/msl188. PubMed PMID:
17135332.

42.    Darlison MG, Pahal I, Thode C. Consequences of the evolution of the GABA(A) receptor
gene family. Cell Mol Neurobiol. 2005;25(3-4):607-24. Epub 2005/08/03. doi: 10.1007/s10571-
005-4004-4. PubMed PMID: 16075381.

43.    Martyniuk CJ, Aris-Brosou S, Drouin G, Cahn J, Trudeau VL. Early evolution of
ionotropic GABA receptors and selective regimes acting on the mammalian-specific theta and
epsilon subunits. PLoS One. 2007;2(9):e894. Epub 2007/09/20. doi:
10.1371/journal.pone.0000894. PubMed PMID: 17878929; PubMed Central PMCID:
PMCPMC1975676.

44.    Sinkkonen ST, Hanna MC, Kirkness EF, Korpi ER. GABA(A) receptor epsilon and theta
subunits display unusual structural variation between species and are enriched in the rat locus
ceruleus. J Neurosci. 2000;20(10):3588-95. Epub 2000/05/11. PubMed PMID: 10804200.

45.    Higashijima S, Mandel G, Fetcho JR. Distribution of prospective glutamatergic,
glycinergic, and GABAergic neurons in embryonic and larval zebrafish. J Comp Neurol.
2004;480(1):1-18. Epub 2004/10/30. doi: 10.1002/cne.20278. PubMed PMID: 15515020.

46.    Kinkhabwala A, Riley M, Koyama M, Monen J, Satou C, Kimura Y, et al. A structural and
functional ground plan for neurons in the hindbrain of zebrafish. Proc Natl Acad Sci U S A.
2011;108(3):1164-9. Epub 2011/01/05. doi: 10.1073/pnas.1012185108. PubMed PMID:
21199947; PubMed Central PMCID: PMCPMC3024665.

47.    Downes GB, Waterbury JA, Granato M. Rapid in vivo labeling of identified zebrafish
neurons. Genesis. 2002;34(3):196-202. PubMed PMID: 12395384.

48.    Kaeser GE, Rabe BA, Saha MS. Cloning and characterization of GABAA alpha subunits
and GABAB subunits in Xenopus laevis during development. Dev Dyn. 2011;240(4):862-73.

Epub 2011/03/09. doi: 10.1002/dvdy.22580. PubMed PMID: 21384470; PubMed Central
PMCID: PMCPMC3071254.

49. Fetcho JR, Higashijima S, McLean DL. Zebrafish and motor control over the last decade.
Brain Res Rev. 2008;57(1):86-93. Epub 2007/09/11. doi: 10.1016/j.brainresrev.2007.06.018.
PubMed PMID: 17825423; PubMed Central PMCID: PMCPMC2237884.

50. Friedrich RW, Genoud C, Wanner AA. Analyzing the structure and function of neuronal
circuits in zebrafish. Front Neural Circuits. 2013;7:71. Epub 2013/05/01. doi:
10.3389/fncir.2013.00071. PubMed PMID: 23630467; PubMed Central PMCID:
PMCPMC3632777.

51. Hale ME, Katz HR, Peek MY, Fremont RT. Neural circuits that drive startle behavior,
with a focus on the Mauthner cells and spiral fiber neurons of fishes. J Neurogenet.
2016;30(2):89-100. Epub 2016/06/16. doi: 10.1080/01677063.2016.1182526. PubMed PMID:
27302612.

52. Griffin A, Krasniak C, Baraban SC. Advancing epilepsy treatment through personalized
genetic zebrafish models. Prog Brain Res. 2016;226:195-207. Epub 2016/06/22. doi:
10.1016/bs.pbr.2016.03.012. PubMed PMID: 27323944.

53. Cunliffe VT. Building a zebrafish toolkit for investigating the pathobiology of epilepsy and
identifying new treatments for epileptic seizures. J Neurosci Methods. 2016;260:91-5. Epub
2015/07/30. doi: 10.1016/j.jneumeth.2015.07.015. PubMed PMID: 26219659.

54. Kralic JE, Sidler C, Parpan F, Homanics GE, Morrow AL, Fritschy JM. Compensatory
alteration of inhibitory synaptic circuits in cerebellum and thalamus of gamma-aminobutyric acid
type A receptor alpha1 subunit knockout mice. J Comp Neurol. 2006;495(4):408-21. Epub
2006/02/18. doi: 10.1002/cne.20866. PubMed PMID: 16485284.

55. Peng Z, Zhang N, Chandra D, Homanics GE, Olsen RW, Houser CR. Altered
localization of the delta subunit of the GABAA receptor in the thalamus of alpha4 subunit
knockout mice. Neurochem Res. 2014;39(6):1104-17. Epub 2013/12/20. doi: 10.1007/s11064-
013-1202-1. PubMed PMID: 24352815; PubMed Central PMCID: PMCPMC4024081.

## Supporting Information

**S1 Fig. RT-PCR results suggest that zebrafish contain only one α3 gene**

(A) Schematic of a *gabra3* cDNA. The portions that are identical to *gabra3*-like and *gabra3*

sequences are shaded green. The white region that connects the two was identified through RT-

PCR. The location of the three primers used for PCR are shown as numbered arrows. (B) RT-PCR

results. Primers 2 and 3 served as a positive control since they amplify sequence from the known

*gabra3* region. Primers 1 and 3 amplify a previously unknown region that links *gabra* 3-like and

*gabra3* sequence, showing that they are contained within same transcript and likely portions of the

same gene.

**S1 Table. GABA$_A$ α subunit antisense RNA probe information.** Note- Sizes are shown in base

pairs. The probe start site is defined here as the start codon of the open reading frame. Negative

values indicate that the probe contains 5' untranslated sequence. Some probes were generated via

RT-PCR with a primer that contains a RNA polymerase promoter and subcloning into vectors wasn't

performed.