Role of dynamic cooperativity in the mechanism of

hexameric M17 aminopeptidases

Nyssa Drinkwater [a,1,2], Wei Yang [a,1], Blake T. Riley [b], Komagal Kannan Sivaraman [a], Itamar Kass [b,c], Ashley M. Buckle [b], Sheena McGowan [a, 2]

[a] Biomedicine Discovery Institute, Department of Microbiology, Monash University, Clayton Melbourne, VIC 3800, Australia

[b] Biomedicine Discovery Institute, Department of Biochemistry and Molecular Biology, Monash University, Clayton Melbourne, VIC 3800, Australia

[c] Victorian Life Sciences Computation Centre, Monash University, Clayton 3800, Victoria, Australia

[1] N.D. and W.Y. contributed equally to this work.

[2] To whom correspondence should be addressed: nyssa.drinkwater@monash.edu; +613 99029371 sheena.mcgowan@monash.edu; +613 990293009

Abstract

M17 aminopeptidases possess a conserved hexameric arrangement throughout all kingdoms

of life. Of particular interest is the M17 from *Plasmodium falciparum* (*Pf*A-M17), which is a

validated antimalarial drug target. Herein we have examined *Pf*A-M17 using an integrated

structural biology and biochemical approach to provide the first description of the fundamental role

of oligomerisation. We found that, rather than operating as discrete units, the active sites of the

*Pf*A-M17 hexamer are linked by a dynamic loop, which operates cooperatively to regulate activity.

Further, we characterised motions in key surface loops that moderate access to the central catalytic

cavity. Based on our new understanding of the dynamics inherent to *Pf*A-M17, we propose a novel

mechanism that would allow exquisite control of enzyme function in response to cellular signals,

and go on to discuss how, through divergent evolution, this mechanism might have developed to

moderate key differences in M17 function across species.

Introduction:

Our appreciation of protein flexibility and dynamics has grown remarkably from the early

static lock-and-key model, to our current understanding that proteins are dynamic entities capable

of extreme flexibility. Dynamics are now recognised as integral to protein function and regulation,

and therefore fundamental to life. For enzymes, flexibility influences substrate binding (Vögeli,

Bibow, & Chi, 2016; Wurm, Holdermann, Overbeck, Mayer, & Sprangers, 2017) and reaction

mechanism (Bhabha et al., 2011; McElheny, Schnell, Lansing, Dyson, & Wright, 2005), allowing

precise control of reactions that if left unregulated, could be harmful to the cell. Despite the clear

importance of enzyme flexibility, there are few cases where dynamics, and how those dynamics

link to function, are truly understood on an atomic level.

Intracellular proteolysis requires precise spatial and temporal control to prevent cleavage of

proteins not destined for destruction. For this purpose, high-molecular weight protease enzymes are

self-compartmentalised, whereby the active sites are enclosed in inner cavities isolated from the

cellular environment. Such an arrangement is often mediated by multimeric self-association, such as that seen for the family of M17 aminopeptidases, otherwise known as PepA or LAP (leucyl aminopeptidases; Clan MF; M17 family) (MEROPS (Rawlings, Barrett, & Finn, 2016)), which are present in all kingdoms of life. Although overall sequence conservation is low, M17 aminopeptidases possess a conserved homohexameric structure wherein a dimer of trimers encloses an inner cavity harbouring the six active sites (Lowther & Matthews, 2002). The sequence and structure of the proteolytic sites themselves, as well as the reaction they catalyse, are highly conserved, all utilising two divalent metal ion cofactors to catalyse the removal of select N-terminal amino acids from short peptide chains (Lowther & Matthews, 2002). This reaction contributes to intracellular protein turnover, a fundamental housekeeping process across all living organisms (Matsui, Fowler, & Walling, 2006). However, a wide range of additional functions beyond aminopeptidase activity have also been attributed to M17 family members. M17 proteases have been shown in plants to function as molecular chaperones to control stress-induced damage (Kumar, Kaur, Chattopadhyay, & Bachhawat, 2015; Scranton, Yee, Park, & Walling, 2012), while in bacteria they form hetero-oligomers (Minh, Devroede, Massant, Maes, & Charlier, 2009; Sträter, Sherratt, & Colloms, 1999) and contribute to site-specific DNA recombination (Alén, Sherratt, & Colloms, 1997; Stirling, Colloms, Collins, Szatmari, & Sherratt, 1989) and transcriptional control (Charlier et al., 1995). Therefore, although the family of M17 aminopeptidases have a highly conserved structure across different organisms, they are multifunctional, capable of performing diverse organism-specific functions far beyond peptide hydrolysis. Our understanding of mammalian M17 aminopeptidases is based on early investigations into the structure and catalytic mechanism of the bovine lens M17 aminopeptidase, which while considerable, consider the six chains within the hexamer as discrete entities. Further, although some flexibility within the active site has been considered (Schurer, Horn, Gedeck, & Clark, 2002), the role/s of the conserved hexameric arrangement of M17 aminopeptidases, as well as motions within this arrangement, have thus far not been investigated.

Of particular interest is the M17 aminopeptidase from *Plasmodium falciparum* (*Pf*A-M17),

the major causative agent of malaria in humans. *Pf*A-M17 has a proposed role in haemoglobin

digestion, the essential process by which parasites break down human haemoglobin to yield the free

amino acids necessary for growth and development (Rosenthal, 2002; Stack et al., 2007). *Pf*A-M17

is essential for the blood stage of the parasite life cycle (Dalal & Klemba, 2007) and inhibition of

*Pf*A-M17 results in parasite death both *in vitro* (Harbut et al., 2011) and *in vivo* (Skinner-Adams et

al., 2007). However, this effect occurs early in the life cycle before haemoglobin digestion is

thought to have initiated, which has led to speculation that *Pf*A-M17 possesses additional unknown

function/s (Harbut et al., 2011). Irrespectively, *Pf*A-M17 is an exciting target for the development

of novel antimalarial therapeutics (Drinkwater, Bamert, Sivaraman, Paiardini, & McGowan, 2015;

Drinkwater et al., 2016; Mistry et al., 2014; Skinner-Adams et al., 2007; Skinner-Adams et al.,

2012). The crystal structure of *Pf*A-M17 shows the homohexameric arrangement characteristic of

M17 aminopeptidase enzymes (Fig. 1A), with the six active sites orientated inwards and accessible

to the central cavity (Fig. 1B and C) (McGowan et al., 2010). We were interested to determine the

mechanistic basis of the conserved hexameric assembly by investigating the dynamics inherent to

*Pf*A-M17 and probing how they contribute to function.

Figure 1. *Pf*A-M17 is a homo hexamer composed of a dimer of trimers. (A) Three chains, *a* (blue), *b* (purple), and *c* (teal) form a 'trimer' (*abc*), which interacts with an identical trimer made up of *a** (yellow), *b** (orange), and *c** (red) to form the hexamer (abc)₂. (B) *Pf*A-M17 (cartoon) possesses a large internal cavity (grey surface representation) that contains the six active sites. (C) A single trimeric face (*abc*) shown in the same orientation as A (front trimer hidden). Orange spheres substrate-analogue bestatin bound in the active sites, which are exposed to the inner cavity.

Results and Discussion:

Does hexamerisation play a functional role?

The hexameric arrangement of M17 aminopeptidases is highly conserved, however, the contribution of this assembly to enzyme function is unknown. The arrangement is characterised as a dimer of trimers (Burley, David, Taylor, & Lipscomb, 1990); three chains (*a*, *b*, and *c*, Fig. 1A) interact via their C-terminal catalytic domains to form a 'trimer' (*abc*), two of which (*abc* and *abc**) then associate to form the hexamer, (*abc*)$_2$ (Fig. 1A). The result is a large central cavity containing the six active sites arranged in two symmetrical disk-like arrangements (Fig. 1B and 1C). Despite the proximity of the active sites, both unliganded and liganded crystal structures of *Pf*A-M17 show they exist as six discrete units (McGowan et al., 2010). Each of the active sites is composed entirely of residues from a single chain and contains its own catalytic machinery that includes two zinc ions and a carbonate ion. Based on the current evidence, hexamerisation does not have a clear role in the catalysis. However, the M17 aminopeptidase from *H. pylori* (*Hp*-M17) exhibits positive cooperativity, which suggests that communication between the *Hp*-M17 active sites does occur (Dong et al., 2005). Therefore, to probe the role of *Pf*A-M17 hexamerisation in catalysis, we first attempted to identify whether any cooperativity between the *Pf*A-M17 active sites exists, or if the sites operate as discrete units. We employed a fluorescence-based aminopeptidase activity assay in substrate saturation experiments, wherein we assessed the relationship between substrate concentration and reaction velocity and analysed the Hill coefficient (SI 1). In contrast to the results of *Hp*-M17, we were unable to detect any evidence of cooperativity during *Pf*A-M17 catalysis (Hill coefficient, n_H = 1.0), which suggests that although both *Hp*-M17 and *Pf*A-M17 share a conserved overall assembly and highly conserved active site structures, they process substrates differently. This result is in contrast to the major assumption currently made within the M17 aminopeptidase literature, that the catalytic mechanism of the enzymes are conserved, and operate as has been described for the M17 enzyme from bovine lens (Lowther & Matthews, 2002).

Gatekeeper loops mediate access to central catalytic cavity

In the absence of evidence showing catalytic cooperativity between the active sites of the *Pf*A-M17 hexamer, we hypothesised that oligomerisation plays a regulatory role. To examine this theory, we turned to all-atom molecular dynamics (MD) simulations to determine the mechanism by which protein dynamics might be moderating *Pf*A-M17 activity. We performed all-atom MD simulations of hexameric *Pf*A-M17 (3 × 400 ns). Analysis of the root mean square deviation (RMSD) over the course of the simulation indicated that the hexamer did not undergo any large conformational changes, nor rigid body movements (average RMSD of Cα atoms = 2.5 ± 0.04 Å, SI 2A). We performed a principal component analysis (PCA) of the simulations on the backbone and zinc atoms of hexameric *Pf*A-M17, and observed that the top PC, PC1, accounts for 62 % of the total variance, while PC2 accounts for only 10 %. Projecting the trajectories onto the top two PCs showed the major motion, described by PC1, is an expansion of the hexamer from 120 Å to 127 Å (average of three measurements between Cα of Asn181[a], Asn181[b] and Asn181[c]) (SI 2B). The expansion is ~ 6%, and likely results primarily from the release of crystal constraints (Gerstein & Chothia, 1996).

The crystal structure of *Pf*A-M17 showed six channels connecting the central catalytic cavity to the protein surface, and identified a ~20 Å flexible loop (residues 246–265) that sits at the entrance to the channels (McGowan et al., 2010). The complete loop was able to be modelled in only one of the six chains, where it occluded the channel entrance, suggesting that the loops may regulate substrate ingress and/or product egress (McGowan et al., 2010). The putative access channels, flexible loops, and any motions they might undergo, were therefore of great interest to us. Initial modelling and energy minimisation of the starting loop conformations resulted in almost complete occlusion of the channel entrances (diameters ~4–5 Å across the bottleneck, Fig 2A, SI 3), in line with the conformation modelled for the single complete chain in the original crystal structure. Over the course of the MD simulation, the diameter of the pores increased, opening some of the channels to external solvent (Fig. 2B). Pore increases were partially mediated by the flexible

loop identified in the crystal structure (residues 246–265), but also by a second loop on the other

side of the channel, which is contributed from the opposing trimer in the hexamer (residues 132–

150). The 'gatekeeper loops' underwent a range of motions, affecting the size of the channels to

varying degrees. While one channel increased in diameter throughout the simulation (channel 6,

increased from 4.9 Å to 8.9 Å, Δ4.0 Å, Fig. 2C, SI 3), another remained occluded throughout the

simulation (channel 3 starting diameter was 4 Å compared to 4.5 Å at completion, Δ0.5 Å, Fig. 2D,

SI 3). Intriguingly, these two pores that sampled the most extreme changes (Δ0.5 Å in channel 3

versus Δ4.0 Å in channel 6) are in close proximity, on opposite sides of the same dimer pair

(channel 3 is formed by $c^{247-266}$ and $c*^{132-150}$, channel 6 by $c^{132-150}$ and $c*^{247-266}$, Fig. 2C and 2D),

which suggests that the channels are linked and operating in opposing concert. Overall, our analysis

of the putative access channels by MD demonstrates that channel opening, and consequently access

to the central catalytic cavity, may be mediated by motions in these key gatekeeper loops. Further,

each channel entrance is controlled by the loops of neighbouring chains of the hexamer, which

presents the first evidence that hexamerisation plays a role in moderating access to the catalytic

core. By regulating the size and/or nature of the access channels to the central cavity,

hexamerisation might also play an important role in substrate discrimination by mediating the type

of substrate allowed access to the inner cavity. Such a role has key implications for understanding

the wide range of different functionalities observed throughout the family of M17 aminopeptidases.

Figure 2. Flexible loops moderate channel size and access to catalytic cavity of hexameric *Pf*A-M17. (A) *Pf*A-M17 conformation at the start of the MD simulation shown in surface representation (transparent grey) and cartoon (*a* in blue, *b* in purple, *c* in teal, *a** in yellow, *b** in orange, *c** in red). Access of the six channels (blue) to external solvent is blocked by gatekeeper loops at the entrance to the channels. Yellow circles indicate the occluded channels (4, 5, and 6) on the front face of the hexamer. (B) During the simulation, the six access channels become accessible, creating a range of pore access sizes from 'open' (8.9 Å, channel 6, $c^{132-150}$:$c*^{247-266}$) shown in (C) to 'closed' (3.9 Å, channel 3, $c^{247-266}$:$c*^{132-150}$) shown in (D).

Hexamerisation is essential for catalytic function

Our MD analysis suggests that hexamerisation serves a regulatory role by mediating ingress and/or egress routes to the central catalytic cavity via the channels described above. If this is the sole purpose of the hexameric arrangement, then oligomerisation would be expected to control reaction rate or specificity, not the ability of the enzyme to perform the hydrolysis reaction itself. We therefore questioned whether trimeric, or indeed monomeric, *Pf*A-M17 would possess aminopeptidase activity. Examination of the *Pf*A-M17 crystal structure identified Trp525 and Tyr533 as key residues that mediate association of *abc* and *abc** into $(abc)_2$ and from this, we hypothesised that their elimination would ablate hexamerisation. *Pf*A-M17(W525A+Y533A) expressed and purified similarly to the wild type enzyme, however, analysis by analytical gel filtration chromatography showed that the mutated enzyme was monomeric in solution (SI 4A). Examination of *Pf*A-M17(W525A+Y533A) activity showed that the monomeric enzyme is inactive, even at high concentrations (SI 4B). It is therefore evident that hexamer formation is essential for *Pf*A-M17 proteolytic activity, and that oligomerisation plays an important functional role beyond regulation of substrate ingress and/or product egress.

Hexameric assembly stabilises and links the *Pf*A-M17 active sites

Having established that monomeric *Pf*A-M7 is unable to function independently, we continued to search for the key functional role of hexamerisation. We performed MD simulations of the monomeric enzyme and compared the output to the simulation of hexameric *Pf*A-M17. Over the time course sampled, the monomer trajectories underwent movements equivalent to an average RMSD of 4.1 ± 0.5 Å, demonstrating that the monomer underwent greater structural changes than the hexameric enzyme (RMSD of 2.5 ± 0.04 Å). Further, in the simulation of monomeric *Pf*A-M17, variation was observed between triplicate runs (SI 5A). This is in contrast to the hexamer simulation where all three runs showed a similar RMSD profile (SI 2A). This difference in the dynamics of hexameric versus monomeric *Pf*A-M17 supports an assertion that the hexameric

assembly stabilises the monomer to enable catalysis. To determine how this stabilisation translates

to the catalytic machinery, we examined the local environment of the active site throughout the

simulation. The catalytic mechanism of M17 aminopeptidases relies on the deprotonation of a

catalytic water for nucleophilic attack on the peptide carbonyl (Schurer et al., 2002; Sträter &

Lipscomb, 1995). In the simulation of monomeric *Pf*A-M17, we did not observe a stable position

for a water molecule within the active site (SI 5D). In contrast, in the hexameric *Pf*A-M17

simulation, we observed a stable water conformation wherein it associates with the site 1 zinc ion

(SI 5E), consistent with the position of a nucleophilic water (Schurer et al., 2002; Yang et al.,

2017). An active site with increased mobility may compromise catalysis, which relies on precise

placement of chemical moieties. We therefore assessed the mobility of individual residues

throughout the simulations via calculation of the root mean square fluctuations (RMSF) of the Cα

atoms (SI 5B and C). The RMSF analysis showed clear differences between the two simulations,

particularly residues 385–391, which lie on a loop flanking the active site on the interior of the

central cavity. In the simulation of monomeric *Pf*A-M17, this loop exhibited a high degree of

flexibility (average RMSF over three runs for residues 385–391 = 4.5 ± 0.4 Å). However, in the

simulation of hexameric *Pf*A-M17, the loop was relatively stable in four chains (chain *a*, *a**, b* and

*c**, average RMSF = 1.6 – 2.1 Å) and only moderately flexible in two chains (chain *b* and *c*,

average RMSF *b* = 2.8 ± 0.4 and *c* = 3.1 ± 0.4 Å). These key differences between the dynamics of

monomeric and hexameric *Pf*A-M17 demonstrate that hexamerisation enables catalysis by shielding

the essential nucleophilic water, and further, stabilises key active site loop structures. Such a

mechanism of preserving active site stability by oligomerisation has been previously described, for

example, DHDPS, which utilises a tetrameric arrangement to stabilise the catalytic dimerization

interface for optimal catalytic efficiency (Reboul et al., 2012).

Early work based on static structures of M17 aminopeptidases, primarily the M17 from

bovine lens, suggested that the nucleophilic water is likely deprotonated by either a bound

carbonate molecule or a conserved active site lysine residue (Sträter & Lipscomb, 1995). Density

function theory was applied, which suggested that the energy barrier for the lysine to act as the

catalytic base is prohibitively high, and therefore implicated the carbonate ion for this role (Zhu et

al., 2012). Conversely, use of a semiempirical quantum mechanical/molecular mechanical hybrid

approach supported the model in which the lysine residue is the most likely candidate as a catalytic

base (Schurer et al., 2002; Zhu et al., 2012). Sequence alignment of bovine M17 and *Pf*A-M17

determined the equivalent lysine to be residue 386 (Lys386), which lies on the active site loop

shown to fluctuate in the *Pf*A-M17 MD. In the crystal structure the loop lines the active site and

extends into the solvent of the inner cavity. PCA showed that in PC2 of the *Pf*A-M17 hexamer

simulation, the loop from one chain of each trimer b and c*($b^{383\text{-}397}$ and $c*^{383\text{-}397}$) moves away from

its own active site, extends across the entrance of the pocket and stretches toward the active site of

the neighbouring chain (Movie 1). In the most extreme conformation, observed in *c** (and *b* to a

slightly lesser extent), the loop occludes the entrance to the pocket and makes contact with the

equivalent loop of the neighbouring chain, which has not undergone the movement. Although the

loop does show some movement in the remaining chains, the extreme motion is observed for one

site per trimeric unit only (*b* and *c**). This analysis suggests communication between active sites

within each of the trimeric faces of *Pf*A-M17, and prompted us to employ further methods to

experimentally examine the atomic detail of alternative *Pf*A-M17 conformations.

Movie 1: Movie showing displacement along PC2 of *Pf*A-M17 hexamer simulation. *Pf*A-M17

trimer with greatest level of movement shown only (*abc**), with chain *a** in yellow, *b** in orange,

*c** in red.

https://www.dropbox.com/s/ir8pzku1he4khwq/Movie1_2880x2160.10Mbps.mp4?dl=0
254

Novel *Pf*A-M17 conformation captured by crystallography

The MD simulations identified a flexible loop capable of linking one *Pf*A-M17 active site to the neighbouring unit of the trimer. This is the first evidence that shows the active sites within the trimer might be linked, and raises the possibility that *Pf*A-M17 might utilise cooperativity during catalysis. If different conformations of *Pf*A-M17 do indeed exist throughout the catalytic reaction, we rationalised that we might be able to capture a previously unobserved state in a novel crystal form and determine its structure by X-ray crystallography. We therefore screened for different *Pf*A-M17 crystallisation conditions and solved a novel structure to 2.3 Å by molecular replacement (SI 6). The structure consisted of two copies of the hexamer in the asymmetric unit, with the overall quaternary structure similar to previous *Pf*A-M17 structures. However, the new structure shows a vastly different active site arrangement to any previously observed M17 structure. In the previously determined *Pf*A-M17 structure, herein referred to as the 'active' conformation, the flexible active site loop (residues 379–391) lines the active site and extends into the solvent of the inner cavity. In the new conformation, this loop is observed to cross the active site, completely occluding its entrance (Fig 3). Further, the loop extends to the active site of the neighbouring chain in the trimer, where it occupies the binding pocket with a key lysine residue (Lys386) (Fig 4, Movie 2). The same ~13 Å loop rearrangement is observed for all binding pockets, resulting in a direct link between the three active sites of each trimer: a^{386} inserts into the binding pocket of b, b^{386} inserts into the binding pocket of c, c^{386} inserts into the binding pocket of a (equivalent movements repeated in *abc**, Fig. 4C–D, Movie 2). To obtain the slack within the loop to adopt this extended, occluded conformation, the secondary structure at both ends of the loop has been disrupted (Fig 3C). This includes disruption of the α-helical structure of nine residues (392–401) and complete disruption of a short beta strand (372–379) (Fig 3D, Fig 4).

Figure 3. Novel *Pf*A-M17 conformation shows the active site occluded by loop (A and B)
Overlay of a single monomer of the 'occluded' *Pf*A-M17 (yellow cartoon) with bestatin-bound
'active' *Pf*A-M17. Bestatin binding mode (orange) indicates the active site. Yellow cartoon
protruding through the surface shows the position of the loop in the 'occluded' conformation. The
loop moves ~13 Å (yellow dash) to occlude the active site. (C) Overlay of 'active' (blue) and
'occluded' (grey and yellow) crystal structures of *Pf*A-M17. Orientation equivalent to B. Black
dash indicates the ~13 Å movement of the loop (yellow), which disrupts an alpha helix and beta
sheet. (D) inverted representation of panel C, wherein 'occluded' *Pf*A-M17 (grey) is shown in
surface representation, with the loop (yellow) completely occluding the active site.
288

289
290

Figure 4. Conformational change from active to inactive *Pf*A-M17 mediated by a flexible regulatory loop. (A) Active site arrangement of active *Pf*A-M17 at the junction of the *abc* trimer; chain *a* (light grey, regulatory loop and zincs yellow), chain *b* (dark grey, regulatory loop and zincs magenta), chain *c* (medium grey, regulatory loop and zincs cyan). Lys386 of all chains shown in stick representation. (B) Zoom of chain *a* active site in active *Pf*A-M17. Orientation equivalent to panel A. Zinc positions 1 and 2 are and active site residues are indicated (subscript denotes chain identifier). (C) Active site arrangement of inactive *Pf*A-M17, showing that the active sites are linked by the regulatory loop. Coloring and orientation consistent with panel A. Lys386 sits in the active site of the neighbouring chain, thereby linking all three active sites within the trimer. (D) Zoom of chain *b* active site in inactive. Disruption of active site α-helix (residues 394–400) is observed, and re-organisation of the active site architecture, including zinc binding positions has occurred.

306   **Movie 2:** Movie showing morph between active and inactive *Pf*A-M17. Chain *a* in blue, *b* in

307   purple, *c* in teal, *a** in yellow, *b** in orange, and *c** in red.

308   https://www.dropbox.com/s/sl82io7vwur62bv/Movie2_2880x2160.10Mbps.mp4?dl=0
309

The active site rearrangement extends beyond protein backbone changes, and includes the catalytic zinc ions absolutely required for enzyme activity. Based on kinetic and biophysical characterisation, the two zinc sites of the M17 aminopeptidases have previously been termed site 1 and site 2, whereby site 1 is that closest to the mouth of the active site (Fig. 4B) (Lowther & Matthews, 2002). Site 1 is readily lost from the active site, as observed by kinetic and crystallographic studies, while site 2 is always occupied in structures, and its removal has been reported to result in irreversible ablation of catalytic activity (Maric et al., 2009). Therefore, site 2 is generally considered the 'tight' binding, catalytic site (Allen, Yamada, & Carpenter, 1983; Maric et al., 2009; McGowan et al., 2010). In the novel conformation described here, we observed re-arrangement of the zinc binding positions (Fig. 4B and 4D). While site 1 is occupied with a zinc ion, the 'catalytic' zinc in site 2 is absent. Further, a zinc ion is bound in a third, previously uncharacterised site, coordinated by the side chains of Asp394 and Asp399, and the main chain oxygen of Met362, as well as two ordered water molecules, changing the coordination from tetrahedral (Fig 4B) to octahedral (Fig 4D). Dialysis in mixed metal buffers (Zn^{2+} and Co^{2+}) has previously shown that metal ion exchange between site 1 and 2 is also possible (Allen et al., 1983), therefore re-arrangement of the active site zinc ions is not unprecedented, though the existence of a third site was not considered. In the new conformation characterised here, the zinc coordination by Asp394 is significant since this residue lies within the flexible loop (residues 375–401), and it is the rotation of Asp394 from the external solvent into the active site that resulted in disruption of the active site alpha helix. There is therefore a direct link between the third zinc binding site and the flexible loop, which has substantial implications for catalysis. To investigate the potential impact of the loop rearrangement on the catalytic activity of *Pf*A-M17, we superposed the previously observed active *Pf*A-M17 conformation with the occluded conformation characterised here. The overlay shows that in the occluded conformation, Lys386 occupies approximately the same space in the neighbouring binding pocket that was previously occupied by the original Lys386 conformation (Fig. 4B and 4D). Therefore, within the trimer, the two conformations are incompatible, and one

chain would not be expected to adopt the active conformation whilst its neighbour is in the occluded conformation. Further, overlay of *Pf*A-M17 in complex with substrate-analog bestatin, shows that the occluded loop conformation occupies the bestatin-binding space. Based on this, we would not expect the occluded conformation to be capable of binding substrate. To confirm this hypothesis, we attempted to soak bestatin into the occluded crystal form; bestatin was never observed to bind. We therefore propose that the occluded conformation of *Pf*A-M17 is unable to bind substrate, and conclude that we have captured an inactive conformation of *Pf*A-M17.

Importantly, the crystal structure of occluded *Pf*A-M17 is consistent with our MD simulations, although the MD simulation showed the loop rearrangement in only two active sites (one of each trimer). The major difference between the MD and crystallographic 'inactive' conformations is the rearrangement of the catalytic zinc ions, which is a key component of the crystallographic conformational change. However, the hybrid 'bonded / non-bonded approach' that was used to model the metal centre in our simulations, required us to fix the positions of the zinc ions (Yang et al., 2017). Since the novel zinc coordination site is directly linked to the flexible loop through Asp394, and the MD simulations, which do not allow zinc rearrangement, show only a partial loop conformational change, we suggest that zinc movement is necessary for the complete rearrangement of the active conformation into the inactive conformation.

The dynamic loop is key to *Pf*A-M17 catalysis

Our MD and crystallographic studies identified a flexible loop in *Pf*A-M17, which has important mechanistic implications. Therefore, to confirm that the characterised conformational changes occur as part of the catalytic mechanism, we sought to (1) probe the role of key loop residue, Lys386, (2) examine how loop flexibility affects catalysis, and (3), determine the functional role of the third zinc binding site. For this purpose, we used site directed mutagenesis to alter key residues, and measured their activity using our fluorescence-based activity assay.

Lys386 has a potential role in the *Pf*A-M17 catalytic mechanism, but is also a key player in

the identified loop re-arrangement. To probe the importance of Lys386 to catalysis, we generated

*Pf*A-M17(K386A). *Pf*A-M17(K386A) showed substantially reduced activity compared to the wild

type enzyme (Table 1). This retardation of activity is due to a decrease in the ability of the enzyme

to accelerate the reaction (decrease in k_{cat}), as opposed to reduced substrate binding (K_M is largely

unchanged). Although this result demonstrates that Lys386 is key to *Pf*A-M17 function, it does not

discriminate between a role in the catalytic reaction versus the loop rearrangement. To specifically

examine the role of loop conformational dynamics in catalysis, we introduced a proline in place of

Ala387, a loop residue that does not form interactions in either the active or occluded structures.

*Pf*A-M17(A387P), similarly to *Pf*A-M17(K386A), showed substantially reduced catalytic activity

(Table 1). Large concentrations of enzyme were required to measure enzyme activity, which

manifested as reduced product turnover (decreased k_{cat}). Therefore, flexibility of the loop that links

the active sites of *Pf*A-M17 is clearly important to catalytic function.

The presence of a third zinc binding site in *Pf*A-M17 was surprising since, to our

knowledge, an equivalent site has never been observed in any M17 aminopeptidase. We were

therefore curious to examine potential functional roles for the site. In the active conformation of

*Pf*A-M17, Asp394 has no clear role. While close to the active site, the side chain is directed to

solvent and makes no direct interactions. Therefore, we disrupted the ability of site 3 to coordinate

zinc with *Pf*A-M17(D394A). The effect of this mutation on enzyme activity was profound and

unexpected. Rather than a reduction in activity observed for the previous mutations, *Pf*A-

M17(D394A) showed greatly increased catalytic ability. Again, this resulted from an altered rate

(increased k_{cat}) rather than substrate binding affinity (K_M unchanged, Table 1). Therefore,

preventing the rearrangement of the catalytic zinc ions by removal of zinc binding site 3 greatly

increases the catalytic efficiency of *Pf*A-M17.

Table 1. Kinetic analysis of mutant *Pf*A-M17 enzymes.
387

	[E] (nM)	$K_M \pm$ S.E.M (μM)	$k_{cat} \pm$ S.E.M. (FU/min)
Wild type	150	22.8 ± 1.6	27 ± 1
*Pf*A-M17(K386A)	1000	21.6 ± 4.2	4.9 ± 0.3
*Pf*A-M17(A387P)	1000	34.8 ± 4.4	7.7 ± 0.3
*Pf*A-M17(D394A)	20	17.3 ± 0.2	164 ± 6

A dynamic, cooperative mechanism for regulation of *Pf*A-M17 catalysis

We have demonstrated that the characterised loop motion linking the *Pf*A-M17 active sites has an important role in catalysis, but what is that role? For proteases, regulation of the catalytic turnover rate is crucial, particularly under conditions in which substrate and/or product concentrations are variable, or aberrant proteolytic activity is undesirable. The loop movement is coupled with the rearrangement of the active site zinc ions, which we propose is key to the mechanism. We therefore suggest that the inactive conformation we have characterised is the 'off' switch in a dynamic regulatory mechanism. Based on current evidence, there are two potential mechanisms by which the loop motion might regulate activity: (1) concerted activation, wherein all chains transition between the active and inactive states concurrently, or (2) sequential activation, where only one active site of each trimer is active at any one time, and turnover in one site triggers the neighbouring site in a continual cycle. A mechanism of concerted activation is supported by our crystal structures, which show that all six chains can adopt the active or inactive conformations concurrently. Our MD simulations however, show only a single chain of each trimer initiating the loop motion. Further, in a concerted activation mechanism, we would expect kinetic assays to show cooperativity similarly to *Hp*-M17, which possesses a Hill coefficient of 2.3 (Dong et al., 2005). Detection of cooperativity by the Hill coefficient relies on the assumption that ligand molecules bind simultaneously; therefore, a Hill coefficient of 1.0 for *Pf*A-M17 supports a sequential mechanism.

Our current data lend support to both concurrent and sequential activation mechanisms. Although it is possible that we have not yet identified the specific conditions to differentiate between the two mechanisms, we propose the more likely scenario, is that *Pf*A-M17 functions by a combination of the two (Fig. 5). In this mechanism, the enzyme is capable of sampling three major states: (1) an inactive state wherein all chains adopt the occluded conformation and no catalysis occurs (Fig. 5A), (2) an active state which allows catalysis to occur in all active sites concurrently (Fig. 5C), and (3) an intermediate, or sequential activation state, which is moderated by the

regulatory loop to allow proteolysis in only one active site per trimeric unit (Fig. 5B). This 'combined' mechanism would allow exceptionally fine control of catalytic rate when fast (active state), moderate (sequential activation state), or no (inactive state) proteolysis is required. Dynamic ensembles are known to play an important role in enzyme catalysis (Boehr, Nussinov, & Wright, 2009; Nagel & Klinman, 2009). We propose a dynamic system in which the conformational space available to *Pf*A-M17 is influenced by the presence of specific environmental signals that alter the conformational energy landscape to favour certain activation states depending on the rate of proteolysis required. Candidates for the signal molecules that regulate the rate of catalysis include: substrate/product, co-factor, or external signal molecule. While *Hp*-M17 identified an allosteric link between substrate concentration and activity, we were unable to find a substrate-based link for *Pf*A-M17 using a similar substrate. However, M17 aminopeptidases from different organisms show altered activity when different metal ions are bound (Allen et al., 1983; Cappiello et al., 2006; Carroll et al., 2013; Maric et al., 2009; Zhu et al., 2012). For this reason, many studies have focused on examination of the identity and role of the two metal ions; however, have never considered that the positions themselves might be dynamic and playing a regulatory role. Our studies described herein have shown that re-arrangement of active site metal ions play a key role in the mechanism of *Pf*A-M17. The metal co-factor is therefore a prime candidate to act as a signalling factor in the regulation of *Pf*A-M17 activity. The concentrations of different divalent metal cations are known to fluctuate throughout the parasite life cycle (Marvin et al., 2012). Therefore, environmental metal ion concentration could be serving to moderate *Pf*A-M17 activity according to different metabolic demands throughout the complex parasite life cycle.

Figure 5. Schematic representation of the proposed *Pf*A-M17 regulatory mechanism. In the proposed regulatory mechanism, each trimer operates independently (trimer *abc* is shown with *a* in yellow, *b* in pink, and *c* in cyan). Environmental conditions alter the conformational space to favour (A) an inactive state in which proteolysis does not occur, (B) an intermediate state that is capable of hydrolysing substrate in one active site only, and (C) an active state wherein all sites are active concurrently.

Divergence of M17 function between species

M17 aminopeptidases have vital roles in a wide range of physiological processes throughout all kingdoms of life. We therefore sought to determine whether the functionally important communication between the active sites is a conserved feature of the conserved hexameric assembly of M17 aminopeptidases. We performed a sequence alignment of M17 aminopeptidases from a range of different organisms and compared the region of the flexible loop (SI 7). The alignment showed that while Lys386 is highly conserved, neither the flexible loop nor the aspartic acid residue that defines the third zinc binding site are conserved. Closer examination showed that the loop regions could be classified into two groups; those with loops of similar length and nature to *Pf*A-M17 as well as zinc site 3 (Asp or Glu), and those that have a shortened loop, often coupled with increased proline content and absence of zinc site 3 (though Asp394 is still present in some of these enzymes). Due to the predicted increase in rigidity and decreased length of the loop in this latter group, we speculate that these enzymes are incapable of the loop dynamics described for *Pf*A-M17 here. Interestingly *Hp*-M17, the only M17 aminopeptidase of our knowledge to have identified cooperativity of substrate binding (Dong et al., 2005), fits into the latter class. The flexible loop, and the dynamic regulatory mechanism that it moderates, might therefore be key to explaining the differences between the mechanisms of *Hp*-M17 and *Pf*A-M17. It has been suggested that flexible loops in proteins can facilitate the emergence of novel functions, which results in divergent evolution (Bhabha et al., 2013; Campbell et al., 2017; Tokuriki & Tawfik, 2009). The dynamic regulatory mechanism that we have described is therefore a prime candidate for role of mediating different functionalities throughout the M17 enzyme family.

Conclusion:

The M17 aminopeptidases have been of interest as key players in a wide range of physiological processes for over 50 years. From a protein mechanics perspective, the conserved hexameric assembly containing six discrete active sites, represents an exciting system in which to

investigate the role of dynamic cooperativity within a large oligomer. Herein, we have shown that

the hexameric assembly is absolutely essential for the proteolytic activity of *Pf*A-M17. The

arrangement stabilises and preserves key catalytic machinery, and further, regulates access routes to

the catalytic cavity through movements in surface loops that moderate the pore size of key

channels. Within the hexamer, each of the two disc-like trimers operate independently; a flexible

loop links each of the three active sites to its neighbour, thereby operating cooperatively to convert

the enzyme between inactive and active states. Further, movement of the regulatory loop is coupled

with a rearrangement of active site metal cofactors. Our studies show that not only do metal ion

dynamics exist, but that the positions of the metals are manipulated to control the activity of the

enzyme. We therefore propose a dynamic regulatory mechanism for *Pf*A-M17 that is moderated by

changes in the physiological metal environment. Characterisation of the function, and regulation of

that function, described herein for *Pf*A-M17 thereby provides insight into how the M17

aminopeptidases can operate in such varying capacities throughout all kingdoms of life.

Materials and Methods:

Bacterial strains and plasmids.

The cloning of the truncated *Pf*A-M17 gene (encoding amino acids 85 – 605) into

pTrcHis2B has been previously reported (McGowan et al., 2010). Site directed mutagenesis was

performed by PCR and confirmed by DNA sequencing. *Escherichia coli* strains DH5α and K-12

were used for DNA manipulation, and BL21(DE3) used for protein expression. Recombinant His$_6$-

tagged wild type and mutant *Pf*A-M17 were expressed using an autoinduction method as previously

described for *wild type Pf*A-M17 (McGowan et al., 2010).

Protein purification and analysis by gel filtration chromatography.

Proteins were purified as has previously been described for wild type *Pf*A-M17 (McGowan

et al., 2010) using a two-step purification procedure of Ni–NTA agarose column followed by gel

filtration chromatography on a Superdex S200 10/300 column in 50 mM HEPES pH 8.0, 150 mM

NaCl buffer. Analytical gel filtration chromatography was performed on a Superdex S200 Increase

10/300 in 50 mM HEPES pH 8.0, 150 mM NaCl. Approximate molecular weight of eluate was

calculated by interpolation of a standard curve constructed with appropriate molecular weight

standards.

Aminopeptidase assays and analysis.

Aminopeptidase activity was determined by measuring the release of the fluorogenic-

leaving group, L-Leucine-7-amido-4-methylcoumarin hydrochloride (Sigma-Aldrich, L2145)

(NHMec), from the fluorogenic peptide substrates H-Leu-NHMec as described previously (Stack et

al., 2007). Briefly, assays were performed in triplicate and carried out in 50 μL total volume in 100

mM Tris-HCl, pH 8.0, 1 mM $MnCl_2$ at 37 °C and activity was monitored until steady-state was

achieved. Fluorescence was measured using a FluoroStar Optima plate reader (BMG Labtech), with

excitation and emission wavelengths of 355 nm and 460 nm, respectively. Activity of the

association mutant PfA-M17(W525A,Y533A) was assessed with 10 μM H-Leu-NH-Mec, up to an

enzyme concentration of 1000 nM with no detectable activity.

For active enzymes, the Michaelis constant, K_M, was calculated from the initial rates over a

range of substrate concentrations (0.5 – 500 μM) with enzyme concentrations fixed at 150 nM for

the *wild type Pf*A-M17, 20 nM for PfA-M17(D394A), and 1000 nM for PfA-M17(A387P) and PfA-

M17(K386A). Gain was fixed at 800 for all Michaelis-Menten assays. Kinetic parameters

including, K_M, k_{cat}, and n_H were calculated with non-linear regression protocols by using GraphPad

Prism 7.

Crystallisation, data collection, structure determination and refinement.

PfA-M17 was concentrated to 10 mg/mL in 50 mM HEPES pH 8.0, 150 mM NaCl for

crystallisation. Crystals were grown by hanging drop vapour diffusion, in 20 % PEG3350, 0.2 M

calcium acetate, with drops composed of 2 μL protein plus 1 μL precipitant. Crystals grew to large

plates in 7 days. For soaking experiments, crystals were transferred to fresh drop composed of

crystallisation solution supplemented with 1 mM of bestatin (Sigma-Aldrich, B8385) for 24 hours

prior to cryoprotection. Crystals were cryoprotected in mother liquor supplemented with 15 % 2-

Methyl-2,4-pentanediol for 30 s before flash cooling in liquid nitrogen.

Data were collected at 100 K using synchrotron radiation at the Australian Synchrotron

using the micro crystallography MX2 beamline 3ID1. Data were collected from two wedges of the

same crystal, which were merged after integration. Data were processed using iMosflm (Battye,

Kontogiannis, Johnson, Powell, & Leslie, 2011) and Aimless (Evans & Murshudov, 2013) as part

of the CCP4i suite (Winn et al., 2011). The structure was solved by molecular replacement in

Phaser using the structure of unliganded _Pf_A-M17 (RCSB ID 3KQZ) as the search model

(McGowan et al., 2010). Refinement was carried out with iterative rounds of model building in

Coot (Emsley, Lohkamp, Scott, & Cowtan, 2010) and refinement using Phenix (Adams et al.,

2010). Water molecules and metal ions were placed manually based on the presence of $F_o–F_c$ and

$2F_o–F_c$ electron density of appropriate signal. The structure was validated with MolProbity (Chen et

al., 2010) and figures generated using PyMOL version 1.8.23. Final structure coordinates were

deposited in the protein databank (RCSB ID 6BVO), and a summary of data collection and

refinement statistics is provided in Supplementary Information 8. For clarity and consistency, the

chains in the deposited structure (PDB ID 6VBO) are numbered according to the model of

unliganded _Pf_A-M17 (PDB ID 3KQZ), whereby chain _a_, _b_, and _c_ are equivalent to A, B, and C

respectively, and _a_*, _b_*, _c_* are equivalent to D, F, and E respectively.

MD system setup and simulation protocol.

The starting _Pf_A-M17 model for MD simulations was based on the unliganded crystal

structure, 3KQZ. To prepare the model, missing atoms and residues ($a^{84, \ 257-261}$, $b^{84-85, \ 255-262}$, $c^{84-85,}$

$^{255-259}$, $a*^{84, \ 255-259}$, $b*^{84-85, \ 136, \ 255-261}$, $c*^{84-85, \ 152}$) were rebuilt using Modeller v9.11 (Fiser & Sali,

2003), protonated according to their states at pH 7.0 using the PDB2PQR server (Dolinsky et al., 2007) and subjected to energy minimisation using Modeller (Fiser & Sali, 2003). The hexameric M17 system consisted of ~ 294 000 atoms with a periodic box of 167 Å × 166 Å × 121 Å and the monomeric M17 system consisted of ~ 81,000 atoms with a periodic box of 120Å × 85 Å × 94 Å (after solvation). Periodic boundary conditions (pbc) were used at all simulations. System charges were neutralised with sodium counter ions. Proteins and ions were modelled using the AMBER force field FF12SB (Case et al., 2005) , the metal centre was defined as described previously (Yang et al., 2017) and waters represented using the 3-particle TIP3P model (Jorgensen, Chandrasekhar, Madura, Impey, & Klein, 1983). All atom MD simulations were performed using NAMD 2.9 on an IBM Blue Gene/Q cluster (monomer simulations) or x86 (hexamer simulations). Equilibration was performed in three stages. First, potential steric clashes in the initial configuration were relieved with 50000 steps of energy minimization. Initial velocities for each system were then assigned randomly according to a Maxwell–Boltzmann distribution at 100 K. Each system was then heated to 300 K over 0.1 ns, under the isothermal-isometric ensemble (NVT) conditions, with the protein atoms (excluding hydrogens) harmonically restrained (with a force constant of 10 kcal mol^{-1} A^{-2}). Following this, each system was simulated for 100 ps under the isothermal-isobaric ensemble (NPT) with heavy atoms restrained. The harmonic restrained used were reduced from 10 to 2 kcal mol^{-1} A^{-2} during the simulations. The above equilibration process was performed three times from the same starting structure in order to initiate three production simulations with different initial velocities. For production simulations, the time step was set to 2 fs and the SHAKE algorithm was used to constrain all bonds involving hydrogen atoms. All simulations were run at constant temperature (300 K) and pressure (1 atm), using a Langevin damping coefficient of 0.5 fs^{-1}, and a Berendsen thermostat relaxation time of τ_P = 0.1 ps. The Particle-Mesh Ewald (PME) method was used to set the periodic boundary conditions (PBC) that were used for long-range electrostatic interactions and a real space cut-off of 10 Å was used. Conformations were sampled every 10 ps for subsequent analysis. All frames with time interval of 10 ps were saved to disk.

MD Analysis.

Simulation trajectories were analysed using the GROMACS 5.14 simulation package. For

principle component analysis (PCA), 3N*3N atom covariance matrices of the protein displacement

in simulations were generated based on backbone atoms (N, Cα, C, O) of the *Pf*A-M17 crystal

structure. Principle Components (PCs), that taken together accounted for more than 50% of the

overall covariance, were chosen for essential dynamics analysis. The GROMACS 5.14 simulation

package was used to project the trajectory onto the top PCs. Graphs and plots were produced with

Xmgrace and GraphPad Prism7. Molecular graphics were prepared with PyMOL 1.8.23 and

VMD1.9.3.

Channel analysis.

The channels to the interior of the *Pf*A-M17 hexamer were examined using Caver

(Chovancova et al., 2012). Channel dimensions were assessed at the start of the simulation, which

represented the crystallographic conformation of *Pf*A-M17 after the missing loop residues were

modelled and subjected to energy minimisation, and the end of the simulation. Values reported are

for the channel 'bottleneck', which represent the diameter of each channel at the narrowest point. If

multiple forks off a single channel were observed, the dimensions for the larger channel (which

represent the most likely access points) were reported. Figures of channels were created using

Caver (Chovancova et al., 2012) and visualised using the PyMOL plugin.

Sequence alignment and analysis.

ClustalOmega (https://www.ebi.ac.uk/Tools/msa/clustalo/) was used to align sequences of

M17 aminopeptidases (*C. elegans* P34629.1; *B. taurus* AAB28170.1; *H. sapiens* AAD17527.1, *M.*

musculus AAK13495.1; *R. norvegicus* AAH79381.1; *D. melanogaster* AAF50390.2; *A. gambiae*

XP_321111.3; *A. thaliana* Q944P7; *S. Lycopersicum* (LAP2) XP_006350101.1; *S. Lycopersicum*

(LAP1) NP_001233862.2; *S. tuberosum* CAA48038.1; *T. annulata* CAI76586.1; *T. gondii*

XP_018636441.1; *P. falciparum* XP_001348613.1; *C. hominis* XP_667960.1; *C. parvum*

XP_626197.1; *L. major* AAL16097.1; *H. pylori* WP_064816828.1; *S. aureus* WP_075108680.1; *E.*

coli P68767; *R. typhi* AAU03616.1; *C. tetani* WP_011100044.1; *B. cereus* AAP11794.1).

Abbreviations:

*Pf*A-M17, *Plasmodium falciparum* M17 aminopeptidase*; Hp*-M17, *Helicobacter pylori*

M17 aminopeptidase; MD, molecular dynamics; PCA, principle component analysis; RDF, radial

distribution function.

Author Contributions:

S.M. and N.D. designed the research; N.D., W.Y., K.K.S., B.T.R., and I.K. performed the

research and analysed the data; A.M.B. and S.M. supervised the research; N.D. and S.M. wrote the

manuscript with contributions from all other authors.

Acknowledgements:

We thank the National Health and Medical Research Council (Project Grant 1063786 to S.M. and

P.J.S.; Fellowship 1022688 to A.M.B.) and the Australian Research Council (Fellowship

FT100100690 to S.M.) for funding support. This work was supported by the Victorian Life

Sciences Computation Initiative (VLSCI) and National Computational Infrastructure (NCI). This

research was undertaken in part using the MX2 beamline at the Australian Synchrotron, part of

ANSTO.

References:

Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J., Hung, L.-W., Kapral, G. J., Grosse-Kunstleve, R. W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D., 66*, 213-221. doi: 10.1107/s0907444909052925

Alén, C., Sherratt, D. J., & Colloms, S. D. (1997). Direct interaction of aminopeptidase A with recombination site DNA in Xer site-specific recombination. *EMBO J., 16*(17), 5188-5197. doi: 10.1093/emboj/16.17.5188

Allen, M. P., Yamada, A. H., & Carpenter, F. H. (1983). Kinetic-parameters of metal-substituted leucine aminopeptidase from bovine lens. *Biochemistry, 22*(16), 3778-3783. doi: 10.1021/bi00285a010

Battye, T. G. G., Kontogiannis, L., Johnson, O., Powell, H. R., & Leslie, A. G. W. (2011). iMOSFLM: a new graphical interface for diffraction-image processing with MOSFLM. *Acta Crystallogr. D., 67*, 271-281. doi: 10.1107/s0907444910048675

Bhabha, G., Ekiert, D. C., Jennewein, M., Zmasek, C. M., Tuttle, L. M., Kroon, G., Dyson, H. J., Godzik, A., Wilson, I. A., & Wright, P. E. (2013). Divergent evolution of protein conformational dynamics in dihydrofolate reductase. *Nature Structural & Molecular Biology, 20*(11), 1243-U1262. doi: 10.1038/nsmb.2676

Bhabha, G., Lee, J., Ekiert, D. C., Gam, J., Wilson, I. A., Dyson, H. J., Benkovic, S. J., & Wright, P. E. (2011). A Dynamic Knockout Reveals That Conformational Fluctuations Influence the Chemical Step of Enzyme Catalysis. *Science, 332*(6026), 234-238. doi: 10.1126/science.1198542

Boehr, D. D., Nussinov, R., & Wright, P. E. (2009). The role of dynamic conformational ensembles in biomolecular recognition. *Nature Chemical Biology, 5*(11), 789-796. doi: 10.1038/nchembio.232

Burley, S. K., David, P. R., Taylor, A., & Lipscomb, W. N. (1990). Molecular structure of leucine aminopeptidase at 2.7 Å resolution *Proc. Natl. Acad. Sci., 87*(17), 6878-6882. doi: 10.1073/pnas.87.17.6878

Campbell, E. C., Correy, G. J., Mabbitt, P. D., Buckle, A. M., Tokuriki, N., & Jackson, C. J. (2017). Laboratory evolution of protein conformational dynamics. *Curr Opin Struct Biol, 50*, 49-57. doi: 10.1016/j.sbi.2017.09.005

Cappiello, M., Alterio, V., Amodeo, P., Del Corso, A., Scaloni, A., Pedone, C., Moschini, R., De Donatis, G. M., De Simone, G., & Mura, U. (2006). Metal ion substitution in the catalytic site greatly affects the binding of sulfhydryl-containing compounds to leucyl aminopeptidase. *Biochemistry, 45*(10), 3226-3234. doi: 10.1021/bi052069v

Carroll, R. K., Veillard, F., Gagne, D. T., Lindenmuth, J. M., Poreba, M., Drag, M., Potempa, J., & Shaw, L. N. (2013). The Staphylococcus aureus leucine aminopeptidase is localized to the bacterial cytosol and demonstrates a broad substrate range that extends beyond leucine. *Biol. Chem., 394*(6), 791-803. doi: 10.1515/hsz-2012-0308

Case, D. A., Cheatham, T. E., Darden, T., Gohlke, H., Luo, R., Merz, K. M., Onufriev, A., Simmerling, C., Wang, B., & Woods, R. J. (2005). The Amber biomolecular simulation programs. *J. Comput. Chem., 26*(16), 1668-1688. doi: 10.1002/jcc.20290

Charlier, D., Hassanzadeh, G., Kholti, A., Gigot, D., Pierard, A., & Glansdorff, N. (1995). *carP*, Involved in Pyrimidine Regulation of the *Escherichia coli* Carbamoylphosphate Synthetase Operon Encodes a Sequence-specific DNA-binding Protein Identical to XerB and PepA, also Required for Resolution of ColEl Multimers. *Journal of Molecular Biology, 250*(4), 392-406. doi: 10.1006/jmbi.1995.0385

Chen, V. B., Arendall, W. B., Headd, J. J., Keedy, D. A., Immormino, R. M., Kapral, G. J., Murray, L. W., Richardson, J. S., & Richardson, D. C. (2010). MolProbity: all-atom structure

validation for macromolecular crystallography. *Acta Crystallogr. D., 66*, 12-21. doi:
10.1107/s0907444909042073

Chovancova, E., Pavelka, A., Benes, P., Strnad, O., Brezovsky, J., Kozlikova, B., Gora, A., Sustr, V.,
Klvana, M., Medek, P., Biedermannova, L., Sochor, J., & Damborsky, J. (2012). CAVER
3.0: A Tool for the Analysis of Transport Pathways in Dynamic Protein Structures. *Plos*
Comput. Biol., 8(10). doi: 10.1371/journal.pcbi.1002708

Dalal, S., & Klemba, M. (2007). Roles for two aminopeptidases in vacuolar hemoglobin
catabolism in Plasmodium falciparum. *J. Biol. Chem., 282*(49), 35978-35987. doi:
10.1074/jbc.M703643200

Dolinsky, T. J., Czodrowski, P., Li, H., Nielsen, J. E., Jensen, J. H., Klebe, G., & Baker, N. A. (2007).
PDB2PQR: expanding and upgrading automated preparation of biomolecular
structures for molecular simulations. *Nucleic Acids Res., 35*, W522-W525. doi:
10.1093/nar/gkm276

Dong, L., Cheng, N., Wang, M. W., Zhang, J. F., Shu, C., & Zhu, D. X. (2005). The leucyl
aminopeptidase from Helicobacter pylori is an allosteric enzyme. *Microbiology, 151*,
2017-2023. doi: 10.1099/mic.0.27767-0

Drinkwater, N., Bamert, R. S., Sivaraman, K. K., Paiardini, A., & McGowan, S. (2015). X-ray
crystal structures of an orally available aminopeptidase inhibitor, Tosedostat, bound to
anti-malarial drug targets PfA-M1 and PfA-M17. *Proteins, 83*(4), 789-795. doi:
10.1002/prot.24771

Drinkwater, N., Vinh, N. B., Mistry, S. N., Bamert, R. S., Ruggeri, C., Holleran, J. P., Loganathan, S.,
Paiardini, A., Charman, S. A., Powell, A. K., Avery, V. M., McGowan, S., & Scammells, P. J.
(2016). Potent dual inhibitors of Plasmodium falciparum M1 and M17
aminopeptidases through optimization of S1 pocket interactions. *Eur. J. Med. Chem.,*
110, 43-64. doi: 10.1016/j.ejmech.2016.01.015

Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010). Features and development of Coot.
Acta Crystallogr. D., 66, 486-501. doi: 10.1107/s0907444910007493

Evans, P. R., & Murshudov, G. N. (2013). How good are my data and what is the resolution?
Acta Crystallogr. D., 69, 1204-1214. doi: 10.1107/s0907444913000061

Fiser, A., & Sali, A. (2003). MODELLER: Generation and refinement of homology-based protein
structure models. In C. W. Carter & R. M. Sweet (Eds.), *Methods Enzymol.* (Vol. 374, pp.
461-491).

Gerstein, M., & Chothia, C. (1996). Packing at the protein-water interface. *Proceedings of the*
National Academy of Sciences of the United States of America, 93(19), 10167-10172. doi:
10.1073/pnas.93.19.10167

Harbut, M. B., Velmourougane, G., Dalal, S., Reiss, G., Whisstock, J. C., Onder, O., Brisson, D.,
McGowan, S., Klemba, M., & Greenbaum, D. C. (2011). Bestatin-based chemical biology
strategy reveals distinct roles for malaria M1-and M17-family aminopeptidases. *Proc.*
Natl. Acad. Sci., 108(34), E526-E534. doi: 10.1073/pnas.1105601108

Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W., & Klein, M. L. (1983).
Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.,*
79(2), 926-935. doi: 10.1063/1.445869

Kumar, S., Kaur, A., Chattopadhyay, B., & Bachhawat, A. K. (2015). Defining the cytosolic
pathway of glutathione degradation in Arabidopsis thaliana: role of the ChaC/GCG
family of gamma-glutamyl cyclotransferases as glutathione-degrading enzymes and
AtLAP1 as the Cys-Gly peptidase. *Biochemical Journal, 468*, 73-85. doi:
10.1042/bj20141154

Lowther, W. T., & Matthews, B. W. (2002). Metalloaminopeptidases: Common functional
themes in disparate structural surroundings. *Chem. Rev., 102*(12), 4581-4607. doi:
10.1021/cr0101757

Maric, S., Donnelly, S. M., Robinsin, M. W., Skinner-Adams, T., Trenholme, K. R., Gardiner, D. L., Dalton, J. P., Stack, C. M., & Lowther, J. (2009). The M17 Leucine Aminopeptidase of the Malaria Parasite Plasmodium falciparum: Importance of Active Site Metal Ions in the Binding of Substrates and Inhibitors. *Biochemistry, 48*(23), 5435-5439. doi: 10.1021/bi9003638

Marvin, R. G., Wolford, J. L., Kidd, M. J., Murphy, S., Ward, J., Que, E. L., Mayer, M. L., Penner-Hahn, J. E., Haldar, K., & O'Halloran, T. V. (2012). Fluxes in "Free" and Total Zinc Are Essential for Progression of Intraerythrocytic Stages of Plasmodium falciparum. *Chem. Biol., 19*(6), 731-741. doi: 10.1016/j.chembiol.2012.04.013

Matsui, M., Fowler, J. H., & Walling, L. L. (2006). Leucine aminopeptidases: diversity in structure and function. *Biol. Chem., 387*(12), 1535-1544. doi: 10.1515/bc.2006.191

McElheny, D., Schnell, J. R., Lansing, J. C., Dyson, H. J., & Wright, P. E. (2005). Defining the role of active-site loop fluctuations in dihydrofolate reductase catalysis. *Proc. Natl. Acad. Sci., 102*(14), 5032-5037. doi: 10.1073/pnas.0500699102

McGowan, S., Oellig, C. A., Birru, W. A., Caradoc-Davies, T. T., Stack, C. M., Lowther, J., Skinner-Adams, T., Mucha, A., Kafarski, P., Grembecka, J., et al. (2010). Structure of the Plasmodium falciparum M17 aminopeptidase and significance for the design of drugs targeting the neutral exopeptidases. *Proc. Natl. Acad. Sci., 107*(6), 2449-2454. doi: 10.1073/pnas.0911813107

Minh, P. N. L., Devroede, N., Massant, J., Maes, D., & Charlier, D. (2009). Insights into the architecture and stoichiometry of Escherichia coli PepADNA complexes involved in transcriptional control and site-specific DNA recombination by atomic force microscopy. *Nucleic Acids Research, 37*(5), 1463-1476. doi: 10.1093/nar/gkn1078

Mistry, S. N., Drinkwater, N., Ruggeri, C., Sivaraman, K. K., Loganathan, S., Fletcher, S., Drag, M., Paiardini, A., Avery, V. M., Scammells, P. J., & McGowan, S. (2014). Two-Pronged Attack: Dual Inhibition of Plasmodium falciparum M1 and M17 Metalloaminopeptidases by a Novel Series of Hydroxamic Acid-Based Inhibitors. *J. Med. Chem., 57*(21), 9168-9183. doi: 10.1021/jm501323a

Nagel, Z. D., & Klinman, J. P. (2009). A 21(st) century revisionist's view at a turning point in enzymology. *Nat. Chem. Biol., 5*(8), 543-550. doi: 10.1038/nchembio.204

Rawlings, N. D., Barrett, A. J., & Finn, R. (2016). Twenty years of the MEROPS database of proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Res., 44*(D1), D343-D350. doi: 10.1093/nar/gkv1118

Reboul, C. F., Porebski, B. T., Griffin, M. D. W., Dobson, R. C. J., Perugini, M. A., Gerrard, J. A., & Buckle, A. M. (2012). Structural and Dynamic Requirements for Optimal Activity of the Essential Bacterial Enzyme Dihydrodipicolinate Synthase. *Plos Comput. Biol., 8*(6). doi: 10.1371/journal.pcbi.1002537

Rosenthal, P. J. (2002). Hydrolysis of erythrocyte proteins by proteases of malaria parasites. *Curr. Opin. Hematol., 9*(2), 140-145. doi: 10.1097/00062752-200203000-00010

Schurer, G., Horn, A. H. C., Gedeck, P., & Clark, T. (2002). The reaction mechanism of bovine lens leucine aminopeptidase. *J. Phys. Chem. B, 106*(34), 8815-8830. doi: 10.1021/jp025575s

Scranton, M. A., Yee, A., Park, S. Y., & Walling, L. L. (2012). Plant Leucine Aminopeptidases Moonlight as Molecular Chaperones to Alleviate Stress-induced Damage. *J. Biol. Chem., 287*(22), 18408-18417. doi: 10.1074/jbc.M111.309500

Skinner-Adams, T. S., Lowther, J., Teuscher, F., Stack, C. M., Grembecka, J., Mucha, A., Kafarski, P., Trenholme, K. R., Dalton, J. P., & Gardiner, D. L. (2007). Identification of phosphinate dipeptide analog inhibitors directed against the Plasmodium falciparum M17 leucine aminopeptidase as lead antimalarial compounds. *J. Med. Chem., 50*(24), 6024-6031. doi: 10.1021/jm070733v

Skinner-Adams, T. S., Peatey, C. L., Anderson, K., Trenholme, K. R., Krige, D., Brown, C. L., Stack,
C., Nsangou, D. M. M., Mathews, R. T., Thivierge, K., Dalton, J. P., & Gardinerd, D. L.
(2012). The Aminopeptidase Inhibitor CHR-2863 Is an Orally Bioavailable Inhibitor of
Murine Malaria. *Antimicrob. Agents Ch., 56*(6), 3244-3249. doi: 10.1128/aac.06245-11
Stack, C. M., Lowther, J., Cunningham, E., Donnelly, S., Gardiner, D. L., Trenholme, K. R.,
Skinner-Adams, T. S., Teuscher, F., Grembecka, J., Mucha, A., Kafarski, P., Lua, L., Bell, A.,
& Dalton, J. P. (2007). Characterization of the Plasmodium falciparum M17 leucyl
aminopeptidase - A protease involved in amino acid regulation with potential for
antimalarial drug development. *J. Biol. Chem., 282*(3), 2069-2080. doi:
10.1074/jbc.M609251200
Stirling, C. J., Colloms, S. D., Collins, J. F., Szatmari, G., & Sherratt, D. J. (1989). xerB, an
Escherichia coli gene required for plasmid ColE1 site-specific recombination, is
identical to pepA, encoding aminopeptidase A, a protein with substantial similarity to
bovine lens leucine aminopeptidase. *Embo Journal, 8*(5), 1623-1627.
Sträter, N., & Lipscomb, W. N. (1995). 2-Metal ion mechanism of bovine lens leucine
aminopeptidase - active site solvent structure and binding mode of L-leucinal, a gem-
diolate transition-state analog, by X-ray crystallography. *Biochemistry, 34*(45), 14792-
14800. doi: 10.1021/bi00045a021
Sträter, N., Sherratt, D. J., & Colloms, S. D. (1999). X-ray structure of aminopeptidase A from
Escherichia coli and a model for the nucleoprotein complex in Xer site-specific
recombination. *EMBO J., 18*(16), 4513-4522. doi: 10.1093/emboj/18.16.4513
Tokuriki, N., & Tawfik, D. S. (2009). Protein Dynamism and Evolvability. *Science, 324*(5924),
203-207. doi: 10.1126/science.1169375
Vögeli, B., Bibow, S., & Chi, C. N. (2016). Enzyme Selectivity Fine-Tuned through Dynamic
Control of a Loop. *Angewandte Chemie-International Edition, 55*(9), 3096-3100. doi:
10.1002/anie.201511476
Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., Keegan, R. M.,
Krissinel, E. B., Leslie, A. G. W., McCoy, A., et al. (2011). Overview of the CCP4 suite and
current developments. *Acta Crystallogr. D., 67*, 235-242. doi:
10.1107/s0907444910045749
Wurm, J. P., Holdermann, I., Overbeck, J. H., Mayer, P. H. O., & Sprangers, R. (2017). Changes in
conformational equilibria regulate the activity of the Dcp2 decapping enzyme. *Proc.
Natl. Acad. Sci., 114*(23), 6034-6039. doi: 10.1073/pnas.1704496114
Yang, W., Riley, B. T., Lei, X., Porebski, B. T., Kass, I., Buckle, A. M., & McGowan, S. (2017).
Generation of AMBER force field parameters for zinc centres of M1 and M17 family
aminopeptidases. *J. Biomol. Struct. Dyn.*, 1-10. doi: 10.1080/07391102.2017.1364669
Zhu, X. X., Barman, A., Ozbil, M., Zhang, T. T., Li, S. H., & Prabhakar, R. (2012). Mechanism of
peptide hydrolysis by co-catalytic metal centers containing leucine aminopeptidase
enzyme: a DFT approach. *J. Biol. Inorg. Chem., 17*(2), 209-222. doi: 10.1007/s00775-
011-0843-2
814

Supplementary Information 1: Analysis of the Hill coefficient (n_H) by substrate-saturation

experiment showed no evidence of cooperativity in *Pf*A-M17 catalysis. Data shown are means ± SD

($n = 3$).

Supplementary Information 2: (A) RMSD plot of hexameric *Pf*A-M17 over course of simulation.

(B) Graphical representation of PC1 (black arrows, stretching motion).

Supplementary Information 3: Table of change in pore dimensions over course of the simulation

(measurement is of the bottleneck of each of the channels as calculated by Caver).

Channel	Defined by loops	Channel diameter at bottleneck (Å)		
		Start	*End*	*Change*
1	$a^{247–266}{:}a*^{132–150}$	3.8	6.5	2.7
2	$b^{247–266}{:}b*^{132–150}$	4.0	5.9	1.9
3	$c^{247–266}{:}c*^{132–150}$	3.9	4.5	0.6
4	$a^{132–150}{:}a*^{247–266}$	4.6	6.5	1.9
5	$b^{132–150}{:}b*^{247–266}$	4.4	7.1	2.7
6	$c^{132–150}{:}c*^{247–266}$	4.9	8.9	4.0

Supplementary Information 4: *Pf*A-M17(Y525A+Y533A) is monomeric in solution as

determined by analytical size exclusion chromatography. Absorbance trace (A) shows that the

association mutant elutes at 14.2 mL, which corresponds to an approximate molecular weight of

68 kDa (*Pf*A-M17 molecular weight = 59 kDa). The mutant therefore represents monomeric *Pf*A-

M17. Assessment of the catalytic activity of *Pf*A-M17(W525A+Y533A) under *wild type*

conditions, showed that the monomeric enzyme is inactive (B).

Supplementary Information 5: MD simulations of monomer versus hexamer (A) RMSD analysis

of monomer throughout the simulation. (B) RMSF analysis of monomer throughout simulation. (C)

RMSF analysis of individual six chains in hexamer throughout simulation. Radial distribution

function (RDF) plot of water molecules in the zinc environment from the MD simulations of (D)

the monomer and (E) the hexamer. The RDF was calculated by the average of all six sites in the

hexamer. Probability percentage was calculated by the occupancy of the water divided by the

number of waters (as measured by oxygen atom of the waters) at any given distance to the zinc

ions. The plots show that the catalytic water in simulation of monomer is unstable, however, the

hexamer has a single water molecule in the catalytic position throughout the simulation.

Supplementary Information 6: Crystallographic Data Collection and Refinement Statistics

Data collection	
Space Group	P 21 21 21
Cell dimensions:	
a, b, c (Å)	172.73, 178.53, 222.79
$\alpha = \beta = \gamma$ (°)	90.0
Resolution range (Å)	48.38 – 2.30 (2.382 – 2.30)
Total Observations	2,540,053 (123,225)
Unique Observations	301,951 (14,522)
Multiplicity	8.4 (8.5)
Completeness (%)	99.4 (97.4)
$<I/\sigma I>$	3.6 (0.9)
CC(1/2)	0.987 (0.451)
R_{pim} (%)	14.9 (100)
PDB	6VBO
Structure refinement	
Non hydrogen atoms	
Protein	47,419
Water	3235
Metal (Zn)	12
Ligand	111
R_{free} (%)	23.9
R_{cryst} (%)	28.9
RMS deviations	
Bond lengths (Å)	0.002
Angles (°)	0.57
Ramachandran plot	
Favoured(%)	96.3
Outliers (%)	0.2
B factors ($Å^2$)	
Protein	37.1
Solvent	40.5
Ligand	52.3
Molprobity	
Clashscore	3.2 (100[th] percentile)
MolProbity score	1.4 (99[th] percentile)

Supplementary Information 7: Sequence alignment of verified M17 aminopeptidases from a

range of organisms performed with Clustal Omega. Conserved lysine residue (Lys386 in *Pf*A-M17)

is highlighted in green and loop residues are highlighted in pink. Yellow highlight shows that

Asp394 of *Pf*A-M17, representative of zinc binding site 3, is not conserved throughout the enzyme

family. Blue box indicates the M17 aminopeptidases that possess a loop of similar length and nature

to *Pf*A-M17.

```
C. elegans             IYHVGK-AG--PTPPAFVVLSHEVP-G-STEHIALVGKGVVYDTGGLQIKT--KTGMPNM  272
B. taurus              FLSVAK-GS--EEPPVFLEIHYKGSPNASEPPLVFVGKGITFDSGGISIKA--AANMDLM  296
H. sapiens             FLSVAK-GS--DEPPVFLEIHYKGSPNANEPPLVFVGKGITFDSGGISIKA--SANMDLM  302
M. musculus            FLSVAK-GS--EEPPVFLEIHYMGSPNATEAPLVFVGKGITFDSGGISIKA--SANMDLM  302
R. norvegicus          FLSVAK-GS--EEPPVFLEIHYTGSPNATEAPLVFVGKGITFDSGGISIKA--SANMDLM  302
D. melanogaster        FLAVGK-AS--CEPPIFLELSYYGT-CAEERPIVLVGQGITYDAGGLCLKK--KKELFNM  331
A. gambiae             FLTVAK-GS--CEPPIFLELSYYGT-NTKERPIVLIGQGNTFDSGGLCLKT--LEALTDM  315
A. thaliana            YLAVAA-AS--ANPPHFIHLIYKPSSGPVKTKLALVGKGLTFDSGGYNIKTGPGCLIELM  373
S. Lycopersicum LAP2   YLRVAA-AS--ANPAHFIHLCYKPSSGEIKKKIALVGKGLTFDSGGYNIKTGPGCSIELM  361
S. tuberosum           YLGVAAAAT--ENPPYFIHLCFKTNSRERKTKIALVGKGLTFDSGGYNLKTGAGSKIELM  345
S. Lycopersicum LAP1   YLAVAAAAT--ENPPYFIHLCFKTPTKERKTKLALVGKGLTFDSGGYNLKVGAGSRIELM  364
T. annulata            YLSVCQ-GS--KFPPKFLHAVYKP-DGEVKKTLAFVGKGITFDAGGYNVKN-SSTMIHYM  318
T. gondii              YLAVAK-GS--LFPPQFIHLTYKS-SN-PKRRVAFVGKGICFDAGGYNLKT-GGAQIELM  566
P. falciparum          YLSVGK-GS--MYPNKFIHLTYKS-KGDVKKKIALVGKGITFDSGGYNLKAAPGSMIDLM  396
C. hominis             FLAVAQ-GS--KSPAQFVHLTYKP-KGEIKKRIALVGKGITMDTGGYNIKH---QMIHFM  315
C. parvum              FLAVAQ-GS--KSPAQFVHLTYKP-KGEIKKRIALVGKGITMDTGGYNIKH---QMIHFM  341
L. major               MYNVGR-GS--RYEPYLVVLEYIGNPR-SSATTAIVGKGVTFDCGGLNIKP--YTSMETM  345
H. pylori              FLAVNK-ASLSVNPPRLIHLVYKPKKA--KKKIALVGKGLTYDCGGLSLKP--ADYMVTM  284
S. aureus              IHAVGK-GS--EHPPVVITMTYNGGDS-NDAPIALVGKGITYDSGGYSIKS--KIGMQTM  280
E. coli                YLAVGQ-GS--QNESLMSVIEYKGNASEDARPIVLVGKGLTFDSGGISIKP--SEGMDEM  290
R. typhi               LLGVGQ-GS--QNESKLVVMEYKGGSR-DDSTIALVGKGVIFDTGGISLKP--SSNMHLM  285
C. tetani              FLAVSK-GS--FEESQLIVMNYKGNSN-SDKKLALVGKGLTYDSGGYSIKP--TSSMINM  267
B. cereus              LLAVNQ-GS--VEPPKMIALIYKGKEE-WKDVIGFVGKGITYDTGGYSLKP--REGMVGM  280
                          *   .      .     .        ::*:*    *  **  :*            :   *
```