1

Oscillating bacterial expression states generate herd immunity to viral infection

Christopher J. R. Turkington [a,c], Andrey Morozov [b], Martha R. J. Clokie[c],

Christopher D. Bayliss[a]#

Department of Genetics and Genome Biology, University of Leicester, Leicester, United

Kingdom[a]; Department of Mathematics, University of Leicester, Leicester, United

Kingdom[b]; Department of Infection, Immunity, and Inflammation, Leicester, United

Kingdom[c]

Running Head: Bacterial herd immunity against viral infection

#Address correspondence to Christopher D. Bayliss, cdb12@le.ac.uk

Abstract word count = 184

Main text word count = 4527

Abstract:

Hypermutable loci are widespread in bacteria as mechanisms for rapid generation of

phenotypic diversity, enabling individual populations to survive fluctuating, often

antagonistic, selection pressures. As observed for adaptive immunity, hypermutation may

facilitate survival of multiple, spatially-separated bacterial populations. We developed an

'oscillating prey assay' to examine bacteriophage (phage) spread through populations of

Haemophilus influenzae whose phage receptor gene, *lic2A*, is switched 'ON' and 'OFF'

by mutations in a hypermutable tetranucleotide repeat tract. Phage extinction was

frequently observed when the proportion of phage-resistant sub-populations exceeded

34%. *In silico* modelling indicated that phage extinction was interdependent on phage

loss during transfer between populations and the frequency of resistant populations. In a

fixed-area oscillating prey assay, heterogeneity in phage resistance was observed to

generate vast differences in phage densities across multiple bacterial populations

resulting in protective quarantining of some populations from phage attack. We conclude

that phase-variable hypermutable loci produce bacterial 'herd immunity' with resistant

intermediary-populations acting as a barricade to reduce the viral load faced by phage-

sensitive sub-populations. This paradigm of meta-population protection is applicable to

evolution of hypermutable loci in multiple bacteria-phage and host-pathogen interactions.

Importance

Herd immunity is a survival strategy wherein populations are protected against invading

pathogens by resistant individuals within the population acting as a barrier to spread of

the infectious agent. Although, this concept is normally only applied to higher

eukaryotes, prokaryotic organisms also face invasion by infectious agents, such as

bacterial viruses, bacteriophage (phage). Here we use novel experimental approaches and

mathematical modelling, to show that bacteria exhibit a form of herd immunity through

stochastically generated resistant variants acting as barricades to phage predation of

sensitive cells. With hypermutable loci found in many prokaryotic systems, this

phenomenon may be widely applicable to phage-bacteria interactions and could even

impact phage-driven evolution in bacteria.

Introduction

Hypermutable loci as mediators of survival against constantly fluctuating

selection pressures are a predictable outcome of the evolution of evolvability as stated in

the Red Queen hypothesis (1, 2). Fluctuating selection pressures are regularly faced by

bacteria during persistence in human hosts, where bacteria adhere to host surfaces while

contending with varying nutrient concentrations, frequent exposure to immune effectors

and predation by bacteriophages. These fluctuations often select for and against opposing

gene expression states of single loci leading to evolution of localised hypermutable

mechanisms that produce frequent switches in single-gene expression states.

One class of hypermutable loci facilitate survival of conflicting selective

pressures by pre-emptive, frequent, and reversible 'ON/OFF' generation of adaptive

variants, in a process known as 'phase variation' (PV; 2–5). A major mechanism of PV

involves increases and decreases in identical, tandemly-arranged DNA repeats

(microsatellites) by slipped strand mispairing during DNA replication (3, 5, 6). The

obligate human respiratory commensal and pathogen *H. influenzae* contains an expansive

array of repeat driven phase variable loci (7–13). Several of these loci are required for

addition of sugar molecules onto the surface-exposed outer-core of the

lipooligosaccharide (LOS) (5, 7). PV of the UDP-galactose-LOS-galactosyltransferase

encoding gene, *lic2A*, is mediated by a 5'CAAT repetitive tract present in the open-

reading frame (14). Phage HP1c1 attaches to an LOS epitope of *H. influenzae* strain Rd

that contains the galactose sugar added by *lic2A* (15). PV of *lic2A* causes switching

between phage sensitive (*lic2A* ON) and phage resistant states (*lic2A* OFF) (15, 16).

Partial resistance to phage HP1c1 in strain Rd is also mediated by PV of a Type I

restriction-modification (RM) system. PV of surface receptors and RM systems generates

resistance to phage infection in several bacterial species (17–22).

While phage-receptor PV prevents viral propagation in individual populations, the

frequency and distribution of resistant variants within larger meta-populations may also

impose inhibitory effects on phage spread through multiple spatially-linked sub-

populations. In order to explore this potential benefit, we examined how diversity in

phage resistance/sensitivity phenotypes generated by one hypermutable locus, *lic2A*,

alters spread of phage HP1c1.

RESULTS

Low numbers of resistant bacterial populations significantly restrict phage spread

and densities. PV can generate high levels of ON and OFF variants of single genes

within individual populations but also has the potential to generate population-to-

population variation within a meta-population. For example, *lic2A*-positive *H. influenzae*

colonies on agar plates will have most cells in a *lic2A* ON expression state but will also

contain small, but significant, numbers of *lic2A* OFF variants. Similarly, an *H. influenzae*

meta-population may consist of multiple sub-populations with one fraction being *lic2A*-

positive and another fraction being *lic2A*-negative. Such population-to-population

variation is observed for *H. influenzae* colonies on agar plates and for *H. influenzae*

populations isolated from artificially-inoculated animals and asymptomatic human

carriers (23–27). Proportions of ON and OFF sub-populations in a meta-population will

depend on levels of selection for each expression state and on 'founder' effects that

influence the starting state of each sub-population. Thus, any phage invading a bacterial

meta-population where there is PV of the receptor must contend with highly variable

distributions of resistance and sensitive sub-populations.

To simulate the effect of phage-receptor PV on spread of phage through bacterial

meta-populations we developed the 'oscillating prey assay' (Fig. 1). This assay involves

continual cycling of phage through *H. influenzae* strain Rd cultures with either a majority

lic2A ON or OFF phenotype. Each cycle allows for one round of phage replication after

which the phage-containing supernatant is recovered, filtered, and transferred to a new

culture of either *lic2A* ON or OFF *H. influenzae* cells. During transfer, the supernatant is

subject to a 10-fold dilution. This arbitrary dilution factor simulates loss of phage during

population-to-population transmission.

Six population structures were examined in the oscillating prey assay: [1] 100 %

ON (HP1c1 cycled only through *lic2A* ON populations; S100), [2] 66 % ON (2:1 *lic2A*

ON:OFF; S66), [3] 50 % ON (1:1 *lic2A* ON:OFF, starting with an ON culture; S50), [4]

50 % OFF (1:1 *lic2A* ON:OFF, starting with an OFF culture; R50), [5] 66 % OFF (1:2

lic2A ON:OFF; R66), and [6] 100 % OFF (HP1c1 cycled only through *lic2A* OFF

populations; R100). Survival and propagation of phage was dependent on the proportion

of phage-resistant sub-populations in each series of 20 cycles (Fig. 2). Survival of phage

to the final cycle was only observed when the proportion of phage-resistance populations

was ≤ 34% (i.e. the S66 and S100 populations; Fig. 2). Extinction events occurred within

5 to 16 cycles in all other heterogeneous and homogeneous populations at a rate that was

dependent on the proportion of resistant sub-populations. Despite survival of phage

through to cycle 20 in the S66 population series, phage densities were significantly

decreased by this cycling regime (Fig. 2; paired *t*-test: $t = 4.97$, $P < 0.05$; mean ± SEM

PFU/mL values for phage densities at cycle 0 = $7.16 \pm 0.26 \times 10^5$ and cycle 20 =

$2.82 \pm 0.9 \times 10^4$), indicating that further passages with a similar pattern would have

resulted in phage extinction. Contrastingly, phage density increased when all sub-

populations were phage sensitive (S100), with phage densities plateauing after ~11

cycles. Thus, both phage survival and density were limited by the frequency of

encounters with phage-resistant *lic2A* OFF phase variants during passage through a linear

series of sub-populations.

Detection of combinatorial effects of population structure and dilution rate

on phage extinction using an *in-silico* model of phage spread. Our experimental data

indicated that meta-population structure had a major impact on phage survival and

spread. In order to explore a wider range of linear and non-linear cycling patterns and the

effects of dilution rate, we developed a mathematical model of the oscillating prey assay.

This model utilized key phage parameters for adsorption rate, replication time, burst size

and stability of phage HP1c1 (see Fig. S1). The number of oscillations was extended to

105 phage replicative cycles, while cycling patterns were randomised for each specific

overall proportion of *lic2A* ON/OFF sub-populations. The mathematical model produced

comparable findings to the experimental setting (Fig. S2). Multiple runs of this model

exhibited stochastic variation in phage densities and extinction events as observed in the

experimental model but with extinction events occurring over a wider band of replication

cycles (Fig. 3A-B). This model demonstrates that random patterns of ON/OFF expression

states for a phage receptor can limit phage spread.

Multiple simulations (n = 200) were performed for each combination of R (the

percentage of phage-resistant *lic2A* OFF sub-populations) and C (the inverse of the

dilution coefficient). Average phage densities were measured for all cycles and runs of

each combination of R and C, and phage extinction was defined to occur whenever the

density fell below 100 PFU/mL. Phage extinction was always observed when the number

of resistant states exceeded 70% of sub-populations, even at a low dilution rate of 1 in 2

(C=0.5; Fig. 3C). Similarly, when phage loss was $\geq$ 98 % at each transfer (i.e. C=0.02),

phage extinction was observed for all meta-population structures except those consisting

of >95% phage-sensitive sub-populations (Fig. 3C). Between these extremes, there was

an accelerating trade-off between R and C with respect to phage extinction and average

phage density; decreases in dilution rate were countered by a high prevalence of resistant

sub-populations enabling bacterial populations to survive even when phage dispersal was

low (Fig. 3C). This observation suggests that on-going evolution of localised

hypermutability for a surface-exposed bacterial epitope could be tuned to the prevalence

and density of phages capable of using the phase-variable epitope as a receptor for

surface attachment.

Localised hypermutation-driven herd-immunity produces regional variations

in phage densities within bacterial meta-populations. Both our experimental and *in*

silico oscillating prey assays demonstrated that phage spread was influenced by the linear

pattern of phase-variant sub-populations. However, the distribution of phase variants

across a surface (e.g. microcolonies on upper respiratory tract surfaces for *H. influenzae*)

is anticipated to be random and hence to result in spatial effects on phage spread. Indeed,

spatially structured environments are known to restrict phage spread (28). Spatial

structuring was explored by examining phage transmission across meta-populations of

fixed dimensions but with varying proportions of each *lic2A* expression state (Fig. S3).

This fixed-area oscillating prey assay was initiated by inoculating phage into one well of

a 96-well plate and then expanding outward by seeding each subsequent replicative cycle

into neighbouring wells (see Fig. S4).

Passage of HP1c1 through heterogeneous populations of the fixed-area oscillating

prey assay resulted in uneven phage densities across the meta-populations consisting of

50-66% phage-sensitive populations (Fig. 4B-4D). Conversely, homogenous densities

were observed for high (Fig. 4A) or low (Fig. 4E and 4F) phage-sensitive distributions.

When 66% of populations were phage-sensitive (Fig. 4B), densities ranged from 10^3 to

10^{10} PFU/mL, whereas densities always exceeded 10^8 PFU/mL if all population were

sensitive (Fig. 4A). Phage densities in specific regions of heterogeneous populations were

dependent on the direction of propagation with passage through multiple resistant or

sensitive sub-populations resulting in low or high phage densities, respectively. This

model shows how spatial meta-population heterogeneity could prevent equal

dissemination of phage through host populations with phase-variable phage-receptors and

aid survival of phage-sensitive sub-populations whose phenotypes may be beneficial for

bacterial survival against other selective pressures.

Observed PV rates generate populations with phase-variant proportions

capable of herd-immunity. The fixed-area oscillating prey assay outputs showed how

phage extinction was dependent on the proportion of phage-resistant sub-populations. *H.*

influenzae normally resides in the upper respiratory tract where selection is likely to act

on both PV states of a locus. Thus, the proportion of resistant populations depends on

both selection strength for/against the phage-resistance phenotype, and the ON/OFF

switching rate. Switching rates of *H. influenzae* phase-variable genes are malleable due to

changes in repeat number and can evolve in response to alternating selection pressures

(28, 29).

A mathematical model was developed to examine the impact of different

switching rates and immune selection on the proportions of *lic2A* expression states (Fig. 5

and Supplementary Data S1). Dixon *et al.* (30) found that the *lic2A* ON-to-OFF (S-to-R;

where R and S are the phage-resistant and phage-sensitive states, respectively) switching

rate was 1.7-fold higher than the *lic2A* OFF-to-ON (R-to-S) switching rate. The

mathematical model assumed that these proportions were maintained for three 10-fold

differences in overall mutation rate representing low, intermediate and high repeats

numbers (Fig. 5A-5C). In the absence of any selective difference between the S and R

states ($m=1$), a high, steady-state proportion (>63%) of R, was observed for all mutation

rates but with minor differences in the rate of approach to the steady state and absolute

amounts of R. The *lic2A* OFF state (i.e. R) is known to be more immune sensitive than

the ON state (i.e. S), we therefore imposed a selection against R. Even with strong

selection (m=0.99), high levels of R were maintained by medium to high switching rates

(Fig. 5A and 5C, respectively). In contrast, when switching rates were low (Fig. 5B),

even weak selection (m=0.999) drives resistant variants to <1% (Fig. 5B). Our other *in*

silico models indicated that phage spread was inhibited when the probability of

encountering R variants was between 10-70% for dilution rates of 0.02 to 0.6. Thus, the

immune model shows that observed repeat numbers and switching rates for the *lic2A*

gene of *H. influenzae* strains can maintain sufficient phage-resistant variants for

restricting phage spread even when there is immune selection against this state.

DISCUSSION

Hypermutable loci have well-documented roles in facilitating survival of

individual bacterial populations against phage predation through generation of phage-

resistant cells. An unexplored concept is that hypermutation driven heterogeneity in

phage resistance across the wider population also facilitates bacterial survival of phage

predation.

Phase-variable loci can generate herd immunity in bacterial meta-

populations. The concept of herd immunity was derived to explain the protection of

susceptible individuals in populations with high levels of immunity to an infectious agent

as observed for measles in Baltimore (31). Ordinarily applied to naturally or vaccine-

acquired immunity to infectious agents in human populations (32), we show, herein, that

repeat-mediated PV of a phage-receptor provides a form of 'bacterial herd-immunity' at

the population level (Fig. 6). Thus, phage spread is retarded by resistant sub-populations

creating barriers between the phage and phage-sensitive bacterial sub-populations as

shown in our one- and multidirectional experimental models (Fig. 2 and Fig. 5), and an *in*

silico model with randomized patterns of phage sensitivity (Fig. 3). Key features of these

models were that: the chance of phage survival was inversely proportional to the linear

pattern of resistance faced by the phage population; random distributions of phage-

resistant/sensitive populations result in large variations in viral numbers across a meta-

population with some regions being completely free of phage; and dilution during

transfer of phage between populations modulates the number of phage-resistant

populations required to retard phage spread. Thus, localised hypermutation of a phage-

receptor generates herd-immunity whereby phage-sensitive sub-populations can be

maintained at high levels within bacterial meta-populations.

PV of phage receptors or RM systems is an established phenomenon observed

across numerous species and occurring by multiple mechanism. Repeat-mediated

switching due to hypermutation of polyG tracts in phage receptor genes of *C. jejuni* strain

NCTC11168 abrogates infection by phage F336 (19). Similar polyG hypermutation

controls the phage growth limitation system of *Streptomyces coelicor* A3(2) and confers

resistance to infection with phage φC31 (33). Epigenetic PV of Ag43 in *E. coli* or the

gtr^{P22} operons that control O-antigen modification in *Salmonella* are known or proposed

to modulate phage infection (34, 35). High frequency, but not hypermutable, mutations in

short polyG or polyA tracts produce resistance to phages in both *Bordetella pertussis* (36)

and *Vibrio cholerae* (17). Although these mechanisms vary in the rates of generation of

phage resistant variants and in the strength of phage resistance (e.g. high for receptor PV

and low for RM PV), they all have the potential to generate spatially-structured

populations and hence herd immunity to phage infection.

Additionally, there is potential for evolution of the herd immunity state. PV rates

can evolve through changes in the mutable mechanism. Thus repeat-mediated PV rates

increase as a function of tract length. We anticipate that frequent exposure to phage

would select for a greater capacity to form heterogeneous meta-populations through

secondary selection for an increase in mutability of the phage receptor.

Evidence for immune-driven selection of lic2A ON expression states. One

rationale for the existence of PV-driven herd immunity is that protection of the phage-

sensitive state is required because this state is advantageous under certain circumstances.

For *lic2A*, the ON state in *H. influenzae* strain Rd is phage-sensitive and this states leads

to extension of the LOS side-chain from the third heptose with a single galactose, a

digalactose or more complex sugars. In *in vitro* studies, *lic2A*-dependent epitopes aid in

survival against human immune responses (16), with the LOS extensions associated with

lic2A expression encoding epitopes also present on the human P blood group antigens

(37). Although human volunteer studies of colonisation with *H. influenzae* have found

that expression of *lic2A* was not essential for human nasopharyngeal colonisation (26),

the *lic2A* ON state has been associated with disease states, including non-typeable *H.*

influenzae pneumonia (24). We performed an analysis of 104 *H. influenzae* genome

sequences and found that *lic2A* is in an ON state in ~63% of all isolates and is

predominantly in the ON state for multiple disease conditions, suggesting selection for

expression of *lic2A* occurs across a wide range of niches for *H. influenzae* (Fig. S5).

These observations are consistent with a scenario where phage drive evolution of

hypermutability rates as the *lic2A* gene oscillates between selection for/against the

immune-resistant/phage-sensitive and immune-sensitive/phage-resistant states. A caveat

to these conclusions is phage-specificity as phage HP1c1 may be specific to extension of

the third heptose of *H. influenzae* LOS whereas Lic2A can, in the appropriate genetic

context, generate extensions from any of the three heptoses of the LOS inner core.

Further work is required to determine whether HP1c1 is specific for extension from the

third heptose and if other phages can target this epitope in *H. influenzae* strains where

extension is from the first or second heptose.

Transmission and the 'phage loss' phenomenon. Our mathematical model of

herd immunity indicates that phage spread is interdependent on population structure and

the rate of phage loss from the environment by dilution. Thus, when dilution rates are

high, phage extinction events are frequent despite high levels of sensitivity within the

bacterial population. While natural rates of phage loss from respiratory environments is

unknown, a number of factors have the capacity to impact phage loss, such as humidity,

salinity and immune responses (38–41). For phage infections of human commensals or

pathogens, such as *H. influenzae*, more extreme environmental selection pressures will

apply as phages are transmitted between carriers of target bacterial species.

There are two potential extreme scenarios where the herd immunity model is

applicable and 'phage loss' is either low or high. These scenarios are elaborated for *H.*

influenzae but are relevant to other bacterial commensals/pathogenic bacteria. Firstly,

colonisation of asymptomatic carriers by *H. influenzae* is likely to involve a series of

microcolonies, a meta-population, distributed across the nasopharyngeal surface rather

than one continuous population. Phage will therefore have to transmit between

microcolonies thereby imposing a low potential for phage loss such that only high

numbers of phage-resistant populations will provide protection for any phage-sensitive

microcolonies. A second scenario is a population of *H. influenzae* carriers, in this case

phage transmission between carriers is likely to result in significant phage loss and hence

low numbers of carriers colonized by phage-resistant populations could prevent phage

spread to all carriers. These scenarios illustrate the central role of transmission in shaping

bacterial herd immunity and in the impact of this fitness trait on localised hypermutation

of the phage receptor.

Bacterial herd immunity could impact on phage-driven evolution. We

observed that phage densities were highly variable across spatially-structured bacterial

populations with some regions exhibiting a complete absence of phage (Fig. 6). This

imposition of spatially-discrete levels of phage selection could select for alternative

adaptive traits within the bacterial host. In studies of *Caulobacter crescentus*, low phage

selection led to isolation of >200 phage resistance mechanisms, while only ~60 distinct

resistance forms were isolated during high phage selection (42). Thus, bacterial herd-

immunity may prevent uniformity in phage selection pressures across bacterial meta-

populations leading to evolution of distinct phage-resistance mechanisms within a single

clonal lineage.

In summary, our demonstration of a hypermutable locus retarding spread of an

infectious agent within a prokaryotic meta-population suggests that the herd immunity

phenomenon may be applicable to a wide variety of interacting biological organisms and

have deep evolutionary roots. Our conceptual framework could be utilised to explore

whether somatic hypermutation, a key example of localised hypermutation in eukaryotes,

evolved through selection for sub-population heterogeneity linked to pathogen resistance.

MATERIALS AND METHODS

Phage and bacterial strains used in this study. Phage HP1c1, and the *lic2A* ON (Rd 30S) and OFF (Rd 30R) phase variants of *H. influenzae* were obtained from A. Piekarowicz (University of Warsaw, Poland). Phage HP1c1 is maintained in the lysogenic state within *H. influenzae* RM118-L. *H. influenzae* strains were cultured overnight at 37°C on 1% BHI agar supplemented with 10% Leventhal's supplement and 2 µg/mL nicotinamide adenine dinucleotide (NAD) or in 10 mL BHI broth supplemented with 2 µg/mL NAD and 10 µg/mL hemin (sBHI).

Phage HP1c1 stocks. Mitomycin C was added to a concentration of 300 ng/mL to an OD_{600} 0.1 culture of RM118-L in 10 ml of sBHI. After incubation for 8 hours, the culture was centrifuged (4,946xg, 4°C, 10 minutes) and then the supernatant was passed through a 0.22 µm syringe filter to obtain a phage suspension, which was stored at 4°C.

High titer phage stocks were propagated in *H. influenzae* Rd 30S by adding 100 µL of induced phage to mid-log phase cultures diluted to an OD_{600} of 0.01 in 10 ml sBHI and incubating for 8 hours. Cultures were processed as described above.

Determination of phage titers. Phage titers were determined using the small-drop plating assay (43). Briefly, 150 µl of a mid-log phase Rd 30S culture, OD_{600} 0.1, was added to 3 mL of 0.3 % BHI agar (supplemented with 2 µg/mL NAD and 10 µg/mL hemin), mixed by inversion, and poured onto 1% BHI agar plates supplemented with 10% Leventhal's media and 2 µg/mL NAD. Ten-fold serial dilutions were spotted in triplicate 10 µl drops onto the soft agar (the minimum detection threshold is 33.3 PFU/mL).

Oscillating prey assay. Two phase variants, namely, Rd 30S (*lic2A* ON, phage

sensitive variant) and Rd 30R (*lic2A* OFF, phage resistant variant) were utilized to

generate the following three-repeat host population-cycling patterns: S100, ON-ON-ON;

S66, ON-ON-OFF; S50, ON-OFF-ON; R50, OFF-ON-OFF; R66, OFF-OFF-ON; R100,

OFF-OFF-OFF. *H. influenzae* Rd 30S and Rd 30R were sub-cultured as described above

in 20 mL of sBHI to an OD_{600} of 0.1. A 5 mL aliquot of the relevant strain was inoculated

with HP1c1 at MOI ~0.01. Cultures were adjusted to 6 mL, mixed by inversion and 1 mL

was removed for filtration and determination of the T = 0 phage titre. The remaining 5

mL was incubated for 50 minutes (i.e. one viral replication cycle) at 37°C with shaking

and then filtered for phage quantification. Subsequent cycles were initiated by

transferring 600 µl of filtrate to a fresh 5 mL culture of relevant *lic2A* phase variant. Five

phage transfers were conducted per day with phage-containing filtrates being stored

overnight at 4°C.

Fixed-area oscillating prey assay. Six cycling frequencies were tested (Fig. 5).

Allocation of each phase variant (i.e. Rd 30S or Rd 30R) to specific wells was

determined by numbering wells from 1 to 631 followed by randomization of these

numbers into two sets using R (see Fig. S3). One phase variant was added to the first set

of numbered wells and the other phase variant to the second set.

Rd 30S and Rd 30R were sub-cultured to OD_{600} 0.1 and then 250 µL of appropriate

culture was added to the starter well. Phage HP1c1 was added at a final concentration of

~1 x 10^6 PFU/mL to this well and the volume was adjusted to 300 µL with fresh sBHI

broth. After mixing by tituration, 100 µL was removed for phage titration. The plate was

incubated at 37°C with shaking for 70 minutes (a longer incubation time was required in

this miniaturised oscillating prey assay for completion of one phage replication cycle

[data not shown]). Following incubation, the plate was centrifuged at 1500 x *g* for 4

minutes to pellet bacterial cells and then 20 µl of supernatant was transferred to

surrounding wells (see Fig. S4). Remaining supernatant was harvested for phage titration.

Newly-inoculated wells received 167 µl of a fresh OD_{600} 0.1 culture of either Rd 30S or

Rd 30R, depending on the cycling pattern, followed by repetition of all previously

described steps. Ten cycles were performed in the left, right, and downward directions,

and 20 cycles in the upward direction (see Fig. S3 and S4). Five transfers and cycles were

conducted each day with the 96-well plates being wrapped in paraffin film and stored at

4°C overnight. This assay was conducted once for each population structure.

Survey of Lic2A expression state in multiple *H. influenzae* genomes. A set of

126 genomes available in GenBank as of 06-09-2016 were screened with blast2seq for

the presence of genes homologous to the *lic2A* gene of *H. influenzae* strain Rd KW20.

Lic2A homologues were found in 121 of the 126 genomes but only 104 could be analysed

due to incomplete sequence coverage in 17 genomes. The ON state was identified by the

presence of a full length amino acid product (~300 amino acids). The 5'CAAT repeat

numbers were determined by visual inspection of aligned sequences. All metadata, where

available, was collated from the GenBank entry for each genome, or references

associated with each strain.

Mathematical model and simulations. We describe bacteria-phage interaction

using the conventional modelling approach (45, 46). Dynamics of bacteria and phage

densities in each experimental cycle (transfer) of number *n* within time T_0=40min (i.e.

before the start of mass replication of phages) is given by

$$\frac{dB_0}{dt} = -KPB_0,$$

$$\frac{dB_i}{dt} = KPB_{i-1} - KPB_i, \quad i = 1, ..., N-1$$

$$\frac{dB_N}{dt} = KPB_{N-1},$$

$$\frac{dP}{dt} = -PK \sum_{i=0}^{N-1} B_i - mP, \quad nT < t < Tn + T_0,$$

where B_0 is the number/density of phage-free bacteria; B_i is the number/density of bacteria with i phage attachments; P is the number/density of free phages. The maximal number of phage attachments for an individual bacterial cell is given by N. In the model, we assume that the injection rate is very fast (i.e. instantaneous), so that attachment of one phage immediately results in an infection.

For simplicity, we assume that there is no bacterial growth. We also assume that all phage attachments occurring within the first 10 minutes lead to replication (each phage produces b new phages) whereas later attachments result in phage loss without replication. We neglect binding of newly replicated phages within the last $\Delta = 10$ min of each cycle. The other model parameters are: K-, phage adsorption constant (note that this constant is assumed to be independent from the number of bound phages and that $K=0$ for phage-resistant bacteria); $m,$ natural mortality of phages; b, phage burst size.

At the start of each experimental cycle (i.e. just after dilution) all bacteria are phage-free and their number is always equal to B_S, in other words,

$$B_0\left(t = (Tn)^+\right) = B_s, \quad B_i\left(t = (Tn)^+\right) = 0, \quad B_i(t = (Tn)^+) = 0, \quad i=1,\ldots, N,$$

Here the symbol '+' denotes the time just after the n^{th} dilution, i.e. just prior to the

cycle ($n+1$); the symbol '−' denotes the time prior to the n^{th} dilution, i.e. at the very end

of cycle n.

The phage density is obtained from the final density in each cycle multiplied by the

dilution coefficient C_n in cycle #n (this value can vary between experiments).

$$P(t = (Tn)^+) = P(t = (Tn)^-)C_n$$.

The density of phages just prior to dilution n (i.e. at the end of cycle #n) is

determined by

$$P(t = (Tn)^-) = \sum_{i=1}^{N} B_i (Tn - T_0)b + P_S(t = (Tn)^-)$$,

where P_S are non-attached phages that survived to the end of the cycle; i.e. the phage

number at the cycle end, prior to dilution, is given by the number of infected bacteria at

time $T - T_0$ multiplied by the burst size b plus the number of surviving phages P_s.

Susceptible bacteria are characterized by $K>0$, whereas for resistant bacteria we

have $K=0$. The value of K is kept constant across each cycle of 50 minutes.

Model parameters and verification were derived from experimental settings or

findings. Direct observations indicated that $B_S=1.75\pm0.25*10^8$ cell/ml, $T=50$ min, and

$T_0=40$ min. The adsorption constant of $K=7\pm3*10^{-10}$ mL/(cell min) was estimated from

an adsorption assay (Fig. S1). Other parameters were estimated directly from the

oscillating prey assay by model fitting (see Fig. S2): $b=42\pm5$ (burst size); and

$m=0.006\pm0.003$ 1/min. The parameter N had a minor effect in our computation and hence

we utilised $N=20$ in all subsequent models.

430   Further simulations (e.g. Fig. 3c) considered the phage-bacterial dynamics across

431  105 cycles. Variations in the dilution rate C were simulated by changing the value of C

432  according to $C=C^*(1+\varepsilon)$, where ε is a normally distributed random variable with a mean

433  of 0. and variance of 0.3^2 . In each cycle, K was randomly switched from $K= 7*10^{-10}$

434  (susceptible bacteria) to $K=0$ (resistant bacteria). The frequency of switching was

435  determined by the probability p, which gives the probability of encountering susceptible

436  bacteria. Examples simulations, shown in Fig. 3a and 3b, were constructed for $C=0.1$,

437  $p=0.75$ (A) $p=0.85$ (B). Fig. 3c was obtained by repeating simulations across 105 for

438  variable parameters C and p. Numerical simulation was based on the standard Runge-

439  Kutta integration method of order 4 using MATLAB software. When phage density

440  dropped to or below the low value threshold of $P_0=100$, we considered that this was

441  equivalent to $P=0$ and stopped further simulations. The initial density of phages at time

442  $t=0$ was $4.28*10^7$ PFU/mL.

444  **FUNDING INFORMATION**

445  C.J.R.T. was supported by the Biotechnology and Biological Sciences Research Council

446  (BBSRC) [BB/J014532/1] and the College of Life Sciences of the University of

447  Leicester.

449  **REFERENCES**

450  1.  van Valen L. 1973. A new evolutionary law. Evol Theory 1:1–30.

451  2.  Moxon ER, Rainey PB, Nowak MA, Lenski RE. 1994. Adaptive evolution of

452    highly mutable loci in pathogenic bacteria. Curr Biol 4:24–33.

3.   Bayliss CD. 2009. Determinants of phase variation rate and the fitness

implications of differing rates for bacterial pathogens and commensals. FEMS

Microbiol Rev 33:504–20.

4.   Rainey PB, Beaumont HJE, Ferguson GC, Gallie J, Kost C, Libby E, Zhang X-X.

2011. The evolutionary emergence of stochastic phenotype switching in bacteria.

Microb Cell Fact 10 Suppl 1:S14.

5.   Moxon ER, Bayliss CD, Hood DW. 2006. Bacterial contingency loci: the role of

simple sequence DNA repeats in bacterial adaptation. Annu Rev Genet 40:307–33.

6.   van der Woude MW, Bäumler AJ. 2004. Phase and antigenic variation in bacteria.

Clin Microbiol Rev 17:581–611, table of contents.

7.   Power PM, Sweetman W a, Gallacher NJ, Woodhall MR, Kumar G a, Moxon ER,

Hood DW. 2009. Simple sequence repeats in *Haemophilus influenzae*. Infect

Genet Evol 9:216–28.

8.   Bayliss CD, Field D, Moxon ER. 2001. The simple sequence contingency loci of

Haemophilus influenzae and *Neisseria meningitidis*. J Clin Invest 107:657–62.

9.   Kroll JS, Moxon ER. 1988. Capsulation and gene copy number at the cap locus of

Haemophilus influenzae type b. J Bacteriol 170:859–64.

10.   Ren Z, Jin H, Whitby PW, Morton DJ, Stull TL. 1999. Role of CCAA nucleotide

repeats in regulation of hemoglobin and hemoglobin-haptoglobin binding protein

genes of *Haemophilus influenzae*. J Bacteriol 181:5865–70.

11.   Dawid S, Barenkamp SJ, St Geme JW. 1999. Variation in expression of the

Haemophilus influenzae HMW adhesins: a prokaryotic system reminiscent of

eukaryotes. Proc Natl Acad Sci U S A 96:1077–82.

12.   Glover SW, Piekarowicz A. 1972. Host specificity of DNA in *Haemophilus*

477   *influenzae*: restriction and modification in strain Rd. Biochem Biophys Res

478   Commun 46:1610–7.

479   13.   De Bolle X, Bayliss CD, Field D, Van De Ven T, Saunders NJ, Hood DW, Moxon

480   ER. 2000. The length of a tetranucleotide repeat tract in *Haemophilus influenzae*

481   determines the phase variation rate of a gene with homology to type III DNA

482   methyltransferases. Mol Microbiol 35:211–222.

483   14.   High NJ, Deadman ME, Moxon ER. 1993. The role of a repetitive DNA motif (5'-

484   CAAT-3') in the variable expression of the *Haemophilus influenzae*

485   lipopolysaccharide epitope alpha Gal(1-4)beta Gal. Mol Microbiol 9:1275–82.

486   15.   Zaleski P, Wojciechowski M, Piekarowicz A. 2005. The role of Dam methylation

487   in phase variation of *Haemophilus influenzae* genes involved in defence against

488   phage infection. Microbiology 151:3361–9.

489   16.   Clark SE, Eichelberger KR, Weiser JN. 2013. Evasion of killing by human

490   antibody and complement through multiple variations in the surface

491   oligosaccharide of *Haemophilus influenzae*. Mol Microbiol 88:603–18.

492   17.   Seed KD, Faruque SM, Mekalanos JJ, Calderwood SB, Qadri F, Camilli A. 2012.

493   Phase variable O antigen biosynthetic genes control expression of the major

494   protective antigen and bacteriophage receptor in *Vibrio cholerae* O1. PLoS Pathog

495   8:e1002917.

496   18.   Cota I, Blanc-Potard AB, Casadesús J. 2012. STM2209-STM2208 (opvAB): a

497   phase variation locus of *Salmonella enterica* involved in control of O-antigen

498   chain length. PLoS One 7:e36863.

19.   Sørensen MCH, van Alphen LB, Harboe A, Li J, Christensen BB, Szymanski CM,

Brøndsted L. 2011. Bacteriophage F336 recognizes the capsular phosphoramidate

modification of *Campylobacter jejuni* NCTC11168. J Bacteriol 193:6742–6749.

20.   Anjum A, Brathwaite KJ, Aidley J, Connerton PL, Cummings NJ, Parkhill J,

Connerton I, Bayliss CD. 2016. Phase variation of a Type IIG restriction-

modification enzyme alters site-specific methylation patterns and gene expression

in *Campylobacter jejuni* strain NCTC11168. Nucleic Acids Res gkw019.

21.   Sekulovic O, Ospina Bedoya M, Fivian-Hughes AS, Fairweather NF, Fortier L-C.

2015. The *Clostridium difficile* cell wall protein CwpV confers phase-variable

phage resistance. Mol Microbiol 98:329–342.

22.   Latino L, Midoux C, Hauck Y, Vergnaud G, Pourcel C. 2016. Pseudolysogeny and

sequential mutations build multiresistance to virulent bacteriophages in

Pseudomonas aeruginosa. Microbiology 162:748–63.

23.   High NJ, Jennings MP, Moxon ER. 1996. Tandem repeats of the tetramer 5'-

CAAT-3' present in *lic2A* are required for phase variation but not

lipopolysaccharide biosynthesis in *Haemophilus influenzae*. Mol Microbiol

20:165–74.

24.   Weiser JN, Pan N. 1998. Adaptation of *Haemophilus influenzae* to acquired and

innate humoral immunity based on phase variation of lipopolysaccharide. Mol

Microbiol 30:767–75.

25.   Hosking SL, Craig JE, High NJ. 1999. Phase variation of *lic1A*, *lic2A* and *lic3A* in

colonization of the nasopharynx, bloodstream and cerebrospinal fluid by

Haemophilus influenzae type b. Microbiology 145 (Pt 1:3005–11.

26.   Poole J, Foster E, Chaloner K, Hunt J, Jennings MP, Bair T, Knudtson K,

Christensen E, Munson RS, Winokur PL, Apicella MA. 2013. Analysis of

nontypeable *Haemophilus influenzae* phase-variable genes during experimental

human nasopharyngeal colonization. J Infect Dis 208:720–7.

27.   Fox KL, Atack JM, Srikhanta YN, Eckert A, Novotny L a, Bakaletz LO, Jennings

MP. 2014. Selection for phase variation of LOS biosynthetic genes frequently

occurs in progression of non-typeable *Haemophilus influenzae* infection from the

nasopharynx to the middle ear of human patients. PLoS One 9:e90505.

28.   Brockhurst M, Buckling A, Rainey PB. 2006. Spatial heterogeneity and the

stability of host-parasite coexistence. J Evol Biol 19:374–9.

29.   Palmer M, Lipsitch M, Moxon ER, Bayliss CD. 2013. Broad conditions favor the

evolution of phase-variable loci. MBio.

30.   Dixon K, Bayliss CD, Makepeace K, Moxon ER, Hood DW. 2007. Identification

of the functional initiation codons of a phase-variable gene of *Haemophilus

influenzae*, *lic2A*, with the potential for differential expression. J Bacteriol

189:511–21.

31.   Hedrich AW. 1930. The corrected average attack rate from measles among city

children. Am J Epidemiol 11:576–600.

32.   Brisson M, Bénard É, Drolet M, Bogaards JA, Baussano I, Vänskä S, Jit M, Boily

M-C, Smith MA, Berkhof J, Canfell K, Chesson HW, Burger EA, Choi YH, De

Blasio BF, De Vlas SJ, Guzzetta G, Hontelez JAC, Horn J, Jepsen MR, Kim JJ,

Lazzarato F, Matthijsse SM, Mikolajczyk R, Pavelyev A, Pillsbury M, Shafer LA,

Tully SP, Turner HC, Usher C, Walsh C. 2016. Population-level impact, herd

immunity, and elimination after human papillomavirus vaccination: a systematic

review and meta-analysis of predictions from transmission-dynamic models.

Lancet Public Heal 1:e8–e17.

33.   Sumby P, Smith MCM. 2003. Phase variation in the phage growth limitation

system of *Streptomyces coelicolor* A3 (2). J Bacteriol 185:4558–4563.

34.   Gabig M, Herman-Antosiewicz A, Kwiatkowska M, Los M, Thomas MS,

Wegrzyn G. 2002. The cell surface protein Ag43 facilitates phage infection of

Escherichia coli in the presence of bile salts and carbohydrates. Microbiology

148:1533–42.

35.   Broadbent SE, Davies MR, van der Woude MW. 2010. Phase variation controls

expression of *Salmonella* lipopolysaccharide modification genes by a DNA

methylation-dependent mechanism. Mol Microbiol 77:337–53.

36.   Liu M. 2002. Reverse Transcriptase-Mediated Tropism Switching in *Bordetella*

Bacteriophage. Science (80-) 295:2091–2094.

37.   Virji M, Weiser JN, Lindberg AA, Moxon ER. 1990. Antigenic similarities in

lipopolysaccharides of *Haemophilus* and *Neisseria* and expression of a

digalactoside structure also present on human cells. Microb Pathog 9:441–50.

38.   Ehrlich R, Miller S, Idoine LS. 1964. Effects of environmental factors on the

survival of airborn T-3 coliphage. Appl Microbiol 12:479–82.

39.   Dubovi EJ, Akers TG. 1970. Airborne stability of tailless bacterial viruses S-13

and MS-2. Appl Microbiol 19:624–628.

40.   Trouwborst T, de Jong JC. 1973. Interaction of some factors in the mechanism of

inactivation of bacteriophage MS2 in aerosols. Appl Microbiol 26:252–7.

41. Łusiak-Szelachowska M, Zaczek M, Weber-Dąbrowska B, Międzybrodzki R, Kłak

M, Fortuna W, Letkiewicz S, Rogóż P, Szufnarowski K, Jończyk-Matysiak E,

Owczarek B, Górski A. 2014. Phage neutralization by sera of patients receiving

phage therapy. Viral Immunol 27:295–304.

42. Christen M, Beusch C, Bösch Y, Cerletti D, Flores-Tinoco CE, Del Medico L,

Tschan F, Christen B. 2016. Quantitative Selection Analysis of Bacteriophage

φCbK Susceptibility in Caulobacter crescentus. J Mol Biol 428:419–430.

43. Mazzocco A, Waddell TE, Lingohr E, Johnson RP. 2009. Enumeration of

bacteriophages using the small drop plaque assay system. Methods Mol Biol

501:81–85.

44. Kropinski AM. 2009. Measurement of the rate of attachment of bacteriophage to

cells, p. 151–155. *In* Methods in molecular biology (Clifton, N.J.).

45. Cairns BJ, Timms AR, Jansen V a a, Connerton IF, Payne RJH. 2009. Quantitative

models of in vitro bacteriophage-host dynamics and their application to phage

therapy. PLoS Pathog 5:e1000253.

46. Krysiak-Baltyn K, Martin GJO, Stickland AD, Scales PJ, Gras SL. 2016.

Computational models of populations of bacteria and lytic phage. Crit Rev

Microbiol 42:942–68.

FIGURE LEGENDS

FIG 1. Graphic representation of the oscillating prey assay. This diagram illustrates the

methodology for experimental simulation of phage HP1c1 transmission through a 50 %

ON meta-population structure. Green circles represent a phage-sensitive Rd 30S culture

(*lic2A* ON) while pink circles represent a phage-resistant Rd 30R culture (*lic2A* OFF).

(A) Phage are added to an exponential phase culture of the bacterial phase variant with an

OD_{600} of 0.01 to a final MOI of 0.01 (i.e. ~1 x 10^6 PFU/mL). (B) Incubation of phage

with host at 37°C for 50 minutes to allow for one viral replicative cycle. (C) Filtration of

the phage-bacteria mix through a 0.22 μm filter and titration to determine the phage

density at the end of the cycle. (D) Transfer of 600 µL of filtered culture-suspension to

5.4 mL of fresh culture of the appropriate phase variant (in this example Rd 30R). (E)

Incubation of this new culture at 37°C for 50 minutes beyond which the process is

repeated until a total of 20 cycles have been completed.

FIG. 2. Oscillating prey assay for phage HP1c1 infections of *H. influenzae* strain Rd

populations with varying population structures for *lic2A* expression. Each line represents

cycling of the phage through a defined series of cultures of two *H. influenzae* strain Rd

phase variants, namely Rd30S (S; *lic2A* ON; phage sensitive) and Rd 30R (R; *lic2* OFF;

phage resistant). The population structures are indicated in the legend (e.g. S100, all S;

S66, S-S-R-S-S-R; S50, S-R-S-R; etc.). The phage HP1c1 concentration was determined

at the end of each cycle. Circles represent the phage titre observed from each of five

biological replicates, with the line showing the mean of these replicates.

FIG. 3. Mathematical model of the impact of sub-population phage resistance/sensitivity

composition and dilution rate on phage extinction events. This model simulates

transmission of phage HP1c1 through meta-populations of *H. influenzae* strain Rd

comprising either phage-sensitive (S, *lic2A* ON) or phage-resistant (R, *lic2A* ON)

populations for 105 cycles. In these simulations, the order with which phage met either S

or R populations was random but dependent on the probability of encountering a resistant

population (R). Note that the phage population is classified as extinct if the phage titre

falls below the extinction threshold ($P_0 = 100$) with the density being set to 0 for all

remaining subsequent cycles. Panels A and B show examples of model outputs, which

are quantified as phage densities log(P). These graphs show six iterations of meta-

populations comprising either 75% (i.e. $R = 0.25$; panel A) or 85 % (i.e. $R = 0.15$; panel

B) phage-sensitive populations. Panel (C) shows the mean proportion of phage extinction

events occurring in 200 lineages for 200 combinations of probabilities to encounter R

(i.e. 1 = 100 % resistant, 0 = 0 % resistant) and rates of phage loss during transmission

after each cycle (inverse dilution rate, C; 0.02 = 2 % of phage carried through in each

transfer). Colour correlations are shown on the bar to the right of the graph with 1

representing no extinction events while 0 is extinction in all lineages.

FIG. 4. Illustration of phage survival in spatially-structured sub-populations of *lic2A*

phase variants of *H. influenzae*. Phage HP1c1 was passaged through a two-dimensional

array of phage-resistant (R) and phage-sensitive (S) populations. The proportion of phage

sensitive sub-populations (*lic2A* ON) in each fixed area structure were as follows: (A)

100 % *lic2A* ON (S100); (B) 66 % *lic2A* ON (S66); (C) 50 % *lic2A* ON (S50); (D) 50 %

lic2A ON (R50); (E) 33 % *lic2A* ON (R66); and (F) 0 % *lic2A* ON (R100). Each node

represents a well in which phage density was measured. The colour of each node

indicates the concentration of bacteriophage detected in a specific well. Lines indicate the

route taken starting from a central initiator well with line length being proportional to the

number of cycles between each node. All central initiator wells were inoculated with 10^6

PFU/mL. (G) shows a box-plot of the distribution of phage densities for each tested well

of the fixed-area oscillating prey assays. Densities of phage (PFU/mL) obtained from

each test well are represented by a dot for the six distributions of *lic2A* phase variants (x-

axis). The boxed area indicates the first to third quartile, the line is the median of all

points, and whiskers represent 1.5x the interquartile range. Due to the nature of the small-

drop plating methodology employed for phage enumeration the minimum detection

threshold at any time point is 3.3×10^1 PFU/mL.

FIG. 5 Model of the temporal fluctuations in the relative amounts of phage-sensitive and

phage-resistant phase variants for a range of PV rates and selection pressures. PV of the

lic2A gene results in switching between phage-sensitive (S, *lic2A* ON) and phage-

resistant (R, *lic2A* OFF) phase variants. The *lic2A* ON state is however known to mediate

serum resistance (see text). This model examines how the rate of *lic2A* PV (β, ON-to-

OFF switching; α, OFF-to-ON switching; note that switching rates were obtained from

Dixon *et al*. [30]) and the degree of selection (*m*) against the *lic2A* OFF (i.e. R, the

phage-resistant state) expression state influences the relative amounts of the R and S

states in a population. All panels show changes in the proportion of R. Three different

switching rates were examined: $\beta = 1.89 \times 10^{-4}$, $\alpha = 1.13 \times 10^{-4}$ (A); $\beta = 1.89 \times 10^{-5}$, $\alpha =$

1.13×10^{-5} (B); $\beta = 1.89 \times 10^{-3}$, $\alpha = 1.13 \times 10^{-3}$ (C).

FIG 6. Phase variation of phage resistance genes generates 'bacterial herd-immunity' in

bacterial meta-populations. Phase variation of a phage-receptor can generate

heterogeneous bacterial meta-populations containing a mix of sub-populations that are either susceptible or resistant to phage infection. High proportions of resistant sub-populations can both hinder phage spread and protect susceptible populations from phage attack. In this figure, we show the routes of phage dissemination through four different bacterial meta-populations. Green circles represent phage-sensitive populations while pink circles are phage-resistant populations i.e. *lic2A* ON and OFF populations in our experimental system. Lines with arrowheads show infection events leading to successful phage replication, while lines with barheads represent directions where phage-resistant populations act as a barricade blocking phage spread. Regions free of arrows are those free of phage. Phage spread is shown for: (A) a 100% phage-sensitive (*lic2A* ON) sub-population; (B) 66% phage-sensitive (*lic2A* ON) sub-populations; (C) 66% phage-resistant (*lic2A* OFF) sub-populations; and (D) 100% phage-resistant (*lic2A* OFF) sub-populations.

Supplemental material

Fig. S1 Parameter setting experiments for mathematical modelling.

Fig. S2 Fit of mathematical model to experimental data.

Fig. S3 Distribution of wells receiving a *lic2A* ON or *lic2A* OFF phase variant of *H. influenzae* strain Rd for the 50 % ON (panel A) and 66 % ON (panel B) population structures.

Fig. S4 Sampling and transfer regimes for testing phage expansion over a fixed area.

Text S1. Modelling the dynamics of PV of the *lic2A* gene of *H. influenzae*

Fig. S5 Putative ON/OFF state of the *lic2A* gene from 104 *H. influenzae* strains.

Fig. 1 Graphic representation of the oscillating prey assay.

Fig. 2. Oscillating prey assay for phage HP1c1 infections of *H. influenzae* strain Rd populations with varying population structures for *lic2A* expression.

Fig. 3. Mathematical model of the impact of sub-population phage resistance/sensitivity composition and dilution rate on phage extinction events.

Fig. 4. Illustration of bacteriophage survival in spatially-structured sub-populations of *lic2A* **phase variants of** *H. influenzae***.**

Fig. 5 Model of the temporal fluctuations in the relative amounts of phage-sensitive and phage-resistant phase variants for a range of PV rates and selection pressures.

Fig 6. Phase variation of phage resistance genes generates 'bacterial herd-immunity' in bacterial meta-populations