Variability in β-catenin pulse dynamics in a stochastic cell fate decision in *C. elegans*

Jason R. Kroll[*,1], Jasonas Tsiaxiras[*], and Jeroen S. van Zon[*]

[*]Department of Living Matter, AMOLF, 1098 XG Amsterdam, The Netherlands

[1]Present address: Developmental Biology, Department of Biology, Faculty of Sciences, Utrecht University, 3584 CS Utrecht, Netherlands

Abstract

10   During development, cell fate decisions are often highly stochastic, but with the

frequency of the different possible fates tightly controlled. To understand how signaling

12   networks control the cell fate frequency of such random decisions, we studied the stochastic

decision of the *Caenorhabditis elegans* P3.p cell to either fuse to the hypodermis or assume

14   vulva precursor cell fate. Using time-lapse microscopy to measure the single-cell dynamics of

two key inhibitors of cell fusion, the Hox gene LIN-39 and Wnt signaling through the β-catenin

16   BAR-1, we uncovered significant variability in the dynamics of LIN-39 and BAR-1 levels. Most

strikingly, we observed that BAR-1 accumulated in a single, 1-4 hour pulse at the time of the

18   P3.p cell fate decision, with strong variability both in pulse slope and time of pulse onset. We

found that the time of BAR-1 pulse onset was delayed relative to the time of cell fusion in

20   mutants with low cell fusion frequency, linking BAR-1 pulse timing to cell fate outcome. Overall,

a model emerged where animal-to-animal variability in LIN-39 levels and BAR-1 pulse dynamics

22   biases cell fate by modulating their absolute level at the time cell fusion is induced. Our results

highlight that timing of cell signaling dynamics, rather than its average level or amplitude, could

24   play an instructive role in determining cell fate.

Article summary

We studied the stochastic decision of the *Caenorhabditis elegans* P3.p cell to either fuse to the hypodermis or assume vulva precursor cell fate. We uncovered significant variability in the dynamics of LIN-39/Hox and BAR-1/β-catenin levels, two key inhibitors of cell fusion. Surprisingly, we observed that BAR-1 accumulated in a 1-4 hour pulse at the time of the P3.p cell fate decision, with variable pulse slope and time of pulse onset. Our work suggests a model where animal-to-animal variability in LIN-39 levels and BAR-1 pulse dynamics biases cell fate by modulating their absolute level at the time cell fusion is induced.

Introduction

36   During development, cells robustly obtain the correct cell fate to give rise to a viable

adult organism, despite internal molecular noise and environmental variability. It is commonly

38   assumed that suppressing this variability is essential for successful development. However,

stochastic cell fate decisions, where cells randomly assume one cell fate out of a limited

40   repertoire of different fates, are the cornerstone of many developmental processes (Johnston

and Desplan 2010). For example, heterogeneous and stochastic gene expression between cells

42   in the four-cell mouse embryo influences and restricts their cell fates (Goolam et al. 2016).

Similarly, photoreceptor cells in the human retina randomly express either a red, green or blue

44   photoreceptor gene (Smallwood, Wang, and Nathans 2002; Roorda and Williams 1999). In

these stochastic decision processes, even though each individual outcome is random, the

46   relative frequency of the different cells fates is often tightly controlled.

      Currently, stochastic cell fate decisions are best understood in the context of single-

48   celled organisms, where gene expression noise dominates as the key source of variability

driving stochastic cell fate decisions (Losick and Desplan 2008; Balaban 2004; Maamar, Raj,

50   and Dubnau 2007; Süel et al. 2006). However, it is unclear how stochastic cell fate decisions

are regulated during animal development, as multicellular organisms pose unique constraints

52   compared to single-celled organisms. Here, stochastic cell fate decisions have to be precisely

coordinated with developmental timing, are potentially influenced by neighboring cells and rely

54   on external, long-range signals mediated by a small number of key developmental signaling

pathways. How these canonical signaling pathways drive stochastic cell fate decisions with

56   strong control over cell fate frequencies is an open question.

      Even though *C. elegans* development occurs in a largely invariant manner (Sulston et al.

58   1983), some cell fate decisions occur in a stochastic manner. One such decision is the

specification of the vulval precursor cell (VPC) competence group, beginning early in the L2

60 larval stage (Myers and Greenwald 2007; Gleason, Korswagen, and Eisenmann 2002). This

group consists of six epidermal cells named P3.p-P8.p, which are subsequently patterned to

62 various vulval cell fates by multiple signaling pathways (Gupta, Hanna-Rose, and Sternberg

2012; Hill and Sternberg 1993; D M Eisenmann et al. 1998; Félix 2012; Sternberg and Horvitz

64 1986; Gleason, Korswagen, and Eisenmann 2002). The establishment of the VPC competence

group is partly stochastic, as the P3.p cell assumes VPC fate in roughly 50% of wild-type (N2)

66 hermaphrodites (Fig. 1a), while in the remainder P3.p assumes hypodermal fate by fusing to a

neighboring syncytial hypodermal cell, called hyp7 (Gidi Shemer and Podbilewicz 2002;

68 Sternberg and Horvitz 1986). Moreover, the tendency for the P3.p cell to fuse or not in a given

strain is sensitive to differences in environmental conditions and genetic backgrounds (Braendle

70 and Félix 2008; Pénigault and Félix 2011a).

The Wnt pathway is a highly conserved signaling pathway that regulates many

72 developmental events and cell fates (Ohyama 2006; Hudson et al. 2013; Hirabayashi 2004;

Mucenski et al. 2003; Lindström et al. 2014; Clevers and Nusse 2012). Previous investigations

74 into the P3.p cell fate decision showed that its cell fate frequency is extremely sensitive to the

dosage of Wnt ligands, particularly *cwn-1*, suggesting that variability in the ligand concentration

76 or in the response of the P3.p cell to Wnt ligands could provide the noise source driving the

stochastic fate decision (Pénigault and Félix 2011a; 2011b). In the canonical pathway, the

78 presence of Wnt ligands leads to the accumulation of the transcriptional co-activator BAR-1/β–

catenin, which co-regulates Wnt pathway target genes (Sawa and Korswagen 2013; Hendrik C.

80 Korswagen 2002; D M Eisenmann et al. 1998; H C Korswagen, Herman, and Clevers 2000). In

addition to the Wnt pathway, mutations in the *C. elegans* Hox gene *lin-39* impact the Pn.p cell

82 fate frequencies, by repression of cell fusion and promoting division of VPC fate cells (Koh et al.

2002; Gidi Shemer and Podbilewicz 2002; Maloof and Kenyon 1998; Roiz et al. 2016; Clark,

84 Chisholm, and Horvitz 1993). Both Wnt signaling and LIN-39 inhibit hyp7/fusion fate, with loss-

of-function mutants exhibiting increased frequency of cell fusion, including in the P4.p-P8.p cells

86   that otherwise never assume hyp7/fusion fate (Gleason, Szyleyko, and Eisenmann 2006; Myers

and Greenwald 2007). However, what aspects of Wnt signaling and LIN-39 dynamics control

88   the frequency of hyp7/fusion versus VPC fate in P3.p remains unknown.

Here, we use a novel time-lapse microscopy approach (Gritti et al. 2016) to observe

90   gene expression and signaling dynamics in single Pn.p cells during specification of the VPCs,

allowing us to directly connect variability during the decision process to the final cell fate

92   outcome. Using this approach, we observed animal-to-animal variability in LIN-39 and BAR-1

dynamics that could be linked to cell fate outcome. In particular, we found that BAR-1

94   accumulated in a single, ~1-4 hr pulse in the Pn.p cells at the time of the hyp7/fusion versus

VPC fate decision, with strong variability in pulse slope and timing. Using experiments in

96   mutants combined with quantitative data analysis, we arrived at a model where stochastic

variability in LIN-39 levels and BAR-1 pulse dynamics biases cell fate by modulating their

98   absolute level at the time cell fusion is induced.

Results

Time-lapse microscopy of a stochastic cell fate decision

So far, whether P3.p undergoes fusion or assumes VPC fate in wild-type or mutant

104 animals has been assessed only after the process has completed (Myers and Greenwald 2007;

D M Eisenmann et al. 1998; S Alper and Kenyon 2001; Scott Alper and Kenyon 2002; Pénigault

106 and Félix 2011b; 2011a; Chen and Han 2001). To correlate early stochastic molecular events to

eventual cell fate outcome it is essential to follow these processes directly in time. Here, we

108 utilize a fluorescent time-lapse microscopy approach we developed recently to study single-cell

dynamics inside moving and feeding *C. elegans* larvae for their entire ~40 hr development (Gritti

110 et al. 2016). We tested whether we could directly observe P3.p fusion events in single animals.

We initially used two measures of cell fusion: first, imaging the apical junction protein AJM-1,

112 which localizes on the apical junction of Pn.p cells but is displaced upon cell fusion and has

been used as a cell fusion marker in other studies, including live imaging applications (Brabin,

114 Appleford, and Woollard 2011; G Shemer et al. 2004). Using animals carrying AJM-1::GFP, we

found that the fusion event occurred in 28% of animals during the L2 stage (n= 57 total animals,

116 see Table 1), and could detect disruption of AJM-1 and its subsequent loss, indicating that the

P3.p fusion fate is indeed stochastic between animals in the same strain. The *ajm-1::mCherry*

118 strain showed similar results (Table 1). In both strains, the cell fusion fate of the P3.p cell was

reduced from the canonical 50% ratio, possibly resulting from the fluorescent AJM-1 markers, or

120 the environmental conditions of our time-lapse microscope (Braendle and Félix 2008). Second,

as an independent measure of cell fusion, we directly observed the flow of GFP from the

122 hypodermis into the fused P3.p cell, using animals carrying an extrachromosal array targeting

GFP expression to the hyp7 hypodermal syncytium, while simultaneously monitoring the apical

124 junction marker AJM-1::mCherry. As seen before, the AJM-1::mCherry signal expanded along

the A-P axis during the early L2 larval stage (Supplementary Movies S1-S2, Fig. 1c). In animals

126   with a fusing cell, this was followed by a sudden and pronounced ruffling of the AJM-1:mCherry

signal and a rapid retraction of AJM-1::mCherry towards the posterior, with its fluorescent signal

128   fully disappearing from P3.p within 1 hr (Fig. 1c). Inflow of GFP from the hypodermis into P3.p

was observed as soon as AJM-1::mCherry retraction commenced (Fig. 1c,d), showing that both

130   are accurate markers of (the time of) fusion. Because AJM-1 fluorescence was easily monitored

in combination with other quantification, we used AJM-1 dynamics to establish fate and timing of

132   P3.p fusion for all subsequent experiments.

Even though changes to the frequency of P3.p hyp7/fusion versus VPC fate in mutants

134   are well studied (Pénigault and Félix 2011b; 2011a; Chen and Han 2001; Myers and Greenwald

2007; D M Eisenmann et al. 1998; S Alper and Kenyon 2001; Scott Alper and Kenyon 2002), it

136   was not known how such mutants impact the timing of this decision. We quantified the time of

P3.p fusion in wild-type animals and found that cell fusion occurred in a relatively narrow time

138   window between 40-60% of the L2 larval stage (Fig. 1e). We then examined mutants in which

fusion frequency is increased by removing inhibitory Wnt signaling or LIN-39 (Fig. 1b). We found

140   that in these mutants P3.p fusion occurred within the same time window as wild-type animals,

with only small differences between wild-type and mutant animals in average timing (Fig. 1e).

142   Strikingly, even though the exact time of fusion can vary as much as 2 hrs between animals,

when multiple VPCs fused in a single animal, they typically did at the same time (Fig. 1f). In the

144   absence of key repressors of hyp7/fusion fate, cell fusion frequency is increased independently

of its timing, providing evidence that hyp7/fusion inhibitors (Wnt signaling, LIN-39) do not control

146   timing of fusion, but rather modulate hyp7/fusion frequency. This also suggests that a yet-

unknown signal exists that activates cell fusion at the appropriate time.

Figure 1. **Stochastic cell fate decisions in Pn.p cells.**

(A) Overview of the hyp7/fusion versus vulva precursor cell fate (VPC) decision in the P(3-8).p cells. Cells assuming hyp7/fusion fate fuse (indicated by the dashed line) with the hypodermal syncytium hyp7 and lose the AJM-1 apical junction marker (green). Cell fusion requires the expression of the fusogen EFF-1 and is inhibited by the Hox protein LIN-39 and Wnt signaling through the β-catenin BAR-1. BAR-1 accumulation is induced by binding of Wnt ligands, such as CWN-1 (purple) to Wnt receptors (magenta).

(B) Measured hyp7/fusion frequencies in Pn.p cells in wild-type and mutant backgrounds. Mutants carried the *ajm-1::GFP* reporter except for *lin-39(lf)* which carried *ajm-1::mCherry*. Wild-type animals carried

158 either *ajm-1:: GFP* (shown here) or *ajm-1::mCherry* (Table 1), with no differences in fusion frequencies.

(C) AJM-1 dynamics in non-fusing (top) and fusing (bottom) P3.p cells carrying a nuclear *dpy-7p:mCherry*

160 marker. Cell fusion occurred 6h 20m after the start of L2, as shown by the appearance of GFP from the

hypodermal syncytium hyp7 in the P3.p nuclear area (region enclosed by yellow line). Simultaneously,

162 AJM-1 showed a pronounced ruffling, followed by its removal from P3.p. In contrast, no such AJM-1

dynamics were observed in non-fusing cells assuming VPC fate. **(D)** Comparing GFP inflow from the

164 hyp7 syncytium in fusing and non-fusing cells as a function of time after the start of the L2 larval stage.

Shown is the ratio of GFP fluorescence intensity between P3.p and P4.p in the same animal, where P4.p

166 never fused. The blue and red line corresponds to the non-fusing and fusing cell in (C). Symbols

correspond to the time points shown in (C). Arrow indicates the time of AJM-1 ruffling and coincides

168 exactly with inflow of GFP into the fusing cell. **(E)** Individual cell fusion times and box-and-whisker plots

for P3.p (green) and P4.p cells (magenta) in different genetic backgrounds. Fusion time was determined

170 for AJM-1 dynamics and is expressed as fraction of the L2 larval stage duration T_{L2} (~8-12 hrs for all

backgrounds). Even though small but significant differences exist in average fusion time between strains,

172 (one-way ANOVA to determine a difference between all strains, followed by Student's t-test, * indicates P

<0.05 in the Student's t-test), the full distributions show extensive overlap. **(F)** Distribution of difference in

174 cell fusion time between P3.p and P4.p cells, where both cells fuse (data pooled for all genotypes with

double fusions).

Stochastic *eff-1* induction precedes P3.p fusion

178   The most downstream regulator of the hyp7/fusion versus VPC fate decision is the gene

eff-1, a fusogenic protein whose expression is sufficient to induce cell fusion (Mohler et al. 2002;

180 G Shemer et al. 2004). EFF-1 is a transmembrane protein that is required for most cell fusions

in *C. elegans*, and must be present on both the Pn.p and hyp7 plasma membrane to induce

182 fusion (Zeev-Ben-Mordehai et al. 2014; Smurova and Podbilewicz 2016). To understand if cell

fate frequency is regulated on the level of *eff-1* expression, and how it might be regulated by the

184   Hox/LIN-39 and Wnt signaling pathways, we counted *eff-1* mRNA molecules in Pn.p cells,using

single molecule FISH (smFISH) (Raj et al. 2008). In wild-type animals in the late L1 and early L2

186   stage (230 - 325 μm in body length), stages of development immediately preceding the

hyp7/fusion versus VPC fate decision, we found that all P3.p cells were unfused, as determined

188   by the presence of the AJM-1 signal. They also exhibited low *eff-1* expression in P3.p, <10

molecules (Fig. 2a,d), similar to the P4.p cell that always assumes VPC fate in wild-type

190   animals. However, in older, mid-L2 stage animals (>325 μm in length), corresponding to the

time of fusion, we observed a subset of animals expressing ~30-50 molecules in unfused P3.p

192   cells, something not observed in P4.p (Fig. 2b,d). In fused P3.p cells, we found similar number

of *eff-1* mRNA molecules located in close proximity to the cell nucleus, suggesting that high *eff-*

194   *1* expression is maintained after fusion, before finally disappearing by the end of the L2 stage

(Supplemental Fig. S1a,b). We confirmed that high *eff-1* expression preceded cell fusion by

196   examining a temperature-sensitive loss-of-function point mutation in *eff-1* (Mohler et al. 2002).

Here, we still found high *eff-1* mRNA levels in P3.p at the restrictive temperature, even though

198   cell fusion was blocked (Supplemental Fig. S1c).

        The fraction of wild-type animals showing high (>20 molecules) *eff-1* mRNA levels in

200   P3.p was significantly smaller than the expected fraction of animals where P3.p assumes

hyp7/fusion fate. For our analysis, we randomly sampled animals within the time window we

202   expected fusion to occur. Given the observed variability in the time of fusion (Fig. 1f), it is

expected that some animals with low *eff-1* expression would have ultimately fused at a later

204   point. In particular, the fraction of animals observed with high *eff-1* expression in unfused P3.p

cells should increase with the average duration the cell expresses high *eff-1* before this results

206   in fusion. Hence, our results suggested that induction of high *eff-1* expression was quickly

followed by cell fusion.

Figure 2.**Stochastic *eff-1* expression in Pn.p cells precedes cell fusion.**

(A),**(B)** Example of (A) low and (B) high *eff-1* expression preceding cell fusion in wild-type animals of similar age, as determined by body length. Single *eff-1* mRNA molecules (red) were visualized using smFISH and nuclei with DAPI (blue). Unfused cells were selected based on AJM-1::GFP localization (green), with white brackets indicating intact apical junctions typical of unfused cells. **(C)** Panels (i) and (ii) show details of the P3.p cell region, corresponding to the area delineated by the dashed line in panels (A) and (B). **D)** Number of *eff-1* mRNA molecules in unfused P3.p and P4.p cells, for wild type animals (n=184) and mutants with decreased (*lin-39(++), n=99*) and increased (*cwn-1(0), n=81*) cell fusion frequency. Late L1 and early L2 animals (230-325 μm body length) are at a developmental stage preceding the hyp7/fusion versus VPC fate decision, with fusion occurring only in mid-late L2 animals (>325 μm body length). In each panel, the fusion rate of that cell in the indicated genotype is indicated as a percentage.

To understand how *eff-1* expression is correlated to cell fate frequency, we quantified *eff-1* levels in a strain with a functional LIN-39::GFP insertion that caused low fusion frequency

(*lin-39(++)*, ~2% P3.p fusion frequency). Consistently, we no longer observed unfused P3.p

226   cells expressing high (>20 molecules) *eff-1* levels (Fig. 2d). In contrast, in the *cwn-1(0)* mutant

that lacks the dominant Wnt ligand and exhibited high (>80%) fusion frequency both in P3.p and

228   P4.p, we found that the numbers of unfused cells observed with high *eff-1* expression had

increased substantially, most strikingly in P4.p (Fig. 2d). Together, this indicates that Wnt

230   signaling and LIN-39 controlled cell fate frequency mainly by tuning the fraction of cells in which

high *eff-1* expression is induced.

Bias of cell fate decision by noise in LIN-39 protein level

234   The Hox transcription factor LIN-39 inhibits hyp7/fusion fate by repressing *eff-1*

expression (Gidi Shemer and Podbilewicz 2002), with *lin-39* null mutations causing all Pn.p cells

236   to fuse in the L1 larval stage (Clark, Chisholm, and Horvitz 1993; Wang et al. 1993). Hence,

stochastic variability in LIN-39 protein levels could result in variability in induction of high *eff-1*

238   expression. It was shown previously that LIN-39 levels are similar between P3.p and P4.p in

early L2 larval stage animals prior to cell fusion (Pénigault and Félix 2011a), even though both

240   cells have a different fate frequency (Fig. 1b). However, LIN-39 dynamics in individual cells

were not followed over time and linked to the eventual cell fate. Hence, it remains possible that

242   small differences in LIN-39 between P3.p cells in different animals are sufficient to explain the

outcome. To connect animal-to-animal variability in LIN-39 level with P3.p cell fate, we

244   performed time-lapse microscopy on animals carrying an integrated low-copy *lin-39::GFP*

translational fusion transgene (Sarov et al. 2012) and *ajm-1::GFP* as a cell fusion marker (Fig.

246   3a,b). Since *lin-39::GFP (lin-39(++))* is present as an insertion, it decreased the P3.p fusion rate

from ~30% to ~2%, making it challenging to capture sufficient P3.p fusion events for analysis.

248   For that reason, we further crossed this reporter into the *cwn-1(0)* mutant, which increased the

P3.p and P4.p fusion rates to 20% and 14%, respectively (Table 1).

250   We observed LIN-39 in the P3.p nucleus at the start of the L2 larval stage and it

remained there for the entire larval stage when P3.p assumed VPC fate (Fig. 3a,c). However, in

252   P3.p cells that fused, nuclear LIN-39 levels decreased rapidly after fusion commenced and fully

disappeared within 90 mins (Fig.3b,c), consistent with past observations of loss of LIN-39 in

254   fused Pn.p cells (Pénigault and Félix 2011a). LIN-39 levels were not significantly different in the

cwn-1(0) mutant compared to wild type, indicating that the Wnt ligand acts in a different pathway

256   from LIN-39 to inhibit eff-1 expression. We compared the distribution of LIN-39 levels, averaged

over 3 hrs prior to fusion in P3.p cells that assumed hyp7/fusion fate, with the distribution in

258   P3.p cells that assumed VPC fate, averaged over 3 hrs prior to the average time of P3.p fusion

in this strain (Fig. 3d). We found strong overlap between the two distributions, also when

260   changing the size of the time window (data not shown), making it unlikely that fluctuations in

LIN-39 levels are the main driver of eff-1 induction and cell fusion.

262   These results leave open the question whether the observed variability in LIN-39 has

any effect on the P3.p cell fate outcome. In lin-39::GFP; cwn-1(0) animals, P4.p also assumed

264   hyp7/fusion fate in a stochastic manner, allowing us to test whether differences in LIN-39 levels

between P3.p and P4.p within the same animal are correlated with their eventual fate. First, we

266   established that in this strain LIN-39 distributions for P3.p and P4.p were similar and also

showed substantial overlap between fusing and non-fusing cells. We then selected animals in

268   which one cell, either P3.p or P4.p, fused but the other assumed VPC fate (Fig. 3e). Indeed, in

these animals, absolute LIN-39 level was not predictive of the eventual fate of P3.p or P4.p, but

270   the difference in LIN-39 levels between P3.p and P4.p was predictive of their fate (Fig. 3f), with

fusing cells having lower LIN-39 levels than their non-fusing neighbor cell (P = 0.0095, Fisher's

272   Exact Test). This shows that variability in LIN-39 levels is correlated to the hyp7/fusion versus

VPC fate decision, but likely in conjunction with another source of variability.

276 Figure 3. **Variability in LIN-39 protein level.**

278 **(A), (B)** Image sequence of a cell assuming (A) VPC or (B) hyp7/fusion fate. Nuclear LIN-39 levels are

visualized using a *LIN-39::GFP* protein fusion, while cell fusion is determined using an *ajm-1::GFP*

280 reporter. The red star indicates cell fusion has occurred. Times are relative to the start of the L2 stage. **(C)**

Nuclear LIN-39 fluorescence in fusing (red) and non-fusing (blue) P3.p cells. For fusing P3.p cells, LIN-39

282 levels are variable prior to fusion (dark red) and fall rapidly after fusion (light red). **(D)** Distribution of

nuclear LIN-39 fluorescence in fusing (red) and non-fusing (blue) P3.p cells show strong overlap. In

284 fusing cells, fluorescence was averaged over 3 hour window directly prior to fusion, whereas in non-fusing

cells a time window of 3 hours prior to the average time of cell fusion was used (N = 48 total animals) **(E)**

286 Nuclear LIN-39 fluorescence in P3.p (green) and P4.p (magenta) in an animal in which P3.p but not P4.p

assumes hyp7/fusion fate. The arrow indicates the time window over which LIN-39 fluorescence is

288 averaged in (D) and (F). The dashed line indicates the time of fusion. **(F)** Nuclear LIN-39 fluorescence

levels in P3.p and P4.p for animals in which P3.p assumes hyp7/fusion fate and the P4.p assumes VPC

290 fate (black circle) or the reverse (yellow diamond), see legend for schematic of P3.p and P4.p fates,

dashed circle indicates fusion. Each symbol corresponds to a single animal and is averaged over a 3 h

292 time window prior to cell fusion. In general, LIN-39::GFP levels were lower in the cell that assumed

hyp7/fusion fate.

β-catenin activation dynamics during the cell fate decision

296 Upon activation of Wnt receptors by Wnt ligands, β-catenin accumulates in the cell and

nucleus (Sawa and Korswagen 2013). In the absence of Wnt ligands, β-catenin is degraded by

298 a degradation complex. In P3.p, the presence of the β-catenin BAR-1 is required to inhibit *eff-1*

expression and, hence, inhibit hyp7/fusion fate (D M Eisenmann et al. 1998). In contrast to β–

300 catenins involved in the Wnt asymmetry pathway (Mila et al. 2015; Park and Priess 2003),

dynamics of BAR-1 during canonical Wnt signaling is poorly characterized.

302 We aimed to quantify activation of the Wnt pathway by monitoring the accumulation

dynamics of BAR-1/β-catenin. We first endogenously tagged the C-terminus of BAR-1 with GFP

304 using CRISPR/Cas9 genome editing, (see Supplemental Methods) (Vicencio et al. 2019). The

fusion rate when the strain was crossed to the *ajm-1::mCherry* marker was similar to other

306 controls, (see Table 1), indicating that tagging BAR-1 with GFP has no detrimental effect on its

function. However, no fluorescence was detected in the Pn.p cells with the laser intensities and

308 exposure times required for our time-lapse method. To visualize BAR-1 dynamics, we instead

used a multi-copy integrated BAR-1::GFP transgene, *gaIs45*, which rescues the *bar-1(0)*

310 phenotype (D M Eisenmann et al. 1998) and has been used previously to study BAR-1

dynamics during male hook development (Yu et al. 2009). Similar transgenes have been used

312 extensively to study the dynamics of the *C. elegans* β-catenins SYS-1 (Robertson et al. 2017)

and WRM-1 (Kim et al. 2013). A disadvantage of using a multi-copy insertion is that the

314   elevated BAR-1 level perturbs the observed P3.p fate frequency. However, a key advantage is

its increased fluorescence signal, which was crucial for imaging its dynamics using our time-

316   lapse imaging approach.

Wnt ligand gene expression levels and protein distribution are considered to be largely

318   constant in time (Pani and Goldstein 2018; Coudreuse et al. 2006). Hence, we expected BAR-1

to show constant expression and dynamics in P(3-8).p cells, similar to LIN-39. Instead, we

320   found that BAR-1::GFP levels were strikingly dynamic, with no BAR-1::GFP in P(3-8).p at the

start of the L2 stage, followed by steady accumulation of BAR-1::GFP in P(3-8).p at the mid-L2

322   stage that lasted 1-4 hours and was strongly coordinated between cells (Supplemental Movie

S3, Fig. 4a-c). After the accumulation phase, BAR-1 was rapidly degraded, with the overall

324   dynamics of BAR-1 resembling a single pulse. The protein was detected in both the nucleus and

cytoplasm. We found stochastic variability in the amplitude of the BAR-1 pulse between different

326   Pn.p cells (Fig. 4b,c). It was speculated that P3.p, which is considered most distant to the

source of Wnt ligands, receives a lower Wnt signal than P(4-8).p, thereby resulting in its

328   occasional hyp7/fusion fate (Harterink et al. 2011; Pénigault and Félix 2011b). However, the

BAR-1::GFP accumulation pulse in P3.p was frequently of similar or higher amplitude compared

330   to the other Pn.p cells and, in general, we found no sign of a systematic spatial pattern in

downstream Wnt signaling. We also found significant variability in the amplitude and timing of

332   the BAR-1::GFP pulse in the P3.p cell compared between different animals (Fig. 4d).

Figure 4. **BAR-1 dynamics during cell fate decision.**

(A) Image sequence of BAR-1::GFP dynamics (green) over time in P3.p and P4.p cells of a *bar-1*(++); *ajm-1::mCherry* animal, with both cells assuming VPC fate. **(B),(C)** Dynamics of cellular BAR-1::GFP fluorescence in P(3-6).p cells in two different animals. The animal in (B) corresponds to images in (A). Different cells show similar timing of BAR-1 pulse onset, but vary in amplitude. **(D)-(F)** Examples of BAR-1::GFP dynamics in non-fusing P3.p cells in (D) *bar-1(++)*, (E) *bar-1(++);lin-39(lf)*, (F) *bar-1(++);cwn-1(0)* animals. Traces for different animals are shifted along the vertical axis for clarity. Pulses show strong differences in time of onset and duration, even within the same strain. Traces run until the end of the L2 larval stage. Some *bar-1(++);cwn-1(0)* animals displayed significantly extended L2 stage duration (~15-20hr). Total animals imaged for *bar-1(++)*, *bar-1(++);lin-39(lf),* and *bar-1(++);cwn-1(0)* genotypes are n= 30, 64, and 70 animals, respectively.

346   Since the BAR-1::GFP reporter is integrated as a functional multi-copy transgene, we

expected this strain to act as a BAR-1 overexpression mutant and, indeed, observed no P(3-

348   8).p cell fusions (n=30 animals, Table 1). Hence, we refer to this strain as *bar-1(++)*. To study

how BAR-1 dynamics relates to cell fate frequency, we used different approaches to increase

350   the frequency of hyp7/fusion fate. First, we decreased the level of the inhibitor LIN-39, using the

lin-39(n709) temperature sensitive loss-of-function mutant (Table 1). We found that *bar-*

352   *1(++);lin-39(lf)* animals showed similar BAR-1::GFP dynamics (Fig. 4e). Next, we sought to

lower BAR-1 levels in the *bar-1(++)* background by decreasing activity of the Wnt signaling

354   pathway, using the *cwn-1(0)* mutant that lacks the CWN-1 Wnt ligand (Fig. 1a). In these

animals, the BAR-1::GFP pulse amplitude was lower (Fig. 4f), and in some animals the L2 stage

356   was significantly lengthened. Moreover, we found that the BAR-1::GFP pulse occurred at

significantly later times, as a fraction of larval stage duration, even in animals with a L2 larval

358   stage duration similar to wild-type. The BAR-1 pulse also showed considerable variability in

timing and amplitude in these mutants. Finally, in both strains, fusing Pn.p cells did so during

360   the early accumulation phase of the BAR-1 pulse (Fig. 5a, n=29 cells). Even though the

increased BAR-1 protein levels in the *bar-1(++);lin-39(lf)* and *bar-1(++);cwn-1(0)* mutants do not

362   represent the wild-type condition, these animals still undergo a stochastic hyp7/fusion fate

decision, allowing us to correlate pulse dynamics with cell fate outcomes.

364   ## Variability in β-catenin dynamics

To characterize variability in BAR-1 accumulation dynamics, we used a minimal

366   parameterization of the BAR-1 pulse shape to fit to the experimental data (Fig. 5a, Eq. 3 in

Methods). Briefly, we assume that prior to the BAR-1 pulse, Wnt signaling is inactivated and

368   BAR-1 is degraded. At pulse onset time t_0, Wnt signaling is activated, leading to inhibition of

BAR-1 degradation and hence linear accumulation of BAR-1 in the cell. Linear BAR-1

370   accumulation continues for a pulse duration T in cells that assume VPC fate or until the time of

fusion, t_{fusion}, in cells that assume hyp7/fusion fate. Upon fusion BAR-1 vanishes immediately,

372   as observed experimentally, whereas in cells that assume VPC fate, BAR-1 levels decrease

exponentially once the pulse ends. This fitted the experimental data surprisingly well (Fig. 5a,b).

374   Moreover, it allowed us to describe each BAR-1 pulse by three parameters: pulse onset time t_0,

pulse slope s and pulse duration T for VPC cells or fusion time t_{fusion} for hyp7/fusion cells.

376   First, we compared the distribution of pulse onset time t_0 and linear slope s of each

BAR-1 accumulation pulse between strains. We found that both were highly variable among

378   animals (Fig. 5c,d). We found that the pulse parameters of *bar-1(++)* and *bar-1(++);lin-39(lf)*

were similar, consistent with increase in hyp7/fusion fate frequency in this mutant resulting from

380   the absence of the fusion inhibitor LIN-39 rather than changes in Wnt signaling. In contrast, we

assumed that the increase in frequency of hyp7/fusion fate in *bar-1(++);cwn-1(0)* animals was

382   due to a decrease in BAR-1 level, given that BAR-1 accumulation is thought to be proportional

to the amount of external Wnt ligands. Surprisingly, we found that the pulse slope distribution

384   was highly similar for *bar-1(++)* and *bar-1(++);cwn-1(0)* animals (Fig. 5c) and that the only

different characteristic was the delayed pulse onset (Fig. 5d).

Figure 5. **Variability in BAR-1 dynamics.**

(A) Quantifying BAR-1 pulse dynamics in a *bar-1(++);lin-39(lf)* animal. Symbols are measured BAR-1::GFP levels in P3.p (green) and P4.p (magenta). Here, P3.p, but not P4.p, assumed hyp7/fusion fate. Solid lines represent a model of BAR-1 accumulation dynamics (Eq. 3 in Methods). Using the model, each BAR-1 pulse is described by three key parameters: pulse onset time t_0, pulse slope s and pulse duration T (for non-fusing VPCs) or time of fusion t_{fusion} (for hyp7/fusion cells). **(B)** Animals showing differences in (relative) timing of BAR-1 accumulation pulses. **(C), (D)** Distribution of (C) pulse slope s and (D) pulse onset time t_0 in P3.p cells, in *bar-1(++)* (purple, n= 27 animals), *bar-1(++);lin-39(lf)* (orange, n= 30 animals) and *bar-1(++);cwn-1(0)* mutants (green, n= 17 animals). T_{L2} is the L2 larval stage duration. Strains differ in BAR-1 pulse onset time rather than pulse slope. **(E)** Correlation in pulse onset time t_0 between P3.p and P4.p cells. Each symbol corresponds to a P3.p and P4.p cell in a single animal where both assumed VPC fate. Arrows indicate the animals in (B). **(F)** Correlation between pulse onset time t_0

and pulse duration T in non-fusing P3.p (circles) and P4.p cells (triangles). The dashed line is $T/T_{L2} = 0.66 - t_0/T_{L2}$. BAR-1 pulses that start later in the L2 larval stage have a shorter duration.

402   In general, we found that pulse onset time t_0 was correlated between P3.p and P4.p in

the same animal, meaning that if BAR-1 accumulation started late in the L2 larval stage in P3.p,

404   it was also likely to start late in P4.p (Fig. 5e). However, in *bar-1(++)* and *bar-1(++);lin-39(lf)*

animals we still observed significant variability in pulse onset time between P3.p and P4.p in the

406   same animal, with BAR-1 accumulation in P3.p preceding that in P4.p as often as the reverse

(Fig. 5e). Strikingly, in the *bar-1(++);cwn-1(0)* mutant the variability in pulse onset time between

408   P3.p and P4.p was almost completely removed, with the onset of BAR-1 accumulation occurring

in P3.p and P4.p within 20 min in all animals (Fig. 5e). This result suggests that the Wnt ligand

410   *cwn-1* not only controls the average onset of BAR-1 pulses, but it also induces variability in

pulse onset time between P3.p and P4.p cells in the same animal.

412   We then asked whether the duration of the BAR-1 pulse depended on the time of pulse

onset. Because the duration of the L2 larval stage varied significantly between strains and

414   animals, we examined the pulse onset time t_0/T_{L2} and duration T/T_{L2} relative to the duration of

the larval stage, T_{L2}. In this case, we found a striking anti-correlation, with late pulse onset

416   resulting in shorter pulses (Fig. 5f). In fact, the data for all strains clustered along the line

$T/T_{L2} = 0.66 - t_0/T_{L2}$, meaning that the end of the BAR-1 accumulation occurs at 66% of the

418   L2 larval stage independent of the BAR-1 pulse onset time. This correlation also held for the

bar-1(++);cwn-1(0) mutant, where not only the onset of the BAR-1 pulse was delayed but also

420   the L2 larval stage was much extended in some animals. Overall, this indicated that the BAR-1

pulse ended at specific point in the L2 larval stage, with much less variability in timing compared

422   to start of the pulse.

Bias of cell fate decision by absolute BAR-1/β-catenin level

The observation of BAR-1 pulse dynamics raised the question how this controlled cell fate outcome. It is often assumed that the outcome of Wnt signaling depends on the absolute β-catenin level detected by the cell (Sawa and Korswagen 2013). In this case, both changes to BAR-1 pulse slope and pulse onset could bias hyp7/fusion frequency, by modulating absolute BAR-1 levels at the time the decision to fuse or not is made. However, it has also been suggested that cells respond to fold change in rather than absolute level of β-catenin (Goentoro and Kirschner 2009). In this case, cell fate outcome would depend only on BAR-1 pulse slope. Finally, cells could respond to more complex features of BAR-1 pulse dynamics.

To address this question, we first asked whether the change in hyp7/fusion frequency we observed between different strains was accompanied by systematic changes in pulse dynamics. Indeed, while *bar-1(++);cwn-1(0)* animals exhibited the same slope distribution as *bar-1(++)* animals (Fig. 5c), their BAR-1 pulse started at a later point in larval stage compared to *bar-1(++)* and *bar-1(++);lin-39(lf)* animals (Fig. 5d). This already suggested that the onset of the BAR-1 pulse was delayed compared to the time of fusion in this strain, but left open the possibility that the time of fusion was also delayed. Therefore, we attempted to directly measure the delay between pulse onset and time of fusion in these strains.

444   Figure 6. **Impact of BAR-1 dynamics on cell fate decision.**

(A) Correlation between pulse onset time t_0 in a VPC and fusion time t_{fusion} in a hyp7/fusion cell. Circles

446   correspond to animals where P3.p, but not P4.p, assumed hyp7/fusion fate and triangles to animals

where the cell fate assignment is reversed. The lines correspond to $t_{\text{fusion}}/T_{\text{L2}} = \Delta T + t_0/T_{\text{L2}}$, with

448    $\Delta T = 0.2$ (orange) and $\Delta T = 0.1$ (green). The *bar-1(++);cwn-1(0)* mutant has a shorter time delay Δt

between BAR-1 pulse onset and fusion. (B) Comparing BAR-1 levels at the time of cell fusion between

450   *bar-1(++);lin-39(lf)* (orange, n=17) and *bar-1(++);cwn-1(0)* animals (green, n=6). Data for fusing P3.p and

P4.p cells was pooled. Even though the difference in the mean of the two distributions was not significant

452   due to the limited number of fusion events that could be recorded (Independent t-test, P=0.2), the highest

BAR-1::GFP levels at the time of cell fusion were observed in the *bar-1(++);lin-39(lf)* mutant that lacks the

454   parallel fusion inhibitor LIN-39. This was consistent with LIN-39 and BAR-1 inhibiting cell fusion in an

additive manner through their absolute levels. (C) Number of *eff-1* mRNA molecules, as measured using

456   smFISH, and BAR-1::GFP fluorescence level in unfused P(3-6).p cells of *lin-39(lf);bar-1(++)* animals.

Each marker corresponds to a single cell (n=28 total animals). High *eff-1* expression (>20 mRNA

458   molecules) was only observed for sufficiently low BAR-1::GFP levels.

460   Because cells that fused did so rapidly after the onset of the BAR-1 pulse, it proved

challenging to accurately determine the pulse onset time for these cells from our fitting

462   procedure (see Methods). To circumvent this issue, we selected animals where one Pn.p cell,

either P3.p or P4.p, assumed hyp7/fusion fate whereas the other assumed VPC fate, in *bar-*

464   *1(++);lin-39(lf)* and *bar-1(++);cwn-1(0)* animals where both P3.p and P4.p frequently assumed

hyp7/fusion fate. We then compared within the same animal the fusion time t_{fusion} in the

466   hyp7/fusion cell with the pulse onset time t_0 in the VPC (Fig. 6a). For *bar-1(++);lin-39(lf)*

animals, the time of fusion correlated strongly with the pulse onset time (R=0.86 for P3.p

468   assuming hyp7/fusion and P4.p VPC fate). Specifically, the data clustered along the line

$t_{\text{fusion}}/T_{\text{L2}} = 0.2 + t_0/T_{\text{L2}}$, i.e. cell fusion occurred at a time $0.2T_{\text{L2}}$, or on average ~2 hrs, after the

470   onset of the BAR-1 pulse. However, in *bar-1(++);cwn-1(0)* animals, where the hyp7/fusion

frequency is increased, the delay was halved, with cell fusion now occurring at a time $0.1T_{\text{L2}}$

472   after onset of the BAR-1 pulse (Fig. 6a). Because *bar-1(++);cwn-1(0)* animals generally have

similar BAR-1 pulse slopes as the *bar-1(++)* and *bar-1(++);lin-39(lf)* mutants (Fig. 5c), the main

474   impact of the observed shorter delay between pulse onset and time of fusion would be to lower

inhibitory BAR-1 levels at the time cell fusion is activated. In agreement with this, when we

476   compared BAR-1 levels, as measured by BAR-1::GFP fluorescence, between *bar-1(++);cwn-*

1(0) and *bar-1(++);lin-39(lf)* animals, we found lower BAR-1 levels in fusing P3.p and P4.p cells

478   in the *cwn-1(0)* background at the time of cell fusion (Fig. 6b). Indeed, this observed difference

is predicted when absolute BAR-1 and LIN-39 levels bias cell fate outcome in an additive

480   manner: because the fusion inhibitor LIN-39 is not functional in *bar-1(++);lin-39(lf)* mutants,

higher BAR-1 levels would be required to inhibit fusion.

482   Finally, we examined the relation between absolute BAR-1 levels and *eff-1* expression.

In general, we expected *eff-1* expression to depend both on BAR-1 and LIN-39 level. To avoid

484   confounding the relation between BAR-1 level and *eff-1* expression by variability in LIN-39 level,

we quantified *eff-1* mRNA level in *bar-1(++);lin-39(lf)* animals. In general, we found that unfused

486   P(3-6).p cells with high BAR-1::GFP level had few *eff-1* transcripts, while high *eff-1* levels were

found only in unfused P(3-6).p cells with low BAR-1::GFP levels (Fig. 6c). Overall, these results

488   are consistent with BAR-1 biasing the hyp7/fusion versus VPC fate decision through its absolute

level.

Discussion

Variability in LIN-39/Hox and BAR-1/β-catenin dynamics

492  Here, we combined a novel time-lapse microscopy approach with quantitative analysis to study how the stochastic outcome of the hyp7/fusion versus VPC fate decision in the P3.p cell is

494 controlled by variability in the dynamics of the underlying signaling network (Fig. 7). We found evidence that linked variability in the dynamics of the fusion inhibitors LIN-39/Hox and BAR-1/β-

496 catenin to cell fate outcome. First, we observed that LIN-39::GFP levels exhibited significant variability, both in time and between animals (Fig. 3c,d). In addition, in *lin-39::gfp;cwn-1(0)*

498 animals, where both P3.p and P4.p assume hyp7/fusion fate in a stochastic manner, we found that cell fusion occurred in the cell with the lowest LIN-39::GFP level (Fig. 3f). Second, we found

500 that BAR-1/β-catenin accumulated in P(3-6).p cells in a single, ~1-4hr pulse around the time of the decision between vulva precursor or hyp7/fusion fate (Fig. 4). These BAR-1 pulses showed

502 considerable animal-to-animal variability, both in time of onset and in slope, i.e. the rate of BAR-1 accumulation (Fig. 5a-d). BAR-1 pulses likely impacted cell fate outcome by modulating the

504 absolute BAR-1 level at the time of activation of cell fusion. In particular, two observations suggested that BAR-1 impacts fusion frequency through its absolute level: Pn.p cells in a

506 mutant lacking the parallel fusion inhibitor LIN-39 fused with higher BAR-1::GFP levels at time of cell fusion, compared to a strain where LIN-39 was present (Fig. 6b). Moreover, high expression

508 of *eff-1*, the fusogen protein required for cell fusion, only occurred in Pn.p cells with low BAR-1::GFP levels (Fig. 6c).

512 Figure 6. **Model of cell fate bias by variability in LIN-39 levels and BAR-1 pulse dynamics.**

(A) Overview of interactions. Expression of *eff-1* and cell fusion are induced through an unknown

514 activating signal, while BAR-1/β-catenin and LIN-39/Hox inhibit *eff-1* expression by inducing inhibitors,

such as EGL-18 and ELT-6 (Koh et al. 2002), Developmental timing cues control both activation of *eff-1*

516 expression in the mid-L2 stage and BAR-1 accumulation dynamics, with the BAR-1 accumulation pulse

starting and ending in the early and late L2 stage, respectively. **(B)**, **(C)** Examples linking the

518 experimentally observed variability in LIN-39 levels and BAR-1 pulse dynamics to cell fate outcome. If

combined levels of fusion inhibitors LIN-39 (green line) and BAR-1 (red line) are sufficiently high at the

520 time of *eff-1* activation (blue line), no cell fusion occurs (B). Reduced levels of LIN-39 and BAR-1 allow for

activation of *eff-1* expression, resulting in cell fusion (C). BAR-1 levels can be reduced both by delayed

522 BAR-1 pulse onset time for a constant BAR-1 accumulation rate (C, top panel), or by lowering the BAR-1

pulse slope, corresponding to lower BAR accumulation rate (C, bottom panel).

Temporal regulation

526   Our observations revealed a high degree of temporal regulation of the key events

controlling the decision to fuse or not (Fig. 7a). We found that mutants that strongly impacted

528 hyp7/fusion frequency had little impact on the time of cell fusion (Fig. 1e) and cell fusion

occurred in a synchronized manner in mutants where multiple Pn.p cells assume hyp7/fusion

530   fate (Fig. 1f). Similarly, we found that BAR-1 pulses started and ended at distinct points in the

L2 larval stage, with their timing synchronized between Pn.p cells (Figs. 4, 5e). The onset and

532   end of BAR-1 pulses showed clear differences in terms of variability: whereas BAR-1 pulse

onset varied strongly between animals, BAR-1 pulses ended at a much more specific point in

534   the L2 larval stage, leading to a broad distribution of BAR-1 pulse durations (Fig. 5e,f). Our

findings linked the timing of BAR-1 pulses to the frequency of cell fusion. In particular, the only

536   detectable difference between a *bar-1(++)* mutant, with low fusion frequency, and a *bar-1(++);cwn-1(0)* mutant, with high fusion frequency, was a delay in BAR-1 pulse onset with

538   respect to the time of cell fusion (Figs. 5c,d, 6a). The strong link between the timing of cell

fusion and BAR-1 pulse dynamics with L2 larval stage progression suggests that these events

540   are regulated by a timing program that is independent of LIN-39 and Wnt signaling, either

directly by a body-wide timing program such as the heterochronic pathway (Resnick, McCulloch,

542   and Rougvie 2010) or indirectly, e.g. as function of Pn.p cell cycle progression during the L2

larval stage.

Control of *eff-1* expression

546   LIN-39 and Wnt signaling likely influence the decision to assume hyp7/fusion fate by

controlling expression of the fusogen protein EFF-1 (Fig. 7a). Indeed, we found that *eff-1*

548   expression was generally low (< 10 mRNA molecules per cell), but showed a strong peak (20-

60 mRNA molecules per cell) shortly before fusion (Fig. 2). Moreover, LIN-39 and Wnt signaling

550   mutations that strongly impacted fusion frequency did not lead to an overall increase in *eff-1*

mRNA level, but rather changed the fraction of animals in which we observed an *eff-1*

552   expression peak (Fig. 2d). The change in *eff-1* expression in LIN-39 and Wnt mutants was

consistent with their known function as inhibitors of cell fusion (Gidi Shemer and Podbilewicz

554   2002; D M Eisenmann et al. 1998). However, because LIN-39 and BAR-1 are typically assumed

to act as activators, their inhibitory action is probably mediated by the induction of transcription

556   factors that directly inhibit *eff-1* expression. For LIN-39, this likely occurs through the

transcription factors EGL-18 and ELT-6 (Koh et al. 2002), but whether BAR-1 impacts *eff-1*

558   expression via these same factors remains an open question. Even though we found evidence

linking (variability in) both LIN-39 and BAR-1 levels to cell fate outcome, these results were

560   obtained in strains where LIN-39 and BAR-1 were overexpressed. As a consequence, it remains

an open question how strong their relative contributions to the bias of hyp7/fusion versus vulva

562   precursor cell fate are under wild-type conditions. Finally, no activator of *eff-1* expression is

currently known. However, our observation that timing of cell fusion is not impact by loss of LIN-

564   39 or Wnt signaling (Fig. 1e) implied the presence of one or more unknown activators that

induce expression of *eff-1* at a highly specific point in the L2 larval stage. Overall, our

566   observations suggested that hyp7/fusion fate required sufficiently high *eff-1* expression and that

a mixture of activating and inhibitory signals controlled the fraction of animals in which this

568   threshold was crossed and cell fusion occurred.

570   **Model of stochastic cell fate bias by LIN-39/Hox and BAR-1/β-catenin**

Based on the above observations we currently favor the following model for how variability in

572   LIN-39 and BAR-1 dynamics biases the stochastic decision to assume hyp7/fusion or vulva

precursor fate (Fig. 7b,c): *eff-1* expression is induced by an activating signal at a highly specific

574   time, but simultaneously inhibited by LIN-39 and BAR-1 in a manner proportional to their

absolute levels in the cell. If the combined level of LIN-39 and BAR-1 is sufficiently high, *eff-1*

576   expression remains below threshold required for fusion and the cell assumes vulva precursor

cell fate (Fig. 7b). However, the observed variability in LIN-39 and BAR-1 dynamics can cause

578   their inhibitory levels to fall below what is required to block *eff-1* expression (Fig. 7c). For BAR-1

pulses, this could occur not only by a lower rate of BAR-1 accumulation, i.e. low BAR-1 pulse

580   slope, but also by delaying the onset of the BAR pulse relative to the time of activation of *eff-1*

expression. Hence, in this model it is not the absolute BAR-1 level itself, as is generally

582   assumed, but rather the timing of its accumulation dynamics relative to other developmental

events, that is a key factor determining the stochastic cell fate outcome.

Generation of BAR-1/β-catenin pulses

586   The observation that BAR-1 accumulated in Pn.p cells in precisely timed pulses was

surprising. However, pulsatile β-catenin accumulation dynamics under constant external Wnt

588   levels has been observed in recent years (Murphy et al. 2014; Kafri et al. 2016; Massey et al.

2019). An important question is whether the BAR-1 pulse dynamics we observed here is

590   influenced by the elevated levels of BAR-1 in the integrated transgene in which it was

visualized. While our endogenously tagged BAR-1::GFP line had wild-type rates of hyp7/fusion

592   fate in P3.p cells, we were unable to detect GFP fluorescence in Pn.p cells using our time-lapse

microscope, indicating significantly lower expression levels compared to the BAR-1::GFP

594   transgene. In the canonical Wnt signaling pathway, BAR-1 levels are controlled by their

degradation through the β-catenin destruction complex, with Wnt signaling leading to

596   inactivation of this complex and accumulation of BAR-1 (Sawa and Korswagen 2013). In the

overexpression BAR-1::GFP strain, the absence of BAR-1::GFP prior to the pulse and its rapid

598   disappearance directly afterwards showed that the increased BAR-1 level was still sufficiently

low to not overwhelm the β-catenin destruction complex. Hence, if the BAR-1 accumulation

600   pulse is controlled by the activity of the destruction complex, as is expected, BAR-1 should

exhibit the same pulse dynamics under wild-type conditions.

602   It is an important question how the BAR-1/β-catenin pulse is generated. BAR-1/β-catenin

accumulation could be controlled by changes in the level of the Wnt ligands outside of the cell or

604   rather by changes in the Wnt pathway components inside the cell, such as changes to Wnt

receptor levels or presence/activity of components of the β-catenin destruction complex. We

606   currently favor the latter hypothesis, since neuronal cells close to the Pn.p cells show BAR-

1::GFP expression in the late L1/early L2 stage when Pn.p cells do not, with a significantly

608   decreased BAR-1::GFP signal in those cells in a *cwn-1(0)* background (data not shown). This

suggests that Wnt ligands are already present and able to activate Wnt signaling at this time

610   and position in the body.

As ligand-activated Wnt receptors sequester and thereby inactivate the destruction

612   complex that induces β-catenin degradation, we expected that changing Wnt levels would

predominantly impact the rate of β-catenin accumulation (Sawa and Korswagen 2013; Clevers

614   and Nusse 2012). However, using the *cwn-1(0)* mutant we found that removing a Wnt ligand

instead changed only the timing of the induced BAR-1/β-catenin pulse (Figs. 4d,f and 5c,d),

616   which is difficult to explain based on our current knowledge of the Wnt pathway. Particularly

surprising is that loss of *cwn-1* almost completely removed the variability in BAR-1/β-catenin

618   pulse timing between adjacent Pn.p cells in the same animal (Fig. 5e). In absence of CWN-1,

other Wnt ligands such as EGL-20 repress hypodermal fate, albeit at reduced efficiency

620   (Pénigault and Félix 2011b). Our results indicate that CWN-1 acts in a significantly more

stochastic manner, either on the level of ligand/receptor interaction or the delivery of ligands to

622   the Pn.p cells, than the other Wnt ligands in the body, even though it is the Wnt ligand

expressed closest to the Pn.p cells (David M Eisenmann 2005; Harterink et al. 2011).

Role of β-catenin pulse dynamics in development

626   It is increasingly clear that many of the canonical metazoan signaling pathways control

development using temporal information encoded in their dynamics (Levine, Lin, and Elowitz

628   2013; Shimojo, Ohtsuka, and Kageyama 2008) In particular, pulses in the output of signaling

pathways have now been identified *in vivo* in a number of developmental systems. For example,

630   EGF signaling acts in a pulsatile manner in the VPCs, with signaling strength transmitted in the

frequency of pulses instead of a continuous graded signal (de la Cova et al. 2017). Moreover,

632   time-lapse imaging of oscillatory, rather than pulsatile dynamics of Notch and Wnt signaling

during segmentation of mouse embryos showed that the relative phase of the two oscillations

634   instructs the segmentation process (Sonnen et al. 2018). Although single pulses of β-catenin

accumulation have been observed in cell lines following exogenous application of Wnt ligands

636   (Murphy et al. 2014; Kafri et al. 2016), their function inside in context of development is more

poorly understood. Hence, the BAR-1 pulses we observed here during C. elegans larval

638   development could be an attractive model to study how pulsatile Wnt signaling dynamics

impacts cell fate decisions.

640   It remains an open question what the advantage is of activating BAR-1/β-catenin as a

pulse rather than it being present continuously like the parallel hypodermal fate inhibitor LIN-39.

642   One potential advantage is that BAR-1/β-catenin pulsatile dynamics allows for pulse timing as

an additional control parameter, next to BAR-1/β-catenin accumulation rate, to tune cell fate

644   frequency (Fig. 7b,c). Whether cells receive Wnt input is tightly controlled in space, e.g. by

regulating Wnt receptor expression. Pulsatile dynamics could be a powerful mechanism to

646   control precisely when cells respond to ligands in time. This might be particularly important

because the same signaling pathways are used many times during development, sometimes

648   even in the same cell, to control different developmental events. Reading out these signaling

pathways only at particular time points would allow the reconfiguration of the pathway from

650   executing one developmental decision to another. Interestingly, Wnt signaling is used in VPCs

at the mid-L3 stage, ~10 hrs after the hypodermal versus VPC decision, to control the

652   anteroposterior orientation of their asymmetric divisions (Green, Inoue, and Sternberg 2008).

Here, EGL-20 plays an important role, whereas BAR-1 and CWN-1 have a smaller contribution.

654   The decay of the BAR-1/β-catenin pulse at the end of the L2 stage might be crucial to avoid

temporal crosstalk between the outputs of the Wnt pathway as the VPCs transition from one

656   process to the next.

In conclusion, we have shown here that β-catenin accumulation can be highly dynamic

658   during development, with temporal information instructing development contained in its

dynamics. Many (stochastic) cell fate decisions in organism from nematodes to humans are

660   controlled by Wnt signaling and it will be interesting to see whether pulsatile β-catenin plays a

similar role in biasing cell fate frequencies in those systems. The quantitative approach we

662   employed here, combining in vivo time-lapse imaging of β-catenin dynamics with measurements

of key dynamical parameters such as pulse slope and pulse timing, can provide a template for

664   such future studies.

Materials and Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

668   Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Jeroen van Zon (j.v.zon@amolf.nl).

670   **EXPERIMENTAL DETAILS**

Strains

672   All strains were handled according to the standard protocol on Nematode growth medium (NGM) agar plates with OP50 bacteria (Brenner 1974). Experiments were performed

674   on L2 stage hermaphrodites. Strains were obtained from the CGC unless otherwise indicated. The following mutations were used in this study:

676   LGII: *cwn-1(ok546)* (The C. elegans Deletion Mutant Consortium 2012)*.*

LGIII*: lin-39(gk893)* (The C. elegans Deletion Mutant Consortium 2012)*, lin-39(n709)* (Clark,

678   Chisholm, and Horvitz 1993)*.*

LGX: *bar-1(ga80)* (D M Eisenmann et al. 1998).

680   The following transgenes were used in this study:

ncIs13[ajm-1::GFP] (Liu et al. 2005).

682   *sIs11337[rCesY37A1B.5::GFP + pCeh361)]* (McKay et al. 2003)*.*

*ouIs20 [ajm-1::mCherry + unc-119+] (*gift from Alison Woollard)*.*

684   *itIs37[pie-1p::mCherry::H2B::pie-1 3'UTR + unc-119(+)] IV, stIs10116[his-72(promoter)::his-24::mCherry::let-858 3'UTR + unc-119(+)], stIs10311[lin-39::TGF(3D3)::GFP::TY1::3xFLAG]*

686   (Sarov et al. 2012)*,*

gaIs45[pDE218(bar-1::bar-1::GFP)] (gift from David Eisenmann) (D M Eisenmann et al. 1998),

688   *stIs10226[his-72p::HIS-24::mCherry::let-858 3' UTR + unc-119(+)]* (Sarov et al. 2012).

The presence of the *cwn-1(ok546)* homozygous deletion was confirmed by nested PCR

690   screening. The following primers were used: outer F ('5-TCGTTTCTGACATGGCTCAC-3'),

outer R ('5-ACCCATCCTTTCCCAATCTC-3'), inner F ('5-CGTATCCACGACCACAACAG-3')

692   and inner R (5'-AGAATCTTCACACCAACGGG-3').

694   **Time-lapse imaging**

The microchamber size used in the study was 190 μm x 190 μm, with a depth of 10 μm,

696   made as previously described (Gritti et al. 2016). Using an eyelash attached to a glass pipette,

OP50 bacteria were used as "glue" to transfer eggs into the microchambers using M9 solution to

698   keep the microchamber moist. A Nikon Ti-E inverted microscope with a large chip camera

(Hamamatsu sCMOS Orca v2) and a 60 X magnification objective (NA=1.4 oil immersion) was

700   used for imaging. Transmission imaging was performed using an LED light source (CoolLED

pE-100 615nm), while 488 and 561 nm fluorescence images were acquired using Coherent

702   OBIS LS 488-100 and OBIS LS 561-100 lasers, respectively. Images were acquired in a

temperature controlled room at 19° with a sample temperature of 22°. Exposure time was 10 ms

704   and approximately 25 images were taken with a z-distance of 1 μm. Images were taken every

20 min. Images were 2048 x 2048 pixels and saved in 16-bit TIFF format. Fusion times were

706   determined by *ajm-1::GFP/mCherry* localization and morphology.

708   **Quantitative Image Analysis**

For all experiments, transmitted light images where used to identify molt times. Custom

710   Python scripts (available at Github: https://github.com/jvzonlab/P3p_bar1) and ImageJ were

used to quantitatively analyze the acquired images (Schindelin et al. 2012; Schneider, Rasband,

712   and Eliceiri 2012). First, images to be used for quantitative analysis were corrected for uneven

laser illumination in the following manner: flat field intensity for the particular experiment was

714   obtained by imaging a uniformly fluorescent (488nm) testing slide, and averaging the result of

10 images. We divided each pixel's intensity value in the experimental images by the

716   corresponding flat field pixel's intensity, normalized to the median value of the entire flat field

image. This normalization procedure corrects for position-dependent variation in light intensity

718   due to the Gaussian profile of the laser beam. The region of interest was cropped at this time.

Pn.p cells were manually identified by stereotyped nuclear position location and the domains of

720   *ajm-1::GFP/mCherry* expression, if present. To measure LIN-39::GFP expression, a mask was

manually drawn around the nucleus and the mean fluorescence intensity of the pixels within the

722   mask was calculated. The z-slice closest to the center of the nucleus was used. A background

fluorescence measurement for each image was obtained by creating a mask of the intranuclear

724   space in a region near P3.p and P4.p along the axis of the ventral nerve cord. The mean

background fluorescence value was then subtracted from the mean fluorescence value of the

726   reporter for the same image. To measure BAR-1::GFP expression, a mask was manually drawn

around the Pn.p cytoplasmic region using AJM-1::mCherry signal as a positional guide, with

728   background corrections performed similarly as described above. For the Supplementary Movies,

fluorescence images were computationally straightened and aligned, using the animal's body

730   shape and position of the Pn.p cells as measured from the fluorescent markers.

732   **Single-molecule fluorescence in situ hybridization (smFISH)**

Probe design and smFISH hybridization to visualize *eff-1* mRNA transcripts were

734   performed as previously described (Huelsz-Prince and van Zon 2017; Raj et al. 2008). Custom

probes were designed against the exons of the *eff-1* gene by utilizing the Stellaris® RNA FISH

736   Probe Designer (Biosearch Technologies, Inc., Petaluma, CA). The probes were hybridized with

the dye Cy5 (Huelsz-Prince and van Zon 2017). The sequences of the oligonucleotide probes

738   used in this study are listed in the Supplementary Methods. Animals were collected by washing

plates with M9 and were fixed in 4% formaldehyde in 1 X PBS for 45 min at room temperature.

740   Fixed animals were permeabilized in 70% ethanol at least one night at 4°C. Subsequently,

animals were incubated with the 0.5 µl probes overnight at 30°C in Stellaris® Hybridization

742   Solution containing 10% formamide. The next day, animals were quickly washed two times with

10% formamide and 2 X SSC, followed by an incubation wash for 30 min at 30°C. DAPI was

744   added at 0.01 µg/ml in a final incubation step for 20 min at 30°C. Animals were mounted in Glox

Buffer with catalase and glucose oxidase, and images were acquired with a Nikon Ti-E inverted

746   fluorescence microscope, equipped with a 100X plan-apochromat oil-immersion objective and

an Andor Ikon-M CCD camera controlled by µManager software (Edelstein et al. 2014). Stacks

748   of each animal were taken with a z-distance of 0.33 µm and approximately 30 images were

taken per animal. Cy5 exposure time was 3 s, while DAPI and GFP exposure time were 100 ms

750   and 500 ms, respectively. Animals were then imaged at 40 X to determine their body length,

which was measured using ImageJ by drawing a spline from the tip of the head to the end of the

752   tail. smFISH images were analyzed with a custom Python script using techniques previously

described (Raj et al. 2008). The *ajm-1::GFP* transgene was used to determine the cell fusion

754   status. For Figure 6C, ImageJ was used to quantify GFP fluorescence in the Pn.p cellular region

of P3-P6.p by selecting the slice most central to the nucleus. The cellular region was manually

756   outlined and the mean fluorescence value of the cellular region was quantified using ImageJ. A

background fluorescence measurement near the Pn.p cells for each image was then subtracted

758   to correct for background fluorescence. If the corrected BAR-1::GFP fluorescence resulted in a

negative value, it was manually adjusted to zero.

Parameterization and fitting of BAR-1::GFP dynamics

762   To fit the experimentally measured BAR-1::GFP dynamics, we assume the following

minimal model of BAR-1 production and degradation:

$$\frac{dc_{\text{BAR-1}}}{dt} = k_p - k_d \frac{1}{1 + \alpha S(t)} c_{\text{BAR-1}}, \tag{1}$$

764   where $c_{\text{BAR-1}}$ is the BAR-1 level and k_p is the BAR-1 production rate. In the absence of Wnt

signaling, $S(t)=0$, the degradation complex degrades BAR-1 at a basal rate k_d. However, in the

766   presence of Wnt signaling, $S(t)=1$, the degradation complex is inhibited (David M Eisenmann

2005) and degradation occurs at a reduced rate $\frac{k_d}{1+\alpha}$, with $\alpha > 1$. In the model, we assume the

768   BAR-1 pulse is generated by changes in Wnt signaling level $S(t)$. In particular, we assume Wnt

signaling is activated at a constant level $S(t)=1$ starting at time t_0 and ending at time t_0+T,

770   where T is the pulse duration. For other times, Wnt signaling is not activated, $S(t)=0$. These

assumptions yield the following expression for the BAR-1 dynamics:

$$c_{\text{BAR-1}}(t) = \begin{cases} \dfrac{k_p}{k_d} & t < t_0 \\[2ex] \dfrac{k_p}{k_d}(1+\alpha) - \alpha\dfrac{k_p}{k_d}e^{-k_d\frac{t-t_0}{1+\alpha}} & t_0 \le t < t_0 + T \\[2ex] \dfrac{k_p}{k_d} + \alpha\dfrac{k_p}{k_d}\left(1 - e^{-k_d\frac{T}{1+\alpha}}\right)e^{-k_d(t-t_0-T)} & t_0 + T < t \end{cases} \qquad (2)$$

772   For sufficiently long pulse duration T, the BAR-1 level will reach a steady state $c_{\text{BAR-1}} = \frac{k_p}{k_d}(1 +$

$\alpha)$. However, in the experimentally obtained data we never observed BAR-1::GFP levels

774   reaching a steady state before the end of the pulse. Instead, we found that BAR-1::GFP

accumulation remained approximately linear throughout the full duration of the pulse. Indeed,

776   when the pulse duration is sufficiently short or the Wnt-mediated inhibition of BAR-1 degradation

is sufficiently strong, i.e. $(1 + \alpha) \gg k_d T$, the exponential term in $c_{\text{BAR-1}}$(t) for $t_0 \le t < t_0+T$ reduces

778   to an expression that linearly with time in linear fashion, giving rise to the following expression

used to fit the experimental data:

$$c_{\text{BAR-1}}(t) = \begin{cases} \dfrac{k_p}{k_d} & t < t_0 \\[2ex] \dfrac{k_p}{k_d} + k_p \dfrac{\alpha}{1+\alpha}(t - t_0) & t_0 \leq t < t_0 + T \\[2ex] \dfrac{k_p}{k_d} + k_p \dfrac{\alpha}{1+\alpha}Te^{-k_d(t-t_0-T)} & t_0 + T < t \end{cases} \qquad (3)$$

780   We fitted this expression to the experimental BAR-1::GFP data by least-square fitting, using the

implementation of the Levenberg-Marquardt algorithm, as implemented in the Python function

782   scipy.optimize.curve_fit (Eric Jones, Oliphant, and Peterson 2001) and using k_p, k_d, α, t_0 and T

as fit parameters. We obtained these fit parameters for each Pn.p cell by independent fitting.

784   For fusing cells, we fitted $c_{\text{BAR-1}}(t)$ to the experimental data only for time points until the

experimentally determined time of fusion, t_{fusion}. We found that cell fusion always occurred

786   before the end of the pulse, i.e. $t_0+T>t_{\text{fusion}}$ and, hence, T was not defined for fusing cells. In

general, this fitting procedure provides good fits for most BAR-1::GFP trajectories, but fails to

788   converge to a correct fit for trajectories with very low pulse amplitude or no apparent pulse. In

that case, we assume $c_{\text{BAR-1}}(t)=0$, with t_0 and T not defined. To characterize pulse dynamics

790   for non-fusing cell, we compare pulse onset time t_0, pulse duration T and pulse slope $s=k_p\dfrac{\alpha}{1+\alpha}$.

For fusing cells, pulse duration is not defined and instead we compare the time of fusion.

792   Finally, it proved in practice to determine the time of pulse onset t_0 from the above fitting

procedure in cells that fused. This was because fusion typically occurred only a few time points

794   after the onset of the pulse, when BAR-1::GFP levels had not risen far above the background,

providing too little data for the fitting procedure to converge on a proper fit. Using the

796   observation that both pulse onset time and pulse slope were similar in neighboring Pn.p cells,

we could still estimate the pulse onset tme in fusing cells by using the measured value from a

798   neighboring non-fusing cell.

Generation of *bar-1::GFP* Using CRISPR-Cas9 Genome Editing

800   Wild type N2 *C. elegans* genetic background was used for the generation of the

CRISPR/Cas9 *bar-1::GFP* allele. The endogenous *bar-1* gene was tagged C-terminally using

802   the nested CRISPR technique (Vicencio et al. 2019). The protocol was followed with slight

modifications. For step two of the nested CRISPR technique, instead of a fully double-stranded

804   PCR product serving as the repair template, partially single-stranded dsDNA GFP donors were

used. Briefly, two GFP PCR products were created (see Supplemental Methods for primers),

806   one spanning the entire GFP coding region, while the other spanning the interior sequence. The

PCR products were hybridized together to create a donor cocktail featuring fragments with

808   ssDNA overhangs (56 and 57 bp) (Dokshin et al. 2018). The template for GFP was the plasmid

pJJR82 (gift from Mike Boxem). The two GFP PCR products were annealed by mixing the

810   individual PCR products 1:1, then running the following thermocycler program: 95° 2:00s, 85°

0:10s, 75° 0:10s, 65° 0:10s, 55° 1:00s, 45° 0:30s, 35° 0:10s, 25° 0:10s, and finally 4° hold. All

812   crRNA (2 nmol product) and single-stranded repair templates (4 nmol product) were prepared

commercially (IDT) and were resuspended in 20 or 40 μl, respectively, of nuclease-free duplex

814   buffer to yield a stock concentration of 100 μM. The strand of the single-stranded repair

template was determined by data from (Farboud, Severson, and Meyer 2019). Two crRNAs

816   were used for step one of nested CRISPR, with the double-stranded breaks occurring 15 and 18

bp away from the insertion site. The first injection mix contained 250 ng/μl Cas9 (IDT), 10 ng/μl

818   tracrRNA, 3 ng/μl *bar-1* crRNA #1, 3 ng/μl *bar-1* crRNA #2, 3 ng/μl GFP crRNA, 2 ng/μl *bar-1*

repair ssODN, 400 ng/μl *GFP* PCR product, 1 ng/μl *dpy-10* crRNA, 1 ng/μl *dpy-10(cn64)* repair

820   ssODN, and RNAse free buffer to a final volume of 10 μl. The second injection mix contained

250 ng/μl Cas9 (IDT), 10 ng/μl tracrRNA, 7 ng/μl GFP crRNA, 2 ng/μl *bar-1* repair ssODN, 400

822   ng/μl *GFP* hybrid PCR product, 3 ng/μl *dpy-10* crRNA, 1 ng/μl *dpy-10(cn64)* repair ssODN,

RNAse free buffer to 10 μl. In both steps, young adult N2 hermaphrodites (P0) were injected in

824   the germline using an inverted micro-injection setup (Eppendorf). After injection, *dpy-10* edited

animals (F1) were singled and grown at 20°, and the *bar-1* locus was genotyped for successful

826   co-conversion by PCR amplification with OneTaq polymerase (New England Biolabs) using

primers that annealed to the genomic DNA surrounding the insertion point. Potential GFP

828   insertions were determined visually by amplicon size on agarose gel and were sequenced to

verify in-frame insertions (Macrogen Europe). The *dpy-10* edited co-CRISPR allele was

830   removed from the genetic background from the first nested CRISPR step by picking a wild-type

animal from a heterozygous *dpy-10* edited animal. In the second step, it was removed by

832   outcrossing the edited *dpy-10* homozygous animal to N2 for one generation. Growth,

movement, and fertility were all superficially wild-type in the tagged line.

Data availability statement

836   Strains and plasmids are available upon request. Supplemental data (analyzed time-

lapse and smFISH data for individual animals, fitted pulse parameter for BAR-1) and Python

838   software used for analysis and creating figures are available at http://github.com/jvzonlab.

Acknowledgements

This work is part of the research program of the Netherlands Organisation for Scientific

842   Research (NWO) and was performed at the research institute AMOLF. Some *C. elegans* strains

were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs

844   (P40 OD010440). The work was supported by a European Research Council Starting Grant

(338200-STOCHCELLFATE) awarded to J.S.v.Z.

Author Contributions

848   Conceptualization, J.S.v.Z.; Methodology, J.R.K. and J.S.v.Z.; Formal Analysis: J.R.K., J.T, and

J.S.v.Z.; Investigation, J.R.K., and J.T.; Writing – Original Draft, J.R.K. and J.S.v.Z.; Writing –

850   Review & Editing, J.R.K. and J.S.v.Z.; Funding Acquisition, J.S.v.Z.; Supervision, J.S.v.Z.

852   # Declaration of Interests

The authors have declared that no competing interests exist.

References

Alper, S, and C Kenyon. 2001. "REF-1, a Protein with Two BHLH Domains, Alters the Pattern of Cell Fusion in C. Elegans by Regulating Hox Protein Activity." *Development (Cambridge, England)* 128 (10): 1793–1804. http://www.ncbi.nlm.nih.gov/pubmed/11311160.

Alper, Scott, and Cynthia Kenyon. 2002. "The Zinc Finger Protein REF-2 Functions with the Hox Genes to Inhibit Cell Fusion in the Ventral Epidermis of C. Elegans." *Development (Cambridge, England)* 129 (14): 3335–48. http://dev.biologists.org/content/129/14/3335.short.

Balaban, N. Q. 2004. "Bacterial Persistence as a Phenotypic Switch." *Science* 305 (5690): 1622–25. https://doi.org/10.1126/science.1099390.

Brabin, Charles, Peter J. Appleford, and Alison Woollard. 2011. "The Caenorhabditis Elegans GATA Factor ELT-1 Works through the Cell Proliferation Regulator BRO-1 and the Fusogen EFF-1 to Maintain the Seam Stem-Like Fate." *PLoS Genetics* 7 (8): e1002200. https://doi.org/10.1371/journal.pgen.1002200.

Braendle, Christian, and Marie Anne Félix. 2008. "Plasticity and Errors of a Robust Developmental System in Different Environments." *Developmental Cell* 15 (5): 714–24. https://doi.org/10.1016/j.devcel.2008.09.011.

Brenner, Sydney. 1974. "THE GENETICS OF CAENORHABDITIS ELEGANS." *Genetics* 77 (May): 71–94. https://doi.org/10.1002/cbic.200300625.

Chen, Zhe, and Min Han. 2001. "C. Elegans Rb NuRD and Ras Regulate Lin-39-Mediated Cell Fusion during Vulval Fate Specification." *Current Biology* 11 (23): 1874–79. https://doi.org/10.1016/S0960-9822(01)00596-6.

Clark, Scott G., Andrew D. Chisholm, and H. Robert Horvitz. 1993. "Control of Cell Fates in the Central Body Region of C. Elegans by the Homeobox Gene Lin-39." *Cell* 74 (1): 43–55. https://doi.org/10.1016/0092-8674(93)90293-Y.

Clevers, Hans, and Roel Nusse. 2012. "Wnt/β-Catenin Signaling and Disease." *Cell* 149 (6): 1192–1205. https://doi.org/10.1016/j.cell.2012.05.012.

Coudreuse, Damien Y.M., Roel Giulietta, Marco C. Betist, Olivier Destree, and Hendrik C. Korswagen. 2006. "Wnt Gradient Formation Requires Retromer Function in Wnt-Producing Cells." *Science* 312: 921–24.

Dokshin, Gregoriy A., Krishna S. Ghanta, Katherine M. Piscopo, and Craig C. Mello. 2018. "Robust Genome Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in Caenorhabditis Elegans." *Genetics* 210 (3): 781–87. https://doi.org/10.1534/genetics.118.301532.

Edelstein, Arthur D, Mark A Tsuchida, Nenad Amodaj, Henry Pinkard, Ronald D Vale, and Nico Stuurman. 2014. "Advanced Methods of Microscope Control Using MManager Software." *Journal of Biological Methods* 1 (2): 10. https://doi.org/10.14440/jbm.2014.36.

Eisenmann, D M, J N Maloof, J S Simske, C Kenyon, and S K Kim. 1998. "The Beta-Catenin Homolog BAR-1 and LET-60 Ras Coordinately Regulate the Hox Gene Lin-39 during Caenorhabditis Elegans Vulval Development." *Development (Cambridge, England)* 125: 3667–80.

Eisenmann, David M. 2005. "Wnt Signaling." *WormBook : The Online Review of C. Elegans Biology*, 1–17. https://doi.org/10.1895/wormbook.1.7.1.

Eric Jones, Travis Oliphant, and Pearu Peterson. 2001. "SciPy: Open Source Scientific Tools for Python."

Farboud, Behnom, Aaron F. Severson, and Barbara J. Meyer. 2019. "Strategies for Efficient Genome Editing Using CRISPR-Cas9." *Genetics* 211 (2): 431–57. https://doi.org/10.1534/genetics.118.301775.

906   Félix, Marie Anne. 2012. "Caenorhabditis Elegans Vulval Cell Fate Patterning." *Physical Biology*
          9 (4). https://doi.org/10.1088/1478-3975/9/4/045001.
908   Gleason, Julie E., Elizabeth a. Szyleyko, and David M. Eisenmann. 2006. "Multiple Redundant
          Wnt Signaling Components Function in Two Processes during C. Elegans Vulval
910   Development." *Developmental Biology* 298 (2): 442–57.
          https://doi.org/10.1016/j.ydbio.2006.06.050.
912   Gleason, Julie E, Hendrik C Korswagen, and David M Eisenmann. 2002. "Activation of Wnt
          Signaling Bypasses the Requirement for RTK/Ras Signaling during C. Elegans Vulval
914   Induction." *Genes & Development* 16 (10): 1281–90. https://doi.org/10.1101/gad.981602.
      Goentoro, Lea, and Marc W. Kirschner. 2009. "Evidence That Fold-Change, and Not Absolute
916   Level, of β-Catenin Dictates Wnt Signaling." *Molecular Cell* 36 (5): 872–84.
          https://doi.org/10.1016/j.molcel.2009.11.017.
918   Goolam, Mubeen, Antonio Scialdone, Sarah J L Graham, Thierry Voet, John C Marioni,
          Magdalena Zernicka-goetz, Mubeen Goolam, et al. 2016. "Heterogeneity in Oct4 and Sox2
920   Targets Biases Cell Fate in 4-Cell Mouse Embryos Article Heterogeneity in Oct4 and Sox2
          Targets Biases Cell Fate in 4-Cell Mouse Embryos." *Cell* 165 (1): 61–74.
922   https://doi.org/10.1016/j.cell.2016.01.047.
      Green, Jennifer L., Takao Inoue, and Paul W. Sternberg. 2008. "Opposing Wnt Pathways Orient
924   Cell Polarity during Organogenesis." *Cell* 134 (4): 646–56.
          https://doi.org/10.1016/j.cell.2008.06.026.
926   Gritti, Nicola, Simone Kienle, Olga Filina, and Jeroen Sebastiaan van Zon. 2016. "Long-Term
          Time-Lapse Microscopy of C. Elegans Post-Embryonic Development." *Nature
928   Communications* 7: 12500. https://doi.org/10.1038/ncomms12500.
      Gupta, Bhagwati P., Wendy Hanna-Rose, and Paul W. Sternberg. 2012. "Morphogenesis of the
930   Vulva and the Vulval-Uterine Connection*." *WormBook : The Online Review of C. Elegans
          Biology*, 1–17.
932   Harterink, Martin, Dong Hyun Kim, Teije C Middelkoop, Thang Dinh Doan, Alexander van
          Oudenaarden, and Hendrik C Korswagen. 2011. "Neuroblast Migration along the
934   Anteroposterior Axis of C. Elegans Is Controlled by Opposing Gradients of Wnts and a
          Secreted Frizzled-Related Protein." *Development (Cambridge, England)* 138 (14): 2915–
936   24. https://doi.org/10.1242/dev.064733.
      Hill, R J, and P W Sternberg. 1993. "Cell Fate Patterning during C. Elegans Vulval
938   Development." *Development* 18: 9–18. http://www.ncbi.nlm.nih.gov/pubmed/8049492.
      Hirabayashi, Y. 2004. "The Wnt/β-Catenin Pathway Directs Neuronal Differentiation of Cortical
940   Neural Precursor Cells." *Development* 131 (12): 2791–2801.
          https://doi.org/10.1242/dev.01165.
942   Hudson, Clare, Narudo Kawai, Takefumi Negishi, and Hitoyoshi Yasuo. 2013. "B-Catenin-
          Driven Binary Fate Specification Segregates Germ Layers in Ascidian Embryos." *Current
944   Biology* 23 (6): 491–95. https://doi.org/10.1016/j.cub.2013.02.005.
      Huelsz-Prince, Guizela, and Jeroen Sebastiaan van Zon. 2017. "Canalization of C. Elegans
946   Vulva Induction against Anatomical Variability." *Cell Systems* 4 (2): 219-230.e6.
          https://doi.org/10.1016/j.cels.2017.01.009.
948   Johnston, Robert J., and Claude Desplan. 2010. "Stochastic Mechanisms of Cell Fate
          Specification That Yield Random or Robust Outcomes." *Annual Review of Cell and
950   Developmental Biology* 26 (1): 689–719. https://doi.org/10.1146/annurev-cellbio-100109-
          104113.
952   Kafri, Pinhas, Sarah E. Hasenson, Itamar Kanter, Jonathan Sheinberger, Noa Kinor, Sharon
          Yunger, and Yaron Shav-Tal. 2016. "Quantifying β-Catenin Subcellular Dynamics and
954   Cyclin D1 MRNA Transcription during Wnt Signaling in Single Living Cells." *ELife* 5
          (NOVEMBER2016). https://doi.org/10.7554/eLife.16748.
Kim, S., T. Ishidate, R. Sharma, M. C. Soto, D. Conte, C. C. Mello, and M. Shirayama. 2013.

"Wnt and CDK-1 Regulate Cortical Release of WRM-1/ -Catenin to Control Cell Division Orientation in Early Caenorhabditis Elegans Embryos." *Proceedings of the National Academy of Sciences* 110 (10): E918–27. https://doi.org/10.1073/pnas.1300769110.

Koh, Kyunghee, Sara M Peyrot, Cricket G Wood, Javier A Wagmaister, Morris F Maduro, David M Eisenmann, and Joel H Rothman. 2002. "Cell Fates and Fusion in the C. Elegans Vulval

Primordium Are Regulated by the EGL-18 and ELT-6 GATA Factors -- Apparent Direct Targets of the LIN-39 Hox Protein." *Development* 129: 5171–80.

Korswagen, H C, M a Herman, and H C Clevers. 2000. "Distinct Beta-Catenins Mediate Adhesion and Signalling Functions in C. Elegans." *Nature* 406 (6795): 527–32.

https://doi.org/10.1038/35020099.

Korswagen, Hendrik C. 2002. "Canonical and Non-Canonical Wnt Signaling Pathways in

Caenorhabditis Elegans: Variations on a Common Signaling Theme." *BioEssays* 24 (9): 801–10. https://doi.org/10.1002/bies.10145.

la Cova, Claire de, Robert Townley, Sergi Regot, and Iva Greenwald. 2017. "A Real-Time Biosensor for ERK Activity Reveals Signaling Dynamics during C. Elegans Cell Fate

Specification." *Developmental Cell* 42 (5): 542-553.e4. https://doi.org/10.1016/j.devcel.2017.07.014.

Levine, Joe H, Yihan Lin, and Michael B. Elowitz. 2013. "Functional Roles of Pulsing." *Science* 342 (December): 1193–1200. https://doi.org/10.1126/science.1239999.

Lindström, Nils O., Melanie L. Lawrence, Sally F. Burn, Jeanette A. Johansson, Elvira R.M. Bakker, Rachel A. Ridgway, C. Hong Chang, et al. 2014. "Integrated β-Catenin, BMP,

PTEN, and Notch Signalling Patterns the Nephron." *ELife* 3: e04000. https://doi.org/10.7554/eLife.04000.

Liu, Zhicen, Takashi Fujii, Akira Nukazuka, Rie Kurokawa, Motoshi Suzuki, Hajime Fujisawa, and Shin Takagi. 2005. "C. Elegans PlexinA PLX-1 Mediates a Cell Contact-Dependent

Stop Signal in Vulval Precursor Cells." *Developmental Biology* 282 (1): 138–51. https://doi.org/10.1016/j.ydbio.2005.03.002.

Losick, Richard, and Claude Desplan. 2008. "Stochasticity and Cell Fate." *Science* 320 (5872): 65–68. https://doi.org/10.1126/science.1147888.Stochasticity.

Maamar, Hedia, Arjun Raj, and David Dubnau. 2007. "Noise in Gene Expression Determines Cell Fate in Bacillus Subtilis." *Science* 317 (July): 526–29.

Maloof, J N, and C Kenyon. 1998. "The Hox Gene Lin-39 Is Required during C. Elegans Vulval Induction to Select the Outcome of Ras Signaling." *Development (Cambridge, England)*

125 (2): 181–90. http://www.ncbi.nlm.nih.gov/pubmed/9486792.

Massey, Joseph, Yida Liu, Omar Alvarenga, Teresa Saez, Matthew Schmerer, and Aryeh

Warmflash. 2019. "Synergy with TGFβ Ligands Switches WNT Pathway Dynamics from Transient to Sustained during Human Pluripotent Cell Differentiation." *Proceedings of the*

National Academy of Sciences 116 (11): 4989–98. https://doi.org/10.1073/pnas.1815363116.

McKay, S. J., R. Johnsen, J. Khattra, J. Asano, D. L. Baillie, S. Chan, N. Dube, et al. 2003. "Gene Expression Profiling of Cells, Tissues, and Developmental Stages of the Nematode

C. Elegans." *Cold Spring Harbor Symposia on Quantitative Biology* 68 (January): 159–69. https://doi.org/10.1101/sqb.2003.68.159.

Mila, Danielle, Adriana Calderon, Austin T. Baldwin, Kelsey M. Moore, McLane Watson, Bryan T. Phillips, and Aaron P. Putzke. 2015. "Asymmetric Wnt Pathway Signaling Facilitates

Stem Cell-like Divisions via the Nonreceptor Tyrosine Kinase FRK-1 in Caenorhabditis Elegans." *Genetics* 201 (3): 1047–60. https://doi.org/10.1534/genetics.115.181412.

Mohler, William a, Gidi Shemer, Jacob J del Campo, Clari Valansi, Eugene Opoku-Serebuoh, Victoria Scranton, Nirit Assaf, John G White, and Benjamin Podbilewicz. 2002. "The Type I

Membrane Protein EFF-1 Is Essential for Developmental Cell Fusion." *Developmental Cell* 2 (3): 355–62. https://doi.org/10.1016/S1534-5807(02)00129-6.

1008 Mucenski, Michael L., Susan E. Wert, Jennifer M. Nation, David E. Loudy, Joerg Huelsken, Walter Birchmeier, Edward E. Morrisey, and Jeffrey A. Whitsett. 2003. "β-Catenin Is
1010 Required for Specification of Proximal/Distal Cell Fate During Lung Morphogenesis." *Journal of Biological Chemistry* 278 (41): 40231–38.
1012 https://doi.org/10.1074/jbc.M305892200.

Murphy, Malea M., Alexandra C. Keefe, Jennifer A. Lawson, Steven D. Flygare, Mark Yandell,
1014 and Gabrielle Kardon. 2014. "Transiently Active Wnt/β-Catenin Signaling Is Not Required but Must Be Silenced for Stem Cell Function during Muscle Regeneration." *Stem Cell*
1016 *Reports* 3 (3): 475–88. https://doi.org/10.1016/j.stemcr.2014.06.019.

Myers, Toshia R, and Iva Greenwald. 2007. "Wnt Signal from Multiple Tissues and Lin-3/EGF
1018 Signal from the Gonad Maintain Vulval Precursor Cell Competence in Caenorhabditis Elegans." *Proceedings of the National Academy of Sciences of the United States of*
1020 *America* 104 (51): 20368–73. https://doi.org/10.1073/pnas.0709989104.

Ohyama, T. 2006. "Wnt Signals Mediate a Fate Decision between Otic Placode and Epidermis."
1022 *Development* 133 (5): 865–75. https://doi.org/10.1242/dev.02271.

Pani, Ariel M., and Robert Goldstein. 2018. "Direct Visualization of a Native Wnt in Vivo Reveals
1024 That a Long-Range Wnt Gradient Forms by Extracellular Dispersal." *ELife* 7 (August). https://doi.org/10.7554/eLife.38325.

1026 Park, Frederick D, and James R Priess. 2003. "Establishment of POP-1 Asymmetry in Early C. Elegans Embryos." *Development (Cambridge, England)* 130 (15): 3547–56.
1028 https://doi.org/10.1242/dev.00563.

Pénigault, Jean Baptiste, and Marie Anne Félix. 2011a. "Evolution of a System Sensitive to
1030 Stochastic Noise: P3.p Cell Fate in Caenorhabditis." *Developmental Biology* 357 (2): 419–27. https://doi.org/10.1016/j.ydbio.2011.05.675.

1032 Pénigault, Jean Baptiste, and Marie Anne Félix. 2011b. "High Sensitivity of C. Elegans Vulval Precursor Cells to the Dose of Posterior Wnts." *Developmental Biology* 357 (2): 428–38.
1034 https://doi.org/10.1016/j.ydbio.2011.06.006.

Raj, Arjun, Patrick van den Bogaard, Scott A. Rifkin, Alexander van Oudenaarden, and Sanjay
1036 Tyagi. 2008. "Imaging Individual MRNA Molecules Using Multiple Singly Labeled Probes." *Nature Methods* 5 (10): 877–79. https://doi.org/10.1038/nmeth.1253.

1038 Resnick, Tamar D., Katherine A. McCulloch, and Ann E. Rougvie. 2010. "MiRNAs Give Worms the Time of Their Lives: Small RNAs and Temporal Control in Caenorhabditis Elegans."
1040 *Developmental Dynamics* 22 (2): NA-NA. https://doi.org/10.1002/dvdy.22260.

Robertson, Scott M., Jessica Medina, Marieke Oldenbroek, and Rueyling Lin. 2017. "Reciprocal
1042 Signaling by Wnt and Notch Specifies a Muscle Precursor in the *C. Elegans* Embryo." *Development* 144 (3): 419–29. https://doi.org/10.1242/dev.145391.

1044 Roiz, Daniel, Juan Miguel Escobar-Restrepo, Philipp Leu, and Alex Hajnal. 2016. "The C. Elegans Hox Gene Lin-39 Controls Cell Cycle Progression during Vulval Development."
1046 *Developmental Biology* 418 (1): 124–34. https://doi.org/10.1016/j.ydbio.2016.07.018.

Roorda, Austin, and David R. Williams. 1999. "The Arrangement of the Three Cone Classes in
1048 the Living Human Eye." *Nature* 397 (6719): 520–22. https://doi.org/10.1038/17383.

Sarov, Mihail, John I. Murray, Kristin Schanze, Andrei Pozniakovski, Wei Niu, Karolin
1050 Angermann, Susanne Hasse, et al. 2012. "A Genome-Scale Resource for In Vivo Tag-Based Protein Function Exploration in C. Elegans." *Cell* 150 (4): 855–66.
1052 https://doi.org/10.1016/j.cell.2012.08.001.

Sawa, Hitoshi, and Hendrik C Korswagen. 2013. "Wnt Signaling in C. Elegans." *WormBook*, 1–
1054 30. https://doi.org/10.1895/wormbook.1.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair,
1056 Tobias Pietzsch, Stephan Preibisch, et al. 2012. "Fiji: An Open-Source Platform for Biological-Image Analysis." *Nature Methods* 9 (7): 676–82.
1058 https://doi.org/10.1038/nmeth.2019.

1060 Schneider, Caroline A, Wayne S Rasband, and Kevin W Eliceiri. 2012. "NIH Image to ImageJ: 25 Years of Image Analysis." *Nature Methods* 9 (7): 671–75.

1062 Shemer, G, M Suissa, I Kolotuev, K C Nguyen, D H Hall, and B Podbilewicz. 2004. "EFF-1 Is Sufficient to Initiate and Execute Tissue-Specific Cell Fusion in C. Elegans." *Curr Biol* 14 (17): 1587–91. https://doi.org/10.1016/j.cub.2004.07.059.

1064 Shemer, Gidi, and Benjamin Podbilewicz. 2002. "LIN-39 / Hox Triggers Cell Division and Represses Vulval Cell Fusion." *Genes & Development* 16 (24): 3136–41.
1066 https://doi.org/10.1101/gad251202.Wang.

Shimojo, Hiromi, Toshiyuki Ohtsuka, and Ryoichiro Kageyama. 2008. "Oscillations in Notch
1068 Signaling Regulate Maintenance of Neural Progenitors." *Neuron* 58 (1): 52–64. https://doi.org/10.1016/j.neuron.2008.02.014.

1070 Smallwood, P. M., Y. Wang, and J. Nathans. 2002. "Role of a Locus Control Region in the Mutually Exclusive Expression of Human Red and Green Cone Pigment Genes."
1072 *Proceedings of the National Academy of Sciences* 99 (2): 1008–11. https://doi.org/10.1073/pnas.022629799.

1074 Smurova, Ksenia, and Benjamin Podbilewicz. 2016. "RAB-5- and DYNAMIN-1-Mediated Endocytosis of EFF-Fusogen Controls Cell-Cell Fusion." *Cell Reports* 14 (6): 1517–27.
1076 https://doi.org/10.1016/j.celrep.2016.01.027.

Sonnen, Katharina F., Volker M. Lauschke, Julia Uraji, Henning J. Falk, Yvonne Petersen, Maja
1078 C. Funk, Mathias Beaupeux, Paul François, Christoph A. Merten, and Alexander Aulehla. 2018. "Modulation of Phase Shift between Wnt and Notch Signaling Oscillations Controls
1080 Mesoderm Segmentation." *Cell* 172 (5): 1079-1090.e12. https://doi.org/10.1016/j.cell.2018.01.026.

1082 Sternberg, Paul W., and H.Robert Horvitz. 1986. "Pattern Formation during Vulval Development in C. Elegans." *Cell* 44 (5): 761–72. https://doi.org/10.1016/0092-8674(86)90842-1.

1084 Süel, Gürol M., Jordi Garcia-Ojalvo, Louisa M. Liberman, and Michael B. Elowitz. 2006. "An Excitable Gene Regulatory Circuit Induces Transient Cellular Differentiation." *Nature* 440
1086 (7083): 545–50. https://doi.org/10.1038/nature04588.

Sulston, J. E., E. Schierenberg, J. G. White, and J. N. Thomson. 1983. "The Embryonic Cell
1088 Lineage of the Nematode Caenorhabditis Elegans." *Developmental Biology* 100 (1): 64–119. https://doi.org/10.1016/0012-1606(83)90201-4.

1090 The C. elegans Deletion Mutant Consortium. 2012. "Large-Scale Screening for Targeted Knockouts in the Caenorhabditis Elegans Genome." *G3* 2 (11): 1415–25.
1092 https://doi.org/10.1534/g3.112.003830.

Vicencio, Jeremy, Carmen Martínez-Fernández, Xènia Serrat, and Julián Cerón. 2019. "
1094 Efficient Generation of Endogenous Fluorescent Reporters by Nested CRISPR in Caenorhabditis Elegans ." *Genetics* 211 (April): genetics.301965.2019.
1096 https://doi.org/10.1534/genetics.119.301965.

Wang, Bruce B., Michael M. Müller-Immergluck, Judith Austin, Naomi Tamar Robinson, Andrew
1098 Chisholm, and Cynthia Kenyon. 1993. "A Homeotic Gene Cluster Patterns the Anteroposterior Body Axis of C. Elegans." *Cell* 74 (1): 29–42. https://doi.org/10.1016/0092-
1100 8674(93)90292-X.

Yu, Hui, Adeline Seah, Michael A. Herman, Edwin L. Ferguson, H. Robert Horvitz, and Paul W.
1102 Sternberg. 2009. "Wnt and EGF Pathways Act Together to Induce C. Elegans Male Hook Development." *Developmental Biology* 327 (2): 419–32.
1104 https://doi.org/10.1016/j.ydbio.2008.12.023.

Zeev-Ben-Mordehai, Tzviya, Daven Vasishtan, C Alistair Siebert, and Kay Grünewald. 2014.
1106 "The Full-Length Cell-Cell Fusogen EFF-1 Is Monomeric and Upright on the Membrane." *Nature Communications* 5 (May): 3912. https://doi.org/10.1038/ncomms4912.

1108

Table 1

	Fusion/hyp7cell fate rates (%) [a]						
Genotype	P3.p	P4.p	P5.p	P6.p	P7.p	P8.p	N
ncIs13[ajm-1::GFP] [b]	28	0	0	0	0	0	**57**
ouIs20[ajm-1::mCherry] [b]	37	0	0	0	0	0	**30**
cwn-1(ok546);ncIs13[ajm-1::GFP] **cwn-1(0)**	90	83	0	0	0	0	**41**
bar-1(ga80);ncIs13[ajm-1::GFP] **bar-1(0)**	80	76	4	0	0	4	**25**
lin-39(gk893);ncIs13[ajm-1::GFP] [c] **lin-39(0)**	100	100	100	100	100	100	**17**
lin-39(n709);ouIs20[ajm-1::mCherry] **lin-39(lf)**	69	19	0	0	19	50	**16**
lin-39::GFP;HIS24-H2B::mCherry; ncIs13[ajm-1::GFP] **lin-39(++)**	2	0	0	0	0	0	**48**
cwn-1(ok546);lin-39::GFP;HIS24-H2B::mCherry;ncIs13[ajm-1::GFP] **cwn-1(0); lin-39(++)**	20	14	1	0	0	0	**126**
bar-1::GFP;ouIs20[ajm-1::mCherry] **bar-1(++)**	0	0	0	0	0	0	**30**
bar-1::GFP;cwn-1(ok546);ouIs20[ajm-1::mCherry] **cwn-1(0); bar-1(++)**	6	8	0	0	0	0	**64**
bar-1::GFP;lin-39(n709);ouIs20 [ajm-1::mCherry] **lin-39(lf); bar-1(++)**	24	4	3	0	13	19	**70**
bar-1::GFP[CRISPR]; ouIs20[ajm-1::mCherry] **bar-1::GFP[CRISPR]** [b]	30	0	0	0	0	0	**23**

Pn.p fusion frequencies in different genetic backgrounds with time-lapse microscopy techniques.

 [a] Fusion rates are rounded to the nearest percentage. Fusion events were counted by the loss of *ajm-1* staining during the L2 stage, and non-fusion animals were only counted if the animal reached the L3 ecdysis without a fusion event.

 [b] No statistical difference between P3.p fusion rates in these control strains, (Chi-square statistic =0.68, P= 0.71).

[c] P3.p – P8.p fused prematurely during the L1 stage in the null mutant.

Figure S1

A immediately post-fusion

DAPI

eff-1 mRNA

ajm-1::GFP

P2.p   P3.p   P4.p   P5.p

cwn-1(0)

— 6μm

B immediately post-fusion

P3.p
P4.p

eff-1 spots

ns   ns

wild type   *cwn-1(0)*
n=4   n=16

C

P2.p   P3.p   P4.p

eff-1(hy21)

——— 15μm

Figure S1. **The gene *eff-1* is expressed at a high level before cell fusion.**

(A) Example of a *cwn-1(0)* animal with both P3.p and P4.p showing evidence of recent fusion, due to the currently degrading and ruffled appeared of AJM-1::GFP, marked by dashed-white brackets in the composite image (bottom panel). **(B)** Quantification of the number of *eff-1* spots in cells that have recently fused as in (A). Different genetic backgrounds show similar levels of *eff-1* expression at the time of fusion. Dots correspond to individual cells and are plotted with the mean and standard error. No significant differences are present in the average number of *eff-1* spots (Student's t-test). **(C)** Image of a mid-L2 stage *eff-1(hy21)* temperature-sensitive-mutant with a point mutation that renders the protein non-functional, but allows for mRNA staining with smFISH probes. High levels of *eff-1* expression are seen in P2.p and P3.p, despite them remaining unfused (marked by white brackets).

Supplemental Movies

1132 **Movie S1**: (Top panel) Apical junction marker AJM-1::mCherry and (bottom panel) GFP

expressed in the hypodermal syncytium hyp7 for a P3.p cell assuming hyp7/fusion fate. At the

1134 time of fusion with hyp7, 6h 20m after the start of the L2 larval stage, pronounced ruffling of

AJM-1 is followed by its removal from P3.p. Concomitantly, GFP flows from the hyp7 syncytium

1136 into P3.p, as indicated by the yellow arrow. A *dpy-7p::mCherry* nuclear marker was used for cell

identification (top panel).

1138

Movie S2: (Top panel) Apical junction marker AJM-1::mCherry and (bottom panel) GFP

1140 expressed in the hypodermal syncytium hyp7 for a P3.p cell that assumes vulva precursor cell

fate. AJM-1 is present in P3.p throughout the entire L2 larval stage and no inflow of GFP into

1142 P3.p is observed. A *dpy-7p::mCherry* nuclear marker was used for cell identification (top panel).

1144 **Movie S3**: Pulsatile BAR-1 dynamics in P(3-5).p cells in a single animal. Time is relative to the

start of the L2 larval stage. Each panel corresponds to a single cell. Shown are the apical

1146 junction marker AJM-1 (magenta) and BAR-1::GFP (green). The animal examined corresponds

to that shown in Fig. 4a,b.

1148

DNA sequences used for CRISPR/Cas9 genome editing

Name	Sequence 5' to 3'
dpy-10 co-CRISPR crRNA	GCTACCATAGGCACCACGAG
dpy-10(cn64) co-CRISPR single-stranded repair oligo	CACTTGAACTTCAATACGGCAAGATGAGAATGACTGGAAACCGTACCGCATGCGGTGCCTATGGTAGCGGAGCTTCACATGGCTTCAGACCAACAGCCT
bar-1 crRNA #1	CGTCTGATTATTATCCTTCT
bar-1 crRNA #2	TAAAATCGACTATTCCTAGA
GFP crRNA	CGTCGAGCTCGACGGAGTCA
bar-1 single-stranded repair oligo[1]	AAGTGTGGGGGTCTATGTTATAATTAAAAAATTACTTGTAGAGCTCGTCCATTCCGTGGGTGATTCCGGCAGCGGTGACGAACTCCTTGACTCCGTCGAGCTCGACGAGGATTGGGACGACTCCGGTGAAGAGCTCCTCTCCCTTGGA<u>GAA</u>AC<u>G</u>G<u>GGAG</u>T<u>A</u>C<u>GG</u>GAAGGATAATAATCAGACGAATGACTACTCGGACCA
bar-1 primer (F) for genotyping CRISPR insertions	CTCAGCCAGTACGTCACGAGAA
bar-1 primer (R) for genotyping CRISPR insertions	CTGACGCTACTCCACCTTCGAA
GFP primer (F) full sequence	CCAAGGGAGAGGAGCTCTTCA
GFP primer (R) full sequence	CTTGTAGAGCTCGTCCATTC
GFP primer (F) interior sequence	GGACCATGTGGTCACGCTTCTC
GFP primer (R) interior sequence	ACGTCAACGGACACAAGTTCTC

1150

[1]Underscored nucleotides represent silent mutations to avoid recutting after successful repair.

1152

1154 **smFISH probes used to hybridize with *eff-1* mRNA molecules**

Primer #	Probe (5'-> 3')
1	AACTGGGGAGACCACTCAAA
2	GTAACTGCTAGGAGAAGCAG
3	CTCGAGTGGAAATCCGTAGG
4	GGAAGAGCCCATCGAATTTC
5	TGTCTTGGAACAGTGTGGTG
6	GAGATGTTTGAGCACGGACA
7	ATCTGAAGCAGACTGCAGTG
8	TCATTGATCTCTTGGGATGC
9	TCCAAAAGTGTACCGCTGAG
10	AACTGGCATGAACTTCAGGG
11	ATGTGGCATCACACTCACAG
12	AGATTCTGCGGTACATGTTG
13	CTGGACAAGCGGTAAACTGA
14	GTTTCATCAGACTTATCAGA
15	GAACGTGCGGTAGCATGAAG
16	CGATTGGTGTCTGATTTGGG
17	CAAAGCTTGGGGATGTCGTC
18	TGGCTTGAATCGAACGTCAC
19	GACTGCGAGGAATGTCATAT
20	ACGTTGTAGGTTGTTCTAGT
21	GCTGCGTAGACAAATGTTGC
22	TTTATCTTTTTCCACCCAAT
23	TGTGTTCCACCATCTAATTG
24	CGACGTTTTTGGTCGAGATG
25	GGCAGTTACAGCCAATGAAA
26	CAGTTGATGAGATGCTCGTC

Primer #	Probe (5'-> 3')
27	CTCCATTACTTGTTCTTGAG
28	TTGCATTCTCAGTTCTTCTG
29	ATTGTCTGTAATCTCATTCA
30	ACCATCCAAGACGGTCAAAG
31	ATGACCAGAATCGTCCATTC
32	TCACAACTCCATTATTCACA
33	GCCTTGTGAATATCATCCAT
34	GCTCTTTGCAATTTTTCACT
35	ACTTCAAGTGGACGAGTCAA
36	TCGAGCAGATTGAATCCACG
37	GTGTTACAACAGCTTGTCTG
38	TGAAGATTAGTTCCTTCGGC
39	GACAAGGTTTTGACTTTCCA
40	AGTCACGAATTCTTGATGCA
41	TCAACGATGATGGATCCACT
42	AGCCTCATATACTGTCAAGT
43	CTGATCCATCAATTTTTCCA
44	TCCAAATCCAGTTGACATCT
45	GCAGTGAATGTGTGAATTGT
46	CGGTTTGAAGCATGAAGATC
47	ATGGCTGGCAGTGGAATAAT