1

Dopamine and eye movement control in Parkinson's disease: deficits in corollary discharge signals

Henry Railo* [1,2,3], Henri Olkoniemi[2], Enni Eeronheimo[2], Oona Pääkkönen[2], Juho Joutsa[4,5,6], and Valtteri

Kaasinen[7,8]

[1] Department of Clinical Neurophysiology, University of Turku and Turku University Hospital, FI-
20521, Turku, Finland
[2] Department of Psychology, University of Turku, FI-20014, Turku, Finland
[3] Brain and Mind Centre, University of Turku, FI-20014, Turku, Finland
[4] Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital and
Harvard Medical School, Charlestown, MA 02129
[5] Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center
and Harvard Medical School, Boston, MA 02215
[6] Department of Neurology, University of Turku, FI-20521, Turku, Finland
[7] Division of Clinical Neurosciences, University of Turku and Turku University Hospital, FI-20521
Turku, Finland
[8] Turku PET Centre, University of Turku and Turku University Hospital, FI-20521 Turku, Finland
23

* Corresponding author:
Henry Railo,
Department of Psychology,
University of Turku,
Finland
Email: hmrail@utu.fi
Phone: +358-45-6765227

During motor behavior the brain has to predict the sensory consequences of movements to enable

accurate perception of the world. This process is assumed to rely on corollary discharge signals sent

through the basal ganglia and dopaminergic system, both of which malfunction in Parkinson's disease

(PD). We studied early-stage PD patients (N=14), and age-matched healthy control participants (N=14)

to examine whether PD patients reveal deficits in updating their sensory representations after eye

movements. The participants performed a double-saccade task where, in order to accurately fixate a

second target, the participant must correct for the displacement caused by the first saccade. PD

patients showed systematic deficits in fixating the second target, especially when they made saccades

towards the body side with predominant motor symptoms. Across PD patients, the impairment in

fixating the second target correlated with the integrity of the dopaminergic system as measured with

[^{123}I]FP-CIT SPECT: Patients with lower striatal dopamine transporter binding made larger errors. The

direction of the errors suggests that the corollary discharge indicated a larger movement than was

actually made by the patients. Dopamine transporter binding did not correlate with the accuracy of

the first saccade. This deficit in internal monitoring of movements could aggravate motor symptoms,

and explain frequently occurring visual symptoms in PD.

Keywords: Parkinson's disease, corollary discharge, efference copy, predictive coding, dopamine,

dopamine transporter binding, saccadic eye movement

Motor behavior requires the brain to update sensory representations to take into account

changes that are produced by self-induced movements. How the brain accomplishes this remains

open, but theories assume that the basal ganglia, thalamus, and the dopaminergic system contribute

to this process (Crutcher & DeLong, 1984; Wurtz, 2008; Friston, et al., 2012). The present study

examined this little studied hypothesis in Parkinson's disease (PD), which is a neurodegenerative

disease characterized by a deficiency in dopaminergic neurons which leads to malfunction of the basal

ganglia and thalamus (DeLong & Wichmann, 2007; Albin, et al., 1989). We examined PD patients'

ability to monitor their own eye movements using a *double-saccade task* which has been widely

utilized to examine internal monitoring of movements (Wurtz, 2008; Ostendorf, et al., 2010; Thakkar,

et al., 2015). During eye movements the image on the retina changes place, and yet we precisely

know the locations of objects in the world because changes in perception that are due to the eye

movements have been taken into account by the brain. In the double-saccade task a participant must

take into account the displacement caused by a first saccade to accurately perform a second saccade

(Figure 1B). This is assumed to rely on corollary discharge (CD) signals that relay a copy of the motor

command to sensory areas (Figure 1A) via medial thalamus (Sommer & Wurtz, 2008; Wurtz, 2008).

Dopaminergic deficit in PD (DeLong & Wichmann, 2007; Albin, et al., 1989) results in abnormal

functioning of the structures assumed to relay the CD to cortex, most importantly the thalamus. In

PD, the medial dorsal nucleus of the thalamus displays abnormal bursting activity which is assumed to

disturb the information flow to the cortex (Pifl, et al., 2012). The CD signal can be seen as a part of a

more general mechanism that integrates bottom-up sensory information and contextual information.

This is clearly formalized in the predictive coding framework which assumes that the brain is

Figure 1. A) A simplified presentation of CD signals (red arrow). CD signals are predictions from motor commands sent to the sensory areas. B) The double-saccade task. In the baseline condition the participants made two sequential saccades (white arrows) to targets. In the Memory condition, the they made saccades to memorized target locations. C) The grey circles show the possible positions of the targets. For data analysis, the coordinates of the targets/eye positions were transposed to correspond to the case where the first target was at the upper right corner of the screen and the second target directly below it (arrows represent eye movements). D) Hypothesis concerning the double saccade task. If CD is diminished relative to the strength of the motor command, the participants fail to fully take into account the first eye movement they have performed – hence, their second saccades are biased outward (orange dashed line). Amplified CD (relative to the strength of the motor command) produces the opposite pattern of results: the second saccade is biased inward (red dashed line). E) The efficacy of sensorimotor control can also be examined on a trial-by-trial basis by measuring how well the participants take into account errors in the first saccade (Joiner, et al., 2010). In the example, the participant's first saccade falls short of the target T1. The grey dashed line shows what would be the optimal amplitude (length of arrow) and angle (β) of the second saccade to compensate for the error in the first saccade. The second black arrow shows an example where the angle (α) and amplitude of the second saccade is sub-optimal (i.e. participant overestimates the length of the first saccade, for example, due to amplified CD relative to the actual motor output).

constantly attempting to predict (model) incoming sensory information based on previous

experiences (Hohwy, 2013; Clark, 2016). The CD signal is a prediction about the sensory consequence

of a movement. According to this framework, during (eye) movements the brain compares its current

model of the world (predictions) against incoming sensory information (Friston, et al., 2012). Friston

et al. have hypothesized, and using dopamine depletion simulations provided evidence, that the

dopaminergic system mediates sensorimotor integration by adjusting the how much the brain relies

on the predictions (Friston, et al., 2012). Specifically, they have proposed that the dopaminergic

deficiency in PD causes the patients to rely more on their internal predictions (e.g. CD) than bottom-

up sensory information (Friston, et al., 2012).

Defects in CD have been observed in neurological patients with thalamic lesions (Bellebaum, et

al., 2005; Ostendorf, et al., 2010). Using the double-saccade task, Thakkar et al. (2015) observed that

in schizophrenia patients the errors in the angle of the second saccade suggested reduced CD relative

to motor signals. As schizophrenia has been associated with hyperactive striatal dopamine

neurotransmission (Howes & Kapur, 2009), this raises the question whether PD patients – who

characteristically have nigrostriatal dopaminergic defects (Kordower, et al., 2013) – display a reverse

effect (i.e. amplified CD relative to motor commands). Previous studies have not tested whether the

integrity of the dopaminergic system correlates with performance in the double-saccade task. While

dopamine is not *per se* assumed to relay the CD information, it is assumed to mediate the integration

of bottom-up sensory information and contextual information such as previously performed

movements (e.g. CD) (Friston, et al., 2012). The two possible outcomes of CD deficits (reduced vs.

amplified CD relative to the motor signals) are displayed in Figure 1D. As shown in Figure 1E, CD also

allows for taking into account variation in the endpoint of the first saccade on a trial-to-trial basis to

produce accurate saccades to the second target (Joiner, et al., 2010; Munuera, et al., 2009).

Deficits in sensorimotor integration could lead to problems in visual stability and contribute to

the motor symptoms of PD, such as posture and gait, as the patient's internal model of the world

does not accurately reflect world (Friston, et al., 2012). Abnormal CD could also contribute to visual

symptoms observed in PD (Weil, et al., 2016), such as hallucinations (Diederich, et al., 2014; Thakkar,

et al., 2017). For the present study, participants at an early stage of PD without visual symptoms were

recruited to examine if deficits in visuomotor integration can be observed even when the patients do

not themselves report experiencing visual deficits.

Methods

Participants

Fourteen PD patients (8 women; M_{age} = 62, range: 45–75 years, 13 right-handed and one

ambidextrous) and fourteen right-handed neurologically healthy control participants (7 women; M_{age}

= 67, range 44–78) took part in the experiment. PD patients were diagnosed using the UK Brain Bank

criteria, all diagnoses were supported by marked striatal defects in dopamine transporter SPECT and

none of the patients presented features suggestive of atypical parkinsonism. All participants had

normal or corrected-to-normal vision. Three control participants had to be excluded from the analysis

due to technical problems during testing. In most cases the control participants were the spouses of

the PD patients. The duration of motor symptoms in PD patients was on average 24 months (range:

3–72 months), and the Hoehn-Yahr stages of the patients were as follows: Six patients at stage I,

seven patients at stage II, and one patient at stage III. The patients had a mean motor MDS-UPDRS

score of 37 (range 11–95, SD = 20). In eight patients the motor symptoms were primarily on the left

side, in four patients in the right side, and two patients had nearly symmetrical motor symptoms.

None of the patients had depression (mean BDI = 3.8, range 1-12, SD = 2.8), and none showed signs of

abnormal cognitive abilities as measured by the Mini-Mental State Examination (mean 27.6, range 26-

30).

Eleven patients underwent a voluntary break in medication before the experiment, and did not

take medication on the morning of the test session. One patient did not have any medication, and

two participants did not go through a break in medication. Three patients were on levodopa, nine

patients were treated with MAO-B inhibitors and/or dopamine agonists and medication was unknown

for one patient.

The experiment was approved by the Ethics committee of the hospital district of South-West

Finland, and a written informed consent was obtained prior the study. The study was conducted

according to the principle of the Declaration of Helsinki.

Tasks, stimuli and procedure

In addition to performing the double saccade task, we also measured the strength of saccadic

suppression in the participants by measuring visual thresholds during saccades and fixation. Previous

studies have reported increased visual threshold in PD (Silva, et al., 2005; Diederich, et al., 2002), and

saccadic suppression has been hypothesized to be triggered by extra-retinal (e.g. CD) signals (Ross, et

al., 2001). Finally, the participants were queried about their subjective experience of visual stability

using a questionnaire described below. We assumed that deficits in CD and visual updating may lead

to subjectively experienced symptoms of unstable vision.

To assess whether the participants experience symptoms of unstable vision they were asked

whether they perceived motion, experienced vertigo, or had feelings that the world has "changed

position" during saccadic or smooth pursuit eye movements. Smooth pursuit eye movements were

included in the questionnaire as control questions, since we hypothesized that possible impairments

in visual stability selectively affect saccades. In addition participants were asked if they had trouble

estimating the positions of objects, had experienced visual hallucinations, or had experienced any

other visual symptoms. Answers were given on a four step scale (0 = never … 3 = often).

In the double saccade task a participant is asked to shift her gaze to two briefly presented

targets. In the traditional form of the task the targets are presented so briefly that when the

participant initiates the first saccade, both visual targets have disappeared from the screen. Thus the

participant must use her internal monitoring signals (i.e. CD) to take into account the spatial change

produced by the first saccade in order to correctly fixate the second target (Hallett & Lightstone,

1976; Sommer & Wurtz, 2008; Thakkar, et al., 2015). Pilot experiments for the present study showed

that the traditional double saccade task was too challenging for the elderly participants (the

participants had trouble perceiving and making saccades to the locations of the briefly presented

target stimuli). In order to make the task easier for the participants, we modified the task so that each

target stimulus is presented for a longer duration, but the participants were instructed to initiate the

saccades only after both targets had disappeared from the screen (Memory condition; Figure 1B).

While this means that short-term visuospatial memory may contribute to performance more in our

modified version than in the traditional form of the task (which taxes the sensory memory more), the

key idea behind the double saccade task remains unchanged: the participants need to take into

account the spatial change produced by the first saccade in order to correctly fixate the second

target. In the Baseline condition (Fig. 1B), which was physically identical to the experimental

condition, the participants were not required to wait until the targets have disappeared and could

initiate their eye movements voluntarily.

For the double saccade task, round (.6^o) white target stimuli were presented on a black

background. The double saccade task began with a nine point calibration of the eye-tracker (i.e.

whole screen was calibrated). After calibration, each trial began with a white fixation point presented

at the center of the screen. This fixation point was also used as a (eye-tracking) drift-correction point

in the beginning of each trial. After the participants correctly fixated the center of the screen for 1.2-

1.5 sec, the two target stimuli were presented. The first stimulus was always presented in one of the

four quadrants of the screen (e.g. upper right quadrant as in Fig. 1B) for 500 msec ~16 degrees from

the fixation point. Right after the first target, a second target was presented for 130 msec on the

horizontal meridian at the same side of the screen as the first target. However, the exact horizontal

position of the second target varied from trial to trial (there were four equally likely positions,

separated by a minimum of ~2.4 degrees). The possible locations of the saccade targets are presented

in Figure 1C. Each participant performed the baseline and memory conditions once. Thirty trials were

collected for each condition. Before the experimental runs, each participant performed at least 10

practice trials without eye-tracking.

Visual 81% thresholds were measured by presenting a horizontally oriented, low-frequency,

luminance modulated Gabor grating (diameter ~12°, 0.33 cycles per degree, phase varied randomly

on each trial) in the upper or lower visual field (center of the Gabor was 7 degrees from the center of

the screen) for one screen refresh (11.8 msec). The stimuli were presented on mid-grey background.

Participants' task was to report whether the Gabor stimulus was presented in the upper or lower

visual field using the up and down arrow keys. The detection threshold was determined by adaptively

varying the Michelson contrast of the Gabor using a Bayesian staircase method QUEST (Watson &

Pelli, 1983). This procedure is similar to previously reported means of measuring saccadic suppression

(Burr, et al., 1994). Each participant performed three different conditions during which visual

thresholds were measured: during fixation, during saccade, and right after saccade. When the

thresholds were measured during fixations participants kept fixating to a fixation dot, presented in

the middle of the screen. In the 'saccade condition', the visual threshold was determined for Gabor

stimuli presented while participants performed ~19 degree saccades from a fixation point on the left

side of the screen to a black dot at the center of the screen. In the 'after saccade' condition the Gabor

stimulus was presented ~100 msec after the saccade had ended. Participants always initiated the

saccades voluntarily after a brief period of fixation during which the eye tracker was drift corrected.

To enable accurate tracking of horizontal eye-position, the eye-tracker was calibrated by three

calibration points presented on the horizontal meridian (left side, right side and center of the screen).

In both tasks (double saccade and visual threshold) the stimuli were presented using

Psychtoolbox (Brainard, 1997) running on Matlab 2014b. Eye movements were recorded using

EyeLink 1000 eye-tracker (SR Research) that was operated using the EyeLink Toolbox (Cornelissen, et

al., 2002). The eye movement registration was done monocularly, typically for the right eye, using

1000 Hz sampling frequency. The stimuli were presented on a 19˝ CRT screen with a screen resolution

of 1024 $\times$ 768 pixels and 85Hz refresh rate. Participants were seated 70 cm from the screen and a

head rest was used to stabilize the head.

SPECT imaging

To image dopamine transporter (DAT) binding, all PD patients underwent brain [123I]FP-CIT

([123I]N- ω-fluoropropyl-2β-carbomethoxy-3β-(4-iodophenyl) nortropane) SPECT. The interval

between the scanning and behavioural testing (e.g. double saccade task) was 8-29 months. Prior to

scanning, thyroidal update of the ligand was blocked by administering oral potassium perchlorate

(250 mg) 60 minutes before the injection of [123I]FP-CIT. An 185 MBq bolus of [123I]FP-CIT was

administered and scanning performed 3 hours after the injection. Imaging was performed as

described earlier (Kaasinen, et al., 2014). The images were reconstructed and analysed using BRASS

(Hermes Medical Solutions AB, Stockholm, Sweden). Specific binding rations (SBRs) for the caudate

nucleus and the putamen were quantified using occipital cortex as the reference region SBR_{ROI} = (ROI

– OCC) / OCC, where ROI refers to the uptake in the region of interest and OCC in the occipital cortex.

Although occipital cortex may contain minute concentrations of DAT, it has been shown to be a valid

reference region in [123I]FP-CIT imaging (Joutsa, et al., 2015). Mean striatal values (caudate, anterior

putamen and posterior putamen) in the left and right hemisphere were used for correlation analyses.

In statistical analysis of the relationship between DAT binding and behavioural performance, the

analyses always included a factor that indicated whether saccades were made ipsilaterally or

contralaterally with respect to the hemisphere with lower binding.

Data analysis

Single-trial eye movement data (accuracy of fixations) was analyzed using linear mixed-effects

models using the lme4 package (Bates, et al., 2014) in R statistical software (R Development Core

Team , 2014). Because the analysis is performed on a single-trial basis rather than on aggregated

means, mixed-effects models enable taking into account individual differences, in addition to group-

level effects. Participants were included in the model as random variables, that is, each individual

participant's data was fitted with individual intercepts and slopes (concerning the Baseline/Memory

condition). The fixed-effect predictors included in the models are described in the Results section. The

models were pruned by removing fixed-effects predictors that did not significantly ($t < |2|$) contribute

to the model. The visual detection thresholds (saccadic suppression) were analyzed using linear mixed

effects models with separate intercepts for each subject. As there is no standardized way to calculate

the degrees of freedom and the corresponding p values in mixed-effects models (Baayen, et al.,

2008), we do not report p values. However, the statistical significance of an effect can be checked

from the confidence intervals (CI): whenever the value zero is included in the CI, the effect is not

considered statistically significant. Another reason to not report p values is to draw the reader's

attention to the size of the effect (Cumming, 2012). Error bars in all the figures are 95% CIs calculated

from 1000 bootstrap samples.

In the double-saccade task, separate models were built for horizontal and vertical errors

(relative to the target location). Only trials where the first saccade was directed towards the first

target and where the participants made a following saccade towards the second target were included

in the analysis (5.8% of trials were excluded). For each participant, the first fixation to the first and

second targets were included in the analysis, except for cases were the first fixation was very brief

and rapidly followed by a corrective saccade towards the correct location. Before the analysis of

fixation accuracy, the data was transposed to correspond to the case where the first target was at the

upper right corner of the screen and the second target directly below it (Fig. 1C). These preprocessing

steps were performed in Matlab 2014b.

Datasets can be downloaded from the Open Science Framework (https://osf.io/bqnzh).

Results

Visual symptoms

PD patients reported more visual symptoms compared to healthy controls, although this

difference was statistically non-significant (Mann-Whitney test; Table 1). The patient group reported

numerically more symptoms especially concerning general confusion about locations of objects, and

experiencing spatial shifts or jerks during saccades.

Table 1. Visual symptoms (mean and SD)

		Controls	PD	Z	p
General	Nausea	0.07 (0.25)	0.06 (0.24)	-0.05	0.96
	Spatial confusion	0.00 (0.00)	0.13 (0.34)	-1.39	0.16
	Hallucinations	0.00 (0.00)	0.00 (0.00)	0.00	1.00
Saccade	Vertigo	0.14 (0.51)	0.13 (0.33)	-0.45	0.65
	Spatial shift	0.00 (0.00)	0.13 (0.33)	-1.39	0.16
	Motion	0.07 (0.25)	0.13 (0.33)	-0.57	0.36
Smooth pursuit	Vertigo	0.00 (0.00)	0.00 (0.00)	0.00	1.00
	Spatial shift	0.00 (0.00)	0.00 (0.00)	0.00	1.00
	Motion	0.00 (0.00)	0.00 (0.00)	0.00	1.00
Total		0.03 (0.09)	0.07 (0.11)	-1.13	0.26

280
281
282
Saccadic suppression

Visual thresholds did not differ between the control group and the PD group during fixation (t =

-.13), saccade (t = .15), or after saccade (t = -.91). These factors were hence removed from the linear

model used to analyze visual threshold. The pruned model showed that both groups displayed clear

saccadic suppression as visual thresholds increased during saccades (β = 1.96, 95% CI = [1.48, 2.44], t

= 8.06) and right after saccades (β = 0.78, CI = [0.30, 1.25], t = 3.21) when compared to thresholds

during fixation (see Figure 2). This indicates that the patients' visual sensitivity was similar to the

control group.

291
292 Figure 2. Visual thresholds during fixation, saccade, and after saccade for each group. The unit of the y
293 axis is the logarithm of Michelson contrast of the stimulus.
294
295

Double-saccade task

<u>*Horizontal and vertical errors*</u>

Figure 3A shows the participants' accuracy in fixating the first and second targets. The data was

analyzed using linear mixed-effects models that included the predictors group (Control vs. PD),

condition (Baseline vs. Memory), group × condition interaction, and the running trial number

(centered). The models were pruned by removing predictors that did not significantly (t < |2|)

contribute to the model. Horizontal and vertical errors were modelled separately.

Saccades to the first target. Although the fixations to the first target concentrated around the

actual target position (Fig. 3A upper panels), saccades towards the first target were, on average,

somewhat hypometric (i.e. they did not quite reach the target) in the Baseline condition in both

groups (*x error*: β = -23. 49, CI = [-31.19, -15.83], t = -6.11; *y error*: β = -48.19, CI = [-64.33, -32.00], t = -

5.82; in the results, the β values show the size of the average effect in pixels). In the Memory

condition, fixations to the first target became horizontally more accurate in both groups (β = 12.26, CI

= [0.59, 23.76], t = 2.11). However, in PD patients the saccades became vertically more hypometric in

the Memory condition (y error, PD × Memory: β = -42.47, CI = [-78.05, -7.00], t = 2.33), whereas in

control group vertical errors somewhat decreased in the Memory condition (t = 1.61). This means that

when the saccades were performed without visual guidance (Memory condition), the patients' first

saccades were biased towards the second target location.

Figure 3. A) Fixation maps for the first target (T1, upper panels) and second target (T2, lower panels),
separately for control participants (left) and patients (right). The red and blue dots denote individual
fixations in the Baseline and Memory conditions, respectively. The red and blue crosses show the
respective group averages; the lengths of the lines are 95% confidence intervals of the mean. B)
Amount of error as a function of laterality in the PD group. Ipsilateral refers to targets that were
presented at the same side as the motor symptoms. Error bars are the 95% CIs of means. Unit of error
is pixels (22 pixels is ~1⁰).

As motor symptoms in PD are often asymmetric, we analyzed if the accuracy of fixations was

modulated by whether the targets were presented ipsi- or contralaterally with respect to the

predominant motor symptoms. This analysis was restricted to PD patients who showed asymmetric

symptoms (N=12). The results, displayed in Figure 3B, revealed that fixations to the first target were

more inaccurate (horizontally and vertically) when the target was ipsilateral with respect to symptoms

(x error: β = -11.71, CI = [-21.48, -1.93], t = -2.34; y error: β = -19.71, CI = [-19.85, -1.55], t = -2.29).

Moreover, as shown in Fig. 3B (upper panel), in the Memory condition horizontal error was reduced

when the targets were contralateral to motor symptoms (Memory × Laterality: β = -18.91, CI = [-

33.95, -3.87], t = -2.46). Vertical error revealed also an interaction between Memory condition and

target laterality: In the Baseline condition vertical errors increased at ipsilateral locations, whereas

the effect reversed in the Memory condition (β = 16.36, CI = [2.27, 30.40], t = 2.27), but as seen from

Fig. 3B (lower panel) the size of this effect is negligible. In summary, these results show that PD

patients have selective deficits especially in making horizontal saccades to memorized targets that are

at the same side as their primary motor symptoms.

Saccades to the second target. Fixations to the second target (Fig. 3A lower panels) were on

average biased towards the center of the screen and somewhat hypermetric in the Baseline condition

in both groups (x error: β = -15. 48, CI = [-23.87, -7.14], t = -3.68; y error: β = -26.10, CI = [-44.91, -

32.00], t = -7.33). During the Memory condition, fixations were on average biased away from the

center of the screen in the horizontal direction in both groups (β = 18.32, CI = [0.94, 35.70], t = 2.10).

Although Fig. 3A suggests that this effect was smaller in PD patients, this Group × Condition

interaction did not reach statistical significance (t = -1.83) in the linear model. A further analysis

performed on the patient group showed that when the targets were presented at the same side as

motor symptoms, fixations to the second target were on average biased towards the center of the

screen in horizontal direction in both conditions (β = -14.44, CI = [-22.46, -6.38], t = -3.52). This means

that the PD patients show errors whose direction is consistent with amplified CD (relative to motor

command) when they make saccades towards the side with primary motor symptoms. Furthermore,

the bias observed in the fixations to the second target could be in part caused by biases in fixating the

first target. To control for this possible confound, the analyses concerning the bias in the second

saccade were also performed in trial-by-trial basis (see Fig. 1E) in section *Compensation in angle and*

amplitude following the first saccade.

Modulation by DAT binding. Finally, we wanted to see if the accuracy of fixations correlated

with striatal dopamine transporter (DAT) binding in PD patients. This analysis focused on mean striatal

DAT binding. In addition to the condition (Baseline vs. Memory) and trial number factors, the model

included a predictor labelling whether or not the targets were presented contralaterally with respect

to the hemisphere with smaller DAT binding (Contralateral factor), the overall DAT binding in the

more affected hemisphere (i.e. the one with lower binding), the motor MDS-UPDRS score of the

patient, and the interactions of these four predictors.

Horizontal or vertical errors of fixations to the *first target* were not modulated by DAT.

Horizontal errors of fixations to the *second target* become somewhat more accurate as mean

striatal DAT binding increased. This main effect was not statistically significant (β = 24.84, CI = [-3.30,

52.0499], t = 1.7). Interaction between laterality and DAT suggested that it was only observed for

fixations performed to targets presented ipsilaterally to the more affected hemisphere (contralateral

× DAT: $β$ = -23.94, CI = [-44.65, -3.23], t = -2.3). However, although the interaction reached statistical

significance, due to the small size of this correlation this result is not discussed further (DAT binding

vs. horizontal error for ipsilateral targets, $β$ = .9, when the main effect of DAT binding is taken into

account).

Figure 4. Vertical error was modulated by DAT binding, and this effect was stronger when the patients
made fixations to memorized targets (blue = Memory). DAT binding variable has been centered (i.e.
zero represented the mean). The dots represent the observed data, and the lines represent the
modelled results.

Vertical errors in fixations to the *second target* were more clearly modulated by DAT binding, as

shown in Figure 4B. In the Memory condition the vertical accuracy of fixations increased as DAT

binding increased (Memory × DAT: $β$ = 44.00, CI = [13.63, 74.38], t = 2.8). No such effect was observed

in the Baseline condition (t = 0.4). Target laterality did not modulate this effect, and therefore Fig. 4B

shows the average of ipsi- and contralateral data. Again, UPDRS scores did not significantly predict

fixation accuracy (t < |1|), suggesting that the observed modulation by DAT does not simply reflect

the severity of the PD. Altogether these results suggest that dopamine specifically contributed to the

accuracy of the *second* saccades. This effect was most clearly observed in vertical errors, and in the

Memory condition (Fig. 4B) where the saccades are performed without external visual guidance (i.e.

they are based on CD). The direction of the errors is consistent with the interpretation that (perhaps

due to abnormal CD) the participants overestimated the size of the first saccade (see Fig. 1E).

However, as already noted, this effect could also be caused by biases in fixating the first target. Thus,

following analyses examined how the participants compensated for the variability in the endpoint of

the first saccade when making the second saccade.

Compensation in angle and amplitude following the first saccade

As explained in Fig. 1E we also examined how well the participants took into account the trial-

by-trial variability in the first saccade when executing the second saccade. In the regression models,

the predictor variables were the centered optimal angle/amplitude of the saccade (see Fig. 1E), group,

condition, all interaction between these three variables, and running trial number (centered). Models

were pruned by removing non-significant regressors.

As shown in Figure 5A-B, the trial-to-trial variability in the *angle* of the second saccade was

strongly predicted by the optimal angle required to perfectly fixate the second target (β = 0.71, CI =

[0.67, 0.76], t = 29.62). In other words, the participants clearly had information about the error in the

endpoint of the first saccade which they could use to correctly aim to the second target. In the

Memory condition, the angle of the second saccade was smaller than in the Baseline condition. That

is, participants' saccades were biased towards the center of the screen (i.e. the initial fixation point; β

= -4.69, CI = [-7.75, -1.67], t = -3.09). Furthermore, in the Memory condition, the larger the optimal

angle, the larger was the bias towards the center of the screen (interaction: β = -0.13, CI = [-0.20, -

0.02], t = -4.32). PD patients were not statistically significantly different from control participants (t ≤

|1|).

Figure 5. Actual saccade angles (A-B) and amplitudes (C-D) as a function of ideal angle/amplitude. Left
panels show the baseline condition, and the right panel the Memory condition. Red color denotes the
healthy control participants, and the blue color the patient group. Dots show the observed data
(individual saccades) and lines the fitted linear models. Note that angle/amplidute variables are
centered, i.e. negative values show that angle/amplitude was smaller than average. Unit is pixels (22
pixels is ~1⁰).

As with previous analyses, we examined whether the laterality of the motor symptoms

correlated with the angles/amplitudes of the second saccades. As shown in Figure 6, when the targets

were presented at the same side as the motor symptoms the compensation in saccade angle was

worse than when the targets were presented on the side with less motor symptoms – the second

saccades become biased towards the center of the screen, consistent with amplified CD (β = -0.11, CI

= [-0.19, -0.033], t = -2.78). In other words, although the PD group did not in general as a group differ

from healthy controls in compensating for variability in the endpoint of the first saccade, such an

effect was observed when the laterality of the symptoms was considered. In other words, this means

that since in some patients the symptoms are predominantly on the right side, and in other patients

on the left side, the deficit is only observed when the laterality of the symptoms is taken into account.

430
431   Figure 6. The angle of the second saccade as a function of ideal angle. Compensation for variation in
432   the endpoint of the first fixation is worse for targets that are at the predominant side of motor
433   symptoms.
434

Also the *amplitude* of second saccade strongly correlated with the optimal amplitude (β = 0.87,

CI = [0.78, 0.96], t = 19.46). Moreover, although the amplitudes of the second saccades did not differ

between groups (t = 0.04), there was an interaction between group and optimal amplitude: Patients'

second saccades become more hypometric as the optimal amplitude required to reach the target

increased (Fig. 5C-D, blue line: β = -0.15, CI = [-0.27, -0.04], t = 2.61). In other words, the patients had

trouble making long saccades.

Target side relative to motor symptoms did not modulate the precision of compensation for

saccade *amplitudes*.

Modulation by DAT binding. We also examined whether the success of compensating for variability in

the endpoint of the first fixation correlated with dopamine transporter binding in PD patients. As

previously, the model included condition, target laterality (with respect to the more affected

hemisphere), the amount of DAT binding, the optimal angle to perfectly fixate the second target, and

the motor MDS-UPDRS score of the participant (and the interactions between these regressors).

Figure 7. Modulation of compensation in the amplitude of the second saccade. A) Compensation for variation in the endpoint of the first fixation is modulated by DAT binding (blue dashed line) and MDS-UPDRS (red dotted line). The visualized effects represent 2 SD decrease (DAT) or increase (MDS-UPDRS) relative to the patient group average, which is plotted as a grey solid line. The circles represent the observed data. B) Correlation between the amplitude of the second saccade and DAT binding. The red and blue circles represent the (average) observed data in the Baseline and Memory condition respectively. The grey line shows the modelled results (Baseline and Memory conditions pooled, as no Condition × DAT interaction was observed).

As previously stated the actual *angles* of the saccades were strongly predicted by the optimal angle required to reach the second target (β = 0.74, CI = [0.67, 0.81], t = 20.3). As shown in Figure 7A (blue dashed line) patients with lower DAT transporter binding values were more inefficient in taking into account for the trial-to-trial variability in the endpoint of the first saccade and their saccades become more biased towards the center of the screen, consistent with stronger reliance on CD than bottom-up visual information (Ideal angle × DAT: β = 0.17, CI = [0.070, 0.28], t = 3.2). Furthermore, the modulation of the angle of the second saccade by DAT was not explained by the severity of PD as the motor MDS-UPDRS score revealed similar modulatory effect (Ideal angle × UPDRS: β = 0.0030, CI =

[0.00081, 0.0053], t = 2.6). Note that although the *β* coefficient of the Ideal angle × UPDRS interaction

(0.0030) is much smaller than the *β* coefficient of the Ideal angle × DAT interaction (0.17), the relative

size of this effect is in fact very similar (due to the fact that the range of motor MDS-UPDRS scores is

much higher than DAT values). The modulatory effect by motor MDS-UPDRS is visualized in Figure 7A

as the red dotted line which overlaps the blue dashed line that depicts modulation by DAT. In addition

to the above mentioned two-way interactions, a Ideal angle × motor MDS-UPDRS × DAT interaction

was observed (*β* = 0.014, CI = [0.004, 0.025], t = 2.8). This three-way interaction indicates that the

modulation by DAT binding was stronger in participants with more severe PD (i.e. higher UPDRS

scores).

476   Next, we examined whether DAT binding was associated with the ability to modulate the

amplitude of the second saccade to compensate for trial-to-trial variability in the first saccade.

Although patients were in general successful in making a second saccade whose amplitude closely

resembled the ideal amplitude required to perfectly fixate the second target (*β* = 0.73, CI = [0.65,

0.81], t = 17.7), patients with lower DAT binding made more hypometric saccades (*β* = 53.94, CI =

[19.23, 88.65], t = 3.0), as shown in Figure 7B. UPDRS score did not statistically significantly modulate

saccade amplitudes in general (main effect of UPDRS: t = -0.3). However, patients with higher motor

MDS-UPDRS scores were more inefficient in making a close-to-ideal amplitude saccade to the second

target (Ideal amplitude × motor MDS-UPDRS: *β* = -0.010, CI = [-0.015, -0,0050], t = -3.7). Surprisingly,

this effect was only observed when the targets were presented ipsilaterally with respect to the

hemisphere with lower DAT binding (Contralateral × Ideal amplitude × motor MDS-UPDRS: *β* = 0.012,

CI = [0.0048, 0,020], t = 3.1).

Discussion

We examined how well early PD patients can update their spatial representations during eye

movements, and whether this ability correlates with the integrity of the dopaminergic system. The

participants were asked to make eye movements to two targets presented in succession. When

compared to age-matched control participants, the patients' second saccades were biased towards

the center of the screen. Similar effects were observed when examining how well the participants

took into account trial-by-trial variation in the endpoint of the first saccade. In patients, the observed

effects were most prominent when the patients made saccades towards the side with primary motor

symptoms. The spatial error in the second saccade correlated with the integrity of the dopaminergic

system in the PD group.

The present results suggest that the brain systems that internally monitor eye movements are

impaired in PD patients, and that this impairment correlates with dopaminergic deficiency. Crucially,

correlations between dopamine transporter binding and saccade performance were observed only for

fixations to the *second* target, implying that the dopaminergic system contributes specifically to

spatial remapping and not to performing saccades in general. This suggests that the dopaminergic

system may contribute to taking into account previous movements and contextual information when

making sequential saccades. This could be caused by a mismatch between the motor output

commands and the CD signal (previous saccade was shorter than suggested by CD), or a more general

impairment in integrating bottom-up sensory information with top-down contextual information

(Friston, et al., 2012). The present results do not imply that the dopamine system per se carries the

CD, but that dopamine modulates the functioning of the network of brain regions that mediate

visuomotor integration. According to Friston's theory, dopaminergic deficiency leads to reduced

precision in bottom-up sensory information (prediction errors), which causes the brain to rely more

on top-down contextual information (i.e. predictions such as CD signals) than bottom-up visual

information (Friston, et al., 2012). This essentially means that the CD signals start to dominate

sensorimotor integration, and hence motor performance, over motor commands, which is in line with

what we observed.

Previous research has shown that deficits in CD are associated with *reduced* CD relative to

motor signals, that is, converse to what we observed here (Sommer & Wurtz, 2002; Thakkar, et al.,

2015). However, there is no reason to assume that abnormal CD always results in reduction of the CD

signal (although this is obviously the case when an area that conveys the CD signal is lesioned;

(Sommer & Wurtz, 2008)). The present findings are consistent with the finding that schizophrenia

patients reveal the reverse pattern of results when performing the double-saccade task: in contrast to

the present results, schizophrenia patients underestimate the size of the first saccade (Thakkar, et al.,

2015).

The present study employed a modified version of the double-saccade task. In the classical

version (Hallett & Lightstone, 1976; Sommer & Wurtz, 2002; Thakkar, et al., 2015) the visual targets

are presented so briefly that they are removed from the screen before the eye movements begin;

hence the saccades cannot be guided by sensory information. Because the classical double-saccade

task-set up proved too demanding for elderly subjects, we used a modified task where the targets

were presented for a longer duration. We expected that the PD patients would show deficits

especially in the Memory condition, where the saccades to the two target locations are performed

solely based on memory. However, the present results show that PD patients are impaired in their

second saccade also when the first saccade could be directed based on visual information, suggesting

that the impairement is associated with integrating incoming sensory information and top-down

contextual information (Friston, et al., 2012). Because previous studies using the double-saccade task

have focused on the condition where the saccades are directed without visual information, it is

difficult to estimate whether this also applies to previous studies.

Previous studies have emphasized the role of kinesthetic feedback in PD (Jobst, et al., 1997;

Klockgether, et al., 1995), and it could be argued that the errors in fixating the second target are due

to impaired kinesthetic feedback from the extra-ocular muscles, instead of impaired CD. While our

interpretation of the results suggests that the dopaminergic system mediates visuomotor integration,

we cannot completely rule out the explanation that the present results reflect purely deficits motor

behavior as opposed to visuomotor integration. One could argue that the observed correlations

between fixation errors and dopamine partially reflect e.g. combining motor commands to sequences.

This interpretation may in part explain correlations between the *amplitude* of the second saccade and

DAT binding, but it cannot explain why the *angle* of the second saccade is spatially biased towards the

initial fixation. We also wish to note that the present conclusions are restricted by the correlative

nature of our study; more conclusive evidence could be acquired by experimentally manipulating

dopaminergic activity in healthy humans.

PD patients are known to undershoot targets when performing eye movements (Terao, et al.,

2011; Crawford, et al., 1989). In the present study the main difference between the two groups in

making the first saccade was that the PD patients' performance deteriorated in the Memory

condition, which is consistent with previous reports (Le Heron, et al., 2005; Terao, et al., 2011;

Crawford, et al., 1989). Our results also replicate the finding that deficits in saccades are typically

worse on the side with larger primary motor symptoms (Choi, et al., 2011). Importantly, the deficits in

spatial remapping were also dependent on the laterality of the primary motor symptoms: the

hypothesized deficits in the trial-to-trial variability in the angle of the second saccade were only

observed when the laterality of motor symptoms was taken into account (see Fig. 6). This is consistent

with the observation that thalamic lesions (that interfere with the CD) produce lateralized defects

(e.g. (Sommer & Wurtz, 2006)).

The PD patients in the present study were at the early stages of PD, generally did not report

visual symptoms, and the visual sensitivity of the patients was similar to the control group. It is

important to note that with the double-saccade task we were able to detect subclinical deteriorating

visual functions that the patients were unaware of. These disturbances, together with the shown

associations with DAT binding suggest that, as the disease progresses, and the dopaminergic deficit

deepens, relevant clinical CD-associated motor and visual symptoms will emerge. Advanced PD

patients have many visual symptoms (Weil, et al., 2016) and abnormal CD may lead to deficits in

experienced visual stability in PD. More generally, the present study suggests that motor deficits in PD

are partly produced by difficulties in internal monitoring of movements. Studies with patients with

more progressed PD and other dopaminergic diseases are required to confirm these findings.

Acknowledgements

HR was funded by Turku Institute of Advanced Studies and Academy of Finland (grant #308533), and

received a grant from the Finnish Cultural Foundation. HO was funded by the Finnish Cultural

Foundation. JJ was funded by the Academy of Finland (grant # 295580), the Finnish Medical

Foundation and the Orion Research Foundation, and has received travel grants from Orion and

Abbvie, and a research grant from Lundbeck. VK was funded by the Turku University Hospital (ERVA-

funds).

References

Albin, R. L., Young, A. B. & Penney, J. B., 1989. The functional anatomy of basal ganglia disorders.
Trend in Neurosciences, 12(10), pp. 366-375.

Baayen, R. H., Davidson, D. J. & Bates, D. M., 2008. Mixed-effects modeling with crossed random effects for
subjects and items. *Journal of Memory and Language,* Volume 59, pp. 390-412.

Bates, D., Maechler, M., Bolker, B. M. & Walker, S., 2014. lme4: Linear mixed-effects models using Eigen and
S4. *R package version 1.1-7, URL: http://CRAN.R-project.org/package=lme4.*

Bellebaum, C. et al., 2005. The role of the human thalamus in processing corollary discharge. *Brain,* Volume
128, pp. 1139-1154.

Brainard, D. H., 1997. The psychophysics toolbox. *Spatial Vision,* Volume 10, pp. 433-436.

Burr, D. C., Morrone, M. C. & Ross, J., 1994. Selective suppression of the magnicellular visual pathway during
saccadic eye movements. *Nature,* Volume 371, pp. 511-515.

Choi, S.-M.et al., 2011. Directional asymmetries of saccadic hypometria in patients with early Parkinson's
disease and unilateral symptoms. *Europen Neurology,* Volume 66, pp. 170-174.

Clark, A., 2016. *Surfing Uncertainity.* 1 st ed. New York: Oxford University Press.

Cornelissen, F. W., Peters, E. M. & Palmer, J., 2002. The Eyelink toolbox: Eye tracking with MATLAB and the
Psychophysics toolbox. *Behavior Research Methods, Instruments, & Computers,* 34(4), pp. 613-617.

Crawford, T. J., Henderson, L. & Kennard, C., 1989. Abnormalities of nonvisually-guided eye movements in
Parkinson's disease. *Brain,* Volume 112, pp. 1573-1586.

Crutcher, M. D. & DeLong, M. R., 1984. Single cell studies of the primate putamen. II. Relations to direction of
movement and pattern of muscular activity.. *Experimental Brain Research,* 53(2), pp. 244-258.

Cumming, G., 2012. *Understanding The New Statistics: Effect Sizes, Confidence Intervals, and Meta-Analysis.*
New York(NY): Routledge.

DeLong, M. R. & Wichmann, T., 2007. Circuits and circuit disorders of the basal ganglia. *Neurological Review,*
Volume 64, pp. 20-24.

Diederich, N. J., Raman, R., Leurgans, S. & Goetz, C. G., 2002. Progressive worsening of spatial and chromatic
processing deficits in Parkinson disease. *Archievs of Neurology,* 59(8), pp. 1249-1252.

Diederich, N. J., Stebbins, G., Schiltz, C. & Goetz, C. G., 2014. Are patients with Parkinson's disease blind to
blindsight?. *Brain,* Volume 137, pp. 1838-1849.

Friston, K., Adams, R. A., Perrinett, L. & Breakspear, M., 2012. Perceptions as hypotheses: Saccades as
experiments. *Frontiers in Psychology,* 3(151).

Friston, K. J. et al., 2012. Dopamine, affordance, and active inference. *PLoS Computational Biology,* Volume 8,
p. e1002327.

Hallett, P. E. & Lightstone, A. D., 1976. Saccadic eye movements to flashed targets. *Vision Research,* Volume 16,
pp. 107-114.

Hohwy, J., 2013. *The Predictive Mind.* 1 st ed. Oxford: Oxford University Press.

Howes, O. D. & Kapur, S., 2009. The dopamine hypothesis of schizophrenia: Version III - The final common
pathway. *Schizophrenia Bulletin,* 35(3), pp. 549-562.

Jobst, E. E. et al., 1997. Sensory perception in Parkinson's disease. *Archives of Neurology,* Volume 54, pp. 450-
454.

Joiner, W. M., FitzGibbon, E. J. & Wurtz, H. R., 2010. Amplitudes and directions of individual saccades can be
adjusted by corollary discharge. *Journal of Vision,* 10(2), pp. 1-12.

Joutsa, J., Johansson, J. & Kaasinen, V., 2015. Is occipital cortex a valid reference region in 123I-FP-CIT SPECT
imaging?. *Clinical Nuclear Medicine,* 40(7), pp. 615-616.

Kaasinen, V. et al., 2014. Differences in striatal dopamine transporter density between tremor dominant and
non-tremor Parkinson's disease. *European Journal of Nuclear Medicine and Molecular Imaging,* 41(10), pp.
1931-1937.

Klockgether, T. et al., 1995. A defect of kinesthesia in Parkinson's disease. *Movement Disorders,* 10(4), pp. 460-
465.

Kordower, J. H. et al., 2013. Disease duration of the integrity of the nigrostriatal system in Parkinson's disease.
Brain, Volume 136, pp. 2419-2431.

Le Heron, C. J., MacAskill, M. R. & Anderson, T. J., 2005. Memory-guided saccades in Parkinson's disease: long
delays can improve performance. *Experimental Brain Research,* Volume 161, pp. 293-298.

Munuera, J., Morel, P., Duhamel, J.-R. & Deneve, S., 2009. Optimal sensorimotor control in eye movement
sequences. *The Journal of Neuroscience,* 29(10), pp. 3026-3035.

Ostendorf, F., Liebermann, D. & Ploner, C. J., 2010. Human thalamus contributes to perceptual stability across
eye movements. *Proceedings of the National Academy of Sciences, B,* 107(3), pp. 1229-1234.

Pifl, C., Kish, S. J. & Hornykiewicz, O., 2012. Thalamic noradrenaline in Parkinson's disease: deficits suggest role
in motor and non-notor symptoms. *Movement Disorders,* 27(13), pp. 1618-1224.

R Development Core Team , 2014. *R: A language and environment for statistical computing.* Vienna(Austria): R
Foundation for Statistical Computing. Available from http://www.R-project.org/.

Ross, J., Morrone, M. C., Goldberg, M. E. & Burr, D. C., 2001. Changes in visual perception at the time of
saccades. *Trends in Cognitive Sciences,* Volume 24, pp. 113-121.

Silva, M. F. et al., 2005. Independent patterns of damage within magno-, parvo- and koniocellular pathways in
Parkinson's disease. *Brain,* 128(10), pp. 2260-2271.

Sommer, M. A. & Wurtz, R. H., 2002. A pathway in primate brain for internal monitoring of movements.
Science, Volume 296, pp. 1480-1482.

Sommer, M. A. & Wurtz, R. H., 2006. Influence of thalamus on spatial visual processing in frontal cortex.
Nature, Volume 444, pp. 374-377.

Sommer, M. A. & Wurtz, R. H., 2008. Brain circuits for the internal monitoring of movements. *Annual Reviews*
of Neuroscience, Volume 31, pp. 317-338.

Terao, Y. et al., 2011. Initiation and ihibitory control of saccades with the progression of Parkinson's disease -
Changes in three major drives converging on the the superior colliculus. *Neuropsychologia,* Volume 49, pp.
1794-1806.

Thakkar, K. N., Diwadkar, V. A. & Rolfs, M., 2017. Oculomotor prediction: A window into the psychotic mind.
Trend in Cognitive Sciences, 21(5), p. 344–356.

Thakkar, K. N., Schall, J. D., Heckers, S. & Park, S., 2015. Disrupted saccadic corollary discharge in schizophrenia.
Journal of Neuroscience, 35(27), pp. 9935-9945.

Watson, A. B. & Pelli, D. G., 1983. QUEST: A Bayesian adaptive psychometric method. *Perception &*
Psychophysics, 33(2), pp. 113-120.

Weil, R. S. et al., 2016. Visual dysfunction in Parkinson's disease. *Brain,* Volume 139, pp. 2827-2843.

Wurtz, R. H., 2008. Neuronal mechanisms of visual stability. *Vision Research,* 48(20), pp. 2070-2089.