1

**Impairment in updating spatial representations in Parkinson's disease correlates with dopaminergic**

**deficiency**

Henry Railo* [1,2,3], Henri Olkoniemi[2], Enni Eeronheimo[2], Oona Pääkkönen[2], Juho Joutsa[4,5,6], and Valtteri

Kaasinen[7,8]

[1] Department of Clinical Neurophysiology, University of Turku and Turku University Hospital, FI-
20521, Turku, Finland
[2] Department of Psychology, University of Turku, FI-20014, Turku, Finland
[3] Brain and Mind Centre, University of Turku, FI-20014, Turku, Finland
[4] Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital and
Harvard Medical School, Charlestown, MA 02129
[5] Berenson-Allen Center for Noninvasive Brain Stimulation, Beth Israel Deaconess Medical Center
and Harvard Medical School, Boston, MA 02215
[6] Department of Neurology, University of Turku, FI-20521, Turku, Finland
[7] Division of Clinical Neurosciences, University of Turku and Turku University Hospital, FI-20521
Turku, Finland
[8] Turku PET Centre, University of Turku and Turku University Hospital, FI-20521 Turku, Finland
24

26
27
28
29
* Corresponding author:
Henry Railo,
Department of Psychology,
University of Turku,
Finland
Email: hmrail@utu.fi
Phone: +358-45-6765227

Patients with Parkinson's disease (PD) depend on visual feedback in their behavior more than

healthy individuals, suggesting deficits in monitoring their own movements. During eye movements,

internal monitoring signals (corollary discharge, CD) are relayed to the cortex via subcortical

structures whose function is compromised in PD due to deficits in the dopaminergic system. We

studied early-stage PD patients (N=14), and age-matched healthy control participants (N=14) using a

double-saccade task. In the task, participant fixates two targets in succession. In order to accurately

fixate the second target, the participant must correct for the displacement caused by the first

saccade. PD patients showed systematic deficits in fixating the second target although their ability to

fixate the first target was not significantly impaired. The deficit in fixating the second target was

observed especially when the patients made saccades towards the side with primary motor

symptoms. Across patients, the impairment in fixating the second target correlated with the integrity

of the dopaminergic system: Patients with lower striatal dopamine transporter binding made larger

errors. The direction of the errors suggests that the CD signal may be amplified in PD patients relative

to the motor signal. Furthermore, the results suggest that deficits in the dopaminergic system may

contribute to updating of spatial representations after movement. This could in part aggravate motor

symptoms, and explain frequently occurring visual symptoms in PD.

Keywords: Parkinson's disease, corollary discharge, spatial remapping, dopamine, dopamine

transporter binding, saccadic eye movement

Motor behavior requires the brain to update spatial representations to take into account

changes that are produced by self-induced movements. This requirement is apparent in eye

movements: several times per second we sample the environment by rapidly relocating our eyes

using saccadic eye movements during which the image on the retina is displaced. Yet we experience a

stable visual world and precisely know the locations of objects in the world because changes in

perception that are due to the eye movements have been eliminated by the brain. This internal

monitoring of movements and updating of the spatial representation is assumed to rely on corollary

discharge (CD) signals that relay a copy of the motor command to sensory areas (Figure 1A; 1).

Patients with Parkinson's disease (PD) are known to depend on visual feedback in controlling

their movements, more so than healthy individuals, suggesting that the brain mechanisms that

internally monitor movements (e.g. CD) are deficient: When visual feedback is restricted PD patients

tend to undershoot the target 2,3. Previous research has hypothesized that PD patients have defects

in kinesthesia 2,3, and CD 4,5, but studies have not examined this issue in detail. We examined PD

patients' ability to monitor their own eye movements using a *double-saccade task* which has been

widely utilized to examine CD and spatial updating (e.g. 1,6,7). In the task, a participant must take into

account the image displacement caused by a first saccade to accurately perform a second saccade

(Figure 1B).

Saccadic eye movements are triggered by neurons in the superior colliculus, which projects to

oculomotor burst neurons in the pons 8. The superior colliculus receives driving input from various

cortical areas that mediate voluntary control of eye movements, and nuclei of the basal ganglia that

filter out appropriate saccade targets 9. During the initiation of eye movements, the superior

colliculus is assumed to relay the CD to the cortex via medial thalamus 1,10. In other words, as

saccades are triggered by the superior colliculus, cortical systems are updated by the CD signal

through the thalamus. The CD shifts the receptive fields of visually responsive neurons to the location

where the saccade is going to land 1,11. Studies done on monkeys demonstrate that inactivation of

the medial thalamus both disrupts the shift in the receptive field 12 and impairs the monkey's ability

to monitor its own eye movements 13.

Figure 1. A) A simplified presentation of CD signals (red arrow). CD signals are predictions from motor
commands sent to the sensory areas. B) The double-saccade task. In the baseline condition the
participants made two sequential saccades (white arrows) to targets. In the Memory condition, the
they made saccades to memorized target locations. C) The grey circles show the possible positions of
the targets. For data analysis, the coordinates of the targets/eye positions were transposed to
correspond to the case where the first target was at the upper right corner of the screen and the
second target directly below it (arrows represent eye movements). D) Predictions concerning the
double saccade task. If CD is diminished relative to the strength of the motor signal, the participants
fail to fully take into account the first eye movement they have performed – hence, their second
saccades are biased outward (orange dashed line). Amplified CD (relative to the strength of the motor
signal) produces the opposite pattern of results: the second saccade is biased inward (red dashed
line). Note that deficits in CD may affect both the angle and the amplitude of the second saccade. E)
The efficacy of sensorimotor control can also be examined on a trial-by-trial basis by measuring how
well the participants take into account errors in the first saccade 14. In the example, the participant's
first saccade falls short of the target T1. The grey line shows what would be the optimal amplitude
(length of arrow) and angle (β) of the second saccade to compensate for the error in the first saccade.

The second black arrow shows an example where the amplitude (length) and angle (α) of the second
saccade are sub-optimal (i.e. participant overestimates the length of the first saccade, for example,
due to amplified CD).
105
Defects in CD have been observed in neurological patients with lesions to the CD system [15,7].

Using the double-saccade task, Thakkar et al. [6] observed that in schizophrenia patients the errors in

the angle of the second saccade suggested reduced CD relative to motor signals. As schizophrenia has

been associated with hyperactive striatal dopamine neurotransmission [16], this raises the question

whether PD patients – who characteristically have nigrostriatal dopaminergic defects [17] – display a

reverse effect (i.e. amplified CD relative to motor signals). Thus, assuming that the dopaminergic

system contributes to the CD, PD patients should show signs of amplified CD when assessed with the

double-saccade task. While dopamine is not *per se* assumed to relay the CD information, it is crucial in

modulating the activity of basal ganglia and thereby motor behavior [18], and it is hypothesized to

meditate the integration of bottom-up sensory information and contextual information such as

previously performed movements (e.g. CD) [19,20]. The two possible outcomes of CD deficits (reduced

vs. amplified CD relative to the motor signals) are displayed in Figure 1D. Furthermore, as shown in

Figure 1E, impaired CD should also lead to systematic variation in the angle of the second saccade on

trial-to-trial basis [14,20].

Abnormal CD is also predicted by the site of neuronal dysfunction in PD. The decrease of

dopaminergic projections from the substantia nigra pars compacta (SNc) leads to an excessive

inhibition in basal ganglia, thalamus, superior colliculus, and cortex [21,22,23]. This is assumed to

explain in large part the characteristic motor symptoms of PD [18], which often also affects eye

movements [24,25,26]. Dopaminergic deficit in PD also results in abnormal functioning of the

structures assumed to relay the CD to cortex, most importantly the thalamus. In PD, the medial dorsal

nucleus of the thalamus – which is assumed to relay the saccade CD signal to the cortex 12 – displays

abnormal bursting activity, which is assumed to disturb the information flow to the cortex 27. It is also

possible that in PD the strength of the CD signal may be amplified *relative to the strength of the motor*

*signals*, due to inhibition in the neural mechanisms that produce eye movements in PD. That is, one

may assume that, when compared to the inhibition in the actual motor signals, the neural

mechanisms that mediate the CD signals are less affected in PD.

Deficits in CD could lead to problems in visual stability and contribute to the motor symptoms of

PD, such as posture and gait, as the patient's internal spatial model does not accurately reflect the

world. Abnormal CD could also contribute to visual symptoms observed in PD 28, such as

hallucinations 5,29.

**Results**

*Visual symptoms*

PD patients reported more visual symptoms compared to healthy controls, although this

difference was statistically non-significant (Mann-Whitney test; Table 1). The patient group reported

numerically more symptoms especially concerning general confusion about locations of objects, and

experiencing spatial shifts or jerks during saccades.

**Table 1. Visual symptoms (mean and SD)**

|  |  | Controls | PD | Z | p |
|---|---|---|---|---|---|
| General | Nausea | 0.07 (0.25) | 0.06 (0.24) | -0.05 | 0.96 |
|  | Spatial confusion | 0.00 (0.00) | 0.13 (0.34) | -1.39 | 0.16 |
|  | Hallucinations | 0.00 (0.00) | 0.00 (0.00) | 0.00 | 1.00 |
|  |  |  |  |  |  |
| Saccade | Vertigo | 0.14 (0.51) | 0.13 (0.33) | -0.45 | 0.65 |
|  | Spatial shift | 0.00 (0.00) | 0.13 (0.33) | -1.39 | 0.16 |
|  | Motion | 0.07 (0.25) | 0.13 (0.33) | -0.57 | 0.36 |
|  |  |  |  |  |  |
| Smooth pursuit | Vertigo | 0.00 (0.00) | 0.00 (0.00) | 0.00 | 1.00 |
|  | Spatial shift | 0.00 (0.00) | 0.00 (0.00) | 0.00 | 1.00 |
|  | Motion | 0.00 (0.00) | 0.00 (0.00) | 0.00 | 1.00 |
| Total |  | 0.03 (0.09) | 0.07 (0.11) | -1.13 | 0.26 |

149
150
151
*Saccadic suppression*

Visual thresholds did not differ between the control group and the PD group during fixation (t =

-.13), saccade (t = .15), or after saccade (t = -.91). These factors were hence removed from the linear

model used to analyze visual threshold. The pruned model showed that both groups displayed clear

saccadic suppression as visual thresholds increased during saccades ($\beta$ = 1.96, 95% CI = [1.48, 2.44], t

= 8.06) and right after saccades ($\beta$ = 0.78, CI = [0.30, 1.25], t = 3.21) when compared to thresholds

during fixation (see Figure 2). This indicates that the patients' visual sensitivity was similar to the

control group.

Figure 2. Visual thresholds during fixation, saccade, and after saccade for each group. The unit of the y axis is the logarithm of Michelson contrast of the stimulus.

*Double-saccade task*

<u>*Horizontal and vertical errors*</u>

Figure 3A shows the participants' accuracy in fixating the first and second targets. The data was

analyzed using linear mixed-effects models that included the predictors group (Control vs. PD),

condition (Baseline vs. Memory), group × condition interaction, and the running trial number

(centered). The models were pruned by removing predictors that did not significantly (t < |2|)

contribute to the model. Horizontal and vertical errors were modelled separately.

*Saccades to the first target.* Although the fixations to the first target concentrated around the

actual target position (Fig. 3A upper panels), saccades towards the first target were, on average,

somewhat hypometric (i.e. they did not quite reach the target) in the Baseline condition in both

groups (*x error*: $\beta$ = -23. 49, CI = [-31.19, -15.83], t = -6.11; *y error*: $\beta$ = -48.19, CI = [-64.33, -32.00], t = -

5.82; in the results, the $\beta$ values show the size of the average effect in pixels). In the Memory

condition, fixations to the first target became horizontally more accurate in both groups ($\beta$ = 12.26, CI

= [0.59, 23.76], t = 2.11). However, in PD patients the saccades became vertically more hypometric in

the Memory condition (y error, PD × Memory: $\beta$ = -42.47, CI = [-78.05, -7.00], t = 2.33), whereas in

control group vertical errors somewhat decreased in the Memory condition (t = 1.61).

Figure 3. A) Fixation maps for the first target (T1, upper panels) and second target (T2, lower panels),
separately for control participants (left) and patients (right). The red and blue dots denote individual
fixations in the Baseline and Memory conditions, respectively. The red and blue crosses show the
respective group averages; the lengths of the lines is 95% confidence interval of the mean. B) Amount
of error as a function of laterality in the PD group. Ipsilateral refers to targets that were presented at
the same side as the motor symptoms. Error bars are the 95% CIs of means. Unit of error is pixels (22
pixels is ~1⁰).

As motor symptoms in PD are often asymmetric, we analyzed if the accuracy of fixations was

modulated by whether the targets were presented ipsi- or contralaterally with respect to the

predominant motor symptoms. This analysis was restricted to PD patients who showed asymmetric

symptoms (N=12). The results, displayed in Figure 3B, revealed that fixations to the first target were

more inaccurate (horizontally and vertically) when the target was ipsilateral with respect to symptoms

(*x error*: $\beta$ = -11.71, CI = [-21.48, -1.93], t = -2.34; *y error*: $\beta$ = -19.71, CI = [-19.85, -1.55], t = -2.29).

Moreover, as shown in Fig. 3B (upper panel), in the Memory condition horizontal error was reduced

when the targets were contralateral to motor symptoms (Memory × Laterality: $\beta$ = -18.91, CI = [-

33.95, -3.87], t = -2.46). Vertical error revealed also an interaction between Memory condition and

target laterality: In the Baseline condition vertical errors increased at ipsilateral locations, whereas

the effect reversed in the Memory condition ($\beta$ = 16.36, CI = [2.27, 30.40], t = 2.27), but as seen from

Fig. 3B (lower panel) the size of this effect is negligible. In summary, these results show that PD

patients have selective deficits especially in making horizontal saccades to memorized targets that are

at the same side as their primary motor symptoms.

*Saccades to the second target.* Fixations to the second target (Fig. 3A lower panels) were also on

average hypometric in the Baseline condition in both groups (*x error*: $\beta$ = -15. 48, CI = [-23.87, -7.14], t

= -3.68; *y error*: $\beta$ = -26.10, CI = [-44.91, -32.00], t = -7.33). During the Memory condition, fixations

became hypermetric, on average, i.e. they were biased away from the center of the screen in the

horizontal direction in both groups ($\beta$ = 18.32, CI = [0.94, 35.70], t = 2.10). Although Fig. 3A suggests

that this effect was smaller in PD patients, this Group × Condition interaction did not reach statistical

significance (t = -1.83). A further analysis performed on the patient group showed that when the

targets were presented at the same side as motor symptoms, fixations to the second target were on

average biased towards the center of the screen in horizontal direction in both conditions ($\beta$ = -14.44,

CI = [-22.46, -6.38], t = -3.52). This means that the PD patients show errors consistent with amplified

CD when they make saccades towards the side with primary motor symptoms. However, these results

should be taken with caution as the bias observed in the fixations to the second target could be in

part caused by biases in fixating the first target. To control for this possible confound, the analyses

concerning the bias in the second saccade were also performed in trial-by-trial basis (see Fig. 1E) in

section *Compensation in angle and amplitude following the first saccade.*

*Modulation by DAT binding.* Finally, we wanted to see if the accuracy of fixations correlated

with striatal dopamine transporter (DAT) binding in PD patients. This analysis focused on mean striatal

DAT binding. In addition to the condition (Baseline vs. Memory) and trial number factors, the model

included a predictor labelling whether or not the targets were presented contralaterally with respect

to the hemisphere with smaller DAT binding (Contralateral factor), the overall DAT binding in the

more affected hemisphere (i.e. the one with lower binding), the motor MDS-UPDRS score of the

patient, and the interactions of these four predictors.

Horizontal or vertical errors of fixations to the *first target* were not modulated by DAT.

Horizontal errors of fixations to the *second target* become somewhat more accurate as mean

striatal DAT binding increased. This main effect was not statistically significant ($\beta$ = 24.84, CI = [-3.30,

52.0499], t = 1.7), but as shown in Figure 4A, the interaction between laterality and DAT suggested

that it was only observed for fixations performed to targets presented ipsilaterally to the more

affected hemisphere (contralateral × DAT: $\beta$ = -23.94, CI = [-44.65, -3.23], t = -2.3). Figure 4A shows

the averaged result across both memory and baseline condition, because no statistically significant

effect was observed for condition. UPDRS scores did not significantly modulate fixation errors (t <

|1|), and the modulatory effects by DAT binding were observed even though UPDRS main effect and

the contralateral × UPDRS interaction were also included in the model. A further three-way

interaction (Condition × UPDRS × DAT) indicated that in the Memory condition, the modulatory effect

by DAT binding (Fig. 4A), was reduced in participants with higher UPDRS scores ($β$ = -4.20, CI = [-8.17, -

0.24], t =-2.1).

Figure 4. Relationship between average striatal dopamine transporter binding and accuracy of
fixations to the second target. A) Horizontal error was modulated by average DAT binding for targets
ipsilateral (red line) to the more affected hemisphere (blue line = contralateral). B) Vertical error was
modulated by DAT binding, and this effect was stronger when the patients made fixations to
memorized targets (blue = Memory). DAT binding variable has been centered (i.e. zero represented
the mean). The dots represent the observed data, and the lines represent the modelled results.

Vertical errors in fixations to the *second target* were more clearly modulated by DAT binding, as

shown in Figure 4B. In the Memory condition the vertical accuracy of fixations increased as DAT

increased (Memory × DAT: $β$ = 44.00, CI = [13.63, 74.38], t = 2.8). No such effect was observed in the

Baseline condition (t = 0.4). Target laterality did not modulate this effect, and therefore Fig. 4B shows

the average of ipsi- and contralateral data. Again, UPDRS scores did not significantly predict fixation

accuracy (t < |1|), suggesting that the observed modulation by DAT does not simply reflect the

severity of the PD. Altogether these results suggest that dopamine specifically contributed to the

accuracy of the *second* saccades. This effect was most clearly observed in vertical errors, and in the

Memory condition (Fig. 4B) where the saccades are performed without external visual guidance (i.e.

they are based on CD).

*Compensation in angle and amplitude following the first saccade*

As explained in Fig. 1E we also examined how well the participants took into account the trial-

by-trial variability in the first saccade when executing the second saccade. In the regression models,

the predictor variables were the centered optimal angle/amplitude of the saccade (see Fig. 1E), group,

condition, all interaction between these three variables, and running trial number (centered).

As shown in Figure 5A-B, the trial-to-trial variability in the *angle* of the second saccade was

strongly predicted by the optimal angle required to perfectly fixate the second target ($\beta$ = 0.71, CI =

[0.67, 0.76], t = 29.62). In other words, the participants clearly had information about the error in the

endpoint of the first saccade which they could use to correctly aim to the second target. In the

Memory condition, the angle of the second saccade was smaller than in the Baseline condition. That

is, participants' saccades were biased towards the center of the screen (i.e. the initial fixation point; $\beta$

= -4.69, CI = [-7.75, -1.67], t = -3.09). Furthermore, in the Memory condition, the larger the optimal

angle, the larger was the bias towards the center of the screen (interaction: $\beta$ = -0.13, CI = [-0.20, -

0.02], t = -4.32). PD patients were not statistically significantly different from control participants (t $\leq$

|1|).

Also the *amplitude* of second saccade strongly correlated with the optimal amplitude ($\beta$ = 0.87,

CI = [0.78, 0.96], t = 19.46). Moreover, although the amplitudes of the second saccades did not differ

between groups (t = 0.04), there was an interaction between group and optimal amplitude: Patients'

second saccades become more hypometric as the optimal amplitude required to reach the target

increased (Fig. 5C-D, blue line: $\beta$ = -0.15, CI = [-0.27, -0.04], t = 2.61).

Figure 5. Actual saccade angles (A-B) and amplitudes (C-D) as a function of ideal angle/amplitude. Left
panels show the baseline condition, and the right panel the Memory condition. Red color denotes the
healthy control participants, and the blue color the patient group. Dots show the observed data
(individual saccades) and lines the fitted linear models. Note that angle/amplidute variables are
centered, i.e. negative values show that angle/amplitude was smaller than average. Unit is pixels (22
pixels is ~1º).

As with previous analyses, we examined whether the laterality of the motor symptoms

correlated with the angles/amplitudes of the second saccades. As shown in Figure 6, when the targets

were presented at the same side as the motor symptoms the compensation in saccade angle was

worse than when the targets were presented on the side with less motor symptoms – the second

saccades become biased towards the center of the screen, consistent with amplified CD ($\beta$ = -0.11, CI

= [-0.19, -0.033], t = -2.78). In other words, although the PD group did not in general as a group differ

from healthy controls in compensating for variability in the endpoint of the first saccade, such an

effect was observed when the laterality of the symptoms was considered. In other words, this means

that since in some patients the symptoms are predominantly on the right side, and in other patients

on the left side, the deficit is only observed when the laterality of the symptoms is taken into account.

Target side relative to motor symptoms did not modulate the precision of compensation for

saccade *amplitudes*.

304
305   Figure 6.  The angle of the second saccade as a function of ideal angle. Compensation for variation in
306    the endpoint of the first fixation is worse for targets that are at the same side as motor symptoms.
307
308

*Modulation by DAT binding.* We also examined whether the success of compensating for variability in

the endpoint of the first fixation correlated with dopamine transporter binding in PD patients. As

previously, the model included condition, target laterality (with respect to the more affected

hemisphere), the amount of DAT binding, the optimal angle to perfectly fixate the second target, and

the motor MDS-UPDRS score of the participant (and the interactions between these regressors).

As previously stated the actual *angles* of the saccades were strongly predicted by the optimal

angle required to reach the second target ($\beta$ = 0.74, CI = [0.67, 0.81], t = 20.3). As shown in Figure 7A

(blue dashed line) patients with lower DAT transporter binding values were more inefficient in taking

into account for the trial-to-trial variability in the endpoint of the first saccade and their saccades

become more biased towards the center of the screen (Ideal angle × DAT: $\beta$ = 0.17, CI = [0.070, 0.28],

t = 3.2). Furthermore, the modulation of the angle of the second saccade by DAT is not explained by

the severity of PD as the motor MDS-UPDRS score revealed similar modulatory effect (Ideal angle ×

UPDRS: $\beta$ = 0.0030, CI = [0.00081, 0.0053], t = 2.6). Note that although the $\beta$ coefficient of the Ideal

angle × UPDRS interaction (0.0030) is much smaller than the $\beta$ coefficient of the Ideal angle × DAT

interaction (0.17), the relative size of this effect is in fact very similar (due to the fact that the range of

motor MDS-UPDRS scores is much higher than DAT values). The modulatory effect by motor MDS-

UPDRS is visualized in Figure 7A as the red dotted line which overlaps the blue dashed line that

depicts modulation by DAT. In addition to the above mentioned two-way interactions, a Ideal angle ×

motor MDS-UPDRS × DAT interaction was observed ($\beta$ = 0.014, CI = [0.004, 0.025], t = 2.8). This three-

way interaction indicates that the modulation by DAT binding was stronger in participants with more

severe PD (i.e. higher UPDRS scores).

Next we examined whether DAT binding was associated with the ability to modulate the

*amplitude* of the second saccade to compensate for trial-to-trial variability in the first saccade.

Although patients were in general successful in making a second saccade whose amplitude closely

resembled the ideal amplitude required to perfectly fixate the second target ($\beta$ = 0.73, CI = [0.65,

334 0.81], t = 17.7), patients with lower DAT binding made more hypometric saccades (β = 53.94, CI =

335 [19.23, 88.65], t = 3.0), as shown in Figure 7B. UPDRS score did not statistically significantly modulate

336 saccade amplitudes in general (main effect of UPDRS: t = -0.3). However, patients with higher motor

337 MDS-UPDRS scores were more inefficient in making a close-to-ideal amplitude saccade to the second

338 target (Ideal amplitude × motor MDS-UPDRS: β = -0.010, CI = [-0.015, -0,0050], t = -3.7). Surprisingly,

339 as shown in Figure 7C, this effect was only observed when the targets were presented ipsilaterally

340 with respect to the hemisphere with lower DAT binding (Contralateral × Ideal amplitude × motor

341 MDS-UPDRS: β = 0.012, CI = [0.0048, 0,020], t = 3.1).

Figure 7. Modulation of compensation in the amplitude of the second saccade. A) Compensation for variation in the endpoint of the first fixation is modulated by DAT binding (blue dashed line) and MDS-UPDRS (red dotted line). The visualized effects represent 1 SD decrease (DAT) or increase (MDS-UPDRS) relative to the patient group average, which is plotted as a grey solid line. The circles represent the observed data. B) Correlation between the amplitude of the second saccade and DAT binding. The red and blue circles represent the (average) observed data in the Baseline and Memory condition respectively. The grey line shows the modelled results (Baseline and Memory conditions pooled, as no Condition × DAT interaction was observed). C) Participants with higher MDS-UPDRS scores were more inefficient in compensating for the trial-to-trial variability in the endpoint of the first saccade, but this effect was only observed when the patients made saccades to targets presented ipsilaterally with respect to the hemisphere with lower DAT binding (dotted red line). The grey solid line shows the modelled performance of the patients on average.

**Discussion**

We examined how well early PD patients can update their spatial representations during eye movements, and whether this ability correlates with the integrity of the dopaminergic system. The participants were asked to make eye movements to two targets presented in succession. The results showed that while the patients' general ability to fixate the first target was relatively intact, they showed systematic deficits in shifting their gaze to the second target when compared to age-matched neurologically healthy controls. This effect was most prominent when the patients made saccades towards the side with primary motor symptoms. The deficit in fixating the second target correlated with the integrity of the dopaminergic system: patients with lower dopamine transporter binding showed more marked deficits in fixating the second target, and their eye movements were biased towards the center of the screen (i.e. towards the starting point of the first saccade).

The results are consistent with the assumption that PD patients show an amplified CD relative to the motor signals: the brain systems that internally monitor the eye movements of the patients signal that the extent of the movement was larger than what was actually the case. Hence, when the patients make a second saccade quickly after the first one, they tend to undershoot the target of the second saccade. The present findings are especially interesting as previous research has shown that deficits in CD are associated with *reduced* CD relative to motor signals 13,6. However, there is no reason to assume that abnormal CD always results in reduction of the CD signal (although this is obviously the case when an area that conveys the CD signal is lesioned; 1).

The interpretation that PD patients show an amplified CD which correlates with the integrity of the dopaminergic system is consistent with the finding that schizophrenia patients reveal the reverse pattern of results when performing the double-saccade task: in contrast to the present results,

schizophrenia patients underestimate the size of the first saccade 6. Because schizophrenia has been

associated with increased basal ganglia dopamine neurotransmission 16, PD patients (who have

dopamine deficiency) should show an opposite effect assuming that the dopaminergic system

contributes to the CD. Previous studies have not tested whether the integrity of the dopaminergic

system correlates with performance in the double-saccade task. Our results suggest that the

dopamine system contributes to spatial remapping. Crucially, correlations between dopamine

transporter binding and saccade performance were observed only for fixations to the *second* target,

implying that the dopaminergic system contributes specifically to spatial remapping and not to

performing saccades in general. Furthermore, the modulation by DAT binding was observed even

when motor symptom severity of PD was controlled by including the motor MDS-UPDRS score to the

model, suggesting that the effect is not simply due to variation in the motor symptoms of PD, but

rather a specifically dopaminergic phenomenon.

The present results do not imply that the dopamine system per se carries the CD, but that

dopamine modulates the functioning of the network of brain regions that mediate the CD, or

otherwise mediates spatial remapping after saccades. The present study does not specify whether the

possible deficits in the CD system result from 1) the amplification of the CD signal, or 2) the decrease

in the strength of the "output" motor signal relative to CD. The crucial claim is that there may be a

mismatch between internal monitoring signals (CD), and output motor signals in PD. According to a

widely held view, the degeneration of neurons in the SNc 30,31 leads to an inhibition in subcortical

structures 21,23 that take part in relaying the CD to the cortex (basal ganglia, thalamus, superior

colliculus; 1). However, there could also be independent striatal changes in the dopamine system as

the nigral cell counts do not seem to correlate with striatal DAT binding 32. Also, cortical systems and

cortico-subcortical-systems in general, are affected in PD 33, and the spatial remapping after

movement are assumed to rely on computations performed on cortical level 11,34.

Amplified CD signal is not the only possible explanation for the present results. Previous studies

have emphasized the role of kinesthetic feedback in PD 3,2, and it could be argued that the errors in

fixating the second target are due to impaired kinesthetic feedback from the extra-ocular muscles. As

with CD, this interpretation assumes that kinesthetic feedback is amplified. Future studies should

attempt to disentangle the contributions of CD and kinesthetic feedback to spatial remapping in PD.

Furthermore, because the present study focused on the spatial accuracy of fixations, it remains to be

tested whether the proposed deficit in CD is also reflected in saccadic latencies, or whether spatial

biases are accompanied by changes in saccade latencies.

The present study employed a modified version of the double-saccade task. In the classical

version (e.g. 35,13,6) the visual targets are presented so briefly that they are removed from the

screen before the eye movements begin; hence the saccades cannot be guided by sensory

information. Because the classical double-saccade task-set up proved too demanding for elderly

subjects, we used a modified task where the targets were presented for a longer duration. We

expected that the PD patients would show deficits especially in the Memory condition, where the

saccades to the two target locations are performed solely based on memory. However, the present

results show that PD patients are impaired in their second saccade also when the first saccade could

be directed based on visual information. This observation speaks against the interpretation that the

observed results are solely due to impaired CD or kinesthetic feedback. On the other hand, it could be

argued that if CD or kinesthesia is impaired, the bias in their internal spatial representation may

contribute to performance even when sensory information can be used to guide behavior because

behavior and perception are based on the interplay between incoming sensory information and

internal models (predictions such as the CD). The fact that performance was deteriorated both in

Baseline and in Memory condition suggest that the impairment in PD may be related to more general

difficulties in updating spatial representations, or binding sensory information with current internal

spatial representation. The dopamine system may thus be part of a more general prediction system

that estimates the uncertainty of internal models of the world 19,20. Consistent with the present

findings, dopamine contributes to encoding and updating spatial representations in hippocampal

neurons 36,37.

PD patients are known to undershoot targets when performing eye movements 25,24, and

studies have also reported that PD affects saccadic response times 38. The present results showed

that the patient group was relatively unimpaired in making spatially accurate saccades to the first

target. This may be due to the fact that the patients were at an early stage of PD. The main difference

in the performance in the two groups was that the PD patients' performance (in making the first

saccade) deteriorated in the Memory condition, which is consistent with previous reports 39,25,24.

Altogether, this suggests that the patients' deficits in the present task were not simply related to

difficulties in executing eye movements in general. Our results replicate the finding that deficits in

saccades are typically worse on the side with larger primary motor symptoms 26. Importantly, the

deficits in spatial remapping were also dependent on the laterality of the primary motor symptoms:

the hypothesized deficits in the trial-to-trial variability in the angle of the second saccade were only

observed when the laterality of motor symptoms was taken into account (see Fig. 6). This is consistent

with the observation that thalamic lesions (that interfere with the CD) produce lateralized defects

(e.g. 12). However, we also observed some more surprising – and perhaps counterintuitive – findings

related to lateralization. First, the modulation of the horizontal error in fixations to the second target

by dopamine transporter binding was observed only for saccades *ipsilateral* to the hemisphere with

lower DAT binding (Fig. 4A; although note that this effect was weak). Secondly, PD patients with

higher motor symptoms severity scores made larger errors in fixating the second target, but again this

only applied to saccades *ipsilateral* to the hemisphere with lower DAT binding (Fig. 7C). While these

findings could be false positive findings due to limited sample size (Fig. 4A), or they may also suggest

that the laterality effects of dopaminergic deficits are not straightforward (Fig. 7C). PD patients

sometimes display motor symptoms ipsilateral to the dopaminergic deficit 40,41. Another possibility

is that while DAT function in the contralateral striatum is affected severely already in early PD

(possibility due to floor effect), correlations can be more easily detected in the ipsilateral side.

The PD patients in the present study were at the early stages of PD, generally did not report

visual symptoms, and the visual sensitivity of the patients was similar to the control group. It is

important to note that with the double-saccade task we were able to detect subclinical deteriorating

visual functions that the patients were unaware of. These disturbances, together with the shown

associations with DAT binding suggest that, as the disease progresses and the dopaminergic deficit

deepens, relevant clinical CD-associated motor and visual symptoms will emerge. Advanced PD

patients have many visual symptoms 28 and abnormal CD may lead to deficits in experienced visual

stability in PD. More generally, the present study suggests that motor deficits in PD are partly

produced by difficulties in internal monitoring of movements. Studies with patients with more

progressed PD and other dopaminergic diseases are required to confirm these findings.

**Method**

*Participants*

Fourteen PD patients (8 women; $M_{age}$ = 62, range: 45–75 years, 13 right-handed and one

ambidextrous) and fourteen right-handed neurologically healthy control participants (7 women; $M_{age}$

= 67, range 44–78) took part in the experiment. PD patients were diagnosed using the UK Brain Bank

criteria, all diagnoses were supported by marked striatal defects in dopamine transporter SPECT and

none of the patients presented features suggestive of atypical parkinsonism. All participants had

normal or corrected-to-normal vision. Three control participants had to be excluded from the analysis

due to technical problems during testing. In most cases the control participants were the spouses of

the PD patients. The duration of motor symptoms in PD patients was on average 24 months (range:

3–72 months), and the Hoehn-Yahr stages of the patients were as follows: Six patients at stage I,

seven patients at stage II, and one patient at stage III. The patients had a mean motor MDS-UPDRS

score of 37 (range 11–95, SD = 20). In eight patients the motor symptoms were primarily on the left

side, in four patients in the right side, and two patients had nearly symmetrical motor symptoms.

None of the patients had depression (mean BDI = 3.8, range 1-12, SD = 2.8), and none showed signs of

abnormal cognitive abilities as measured by the Mini-Mental State Examination (mean 27.6, range 26-

30).

Eleven patients underwent a voluntary break in medication before the experiment, and did not

take medication on the morning of the test session. One patient did not have any medication, and

two participants did not go through a break in medication. Three patients were on levodopa, nine

patients were treated with MAO-B inhibitors and/or dopamine agonists and medication was unknown

for one patient.

488  The experiment was approved by the Ethics committee of the hospital district of South-West

489 Finland, and a written informed consent was obtained prior the study.  The study was conducted

490 according to the principle of the Declaration of Helsinki.

492  *Tasks, stimuli and procedure*

493  In addition to performing the double saccade task, we also measured the strength of saccadic

494 suppression in the participants by measuring visual thresholds during saccades and fixation. Previous

495 studies have reported increased visual threshold in PD 42,43, and saccadic suppression has been

496 hypothesized to be triggered by extra-retinal (e.g. CD) signals 44. Finally, the participants were

497 queried about their subjective experience of visual stability using a questionnaire described below.

498 We assumed that deficits in CD and visual updating may lead to subjectively experienced symptoms of

499 unstable vision.

500  To assess whether the participants experience symptoms of unstable vision they were asked

501 whether they perceived motion, experienced vertigo, or had feelings that the world has "changed

502 position" during saccadic or smooth pursuit eye movements. Smooth pursuit eye movements were

503 included in the questionnaire as control questions, since we hypothesized that possible impairments

504 in visual stability selectively affect saccades. In addition participants were asked if they had trouble

505 estimating the positions of objects, had experienced visual hallucinations, or had experienced any

506 other visual symptoms. Answers were given on a four step scale (0 = never … 3 = often).

507  In the double saccade task a participant is asked to shift her gaze to two briefly presented

508 targets. In the traditional form of the task the targets are presented so briefly that when the

509 participant initiates the first saccade, both visual targets have disappeared from the screen. Thus the

participant must use her internal monitoring signals (i.e. CD) to take into account the spatial change

produced by the first saccade in order to correctly fixate the second target 35,1,6. Pilot experiments

for the present study showed that the traditional double saccade task was too challenging for the

elderly participants (the participants had trouble perceiving and making saccades to the locations of

the briefly presented target stimuli). In order to make the task easier for the participants, we modified

the task so that each target stimulus is presented for a longer duration, but the participants were

instructed to initiate the saccades only after both targets had disappeared from the screen (Memory

condition; Figure 1B). While this means that short-term visuospatial memory may contribute to

performance more in our modified version than in the traditional form of the task (which taxes the

sensory memory more), the key idea behind the double saccade task remains unchanged: the

participants need to take into account the spatial change produced by the first saccade in order to

correctly fixate the second target. In the Baseline condition (Fig. 1B), which was physically identical to

the experimental condition, the participants were not required to wait until the targets have

disappeared and could initiate their eye movements voluntarily.

For the double saccade task, round (.6$^o$) white target stimuli were presented on a black

background. The double saccade task began with a nine point calibration of the eye-tracker (i.e.

whole screen was calibrated). After calibration, each trial began with a white fixation point presented

at the center of the screen. This fixation point was also used as a (eye-tracking) drift-correction point

in the beginning of each trial. After the participants correctly fixated the center of the screen for 1.2-

1.5 sec, the two target stimuli were presented. The first stimulus was always presented in one of the

four quadrants of the screen (e.g. upper right quadrant as in Fig. 1B) for 500 msec ~16 degrees from

the fixation point. Right after the first target, a second target was presented for 130 msec on the

horizontal meridian at the same side of the screen as the first target. However, the exact horizontal

position of the second target varied from trial to trial (there were four equally likely positions,

separated by a minimum of ~2.4 degrees). The possible locations of the saccade targets are presented

in Figure 1C. Each participant performed the baseline and memory conditions once. Thirty trials were

collected for each condition. Before the experimental runs, each participant performed at least 10

practice trials without eye-tracking.

Visual 81% thresholds were measured by presenting a horizontally oriented, low-frequency,

luminance modulated Gabor grating (diameter ~12°, 0.33 cycles per degree, phase varied randomly

on each trial) in the upper or lower visual field (center of the Gabor was 7 degrees from the center of

the screen) for one screen refresh (11.8 msec). The stimuli were presented on mid-grey background.

Participants' task was to report whether the Gabor stimulus was presented in the upper or lower

visual field using the up and down arrow keys. The detection threshold was determined by adaptively

varying the Michelson contrast of the Gabor using a Bayesian staircase method QUEST 45. This

procedure is similar to previously reported means of measuring saccadic suppression 46. Each

participant performed three different conditions during which visual thresholds were measured:

during fixation, during saccade, and right after saccade. When the thresholds were measured during

fixations participants kept fixating to a fixation dot, presented in the middle of the screen. In the

'saccade condition', the visual threshold was determined for Gabor stimuli presented while

participants performed ~19 degree saccades from a fixation point on the left side of the screen to a

black dot at the center of the screen. In the 'after saccade' condition the Gabor stimulus was

presented ~100 msec after the saccade had ended. Participants always initiated the saccades

voluntarily after a brief period of fixation during which the eye tracker was drift corrected. To enable

accurate tracking of horizontal eye-position, the eye-tracker was calibrated by three calibration points

presented on the horizontal meridian (left side, right side and center of the screen).

In both tasks (double saccade and visual threshold) the stimuli were presented using

Psychtoolbox 47 running on Matlab 2014b. Eye movements were recorded using EyeLink 1000 eye-

tracker (SR Research) that was operated using the EyeLink Toolbox 48. The eye movement

registration was done monocularly, typically for the right eye, using 1000 Hz sampling frequency. The

stimuli were presented on a 19˝ CRT screen with a screen resolution of $1024 \times 768$ pixels and 85Hz

refresh rate. Participants were seated 70 cm from the screen and a head rest was used to stabilize the

head.

*SPECT imaging*

To image dopamine transporter (DAT) binding, all PD patients underwent brain [123I]FP-CIT

([123I]N- ω-fluoropropyl-2β-carbomethoxy-3β-(4-iodophenyl) nortropane) SPECT. The interval

between the scanning and behavioural testing (e.g. double saccade task) was 8-29 months. Prior to

scanning, thyroidal update of the ligand was blocked by administering oral potassium perchlorate

(250 mg) 60 minutes before the injection of [123I]FP-CIT. An 185 MBq bolus of [123I]FP-CIT was

administered and scanning performed 3 hours after the injection. Imaging was performed as

described earlier 49. The images were reconstructed and analysed using BRASS (Hermes Medical

Solutions AB, Stockholm, Sweden). Specific binding rations (SBRs) for the caudate nucleus and the

putamen were quantified using occipital cortex as the reference region $SBR_{ROI} = (ROI - OCC) / OCC$,

where ROI refers to the uptake in the region of interest and OCC in the occipital cortex. Although

occipital cortex may contain minute concentrations of DAT, it has been shown to be a valid reference

region in [123I]FP-CIT imaging 50. Mean striatal values (caudate, anterior putamen and posterior

putamen) in the left and right hemisphere were used for correlation analyses. In statistical analysis of

the relationship between DAT binding and behavioural performance, the analyses always included a

factor that indicated whether saccades were made ipsilaterally or contralaterally with respect to the

hemisphere with lower binding.

*Data analysis*

Single-trial eye movement data (accuracy of fixations) was analyzed using linear mixed-effects

models using the lme4 package 51 in R statistical software 52. Because the analysis is performed on a

single-trial basis rather than on aggregated means, mixed-effects models enable taking into account

individual differences, in addition to group-level effects. Participants were included in the model as

random variables, that is, each individual participant's data was fitted with individual intercepts and

slopes (concerning the Baseline/Memory condition). The fixed-effect predictors included in the

models are described in the Results section. The models were pruned by removing fixed-effects

predictors that did not significantly ($t < |2|$) contribute to the model. The visual detection thresholds

(saccadic suppression) were analyzed using linear mixed effects models with separate intercepts for

each subject. As there is no standardized way to calculate the degrees of freedom and the

corresponding p values in mixed-effects models 53, we do not report p values. However, the

statistical significance of an effect can be checked from the confidence intervals (CI): whenever the

value zero is included in the CI, the effect is not considered statistically significant. Another reason to

not report p values is to draw the reader's attention to the size of the effect 54. Error bars in all the

figures are 95% CIs calculated from 1000 bootstrap samples.

In the double-saccade task, separate models were built for horizontal and vertical errors

(relative to the target location). Only trials where the first saccade was directed towards the first

target and where the participants made a following saccade towards the second target were included

in the analysis (5.8% of trials were excluded). For each participant, the first fixation to the first and

second targets were included in the analysis, except for cases were the first fixation was very brief

and rapidly followed by a corrective saccade towards the correct location. Before the analysis of

fixation accuracy, the data was transposed to correspond to the case where the first target was at the

upper right corner of the screen and the second target directly below it (Fig. 1C). These preprocessing

steps were performed in Matlab 2014b.

Datasets can be downloaded from the Open Science Framework (https://osf.io/bqnzh).

**Additional information**

*Competing Interests*

The authors declare that they have no competing interests.

*Author contributions*

H.R. and V.K. designed the research. H.O. and H.R. programmed the eye-tracking experiments.

E.E. and O.P. collected the eye movement and questionnaire data. V.K. and J.J. collected and analyzed

the SPECT data. H.R. analyzed the eye-movement data and wrote the first draft of the manuscript.

V.K., H.O. and J.J. reviewed and edited the manuscript.

*Acknowledgements*

HR was funded by Turku Institute of Advanced Studies and Academy of Finland (grant #308533), and

received a grant from the Finnish Cultural Foundation. HO was funded by the Finnish Cultural

Foundation. JJ was funded by the Academy of Finland (grant # 295580), the Finnish Medical

Foundation and the Orion Research Foundation, and has received travel grants from Orion and

Abbvie, and a research grant from Lundbeck. VK was funded by the Turku University Hospital (ERVA-

funds).

**References**

1. Sommer, M. A. & Wurtz, R. H., Brain circuits for the internal monitoring of movements. *Annual Reviews of Neuroscience* **31**, 317-338 (2008).

2. Klockgether, T., Borutta, M., Rapp, H., Spieker, S. & Dichgans, J., A defect of kinesthesia in Parkinson's disease. *Movement Disorders* **10** (4), 460-465 (1995).

3. Jobst, E. E., Melnick, M. E., Byl, N. N., Dowling, G. A. & Aminoff, M. J., Sensory perception in Parkinson's disease. *Archives of Neurology* **54**, 450-454 (1997).

4. Moore, A. P., Impaired sensorimotor integration in Parkinsonism and dyskinesia: a role for corollary discharge? *Journal of Neurology, Neurosurgery and Psychiatry* **50**, 544-552 (1987).

5. Diederich, N. J., Stebbins, G., Schiltz, C. & Goetz, C. G., Are patients with Parkinson's disease blind to blindsight? *Brain* **137**, 1838-1849 (2014).

6. Thakkar, K. N., Schall, J. D., Heckers, S. & Park, S., Disrupted saccadic corollary discharge in schizophrenia. *Journal of Neuroscience* **35** (27), 9935-9945 (2015).

7. Ostendorf, F., Liebermann, D. & Ploner, C. J., Human thalamus contributes to perceptual stability across eye movements. *Proceedings of the National Academy of Sciences, B* **107** (3), 1229-1234 (2010).

8. Moschovakis, A. K., The superior colliculus and eye movement control. *Current Opinion in Neurobiology* **6** (6), 811-816 (1996).

9. Hikosaka, O., Takikawa, Y. & Kawagoe, R., Role of the basal ganglia in the control of purposive saccadic eye movements. *Physiological Reviews* **80** (3), 953-978 (2000).

10. Wurtz, R. H., Neuronal mechanisms of visual stability. *Vision Research* **48** (20), 2070-2089 (2008).

11. Duhamel, J. R., Colby, C. L. & Goldberg, M. E., The updating of the representation of visual space in parietal cortex by intended eye movements. *Science* **255**, 90-92 (1992).

12. Sommer, M. A. & Wurtz, R. H., Influence of thalamus on spatial visual processing in frontal cortex. *Nature* **444**, 374-377 (2006).

13. Sommer, M. A. & Wurtz, R. H., A pathway in primate brain for internal monitoring of movements. *Science* **296**, 1480-1482 (2002).

14. Joiner, W. M., FitzGibbon, E. J. & Wurtz, H. R., Amplitudes and directions of individual saccades can be adjusted by corollary discharge. *Journal of Vision* **10** (2), 1-12 (2010).

15. Bellebaum, C., Daum, I., Koch, B., Schwarz, M. & Hoffman, K.-P., The role of the human thalamus in processing corollary discharge. *Brain* **128**, 1139-1154 (2005).

16. Howes, O. D. & Kapur, S., The dopamine hypothesis of schizophrenia: Version III - The final common pathway. *Schizophrenia Bulletin* **35** (3), 549-562 (2009).

17. Kordower, J. H. *et al.*, Disease duration of the integrity of the nigrostriatal system in Parkinson's disease. *Brain* **136**, 2419-2431 (2013).

18. Galvan, A. & Wichmann, T., Pathophysiology of parkinsonism. *Clinical Neurophysiology* **119** (7), 1459-1474 (2008).

19. Friston, K. J. *et al.*, Dopamine, affordance, and active inference. *PLoS Computational Biology* **8**, e1002327 (2012).

20. Redgrave, P. & Gurney, K., The short-latency dopamine signal: a role in discovering novel actions? *Nature Neuroscience* **7**, 967-975 (2006).

21. Munuera, J., Morel, P., Duhamel, J.-R & Deneve, S., Optimal sensorimotor control in eye movement sequences. *The Journal of Neuroscience* **29** (10), 3026-3035 (2009).

22. DeLong, M. R. & Wichmann, T., Circuits and circuit disorders of the basal ganglia. *Neurological Review* **64**, 20-24 (2007).

23. Terao, Y., Fukuda, H., Ugawa, Y. & Hikosaka, O., New perspectives on the pathophysiology of Parkinson's disease as assessed by saccade performance: A clinical review. *Clinical Neurophysiology* **124**, 1491-1506 (2013).

24. Albin, R. L., Young, A. B. & Penney, J. B., The functional anatomy of basal ganglia disorders. *Trend in*

*Neurosciences* **12** (10), 366-375 (1989).

25. Crawford, T. J., Henderson, L. & Kennard, C., Abnormalities of nonvisually-guided eye movements in Parkinson's disease. *Brain* **112**, 1573-1586 (1989).

26. Terao, Y. *et al.*, Initiation and ihibitory control of saccades with the progression of Parkinson's disease - Changes in three major drives converging on the the superior colliculus. *Neuropsychologia* **49**, 1794-1806 (2011).

27. Choi, S.-M. *et al.*, Directional asymmetries of saccadic hypometria in patients with early Parkinson's disease and unilateral symptoms. *Europen Neurology* **66**, 170-174 (2011).

28. Pifl, C., Kish, S. J. & Hornykiewicz, O., Thalamic noradrenaline in Parkinson's disease: deficits suggest role in motor and non-notor symptoms. *Movement Disorders* **27** (13), 1618-1224 (2012).

29. Weil, R. S. *et al.*, Visual dysfunction in Parkinson's disease. *Brain* **139**, 2827-2843 (2016).

30. Thakkar, K. N., Diwadkar, V. A. & Rolfs, M., Oculomotor prediction: A window into the psychotic mind. *Trend in Cognitive Sciences* **21** (5), 344–356 (2017).

31. Rinne, J., Nigral degeneration in Parkinson's disease. *Movement Disorders* **8**, S31-S35 (1993).

32. Damier, P., Hirsch, E. C., Agid, Y. & Graybiel, A. M., The substantia nigra of the human brain. II. Patterns of loss of dopamine-containing neurons in Parkinson's disease. *Brain* **122** (8), 1437-1448 (1999).

33. Saari, L. *et al.*, Dopamine transporter imaging does not predict the number of nigral neurons in Parkinson disease. *Neurology* **88** (15), 1461-1467 (2017).

34. Caligiore, D. *et al.*, Parkinson's disease as a system-level disorder. *npj Parkinson's Disease* **2**, 16025 (2016).

35. Duhamel, J.-R., Goldberg, M. E., Fitzgibbon, E. J., Sirigu, A. & Grafman, J., Saccadic dysmetria in a patient with a right frontoparietal lesion. *Brain* **115**, 1387-1402 (1992).

36. Hallett, P. E. & Lightstone, A. D., Saccadic eye movements to flashed targets. *Vision Research* **16**, 107-114 (1976).

37. Tran, A. H. *et al.*, Dopamine D1 Receptor Modulates Hippocampal Representation Plasticity to Spatial Novelty. *Journal of Neuroscience* **28** (50), 13390-13400 (2008).

38. McNamara, C. G., Tejero-Cantero, Á., Trouche, S., Campo-Urriza, N. & Dupret, D., Dopaminergic neurons promote hippocampal reactivation and spatial memory persistence. *Nature Neuroscience* **17** (12), 1658-1660 (2014).

39. Chambers, J. M. & Prescott, T. J., Response times for visually guided saccades in persons with Parkinson's disease: a meta-analytic review. *Neuropsychologia* **48** (4), 887-899 (2010).

40. Le Heron, C. J., MacAskill, M. R. & Anderson, T. J., Memory-guided saccades in Parkinson's disease: long delays can improve performance. *Experimental Brain Research* **161**, 293-298 (2005).

41. Kaasinen, V., Ipsilateral deficits of dopaminergic neurotransmission in Parkinson's disease. *Annals of Clinical and Translational Neurology* **3** (1), 21-26 (2015).

42. Aguirregomozcorta, M. *et al.*, Patients with rest-tremor and scans with ipsilateral dopaminergic deficit. *Journal of Neurology* **260** (4), 1132-1135 (2013).

43. Silva, M. F. *et al.*, Independent patterns of damage within magno-, parvo- and koniocellular pathways in Parkinson's disease. *Brain* **128** (10), 2260-2271 (2005).

44. Diederich, N. J., Raman, R., Leurgans, S. & Goetz, C. G., Progressive worsening of spatial and chromatic processing deficits in Parkinson disease. *Archievs of Neurology* **59** (8), 1249-1252 (2002).

45. Ross, J., Morrone, M. C., Goldberg, M. E. & Burr, D. C., Changes in visual perception at the time of saccades. *Trends in Cognitive Sciences* **24**, 113-121 (2001).

46. Watson, A. B. & Pelli, D. G., QUEST: A Bayesian adaptive psychometric method. *Perception & Psychophysics* **33** (2), 113-120 (1983).

47. Burr, D. C., Morrone, M. C. & Ross, J., Selective suppression of the magnicellular visual pathway during saccadic eye movements. *Nature* **371**, 511-515 (1994).

48. Brainard, D. H., The psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

49. Cornelissen, F. W., Peters, E. M. & Palmer, J., The Eyelink toolbox: Eye tracking with MATLAB and the Psychophysics toolbox. *Behavior Research Methods, Instruments, & Computers* **34** (4), 613-617 (2002).

50. Kaasinen, V., Kinos, M., Joutsa, J., Seppänen, M. & Noponen, T., Differences in striatal dopamine transporter density between tremor dominant and non-tremor Parkinson's disease. *European Journal of Nuclear Medicine and Molecular Imaging* **41** (10), 1931-1937 (2014).

51. Joutsa, J., Johansson, J. & Kaasinen, V., Is occipital cortex a valid reference region in 123I-FP-CIT SPECT imaging? *Clinical Nuclear Medicine* **40** (7), 615-616 (2015).

52. Bates, D., Maechler, M., Bolker, B. M. & Walker, S., lme4: Linear mixed-effects models using Eigen and S4. *R package version 1.1-7, URL: http://CRAN.R-project.org/package=lme4* (2014).

53. R Development Core Team , *R: A language and environment for statistical computing* (R Foundation for

Statistical Computing. Available from http://www.R-project.org/, Vienna, 2014).

54. Baayen, R. H., Davidson, D. J. & Bates, D. M., Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language* **59**, 390-412 (2008).

55. Cumming, G., *Understanding The New Statistics: Effect Sizes, Confidence Intervals, and Meta-Analysis* (Routledge, New York, 2012).

56. Ross, J., Burr, D. & Morrone, C., Suppression of the magnocellular pathways during saccades. *Behavioural Brain Research* **80**, 1-8 (1996).