TuBA: Tunable Biclustering Algorithm Reveals Clinically Relevant Tumor

Transcriptional Profiles in Breast Cancer

Amartya Singh[1,2], Gyan Bhanot[1,2,3], Hossein Khiabanian[1,2,3,4]

[1] Department of Physics and Astronomy, Rutgers University, Piscataway, NJ, USA.

[2] Center for Systems and Computational Biology, Rutgers Cancer Institute, Rutgers

University, New Brunswick, NJ, USA.

[3] Department of Molecular Biology and Biochemistry, Rutgers University, Piscataway,

NJ, USA.

[4] Department of Pathology and Laboratory Medicine, Rutgers Robert Wood Johnson

Medical School, Rutgers University, New Brunswick, NJ, USA.

Emails – AS: as2197@scarletmail.rutgers.edu; GB: gyanbhanot@gmail.com; HK:

h.khiabanian@rutgers.edu

Corresponding addresses

Hossein Khiabanian, PhD

Assistant Professor of Pathology in Medical Informatics

Rutgers Cancer Institute of New Jersey, Rutgers University

195 Little Albany Street, New Brunswick, NJ, 08903-2681

Phone: (732) 235 7554; E-mail: h.khiabanian@rutgers.edu

ABSTRACT

Background: Traditional clustering approaches for gene expression data are not well adapted to address the complexity and heterogeneity of tumors, where small sets of genes may be aberrantly co-expressed in specific subsets of tumors. Biclustering algorithms that perform local clustering on subsets of genes and conditions help address this problem. We propose a graph-based Tunable Biclustering Algorithm (TuBA) based on a novel pairwise proximity measure, examining the relationship of samples at the extremes of genes' expression profiles to identify similarly altered signatures.

Results: TuBA's predictions are consistent in 3,940 Breast Invasive Carcinoma (BRCA) samples from three independent sources, employing different technologies for measuring gene expression (RNASeq and Microarray). Over 60% of biclusters identified independently in each dataset had significant agreement in their gene sets, as well as similar clinical implications. About 50% of biclusters were enriched in the ER-/HER2- (or basal-like) subtype, while more than 50% were associated with transcriptionally active copy number changes. Biclusters representing gene co-expression patterns in stromal tissue were also identified in tumor specimens.

Conclusion: TuBA offers a simple biclustering method that can identify biologically relevant gene co-expression signatures not captured by traditional unsupervised clustering approaches. It complements biclustering approaches that are designed to identify constant or coherent submatrices in gene expression datasets, and outperforms them in identifying a multitude of altered transcriptional profiles that are associated with observed genomic heterogeneity of diseased states in breast cancer, both within and across tumor subtypes, a promising step in understanding disease heterogeneity, and a necessary first step in individualized therapy.

48 **KEYWORDS**

49 Clustering, Gene co-expression, Tumor heterogeneity, Copy number aberrations, Breast

50 Invasive Carcinoma

BACKGROUND

The first step in organizing and analyzing high-throughput gene expression datasets is to group together (cluster) genes, or samples based on some mathematical measure of similarity between the respective entities of interest. Since a priori knowledge about both the relevant genes, and the unique phenotypic characteristics of samples is usually limited, clustering is often performed in an unsupervised manner [1-5]. Quite frequently, measures of similarity (such as the Pearson correlation coefficient, Spearman correlation coefficient, Mutual Information, etc.) are employed to quantify the level of similarity between every pair of genes (or samples) across all samples (or genes). Such an approach is known as global clustering. In case of datasets with a heterogeneous assortment of samples, only a small subset of genes in a fraction of the total set of samples may be co-regulated in specific cellular processes. This is especially true for diseases like cancer that manifest a plethora of diseased phenotypes. In gene expression datasets comprising tumor samples, depending on the heterogeneity of the diseased states, there may be multiple distinct transcriptional alterations that are exhibited by multiple (not necessarily exclusive) subsets of the tumor samples. Moreover, it is well known that even in normal cells, the same genes can regulate and participate in multiple distinct pathways, depending on the context. Therefore, global clustering is not an optimal approach to identify co-expressed sets of genes or samples in gene expression datasets associated with heterogeneous diseases.

To address these concerns, a variety of local biclustering algorithms have been proposed that satisfy the following requirements: 1) a cluster of genes is defined with respect to only a subset of conditions (patient samples) and vice versa, and 2) the clusters are not exclusive and/or exhaustive – i.e. a gene/condition may belong to more than one cluster or to none at all [6-9]. Based on the type of biclusters and the

mathematical formulation used to discover them, biclustering techniques are categorized

by Oghabian *et al* [10] into four classes: (i) correlation maximization methods that

identify subsets of genes and samples where the expression values of genes (or

samples) is highly correlated across samples (or genes) [6]; (ii) variance minimization

methods that identify biclusters where the expression values have low variance among

the selected genes or conditions or both [11]; (iii) two-way clustering methods that

iteratively perform one-way clustering on the genes and samples [12], and (iv)

probabilistic and generative methods that employ stochastic approaches to discover

genes (or samples) that are similarly expressed in subsets of samples (or genes) [13,

14]. Another classification scheme proposed by Pontes et al [15, 16] categorizes the

generated biclusters based on their gene expression patterns into four classes:

(i) biclusters with constant values, (ii) biclusters with constant values on rows (genes) or

columns (conditions), (iii) biclusters with additive and/or multiplicative relationships

between genes and conditions, and (iv) biclusters based on evidence that a subset of

genes is up-regulated or down-regulated across a subset of conditions without taking

into account actual expression values; data in such biclusters does not follow any

mathematical model.

In this paper, we introduce a graph-based method, called the Tunable Biclustering

Algorithm (TuBA), which discovers biclusters consistent with the latter category. TuBA is

based on a novel measure of proximity that identifies aberrantly co-expressed gene sets

within subsets of tumor samples that correspond to the expression extremals for the

genes. A key feature of the proximity measure used in TuBA is that it does not rely

explicitly on the actual gene expression values. We demonstrate the utility of TuBA by

applying it to three large, independent cohorts of breast invasive carcinoma (BRCA)

encompassing 3,940 patients. In addition to detecting known pathways and subtypes

associated with breast cancer, TuBA was able to uncover several novel sets of co-

expressed genes across subtypes that may be relevant as biomarkers for therapeutic

identification and intervention.

METHODS

Proximity measure. TuBA's proximity measure addresses the following question: in a

given gene expression dataset, which genes exhibit higher (or lower) expression levels

in the same subset of samples relative to the rest? In other words, if we only consider

the top (or bottom) x percentile samples for every gene, which gene-pairs share a

significant number of samples between their percentile sets? The number of samples

shared between any pair of percentile sets follows the hypergeometric distribution;

therefore, we can compute the significance (p-value) of overlaps between pairs of

percentile sets based on the numbers of shared samples by using the one-sided Fisher's

exact test. Thus, TuBA's proximity measure between two genes is defined by the

significance of overlaps between their respective percentile sets (**Fig. 1**).

In a real biological dataset, we expect the following two scenarios to arise: (i) subsets of

genes associated with particular biological processes/pathways are co-expressed in all

samples. In this case, it is reasonable to expect a significant agreement between the

sets of samples that exhibit higher (or lower) expression levels of the involved genes.

(ii) Alternatively, subsets of genes may be dysregulated via shared underlying

mechanisms, such that their expression levels are higher (or lower) compared to the rest

of the samples that are not influenced by that mechanism. The latter case is of particular

interest for datasets associated with diseased states, especially cancers, since these

gene co-expression signatures and their underlying mechanisms could help us identify

potential biomarkers with prognostic and/or predictive value. This is the basic motivation

behind standard different differential expression analyses as well [17]. However, unlike

usual differential co-expression analyses, our proximity measure does not rely on any

pre-specification of subtypes.

A salient feature of our proximity measure is that it does not model the distributions of

the measured expression levels of genes across samples. Moreover, it does not rely on

significant differences between the expression levels of genes in samples comprising the

extremal sets versus the rest of the samples. Thus, biologically relevant gene co-

expression signatures can be identified without restricting the analysis exclusively to

genes that exhibit differential expression across subsets of samples. In case of tumor

datasets, this increases the likelihood for identification of gene co-expression signatures

associated with the microenvironment. Another salient feature of our proximity measure

is that there is no penalty for relative changes in ranks of samples in the respective

percentile sets. This is important because, even if the ranks of matching samples are

significantly different in the two percentile sets, there is still valuable information to be

gleaned by virtue of the fact that these subsets of samples exhibit higher (or lower)

expression levels for a given gene-pair compared to all the other samples. This feature

of our proximity measure makes it less sensitive to noise compared to other proximity

measures, such as the Spearman's Rank correlation.

Graph-based Algorithm. For each gene, TuBA identifies samples in the upper-most (or

lower-most) percentile sets. Pairwise comparison between these percentile sets (using

the one-sided Fisher's exact test) identifies gene-pairs that share a statistically

significant number of samples. Each significant gene-pair is illustrated graphically as a

pair of nodes connected by an edge that represents the samples shared between their

percentile sets. The complete set of these pairwise graphs generates large graphs from

which robust gene co-expression signatures are recovered using the following iterative

process (**Fig. 2**):

1. The graph is pruned such that its elementary units are complete subgraphs
(cliques) of size 3 (triangles).

2. The largest clique (i.e. the seed) in the pruned graph is identified using the Bron-
Kerbosch algorithm [18]. In cases where the largest clique is not unique, the
union of all equally large cliques with a non-zero intersection of their nodes is
designated as the seed; the remaining largest cliques are identified as new
seeds in subsequent iterations.

3. The graph is trimmed by removing all the edges that contain any of the nodes in
the seed in step 2. This step significantly reduces the computation time required
to identify all the robust cliques in the graph.

4. Steps 2 and 3 are repeated till the graph has no elementary units left.

5. The seeds identified in steps 1-4 are exclusive in their gene sets, i.e., no two
seeds share a common gene. To create the bicluster, the seeds are reintroduced
sequentially into the original pruned graph from step 2, and nodes that share
edges with at least two nodes in each seed are identified and added to the seed.
The resulting graphs are the final biclusters obtained by TuBA.

Note, the requirement of largest cliques as seeds of our biclusters in step 2 is a key step
in our algorithm that enables the identification of shared altered mechanisms in subsets
of samples that exhibit high (or low) expression levels of these genes, while permitting
the study of sets of co-expressed genes that are associated with functionally related
pathways. Implicit in this requirement is the crucial assumption that the sets of genes
comprising the largest cliques are co-expressed in a subset of samples that comprise
the edges. This assumption is not the same as requiring all gene-pairs comprising the
seed to share identical sets of samples, or assuming that all the samples comprising the
final biclusters co-express all the genes present in the bicluster. Instead, our expectation

is that the samples present in the final biclusters are *enriched* in the top (or bottom)

samples for each gene comprising the biclusters. We have provided supporting evidence

for this expectation in the **Results** section.

Gene enrichment analysis of the gene sets in the biclusters can be used to identify their

functional relevance, and sample enrichment analysis can elucidate potential clinical

subtypes, underlying mechanisms of disease, and possible therapeutic approaches.

Furthermore, within each bicluster, genes can be assigned degrees, which are the total

number of edges that connect them to other genes in the graph. Genes with higher

degrees exhibit co-expression with other genes in a greater proportion of samples in the

bicluster. These could be candidate driver genes.

Tuning TuBA. TuBA has two adjustable parameters:

1. The Percentile cutoff: This parameter controls the number of top or bottom

samples (based on expression levels) considered for comparison between

genes.

2. The Overlap significance cutoff: The p-value threshold used to assess

significance of the overlap of samples between percentile sets for each gene-

pair. This parameter controls the minimum number of samples that must be

shared between percentile sets for an association to be considered significant,

and to be represented in the graph.

It is best to interpret the parameters as "knobs" that can be tuned to probe different

levels of heterogeneity in the population. For a given dataset, the choice of these two

parameters determines the number, as well as the composition of the final biclusters.

The choice of the first parameter determines the level of heterogeneity and/or the extent

of prevalence of genomic alterations in tumors that may be of interest to the investigator.

To illustrate how the choice of percentile cutoff could affect the identification of co-

expressed gene pairs, we consider a hypothetical dataset consisting of 200 samples.

Assume that there is a gene-pair in this dataset that is up regulated in 5% of the samples

such that the top 5% percentile sets (i.e. top 10 samples) are identical for the two genes.

Fig. 3A shows the significance values for overlaps (p-values calculated using one-sided

Fisher's exact test) as a function of the fraction of samples that overlap. At 5% percentile

cutoff, the significance value for an overlap fraction of 1 (complete match/overlap

between percentile sets) is $p = 4.45e\text{-}17$ (dark blue curve). If instead we had chosen an

upper 10% percentile cutoff, we would have an overlap fraction of 0.5 (overlap of 10 out

of 20 samples) corresponding to a significance value for overlap between 1e-10 and 1e-

5. Thus, an increase in the size of the percentile set results in loss of significance for

aberrant co-expression signatures found in smaller subsets of samples. This does not

imply that it is generally better to choose smaller percentile sets. In fact, a reduction in

the size of the percentile set increases the likelihood that a number of samples match

purely by chance. (Note the p-values corresponding to the overlap fraction of 1 for the

three cases in **Fig. 3A**, further demonstrated by permutation tests in Results.) Thus, as

we vary the size of the percentile set, there exists a trade-off between the sensitivity

(identification of altered transcriptional profiles in small subsets of population) on one

hand and the overlap significance on the other.

The choice of the second parameter – the extent of patient/sample overlap between

percentile sets – determines the gene-pairs that will be represented in the graph that is

explored iteratively to identify sets of co-expressed genes. As we lower the significance

of overlap (increasing p-values), new genes and samples get added to the graph

resulting in an increase in the number of edges (**Fig. 3B**). Further lowering of the overlap

significance results in the addition of many more edges to the graph, however this

addition is not accompanied by a proportional increase in the samples or genes added to

the graph. As more edges get added to the graph, the computational effort required for

finding maximal cliques increases. Since the maximal clique problem is NP-hard [19], it

can take exponential time to find all maximal cliques. Thus, the cutoff for the

significance of overlap is informed by the trade-off between the gain of new information

in the biclusters in terms of new samples and genes, and the number of edges added to

the graph that leads to a disproportionate increase in the computational effort. We

propose the following heuristic for choosing the cutoff value: the cutoff for the

significance level of overlap should be such that a decrease in the significance level by

an order of magnitude leads to an 40-60% increase in the number of edges that get

added to the graph (Note the number of edges that get added to the graph at overlap

significance p-values larger than 1e-20 in **Fig. 3B**).

Because the principal goal of TuBA is to identify subsets of genes that are co-expressed

at high (or low) levels within subsets of samples, the exact number of biclusters is not

biologically relevant despite possible small variations in their total number as the

algorithm is tuned. We investigated the consistency of TuBA's biclusters across different

choices of the parameters. We used the hypergeometric test to identify biclusters that

share significant fractions of their genes, and observed that despite a five-fold difference

in the significance level of overlap, there is greater than 80% agreement between the

sets of biclusters obtained for different choices of the overlap significance cutoff. The

results of the analysis are presented and discussed under **Robustness of TuBA's**

Biclusters in **Supplementary Methods** and **Supplementary Table 4**.

Datasets. We applied TuBA to three independent BRCA datasets that employed distinct

methods for measuring transcript levels: (i) TCGA RNASeq gene expression dataset

using the Illumina HiSeq 2000 RNA sequencing platform, (ii) METABRIC gene

expression dataset using the Illumina HT-12 v3 microarray platform, and (iii) six cohorts

with gene expression data from GEO using the Affymetrix HGU133A microarray

platform. To compare results among datasets, we applied TuBA to only their common

gene sets. For clinical association analysis, we prepared two separate datasets for

patients with known recurrence free survival (RFS) status (908 patients) and patients

with known PAM50 subtype annotation (522 patients) respectively. Henceforth, we will

refer to TCGA RFS, METABRIC RFS and GEO RFS datasets simply as TCGA,

METABRIC and GEO, respectively, and PAM50 datasets are indicated specifically.

1) TCGA – BRCA: The $\log_2(x+1)$ transformed RSEM normalized counts of Level 3 data

(2016-08-16 version), the clinical data (including relapse status and PAM50 subtype

annotation from the 2012 Nature study [20]) (2016-04-27 version), and gene-level copy

number variation (CNV) data, as estimated by GISTIC2 [21] (2016-08-16 version) were

downloaded from the UCSC Xena Portal (http://xena.ucsc.edu). Genes with zero

expression in all samples as well as the samples with NAs for any gene were removed

from the analysis.

2) METABRIC: Normalized gene expression data, the clinical file, and the copy number

file for the METABRIC study were downloaded from the cBioPortal

(http://www.cbioportal.org) on 2017-05-14 [22, 23]. Gene expression dataset of 1,970

samples that had both relapse status and PAM50 subtype annotation were used in this

study.

3) GEO: MAS5 normalized gene expression data and the clinical data with relapse

status were downloaded from [24] on 2017-05-10. The dataset comprises samples from

six independent cohorts. After processing, our gene expression dataset consisted of

1,062 patients with relapse status.

4) breastCancerNKI (Bicmix): Gene expression data from breast cancer study

published by van't Veer et al. in 2002 [25], and van de Vijver et al. in 2002 [26] was

downloaded in the form of an eSet using the breastCancerNKI package [27] in R. The

dataset was further processed by removing probes with > 10% missing values, and

imputing the missing values for the included probes [28].

5) ESTIMATE: Scores for the level of stromal cells present and the infiltration level of

immune cells in tumor tissues for 906 out of 908 samples for the TCGA – BRCA RNA-

SeqV2 dataset using the ESTIMATE algorithm were downloaded from

http://bioinformatics.mdanderson.org/estimate on 2017-10-12.

6) GTEx: RNASeq raw counts data from the Genotype-Tissue Expression (GTEx) portal

(www.gtexportal.org/home/) was downloaded on 2017-06-15. The dataset comprised of

all tissue samples currently available in the GTEx database. The 214 breast tissue

samples were identified and normalized using the DESeq package in R [29].

Statistical Analysis. All computations were performed with R 3.3.0 [30]. The igraph

package [31] was used to perform network/graph computations with some data

summary functions performed using the plyr package [32]. The figures with graphs

showing the genes for some of the biclusters were generated using Cytoscape v3.4.0

[33]. Permutation test was performed on the METABRIC dataset (1970 samples) with

upper percentile set size cutoff: 5%. For each gene, we permuted the labels of the

samples prior to ascertainment of the samples that corresponded to the top 5%

respectively. The significance values for overlaps between every pair of genes were

computed using the Fisher's exact test. We performed 100 iterations of these

permutation tests in total. The data.table package was used to handle data files, and

the ggplot2 package was used to make plots [34]. GeneSCF [35] was used to perform

gene set clustering based on functional annotation and to associate biclusters with

specific biological processes. A binary matrix with biclusters along the rows and samples

along the columns was generated to perform hierarchical clustering. Samples belonging

to respective biclusters were assigned a value of 1. The Hamming distance was used to

measure dissimilarity between the biclusters as well as the samples. All tests for

enrichments were done using the Fisher's exact test; where necessary the p-values

were corrected for multiple hypotheses testing using the Benjamini–Hochberg false

discovery rate method [36]. The details of the contingency tables for the tests are

provided in **Supplementary Methods**. All enrichments are reported at FDR < 0.05,

unless specified otherwise.

RESULTS

Benchmarking TuBA's Proximity Measure: TuBA's pairwise proximity measure for

genes is based on overlaps between their extremal subsets of samples. In comparison,

global pairwise linear correlation coefficients such as the Pearson's correlation

coefficient and Spearman's rank correlation coefficient are both susceptible to the

influence of outliers. Even in the absence of outliers, genes that are co-expressed

across all samples are expected to have significant overlaps between the subsets of

samples that correspond to their top (or bottom) percentile sets. Given these

observations, we tested the hypothesis that gene sets identified by global proximity

measures have significant overlap with those identified by TuBA within its biclusters.

We computed the Pearson's correlation coefficients between all possible pairs of genes

in the TCGA and METABRIC datasets, respectively. We shortlisted all gene-pairs with

the correlation coefficient greater than or equal to 0.6. We then employed our graph-

based algorithm to identify gene co-expression modules within the graphs. For TCGA

and METABRIC, we obtained 569 and 298 gene co-expression modules, respectively

(**Supplementary Table 1**). We investigated the association between the gene sets in

biclusters discovered by TuBA's proximity measure versus gene co-expression modules

identified by global correlation metrics by performing a hypergeometric test for gene

overlaps. The null hypothesis was the absence of significant overlap between their

respective sets of genes. More than 89% (316 out of 353 biclusters) of the biclusters

discovered by TuBA in the TCGA dataset comprised gene sets that were enriched in at

least one gene co-expression module (FDR < 0.001), while 86% (293 out of 340

biclusters) of the biclusters discovered by TuBA in the METABRIC dataset were

enriched in at least one module.

We performed a similar analysis using Spearman's rank correlation, with a cutoff of 0.6

for the correlation coefficient. We obtained 524 and 232 gene co-expression modules for

TCGA and METABRIC, respectively. More than 80% (285 out of 353 biclusters) of

TuBA's biclusters in the TCGA dataset comprised gene sets that were enriched in at

least one gene co-expression module (FDR < 0.001), while 73% (249 out of 340

biclusters) of the biclusters discovered by TuBA in the METABRIC dataset were

enriched in at least one module. Overall, we see a significant enrichment of gene sets in

TuBA's biclusters with the co-expression modules obtained by using the two global

proximity measures. This is in concordance with our hypothesis stated earlier.

As noted earlier, due to samples that exhibit aberrant/outlier expression of some genes,

the linear correlation coefficients can often get skewed to reflect greater pairwise

correlations between such genes that our graph-based algorithm can identify. However,

due to the global nature of these proximity measures, the resulting graphs lack any

information on the samples that might be associated with aberrant expression of these

genes. In other words, unlike the case for TuBA, the edges in these graphs do not

represent any subset of samples; they simply reflect an association between the genes

by virtue of their pairwise correlation coefficient being greater than the chosen cutoff.

The novel design of our proximity measure enables precise identification of co-

expressed gene sets, while discerning the subsets of samples that exhibit higher (or

lower) expression levels of these genes relative to the rest of the samples.

Choice of parameters for TuBA. For a given choice of the size of percentile set, TuBA

generates plots that illustrate the number of added genes, added edges, and added

samples as the overlap significance cutoff is varied. These are used to inform the choice

of the overlap significance cutoff based on our proposed heuristic. Since the

experimental platform and the total number of samples were different among the

analyzed datasets, the choice of the overlap significance cutoff varied (**Fig. 4**). For the

respective choices of the knobs, we obtained 353, 340, and 369 biclusters for the TCGA,

METABRIC, and GEO datasets, respectively (**Supplementary Table 2**). Permutation

tests showed that no gene-pairs had p-values less than the cutoffs (**Fig. S1**). Moreover,

after adjusting for multiple hypotheses testing, none of the gene-pair p-values were

statistically significant.

Enrichment of bicluster samples in top (bottom) sample sets of bicluster genes.

As pointed out earlier, the identification of largest cliques as seeds of our biclusters was

based on the expectation that samples present in the final biclusters were enriched

specifically in the up (down)-regulated samples for each gene comprising the cliques.

We tested the hypothesis that the subsets of samples comprising the biclusters were

enriched by the samples that comprise the top (bottom) sets for each gene in the

bicluster. For example, suppose a dataset consists of 1000 samples, wherein on

application of TuBA for high expression, one bicluster is identified to comprise 100

genes and 200 samples. For each of those 100 genes, we identify their top 200 samples

and test whether these 200 samples are enriched in the 200 samples comprising the

bicluster. The null hypothesis is that these two sets of samples are independent, and

therefore we should not expect to see statistically significant associations between them.

We applied this test for each of TuBA's biclusters in the TCGA, METABRIC, and GEO

datasets. For high expression, we observed that all genes in all 353 biclusters from

TCGA showed significant enrichment (hypergeometric test FDR < 0.001). In case of

METABRIC we observed two biclusters (bicluster 2 and bicluster 269) out of 340

biclusters with only 1% of their constituent genes not exhibiting enrichment, while in case

of GEO we observed only one bicluster (bicluster 22) out of 369 with 1% of its

constituent genes not exhibiting enrichment. For the low expression analysis of TCGA,

we observed two biclusters (bicluster 14 and bicluster 165) out of 203 biclusters that

comprised a few genes that did not exhibit enrichment. A crucial observation across all

the datasets was that even in the few biclusters that included a few genes with

enrichment FDR greater than 0.001, none of these genes were constituents of the seeds

of those biclusters. We therefore found it justified to rely on the subsets of genes that

comprise the seeds for future gene ontological enrichment tests. This enables us to

identify the core functional signatures of biclusters.

We performed a similar analysis for the bicluster samples. Following the previous

example of a high-expression bicluster with 100 genes and 200 samples, we aimed to

identify the genes in the bicluster that had a given sample in their top 200 samples

(based on the expression levels of the genes). Therefore, for each sample, we evaluated

whether their corresponding subset of genes had significant overlaps with the complete

set of genes comprising the bicluster. So, we tested the null hypothesis that overlaps

between them were not statistically significant. For high expression, we observed that

95% of biclusters (336 out of 353) from the TCGA, 97% of biclusters (329 out of 340) in

the METABRIC, and 89% of biclusters (328 out of 369) in the GEO databases had >

95% of samples enriched (hypergeometric test FDR < 0.001). For the low expression

analysis of TCGA, we observed that 98% of biclusters (199 out of 203) had 95% of their

samples enriched in the bottom 200 samples for the corresponding genes in the

biclusters. Based on these analyses, the FDR values for each gene (sample) in any

given bicluster can be viewed as their scores – the closer the value of the FDR is to zero

for a gene (sample), the stronger is the association of the gene (sample) to the bicluster.

Consistency of TuBA within a dataset. To investigate whether TuBA could

consistently discover biclusters within the TCGA RFS cohort, the 908 samples were

divided randomly into two groups of 454 samples each. This was done five times to

generate five pairs of datasets. TuBA was applied to each dataset pair using a percentile

set size of 5% and an overlap significance cut-off of FDR ≤ 1e-08. Pairwise comparisons

(between sets of genes) of biclusters from the five trials showed that on average 73%

biclusters from one dataset in each pair were enriched (FDR < 0.001) in at least one

bicluster from the other (**Supplementary Table 3**). We found a significant difference

(Mann-Whitney U-test $p < $ 1e-05) in the number of genes contained in biclusters that

matched among trials, compared to the number of genes in biclusters that did not; while

the median size of biclusters that matched was 20 (range: 3–840), the median size of

biclusters that did not match was 3 (range: 3–18) (note that 3 is the smallest-

sized bicluster generated by TuBA). Overall, TuBA was able to consistently identify

matching sets of co-expressed genes from randomly sampled subsets of data within a

dataset.

Consistency of TuBA's biclusters among independent datasets. Using common

sets of genes, we compared the biclusters obtained from: (i) TCGA and METABRIC, (ii)

TCGA and GEO, and (iii) METABRIC and GEO. Pairwise comparisons of biclusters

obtained from the two datasets were used to identify the biclusters that shared a

significant proportion of their genes (FDR < 0.001). In the TCGA vs. METABRIC

comparison, 64% of biclusters obtained in one dataset were enriched in at least one

bicluster in the other. In the TCGA vs. GEO comparison, 69% of biclusters obtained in

one dataset were enriched in at least one bicluster in the other. Finally, in the

METABRIC vs. GEO comparison, 76% of the biclusters obtained in one dataset were

enriched in at least one bicluster in the other. Once again, we found that the biclusters

that did not match were significantly smaller (median number of genes: 3-5) than the

biclusters that matched (median number of genes: 20-25) between the datasets (Mann-

Whitney U-test $p < 0.001$).

TuBA identifies subtype-specific biclusters. We classified BRCA samples based on

the expression levels of the ESR1 (ER) and ERBB2 (HER2) genes into four subtypes:

(i) ER-/HER2-, (ii) ER+/HER2-, (iii) ER-/HER2+, and (iv) ER+/HER2+ (where +

corresponds to over expressed and − corresponds to under expressed). A substantial

proportion of biclusters were enriched in the ER-/HER2- subtype − 53% for METABRIC

(**Fig. 5A and 5B**), 54% for TCGA (**Fig. 5C and Fig. 5D**), and 40% for GEO (**Fig. S6**)

(**Supplementary Table 5**).

According to the PAM50 classification, there are five subtypes of BRCA: (i) Basal-like,

(ii) Her2-enriched, (iii) Luminal A, (iv) Luminal B, and (v) Normal-like [37]. We observed a

significant fraction of biclusters enriched in the basal-like subtype − 52% for METABRIC

(**Fig. S4A and S4B**) and 55% for TCGA PAM50 (**Fig. S4C and S4D**). Although tumors

of the basal-like or triple negative subtypes accounted for only about 15% of all BRCAs

in the population, most of the altered expression profiles captured by our biclusters were

in tumors of this subtype.

TuBA identifies down-regulated subtype-specific biclusters in RNA-seq data. RNA

sequencing offers a significant advantage over microarray assays. Theoretically, only

the depth of sequencing limits the dynamic range of RNA-seq data [38, 39]. Given that

TCGA's RNA-seq data has adequate sequencing depth, we expected a reliable

quantification of even lowly expressed transcripts. We therefore applied TuBA to the

TCGA datasets to explore transcriptional profiles associated with low expression. We

found that 46% biclusters from TCGA were enriched in the ER-/HER2- subtype (**Fig.**

S5A and S5B), while 48% biclusters from the TCGA PAM50 dataset were enriched in

the basal-like subtype (**Fig. S4E and S4F**). Thus, biclusters associated with low

expression were predominantly enriched in the ER-/HER2- or basal-like subtypes. This

further underscores the tremendous heterogeneity of altered transcriptional profiles

within tumors of this subtype.

TuBA highlights biclusters with proximally located genes. We observed that several

biclusters discovered by TuBA across the three datasets comprise genes that are

proximally located on the chromosomes, suggesting copy number amplification (CNA)

as an underlying mechanism. Copy number data was used to calculate the significance

of the proportion of samples present in each bicluster that exhibited copy number gains.

For each gene in a given bicluster, we computed a p-value for the significance of the

proportion of samples with CNA present in the bicluster. These p-values were then

combined using Fisher's method to yield a single p-value for the bicluster. This showed

that 56% and 64% of biclusters from the METABRIC and TCGA datasets respectively

were enriched for CNA (FDR < 0.001). Closer scrutiny revealed that only 60 (18%)

biclusters from METABRIC were associated exclusively with CNA of proximally located

genes (**Fig. 4A**), the remaining biclusters associated with CNA were enriched in genes

from distant chromosomal locations (**Fig. 4B**). Similarly, 112 (32%) biclusters from

TCGA were associated with CNA of proximally located genes (**Fig. 4C**). Many of these

biclusters were associated with loci previously identified to exhibit copy number gains in

BRCA [40, 41]. In order to explore the association between the biclusters obtained from

the low expression analysis and loss of copy number, we repeated the copy number

analysis described above. We observed that 52% biclusters from the TCGA dataset

were enriched in copy number losses. However, only 21 biclusters contained genes

located on the same chromosome (**Fig. S5A**), the remaining biclusters associated with

copy number loss were enriched in genes from distant chromosomal locations. Similar

analyses for PAM50 subtype enrichment for METABRIC and TCGA are summarized in

Fig. S4.

To compare CNA associated biclusters between TCGA and METABRIC, we prepared

two datasets that contained genes that were common in the two cohorts (17,209 genes).

Pairwise comparison of the set of genes in the CNA enriched biclusters between the two

datasets revealed that ~61% of biclusters from TCGA matched at least one CNA

associated bicluster from METABRIC. On the other hand, 91% of biclusters from

METABRIC were enriched in at least one CNA associated bicluster from TCGA. This

suggests that most of the CNA enriched biclusters identified in the METABRIC

microarray dataset were independently identified in the RNASeq dataset of TCGA.

We also observed some biclusters with proximally located genes that were not

associated with gain in copy number. For TCGA, 14 biclusters out of 353 consisted of

genes located proximally, while 18 biclusters out of 340 for METABRIC consisted of

genes located near each other. Details of the genes and subtype-specific enrichments

for some of these biclusters are summarized in **Supplementary Table 6**. Examples of

biclusters from this category include the biclusters consisting of genes from the Cancer-

Testis antigens family – *MAGEA2*, *MAGEA3*, *MAGEA6*, *MAGEA10*, *CSAG1*, *CSAG2*,

CSAG3 (Xq28)/*CT45A3*, *CT45A5*, *CT45A6* (Xq26.3). These genes are known to be

aberrantly expressed in triple negative breast tumors [42], as well as in a few other

tumor types [43].

TuBA identifies biclusters associated with non-tumor expression signatures. We

also discovered biclusters that appeared to be associated with non-tumor cells. For

instance, biclusters associated with immune response were among the largest identified

independently in all three datasets. The top five Gene Ontology – Biological Processes

(GO-BP) terms for the bicluster associated with immune response were: T cell

co-stimulation, T cell receptor signaling pathway, T cell activation, regulation of immune

response, and positive regulation of T cell proliferation (**Supplementary Table 7**). This

indicates immune cell infiltration in a significant number of tumor samples. To

corroborate this, we stratified TCGA samples based on their ESTIMATE [44] scores for

the infiltration level of immune cells in tumor tissues into three groups – (i) top 25

percentile, (ii) intermediate 50 percentile, and (iii) bottom 25 percentile – and verified that

samples in these biclusters associated with immune response were enriched in samples

with the highest levels of immune infiltration (FDR < 0.001)

For all three datasets, we also observed a bicluster associated with the stromal adipose

tissue. The top 5 GO-BP terms for this bicluster were: response to glucose, triglyceride

biosynthetic process, triglyceride catabolic process, retinoid metabolic process, and

retinol metabolic process. An analysis based on the ESTIMATE scores for the level of

stromal cells present in tumor tissue of TCGA samples confirmed that this bicluster was

enriched within the top 25 percentile samples for stromal cell level. Subtype enrichment

revealed that the bicluster was enriched in ER-/HER2-, basal-like (PAM50), and normal-

like (PAM50) subtypes.

TuBA's proximity measure was applied to gene expression data from 214 normal breast

tissue samples from the Genotype-Tissue Expression (GTEx) public dataset. We

observed that only 6.75% of biclusters obtained for the TCGA versus GTEx comparison

were enriched in gene-pair associations identified in the GTEx dataset. The bicluster

associated with the adipose tissue signature was one of the biclusters found enriched in

GTEx. Another group of biclusters enriched in the three cancer datasets as well as in

GTEx, were those associated with translation and ribosomal assembly. The top 5 GO-

BP terms for these biclusters were: translation, rRNA processing, ribosomal small

subunit biogenesis, ribosomal large subunit assembly, and ribosomal large subunit

biogenesis. These biclusters were enriched in the ER-/HER2- subtype (FDR < 0.001).

TuBA identifies clinically pertinent biclusters. We performed a Kaplan-Meier (KM)

analysis of recurrence free survival (RFS), comparing the patients present in each

bicluster to the rest for METABRIC and GEO. (The number of patients with incidence of

recurrence in TCGA was insufficient for this kind of survival analysis to be statistically

robust.) As expected for METABRIC, patients in the bicluster (bicluster 25) associated

with the HER2 amplicon (17q12) had significantly shorter RFS time compared to the rest

(**Fig. S7**). This is because patients in the METABRIC study were enrolled before the

general availability of trastuzumab [45].

We also observed biclusters associated with CNA at the 8q24.3 locus in all three

datasets (TCGA – biclusters 39 and 113, METABRIC – biclusters 26, 56, and 167, GEO

– biclusters 16, 24, 37, 55, 74, 118, and 302). These patients also had significantly

shorter RFS times compared to those patients whose tumors did not have amplification

of this locus (**Fig. 6A**, **6B** and **6C**). A similar result was obtained when we restricted the

samples to ER+/HER2- tumors, validating an earlier observation that copy number gain

of the 8q24.3 locus may confer resistance to ER targeted therapy [46]. We note,

however, that biclusters with amplification of the 8q24.3 locus were enriched in the ER-

/HER2- subtype ($p < 0.001$). Hence, amplification of this locus may be even more

relevant in determining treatment for patients with ER-/HER2- breast cancers assigned

into an intermediate (ambiguous) risk class by Oncotype DX [46]. Genes at 8q24.3 that

may be considered promising candidates based on their degrees in the biclusters

include *PUF60*, *EXOSC4*, *COMMD5*, and *HSF1*. Specifically, *PUF60* is an RNA-binding

protein known to contribute to tumor progression by enabling increased *MYC* expression

and greater resistance to apoptosis [47].

For both METABRIC and GEO, patients in biclusters associated with copy number gains

of the 8p11.21-p11.23 loci (METABRIC – bicluster 289, GEO – bicluster 25) had

significantly shorter RFS times compared to patients without amplification of this locus

(**Fig. 6D**, **6E** and **6F**). We found that patients in this bicluster were enriched in the

Luminal B subtype, which has poorer prognosis than the Luminal A subtype among

ER+/HER2- tumors [48]. This suggested that amplification of the 8p11.21-p11.23 loci

may be another marker of potential failure of ER targeted therapy.

Similarly, we found that patients whose tumors have copy number gains of the 17q22-

q23.3 locus (METABRIC – biclusters 33 and 119, GEO – biclusters 15 and 160) had

significantly shorter RFS times compared to patients whose tumors do not exhibit such a

copy number gain (**Fig. 6G**, **6H** and **6I**). For METABRIC, this cohort was enriched in the

Luminal B (PAM50), ER+/HER2+, and ER-/HER2+ subtypes (FDR < 0.001). For GEO,

this cohort was enriched in the ER+/HER2+ and ER-/HER2+ subtypes (FDR < 0.05).

This suggests that amplification of this locus may confer additional risk of recurrence in

HER2+ breast cancers.

Note that the biclusters discussed above were not the only ones that exhibited

differential relapse outcomes. For METABRIC, 61 biclusters out of 340 were found to

exhibit differential relapse outcomes for the patients present in the biclusters. Out of

these 61 biclusters, 69% were enriched in the ER-/HER2- subtype (64% for basal-like)

with a significant proportion (67%) of these associated with copy number gains. For

GEO, there were 48 such biclusters (13%) that exhibited differential relapse outcomes,

25% of these were enriched in the ER-/HER2- subtype.

Tests for enrichment of biclusters in tumors of higher grades revealed that 8 biclusters

from TCGA were enriched in tumors of grade 3C. Some of these biclusters were

associated with GO-BP terms related to angiogenesis, vasculogenesis, blood vessel

maturation etc. For METABRIC, 4 biclusters were enriched in tumors of grade 3, out of

which 2 were associated with the HER2 amplicon (17q12). For GEO, 68 biclusters were

enriched in tumors of grade 3, including biclusters associated with CNA at the HER2

amplicon.

We also looked at the lymph node status of patients and observed that 4 biclusters in

TCGA were enriched in samples with positive lymph node status in the corresponding

patients. One was associated with the HER2 amplicon, while the others were associated

with CNA at the 8q22.1-q22.3 loci, 17q23.1-q23.3 loci and the 19q13.43 locus,

respectively. Similarly in METABRIC, we observed 4 biclusters enriched in samples with

positive lymph node status in the corresponding patients - 2 of them were associated

with copy number gains at the HER2 amplicon, the other 2 were associated with copy

number gains at 19q13.11-q13.12 and 1q21.3-q25.1, respectively. Interestingly,

biclusters associated with CNA at 8q24.3, 8p11.21-p11.23, and 17q22-q23.3 that

exhibited poor RFS outcomes were not enriched in tumors of higher grades or in

patients with positive lymph node status in any of the 3 datasets. In case of METABRIC,

we additionally confirmed that none of these biclusters (8q24.3, 8p11.21-p11.23,

17q23.1-q23.3) were among the 36 biclusters enriched in samples with the poorest

expected 5-year survival outcome (Nottingham Prognostic Index (NPI): > 5.4) [49, 50].

This highlights the importance of these altered transcriptomic signatures for

reclassification of patients into the category with higher risk of recurrence.

Hierarchical clustering of biclusters reveals shared mechanisms. Sample

membership based hierarchical clustering of biclusters revealed distinct groups of

biclusters that presumably share common functional mechanisms (**Fig. 7**). These

included clusters associated with cell cycle and proliferation, immune response, cell

adhesion (extracellular matrix), translation, mitochondrial translation, and ribosomal RNA

processing pathways. Since a significant fraction of our biclusters were associated with

copy number alterations, we also found distinct groups of biclusters associated with

significant copy number changes such as the ones associated with the HER2 amplicon,

the 8p11.21-p11.23 loci, or the 8q24.3 locus.

Similarly, we used hierarchical clustering to group samples that were enriched in similar

sets of biclusters, highlighting differential clinical outcomes. In particular, we observed 2

sets of samples enriched in biclusters associated with CNA at the 8q24.3 locus. In one

group, the samples were enriched in biclusters related to immune response; this group

showed significantly lower incidence of recurrence compared to those without

enrichment in immune response-related biclusters. Both of these sets of samples were

enriched in biclusters associated with cell division and proliferation. In contrast, we

observed a cluster of samples enriched in biclusters associated with 8q24.3 copy

number gain and a number of other loci, however these were not enriched in biclusters

associated with cell division and proliferation. This group exhibited low incidence of

recurrence. We also observed a cluster of samples with significantly poor RFS that were

enriched in biclusters associated with CNA at 17q25.1-q25.3, and in biclusters

associated with cell division and proliferation.

TuBA compared to other biclustering methods. TuBA's proximity measure

distinguishes its biclusters from those identified by other algorithms by leveraging the

size of the datasets to identify subsets of tumor samples that co-express subsets of

genes at their most extreme levels (high or low) relative to other samples. We

emphasize that TuBA is designed to identify biclusters with samples that correspond to

the extremals for the corresponding sets of genes, and does not consider other subset of

conditions for the same sets of genes for biclustering. In contrast, most biclustering

methods seek sub-matrices with constant, or coherent gene expression patterns. Given

this key difference, only those biclusters that exhibit such expression patterns in the

extremal (top or bottom) subsets of samples for some subsets of genes, are expected to

have agreement with the biclusters identified by TuBA. Therefore, a direct comparison

between the biclusters discovered by other algorithms, with the ones identified by TuBA

would necessarily be limited.

In their paper on DeBi [51], a novel biclustering method that identifies differentially

expressed biclusters based on a frequent itemset approach, the authors applied their

method to both synthetic and real gene expression datasets, including diffuse large B-

cell lymphoma (DLBCL) data comprising 661 genes and 180 samples [52]. Apart from

DeBi, they applied ISA [53], OPSM [54], QUBIC [55], and SAMBA [56] to this dataset;

we used these results to evaluate TuBA against these methods. In order to ensure a

uniform and unbiased comparison between the enrichment results for biclusters from

different algorithms, we used GeneSCF [35] to perform GO-BP enrichment on the

biclusters obtained by all methods. **Fig. 8A** shows the proportions of GO-BP- enriched

biclusters for five different significance levels (FDRs) − 0.001%, 0.1%, 0.5%, 1%, and

5%. For the FDR cutoff of less than 5%, almost all the biclusters for every algorithm

were enriched in at least one GO-BP term. TuBA had 2 non-enriched biclusters out of

94, SAMBA had 2 non-enriched biclusters out of 128, and QUBIC had 1 bicluster out of

100 that were not enriched in a GO-BP term (Supplementary Table 7). As the FDR cutoff

was lowered, TuBA had lower proportions of enriched biclusters compared to other

algorithms for the corresponding FDR cutoffs. This can be partly attributed to the fact

that the other algorithms discover biclusters that can have arbitrary overlaps between

their genes. Since most of biclusters discovered by other algorithms shared genes with

652 other biclusters, we could expect a certain amount of redundancy in enriched GO-BP

653 terms. In contrast, TuBA precludes any overlap between the genes of the seeds of its

654 biclusters. Moreover, its biclusters often include proximally located genes with aberrant

655 expression due to copy-number changes, which may not show enrichment in GO-BP

656 terms.

657 For a closer examination of the redundancy in the GO terms enrichment, we identified

658 the top-five GO-BP terms for every bicluster obtained by each algorithm (not every

659 bicluster was enriched in five distinct GO-BP terms, some had less than five, while

660 others were not enriched in any term). For each algorithm, we prepared lists of all the

661 unique GO-BP terms for the entire set of biclusters. The ratios of the number of

662 elements in these lists to the total number of biclusters for each algorithm at five different

663 significance levels show that TuBA identified biclusters enriched in a more extensive

664 array of biological process terms (**Fig. 8B**).

665 In addition to the DLBCL dataset, we also analyzed the TCGA dataset with the following

666 biclustering algorithms: (i) BIMAX [57], (ii) ISA, (iii) QUBIC, and (iv) SAMBA, using their

667 respective default parameters (**Supplementary Tables 7 and 8**). For succinct

668 descriptions of each of these algorithms we refer the reader to Prelic et al [8] and Pontes

669 et al [55]. We used the biclust package in R for BIMAX [58], the isa2 package in R for

670 ISA [59], the QUBIC package in R for QUBIC [60], and the Expander software for

671 running SAMBA [61]. **Fig. 8C** shows the proportion of GO-BP terms enriched in

672 biclusters of each algorithm for five different significance levels. TuBA compared

673 favorably with other algorithms, especially when we accounted for the redundancy of the

674 GO terms that were found enriched in the biclusters. We observed again that TuBA's

675 biclusters were enriched in a larger set of distinct biological process terms (**Fig. 8D**). We

676 observed similar results for METABRIC (**Fig. S8A**, **S8B**). The results of the comparative

677 analysis are summarized in **Supplementary Methods**.

In these analyses, the choice of the parameters is a crucial factor in determining the

performance of each biclustering method. It is possible that different results could be

obtained by more prudent choices of parameters for the other algorithms, however a

detailed analysis of optimal parameter choices for each of these algorithms is beyond

the scope of this study. We must point out for TuBA, that for any given dataset, there is

no optimal (or default) choice of its two parameters; the biclusters obtained for any given

choice of the parameters simply satisfy the basic requirements laid down by those

choices. We looked at GO-BP-term enrichments for TuBA's biclusters for five different

choices of the overlap cutoff (**Supplementary Methods**). Although, the total number of

biclusters obtained differed for each choice, the proportion of enriched biclusters at

different significance levels remained similar irrespective of the parameter choice

(**Fig. S9A**). Similarly, the ratio of the number of unique GO-BP terms and the total

number of biclusters was consistent across all five choices of overlap cutoffs (**Fig. S9B)**.

In earlier studies comparing biclustering algorithm [8, 57], synthetic datasets were

generated with constant, shifting, and/or scaling patterns of expression for subsets of

conditions and genes. The algorithms were evaluated based on how well they were able

to identify the known biclusters implanted in these synthetic datasets. Since TuBA is not

based on a mathematical model of the data in its biclusters, a comparison based on

synthetic datasets is not feasible. However, in case of tumor datasets we have the

benefit of complementary genomic data that could provide us with truth-known scenarios

for validation. For example, alterations at the genomic level can directly influence the

expression levels of genes; it is well known that a significant proportion of tumors across

multiple tumor types frequently exhibit genomic alterations such as gains or losses in the

copy numbers of genes. Quite often, these alterations are not limited to a single gene

but include multiple genes located at neighboring chromosomal locations. If such

alterations are located at transcriptionally active sites, then co-expression of the

neighboring genes that are affected by it will be observed. In BRCA for instance,

approximately 15-20% of tumors possess extensive gains in copy numbers of genes at

the 17q12 cytoband locus (includes *ERBB2 (Her2)*, *STARD3*, *GRB7*, *PNMT, PGAP3,*

MED1 etc.). Identification of co-expression of genes at this locus in the subset of

samples that are histologically HER2-positive (HER2+) represents a simple truth-known

scenario that can be used to verify whether a given biclustering algorithm identifies the

co-expression of these genes in the subset of samples that exhibit this alteration. We

identified HER2+ samples in the TCGA dataset, and for each biclustering algorithm

selected those biclusters that were enriched in these samples (hypergeometric test FDR

< 0.001). BIMAX and SAMBA did not discover any, but ISA identified two biclusters

(biclusters 71 and 72) enriched in HER2+ samples. Although the genes from the 17q12

amplicon – *ERBB2, STARD3, GRB7, PNMT, PGAP3, MED1* etc. were present in ISA's

enriched biclusters, they comprised a small subset within the genes in them – bicluster

71 had 639 genes, while bicluster 72 had 539 genes, respectively. QUBIC also had four

biclusters that were enriched in HER2+ samples, however they did not contain any

genes from the HER2 amplicon (including *ERBB2*). In contrast, not only did TuBA

identify a bicluster (bicluster 256) exclusively associated with the HER2 amplicon, it

identified many other biclusters associated exclusively with CNA of genes located near

each other.

In summary, apart from TuBA, only ISA identified co-expression of the genes located at

the HER2 amplicon. However, ISA's co-expression module corresponding to the

amplicon was embedded within much larger sets of genes. In the absence of information

about copy number gain of the *ERBB2* gene, it would be a challenge to explicitly identify

the co-expression module corresponding to the amplicon, and in turn infer the underlying

mechanism for their co-expression. TuBA successfully uncovers those co-expressed

sets of genes that are associated with CNA of neighboring sites on the chromosome,

and is particularly efficient at identifying transcriptionally active copy number gains, as

compared to other algorithms.

The nature of our proximity measure allows us to determine differential co-expression

signatures without the need to specify subsets of samples in advance. Gao *et. al.* [62]

proposed a biclustering method, *Bicmix*, based on a Bayesian statistical model to infer

subsets of co-regulated genes that covary in all samples, or in only a subset of samples.

They also developed a principled method to recover context-specific gene co-expression

networks from the sparse biclustering matrices obtained by Bicmix. They applied Bicmix

to the breastCancerNKI dataset and identified 432 genes that were differentially co-

expressed in ER+, and ER- samples. Out of these 432 genes, 430 were up-regulated in

ER- samples and down-regulated in ER+ samples, while 2 genes are down-regulated in

ER- samples and up-regulated in ER+ samples. We applied TuBA (for high expression)

to the same dataset with the following choice of parameters: (i) Percentile set size: 10%,

and (ii) Overlap significance cutoff: FDR ≤ 1e-08. We obtained 549 biclusters, several of

which comprised solely of probes that correspond to the same gene (Supplementary

Table 5). This is reasonable, since probes corresponding to the same gene are expected

to demonstrate higher expression levels in the same set of samples. We inquired

whether some of the biclusters discovered by TuBA corroborated the differential co-

expression signature between ER+ and ER- samples identified by Bicmix. Using Fisher's

exact test, we determined that the set of 430 genes up-regulated in ER- samples and

down-regulated in ER+ samples were enriched in 30 biclusters discovered by TuBA –

bicluster 5 shows the maximum enrichment (FDR < 1e-165). In fact, the genes that had

the highest degrees in the co-expression network discovered by Bicmix – *CD247*, *CD53*,

IL10RA, and *CXCR3* – were among the ones with highest degrees in bicluster 5

discovered by TuBA. The two genes (*SFRP2* and *COL12A1*) that were up-regulated in

ER+ samples and down regulated in ER- samples were also found to be co-expressed in

a TuBA bicluster (bicluster 115). TuBA also identified biclusters corresponding to

amplicons at 17q12 (HER2), enriched in ER- samples (FDR = 0.02); 8q24.3, enriched in

ER- (FDR = 0.003) samples; 17q25-q25.3, enriched in ER- samples (FDR = 7.09e-05).

Thus, in addition to the differential co-expression network identified by Bicmix, TuBA

recovers biclusters associated with genomic alterations such as CNA, several of which

are differentially expressed between ER+ and ER- samples. Overall, TuBA recovered

144 biclusters enriched in ER- samples, and 31 biclusters enriched in ER+ samples

(FDR < 0.05). This is consistent with our earlier observation that a significant proportion

of biclusters discovered independently in the TCGA, METABRIC, and GEO datasets,

were enriched in the ER-/HER2- subtype.

Runtime Analysis. TuBA's graph-based algorithm relies on the identification of largest

cliques, which is a computationally hard problem. Large graphs (both in terms of the

number of genes, and edges) can potentially lead to long computation times. The size

of our graphs is principally determined by the choice of the cutoff for the second

parameter – the significance level of overlap between percentile sets. We varied the

cutoffs for the TCGA and METABRIC datasets, respectively, such that the total number

of edges in the resultant graphs ranged between 10,000 and 250,000.

For choices of overlap cutoffs consistent with our suggested heuristic, we recorded

TuBA's computation time to generate final biclusters for each dataset (**Fig. 9A**). The

computation time for TCGA increased dramatically as the number of edges in the graphs

went beyond 150,000 edges. In particular, while the computation time for a graph with

200,000 edges for METABRIC was approximately 70 minutes, the computation time for

a graph of similar size for TCGA was approximately 42 hours. Thus, although

METABRIC is the larger dataset with 24,368 genes and 1,970 samples compared to

TCGA's 20,241 genes and 908 samples, more iterations were required to identify all the

largest cliques in the graphs for TCGA given its respective choices of parameters. We

therefore conclude that TuBA's computation time depends on the nature and complexity

of the graphs themselves.

We also investigated the impact of the size of datasets on computation time. We created

new subsets from the TCGA dataset by randomly sampling a fixed number of genes. We

varied the number of genes from 2,000 to 20,000, and created 10 randomly sampled

datasets for each gene number (**Fig. 9B**). For each individual run, we chose an overlap

cutoff consistent with our suggested heuristic, and ensured that comparable numbers of

edges were generated for different datasets.

We also investigated the impact of the number of samples in a dataset. For this, we

created five randomly selected subsets from the TCGA dataset each with 250, 500, 750,

and 908 samples, (**Fig. 9C**). As expected, TuBA's computation time did not depend

strongly on the number of samples in the datasets.

In its current implementation, using a 2.7 GHz Intel Xeon processor, and 48 GB of RAM.

TuBA has longer runtime than most other existing algorithms. Depending on the choice

of the overlap cutoff, the runtimes can vary between 15 and 120 minutes for datasets

with approximately 20,000 genes and 1,000 samples.

DISCUSSION

Global clustering approaches have successfully unveiled distinct disease subtypes in

tumors, prompting the community to look beyond traditional clinico-pathological

signatures to identify relevant disease processes. However, the extensive heterogeneity,

even within tumors of a given subtype, confounds the identification of many altered

transcriptional programs by such unsupervised clustering methods.

In this paper, we introduce an algorithm called TuBA based on a proximity measure

specifically designed to extract gene co-expression signatures that correspond to the

extremes of expression (both high and low for RNASeq data, and high for array-based

platform). This enables us to preferentially identify co-aberrant gene signatures

associated with the disease states of tumors. The identification of altered transcriptional

profiles can be particularly relevant for those tumors that have so far eluded targeted

drug development for therapy. This is exemplified by tumors of the basal-like or triple

negative subtypes for BRCA. Although these tumors account for only ~15% of all BRCAs

in the population, a significant fraction of biclusters identified by TuBA corresponded to

alterations associated with tumors of these subtypes. For each dataset, a simple

estimation of enrichment of samples in a given bicluster within any other bicluster,

revealed that the samples in the biclusters corresponding to CNA at 8p11.21-p11.23 or

17q12 were enriched (FDR < 0.001) independently in ~5% of all biclusters for both

TCGA and METABRIC, respectively. In sharp contrast, 30–40% of all biclusters were

enriched in samples with copy number gains at the 8q24.3 locus (FDR < 0.001).

Additionally, 51% of all biclusters obtained from the low expression analysis of TCGA

were enriched in the samples corresponding to the 8q24.3 bicluster. Previous studies

have also identified the amplicon at 8q24.3 by Representational Difference Analysis as a

location of oncogenic alterations in breast cancer that can occur independent of

neighboring *MYC* amplifications [63]. Although the 8q24.3 bicluster itself is enriched with

ER-/HER2- samples, these observations, together with poor RFS outcome observed

independently in both METABRIC and GEO, highlight this locus as a promising

prognostic marker for BRCA tumors, irrespective of subtype.

We must mention a notable exception in the biclusters discovered by TuBA for all three

datasets – none of the biclusters contained the *ESR1* gene, which codes for estrogen

receptor. Closer inspection revealed that *ESR1* had statistically significant associations

with several genes, however the level of significance of overlap with these genes was

much lower (FDR > 1e-07) than the chosen cut-off for all three datasets. While it is

known that about 70% of BRCA exhibit elevated expression of estrogen receptor, over-

expression of estrogen receptor may not be a sufficient condition to drive co-expression

of genes involved in other pathways [64]. This may explain why co-expression of *ESR1*

with other genes was not as significant as the other associations that were extracted and

summarized in TuBA's biclusters.

Apart from highlighting the heterogeneity of CNA-associated alterations in tumors of the

ER-/HER2- subtype or basal-like subtype, TuBA offered a glimpse into the utility, the

limitations, and the potential pitfalls with the current subtype classification approaches. In

the ER/HER2-based subtype enrichment, we observed a significant proportion of

biclusters that were not specifically enriched in any one of the four subtypes. For

instance, several CNA-associated biclusters from chromosome eight were not subtype-

enriched. In the case of PAM50 subtype classification however, we observed that most

of these biclusters were enriched in the Luminal B subtype for METABRIC (and to a

limited extent for TCGA). While this appears to indicate that PAM50 offers an

improvement on the traditional clinico-pathological approach to subtype classification, it

unfortunately fails to classify several samples associated with overexpression of ERBB2

as HER2-positive. As a consequence, several of our biclusters associated with the

HER2 amplicon and copy number gains in the neighboring locations on chromosome 17

(17q.21.1-q21.2 and 17q21.32-q21.33), for both METABRIC and TCGA, were observed

to be enriched in the Luminal B subtype. This corroborates the modest level of

agreement with PAM50 classification reported in [37], as well as disagreements in later

studies [65]. Given that trastuzumab is a clinically proven therapeutic drug for HER2+

tumors, misclassification of these patients into any other subtype can be highly

disadvantageous.

Change in copy number is often not a sufficient condition for elevated (or suppressed)

expression levels of transcripts, as there are multiple layers of regulation of transcription

in cells [66, 67]. TuBA specifically identifies sets of genes with copy number changes

that are transcriptionally active (or inactive), filtering out the ones that are unlikely to

influence disease progression. Moreover, the graph-based approach allows us to infer

the relative importance of each gene within a bicluster, based on its degree. In the case

of high expression analysis, the degree of each gene is an indicator of how frequently it

is expressed aberrantly at high levels by the subset of samples that comprise any given

bicluster. As an example, consider the CNA-associated bicluster from TCGA

corresponding to gains at the 8q22.1-q22.3 loci. The bicluster exhibited enrichment in

lymph node positive patients (the corresponding bicluster in METABRIC has a

significance level of *FDR* = 0.052 for patients with positive lymph node status). The gene

with the highest degree in the bicluster was *MTDH* (metadherin), which has been shown

to be associated with increased chemo resistance and metastasis in BRCA [68-70].

Clustering analysis of biclusters and samples based on the membership of samples

within biclusters allowed us to identify the sites that were altered concomitantly within the

same subsets of samples. Moreover, we improved our perspective on the tumor

microenvironment in the subsets of samples that exhibit non-tumor associated

signatures (such as immune, extracellular matrix, etc.). Differences in disease

progression due to distinct microenvironments in tumors with similar transcriptional

alterations can help us better understand the potential role of the microenvironment

within the context of tumors harboring these specific alterations. For instance, we

noticed a difference in RFS outcomes between two groups of patients that exhibit copy

number gains at 8q24.3; the group that was additionally associated with an immune

response signature was observed to have better RFS outcomes compared to the group

that did not exhibit a strong association with the immune response.

Unlike most biclustering methods, TuBA does not allow arbitrary overlaps between its

biclusters. This is because it is designed to discover biclusters with samples that

correspond to the extremals for the corresponding gene set; biclusters with other

conditions are not permitted for the same gene set. However, our biclusters are not

exclusive, and some overlap between their genes and samples is permitted. For

example, in case of an ER-/HER2- BRCA sample that exhibits CNA at 8q24.3, because

of high immune-cell infiltration in the tumor, the same sample may also be present in the

biclusters enriched in the sets of genes associated with immune response.

Another limitation of TuBA is that it can only be applied reliably for large datasets that

contain at least 100 samples. Depending on cohort heterogeneity, some of the overlaps

between percentile sets may not be significant in smaller datasets. However, the

deliberate design of our proximity measure leveraging the size of the datasets offers a

significant benefit – it not only enables the identification of the plethora of gene co-

aberrations associated with the tumors, but also enables the estimation of the extent or

prevalence of the identified alterations in the population. This is where the tunable

aspect of TuBA becomes relevant – the two knobs should be viewed as valuable aids

that help estimate the extents of the prevalence of various alterations in the tumor

population and their clinical relevance. Although transcriptomic changes are not the

ultimate determinants of progression, our algorithm holds the promise to improve

therapeutic selection and design by identifying significantly altered transcriptional

patterns associated with tumors.

CONCLUSION

TuBA is quite distinct from other biclustering algorithms, in that, it is designed to identify

biclusters with samples that correspond to the extremals for the corresponding sets of

genes. Most biclustering algorithms are designed to identify nearly constant or coherent

gene expression levels in subsets of genes across subsets of samples. However, we

were able to show that TuBA performs outperforms other algorithms in identification of

co-expressed genes located in transcriptionally active copy number altered sites.

Moreover, from a differential co-expression perspective, TuBA offers an advantage over

other methods, since no prior specification of subsets of samples (context) is necessary;

the nature of our proximity measure ensures that such differential co-expression

signatures are preferentially identified. Given these considerations, TuBA offers great

promise as a biclustering method that can identify biologically relevant gene co-

expression signatures that are not successfully captured by other unsupervised

clustering or biclustering approaches. These signatures, along with the ones identified

by other biclustering methods would enable a comprehensive understanding of the

underlying alterations and shared mechanisms in subsets of tumors.

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent to publish

Not applicable.

Availability of data and materials

TuBA is open-sourced and available in R scripts at https://github.com/KhiabanianLab.

TCGA dataset was obtained from UCSC Xena Portal (http://xena.ucsc.edu). METABRIC

dataset was obtained from the cBioPortal (http://www.cbioportal.org). GEO and

breastCancerNKI datasets were obtained from Gyorffy & Schafer [24], and van't Veer et

al. [25] and van de Vijver et al. [26], respectively.

Competing interests

The authors declare that they have no competing interests.

Funding

HK acknowledges support from the American Cancer Society (IRG-15-168-01). The

funding agencies had no role in the design of the study and collection, analysis, and

interpretation of data or in writing of the manuscript.

Authors' Contributions

AS, GB, and HK conceived the study and designed the algorithm. AS implemented the

algorithm and performed the statistical analyses. All authors contributed to the drafting of

the manuscript and critical discussion of the results. All authors read and approved the

final manuscript.

Acknowledgements

This research was partially supported by the Biomedical Informatics Shared Resource at

Rutgers Cancer Institute of New Jersey (P30CA072720) as well as Rutgers Office of

Advanced Research Computing (NIH 1S10OD012346- 01A1).

REFERENCES

1.   Eisen MB, Spellman PT, Brown PO, Botstein D: **Cluster analysis and display**

of genome-wide expression patterns. *Proc Natl Acad Sci U S A* 1998,

95(25):14863-14868.

2.   Alizadeh AA, Eisen MB, Davis RE, Ma C, Lossos IS, Rosenwald A, Boldrick JC,

Sabet H, Tran T, Yu X *et al*: **Distinct types of diffuse large B-cell lymphoma**

identified by gene expression profiling. *Nature* 2000, **403**(6769):503-511.

3.   Roth FP, Hughes JD, Estep PW, Church GM: **Finding DNA regulatory motifs**

within unaligned noncoding sequences clustered by whole-genome mRNA

quantitation. *Nat Biotechnol* 1998, **16**(10):939-945.

4.   Golub TR, Slonim DK, Tamayo P, Huard C, Gaasenbeek M, Mesirov JP, Coller

H, Loh ML, Downing JR, Caligiuri MA *et al*: **Molecular classification of cancer:**

class discovery and class prediction by gene expression monitoring.

Science 1999, **286**(5439):531-537.

5.   Perou CM, Jeffrey SS, van de Rijn M, Rees CA, Eisen MB, Ross DT,

Pergamenschikov A, Williams CF, Zhu SX, Lee JC *et al*: **Distinctive gene**

expression patterns in human mammary epithelial cells and breast

cancers. *Proc Natl Acad Sci U S A* 1999, **96**(16):9212-9217.

6.   Cheng Y, Church GM: **Biclustering of expression data**. *Proc Int Conf Intell*

Syst Mol Biol 2000, **8**:93-103.

7.   Madeira SC, Oliveira AL: **Biclustering algorithms for biological data analysis:**

a survey. *IEEE/ACM Trans Comput Biol Bioinform* 2004, **1**(1):24-45.

8.   Eren K, Deveci M, Kucuktunc O, Catalyurek UV: **A comparative analysis of**

biclustering algorithms for gene expression data. *Brief Bioinform* 2013,

14(3):279-292.

9.   Tanay A, Sharan R, Shamir R: **Discovering statistically significant biclusters in gene expression data**. *Bioinformatics* 2002, **18 Suppl 1**:S136-144.

10.   Oghabian A, Kilpinen S, Hautaniemi S, Czeizler E: **Biclustering methods: biological relevance and application in gene expression analysis**. *PLoS One* 2014, **9**(3):e90801.

11.   Yoon S, Nardini C, Benini L, De Micheli G: **Discovering coherent biclusters from gene expression data using zero-suppressed binary decision diagrams**. *IEEE/ACM Trans Comput Biol Bioinform* 2005, **2**(4):339-354.

12.   Getz G, Levine E, Domany E: **Coupled two-way clustering analysis of gene microarray data**. *Proc Natl Acad Sci U S A* 2000, **97**(22):12079-12084.

13.   Sheng Q, Moreau Y, De Moor B: **Biclustering microarray data by Gibbs sampling**. *Bioinformatics* 2003, **19 Suppl 2**:ii196-205.

14.   Hochreiter S, Bodenhofer U, Heusel M, Mayr A, Mitterecker A, Kasim A, Khamiakova T, Van Sanden S, Lin D, Talloen W *et al*: **FABIA: factor analysis for bicluster acquisition**. *Bioinformatics* 2010, **26**(12):1520-1527.

15.   Pontes B, Giraldez R, Aguilar-Ruiz JS: **Biclustering on expression data: A review**. *J Biomed Inform* 2015, **57**:163-180.

16.   Pontes B, Girldez R, Aguilar-Ruiz JS: **Quality measures for gene expression biclusters**. *PLoS One* 2015, **10**(3):e0115497.

17.   van Dam S, Vosa U, van der Graaf A, Franke L, de Magalhaes JP: **Gene co-expression analysis for functional classification and gene-disease predictions**. *Brief Bioinform* 2017.

18.   Bron C, Kerbosch J: **Algorithm 457: finding all cliques of an undirected graph**. *Communications of the ACM* 1973, **16**(9):575-577.

19.   Karp RM: **Reducibility among Combinatorial Problems**. *Proceedings of a symposium on the Complexity of Computer Computations* 1972:85-103.

20.   Cancer Genome Atlas N: **Comprehensive molecular portraits of human**

999   **breast tumours**. *Nature* 2012, **490**(7418):61-70.

1000   21.   Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhim R, Getz G:

1001   **GISTIC2.0 facilitates sensitive and confident localization of the targets of**

1002   **focal somatic copy-number alteration in human cancers**. *Genome Biol* 2011,

1003   **12**(4):R41.

1004   22.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A,

1005   Byrne CJ, Heuer ML, Larsson E *et al*: **The cBio cancer genomics portal: an**

1006   **open platform for exploring multidimensional cancer genomics data**.

1007   *Cancer Discov* 2012, **2**(5):401-404.

1008   23.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y,

1009   Jacobsen A, Sinha R, Larsson E *et al*: **Integrative analysis of complex cancer**

1010   **genomics and clinical profiles using the cBioPortal**. *Sci Signal* 2013,

1011   **6**(269):pl1.

1012   24.   Gyorffy B, Schafer R: **Meta-analysis of gene expression profiles related to**

1013   **relapse-free survival in 1,079 breast cancer patients**. *Breast Cancer Res*

1014   *Treat* 2009, **118**(3):433-441.

1015   25.   van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, Peterse HL, van

1016   der Kooy K, Marton MJ, Witteveen AT *et al*: **Gene expression profiling**

1017   **predicts clinical outcome of breast cancer**. *Nature* 2002, **415**(6871):530-536.

1018   26.   van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, Schreiber

1019   GJ, Peterse JL, Roberts C, Marton MJ *et al*: **A gene-expression signature as a**

1020   **predictor of survival in breast cancer**. *N Engl J Med* 2002, **347**(25):1999-2009.

1021   27.   Schroeder M H-KB, Culhane A, Sotiriou C, Bontempi G, et al.:

1022   **breastCancerNKI: Gene expression dataset. R package version 1.0.6**. 2011.

28.   Hastie T TR, Sherlock G, Eisen M, Brown P, et al. : **Imputing missing data for**

gene expression arrays. *Technical report* 1999.

29.   Anders S, Huber W: **Differential expression analysis for sequence count**

data. *Genome Biol* 2010, **11**(10):R106.

30.   R Core Team: **R: A Language and Environment for Statistical Computing**.

2016.

31.   Gabor Csardi and Tamas Nepusz: **The igraph software package for complex**

network research. *InterJournal* 2006, **Complex Systems**:1695.

32.   Wickham H: **The Split-Apply-Combine Strategy for Data Analysis**. *Journal of*

Statistical Software 2011, **40**(1):1-29.

33.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N,

Schwikowski B, Ideker T: **Cytoscape: a software environment for integrated**

models of biomolecular interaction networks. *Genome Res* 2003,

13(11):2498-2504.

34.   Wickham H: **ggplot2: Elegant Graphics for Data Analysis**. *Springer-Verlag*

New York 2009.

35.   Subhash S, Kanduri C: **GeneSCF: a real-time based functional enrichment**

tool with support for multiple organisms. *BMC Bioinformatics* 2016,

17(1):365.

36.   Benjamini Y, Hochberg Y: **Controlling the false discovery rate: a practical**

and powerful approach to multiple testing. *Journal of the Royal Statistical*

Society 1995, **57**(1):289-300.

37.   Parker JS, Mullins M, Cheang MC, Leung S, Voduc D, Vickery T, Davies S,

Fauron C, He X, Hu Z *et al*: **Supervised risk predictor of breast cancer based**

on intrinsic subtypes. *J Clin Oncol* 2009, **27**(8):1160-1167.

38. Marguerat S, Bahler J: **RNA-seq: from technology to biology**. *Cell Mol Life Sci*

2010, **67**(4):569-579.

39. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B: **Mapping and**

quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* 2008,

5(7):621-628.

40. Kallioniemi A, Kallioniemi OP, Piper J, Tanner M, Stokke T, Chen L, Smith HS,

Pinkel D, Gray JW, Waldman FM: **Detection and mapping of amplified DNA**

sequences in breast cancer by comparative genomic hybridization. *Proc*

Natl Acad Sci U S A 1994, **91**(6):2156-2160.

41. Kao J, Salari K, Bocanegra M, Choi YL, Girard L, Gandhi J, Kwei KA,

Hernandez-Boussard T, Wang P, Gazdar AF *et al*: **Molecular profiling of**

breast cancer cell lines defines relevant tumor models and provides a

resource for cancer gene discovery. *PLoS One* 2009, **4**(7):e6146.

42. Curigliano G, Viale G, Ghioni M, Jungbluth AA, Bagnardi V, Spagnoli GC, Neville

AM, Nole F, Rotmensz N, Goldhirsch A: **Cancer-testis antigen expression in**

triple-negative breast cancer. *Ann Oncol* 2011, **22**(1):98-103.

43. Simpson AJ, Caballero OL, Jungbluth A, Chen YT, Old LJ: **Cancer/testis**

antigens, gametogenesis and cancer. *Nat Rev Cancer* 2005, **5**(8):615-625.

44. Yoshihara K, Shahmoradgoli M, Martinez E, Vegesna R, Kim H, Torres-Garcia

W, Trevino V, Shen H, Laird PW, Levine DA *et al*: **Inferring tumour purity and**

stromal and immune cell admixture from expression data. *Nat Commun*

2013, **4**:2612.

45. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, Speed D,

Lynch AG, Samarajiwa S, Yuan Y *et al*: **The genomic and transcriptomic**

architecture of 2,000 breast tumours reveals novel subgroups. *Nature* 2012,

486(7403):346-352.

46.   Bilal E, Vassallo K, Toppmeyer D, Barnard N, Rye IH, Almendro V, Russnes H,

1075   Borresen-Dale AL, Levine AJ, Bhanot G *et al*: **Amplified loci on chromosomes**

1076   **8 and 17 predict early relapse in ER-positive breast cancers**. *PLoS One*

1077   2012, **7**(6):e38575.

1078   47.   Wang J, Liu Q, Shyr Y: **Dysregulated transcription across diverse cancer**

1079   **types reveals the importance of RNA-binding protein in carcinogenesis**.

1080   *BMC Genomics* 2015, **16 Suppl 7**:S5.

1081   48.   Inic Z, Zegarac M, Inic M, Markovic I, Kozomara Z, Djurisic I, Inic I, Pupic G,

1082   Jancic S: **Difference between Luminal A and Luminal B Subtypes According**

1083   **to Ki-67, Tumor Size, and Progesterone Receptor Negativity Providing**

1084   **Prognostic Information**. *Clin Med Insights Oncol* 2014, **8**:107-111.

1085   49.   Haybittle JL, Blamey RW, Elston CW, Johnson J, Doyle PJ, Campbell FC,

1086   Nicholson RI, Griffiths K: **A prognostic index in primary breast cancer**. *Br J*

1087   *Cancer* 1982, **45**(3):361-366.

1088   50.   Galea MH, Blamey RW, Elston CE, Ellis IO: **The Nottingham Prognostic Index**

1089   **in primary breast cancer**. *Breast Cancer Res Treat* 1992, **22**(3):207-219.

1090   51.   Serin A, Vingron M: **DeBi: Discovering Differentially Expressed Biclusters**

1091   **using a Frequent Itemset Approach**. *Algorithms Mol Biol* 2011, **6**(1):18.

1092   52.   Rosenwald A, Wright G, Chan WC, Connors JM, Campo E, Fisher RI, Gascoyne

1093   RD, Muller-Hermelink HK, Smeland EB, Giltnane JM *et al*: **The use of molecular**

1094   **profiling to predict survival after chemotherapy for diffuse large-B-cell**

1095   **lymphoma**. *N Engl J Med* 2002, **346**(25):1937-1947.

1096   53.   Bergmann S, Ihmels J, Barkai N: **Iterative signature algorithm for the analysis**

1097   **of large-scale gene expression data**. *Phys Rev E Stat Nonlin Soft Matter Phys*

1098   2003, **67**(3 Pt 1):031902.

1099   54.   Ben-Dor A, Chor B, Karp R, Yakhini Z: **Discovering local structure in gene**

1100   **expression data: the order-preserving submatrix problem**. *J Comput Biol*

1101   2003, **10**(3-4):373-384.

1102   55.   Li G, Ma Q, Tang H, Paterson AH, Xu Y: **QUBIC: a qualitative biclustering**

1103   **algorithm for analyses of gene expression data**. *Nucleic Acids Res* 2009,

1104   **37**(15):e101.

1105   56.   Tanay A, Sharan R, Kupiec M, Shamir R: **Revealing modularity and**

1106   **organization in the yeast molecular network by integrated analysis of**

1107   **highly heterogeneous genomewide data**. *Proc Natl Acad Sci U S A* 2004,

1108   **101**(9):2981-2986.

1109   57.   Prelic A, Bleuler S, Zimmermann P, Wille A, Buhlmann P, Gruissem W, Hennig

1110   L, Thiele L, Zitzler E: **A systematic comparison and evaluation of**

1111   **biclustering methods for gene expression data**. *Bioinformatics* 2006,

1112   **22**(9):1122-1129.

1113   58.   Sebastian Kaiser RS, Tatsiana Khamiakova, Martin Sill, Roberto Theron, Luis

1114   Quintales, Friedrich Leisch, Ewoud De Troyer: **biclust: BiCluster Algorithms**.

1115   2018.

1116   59.   Csardi G, Kutalik Z, Bergmann S: **Modular analysis of gene expression data**

1117   **with R**. *Bioinformatics* 2010, **26**(10):1376-1377.

1118   60.   Zhang Y, Xie J, Yang J, Fennell A, Zhang C, Ma Q: **QUBIC: a bioconductor**

1119   **package for qualitative biclustering analysis of gene co-expression data**.

1120   *Bioinformatics* 2017, **33**(3):450-452.

1121   61.   Shamir R, Maron-Katz A, Tanay A, Linhart C, Steinfeld I, Sharan R, Shiloh Y,

1122   Elkon R: **EXPANDER--an integrative program suite for microarray data**

1123   **analysis**. *BMC Bioinformatics* 2005, **6**:232.

1124   62.   Gao C, McDowell IC, Zhao S, Brown CD, Engelhardt BE: **Context Specific and**

1125   **Differential Gene Co-expression Networks via Bayesian Biclustering**. *PLoS*

1126   *Comput Biol* 2016, **12**(7):e1004791.

1127   63.   Mu D, Chen L, Zhang X, See LH, Koch CM, Yen C, Tong JJ, Spiegel L, Nguyen

1128   KC, Servoss A *et al*: **Genomic amplification and oncogenic properties of the**

1129   **KCNK9 potassium channel gene**. *Cancer Cell* 2003, **3**(3):297-302.

1130   64.   Planas-Silva MD, Donaher JL, Weinberg RA: **Functional activity of ectopically**

1131   **expressed estrogen receptor is not sufficient for estrogen-mediated cyclin**

1132   **D1 expression**. *Cancer Res* 1999, **59**(19):4788-4792.

1133   65.   Guiu S, Michiels S, Andre F, Cortes J, Denkert C, Di Leo A, Hennessy BT, Sorlie

1134   T, Sotiriou C, Turner N *et al*: **Molecular subclasses of breast cancer: how do**

1135   **we define them? The IMPAKT 2012 Working Group Statement**. *Ann Oncol*

1136   2012, **23**(12):2997-3006.

1137   66.   Lee TI, Young RA: **Transcriptional regulation and its misregulation in**

1138   **disease**. *Cell* 2013, **152**(6):1237-1251.

1139   67.   Lelli KM, Slattery M, Mann RS: **Disentangling the many layers of eukaryotic**

1140   **transcriptional regulation**. *Annu Rev Genet* 2012, **46**:43-68.

1141   68.   Wan L, Kang Y: **Pleiotropic roles of AEG-1/MTDH/LYRIC in breast cancer**.

1142   *Adv Cancer Res* 2013, **120**:113-134.

1143   69.   Song Z, Wang Y, Li C, Zhang D, Wang X: **Molecular Modification of**

1144   **Metadherin/MTDH Impacts the Sensitivity of Breast Cancer to Doxorubicin**.

1145   *PLoS One* 2015, **10**(5):e0127599.

1146   70.   Shi X, Wang X: **The role of MTDH/AEG-1 in the progression of cancer**. *Int J*

1147   *Clin Exp Med* 2015, **8**(4):4795-4807.

1148

 FIGURE LEGENDS

Fig. 1. Schematic representation of TuBA's proximity measure. (A) For each gene,

samples are arranged in increasing order of expression levels and those corresponding

to a fixed percentile set (top or bottom) are compared between each pair of genes as

shown. The gene-pairs that share a significant number of samples are represented as

nodes linked by edges, which represent the samples. (B) The Venn diagram illustrates

the setup of the contingency table for the one-sided Fisher's exact test. The grey

rectangular box represents the set of all samples in the dataset, the red and blue circles

represent the samples in the top (or bottom) percentile sets of gene 1 and gene 2,

respectively.

Fig. 2. TuBA's schematics. (A) Flowchart of the pipeline for TuBA. (B) Schematic

representation of the graph-based approach to discover biclusters.

Fig. 3. Tuning TuBA's parameters. (A) Significance of overlap corresponding to

fraction of overlap between 0 (no matches/overlap) & 1 (all samples match/overlap) for

percentile set size of: (i) top 20% (green), (ii) top 10% (red), and (iii) top 5% (dark blue)

respectively for a hypothetical dataset consisting of 200 samples. (B) Divergence of the

total number of edges in the graph for the TCGA RFS dataset as we lower the cutoff for

the significance of overlap.

Fig. 4. The effect of TuBA's parameters on the number of genes and samples in

the graph. Plots for the number of genes added to the graph for every incremental

decrease in the significance level for overlap ($-\log_{10}(p)$), the number of samples in graph

at different significance levels of overlap and the total number of edges in graph at

different significance levels of overlap corresponding to a percentile set size of 5% for

(A) METABRIC, (B) TCGA, and (C) GEO datasets, respectively.

Fig. 5. Subtype enrichment of CNV-associated and non-CNV biclusters. Enrichment

of biclusters consisting of proximally located genes with copy number gains in the four

subtypes based on ER/HER2 status for (A) METABRIC and (C) TCGA, respectively. The

biclusters are represented by horizontal bars in each panel, color-coded according to the

chromosome number of their constituent genes. Panels (B) and (D) show the remaining

biclusters arranged according to their serial numbers in Supplementary Table 4 for

METABRIC and TCGA, respectively. The ones that are associated with copy number

(CN) gains of genes located at distant chromosomal sites are shown in red, while the

rest are shown in black. Note, the thickness of the bar in each figure depends on the

total number of biclusters displayed in that figure and so does not represent its

chromosomal extent.

Fig. 6. Clinically pertinent biclusters. Kaplan-Meier survival curves for the set of

patients in the bicluster (red) compared to the remaining set of patients (blue) for

METABRIC and GEO datasets, together with the graphs corresponding to the biclusters

for 8q24.3 (A, B, and C), 8p11.22-p11.23 (D, E, and F) and 17q22-q23.3 (G, H, and I).

Fig. 7. Hierarchical clustering of biclusters-samples. Hamming distance was applied

to the biclusters-samples binary matrix for (A) TCGA and (B) METABRIC RFS datasets,

respectively. The clusters of samples marked by green, brown, and cyan on top in panel

(B) exhibit poor recurrence free survival. The green and brown clusters are associated

with copy number gains at 8q24.3, while the cyan cluster is associated with copy number

gains at 17q25.1-q25.3. Additionally, all three of them were enriched in gene signatures

associated with cellular division and proliferation.

Fig. 8. TuBA compared to other biclustering methods. Proportions of GO-BP terms

enriched biclusters for each biclustering method at five different significance levels for

(A) the DLBCL dataset, (B) the TCGA BRCA dataset. Ratios of number of unique GO-

BP terms and total number of biclusters at five different significance levels for the (C)

DLBCL dataset, (D) the TCGA BRCA dataset.

Fig. 9. Runtime analysis of TuBA. (A) Computation time taken by TuBA to discover

biclusters for different sizes (number of edges) of the graphs based on different choices

of overlap cutoffs for the TCGA (blue), and METABRIC (red) datasets. (B) Dependence

of computation time on the number of rows (genes) for choices of overlap cutoffs

consistent with our suggested heuristic. Here, we chose overlap cutoffs consistent with

our suggested heuristic, and ensured that comparable numbers of edges were

generated for different datasets; this explains why the computation times for datasets

with upwards of 14,000 genes were quite similar to each other. (C) Dependence of

computation time on the number of columns (samples) for choices of overlap cutoffs

consistent with our suggested heuristic.

1210

Figure 2

Figure 4

Hierarchical clustering of biclusters in TCGA   Hierarchical clustering of biclusters in METABRIC

Neurotransmitter secretion and synapse   Mitochondrial translation and respiration + Copy number gains

Copy number gains at 8q24.3   Copy number gains at 17q12 (Her2 Amplicon)   Copy number gains at 8p11.21-p11.23

Cellular division/mitosis + Copy number gains   Response to hormone stimulus, angiogenesis, vasculogenesis

Cell adhesion, extracellular matrix   Immune response   rRNA processing and translation

Mostly copy number gains   No functional association or copy number change