Genome-wide scans of selection highlight the impact of biotic and abiotic constraints in natural populations of the model grass *Brachypodium distachyon*.

Yann Bourgeois[1], Christoph Stritt[2], Jean-Claude Walser[3], Sean P. Gordon[4], John P. Vogel[4], Anne C. Roulin[2*]

[1] New York University Abu Dhabi, PO Box 129188, Saadiyat Island, Abu Dhabi, United Arab Emirates

[2] Institute of Plant and Microbial Biology, University of Zürich, Zollikerstrasse 107, 8008 Zürich, Switzerland

[3] Genetic Diversity Centre, ETH Zürich, Universitätstrasse 16, Zurich, Switzerland

[4] DOE Joint Genome Institute, Walnut Creek, CA 94598, USA.

* Corresponding author: anne.roulin@botinst.uzh.ch

Summary

Grasses are essential plants for ecosystem functioning. Thus, quantifying the selection pressures that act on natural variation in grass species is essential regarding biodiversity maintenance. In this study, we investigated the selection pressures that act on natural populations of the grass model *Brachypodium distachyon* without prior knowledge about the traits under selection. To do so, we took advantage of whole-genome sequencing data produced for two natural populations of *B. distachyon* and used complementary genome-wide scans of selection (GWSS) methods to detect genomic regions under balancing and positive selection. We show that selection is shaping genetic diversity at multiple temporal and spatial scales in this species and affects different genomic regions across the two populations. Gene Ontology annotation of candidate genes reveals that pathogens may constitute important factors of selection in *Brachypodium distachyon*. We eventually cross-validated our results with QTL data available for leaf-rust resistance in this species and demonstrated that, when paired with classical trait mapping, GWSS can help pinpointing candidate genes for further molecular validation. Our study revealed widespread signatures of natural selection on genes involved in adaptation in *B. distachyon* and suggests that pathogens may constitute an important driving force of genetic diversity and evolution in this system. Thanks to a near-base perfect reference genome and the large collection of freely available natural accessions collected across its natural range, *B. distachyon* appears as a prime system for studies in ecology, population genomics and evolutionary biology.

Introduction

Grasses cover more than 40% of the world land area (Gibson, 2009) and dominate a wide variety of ecosystems, from tropical to temperate regions (Clayton, 1981; Gibson, 2009). Grasses also play a key role in eco- and agrosystem functioning as they provide habitats for many animal species (Groves, 2000) and represent the main source of grain and forage (Stromberg, 2011). Increasing crop production to meet the food and energy requirements of the world's growing population is however putting great pressure on natural grasslands (Wallace, 1997; Helm *et al.*, 2009; Ceballos *et al.*, 2010). Faced with constant deterioration and fragmentation due to anthropic activities (Kiviniemi, 2002), these ecosystems are highly endangered (Ceballos *et al.*, 2010), but little is known about their evolutionary resilience. Assessing the genetic basis of adaptation and quantifying the selection pressures that act on natural variation in grass species is therefore crucial with respect to biodiversity maintenance and food security.

To date, reciprocal transplant experiments have been extensively used to test the effect of selection on adaptive differentiation across populations (for review see Savolainen *et al.*, 2013). Based on a "home vs. foreign" effect on fitness, reciprocal transplants are indeed powerful to unravel overall genotype by environment (GxE) interactions and demonstrated the prevalence of local adaptation in grasses and plants in general (for review see Bischoff *et al.*, 2006; Wadgymar *et al.*, 2017). However, reciprocal transplant experiments use proxy such as survival, vegetative growth or seed production to measure the effect of the habitat on fitness (Bischoff *et al.*, 2006). Hence, they provide little information about the functional and genetic bases of adaptation, unless combined with trait mapping such as quantitative trait locus (QTL) analyses and genome-wide association studies (GWAS) (Latta, 2009). QTL analyses and GWAS, on the other hand, are largely constrained by the effort and time necessary for high-resolution mapping. In grasses, while these trait-by-trait approaches have been valuable to decipher the genetic architecture of important characters with regard to crop genetic improvement (Huang *et al.*, 2002; Barbieri *et al.*, 2012; Morris *et al.*, 2013; Slavov *et al.*, 2014), they remain of limited value to grasp the overall selective forces that act on natural populations.

An efficient alternative to provide insights about evolutionary forces in natural populations consists in identifying genes under various types of selection at a whole

genome scale, then describing their function and the type of selection acting on them

(Mitchell-olds *et al.*, 2007). For instance, new mutations that are beneficial in some

populations will be positively selected and are more likely to quickly increase in

frequency. Such so-called selective sweeps tend to reduce genetic diversity, increase

differentiation among populations, and lead to extended haplotypes in the vicinity of

the locus under selection due to genetic hitchhiking (Nielsen, 2005; Hermisson,

2009). Various genome-wide selection scans (GWSS) methods have been developed

to detect such footprints of positive selection in genomes while taking into account

demographic history (Tang *et al.*, 2007; Gautier *et al.*, 2012; Stamatakis *et al.*, 2013;

Messer, 2015), and thanks to the remarkable progress of sequencing technologies,

GWSS are now emerging as complementary approaches to classical trait mapping.

While local adaptation is commonly associated to positive selection on new

advantageous polymorphisms, recent studies have demonstrated that balancing

selection is also playing an important role in this evolutionary process (Mitchell-Olds

et al., 2007; Rasmussen *et al.*, 2014; Wu *et al.*, 2017). The term balancing selection is

an "umbrella" concept (Fijarczyk & Babik, 2015) which describes the maintenance of

genetic diversity over longer periods of time through adaptation to spatial

heterogeneity, heterozygote advantage and negative frequency-dependent selection

(Mitchell-Olds *et al.*, 2007; Rasmussen *et al.*, 2014). Leading to the recycling of

polymorphisms and to selection on standing variation (Richman, 2000; Turchin *et al.*,

2012), this process is more difficult to detect than positive selection (Fijarczyk &

Babik, 2015) since older alleles had more time to recombine and may lead to narrow

signatures around selected sites. As a consequence, the effect of balancing selection is

still largely overlooked in genome scans, which remain strongly biased towards the

detection of recent positive selection (Hassl & Payseur, 2016).

In this study, we capitalize on the near base-perfect quality of the reference genome of

the Mediterranean grass *Brachypodium distachyon* (https://phytozome.jgi.doe.gov) to

investigate how both positive and balancing selection are shaping diversity in this

species. In the last decade, *B. distachyon* has been developed as a powerful model for

research on temperate grass species as it is closely related to major crop cereals and to

some of the grasses used for biofuel production (The international brachypodium

Consortium, 2010). Entirely sequenced, its small diploid genome (272Mb) is fully

assembled into five chromosomes and has been exhaustively annotated (The

international brachypodium Consortium, 2010). In addition, *B. distachyon* is broadly

distributed around the Mediterranean rime (Dell'Acqua *et al.*, 2014; Gordon *et al.*,

2014; Tyler *et al.*, 2016), providing access to natural populations from contrasting

habitats for which a large collection has been collected. It constitutes therefore a

unique and prime system to investigate the genetic basis of local adaptation in natural

grass populations, opening the way to further fundamental and applied research.

Here, we took advantage of whole-genome sequencing data produced for 44

B. distachyon natural accessions originating mainly from Spain and Turkey (Gordon

et al., 2017). We identified over 6 million SNPs and used four complementary GWSS

methods to detect genomic regions under different regimes of selection (Figure 1).

Namely, we asked i) at what time and geographical scale is selection acting in

B. distachyon populations? ii) what are the selective constrains that shape diversity

and adaptation in these populations? iii) whether positive selection is acting on the

same genomic regions in the two populations or, on the opposite, on distinct loci?

Results

Population structure

In this study, we used whole-genome sequencing data (paired-end; Illumina

technology) with a 86-fold median coverage of 44 *B. distachyion* accessions

originating from Turkey, Iraq, Spain and France (Figure 2A, Table S1, (Gordon *et al.*,

2017). After filtering, we identified 6,204,029 SNPs. An ADMIXTURE analysis,

where K=2 was identified as the best model, highlighted two distinct genetic clusters,

an eastern and a western one, with extremely little admixture between the two (Figure

2B). For the rest of the study, accessions from Turkey and Iraq will be referred to as

the eastern population while accessions from Spain and France will be referred to as

the western population. The western population showed a lower level of nucleotide

diversity (Wilcoxon test; P-value < 2.2e-16, Figure 2C) and haplotype diversity (P-

value < 2.2e-16, data not shown) than the eastern one. Excluding the reference

accession Bd21, which has been artificially inbred before sequencing, the average

level of heterozygosity in these accession is of 8% and ranges from 4 to 17.4% (Table

S1).

Functional clustering of the genome of B. distachyon

GWSS outputs provide information about the likelihood for a given locus to be under

selection. A classical approach applied to analyze the results of GWSS consists in

selecting genomic regions containing top 1% outliers for signals of selection and then

assessing whether some biological functions or processes are significantly over-

represented in the gene sets under selection through a Gene Ontology (GO)

annotation (Kelley *et al.*, 2006; Hancock *et al.*, 2011; Nelson *et al.*, 2017). While

recombination rate is relatively high in *B. distachyon* (Huo *et al.*, 2011), signals of

selection around focal loci may decrease slowly due to locally stronger linkage

disequilibrium and subsequent genetic hitchhiking. 1% outlier regions may thus

contain several adjacent genes. Because genes having the same function or being

involved in the same biological process tend to be physically clustered (Hammond-

kosack & Jones, 1996; Michelmore & Meyers, 1998; Takos *et al.*, 2011; Nutzmann &

Osbourn, 2014; Singh *et al.*, 2015), we anticipated that this non-random organization

of genomes could lead to an over-representation of some biological pathways or

functions in small genomic regions and to an artificial enrichment for some GO terms

in GWSS 1% outlier regions (Pavlidis *et al.*, 2012).

To assess whether the genome of *B. distachyon* harbors such functional clusters of

genes, we first performed a GO annotation for the 32,712 genes annotated in the

reference genome. We then controlled for potential gene clustering by following the

procedure described in (Al-Shahrour *et al.*, 2010). Briefly, we split the genome into

overlapping windows of 50 consecutive genes and performed enrichment analyses on

each window. We identified 272 windows significantly enriched for at least one

biological process (Table S2). Several windows were enriched for processes that may

be associated to adaptation to local environmental conditions such response to stress

and defense response (Table S2). This prompted us to narrow down top 1% outlier

regions by keeping only the genes located at and in the immediate vicinity (-10% of

the peak value) of each of the peaks of selection. With the exception of the

coalescence analysis, which is a window-based approach (see methods), all analyses

subsequent to GWSS reported in the following sections were performed on these

filtered outputs.

Genes under balancing selection due to environmental heterogeneity

Variation in abiotic conditions can drive local adaptation at a fine-grained spatial scale and lead to correlations between genotypes and environment. We first used an environmental association analysis approach to detect loci that may have been repeatedly selected by convergent climatic conditions across the two populations. If alleles at these loci have been recycled in front of environmental conditions common to Spain and Turkey, they should display a detectable signal of association at the scale encompassed by our study. Note that geographically varying selection is sometimes considered as being distinct from balancing selection. While it can also be referred to as local adaptation in the literature (Mitchell-olds *et al.*, 2007), we kept it here under the term of balancing selection.

To detect such loci and to highlight alleles shared between the two populations, we selected seven bioclimatic variables showing variation within both the eastern and western populations but little variation between them (Figure 2D). As all these variables were either associated to temperature or to precipitation and are likely to be correlated, we summarized them with a PCA. The first axis of the PCA explained 62.6% of the variance but did not discriminate populations (Figure 2D) and was used to perform an environmental association analysis with all the SNPs identified across the 44 accessions. We identified 26 genomic regions associated with the first PCA axis. These regions harbored 71 genes (Figure 3A). No significant enrichment for any specific biological process was observed (Table 1).

Genes harboring extremely long coalescence time

Balancing selection can be detected with coalescence approaches, as ancient alleles are associated with older coalescence times (Charlesworth, 2006). To detect additional candidate regions for balancing selection in *B. distachyon*, we used the software ARGWeaver (Rasmussen *et al.*, 2014). Briefly, ARGWeaver models the coalescent process along chromosomes and across non-recombining blocks of sequences to address their evolutionary history. It allows recovering several statistics that describe local genealogies and recombination, such as times since the most recent common ancestor (TMRCA), which should be increased near ancient alleles such as those under ancient balancing selection. By doing so, we identified 72 regions harboring 115 genes under what will be referred to as long-term balancing selection in

the following. We observed a significant enrichment for genes involved in
phosphorylation (Table 1).

Genes under disruptive selection between western and eastern populations
As new mutations providing a selective advantage rise in frequency through positive
selection, neutral mutations that are physically close tend to remain strongly linked to
them. Recent positive selection should therefore lead to a signature of long haplotypes
near selected mutations. We used the Rsb test (Tang *et al.*, 2007) to detect such
signatures of recently or almost completed selective sweeps. This test detects
haplotypes that are positively selected in one population by estimating the length of
haplotypes around each allele at a core SNP, then comparing these lengths between
populations. In contrast to the two approaches presented above, the Rsb test should
detect regions that are genetically differentiated across the two populations (Figure 1).
We identified 312 regions harboring 824 genes and 319 regions harboring 1212 genes
in the eastern and western populations respectively (Figure 3A). The selected regions
contained more genes in the western than in the eastern population (Wilcoxon test, P-
value=0.001; Figure 3B). We observed a significant enrichment in genes involved in
response to stress, particularly for defense response and response to oxidative stress in
the eastern population (Table 1, Table S3). The gene set associated to the process of
defense response contained well-known types of resistance genes (R-genes), i.e genes
with NBS-LRR domains (Table S3). Eventually, we observed a significant
enrichment for genes involved in nitrogen transport in the western population (Table
1, Table S3).

Genes under ongoing selection within population
In the case of an ongoing and partial selective sweep (Figure 1), not all individuals
within the population will display haplotype extension in the region under selection,
which can lead to more subtle patterns that are not detected by Rsb test. We used the
program H-scan to detect such incomplete patterns and genes under ongoing positive
selection. We identified 142 regions harboring 487 genes and 79 regions harboring
463 genes in the eastern and western population respectively (Figure 3A). The gene
set under selection in the eastern population was significantly enriched for genes
involved in stress response, including the processes of defense response and response

to cadmium ion (Table 1). The gene set associated to the process of defense response

also contained many R-genes (Table S3). The gene set under selection in the western

population showed a significant enrichment for genes involved in pyruvate metabolic

process (Table 1, Table S3). As in the previous Rsb analysis, the selected windows

contained more genes in the western than in the eastern population (Wilcoxon test, P-

value=0.003; Figure 3B).

GWSS outliers display allele frequency spectra and coalescence patterns consistent

with expectations

To test whether candidate regions displayed an allele frequency spectrum consistent

with the type of selection they were supposed to detect, we computed different

statistics and compared candidate regions to the rest of the genome. We first

computed Tajima's D (Tajima, 1989), which is a measure of genetic diversity

influenced by both selection and demographic variation. Positive values are

associated with balancing selection and bottlenecks, while negative values suggest

recent expansion or positive/purifying selection. We then computed nucleotide

diversity, genome-wide relative (F_{ST}) and absolute (d_{XY}) measures of population

differentiation (Cruickshank & Hahn, 2014). Note that d_{XY} can be interpreted as $F_{ST,}$

but is correlated to the time to coalescence of alleles from all populations, making it

independent of diversity within populations. In addition to TMRCA values, we finally

extracted relative TMRCA halftime (RTH) values from the output of ARGWeaver.

RHT captures coalescence events are skewed toward the recent past but is

independent of the overall coalescence rate (Rasmussen *et al.*, 2014). Assuming

similar selection strength acting on Rsb and H-scan outliers, RTH is thus expected to

be smaller in regions under ongoing positive selection (H-scan) than in regions under

disruptive selection where one allele already reached near-fixation in one of the two

populations (Rsb).

Regions associated with bioclimatic variables or under long-term balancing selection

displayed significantly higher Tajima's D than the genomic background (Wilcoxon

test, all P-values<$1.0.10^{-10}$, Figure 3C) except for associated loci in Spain for which

Tajima's D values were only marginally higher than in the rest of the genome (P-

value=0.055). These regions also display a significantly higher level of nucleotide

diversity within regional groups (P-values<$1.0.10^{-10}$) than the rest of the genome.

Coalescence times (TMRCA) within regional groups were higher for windows
covering loci associated to environmental variables than for genomic background (P-
values=1.10^{-14} and 0.02 for eastern and western populations respectively). As
expected, regions under long-term balancing selection also displayed higher
TMRCAs as they were specifically selected as belonging to the top 1% outlier for this
statistic. Tajima's D, nucleotide diversity and TMRCA statistics are thus consistent
with our expectations and confirmed that both the environmental association and
coalescence analyses detected older polymorphisms shared across the two
populations.

On the other hand, regions identified with the H-scan and Rsb approaches harbored
negative and significantly lower Tajima's D than the rest of the genome, which is
consistent under positive selection (all P-values < 1.10^{-5}, Figure 3C). Outlier windows
for Rsb also displayed higher relative (F_{ST}, P-values<$2.2.10^{-16}$) and absolute levels of
differentiation (d_{XY}; P-values<$2.2.10^{-16}$) than the rest of the genome. These latter
patterns are coherent with the high frequencies of divergent haplotypes between the
western and eastern population for Rsb outliers, resulting in an increased
differentiation between them (Figure 1). H-scan and Rsb outliers also displayed a
lower RTH compared to genomic background (all P-values<$2.2.10^{-16}$ except for
Turkish Rsb outliers for which P-value = 0.002), which is also consistent with
selective sweeps where selection skews coalescence times towards the recent past for
most (but not necessarily all) lineages (Rasmussen *et al.*, 2014). Eventually, H-scan
outliers displayed lower RTH than Rsb outliers (all P-values<$2.2.10^{-16}$), which
confirms that this test tented to detect more recent sweeps than the Rsb.

Overlap between outputs from the four different approaches
As expected given the specificity of each test, we found little overlap between the
outputs of the four different tests (Figure 3A). Out of a total of 1,262 and 1,617 genes
under positive selection, only 49 and 58 genes were common to the H-scan and Rsb
approaches in the eastern and the western population respectively. Similarly, little
overlap (10 genes in total) was observed between the environmental association
analysis and the Rsb or H-scan tests (Figure 3A). No overlap was observed between
the gene sets detected to be under selection with the coalescence approach and with
any the three other tests.

Genomic regions affected by positive selection

We also tested whether distinct loci/genomic regions were affected by recent positive

selection (Rsb and H-scan outliers combined) in the two populations using linear

models. We found no association between the density of genes under selection in the

eastern and western populations on chromosomes 2 and 3 (p-value = 0.4 and 0.3

respectively), which indicates that, as observed on Figure 3D, distinct regions are

affected by positive selection on these two chromosomes. We found a significant

association between these variables on chromosomes 1, 4 and 5 (p-value = 0.004,

1.07e-11 and 0.02 respectively). In these latter cases, however, R^2 were small (7.366e-

05, 0.0004 and 4.42e-05) suggesting that many regions affected by selection remain

specific to each population on these chromosomes as well, as depicted on Figure 3D.

Identification of candidate genes in known QTL regions

Combining association mapping and analyses of selection constitutes a powerful

approach to identify candidate genes and to address their selective regime. We

identified many candidate regions harboring resistance genes. As a proof of concept,

we aimed at assessing whether regions identified as resistance loci against known

pathogens were also highlighted in our scans of selection. In *B. distachyon*, the

genetic basis of resistance to the rust fungus *Puccinia brachypodii* has been

deciphered through a QTL mapping (Barbieri *et al.*, 2012) which showed that leaf-

rust resistance is controlled by three main QTL located on chromosome 2 (from

nucleotide 37,949,269 to 40,903,216), 3 (from nucleotide 13,943,000 to 14,512,222)

and 4 (from nucleotide 9,649,152 to 10,679,750). For the rest of the study, these

regions will be referred to as QTL_{rust-2}, QTL_{rust-3} and QTL_{rust-4}.

We screened these three regions for evidence of selection and found strong Rsb

signals in QTL_{rust-3} and QTL_{rust-4} (Figure 4A). These regions were not detected as

outliers with any other test but belonged to the top 0.05% p-values of Rsb outliers.

The Rsb signal in QTL_{rust-3} reaches its highest point in a serine/threonine phosphatase

(Bradi3g16320: 14,486,831- 14,488,838; Figure 4A, left panel) and 60 kb upstream

the QTL peak identified in this region (Barbieri *et al.*, 2012). The two adjacent Rsb

signals in QTL_{rust-4} reach their highest point into two NBS-LRR resistance genes

(Bradi4g10153: 9,807,879-9,812,927; and Bradi4g10171: 9,828,236-9,835,003;

Figure 3A, right panel). These two genes are respectively located 6 kb and 22 kb

342 upstream the QTL peak. We observed extended haplotypes in the eastern population

343 in both QTL_{rust-3} and QTL_{rust-4} (Figure 4B). Congruent with the signal detected by the

344 Rsb test, the large majority of the eastern accessions displayed the extended haplotype

345 in these regions, indicating a nearly completed selective sweep, especially in QTL_{rust-4}

346 (Figure 4B).

347 QTL_{rust-4} showed a striking enrichment for genes involved in defense response

348 (displayed by grey boxes in Figure 4A, P-value = 8E-09) and in immune signaling

349 process such as phosphorylation (P-value = 5.6E-06). Among 113 genes covered by

350 QTL rust-$_4$, 40 correspond either to a gene with a NBS-LRR, a receptor-like protein

351 kinase (RLK) or a F-box domains, three types of genes that can confer resistance in

352 plants. The presence of such a large gene clusters, as found in other regions of the

353 genome (Table S2), further demonstrates the importance of narrowing down regions

354 under selection to top outlier genes for unbiased GO annotation.

357 **Discussion**

358 Assessing the time and spatial scales at which selection acts is a key to understand

359 how genetic diversity is maintained or lost through adaptation (Stinchcombe &

360 Hoekstra, 2008; Fuller *et al.*, 2015). In plants, and especially in grasses, this question

361 has been largely restricted to crops, which biases our understanding of evolutionary

362 processes that have shaped genomes in their natural ancestors and extant relatives. In

363 this study, we investigate the selective forces influencing adaptation in two

364 populations consisting of 44 natural accessions of the wild Mediterranean grass

365 *B. distachyon.* We found that ancient balancing and recent positive selection left

366 distinct signatures on specific gene categories, and that positive selection affects

367 distinct loci across the two populations. Importantly, our results support a role for

368 pathogens in driving population differentiation and confirm that GWSS constitute

369 effective approaches to pinpoint candidate genes as a complement to classical trait

370 mapping.

372 *Time- and space-varying selection is shaping diversity in B. distachyon*

373 The ending of the last glaciation period 10,000 years ago led to drastic and recent

374 changes of plant communities in Eurasia (Svenning *et al.*, 2008; Binney *et al.*, 2017).

At that time, climate warmed and species distribution expanded over Europe (Hewitt,
1999). Pollen-based studies show that vegetation expansion was fast, reaching up to
2km per year for some species (Hewitt, 1999). To our knowledge, no fossil pollen
records are available for *B. distachyon*, which prevents reconstructing the
geographical distribution of this species before and during the last ice age. Yet, a
previous study showed that the two populations analyzed here experienced a severe
population size reduction during the last glaciation followed by a rapid expansion
within the last 10,000 years (Stritt *et al*., in press). Even though unraveling the history
of populations in southern peninsulas is more complex than in northern regions
(Hewitt, 2000; Feliner, 2011), these results are congruent with the recent global
postglacial recolonization of Europe by plants (Hewitt, 1999; Svenning *et al.*, 2008;
Binney *et al.*, 2017) and imply that *B. distachyon* populations had to adapt to newly
colonized habitats in the recent past.

Balancing selection associated to spatial heterogeneity (Richman, 2000;
Charlesworth, 2006; Fijarczyk & Babik, 2015; Wu *et al.* 2017) may have maintained
ancestral polymorphisms over long periods of time in natural populations of
B. distachyon. Our environmental association analysis indeed shows that loci
associated to bioclimatic variables display higher diversity, older alleles and more
shared variation between regional groups when compared to genomic background.
We exclusively focused the analysis on bioclimatic variables that displayed variation
across localities and not between (Figure 2D). This approach, together with our
relatively small sample size, could explain why this analysis highlighted fewer genes
than the other methods used in the rest of the study. Our observations nonetheless
suggest that adaptation to climate after recolonization does not necessarily involve *de*
novo mutations, even after strong bottlenecks. As a further support of balancing
selection, we identified an even older set of shared polymorphisms with a coalescence
approach. The environmental heterogeneity encountered by *B. distachyon* populations
(Lopez-Alvarez *et al.*, 2015) thus seems to have provided selective pressure strong
enough to maintain polymorphisms over long periods of time within each population.
This result is congruent with a previous analysis of natural populations of
B. distachyon originating exclusively from Turkey (Dell'Acqua *et al.*, 2014) which
showed, at a smaller geographical scale than the one investigated here, that
populations are adapted to local habitats (Dell'Acqua *et al.*, 2014).

On the other hand, we also found evidence for positive selection acting on younger
polymorphisms with both the Rsb and H-scan tests. These tests also revealed that
positive selection is targeting different loci in the two populations. Interestingly, these
loci appear to be non-randomly distributed along chromosomes (Figure 3D). As
recombination rate is relatively high in *B. distachyon* (Huo *et al.*, 2011)*,* we do not
believe that this pattern is due to extended linkage disequilibrium and the subsequent
process of linked selection along such large genomic regions (Cutter & Payseur,
2013; Slotte, 2014). Rather, the peaks of selection we identified were narrow and
allowed to pinpoint genes (Figure 4), indicating that while *B. distachyon* is primarily
inbreeding, outcrossing events must be frequent enough to limit extended linkage
disequilibrium. This is also congruent with the higher level of heterozygosity we
observed here in *B. distachyon* compared to other selfing plants such as *A. thaliana*
(Platt *et al.*, 2010). Whether these regions form islands of divergence and result from
a complex interaction between recombination rate variation, gene flow and selection
(Renaut *et al.*, 2013; Samuk *et al.*, 2017) remains to be investigated. As we obtained
enrichments for different GO terms in the eastern and western populations, our study
nonetheless shows that contrasting abiotic and biotic factors are shaping population
diversity at different regions of the genome of *B. distachyon* through positive
selection.

B. distachyon occurs exclusively in Mediterranean habitats which may appear at a
first glance to be homogeneous and unlikely to promote local adaptation. Our results
defeat this prediction and revealed that natural selection affected different genes and
genomic regions across populations. Even though we identified more genes under
positive than under balancing selection, it would be daring to conclude that the former
selection regime, less challenging to detect (Delph & Kelly, 2014), is a predominant
process shaping diversity in this species. Rather, we believe that we provide here
genomic evidence that large-scale balancing selection also leads to the adaptation of
B. distachyon populations to local environmental conditions.

Pathogens as a potential driving force of population evolution
Host-pathogen interactions lead to a strong coevolutionary dynamics and are
considered as a major factor shaping diversity (Karasov *et al.*; Fumagalli *et al.*, 2011;
Krattinger & Keller, 2016a). Two main types of interaction have been proposed.

Under an arms race model, repeated innovation from both sides results in repeated
fixation of advantageous alleles (Brown & Tellier, 2011). This interaction can
therefore lead to positive selection that can be detected by tests focusing on extended
haplotypes. The other type of interaction is often referred to as Red Queen dynamics
or trench warfare, where alleles involved in the interaction are recycled by negative-
frequency dependence and can therefore subsist for long periods of time in
populations through balancing selection (Brown & Tellier, 2011).

Plant immune system machinery is complex. On one hand, it is composed of two tiers
of extracellular and intracellular receptors (Krattinger & Keller, 2016a,b) that
efficiently detect the presence of pathogens and constitute a first level of defense (for
review Greeff *et al.*, 2012; Couto & Zipfel, 2016; Eckardt, 2017). Further
mechanisms, such as oxidative bursts which are produced at an early stage in case of
pathogen invasion, can act as additional levels of defense and prevent pathogen
proliferation (Wojtaszek, 1997; Torres *et al.*, 2006; Fones & Preston, 2011; Sewelam
et al., 2016). In this study, we found a significant enrichment of signals of selection at
genes involved in these two levels of defense in the eastern population. More
specifically, we found many of well-characterized R-genes, i.e. genes displaying
NBS-LRR domains (Mchale *et al.*, 2006; Jacob *et al.*, 2013; Liu *et al.*, 2014; Couto &
Zipfel, 2016; Eckardt, 2017; Ooijen *et al.*, 2017), and genes involved into oxidative
stress response to be under ongoing and/or disruptive selection in the eastern
population (Table 1). On the other hand, we also found additional R-genes and genes
involved in phosphorylation, a process especially important for immune signaling
response in plants (for review Park *et al.*, 2012), to be under balancing selection. Our
results are thus consistent with the two classical models of host-pathogens
coevolution (Mondragón-palomino *et al.*, 2002; Mchale *et al.*, 2006; Gos *et al.*, 2012;
Mace *et al.*, 2014; Zhong *et al.*, 2015; Wu *et al.*, 2017). Overall, and as shown in
other organisms (Karasov *et al.* 2014; Fumagalli *et al.*, 2011; Krattinger & Keller,
2016a,b; Bourgeois *et al.*, 2017), our genome-wide approach suggests that pathogens
may constitute an important driving force of population and genome evolution in
B. distachyon.

Surprisingly, the significant enrichments for resistance genes in the Rsb and H-scan
outlier gene sets were only observed in the eastern population. The geographical
origin of the populations could play a role in this pattern. The Middle East is indeed

the center of origin of many grasses, including *B. distachyon*, and of their associated
pathogens (Wyand & Brown, 2003; Stukenbrock *et al.*, 2005; Opanowicz *et al.*, 2008;
Hovmøller *et al.*, 2011). Many studies found that both resistance genes in plants and
effector genes in pathogens can be organized in clusters evolving through an arm race
resulting in a gene birth-and-death process (Michelmore & Meyers, 1998; Dong *et al.*,
2015; Singh *et al.*, 2015). Because centers of origin are usually associated to higher
diversity, it is therefore possible that a higher level of pathogen diversity drove
selection at a larger number of resistance genes in the eastern population.

We can, however, not rule out technical biases inherent to the sampling design and to
the history of the studied populations. The western population indeed experienced a
stronger bottleneck than the eastern one (Stritt *et al.*, in press), which, together with
the smaller geographical area sampled, may be responsible for the reduced nucleotide
and haplotype diversity we observed in this population. In addition, the bottleneck
may have led to longer haplotypes in the western population, which could explain
why windows under positive selection were on average larger and contained more
genes in this population than in the eastern one. Discriminating the effect of selection
from the one of demography remains difficult in such bottlenecked populations (Long
et al., 2013). As a consequence, confounding effects may have resulted in the
detection of more false positives and blurred the GO annotation in the western
population. As we applied stringent filtering criteria and used approaches which are
expected to be robust to demographic history (Tang *et al.*, 2007), we nonetheless
believe that we provide the community with a reliable set of candidate loci, even for
the western population. It is worth noting that, despite no significant enrichment for
defense response, footprints of positive selection at R-genes were also found in the
western population. This indicates that while milder, pathogens may constitute a
selection pressure in this population as well.

GWSS as complementary approach to QTL and GWAS
Disentangling the mechanisms that promote or prevent adaptation requires more
integrated studies, using both experimentations in controlled conditions and methods
to characterize genetic diversity in natural populations (Feder & Mitchell-olds, 2003;
Stinchcombe & Hoekstra, 2008; Flood & Hancock, 2017). Several studies used
GWSS to validate genes functionally characterized or previously identified through

GWAS and to propose stronger hypotheses on the mode of selection operating on
traits relevant for adaptation (Roulin *et al.* 2016; Tang *et al.*, 2007; Fumagalli *et al.*,
2011; Bourgeois *et al.* 2017). Following this idea, we inspected three QTL regions
responsible for the resistance of *B. distachyon* to the leaf-rust fungus *Puccinia*
brachypodii, a natural pathogen of *B. distachyon* expected to exert strong selection on
natural accessions.

For two of these QTL regions, we found strong Rsb outliers and reduced haplotype
diversity in the eastern cluster in the vicinity of the QTL peaks, but no sign of
balancing selection. These results strongly indicate that large-scale positive selection
is shaping rust resistance in *B. distachyon* natural populations, as suggested in other
species (Dodds & Thrall, 2009; Chavan *et al.*, 2015). Interestingly, the QTL region
identified on chromosome 3 displays a strong signal of selection in a serine/threonine
phosphatase, a class of genes known for their role in defense response and stress
signaling (País *et al.*, 2009; Durian *et al.*, 2016). The region identified on
chromosome 4 is more complex and consists of a cluster of resistance and stress
signaling genes. Nevertheless, and while other genes display evidence of positive
selection, a strong peak of selection co-localizes with the peak of the QTL and points
at two R-genes. Such genes have been shown to confer resistance to rust in other
species (Bettgenhaeuser *et al.*, 2014) and constitute prime candidates for further
functional characterization.

B. distachyon is closely related to major crop cereals as well as to grass species used
for biofuel production. Translating research from *B. distachyon* to plants of
agronomical and economical interest will require a deeper understanding of the
genetic architecture of traits involved in the response to environmental stresses. The
molecular basis of tolerance to various abiotic stresses such as drought, salt and cold
has been investigated in this species (Luo *et al.*, 2011; Manzaneda, 2013; Carmo &
Charron, 2014; Gordon *et al.*, 2014; Marais & Juenger, 2015; Sun *et al.*, 2015; Mur &
Bosch, 2016). Here, we also highlighted cadmium pollution as a potential factor of
selection in the eastern population. As pollution with heavy metals including
cadmium has been reported in Turkey in regions where accessions were collected for
this study (Bakirdere & Yaman, 2008; Mor & Ceylan, 2017), our results suggest that
B. distachyon could be used to investigate the tolerance to this stress. As genetic
transformation is highly efficient in this species relative to other grasses, we anticipate

that combining classical trait mapping analyses with GWSS will assist allele mining

for additional eco-responsive traits.

Conclusion

Our results revealed widespread signatures of natural selection at genes involved in

adaptation in *B. distachyon*. We also found that pathogens may constitute an

important driving force of genetic diversity and evolution in this system. While we

limited our analysis to classical point mutations, recent studies showed that copy

number variants (CNVs) and transposable element polymorphisms are abundant

across *B. distachyon* populations (Gordon *et al.*, 2017; Stritt *et al.*, in press). Hence,

the important genomic resources currently developed in this species open new

avenues of research to further investigate the role of structural variation in natural

population evolution and adaptation. To date, *B. distachyon* remains a classical model

for research on grass genomics with a strong orientation towards applied research.

Thanks to the high quality of its reference genome and the existence of large

collections of freely available natural accessions collected from the species native

range, it also constitutes a prime system for studies in ecology, population genomics

and evolutionary biology.

Experimental procedures

SNPs calling, population structure and genetic diversity

We used paired-end Illumina sequencing data generated for 44 accessions of

B. distachyon (Gordon *et al*., 2017) originating from Spain (N=16), France (N=1),

Turkey (N=23) and Iraq (N=4, Table S1 for information about the origin of the

accessions and sequencing effort). Reads were aligned to the reference genome v2.0

with BWA-MEM (standard settings; Li, 2013). After removing duplicates with

Sambamba (Tarasov *et al.*, 2017), single nucleotide polymorphisms (SNPs) were

called with Freebayes (Garrison & Marth, 2016). The output was then filtered by

removing SNPs with more than 10 missing genotypes or more than 2 alleles, a quality

lower than 20, a minor allele frequency of 0.05 and a mean depth lower than 20 or

higher than 200. Data were phased using the software BEAGLE V4 (Browning &
Browning, 2007) using default settings.

We then used the program Admixture (Alexander & Novembre, 2009) to identify the
genetic structure of the two populations. The analysis was run for K values from 1 to
5, and the best model was determined as the model with the lowest cross-validation
error. Summary statistics such as within-population nucleotide or haplotype diversity
were computed with the R package PopGenome (Pfeifer *et al.*, 2014). Nucleotide or
haplotype diversity values were square-root transformed to fit a normal distribution
and a t-test was used to compare the two within-population distributions. Levels of
heterozygosity were calculated with VCFtools (Danecek *et al.*, 2011) .

Detecting balancing selection with an environmental association analysis
Bioclimatic variables were downloaded at a 30 arc-seconds resolution from
http://www.worldclim.org/ and extracted for each locality using the R libraries gdal
and raster. We then picked seven bioclimatic variables (bio6, bio8, bio11, bio12,
bio13, bio16, bio18) that displayed substantial variation within geographical groups,
but little between them. Bio6, bio8, bio11, bio12, bio13, bio16 and bio18 correspond
respectively to minimum temperature of coldest month, mean temperature of wettest
quarter, mean temperature of coldest quarter, annual precipitation, precipitation of
wettest month, precipitation of wettest quarter and precipitation of warmest quarter.
As likely to be correlated, these variables were summarized in a principal components
analysis (PCA) in R. The coordinates of the first axis were used to test for correlation
between allele frequencies and environment in the R package GENABEL (Aulchenko
et al., 2007). We accounted for relatedness between samples using a PCA correction
and used corrected P-values of association.

Detecting balancing selection with ancestral recombination graphs (ARG)
We used the software ARGWeaver to detect additional candidate regions for
balancing selection with a coalescence approach (Rasmussen *et al.*, 2014). We
included in the analysis a subset of 12 accessions with high sequencing depth and
covering the largest geographical range (6 accessions from each population) to limit
computation time. We used a mutation rate of 1.4x10-9/bp/generation, a
recombination rate of 5.9x10-8/bp/generation. Mutation rate was estimated by

aligning the orthologs of 100 genes and using rice as an out-group (divergence

estimated at 40My (The international brachypodium Consortium, 2010)). Note that

we subsequently used an outlier approach to identify the oldest polymorphisms

present in the two populations (top 1% outliers). Therefore, potential biases inherent

to the use of a molecular clock do not affect our analysis. ARGWeaver is also flexible

with regard to recombination rate as it reconstructs ancestral recombination graphs

and accommodates variable recombination rates and genealogies along the genome.

The algorithm was run for 1000 iterations, using 20 discretized time steps, a

maximum coalescence time of 3 million generations and a prior effective population

size of 100,000 individuals.

Detecting disruptive positive selection

We used the Rsb test (Tang *et al.*, 2007) to detect signatures of recent or almost

completed hard sweeps. This test detects haplotypes that are positively selected in one

population by using a second population as a contrast. While the output of the test

provides P-value of significance, it also indicates in which population a given allele is

under selection. Rsb statistics were computed for each SNP with the R package

rehh2.0 (Gautier *et al.*, 2012) with default settings. We further visualized the

extension of haplotypes at candidate regions using the bifurcation.diagram() function

of the rehh.

Detecting ongoing positive selection within populations

We eventually used the software H-scan (Schlamp *et al.*, 2016) to detect incomplete

ongoing positive selection. To do so, we calculated average pairwise haplotypes

lengths using the number of segregating sites spanned by each tract within each

population. The statistics is expected to be larger as the number of extended

haplotypes increases in a population. This method is specifically dedicated to the

detection of ongoing sweeps where one (hard sweep) or several (soft sweep)

haplotypes are under positive selection and provides statistics for each SNP. We ran

the method on Turkish and Spanish accessions independently to detect selective

sweeps within each geographical group.

Testing for functional clustering

We first performed a GO annotation for the 32,712 genes annotated in the reference

genome (version 2.1) with Blast2GO (Conesa *et al.*, 2005). We then controlled for

potential gene clustering by following the procedure described in (Al-Shahrour *et al.*,

2010). The entire gene set of the reference genome was split into windows of 50

consecutive genes. Windows were moved along chromosomes in steps of 25 genes to

allow for half-window overlaps. Enrichment analyses of biological processes were

then performed for all the generated windows with the R package GOstats (Falcon &

Gentleman, 2017) using Fisher's exact test. P-values were subsequently adjusted for

multiple testing with a Benjamin-Hochberg correction. Regions were considered

significantly enriched for a biological process when they displayed a corrected P-

value ≤ 0.01 and also harbored at least five genes associated to the given process.

GWWS subsequent filtering

Both the H-scan and the Rsb tests compute statistics at each SNP. To limit false

positives, we first selected 10 kb windows displaying at least four significant SNPs

within the top 1% outliers. Overlapping significant windows were merged. We

narrowed down the selected windows by keeping only the genes located at and around

(-10% of the peak value) each of the top 1% peaks of selection. For the association

test, we also selected 10 kb windows displaying at least four significant SNPs, i.e.

with a corrected P-value ≤ 0.001 ($-\log_{10}$(P-value) ≥ 3) and narrowed them down in the

same manner. These filtering criteria, however, were not applied to the output of

ARGweaver, which is a window-based approach and for which we only kept the top

1% outlier windows.

Overlap between the different approaches

Venn diagrams were drawn with the R package Vennerable to visualize potential

overlap between the different gene sets under selection. To compare the distribution

along chromosomes of candidate genes for recent positive selection (H-scan and Rsb

candidate genes combined) in each population, we used linear models where the

density of selected genes along each chromosome identified in the eastern and

western population (100,000 bins per chromosome) were entered as variables. We

eventually used the function plotBed of the R package Sushi (Phanstiel, 2015) to

vizualize the density of genes under positive selection as a heat map along each
chromosome. The R package ggplot2 (Wickham, 2009) was used to display the
density of all the annotated genes in the genome along each chromosome as a line.
673

Summary statistics and coalescence characterization at candidate loci
Tajima's D, F_{ST} and d_{XY} were computed in the R package PopGenome (v2.2.3) over
5kb windows across the genome. We then compared values between windows
overlapping with candidate regions and windows outside these regions. We extracted
TMRCA and relative TMRCA halftime (RTH) from the output of ARGWeaver.
Because the coalescence approach is window-based, summary statistics were
averaged across windows including 300 non-recombining blocks (around 10kb
windows). Significant windows were merged. We then compared TMRCA and RTH
values between windows overlapping with candidate regions and those outside.

GO annotation of genes under selection
For each test, we extracted the genes located in the filtered regions with bedtools
(Quinlan & Hall, 2010). We then examined potential enrichment for biological
processes for each of the selected gene sets with the R package GOstats (Falcon &
Gentleman, 2017) using the Fisher's exact test. P-values were subsequently adjusted
for multiple testing with a Benjamin-Hochberg correction. Gene sets were considered
significantly enriched for a biological process when they displayed a P-value ≤ 0.01
and harbored at least five genes associated to the given process. The ancestor and
child terms of each significant process were determined using QuickGO
(http://www.ebi.ac.uk/QuickGO) and used to simplify Fisher test outputs and keep
non-redundant terms.
695

QTL for leaf-rust resistance validation
A QTL analysis performed in *B. distachyon* revealed three genomic regions involved
in the resistance to *P. brachypodii* (Barbieri *et al.*, 2012). The coordinates of these
three QTL were extracted from (Barbieri *et al.*, 2012) from v.2.0 of the *B. distachyon*
reference genome. We then assessed weather those three regions were identified as
outliers in at least one of the tests of selection.

Acknowledgements

We thank the Genetic Diversity Center-ETH Zurich for providing access the Euler

high-performance cluster. We also would like to thank Beat Keller's group as well as

Simon Krattinger and Mahendra Mariadassou for their advises during the elaboration

of the study. This work is supported by the Swiss National Science Foundation

(PZ00P3_154724). The work conducted by the U.S. Department of Energy Joint

Genome Institute, a DOE Office of Science User Facility, is supported under Contract

No. DE-AC02-05CH11231.

Availability of data and materials

The raw outputs of each GWWS will be archive upon acceptance of the manuscript.

All whole-genome sequences data are available at the NCBI Sequence Read archive

(SRA available in Gordon *et al*., 2017).

Conflict of interest

There is no conflict of interest issue related to this work.

Supporting Information

Table S1: Geographical coordinates of the 44 accessions and sequencing effort.

Table S2: Significant GO terms in each 50 gene-window of the reference genome

Table S3: Genes under selection with functional annotation

References

Al-Shahrour, Minguez P, Marques-Bonet T, Gazave E, Navarro A, Dopazo J.
2010. Selection upon Genome Architecture: Conservation of Functional
Neighborhoods with Changing Genes. *PLoS computational biology* **6**: e1000953.

Alexander DH, Novembre J. **2009**. Fast Model-Based Estimation of Ancestry in
Unrelated Individuals. *Genome research* **19**: 1655–1664.

Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. **2007**. GenABEL: an R library
for genome-wide association analysis. *Bioinformatics* **23**: 1294–6.

Bakirdere S, Yaman M. **2008**. Determination of lead , cadmium and copper in
roadside soil and plants in Elazig , Turkey. *environmental monitoring Assessment*
136: 401–410.

Barbieri M, Marcel TC, Niks RE, Francia E, Pasquariello M, Mazzamurro V,
Garvin DF. **2012**. QTLs for resistance to the false brome rust Puccinia brachypodii
in the model grass. *Genome* **163**: 152–163.

Bettgenhaeuser J, Gilbert B, Ayliffe M, Moscou MJ. **2014**. Nonhost resistance to
rust pathogens – a continuation of continua. *frontiers in plant science* **5**: 1–15.

Binney H, Edwards M, Macias-fauria M, Lozhkin A, Anderson P, Kaplan JO,
Andreev A, Bezrukova E, Blyakharchuk T, Jankovska V, *et al.* **2017**. Vegetation
of Eurasia from the last glacial maximum to present: Key biogeographic patterns.
Quaternary Science Reviews **157**: 80–97.

Bischoff A, Crémieux L, Smilauerova M, Lawson CS, Mortimer SR, Dolezal J,
Edwards AR, Brook AJ, Macel M, Leps JAN, *et al.* **2006**. Detecting local
adaptation in widespread grassland species – the importance of scale and local plant
community. *journal of ecology* **94**: 1130–1142.

Bourgeois Y, Roulin AC, Müller K, Ebert D. **2017**. Parasitism drives host genome
evolution: Insights from the *Pasteuria ramosa - Daphnia magna* system. *Evolution*
71:1106–13.

Brown JKM, Tellier A. **2011**. Plant-parasite coevolution: bridging the gap between
genetics and ecology. *Annual Review of Phytopathology* **49**: 345–67.

Browning SR, Browning BL. **2007**. Rapid and Accurate Haplotype Phasing and
Missing-Data Inference for Whole-Genome Association Studies By Use of
Localized Haplotype Clustering. *the american journal of human genetics* **81**: 1084–
1097.

Carmo S Do, Charron J. **2014**. Comparative analysis of the cold acclimation and

freezing tolerance capacities of seven diploid Brachypodium distachyon accessions.

Annals of botany **113**: 681–693.

Ceballos, G., Davidson, A., List, R., Pacheco, J., Manzano-Fisher, P., Santos-

Barrera, G., Cruzado J. **2010**. Rapid Decline of a Grassland System and Its

Ecological and Conservation Implications. *PloS one* **5**: e8562.

Charlesworth D. **2006**. Balancing selection and its effects on sequences in nearby

genome regions. *PLoS Genetics* **2**: e64.

Chavan S, Gray J, Smith SM. **2015**. Diversity and evolution of Rp1 rust resistance

genes in four maize lines. *Theoretical and Applied Genetics* **128**: 985–998.

Clayton WD. 1981. Evolution and Distribution of Grasses. *Annals of the Missouri*

Botanical Garden **68**: 5–14.

Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. **2005**.

Blast2GO: a universal tool for annotation, visualization and analysis in functional

genomics research. *Bioinformatics* **21**: 3674–6.

Couto D, Zipfel C. **2016**. Regulation of pattern recognition receptor signalling in

plants. *nature reviews immunology* **16**: 537–552.

Cruickshank TE, Hahn MW. **2014**. Reanalysis suggests that genomic islands of

speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology*

23: 3133–3157.

Cutter AD, Payseur BA. **2013**. Genomic signatures of selection at linked sites :

unifying the disparity among species. *Nature reviews Genetics* **14**: 262–274.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, Depristo MA, Handsaker

RE, Lunter G, Marth GT, Sherry ST, *et al.* 2011. The variant call format and

VCFtools. *Bioinformatics* **27**: 2156–2158.

Dell'Acqua MD, Zuccolo A, Tuna M, Gianfranceschi L, Pè ME. **2014**. Targeting

environmental adaptation in the monocot model Brachypodium distachyon : a

multi-faceted approach. *BMC Genomics* **15**: 801.

Delph LF, Kelly JK. **2014**. On the importance of balancing selection in plants Lynda.

New phytologist **201**: 1–22.

Des Marais DL, Juenger TE. **2015**. Brachypodium and the Abiotic Environment. In:

Vogel J, editor. *Genetics and Genomics of Brachypodium*, 291–311.

Dodds P, Thrall P. **2009**. Recognition events and host–pathogen co-evolution in

gene-for- gene resistance to flax rust. *Functional plant biology* **36**: 395–408.

Dong S, Raffaele S, Kamoun S. **2015**. ScienceDirect The two-speed genomes of

filamentous pathogens□: waltz with plants. *Current Opinion in Genetics &*
Development **35**: 57–65.

Durian G, Rahikainen M, Alegre S, Brosché M. **2016**. Protein Phosphatase 2A in
the Regulatory Network Underlying Biotic Stress Resistance in Plants. *frontiers in*
immunology **7**: 1–17.

Eckardt NA. **2017**. The Plant Cell Reviews Plant Immunity□: Receptor-Like
Kinases, ROS-RLK Crosstalk, Quantitative Resistance, and the Growth / Defense
Trade-Off. *Plant cell* **29**: 601–602.

Falcon SÃ, Gentleman R. **2017**. Using GOstats to test gene lists for GO term
association. *Bioinformatics* **23**: 257–258.

Feder ME, Mitchell-olds T. **2003**. Evolutionary and ecological functional genomics.
Nature reviews Genetics **4**: 649–655.

Feliner GN. **2011**. Southern European glacial refugia□: A tale of tales. *taxon* **60**:
365–372.

Fijarczyk A, Babik A. **2015**. Detecting balancing selection in genomes□: limits and
prospects. *Molecular ecology* **24**: 3529–3545.

Flood J, Hancock AM. **2017**. ScienceDirect The genomic basis of adaptation in
plants. *Current opinion in plant biology* **36**: 88–94.

Fones H, Preston GM. **2011**. Reactive oxygen and oxidative stress tolerance in plant
pathogenic Pseudomonas. *FEMS Microbiology letter* **1**: 1–8.

Fuller ZL, Niño EL, Patch HM, Bedoya-Reina OC, Baumgarten T, Muli E,
Mumoki F, Ratan A, McGraw J, Frazier M, *et al.* **2015**. Genome-wide analysis
of signatures of selection in populations of African honey bees (Apis mellifera)
using new web-based tools. *BMC genomics* **16**: 518.

Fumagalli M, Sironi M, Pozzoli U, Ferrer-admettla A, Pattini L. **2011**. Signatures
of Environmental Genetic Adaptation Pinpoint Pathogens as the Main Selective
Pressure through Human Evolution. *PLoS genetics* **7**: e1002355.

Garrison E, Marth G. **2016**. Haplotype-based variant detection from short-read
sequencing. Available at:https://arxiv.org/abs/1207.3907

Gautier M, Vitalis R, Baillarguet D, Cedex F-M. **2012**. rehh□: an R package to
detect footprints of selection in genome-wide SNP data from haplotype structure.
Bioinformatics **28**: 1176–1177.

Gibson DJ. **2009**. *Grasses and grassland ecology* (Oxford university press, Ed.).

Gordon SP, Contreras-Moreira B, Woods DP, Des Marais DL, Burgess D, Shu S,

Stritt C, Roulin AC, Schackwitz W, Tyler et al. Extensive gene content hetlvariation in the Brachypodium distachyon pan-genome correlates with phenotypic variation. *Nature Communications* **8**: 2184

Gordon SP, Priest H, Marais DL Des, Schackwitz W, Figueroa M, Martin J, Jennifer N, Tyler L, Lee C, Bryant D, *et al.* 2014. Genome diversity in Brachypodium distachyon : deep sequencing of highly diverse inbred lines. Plant Journal **79**: 361–374.

Gos G, Slotte T, Wright SI. 2012. Signatures of balancing selection are maintained at disease resistance loci following mating system evolution and a population bottleneck in the genus Capsella. *BMC Evolutionary Biology* **12**: 152.

Greeff C, Roux M, Mundy J, Petersen M. 2012. Receptor-like kinase complexes in plant innate immunity. *frontiers in plant scienceontiers in plant science* **3**: 1–7.

Groves RH. 2000. Temperate grasslands of the southern hemisphere. In: Jacobs SWL, Everett J, editors. *Grasses Systematics and Evolution*, 3781–3791.

Hammond-kosack KE, Jones JDG. 1996. Resistance Gene-Dependent Plant Defense Responses. *The Plant cell* **8**: 1773–1791.

Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, Toomajian C, Roux F, Bergelson J. 2011. Adaptation to climate across the Arabidopsis thaliana genome. *Science* **334**: 83–6.

Hassl RJ, Payseur BA. 2016. Detecting selection in natural populations: Making sense of genome scans and towards alternative solutions. *Molecular Ecology* **25**: 5–23.

Helm A, Oja T, Saar L, Takkis K, Talve T, Pa M. 2009. Human influence lowers plant genetic diversity in communities with extinction debt. *journal of ecology* **97**: 1329–1336.

Hermisson J. 2009. Who believes in whole-genome scans for selection ? *Heredity* **103**: 283–284.

Hewitt GM. 1999. Post-glacial re-colonization of European biota. *Biological journal of the Linnean society* **68**: 87–112.

Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.

Hovmøller MS, Sørensen CK, Walter S, Justesen AF. 2011. Diversity of Puccinia striiformis on Cereals and Grasses. *Annual Review of Phytopathology* **49**: 197–220.

Huang J, Pray C, Rozelle S. 2002. Enhancing the crops to feed the poor. *Nature* **418**: 678–684.

Huo N, Garvin DF, You FM, Luo SMM, Gu YQ, Lazo GR, Philip J. **2011**. Comparison of a high-density genetic linkage map to genome features in the model grass Brachypodium distachyon. *Theoretical and Applied Genetics* **123**: 455–464.

Jacob F, Vernaldi S, Maekawa T. **2013**. Evolution and conservation of plant NLR functions. *frontiers in immunology* **4**: 1–16.

Karasov TL, Horton MW, Bergelson J. ScienceDirect Genomic variability as a driver of plant – pathogen coevolution□? *Current Opinion in Plant Biology* **18**: 24–30.

Kelley JL, Madeoy J, Calhoun JC, Swanson W, Akey JM. **2006**. Genomic signatures of positive selection in humans and the limits of outlier approaches. *Genome research*: **206**: 980–989.

Kiviniemi K, Eriksson O. 2002. Size-related deterioration of semi-natural grassland fragments in Sweden. *diversity and distribution* **8**: 21–29.

Krattinger SG, Keller B. **2016a**. Tansley review Molecular genetics and evolution of disease resistance in cereals. *New phytologist*: 320–332.

Krattinger SG, Keller B. **2016b**. Trapping the intruder — immune receptor domain fusions provide new molecular leads for improving disease resistance in plants. *Genome Biology* **17**:23.

Latta RG. **2009**. Testing for local adaptation in Avena barbata□: a classic example of ecotypic divergence. *Molecular Ecology* **18**: 3781–3791.

Li H. **2013**. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. preprint arXiv:1303.3997.

Liu W, Frick M, Huel R, Nykiforuk CL, Wang X, Gaudet DA, Eudes F, Conner RL, Kuzyk A, Chen Q, *et al.* **2014**. The Stripe Rust Resistance Gene Yr10 Encodes an Evolutionary-Conserved and Unique CC – NBS – LRR Sequence in Wheat. *Molecular Plant* **7**: 1740–1755.

Long Q, Rabanal FA, Meng D, Huber CD, Farlow A, Platzer A, Zhang Q, Vilhjálmsson BJ, Korte A, Nizhynska V, *et al.* **2013**. Massive genomic variation and strong selection in Arabidopsis thaliana lines from Sweden. *Nature genetics* **45**: 884–890.

Lopez-Alvarez D, Manzaneda AJ, Rey P, Giraldo P, Benavente E, Allainguillaume J *et al.* **2015**. Environmental niche variation and evolutionary diversifi cation of the Brachypodium distachyon grass complex species in their native. *American journal of botany* **102**: 1073–1088.

Luo N, Liu J, Yu X, Jiang Y. **2011**. Natural variation of drought response in
Brachypodium distachyon. *physiologia plantarum* **141**: 19–29.

Mace E, Tai S, Innes D, Godwin I, Hu W, Campbell B, Gilding E, Cruickshank
A, Prentis P, Wang J, *et al.* 2014. The plasticity of NBS resistance genes in
sorghum is driven by multiple evolutionary processes. *BMC plant biology* **14**: 253.

Manzaneda AJ, Rey PJ, Bastida JM, Weiss-Lehman C, Raskin E, Mitchell-Olds
T. **2013**. Environmental aridity is associated with cytotype segregation and
polyploidy occurrence in Brachypodium distachyon (Poaceae) Antonio. *New*
phytologist **193**: 797–805.

Mchale L, Tan X, Koehl P, Michelmore RW. **2006**. Plant NBS-LRR proteins :
adaptable guards. *Genome Biology* **7**: 212.

Messer PW. **2015**. H-scan : Detecting hard and soft sweeps in population genomic
data. **3**: 1–4.

Michelmore RW, Meyers BC. **1998**. Clusters of Resistance Genes in Plants Evolve
by Divergent Selection and a Birth-and-Death Process. *Genome research* **530**:
1113–1130.

Mitchell-Olds T, Willis JH, Goldstein DB. **2007**. Which evolutionary processes
influence natural genetic variation for phenotypic traits? *Nature reviews Genetics* **8**:
845–56.

Mondragón-palomino M, Meyers BC, Michelmore RW, Gaut BS. **2002**. Patterns
of Positive Selection in the Complete NBS-LRR Gene Family of Arabidopsis
thaliana. *Genome research* **12**: 1305–1315.

Mor F, Ceylan S. **2017**. Cadmium and lead contamination in vegetables collected
from industrial , traffic and rural areas in Bursa Province, Turkey. *Food Additives &*
Contaminants **25**: 611–615.

Morris GP, Ramu P, Deshpande SP, Hash CT, Shah T, Upadhyaya HD. **2013**.
Population genomic and genome-wide association studies of agroclimatic traits in
sorghum. *Proceedings of the National Academy of Sciences* **110**: 453–458.

Mur LAJ, Bosch M. **2016**. Linking Dynamic Phenotyping with Metabolite Analysis
to Study Natural Variation in Drought Responses of Brachypodium distachyon.
frontiers in plant science **7**: 1–15.

Nelson RM, Wallberg A, Sim P, Lawson DJ, Webster MT. **2017**. Genomewide
analysis of admixture and adaptation in the Africanized honeybee. *Molecular*
Ecology **26**: 3603–3617.

Nielsen R. **2005**. Molecular Signatures of Natural Selection. Annual Review of
Genetics. **39**: 197–218.

Nutzmann H, Osbourn A. **2014**. Gene clustering in plant specialized metabolism.
current opinion in biotechnology **26**: 91–99.

Ooijen G Van, Mayr G, Kasiem MMA, Albrecht M, Cornelissen BJC, Takken
FLW. **2017**. Structure – function analysis of the NB-ARC domain of plant disease
resistance proteins. *jouranl of experimental botany* **59**: 1383–1397.

Opanowicz M, Vain P, Draper J, Parker D, Doonan JH. **2008**. Brachypodium
distachyon : making hay with a wild grass. *Trends in Plant Science* **13**: 172–177.

País SM, Téllez-iñón MT, Capiati DA. **2009**. Serine / threonine protein
phosphatases type 2A and their roles in stress signaling. *Plant signaling and*
behaviour **4**: 1013–1015.

Park C, Caddell DF, Ronald PC. **2012**. Protein phosphorylation in plant
immunity : insights into the regulation of pattern recognition receptor-mediated
signaling. *frontiers in plant science* **3**: 1–9.

Pavlidis P, Jensen JD, Stephan W, Stamatakis A. **2012**. A Critical Assessment of
Storytelling : Gene Ontology Categories and the Importance of Validating
Genomic Scans Research article. *Molecular biology and evolution* **29**: 3237–3248.

Phanstiel DH. **2015**. Sushi: Tools for visualizing genomics data. *R package version*
1.14.0.

Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ. **2014**. PopGenome:
An efficient swiss army knife for population genomic analyses in R. *Molecular*
Biology and Evolution **31**: 1929–1936.

Platt A, Horton M, Huang YS, Li Y, Anastasio AE, Mulyati NW, Bossdorf O,
Byers D, Donohue K, Dunning M, *et al.* **2010**. The Scale of Population Structure
in Arabidopsis thaliana. *plos genetics* **6**: e1000843.

Quinlan AR, Hall IM. **2010**. BEDTools : a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**: 841–842.

Rasmussen MD, Hubisz MJ, Gronau I, Siepel A. **2014**. Genome-Wide Inference of
Ancestral Recombination Graphs. *PLoS genetics* **10**: e1004342.

Renaut S, Grassa CJ,Yeaman S, Moyers BT, Lai Z, Kane NC, Bowers JE, Burke
JM, Rieseberg LH. **2013**. Genomic islands of divergence are not affected by
geography of speciation in sunflowers. *Nature Communications* **4**: 1–8.

Richman A. **2000**. Evolution of balanced genetic polymorphism. *Molecular Ecology*

9: 1953–1963.

Roulin A, Bourgeois Y, Stiefel U, Walser J-C, Ebert D. A photoreceptor underlies
natural variation in Daphnia magna diapause induction. *Molecular Biology and*
Evolution. **12**:3194–3204.

Samuk K, Rennison DJ, Owens GL, Delmore KE, Miller SE, Schluter D. **2017**.
Gene flow and selection interact to promote adaptive divergence in regions of low
recombination. *Molecular Ecology* **26**: 4378–4390.

Savolainen O, Lascoux M, Merilä J. **2013**. Ecological genomics of local adaptation.
Nature reviews Genetics **14**: 807–820.

Schlamp F, Van Der Made J, Stambler R. **2016**. Evaluating the performance of
selection scans to detect selective sweeps in domestic dogs. *Molecular Ecology* **25**:
342–356.

Sewelam N, Kazan K, Schenk PM. **2016**. Global Plant Stress Signaling : Reactive
Oxygen Species at the. **7**: 1–21.

Singh S, Chand S, Singh NK, Sharma TR. **2015**. Genome-Wide Distribution ,
Organisation and Functional Characterization of Disease Resistance and Defence
Response Genes across Rice Species. *Plos one* **10**:e0125964

Slavov GT, Nipper R, Robson P, Farrar K, Allison GG, Bosch M, Clifton-brown
JC, Donnison IS, Jensen E. **2014**. Genome-wide association studies and prediction
of 17 traits related to phenology, biomass and cell wall composition in the energy
grass Miscanthus sinensis. *New phytologist* **201**: 1227–1239.

Slotte T. **2014**. The impact of linked selection on plant genomic variation. *briefings*
in functional genomics **13**: 268–275.

Stamatakis A, Alachiotis N, Pavlidis P, Daniel Z. **2013**. SweeD : Likelihood-
Based Detection of Selective Sweeps in Thousands of Genomes. *Molecular biology*
and evolution **30**: 2224–2234.

Stinchcombe JR, Hoekstra HE. **2008**. Combining population genomics and
quantitative genetics: finding the genes underlying ecologically important traits.
Heredity **100**: 158–170.

Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC. Transposable elements as a
major source of genetic diversity: insights from genome-wide analyses of
Brachypodium distachyon natural accessions. *Genome biology and evolution*. In
press. Available at https://doi.org/10.1093/gbe/evx276.

Stromberg C. **2011**. Evolution of Grasses and Grassland Ecosystems. *Annual review*

of earth and planetary science **39**: 517–544.

Stukenbrock EH, Banke S, Javan-nikkhah M, Mcdonald BA. **2005**. Origin and

Domestication of the Fungal Wheat Pathogen Mycosphaerella graminicola via

Sympatric Speciation. *Molecular biology and evolution* **24**: 398–411.

Sun J, Hu W, Zhou R, Wang L. **2015**. The Brachypodium distachyon BdWRKY36

gene confers tolerance to drought stress in transgenic tobacco plants. *Plant cell*

reports **34**: 23–35.

Svenning J, Normand S, Kageyama M. **2008**. Glacial refugia of temperate trees in

Europe : insights from species distribution modelling. *journal of ecology* **96**: 1117–

1127.

Tajima F. **1989**. Statistical method for testing the neutral mutation hypothesis by

DNA polymorphism. *Genetics* **123**: 585–95.

Takos AM, Knudsen C, Lai D, Kannangara R, Mikkelsen L, Motawia MS, Olsen

CE, Sato S, Tabata S, Jørgensen K, *et al.* **2011**. Genomic clustering of cyanogenic

glucoside biosynthetic genes aids their identification in Lotus japonicus and

suggests the repeated evolution of this chemical defence pathway. *Plant Journal* **68**:

273–286.

Tang K, Thornton KR, Stoneking M. **2007**. A New Approach for Using Genome

Scans to Detect Recent Positive Selection in the Human Genome. *PLoS biology*

5:e171

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. **2017**. Genome analysis

Sambamba : fast processing of NGS alignment formats. *Bioinformatics* **31**: 2032–

2034.

The international brachypodium Consortium. **2010**. Genome sequencing and

analysis of the model grass Brachypodium distachyon. *Nature* **463**: 763–8.

Torres MA, Jones JDG, Dangl JL. **2006**. Reactive Oxygen Species Signaling in

Response to Pathogens. **141**: 373–378.

Turchin MC, Chiang CWK, Palmer CD, Sankararaman S, Reich D,

Investigation G, Giant T, Hirschhorn JN. **2012**. Evidence of widespread selection

on standing variation in Europe at height-associated SNPs. *Nature Genetics* **44**:

1015–1019.

Tyler AL, Lee SJ, Young ND, Deiulio GA, Benavente E, Reagon M, Sysopha J,

Baldini RM, Troìa A, Hazen SP, *et al.* **2016**. Population structure in the model

grass Brachypodium distachyon is highly correlated with flowering differences

across broad geographic areas. *The Plant Genome* **9**:1–55.

Wadgymar SM, Lowry DB, Gould BA, Byron CN, Mactavish RM, Anderson JT. **2017**. Identifying targets and agents of selection: innovative methods to evaluate the processes that contribute to local adaptation. *Methods in Ecology and Evolution* **8**: 738–749.

Wallace JS, Batchelor CH. **1997**. Managing water resources for crop production. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **352**: 937–947.

Wickham H. **2009**. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wojtaszek P. **1997**. Oxidative burst□: an early plant response to pathogen infection. *biochemical journal* **692**: 681–692.

Wu Q, Han T, Chen X, Chen J, Zou Y, Li Z, Xu Y. **2017**. Long-term balancing selection contributes to adaptation in Arabidopsis and its relatives. *Genome biology* **18**: 217.

Wyand RA, Brown K. **2003**. Genetic and forma specialis diversity in Blumeria graminis of cereals and its implications for host-pathogen co-evolution. *Molecular plant pathology* **4**: 187–198.

Zhong Y, Li Y, Huang K, Cheng Z. **2015**. Species-specific duplications of NBS-encoding genes in Chinese chestnut (Castanea mollissima). *Scientific reports* **5:** 16638.

1065 **Figures**
1066

Figure 1: Schematic representation of the regimes of selection analyzed in this
study

In the case of association between SNPs and environmental variables displaying
variation within population, shared polymorphisms will be maintained across the two
populations. The same outcome is expected under long-term balancing selection. The
Rsb test, on the other hand, detects mutations that rose in frequency and reached near
fixation in one of the two populations. The H-scan test targets mutations currently
under selection and rising in frequency in one of the two populations.

1075
Figure 2: Geographical origin and genetic diversity of the 44 accessions

A) Geographical origin of the 44 accessions used in the study. Blue and orange dots
represent accessions from the western and eastern population respectively. The
asterisk indicates the origin of the reference accession Bd21 B) Structure plot. Each
bar represents the genetic data of one accession. An accession presents some
admixture when the bar displays different colors. The high of the bar is proportional
to the admixture level C) Nucleotide diversity (in 5 kb windows) with the eastern and
western population D) PCA summarizing bioclimatic variables varying within the
eastern and western population.

1085
Figure 3: Venn diagram and descriptive statistics for each test of selection

A) Overlap between environmental association, H-scan and Rsb results in the eastern
and western populations, in number of genes B) Number of genes per window after
keeping the genes located at and in the vicinity of each peak of selection C)
Comparison between genome-wide (GW) Tajima's D and Tajima's D in regions
under the different selection regimes tested D) Overall gene density (line) and heat
maps displaying the density of genes under positive selection (Rsb + H-scan) along
the five chromosomes in the eastern (E) and western (W) populations.

1094

Figure 4: QTL involved in resistance to rust

A) Rsb signals observed in QTL_{rust-3} (left panel) and QTL_{rust-4} (right panel). The
dashed line indicates the 0.1% $-\log_{10}$(Rsb P-values) outlier threshold. Note that in
order to reduce the size of the figure, points with $-\log_{10}$(Rsb P-values) values <1 are
not displayed. Grey boxes display the position of gene potentially involved in defense

response (genes with NBS-LRR, F-box or RLK domains) if present in the region B)

Haplotype bifurcation diagram in sub-regions displaying strong Rsb outliers. Left

panel: zoom in on chromosome 3; right panel: zoom in on chromosome 4. The

diagrams visualize the breakdown of LD at increasing distances from the selected

focal SNP, displayed by a vertical black line. Each horizontal blue line represents a

haplotype. Each accession is represented by two lines, one for each haplotype.

Horizontal blue lines are merged when two accessions share the same haplotype. The

thickness of the line is therefore correlated to the number of accessions sharing the

same haplotype.

1109

1110

1111 **Tables**

1112 **Table 1: Significant GO term associated to each test of selection**

1113

1114

A

B

C

D

A

B

Table 1: significant GO term associated to each test of selection

Test	Significant GO process	Pvalue
Association	None	NA
TMRCA	Phosphorylation	3.13E-08
Rsb East	Response to stress including:	3.00E-05
	- defense response	2.98E-03
	- response to oxidative stress	2.98E-03
RsbWest	Nitrogen compound transport	4.70E-03
H-scan East	Response to stress including:	2.66E-04
	- defense response	7.82E-05
	- response to cadmium ion	2.66E-04
H-scan West	Pyruvate metabolic process	4.00E-04