Characterization of *P. falciparum* dipeptidyl aminopeptidase 3 specificity reveals structural factors responsible for differences in amino acid preferences between peptide-based substrates and covalent inhibitors.

Laura E. de Vries[1,§], Mateo I. Sanchez[2,§], Katarzyna Groborz[3,§], Laurie Kuppens[4,§], Marcin Poreba[3], Christine Lehmann[4], Fang Yuan[5], Shirin Arastu-Kapur[5], Martin Horn[6], Michael Mares[6], Matthew Bogyo[5], Marcin Drag[3], and Edgar Deu[4,§,*] .

[1]Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, Netherlands.

[2]Department of Genetics, Stanford School of Medicine, Stanford, California, United States.

[3]Division of Bioorganic Chemistry, Faculty of Chemistry, Wroclaw University of Technology, Wroclaw, Poland.

[4]Chemical Biology Approaches to Malaria Laboratory, The Francis Crick Institute, London, England, United Kingdom.

[5]Department of Pathology, Stanford University School of Medicine, Stanford, United States.

[6]Institute of Organic Chemistry and Biochemistry, Czech Academy of Sciences, Prague, Czech Republic.

§ These authors contributed equally to this work.

*Corresponding author:

e-mail: edgar.deu@crick.ac.uk

Tel: +44 (0) 20 3796 1412

Abstract

Malarial dipeptidyl aminopeptidases (DPAPs) are cysteine proteases important for parasite development thus making them attractive drug targets. In order to develop inhibitors specific to the parasite enzymes it is necessary to map the determinants of substrate specificity of the parasite enzymes, and its mammalian homologue cathepsin C (CatC). Here, we screened peptide-based libraries of substrates and covalent inhibitors to characterize the differences in specificity between parasite DPAPs and CatC and used this information to develop highly selective DPAP1 and DPAP3 inhibitors. Interestingly, while the primary amino acid specificity of a protease is often used to develop potent inhibitors, we show that equally potent and highly specific inhibitors can be developed based on the sequences of non-optimal peptide substrates. Overall, this study illustrates that focusing the development of protease inhibitors solely on substrate specificity might overlook important structural features that can be exploited to develop highly potent and selective compounds.

Introduction

Malaria is a devastating infectious parasitic disease causing nearly half a million deaths every year(World Health Organization 2017). Malaria is caused by parasites of the *Plasmodium* genus and is transmitted by *Anopheles* mosquitoes during a blood meal. Within the mosquito midgut, parasites reproduce sexually, multiply, and travel to the salivary glands from where they are transmitted to the human host. Upon infection, the parasites first establish an asymptomatic infection in the liver, followed by an exponential asexual replication in the blood stream, through multiple rounds of red blood cell (RBC) invasion, intracellular replication and egress from infected RBCs, this erythrocytic cycle is responsible for the symptoms and pathology of this disease. Over the last 15 years the world has seen a very significant drop in malaria incidence, mainly due to the global distribution of insecticide-impregnated bed nets and the use of artemisinin-based combination therapies as the standard of care for uncomplicated malaria(Bhatt et al. 2015). However, malaria remains a major global health burden with half of the world population at risk and around 200 million clinical cases per year. Unfortunately, mosquitoes are becoming increasingly resistant to insecticides(Ranson & Lissenden 2016) and artemisinin resistance is on the rise(Ashley et al. 2014), thus making the identification of antimalarial targets and the development of drugs with novel mechanism of action extremely urgent(Wells et al. 2015).

Dipeptidyl aminopeptidases (DPAPs) are papain-fold cysteine proteases that are expressed at all stages of parasite development(Bozdech et al. 2003; Young et al. 2005) and might therefore be viable drug targets to treat malaria and prevent its transmission. DPAPs recognize the free N-terminus of protein substrates and cleave N-terminal dipeptides(GUTMANN & FRUTON 1948; WIGGANS et al. 1954). The mammalian homologue cathepsin C (CatC) is the best studied DPAP(Korkmaz et al. 2018). In most cells, CatC plays a catabolic lysosomal function. However, in immune cells it is responsible for

activating various granule serine proteases involved in the immune response and inflammation such as neutrophil elastase, chymase, granzyme A and B, or cathepsin G(McGuire et al. 1993; Kummer et al. 1996; Pham & Ley 1999; Adkison et al. 2002). Because of its role in activating pro-inflammatory proteases, CatC has been pursued as a potential target for chronic inflammatory diseases(Guay et al. 2009; Lainé et al. 2011; Furber et al. 2014), and phase I clinical trials with CatC inhibitors have been performed by GSK (GSK2793660)(Miller et al. 2017) and Astrazeneca (AZD7986)(Palmér et al. 2018), thus proving that DPAPs can be targeted with small drug-like molecules.

Three DPAPs are conserved across *Plasmodium* species but very little is known about their molecular functions. In *P. falciparum*, the most virulent *Plasmodium* specie responsible for 90% of malaria mortality, attempts to directly knockout (KO) DPAP1(Klemba et al. 2004) or DPAP3(Lehmann et al. 2017) have been unsuccessful, suggesting that they are important for parasite replication. Also, in the *P. berghei* murine model of malaria, KO of DPAP1 or DPAP3 results in a significant decrease in parasite replication(Lin et al. 2015; Capuccini et al. 2016; Schwach et al. 2015). DPAP1 localizes mainly in the digestive vacuole(Klemba et al. 2004), an acidic organelle where degradation of haemoglobin takes place. This proteolytic pathway provides a source of amino acids for protein synthesis and liberates space within the RBC for parasites to grow. DPAP1 has been proposed to play an essential role at the bottom of this catabolic pathway(Klemba et al. 2004)·(Deu, Leyva, et al. 2010), however, this function has not yet been confirmed genetically. Previously published inhibition studies suggested that DPAP3 was at the top of the proteolytic cascade that controls parasite egress form iRBCs(Arastu-Kapur et al. 2008). However, our recent conditional KO studies have disproven this hypothesis and shown that DPAP3 is instead critical for efficient RBC invasion(Lehmann et al. 2017). Finally, DPAP2 is only expressed in sexual stages and has been shown to be important for gamete egress from iRBCs, thus making it a potential target to block malaria transmission(Suárez-

Cortés et al. 2016)·(Tanaka et al. 2013). Overall, a pan-DPAP inhibitor will target the parasite at different stages of development, thus potentially slowing down the emergence of resistance.

A clear understanding of the determinants of substrate specificity of *Plasmodium* DPAPs and CatC will be required in order to develop pan-DPAP inhibitors with minimal off target effects on host CatC, and to design highly specific inhibitors to study the biological function of DPAP1 and DPAP3. A general approach to determine the specificity of proteases upstream of the scissile bond (non-prime pockets) is the use of positional scanning substrate libraries where a fluorophore is conjugated to the C-terminus of a peptide library *via* an amide bond. Proteolytic cleavage of this bond results in a significant increase in fluorescence intensity allowing accurate measurement of substrate turnover. The most common libraries used for this purpose are positional scanning synthetic combinatorial libraries (PS-SCL)(Thornberry et al. 1997; Rano et al. 1997; Harris et al. 2000). PS-SCL are composed of multiple sub-libraries designed to determine the specificity of each non-prime binding pocket in a protease. In each sub-library, the amino acid (AA) at a specific position is varied while a stoichiometric mixture of all natural AAs is used in all other positions. PS-SCL thus provide the substrate specificity at each site in the context of all possible combination of AAs at all other positions. Alternatively, the specificity of a given binding pocket can be determined by varying the identity of the AA at that position while fixing the rest of the peptide to residues known to be recognized by the protease of interest. This approach has been used to fingerprint the specificity of amino exopeptidases such as aminopeptidases(Drag et al. 2010; Poreba et al. 2012) or DPAPs(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b), which only recognize one or two AAs upstream of the scissile bond, respectively. PS-SCL have also been applied to protease inhibitor libraries by replacing the fluorophore with a reversible or irreversible warhead(Nazif & Bogyo 2001). Optimum substrates and inhibitors are then designed by combining the best residues in each position. Importantly, the recent

incorporation of non-natural AAs into these libraries has significantly increased the chemical space that can be explored to characterize the specificity of proteases and has allowed the design of substrates and inhibitors with enhanced selectivity over compounds that contain only natural amino acids(Greenbaum, Arnold, et al. 2002; Kasperkiewicz et al. 2014; Kasperkiewicz et al. 2017).

Structure-activity relationship (SAR) studies with positional scanning substrate and inhibitor libraries have been performed both on DPAP1 and CatC(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014a; Wang et al. 2011; Deu, Leyva, et al. 2010), but to a much lesser extent on DPAP3(Arastu-Kapur et al. 2008). Here, we used libraries of peptide-based substrates and inhibitors to determine the specificity of *P. falciparum* DPAP3 at the P1 and P2 positions. Importantly, the libraries used in this study have been previously screened against DPAP1 and CatC and are therefore ideal to compare the specificities of these three proteases(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b). Our studies show that DPAP3 preferentially cleaves after basic and large aromatic residues (P1 position) and that it prefers substrates having N-terminal aliphatic residues (P2 position). We also identified several non-natural P2 residues that are exclusively recognized by either DPAP1 or DPAP3. By combining the SAR information obtained from these substrate and inhibitor screens we developed specific DPAP1 and DPAP3 inhibitors that remain selective in live parasites. Interestingly, while SAR information obtained from positional scanning substrate libraries is often used to develop potent protease inhibitors(Kasperkiewicz et al. 2017), we have identified significant differences in specificity between substrates and inhibitors, particularly in the case of DPAP3 and to a lesser extent for CatC and DPAP1. Surprisingly, we also observed significant discrepancies when we compared previously published specificity data about cysteine cathepsins and caspases obtained from PS-SCL of substrates and inhibitors.

Overall, our study shows that while highly potent inhibitors can be designed based on the sequence of optimal substrates, equally potent and specific inhibitors can be developed using sequences of non-optimal substrates. This work also illustrates how positional scanning substrate libraries might not always identify optimal AAs for inhibitor development.

Results

DPAP3 Substrate Specificity. A positional scanning library of 96 substrates (Figure 1A), composed of a P1 sub-library of 39 substrates (P2 fixed to Met) and a P2 sub-library of 57 substrates (P1 fixed to homophenylalanine (hPhe)), was screened at 1 μM against recombinant DPAP3 (DPAP3; Figure 1B-C). The heat map shown in Figure 1B compares the specificities of DPAP3 with those previously obtained for DPAP1 and CatC(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b) at the same substrate concentration. Note that *D*-Phg is the only *D*-AA in P2 that is cleaved by DPAP3, albeit poorly. The remaining 17 substrates containing *D*-AAs in P2 (*D*-hPhe and all natural *D*-AAs except *D*-Cys and *D*-Met) were not cleaved by DPAP3, nor by DPAP1 or CatC. To simplify the figure, *D*-Phg is the only substrate containing a *D*-AA that is shown in Figure 1.

All three DPAPs show broad and similar P1 specificity, which is not surprising because in clan CA proteases the P1 residue side chain is solvent exposed. For all DPAPs, a general preference for long basic, aliphatic, and aromatic P1 residues was observed—basic: Lys, Arg, homoarginine (hArg), and nitroarginine (Arg (NO$_2$)); aliphatic: Met, norvaline (nVal) and Leu; aromatic: hPhe, (4-benzothiazol-2-yl)homoalanine (hAla (Bht)), 6-benzyloxynorleucine (nLeu(o-Bzl)), glutamic acid benzyl ester (Glu (Bzl)), and homoserine-*O*-benzyl (hSer (Bzl))—. Interestingly, DPAP1 differs from CatC and DPAP3 in that it does not accept large hydrophobic groups such as cyclohexylalanine (Cha), 2-naphthalene (2Nal), biphenylalanine (Bip), 4-benzoyl-phenylalanine (Bpa), tyrosine-*O*-benzoyl (Tyr (Bzl)), or 4-guanidino-

phenylalanine (Phe(guan)) in P1. Unfortunately, we could not identify any P1 residue that was recognized by DPAP1 and/or DPAP3 but not by CatC.

Figure 1. DPAP3 substrate specificity. (A) Structure of P1 and P2 substrate libraries. **(B)** Heat map comparing relative turnover rates for the different DPAPs at 1 μM substrate. For each enzyme and substrate turnover rates were normalized relative to the best natural AA (dark green): Arg in P1 for all DPAPs; Met, Val, and Leu in P2 for CatC, DPAP1, and DPAP3, respectively. Red indicates substrates that are turned over better than the best natural AA. White represent no activity, and grey substrates that were only tested on DPAP3. **(C)** Structure of non-natural AAs used in the substrate library. **(D)** Steady-state Michaelis Menten parameters for DPAP3 determined for selected substrates.

Clear differences in specificity were observed between the three DPAPs at the P2 position (Figure 1B). DPAP3 seems to have a narrower P2 specificity for natural AAs than DPAP1 or CatC, probably reflecting its more specific biological function in RBC invasion compared to the catabolic functions of DPAP1 and CatC. Both DPAP1 and DPAP3 have a strong preference for long aliphatic residues such as Leu, Ile, norleucine (nLeu), Met, or norvaline (nVal). Some non-natural AAs seem to be exclusively cleaved by DPAP3 such as cyclohexylglycine (Chg), Phg, or 4-methyl-phenylalanine (Phe(Me)). Surprisingly, Phe(Me) is the only substrate in the library with an aromatic P2 residue that is accepted by DPAP3 even though vinyl sulfone (VS) inhibitors with P2 aromatic residues such as Tyr, Trp, or nitrotyrosine (Tyr (NO$_2$)) have been shown to be potent DPAP3 inhibitors(Arastu-Kapur et al. 2008; Lehmann et al. 2017). Interestingly, an Ile in P2 is efficiently cleaved by both *Plasmodium* DPAPs but not by CatC.

Development of optimum DPAP3 substrates. To determine how P1 and P2 side chains influence k_{cat} and K_m values for DPAP3, we performed Michaelis Menten studies on selected substrates from the P1 and P2 libraries. In addition, we synthesized a series of substrates that combine optimal natural and non-natural AAs for DPAP3: Arg, hPhe, nLeu(o-Bzl), and Bpa in P1, and Leu, Val, nVal, and Phg in P2. We also tested substrates predicted to be DPAP1-selective (Pro-Arg-ACC and hPro-hPhe-ACC) against DPAP3. Finally, because we were surprised by the lack of activity observed for substrates with aromatic P2 residues, we measured Michaelis Menten parameters for Phe-Arg-ACC, Trp-hPhe-ACC, and Tyr (NO$_2$)-hPhe-ACC. The sequence of the last substrate is based on the structure of SAK1 (Tyr (NO$_2$)-hPhe-VS), which is the most potent DPAP3 inhibitor identified so far (see below). Table 1 and Figure 1D report the Michaelis Menten parameters determined for all these substrates (Michaelis Menten curves are shown in Figure S1).

Table 1. Steady-state Michaelis Menten parameters for DPAP3.

Substrate	N	k_{cat} (s^{-1})	K_m (µM)	$k_{cat}/K_m \cdot 10^{-3}$ (M^{-1}s^{-1})
Met-His-ACC	4	0.23 ± 0.02	42 ± 10	5.4 ± 1
Met-nVal-ACC	4	0.54 ± 0.04	43 ± 5	13 ± 1
Met-hPhe-ACC	4	0.5 ± 0.02	20 ± 2	24 ± 2
Met-Met-ACC	4	0.54 ± 0.02	21 ± 2	26 ± 1
Met-Lys-ACC	4	1.00 ± 0.04	34 ± 2	29 ± 1
Met-Bpa-ACC	3	0.48 ± 0.04	14 ± 3	34 ± 4
Met-hArg-ACC	4	0.92 ± 0.02	27 ± 2	35 ± 1
Met-Arg-ACC	4	1.26 ± 0.06	31 ± 3	41 ± 2
Met-Glu (Bzl)-ACC	4	0.56 ± 0.04	13 ± 2	46 ± 6
Met-Arg (NO$_2$)-ACC	4	1.90 ± 0.08	23 ± 2	84 ± 4
Met-nLeu(o-Bzl)-ACC	10	1.66 ± 0.02	8.8 ± 0.4	186 ± 6
hPro-hPhe-ACC[1]	3	0.007 ± 0.001	26 ± 6	0.30 ± 0.04
Trp-hPhe-ACC	3	0.017 ± 0.002	7 ± 1	2.5 ± 0.3
Phe-hPhe-ACC	3	0.032 ± 0.008	20 ± 8	1.6 ± 0.2
Igl-hPhe-ACC	3	0.044 ± 0.004	4.7 ± 0.9	9.4 ± 1
hAla-hPhe-ACC	3	0.0014 ± 0.002	14.7 ± 0.8	9.8 ± 1
Ile-hPhe-ACC	4	0.12 ± 0.01	12 ± 3	10 ± 1
Phg-hPhe-ACC	4	0.110 ± 0.004	10 ± 1	11 ± 1
Phe(Me)-hPhe-ACC	3	0.072 ± 0.004	6 ± 1	12 ± 1
Tyr (NO$_2$)-hPhe-ACC	3	0.134 ± 0.002	9.9 ± 0.4	13.6 ± 0.4
Gln-hPhe-ACC	3	0.052 ± 0.001	19 ± 10	2.7 ± 0.9
Glu-hPhe-ACC	3	0.024 ± 0.002	1.7 ± 0.5	13 ± 3
Val-hPhe-ACC	3	0.24 ± 0.04	14 ± 4	18 ± 2
nLeu-hPhe-ACC	3	0.28 ± 0.02	5.9 ± 1.3	46 ± 6
nVal-hPhe-ACC	4	0.164 ± 0.005	1.30 ± 0.08	128 ± 6
Leu-hPhe-ACC	3	0.100 ± 0.002	0.50 ± 0.03	210 ± 10
Phg-nLeu(o-Bzl)-ACC	8	0.80 ± 0.02	7.8 ± 0.8	102 ± 6
Leu-nLeu(o-Bzl)-ACC	5	0.308 ± 0.006	0.5 ± 0.1	640 ± 100
nVal-nLeu(o-Bzl)-ACC	8	1.28 ± 0.02	1.4 ± 0.1	920 ± 60
Leu-Bpa-ACC	4	0.170 ± 0.004	0.61 ± 0.06	280 ± 20
Phe-Arg-ACC	4	0.018 ± 0.006	20 ± 10	0.9 ± 0.2
Pro-Arg-AMC[2]	3	0.102 ± 0.004	88 ± 5	1.16 ± 0.04
Val-Arg-ACC[3]	3	1.08 ± 0.06	63 ± 5	17.0 ± 0.4
Leu-Arg-ACC	8	0.330 ± 0.006	1.8 ± 0.2	185 ± 15
nVal-Arg-ACC	8	0.346 ± 0.004	0.72 ± 0.05	480 ± 30
Phe-Arg-BNA	*4*	*0.96 ± 0.04*	*51 ± 5*	*19 ± 1*

[1]For DPAP1, $k_{cat} = 0.22 ± 0.004$ s^{-1}, $K_m = 0.67 ± 0.14$ µM and $k_{cat}/K_m = 320,000 ± 50,000$ M^{-1}s^{-1}; for CatC, $k_{cat} = 10.8 ± 0.2$ s^{-1}, $K_m = 91 ± 5$ µM and $k_{cat}/K_m = 116,000 ± 11,000$ M^{-1}s^{-1}.(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b)

[2]For DPAP1, $k_{cat} = 6.2 ± 0.4$ s^{-1}, $K_m = 84 ± 9$ µM and $k_{cat}/K_m = 74,000 ± 2000$ M^{-1}s^{-1}; for CatC, $k_{cat} = 490 ± 10$ s^{-1}, $K_m = 130 ± 10$ µM and $k_{cat}/K_m = 3,600,000 ± 300,000$ M^{-1}s^{-1}.(Wang et al. 2011)

[3]For DPAP1, $k_{cat} = 3.5 ± 0.1$ s^{-1}, $K_m = 21 ± 2$ µM and $k_{cat}/K_m = 170,000 ± 10,000$ M^{-1}s^{-1}; for CatC, $k_{cat} = 180 ± 10$ s^{-1}, $K_m = 51 ± 8$ µM and $k_{cat}/K_m = 3,600,000 ± 300,000$ M^{-1}s^{-1}.(Wang et al. 2011)

P1 residues have a significant influence in k_{cat}, with nLeu(o-Bzl) and positively charged residues having the highest values. A positive charge on the δ position (Arg (NO_2) and Arg) is favoured over the ε position (Lys and hArg). Elongated aliphatic and hydrophobic residues in P1 decrease K_m, especially when aromatic groups are distant from the peptide backbone. This is evident by the decreasing K_m values between nVal, Met, hPhe, Bpa, Glu (Bzl), and nLeu(o-Bzl). This tendency was also observed for CatC and DPAP1, and might suggest the presence of a distal binding pocket (Figure 1B), potentially an exosite, since P1 residues are usually solvent exposed in clan CA proteases.

P2 residues have a bigger influence on K_m than P1, with Leu and nVal being optimal P2 residues for DPAP3. Beta branched residues are not optimal for DPAP3 as can be observed by an increase in K_m between nVal and Val or nLeu and Ile. However, the γ–branched AA Leu has the lowest K_m. Substrates with aliphatic P2 side chains that extend to the δ position (Met and nLeu) result in higher K_m values than slightly shorter ones (Leu and nVal) but also higher k_{cat} values. Overall, combining optimal P1 (nLeu(o-Bzl) and Arg) and P2 (nVal and Leu) residues results in improved k_{cat}/K_m values (Table 1).

Interestingly, although substrates with Phg and indanyl-glycine (Igl) in P2, or Bpa in P1, are not the preferred AAs at these positions, these non-natural residues are structurally very different from natural AAs and are turned over quite efficiently by DPAP3 when combined with optimal P1 or P2 residues, respectively, i.e. Leu-Bpa-ACC or Phg-nLeu(o-Bzl)-ACC. Finally, the optimal substrate for DPAP1, hPro-hPhe-ACC(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b), is very poorly turned over by DPAP3 (> 200-fold difference in k_{cat}/K_m). We think that substrates containing these non-natural AAs could be used as specific tools to measure DPAP1 or DPAP3 activity in biological samples, i.e. parasite lysates or live parasites, an application we are currently investigating.

Finally, our studies show that substrates with aromatic P2 residues are poorly cleaved by DPAP3 compared to optimal substrates, i.e. 100 to 1000-fold lower k_{cat}/K_m. This is surprising since vinyl sulfone inhibitors containing aromatic P2 residues such as Tyr (NO$_2$) or Trp, are potent and selective DPAP3 inhibitors(Arastu-Kapur et al. 2008; Lehmann et al. 2017). Because these two AA side chains have fluorogenic properties, we investigated whether the low turnover rate measured for Tyr (NO$_2$)-hPhe-ACC and Trp-hPhe-ACC might be due to quenching effects. The emission of free ACC (0, 1, or 5 μM) in assay buffer was measured in the presence of 0-100 μM of these substrates (Figure S2A-B). No significant decrease the ACC emission signal was observed even when substrates were present in 100-fold excess, thus indicating that the low turnover rates measured for these substrates are not due to quenching effects.

As an alternative method to confirm that Tyr (NO$_2$)-hPhe-ACC and Trp-hPhe-ACC bind relatively poorly to DPAP3, we performed substrate competition assays using the (PR)$_2$Rho substrate ($\lambda_{ex} = 492$ nm, $\lambda_{em} = 523$ nm), which emits at much higher wavelengths than ACC ($\lambda_{ex} = 355$ nm, $\lambda_{ex} = 460$ nm), and thus allowing us to simultaneously measure the turnover of (PR)$_2$Rho and ACC substrates without quenching interference. (PR)$_2$Rho was initially designed as a DPAP1 specific substrate to directly measure the activity of this protease in crude parasite extracts(Deu, Yang, et al. 2010), but it is also cleaved by DPAP3 with a $K_{m,app}$ of 40 μM (Figure S2C). This substrate is cleaved twice by DPAPs, releasing two Pro-Arg dipeptides and the rhodamine 110 fluorophore. In this assay, we simultaneously measured inhibition of (PR)$_2$Rho turnover by ACC substrates (IC$_{50}$ values) as well as the apparent K_m values of these ACC substrates in the presence of 40 μM (PR)$_2$Rho ($K_{m,app}$ values). As shown in Figure S2D, the IC$_{50}$ and $K_{m,app}$ values obtained are within experimental error and, as expected, slightly higher than the K_m values reported in Table 1 due to the substrate competition effect. Overall, the lack of quenching effect between Trp or Tyr (NO$_2$) and ACC, and the good agreement

between IC_{50} and $K_{m,app}$ measured under substrate competition conditions indicate that the Michaelis Menten parameters reported in Table 1 are accurate and that substrates containing aromatic P2 AAs are indeed relatively poor DPAP3 substrates compared to those containing optimal aliphatic P2 residues. These results raise the question of why vinyl sulfone inhibitors with aromatic P2 residues are among the most potent DPAP3 inhibitors identified so far. To better understand this discrepancy, we measured the kinetics of inactivation of DPAP3 by the previously published vinyl sulfone inhibitor library(Arastu-Kapur et al. 2008; Deu, Leyva, et al. 2010).

DPAP3 Inhibitor Specificity. Time- and concentration-dependent inactivation of DPAP3 by the P2 library of vinyl sulfone inhibitors (P1 fixed to hPhe) was measured using a continuous assay at 2.2 μM of Met-nLeu(o-Bzl)-ACC ($0.25 \times K_m$). For most compounds, the mechanism of inhibition was consistent with a two-step irreversible inhibition model (Eq. 1, Figure 2A).

$$E + I \overset{K_i}{\rightleftarrows} E{:}I \overset{k_{inact}}{\rightarrow} E-I$$

Eq. 1

K_i is the inhibition equilibrium constant, and k_{inact} the rate of covalent modification of the catalytic Cys.

For a few inhibitors only k_{inact}/K_i values could be obtained, i.e. no saturation was achieved in the k_{obs} *vs.* [I] graph (Figure 2B). The inhibition constants are reported in Figure 2C and Table 2, and the curve fits shown in Figure S3. Only one inhibitor, the one with an amino-methyl-benzyl (Amb) group in P2, was not able to inhibit DPAP3 in a time-dependent manner under our assay conditions (Figure 2D). This is probably due to the fact that this extended and rigid P2 AA (Figure 2E) might prevent proper positioning of the vinyl sulfone group into the active site of DPAP3 to allow covalent modification of the catalytic Cys. For this compound, a K_i value for reversible inhibition was measured.

Figure 2. DPAP3 inhibitor specificity. (A) Representative data showing time-dependent inhibition of DPAP3 by SAK1 (Tyr (NO$_2$)-hPhe-VS). Each progress curve (FU vs. time) was fitted to Eq. 5 to obtain k_{obs} values (top graph). These were then fitted to Eqs. 6 and 7 to obtain k_{inact}, K_i and k_{inact}/K_i. **(B)** Example of an inhibitor where no inhibitor saturation was observed (white bars in **C**). In this case, k_{obs} values were fitted to a linear model to obtain k_{inact}/K_i. **(C)** Inhibition constants determined for DPAP3. The general structure of the inhibitors is shown below the graphs. White bars correspond to inhibitors for which only k_{inact}/K_i values could be determined. The grey bar corresponds to the only inhibitor that showed a reversible mechanism of inhibition, i.e. only a K_i value could be determined. The inhibitor corresponding to SAK1 is indicated with an arrow. **(D)** No time-dependence inhibition of DPAP3 was observed with an inhibitor with a P2 Amb. Initial turnover rates (V$_0$) were fitted to a reversible binding model to obtain K_i. **(E)** Structure of non-natural AAs present in the inhibitor library but not in the substrate library.

Table 2. DPAP3, DPAP1 and CatC inhibition constants for the vinyl sulfone library.

P2	DPAP3			DPAP1			CatC		
	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})
Gly	4.2 ± 0.3	3,300 ± 750	1.30 ± 0.25	3.0 ± 0.2	3,300 ± 750	0.9 ± 0.1	3.0 ± 0.2	70 ± 9	43 ± 4
Ala	2.4 ± 0.1	88 ± 13	27 ± 3	3.03 ± 0.09	90 ± 7	34 ± 2	3.00 ± 0.06	9.7 ± 0.4	310 ± 8
Acpc	3.2 ± 0.1	47,000 ± 3,000	0.0677 ± 0.0001	6.2 ± 0.3	49,000 ± 4,000	0.127 ± 0.006	1.03 ± 0.06	3,200 ± 700	0.33 ± 0.06
hAla	2.2 ± 0.3	13.5 ± 6	166 ± 50	5.2 ± 0.7	19 ± 4	267 ± 26	2.9 ± 0.2	5.9 ± 0.9	500 ± 50
Aib	N.S.	N.S.	0.0363 ± 0.0006	8.9 ± 0.5	18,000 ± 2,000	0.49 ± 0.03	1.20 ± 0.07	2,100 ± 300	0.57 ± 0.06
Val	3.2 ± 0.2	5.7 ± 1.1	554 ± 84	3.6 ± 0.2	2.6 ± 0.4	1,380 ± 140	2.9 ± 0.1	14 ± 2	200 ± 15
Ile	3.4 ± 0.4	23 ± 9	148 ± 45	3.2 ± 0.1	92 ± 8	34 ± 3	3.7 ± 0.8	1,300 ± 400	2.9 ± 0.3
nLeu	3.7 ± 0.2	31 ± 3	120 ± 7.5	4.4 ± 0.3	14 ± 2	305 ± 24	2.40 ± 0.05	7.0 ± 0.5	340 ± 20
Cba	2.5 ± 0.1	140 ± 25	17 ± 2.5	5.4 ± 0.4	13,000 ± 2,000	0.41 ± 0.03	1.29 ± 0.03	2,400 ± 140	0.53 ± 0.02
Thr	2.0 ± 0.1	36 ± 8	55 ± 10	4.4 ± 0.2	112 ± 10	39 ± 2	2.38 ± 0.03	65 ± 3	36 ± 1
Asn	7.4 ± 0.8	2,100 ± 1,000	3.5 ± 1.4	9.9 ± 0.5	10,500 ± 750	0.95 ± 0.02	2.40 ± 0.07	260 ± 30	9.2 ± 0.7
Gln	2.9 ± 0.2	80 ± 20	35 ± 8	4.7 ± 0.2	50 ± 3	93 ± 3	2.3 ± 0.1	15 ± 3	160 ± 25
Asp	2.9 ± 0.2	1,900 ± 400	1.5 ± 0.2	5.4 ± 0.2	2,100 ± 200	2.6 ± 0.10	3.0 ± 0.1	2,200 ± 200	1.4 ± 0.2
Glu	2.7 ± 0.1	51 ± 8	53 ± 6	5.0 ± 0.4	650 ± 100	7.6 ± 0.75	2.6 ± 0.1	140 ± 20	18 ± 2
His	3.7 ± 0.3	100 ± 25	35.5 ± 7.5	4.3 ± 0.3	150 ± 20	29 ± 3	2.1 ± 0.1	0.6 ± 0.1	3,350 ± 500
Lys	N.S.	N.S.	0.020 ± 0.001	N.S.	N.S.	0.081 ± 0.004	1.49 ± 0.05	3,600 ± 200	0.42 ± 0.02
Arg	N.S.	N.S.	0.065 ± 0.006	N.S.	N.S.	0.084 ± 0.002	4.0 ± 0.5	11,000 ± 2,000	0.38 ± 0.02
Phg	N.S.	N.S.	0.0357 ± 0.0007	8.3 ± 0.5	7,000 ± 650	1.19 ± 0.05	2.5 ± 0.2	86 ± 20	29 ± 5
2fa	3.8 ± 0.3	39 ± 10	97 ± 18	5.2 ± 0.4	11 ± 3	470 ± 90	3.7 ± 0.2	2.1 ± 0.2	1,720 ± 90
2ta	3.4 ± 0.3	29 ± 7	118 ± 23	3.8 ± 0.3	82 ± 20	47 ± 7	3.3 ± 0.3	6.5 ± 1	510 ± 55
Cha	2.7 ± 0.1	90 ± 20	31.6 ± 5.5	N.S.	N.S.	0.020 ± 0.001	2.3 ± 0.1	300 ± 40	7.7 ± 0.8
Phe	2.6 ± 0.2	45 ± 13	58 ± 13	N.S.	N.S.	0.40 ± 0.01	2.18 ± 0.08	1.4 ± 0.2	1,600 ± 200
Phe(Me)	5.4 ± 0.4	9,200 ± 300	0.580 ± 0.025	N.S.	N.S.	0.314 ± 0.005	2.8 ± 0.2	1,300 ± 200	2.2 ± 0.3
Phe(I)	3.8 ± 0.3	110 ± 30	36 ± 10	7.0 ± 0.3	47,000 ± 3,500	0.151 ± 0.006	1.02 ± 0.06	185 ± 40	5.5 ± 0.9
Phe(NH2)	3.1 ± 0.2	39 ± 7	80 ± 12	5.0 ± 0.4	22,000 ± 2500	0.23 ± 0.01	1.01 ± 0.02	5.3 ± 0.3	191 ± 9
Tyr	2.18 ± 0.07	12.7 ± 1.5	172 ± 16	N.S.	N.S.	0.310 ± 0.005	2.10 ± 0.06	11.5 ± 0.9	182 ± 10
Tyr (NO$_2$)	7.2 ± 0.9	250 ± 40	28.8 ± 0.8	N.S.	N.S.	0.023 ± 0.002	N.S.	N.S.	0.147 ± 0.008
hPhe	2.36 ± 0.08	310 ± 30	7.6 ± 0.7	N.S.	N.S.	0.029 ± 0.002	N.S.	N.S.	0.103 ± 0.005
Trp	2.5 ± 0.1	30 ± 6	84 ± 14	N.S.	N.S.	0.074 ± 0.002	1.98 ± 0.03	61 ± 4	32 ± 2
Igl	2.2 ± 0.1	29 ± 5	78 ± 12	N.S.	N.S.	0.313 ± 0.007	2.3 ± 0.1	240 ± 40	10 ± 1
2Nal	3.7 ± 0.3	7,100 ± 1,000	0.510 ± 0.04	5.3 ± 1.3	15,200 ± 6,500	0.35 ± 0.07	2.6 ± 0.2	1,800 ± 400	1.5 ± 0.2
Bip	N.S.	N.S.	0.0314 ± 0.0004	N.S.	N.S.	0.0178 ± 0.0004	2.53 ± 0.09	6,200 ± 800	0.41 ± 0.04
Inp	5.2 ± 0.4	65,000 ± 800	0.079 ± 0.004	N.S.	N.S.	0.03 ± 0.001	1.05 ± 0.06	5,600 ± 900	0.19 ± 0.02
Amc	5.7 ± 0.3	75,000 ± 7,000	0.076 ± 0.003	5.8 ± 0.3	27,000 ± 2500	0.213 ± 0.009	0.97 ± 0.04	1000 ± 200	0.94 ± 0.17
3Abz	3.4 ± 0.4	3,100 ± 600	1.1 ± 0.1	4.8 ± 0.3	3000 ± 500	1.6 ± 0.2	1.08 ± 0.07	55 ± 12	20 ± 4
Amb	N/A	3,200 ± 1000	N/A	N.S.	N.S.	0.140 ± 0.007	1.6 ± 0.1	2100 ± 300	0.74 ± 0.06

N.S. No saturation, only k_{inact}/K_i could be obtained. N/A, no time dependent inactivation was observed, data consistent with reversible inhibition.

Overall, changes in P2 do not have a big influence in k_{inact} with the exceptions of Asn, Phe(Me), and Tyr (NO_2), which significantly increase k_{inact}. Intriguingly, substrates containing the latter two P2 residues were the only substrates with a P2 aromatic residue that could be cleaved by DPAP3 with $k_{cat}/K_m > 2{,}000$ $M^{-1}s^{-1}$ (Table 1). In terms of K_i, we observed some SAR similarities between substrates and inhibitors: DPAP3 does not bind inhibitors with an N-terminal basic residue (Arg or Lys), but it is strongly inhibited ($K_i < 25$ nM) by aliphatic residues (Leu, nLeu, hAla, or Val). However, we measured potent inhibition of DPAP3 with aromatic residues in P2 such as Phe, Tyr or Trp ($k_{inact}/K_i \geq 60{,}000$ $M^{-1}s^{-1}$). Substrates with these P2 residues show relatively poor substrate turnover ($k_{cat}/K_m \leq 2{,}000$ $M^{-1}s^{-1}$). We also observed other discrepancies between substrates and inhibitors. For example, Thr in P2 results in a poor substrate (Figure 1B) but a relatively potent inhibitor ($k_{inact}/K_i = 29{,}000$ $M^{-1}s^{-1}$). Inversely, DPAP3 cleaves Phg-hPhe-ACC and Phe(Me)-hPhe-ACC with a k_{cat}/K_m of 11,000 and 12,000 $M^{-1}s^{-1}$, respectively, but the k_{inact}/K_i for the respective inhibitors are only 36 and 420 $M^{-1}s^{-1}$. As shown in Figure 3A, there is not a clear correlation between k_{cat}/K_m and k_{inact}/K_i for DPAP3 as a function of the P2 residue, however, P2 residues that make optimal substrates also make good inhibitors (Val, nVal, nLeu).

Correlation between substrate turnover and inhibition for DPAP1 and CatC. To determine whether the lack of correlation between k_{cat}/K_m and k_{inact}/K_i observed for DPAP3 is a common feature in DPAPs, we calculated apparent k_{cat}/K_m values for DPAP1 for the P2 substrate library based on the activity measurements previously reported at 1 μM and the k_{cat}/K_m for hPro-hPhe-ACC(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014a). We also measured accurate Michaelis Menten parameters for CatC for the P2 substrate library (Table S1 and Figure S4), and k_{inact} and K_i values for DPAP1 and CatC for the P2 vinyl sulfone library (Table 2 and Figure S3). As shown in Figure 3B, we observed a good

correlation between substrate turnover and k_{inact}/K_i for DPAP1, but for CatC we observed some

discrepancies (Figure 3C), albeit not as pronounced as in the case of DPAP3. For example, the

CatC k_{cat}/K_m for Phe(Me)-hPhe-ACC is 30-fold higher than for Phg-hPhe-ACC while the k_{inact}/K_i

for Phe(Me)-hPhe-VS is 13-fold lower than for Phg-hPhe-VS, thus resulting in a 400-fold

discrepancy in the changes in k_{cat}/K_m and k_{inact}/K_i values between these two P2 residues.

Figure 3. Comparison of catalytic efficiency and second order inhibition constants. (A-C) Comparison of k_{inact}/K_i and k_{cat}/K_m for DPAP3 (**A**), DPAP1 (**B**) and CatC (**C**) between vinyl sulfone inhibitors and P2 substrate library. For DPAP1, apparent k_{cat}/K_m ($k_{cat}/K_{m,app}$) were calculated based on previously reported turnover rates at 1 μM(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b). $k_{cat}/K_{m,app}$ were calculated similarly for DPAP3 and CatC for P2 substrates whose activity was too low to obtain accurate Michaelis Menten parameters (i.e., substrates not present in Tables 1 and S2). Filled circles correspond to compounds belonging to the vinyl sulfone library (compounds in Table 2), and empty circles to inhibitors having a phenyl group in P1' (Table 3). The P2 residue is labelled next to each data point. Pearson correlation coefficients (ρ) are shown for each protease. (**D-F**) Comparison of changes in substrate turnover relative to inhibitors potency for any pair of P2 residues. The log value of ΔACT/ΔINH (Eq. 2) calculated for DPAP3 (**D**), DPAP1 (**E**) and CatC (**F**) are shown has a heat map with values above and below zero in red and blue, respectively. Each pairwise value showing more than a 100-fold discrepancy between activity and inhibition (ΔACT/ΔINH > 100 or < 0.01) is highlighted in bold.

A systematic way to visualize discrepancies between substrate turnover and inhibition is

to compare the fold different in k_{cat}/K_m with that in k_{inact}/K_i for any two P2 residues (Eq. 2).

$$\frac{\Delta \text{ACT}}{\Delta \text{INH}} = \frac{(k_{\text{cat}}/K_{\text{m}})_{\text{AA1}}/(k_{\text{cat}}/K_{\text{m}})_{\text{AA2}}}{(k_{\text{inact}}/K_{\text{i}})_{\text{AA1}}/(k_{\text{inact}}/K_{\text{i}})_{\text{AA2}}} \qquad \text{Eq. 2}$$

We performed these pairwise calculations for each of the DPAPs studied here and have presented the results as heat maps in Figure 3D-F. We observed significant and numerous discrepancies between substrates and inhibitors for DPAP3 (30 % of pairwise $\Delta \text{ACT}/\Delta \text{INH} > 100$ or < 0.01), almost no discrepancies for DPAP1 (only 1% $\Delta \text{ACT}/\Delta \text{INH} > 10^2$ or $< 10^{-2}$), and only a few for CatC (4% of $\Delta \text{ACT}/\Delta \text{INH} > 100$ or < 0.01). Overall, this study indicates that the level of correlation between $k_{\text{cat}}/K_{\text{m}}$ and $k_{\text{inact}}/K_{\text{i}}$ is protease dependent.

Development of DPAP1 and DPAP3-selective inhibitors. We next synthesized several inhibitors to determine whether the optimal nLeu(o-Bzl) P1 residue identified from the substrate screen could be used to increase the potency and specificity of inhibitors towards DPAP1 or DPAP3. We selected P2 AAs that were predicted to provide specificity towards DPAP1 (Pro and hPro) or DPAP3 (aromatic residues: Tyr (NO$_2$), Trp, Igl and 2Nal) based on our substrate and inhibitor screening results. We also included in our analysis the previously synthesized compound JCP410 (nVal-hPhe-VS)(Arastu-Kapur et al. 2008), since nVal is one of the best P2 residues identified from the substrate screen. We determined the inhibition constants of these compounds for DPAP1, DPAP3, and CatC (Table 3 and Figures 2 and S3). The major structural difference between these compounds and the inhibitor library is that they have a phenyl group in P' instead of a long aliphatic linker (Figure 2A). This change usually increases the potency of compounds except in the context of a P2 Trp for DPAP3 or P2 Tyr (NO$_2$)/2Nal for CatC (Tables 2 and 3). These exceptions indicate some level of interdependence between the prime and non-prime sites of DPAPs.

Table 3. Inhibition constants for optimal vinyl sulfone inhibitors

Inhibitor Name	DPAP3			DPAP1			CatC			Specificity Ratio[a]		
	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})	$k_{inact} \cdot 10^{-3}$ (s^{-1})	K_i (nM)	$k_{inact}/K_i \cdot 10^{-3}$ (M^{-1}s^{-1})	DPAP1/ DPAP3	DPAP1/ CatC	DPAP3/ CatC
Pro-hPhe-VS	3.5 ± 0.2	860 ± 100	4.1 ± 0.3	2.1 ± 0.2	4.6 ± 0.7	470 ± 35	1.45 ± 0.07	34 ± 5	42 ± 5	**115**	11	0.10
hPro-hPhe-VS	2.1 ± 0.1	30 ± 7	69 ± 13	4.8 ± 0.6	3.9 ± 1.5	1,230 ± 400	1.67 ± 0.08	22 ± 3	76 ± 0.8	18	16	0.91
hPro-nLeu(o-Bzl)-VS	2.0 ± 0.1	39 ± 6	53 ± 6.5	2.9 ± 0.3	0.8 ± 0.2	3,560 ± 150	1.67 ± 0.03	11.2 ± 0.6	150 ± 6	68	24	0.35
Tyr (NO$_2$)-hPhe-VS	2.26 ± 0.05	2.4 ± 0.2	950 ± 74	3.8 ± 1	25,000 ± 1,400	0.15 ± 0.05	3.2 ± 0.6	58,000 ± 20,000	0.054 ± 0.008	**0.00016**	2.8	**17,600**
Tyr (NO$_2$)-nLeu(o-Bzl)-VS	2.7 ± 0.2	53 ± 10	51 ± 10	2.6 ± 0.3	2,400 ± 900	1.1 ± 0.3	2.9 ± 0.2	7,700 ± 1,400	0.38 ± 0.05	0.032	2.9	89
2Nal-hPhe-VS	2.59 ± 0.07	254 ± 16	10.2 ± 0.5	6.7 ± 0.2	33,000 ± 2,000	0.2 ± 0.005	3.7 ± 0.2	4,000 ± 700	0.9 ± 0.1	0.02	0.22	11
2Nal-nLeu(o-Bzl)-VS	6.0 ± 0.7	2,800 ± 600	2.2 ± 0.3	5.0 ± 1.7	27,000 ± 16,000	0.19 ± 0.05	2.4 ± 0.6	1,740 ± 140	1.37 ± 0.09	0.14	0.14	1.0
Igl-hPhe-VS	2.1 ± 0.7	10 ± 1	205 ± 22	2.7 ± 0.1	5,400 ± 500	0.50 ± 0.03	5.8 ± 0.3	320 ± 70	18 ± 3	**0.0024**	0.028	11.4
Igl-nLeu(o-Bzl)-VS	5.6 ± 0.4	160 ± 30	35 ± 4.5	4.1 ± 0.2	4,800 ± 500	0.86 ± 0.05	1.48 ± 0.06	11 ± 2	135 ± 14	0.037	**0.0064**	0.17
L-Trp-hPG-VS	3 ± 1	330 ± 150	9.1 ± 1.5	2.1 ± 0.2	2,800 ± 450	0.74 ± 0.07	1.73 ± 0.03	28 ± 2	61 ± 4	0.12	0.012	0.10
L-Trp-hPhe-VS	3.3 ± 0.4	65 ± 15	52 ± 7	1.7 ± 0.1	540 ± 80	3.20 ± 0.3	2.2 ± 0.09	16 ± 3	131 ± 15	0.094	0.024	0.26
L-Trp-nLeu(o-Bzl)-VS	3.2 ± 0.3	4 ± 1	826 ± 193	2.2 ± 0.1	380 ± 60	5.8 ± 0.6	1.74 ± 0.03	4.9 ± 0.3	355 ± 18	**0.0070**	0.016	2.3
D-Trp-hPhe-VS	1.7 ± 0.6	5,500 ± 3,000	0.32 ± 0.08	2.7 ± 0.2	105,000 ± 10,000	0.026 ± 0.001	2.6 ± 0.1	4,200 ± 600	0.62 ± 0.07	0.12	0.042	0.34
D-Trp-nLeu(o-Bzl)-VS	3.6 ± 0.2	7,600 ± 600	0.47 ± 0.01	5.5 ± 0.7	127,000 ± 20,000	0.044 ± 0.002	1.97 ± 0.06	1,100 ± 100	1.7 ± 0.1	0.094	0.026	0.28
nVal-hPhe-VS	2.6 ± 0.4	2.8 ± 0.9	940 ± 160	2.0 ± 0.1	0.6 ± 0.2	3,160 ± 750	2.2 ± 0.1	0.38 ± 0.08	5,830 ± 970	3.36	0.54	0.16

[a]Specificity ratio was calculated based on k_{inact}/K_i values; numbers highlighted in bold indicate more than 100-fold selectivity for the indicated enzymes.

Compared to hPhe, P1 nLeu(o-Bzl) decreases k_{inact}/K_i value for DPAP3 by 7 to 10-fold except in the context of a P2 Trp where it increases it by 24-fold (Table 3). These differences are mainly due to changes in K_i rather than k_{inact} and likely reflect cooperativity between the S1 and S2 pockets of DPAP3. However, in the case of DPAP1 and CatC, replacement of P1 hPhe with nLeu(o-Bzl) decreases K_i. Because of this P1-P2 interdependence, the most potent inhibitors for DPAP3 are either a combination of P2 Tyr (NO$_2$) and P1 hPhe, i.e. SAK1, or P2 Trp and P1 nLeu(o-Bzl), resulting in k_{inact}/K_i values close to 10^6 M^{-1}s^{-1}. Importantly, nVal-hPhe-VS is as potent as these two inhibitors (k_{inact}/K_i = 940,000 M^{-1}s^{-1}), confirming that optimal inhibitors can be designed based on the structure of optimal substrates. These high second order rate constants are mainly driven by low K_i values (< 5 nM). While Trp-nLeu(o-Bzl), Igl-hPhe-VS and Tyr (NO$_2$)-hPhe-VS show more than 100-fold selectivity for DPAP3 over DPAP1, only the latter is selective for DPAP3 compared to CatC (Table 3). On the other hand, nVal-hPhe-VS is equally potent for all DPAPs (k_{inact}/K_i = 1·10^6, 3.2·10^6 and 5.8·10^6 M^{-1}s^{-1} for DPAP3, DPAP1 and CatC, respectively) making it a highly potent but non-selective pan-DPAP inhibitor

Pro-hPhe-VS (SAK2) is 100-fold more selective towards DPAP1 than DPAP3, but only shows a 10-fold selectivity for DPAP1 compared to CatC. While replacing the P2 Pro of SAK2 with hPro increases the potency of the inhibitors towards DPAP1, this also results in some loss of specificity (Table 3). Overall, we were able to increase the potency of SAK2 (Pro-hPhe-ACC) towards DPAP1 by 7-fold by using the optimal P1 (nLeu(o-Bzl)) and P2 (hPro) residues identified from the substrate library screen(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b), making hPro-nLeu(o-Bzl)-VS the most potent DPAP1 inhibitor identified so far (k_{inact}/K_i = 3.6·10^6 M^{-1}s^{-1})(Deu, Leyva, et al. 2010; Arastu-Kapur et al. 2008; Mahajan et al. 2011). However, this 7-fold increase in inhibitor potency between Pro-hPhe-VS and hPro-nLeu(o-Bzl)-VS was much lower than expected based on the 20- and 30-fold increase in susbtrate turnover reported between P1 hPhe and P1 nLeu(o-

Bzl), and between P2 Pro and P2 hPro, respectively(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b).

Testing inhibitors specificity in live parasites. We then tested the potency and selectivity of our DPAP1 and DPAP3 specific inhibitors in live parasites using the FY01 activity-based probe (ABP) in a competition labelling assay. ABPs are small molecule reporters of activity that use the catalytic mechanism of the targeted enzyme to covalently modify its active site. A reporter tag, usually a fluorophore or a biotin, allows visualization and quantification of the labelled enzyme in a gel-based assay(Sanman & Bogyo 2014). FY01 is a cell-permeable fluorescent ABP that was initially developed for CatC(Yuan et al. 2006) but it also labels *Plasmodium* DPAPs and the falcipains in live parasites(Arastu-Kapur et al. 2008; Deu, Leyva, et al. 2010). The falcipains (FPs) are clan CA Cys proteases involved in haemoglobin degradation (FP2, FP2' an FP3)(Sijwali et al. 2006) and possibly RBC invasion (FP1)(Sijwali et al. 2004; Greenbaum, Baruch, et al. 2002). Binding of inhibitors into the active site of any of these Cys proteases prevents probe labelling resulting in the disappearance of a fluorescent band a SDS-PAGE gel.

Live parasites were treated with different concentrations of inhibitor for 30 min, and the residual level of DPAPs and FPs activities labelled with FY01 (Figure S5) and quantified by densitometry. Dose response curves are shown in Figure 4 and IC_{50} values reported in Table 4. Inhibitors with a P2 Pro or hPro are equally potent and inhibit DPAP1 at mid nanomolar concentrations. However, the P2 Pro makes the inhibitor more selective by blocking inhibition of the FPs. Compounds with a P2 Trp or Tyr (NO_2) are highly specific for DPAP3 in intact parasites. Surprisingly, Trp-hPG-VS is by far the most potent DPAP3 inhibitor in intact parasites (IC_{50} = 1.4 nM) despite being 5 to 100-fold less potent than Trp-nLeu(o-Bzl)-VS or Trp-hPhe-VS against DPAP3 (see k_{inact}/K_i values in Table 3). This suggests that either this

compound is metabolically more stable and/or that decreasing the hydrophobicity of the P1 residue enhances the cell permeability of the compound. Indeed, compounds need to cross four membranes to reach DPAP3: the RBC, parasitophorous vacuole, and parasite plasma membranes, plus the membrane of the apical organelle where DPAP3 resides(Lehmann et al. 2017). (The parasitophorous vacuole is a membrane bound structure within which the parasite growth and multiply isolated from the RBC cytosol.) It is also likely that the apparent increased potency of Trp-hPG-VS in live parasites might be due to its accumulation into the DPAP3 acidic organelle via protonation of its free amine. However, we predict that this lysosomotropic effect likely occurs for all DPAP inhibitors presented here.

Figure 4. Selective inhibition of DPAP1 or DPAP3 in live parasites. Intact mature schizonts were pre-treated for 30 min with a dose response of inhibitor followed by 1 h treatment with FY01 to label the residual level of DPAP1, DPAP3, FP1 and FP2/3 activities. Fluorescents bands corresponding to the different cysteine proteases labelled by the probe (Figure S5) were quantifies by densitometry using ImageJ, and the fluorescence values normalized to the DMSO control. IC_{50} values are reported in Table 4. Note that control inhibitors containing *D*-Trp are more than 1000-fold less potent than those with *L*-Trp.

Table 4. Cysteine proteases IC$_{50}$ values in live parasites.

Inhibitor	IC$_{50,DPAP3}$ (nM)	IC$_{50,DPAP1}$ (nM)	IC$_{50,FP1}$ (nM)	IC$_{50,FP2/3}$ (nM)
hPro-hPhe-VS	160 ± 50	80 ± 7	350 ± 80	3,300 ± 1,000
Pro-hPhe-VS	> 10,000	275 ± 25	~ 10,000	> 10,000
hPro-nLeu(o-Bzl)-VS	900 ± 100	62 ± 7	300 ± 100	1,900 ± 500
Tyr (NO$_2$)-hPhe-VS	190 ± 20	> 10,000	> 10,000	> 10,000
Tyr (NO$_2$)-nLeu(o-Bzl)-VS	730 ± 170	> 10,000	> 10,000	> 10,000
L-Trp-hPG-VS	1.4 ± 0.4	> 10,000	> 10,000	> 10,000
L-Trp-hPhe-VS	130 ± 60	> 10,000	> 10,000	> 10,000
L-Trp-nLeu(o-Bzl)-VS	760 ± 150	~ 10,000	~ 10,000	~ 10,000
D-Trp-hPhe-VS	6,800 ± 1,600	> 10,000	> 10,000	> 10,000
D-Trp-nLeu(o-Bzl)-VS	2,700 ± 1,600	> 10,000	> 10,000	> 10,000

Correlation between PS-SCL of substrates and inhibitors for other Cys proteases. To determine whether the discrepancy observed between substrate and inhibitor specificities for DPAP3 (Figure 3) is a phenomenon observed in other proteases, we look at previously published literature about two of the most commonly studies families of Cys proteases, i.e. caspases and cathepsins. Several studies on caspases(Berger, Sexton, et al. 2006; Poreba et al. 2018) and cathepsins(Choe et al. 2006) have shown that inhibitors designed based on the structure of specific substrates do not retain their selectivity. Also, the sequence of an optimal substrate or inhibitor might render the equivalent inhibitor or substrate completely inactive[51](Choe et al. 2006). We therefore compared previously published specificity data obtained from PS-SCL libraries of ACC substrates and covalent inhibitors for these two protease families (Figure 5).

To simplify our analysis, we only compared data obtained from natural amino acids. For Cys cathepsins (CatB, CatF, CatK, CatL, CatS, and CatV), we compared the substrate specificity data obtained from the Craik lab(Choe et al. 2006) with the inhibitor specificity data obtained from the Bogyo lab using an epoxide library(Greenbaum, Arnold, et al. 2002) (Figure 5A-B). For caspases, we compared the specificity obtained for caspases 3, 8 and 9 (Casp3, Casp8 and Casp9, respectively) using a library of acyloxymethyl ketone inhibitors from the Bogyo lab(Berger, Witte, et al. 2006), with the substrate specificity results from the Drag

lab (Poreba, Kasperkiewicz, et al. 2014) (Figure 5C-D). For each protease, we normalized the level of activity and inhibition to that of the best natural amino acid at each P2-P4 position. As shown in Figure 5, there is very poor correlation between the levels of activity and inhibition. Indeed, for most proteases and P2-P4 positions, the Pearson coefficients are below 0.7. Similar to what we observed for DPAPs (Figure 3), the peptide sequence of inhibitors that is equivalent to that of optimal substrates generally yields very potent inhibitors. This is true for most proteases and for each position (with the exception of P2 in Casp9). However, for all proteases, 90-100 % inhibition was achieved with peptide sequences that were poorly cleaved as substrates (< 10 % activity). Also, at each P2-P4 position we could observe good inhibition with sequences belonging to poor substrates. Interestingly, for most proteases only a few peptide sequences showed good turnover rates but poor inhibition, with the exception of Casp8, and the P3 position for CatF. There are also a few striking individual residues, such as P2 His for caspases 8 and 9, or P2 Thr for Casp3 and P2 Val for CatS, that show good activity but no inhibition. Finally, we would also like to draw attention to the fact that the P2 and P4 positions are particularly important for modulating the potency of inhibitors against Casp9 and Casp3, respectively. However, these positions have little effect on substrate turnover. A possible explanation about the difference in specificity at the P4 position for caspases is that the P4 capping group is different between the substrate (acetyl group) and inhibitor (2-(4-hydroxy-3-nitrophenyl)-acetyl group) PS-CSLs, and might therefore strongly influence how the side chain of P4 residues bind into the S4 pocket. Indeed, using different capping group in PS-CSL of substrates has been shown to result in different amino acid preference in the S4 pocket of the Zika virus NS2B-NS3 protease(Rut et al. 2017; Gruba et al. 2016). Also, the difference in specificity might be due to cooperativity between the prime and non-prime sites (especially at the P2 position). In this case the choice of electrophile used might change the amino acid preference.

Figure 5. Comparison of substrate and inhibitor specificity for cathepsins and caspases.
(**A** and **C**) Structure of the PS-SCL of ACC substrates and inhibitors used to determine the specificity of cathepsins and caspases. For the cathepsins, substrates were screened at 250 μM, and the inhibitors at 10 μM using an ABP competition assay. For caspases, substrates were screened at 50 μM and inhibitors at 50-500 nM. (**B** and **D**) Correlation between substrate turnover and inhibition. For each protease and P2-P4 position, the maximum level of activity and inhibition at each AA position was normalized to 100 %. Pearson correlation coefficients (ρ) are shown next to the legend of each graph for each position. The lower ρ value was obtained by combining all the P2-P4 data points for each protease. Residues showing maximum activity or inhibition are labelled at each position along with specific AAs that illustrate particularly bad correlation between the levels of activity and inhibition.

Discussion

This study provides the first characterization of the specificity of DPAP3, a cysteine protease important for efficient invasion of RBCs by the malaria parasite(Lehmann et al. 2017). DPAP3 is a highly efficient proteolytic enzyme showing similar k_{cat} and k_{cat}/K_m values than either DPAP1 or CatC when optimal substrates are used (Table 1). In general, DPAP3 has a narrower substrate specificity than either DPAP1 or CatC, and it preferentially cleaves substrates with aliphatic residues at the N-terminus. Our study also shows similar P1 substrate specificity across all DPAPs, i.e. a strong preference for basic or long aromatic residues. DPAP3 was the only DPAP able to cleave some substrates with aromatic P2 residues (Tyr, Phe(Me), Phg), albeit with relatively low turnover rates. Surprisingly, vinyl sulfone inhibitors with aromatic residues are as potent as compounds with optimal P2 residues identified from the substrate screen, but in addition, these compounds are highly specificity for DPAP3 compared to DPAP1, CatC, or other malarial cysteine proteases (FPs). The previously described DPAP1 inhibitor SAK2 (Pro-hPhe-VS) shows the greatest specificity for DPAP1 in live parasites. Incorporation of optimal P1 (nLeu(o-Bzl)) and P2 (hPro) residues identified from the substrate screens improves the potency of DPAP1 inhibitors both *in vitro* and in live parasites, but also results in some loss in specificity (Table 3 and Figure 4).

Despite being highly potent and specific DPAP1 and DPAP3 inhibitors, these compounds only show antiparasitic activity at mid to high micromolar concentrations(Deu, Leyva, et al. 2010; Arastu-Kapur et al. 2008; Lehmann et al. 2017), probably due to metabolic stability issues. Indeed, we have previously shown that this is the case for Pro-hPhe-VS, which is not able to sustain target inhibition in live parasites(Deu, Leyva, et al. 2010). A possible cause for this instability is the presence of multiple aminopeptidases in the malaria parasite that might cleave the amide bond of these compounds(Deu 2017), thus preventing them to bind into the DPAPs active sites. Nonetheless, this study provides a very strong SAR foundation to

develop potent non-peptidic inhibitors able to sustain DPAPs inhibition. From a drug development point of view, our SAR studies indicate that inhibitors with short aliphatic P2 residues strongly inhibit both DPAPs (Figure 3E), indicating that potent pan-DPAP inhibitors can easily be developed. Unfortunately, we did not identify any clear P1 or P2 residue that would discriminate malarial DPAPs from host CatC. Therefore, further studies are required to determine whether differences in specificity in the prime binding pockets can be exploited to develop parasite specific inhibitors.

It is however important to point out that given the short-term treatment requirement for antimalarial therapy (single dose or less than 3 doses in 3 days), we think it is unlikely that inhibition of CatC would lead to adverse side effects. Firstly, highly specific DPAP inhibitors might not be necessary given that a high level (> 95 %) of sustained CatC inhibition is required to see a decrease in the activity of serine proteases activated by CatC(Methot et al. 2008). Secondly, activation of granule serine proteases by cathepsin C takes place during cell differentiation in the bone marrow, and a decrease in the level of serine proteases activity in circulating immune cells is only achieved after more than 2 weeks of daily treatment with CatC inhibitors(Palmér et al. 2018). And thirdly, no signs of toxicity were observed in phase I clinical trials when volunteers were treated daily for more than 3 weeks with CatC inhibitors, albeit some on-target side effects such as plantar and palmar epithelial desquamation were observed in some instances(Miller et al. 2017; Palmér et al. 2018). However, these side effects were not observed in volunteers that received a single dose of CatC inhibitor, nor within the first week of daily treatments.

Positional scanning substrate libraries have been used over the last 20 years(Thornberry et al. 1997; Rano et al. 1997; Harris et al. 2000) to determine the specificity of proteases and guide the synthesis of inhibitors. Here we have shown that vinyl sulfone inhibitors containing

P1 and P2 residues corresponding to optimal DPAP1 or DPAP3 substrates result in extremely potent inhibitors. However, our studies clearly identified differences in specificity between substrates and inhibitors, especially for DPAP3, but also for CatC and more generally in caspases and cysteine cathepsins (Figures 3 and 5). A reason why this phenomenon might not have been more broadly reported in the literature is because, in general, either substrate or inhibitor libraries are used to determine the specificity of a protease, but not both. Also, inhibitors are not usually designed based on the structure of non-optimal substrates. That said, there are multiple reasons that can account for this discrepancy in specificity between substrates and inhibitors:

First, although the substrate and inhibitor libraries used in this study have equivalent P1 and P2 residues, the structural features that bind into the S' pockets are quite different. Therefore, if the specificity of a protease shows interdependence between its prime and non-prime binding pockets, it might explain the difference in specificity. This is likely the case for DPAPs since we observed a 50-fold increase in k_{cat} between the Phe-Arg-ACC and Phe-Arg-ⱯNA substrates, which only differ in the structure of the fluorophore that binds in the S1' pocket (Table 1). Also, while we observed very good correlation between k_{cat}/K_m and k_{inact}/K_i of DPAP1 for the P2 substrate and VS library (compounds in Table 2), in the context of a phenyl group in P1' (inhibitors in Table 3), we observed clear discrepancies between substrates and inhibitors (Figure 3B).

Second, the position of the electrophilic warhead within the active site might differ from that of the scissile bond in a substrate, especially in terms of distance and orientation relative to the catalytic Cys. This positioning might be differently affected by changes in the P1 and P2 residues of substrates and inhibitors. Indeed, a recent study on caspases has shown that acyloxymethyl ketone covalent inhibitors might act through a reversible mechanism even if they are designed based on the sequence of optimal substrates(Poreba et al. 2018). Also, ABPs

designed to profiled deubiquitinating proteases by conjugating an electrophile to the C-terminal of ubiquitin have been shown to label different subset of enzymes depending on the warhead used (Borodovsky et al. 2002).

Third, substrate turnover by Cys and Ser proteases requires two different chemical steps: First, nucleophilic attack of the peptide bond by the catalytic residue to form the acyl intermediate and release of the N-terminal product of proteolysis; and second, hydrolysis of the acyl intermediate by an activated water molecule to reconstitute the free enzyme and release the C-terminal product of the reaction (Scheme 1). Therefore, peptide sequences that are poorly turned over because of a very slow acyl intermediate hydrolysis step might still be very good to design covalent inhibitors. For example, a substrate that displaces the catalytic water from the active site upon formation of the acyl intermediate will be very poorly turned over.

$$\text{E} + \text{I} \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\rightleftarrows}} \text{E:I} \xrightarrow{k_{\text{inact}}} E-I$$

$$K_{\text{i}} = \frac{k_{\text{off}}}{k_{\text{on}}} \; ; \; \frac{k_{\text{inact}}}{K_{\text{i}}} = \frac{k_{\text{on}} \cdot k_{\text{inact}}}{k_{\text{off}}}$$

$$\text{E} + \text{S} \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\rightleftarrows}} \text{E:S} \xrightarrow{k_{\text{Ac}}} \text{P}_{\text{C}} + \text{E} - \text{Ac} \xrightarrow{k_{\text{Dac}}} \text{P}_{\text{N}} + \text{E}$$

$$k_{\text{cat}} = \frac{k_{\text{Ac}} \cdot k_{\text{Dac}}}{k_{\text{Ac}} + k_{\text{Dac}}} \; ; \; K_{\text{m}} = \frac{k_{\text{Dac}} \cdot (k_{\text{off}} + k_{\text{Ac}})}{k_{\text{on}} \cdot (k_{\text{Ac}} + k_{\text{Dac}})} \; ; \; \frac{k_{\text{cat}}}{K_{\text{m}}} = \frac{k_{\text{on}} \cdot k_{\text{Ac}}}{(k_{\text{off}} + k_{\text{Ac}})}$$

Scheme 1. Comparison between irreversible inhibition and substrate turnover kinetic constants. E, I, E:I and E-I represent free enzyme, inhibitor, inhibitor associated with the enzyme, and enzyme covalently modified by the inhibitor, respectively. S, E:S, E-Ac, P_{C} and P_{N} represent the substrate, the Michaelis Menten enzyme-substrate complex, the acyl intermediate, and the C- and N-terminal products of proteolysis, respectively; k_{on} and k_{off} the association and dissociation rate constant for substrates or inhibitors; and k_{Ac} and k_{Dac} the kinetic constants for formation of the acyl intermediate and its hydrolysis, respectively.

And fourth, the reaction mechanism between covalent inhibition and substrate turnover are quite different making k_{cat}, K_{m}, and $k_{\text{cat}}/K_{\text{m}}$ not directly comparable with k_{inact}, K_{i}, and $k_{\text{inact}}/K_{\text{i}}$

(Scheme 1). k_{cat}, K_m, and k_{cat}/K_m are empirical parameters obtained under steady state conditions while k_{inact}, K_i, and k_{inact}/K_i are real kinetic and thermodynamic constants that cannot be measured under steady state conditions since covalent inhibitors deplete the concentration of free enzyme over time. k_{cat} might be equivalent to k_{inact} only if $k_{Ac} \ll k_{Dac}$, i.e. formation of the acyl intermediate is the rate limiting step; K_m might be equivalent to K_i only if $k_{Ac} \ll k_{off}$ and $k_{Ac} \ll k_{Dac}$; and k_{cat}/K_m might be equivalent to k_{inact}/K_i only if $k_{off} \gg k_{Ac}$. Although these conditions might be met for certain substrates, the relative magnitudes of all these rate constants depend on the substrate sequence. For example, detailed pre-steady state kinetics and kinetic isotope effects studies on CatC have shown that the rate limiting step in the turnover of dipeptidic AMC substrates can be either formation of the acyl intermediate, its hydrolysis, or a contribution of both, depending on the nature of the P1 residue(Schneck et al. 2008; Rubach et al. 2012).

Although several studies have compared the potency of inhibitors to the turnover of equivalent substrates for selected peptide sequences, to the best of our knowledge this is the first study that systematically compares the potency of a peptide-based covalent inhibitor library to the turnover efficiency of an equivalent substrate library. We think that the discrepancies observed here between k_{cat}/K_m and k_{inact}/K_i are likely to be present in other proteases, as shown for cysteine cathepsins and caspases, but the level of discrepancy will be dependent on the protease studied as well as on the type of substrate and covalent inhibitor that are being compared especially if a there is a high level of cooperativity between the prime and non-prime binding pockets.

Overall, our detailed study on DPAPs specificity and our analysis of specificity results obtained from PS-SCL clearly indicate that there are very significant differences in specificity between substrates and covalent inhibitors. Although it is now well established that highly

potent inhibitors can be developed based on the structure of optimal substrates, this might sometimes result in some loss of specificity. This study clearly demonstrates that optimal inhibitors with improved specificity can be developed based on the structure of relatively poor substrates. Therefore, we strongly recommend using PS-SCL libraries of inhibitors rather than substrates if the goal is to design specific inhibitors and ABPs, and use PS-SCL substrate libraries to determine the substrate specificity of proteases and to develop highly selective substrates as high throughput dynamic tools to measure protease activity in complex biological samples.

Materials and Methods

Reagents. The syntheses of the DPAP substrate library and that of Val-Arg-ACC, Phe-Arg-ACC, Tyr (NO_2)-hPhe-ACC(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014a) and $(PR)_2$Rho(Wang et al. 2011) have been previously described. The syntheses of additional substrates used in this study are described in the supplementary methods and were synthesized following previously published methods(Poreba, Mihelic, Krai, Rajkovic, Krezel, Pawelczak, Klemba, D. Turk, B. Turk, Latajka & Drag 2014b)˙(Poreba, Szalek, et al. 2014). The syntheses of the vinyl sulfone inhibitor library, SAK2, SAK1, *L*-WSAK and *D*-WSAK were also previously described(Lehmann et al. 2017)˙(Arastu-Kapur et al. 2008). The additional DPAP inhibitors used in this study were synthesized following similar methods(Yang et al. 2012). The synthesis and characterization of these inhibitors and their synthetic intermediates are described in detail in the supplementary methods. The Phe-Arg-βNA and Gly-Arg-AMC substrates were purchased from Sigma. Recombinant DPAP3 was expressed in insect cells using the bacculovirus system as recently described(Lehmann et al. 2017). Bovine CatC was purified to homogeneity from spleen by modification of a method described previously(Anon 2018)˙(Horn et al. 2002).

Recombinant DPAP3 active site titration. Recombinant DPAP3 was expressed in SF9 insect cells and purified from the culture supernatant by sequential ion exchange, Ni-NTA, and size exclusion chromatography as previously described(Lehmann et al. 2017). To accurately determine the concentration of active DPAP3 in our enzyme stocks we used the FY01 ABP. Because ABPs only react with the active form of an enzyme and covalently modify its active site, in this case the catalytic Cys, they can be used to perform accurate active site titrations.

Our stock of DPAP3 was diluted 20-fold in assay buffer (100 mM sodium acetate, 100 mM NaCl, 5 mM $MgCl_2$, 1 mM EDTA, 0.1 % CHAPS, and 5 mM DTT at pH 6), pre-treated for 30 min with DMSO or 1 µM SAK1 (Tyr (NO_2)-hPhe-VS), and DPAP3 labelled with 1 µM FY01 for 1 h at RT. These samples were run on a SDS-PAGE gel along with a serial dilution of free probe (1.5-100 nM). In-gel fluorescence was measured using a Bio-Rad PharosFX flat-bed scanner and the intensity of the fluorescent bands quantified using ImageJ. The fluorescent signal from the free probe was used as a calibration curve and compared to the difference in signal between the DMSO and SAK1 treated DPAP3 (Figure S6). Using this method, we determined that our DPAP3 stock contained 840 nM of active protease.

Substrate turnover assay. The substrate library was screened in triplicate at 1 µM substrate and 1 nM DPAP3 in assay buffer. Substrate turnover was measured over 30 min at RT using a SpectroMax M5e plate reader: $\lambda_{ex} = 355$ nm, $\lambda_{em} = 460$ nm, emission filter 455 nm, for ACC or AMC (7-amino-4-methylcoumarin) substrates; $\lambda_{ex} = 315$ nm, $\lambda_{em} = 355$ nm, emission filter 420 nm, for Phe-Arg-βNA); and $\lambda_{ex} = 492$ nm, $\lambda_{em} = 523$ nm, emission filter 520 nm for $(PR)_2$Rho. Calibration curves of free βNA (β-napthylamide) and ACC (0-500 nM) under the same assay conditions were performed to convert the turnover rate measured as fluorescent units per second into moles per second. The determine k_{cat} and K_m values, substrate turnover was measured at different substrate concentrations in assay buffer using 1 nM of DPAP3 or

1 nM of bovine CatC. The initial velocities were then fitted with Prism to the Michaelis Menten Eqs. 3 or 4 to obtain accurate k_{cat} and K_m or k_{cat}/K_m values, respectively. A minimum of three replicates were performed for each substrate.

$$V_i = \frac{k_{cat} \cdot Et \cdot [S]}{K_m + [S]}$$
Eq. 3

$$V_i = \frac{(k_{cat}/K_m) \cdot Et \cdot [S]}{1 + \frac{[S]}{K_m}}$$
Eq. 4

V_i is the initial velocity and Et the total concentration of active protease.

Irreversible inhibition assay. To determine the inhibition constants of vinyl sulfone inhibitors against DPAP3, 2.2 µM of Met-nLeu(o-Bzl)-ACC (0.25 x K_m) was mixed with a dose response of inhibitor in assay buffer, and the turnover rate measured over 40 min at RT after addition of 0.2 nM DPAP3. The data was analyzed according to the irreversible inhibition model shown in Eq. 1. First, the progress curves (fluorescent units vs. time) were fitted to Eq. 5, where F is the measured fluorescence at time t, F_0 the initial fluorescence, V_0 the initial turnover rate, and k_{obs} the observed second order rate constant of inhibition measured at each inhibitor concentration(Morrison & Walsh 1988).

$$(F - F_0) = V_0 \frac{1 - e^{-k_{obs} \cdot t}}{k_{obs}}$$
Eq. 5

Second, k_{obs} values were fitted to Eqs. 6 or 7. to obtain the k_{inact} and K_i or k_{inact}/K_i values, respectively.

$$k_{obs} = \frac{k_{inact} \cdot [INH]}{K_i + [INH]}$$
Eq. 6

$$k_{\text{obs}} = \frac{(k_{\text{inact}}/K_i)\cdot[\text{INH}]}{1+\frac{[\text{INH}]}{K_i}} \qquad \text{Eq. 7}$$

Inhibition constants for DPAP1 and CatC were determined in a similar way but using 0.25 x K_{m} concentrations of the Pro-Arg-AMC (20 µM) and Gly-Arg-AMC (15 µM), respectively. Inhibition of DPAP1 was directly measured in parasite lysates diluted 100-fold in assay buffer using the DPAP1-selective substrate Pro-Arg-AMC(Deu, Yang, et al. 2010); 0.2 nM of CatC was used for these inhibition studies.

Measurement of inhibitor specificity in biological samples. Inhibition of *Plasmodium* DPAPs and the falcipains in intact parasites was measured using the FY01 ABP in competition assays as previously described(Arastu-Kapur et al. 2008; Deu, Leyva, et al. 2010). Because DPAP3 is maximally expressed in very mature schizonts(Bozdech et al. 2003), labelling was performed after treating parasites with 1 µM of Compound 2, a cGMP-dependent protein kinase inhibitor that arrest parasite development 15-30 min before they egress from the infected RBC(Collins et al. 2013). For each sample, 5 µL of percoll-purified schizonts were diluted in 45 µL of RPMI (Gibco), pre-treated for 30 min with a dose response of inhibitor, and labelled for 1 h with 1 µM FY01. Samples were then boiled for 10 min in loading buffer and run on a 12 % SDS-PAGE gel. Fluorescently labelled proteases in the gel were detected using a Bio-Rad PharosFX flatbed fluorescence scanner.

Acknowledgement

We would like to thank the Wellcome Trust and Royal Society for funding this research through the Sir Henry Dale Fellowship 099950. We would also like to thank Erasmus for funding L.E.d.V. and L.K. internship in Dr. Edgar Deu's laboratory.

References

Adkison, A.M. et al., 2002. Dipeptidyl peptidase I activates neutrophil-derived serine proteases and regulates the development of acute experimental arthritis. *Journal of Clinical Investigation*, 109(3), pp.363–371.

Anon, 2018. Cathepsins. *Methods in Enzymology*, 19, pp.285–315.

Arastu-Kapur, S. et al., 2008. Identification of proteases that regulate erythrocyte rupture by the malaria parasite Plasmodium falciparum. *Nature Chemical Biology*, 4(3), pp.203–213.

Ashley, E.A. et al., 2014. Spread of artemisinin resistance in Plasmodium falciparum malaria. *The New England journal of medicine*, 371(5), pp.411–423.

Berger, A.B., Sexton, K.B. & Bogyo, M., 2006. Commonly used caspase inhibitors designed based on substrate specificity profiles lack selectivity. *Cell research*, 16(12), pp.961–963.

Berger, A.B., Witte, M.D., et al., 2006. Identification of early intermediates of caspase activation using selective inhibitors and activity-based probes. *Molecular cell*, 23(4), pp.509–521.

Bhatt, S. et al., 2015. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. *Nature*, 526(7572), pp.207–211.

Borodovsky, A. et al., 2002. Chemistry-based functional proteomics reveals novel members of the deubiquitinating enzyme family. *Chemistry & Biology*, 9(10), pp.1149–1159.

Bozdech, Z. et al., 2003. The Transcriptome of the Intraerythrocytic Developmental Cycle of Plasmodium falciparum Gary Ward, ed. *PLoS biology*, 1(1), pp.e5–16.

Capuccini, B. et al., 2016. Transcriptomic profiling of microglia reveals signatures of cell activation and immune response, during experimental cerebral malaria. *Nature Publishing Group*, 6(1), p.39258.

Choe, Y. et al., 2006. Substrate profiling of cysteine proteases using a combinatorial peptide library identifies functionally unique specificities. *Journal of Biological Chemistry*, 281(18), pp.12824–12832.

Collins, C.R. et al., 2013. Malaria parasite cGMP-dependent protein kinase regulates blood stage merozoite secretory organelle discharge and egress. L. D. Sibley, ed. *PLoS Pathogens*, 9(5), p.e1003344.

Deu, E., 2017. Proteases as antimalarial targets: strategies for genetic, chemical, and therapeutic validation. *The FEBS journal*, 284(16), pp.2604–2628.

Deu, E., Leyva, M.J., et al., 2010. Functional Studies of Plasmodium falciparum Dipeptidyl Aminopeptidase I Using Small Molecule Inhibitors and Active Site Probes. *Chemistry & Biology*, 17(8), pp.808–819.

Deu, E., Yang, Z., et al., 2010. Use of activity-based probes to develop high throughput screening assays that can be performed in complex cell extracts. F. Romesberg, ed. *PloS*

one, 5(8), p.e11985.

Drag, M. et al., 2010. Aminopeptidase fingerprints, an integrated approach for identification of good substrates and optimal inhibitors. *The Journal of biological chemistry*, 285(5), pp.3310–3318.

Furber, M. et al., 2014. Cathepsin C inhibitors: property optimization and identification of a clinical candidate. *Journal of Medicinal Chemistry*, 57(6), pp.2357–2367.

Greenbaum, D.C., Arnold, W.D., et al., 2002. Small molecule affinity fingerprinting. A tool for enzyme family subclassification, target identification, and inhibitor design. *Chemistry & Biology*, 9(10), pp.1085–1094.

Greenbaum, D.C., Baruch, A., et al., 2002. A role for the protease falcipain 1 in host cell invasion by the human malaria parasite. *Science (New York, N.Y.)*, 298(5600), pp.2002–2006.

Gruba, N. et al., 2016. Substrate profiling of Zika virus NS2B-NS3 protease. *FEBS Letters*, 590(20), pp.3459–3468.

Guay, D. et al., 2009. Design and synthesis of dipeptidyl nitriles as potent, selective, and reversible inhibitors of cathepsin C. *Bioorganic & Medicinal Chemistry Letters*, 19(18), pp.5392–5396.

GUTMANN, H.R. & FRUTON, J.S., 1948. On the proteolytic enzymes of animal tissues; an intracellular enzyme related to chymotrypsin. *Journal of Biological Chemistry*, 174(3), pp.851–858.

Harris, J.L. et al., 2000. Rapid and general profiling of protease specificity by using combinatorial fluorogenic substrate libraries. *Proceedings of the National Academy of Sciences*, 97(14), pp.7754–7759.

Horn, M. et al., 2002. Free-thiol Cys331 exposed during activation process is critical for native tetramer structure of cathepsin C (dipeptidyl peptidase I). *Protein Science*, 11(4), pp.933–943.

Kasperkiewicz, P. et al., 2014. Design of ultrasensitive probes for human neutrophil elastase through hybrid combinatorial substrate library profiling. *Proceedings of the National Academy of Sciences of the United States of America*, 111(7), pp.2518–2523.

Kasperkiewicz, P. et al., 2017. Emerging challenges in the design of selective substrates, inhibitors and activity-based probes for indistinguishable proteases. *The FEBS journal*, 284(10), pp.1518–1539.

Klemba, M., Gluzman, I. & Goldberg, D.E., 2004. A Plasmodium falciparum dipeptidyl aminopeptidase I participates in vacuolar hemoglobin degradation. *The Journal of biological chemistry*, 279(41), pp.43000–43007.

Korkmaz, B. et al., 2018. Therapeutic targeting of cathepsin C: from pathophysiology to treatment. *Pharmacology & therapeutics*.

Kummer, J.A. et al., 1996. Expression of human recombinant granzyme A zymogen and its

activation by the cysteine proteinase cathepsin C. *Journal of Biological Chemistry*, 271(16), pp.9281–9286.

Lainé, D. et al., 2011. Discovery of novel cyanamide-based inhibitors of cathepsin C. *ACS medicinal chemistry letters*, 2(2), pp.142–147.

Lehmann, C. et al., 2017. Plasmodium falciparum dipeptidyl aminopeptidase 3 activity is important for efficient erythrocyte invasion by the malaria parasite. *bioRxiv*, pp.1–39.

Lin, J.-W. et al., 2015. Replication of Plasmodiumin reticulocytes can occur without hemozoin formation, resulting in chloroquine resistance. *The Journal of Experimental Medicine*, 212(6), pp.893–903.

Mahajan, S.S. et al., 2011. A fragmenting hybrid approach for targeted delivery of multiple therapeutic agents to the malaria parasite. *ChemMedChem*, 6(3), pp.415–419.

McGuire, M.J., Lipsky, P.E. & Thiele, D.L., 1993. Generation of active myeloid and lymphoid granule serine proteases requires processing by the granule thiol protease dipeptidyl peptidase I. *Journal of Biological Chemistry*, 268(4), pp.2458–2467.

Methot, N. et al., 2008. In vivo inhibition of serine protease processing requires a high fractional inhibition of cathepsin C. *Molecular pharmacology*, 73(6), pp.1857–1865.

Miller, B.E. et al., 2017. Epithelial desquamation observed in a phase I study of an oral cathepsin C inhibitor (GSK2793660). *British journal of clinical pharmacology*, 83(12), pp.2813–2820.

Morrison, J.F. & Walsh, C.T., 1988. The behavior and significance of slow-binding enzyme inhibitors. *Advances in enzymology and related areas of molecular biology*, 61, pp.201–301.

Nazif, T. & Bogyo, M., 2001. Global analysis of proteasomal substrate specificity using positional-scanning libraries of covalent inhibitors. *Proceedings of the National Academy of Sciences*, 98(6), pp.2967–2972.

Palmér, R. et al., 2018. Dipeptidyl Peptidase 1 Inhibitor AZD7986 Induces a Sustained, Exposure-Dependent Reduction in Neutrophil Elastase Activity in Healthy Subjects. *Clinical pharmacology and therapeutics*.

Pham, C.T. & Ley, T.J., 1999. Dipeptidyl peptidase I is required for the processing and activation of granzymes A and B in vivo. *Proceedings of the National Academy of Sciences*, 96(15), pp.8627–8632.

Poreba, M. et al., 2018. Caspase selective reagents for diagnosing apoptotic mechanisms. *Cell death and differentiation*, 9(Pt 10), p.231.

Poreba, M. et al., 2012. Fingerprinting the Substrate Specificity of M1 and M17 Aminopeptidases of Human Malaria, Plasmodium falciparum P. Sinnis, ed. *PloS one*, 7(2), pp.e31938–8.

Poreba, M., Kasperkiewicz, P., et al., 2014. Unnatural amino acids increase sensitivity and provide for the design of highly selective caspase substrates. *Cell death and differentiation*,

21(9), pp.1482–1492.

Poreba, M., Mihelic, M., Krai, P., Rajkovic, J., Krezel, A., Pawelczak, M., Klemba, M., Turk, D., Turk, B., Latajka, R. & Drag, M., 2014a. Unnatural amino acids increase activity and specificity of synthetic substrates for human and malarial cathepsin C. *Amino Acids*, 46(4), pp.931–943.

Poreba, M., Mihelic, M., Krai, P., Rajkovic, J., Krezel, A., Pawelczak, M., Klemba, M., Turk, D., Turk, B., Latajka, R. & Drag, M., 2014b. Unnatural amino acids increase activity and specificity of synthetic substrates for human and malarial cathepsin C. *Amino Acids*, 46(4), pp.931–943.

Poreba, M., Szalek, A., et al., 2014. Positional scanning substrate combinatorial library (PS-SCL) approach to define caspase substrate specificity. *Methods in molecular biology (Clifton, N.J.)*, 1133, pp.41–59.

Rano, T.A. et al., 1997. A combinatorial approach for determining protease specificities: application to interleukin-1beta converting enzyme (ICE). *Chemistry & Biology*, 4(2), pp.149–155.

Ranson, H. & Lissenden, N., 2016. Insecticide Resistance in African Anopheles Mosquitoes: A Worsening Situation that Needs Urgent Action to Maintain Malaria Control. *Trends in Parasitology*, 32(3), pp.187–196.

Rubach, J.K. et al., 2012. The amino-acid substituents of dipeptide substrates of cathepsin C can determine the rate-limiting steps of catalysis. *Biochemistry*, 51(38), pp.7551–7568.

Rut, W. et al., 2017. Extended substrate specificity and first potent irreversible inhibitor/activity-based probe design for Zika virus NS2B-NS3 protease. pp.1–7.

Sanman, L.E. & Bogyo, M., 2014. Activity-Based Profiling of Proteases. *Annual Review of Biochemistry*, 83(1), pp.249–273.

Schneck, J.L. et al., 2008. Chemical mechanism of a cysteine protease, cathepsin C, as revealed by integration of both steady-state and pre-steady-state solvent kinetic isotope effects. *Biochemistry*, 47(33), pp.8697–8710.

Schwach, F. et al., 2015. PlasmoGEM, a database supporting a community resource for large-scale experimental genetics in malaria parasites. *Nucleic acids research*, 43(Database issue), pp.D1176–82.

Sijwali, P.S. et al., 2006. Gene disruptions demonstrate independent roles for the four falcipain cysteine proteases of Plasmodium falciparum. *Molecular & Biochemical Parasitology*, 150(1), pp.96–106.

Sijwali, P.S. et al., 2004. Plasmodium falciparum cysteine protease falcipain-1 is not essential in erythrocytic stage malaria parasites. *Proceedings of the National Academy of Sciences*, 101(23), pp.8721–8726.

Suárez-Cortés, P. et al., 2016. Comparative Proteomics and Functional Analysis Reveal a Role of Plasmodium falciparum Osmiophilic Bodies in Malaria Parasite Transmission. *Molecular & cellular proteomics : MCP*, 15(10), pp.3243–3255.

Tanaka, T.Q. et al., 2013. Plasmodium dipeptidyl aminopeptidases as malaria transmission-blocking drug targets. *Antimicrobial agents and chemotherapy*, 57(10), pp.4645–4652.

Thornberry, N.A. et al., 1997. A combinatorial approach defines specificities of members of the caspase family and granzyme B. Functional relationships established for key mediators of apoptosis. *Journal of Biological Chemistry*, 272(29), pp.17907–17911.

Wang, F. et al., 2011. Biochemical characterization of Plasmodium falciparum dipeptidyl aminopeptidase 1. *Molecular & Biochemical Parasitology*, 175(1), pp.10–20.

Wells, T.N.C., Hooft van Huijsduijnen, R. & Van Voorhis, W.C., 2015. Malaria medicines: a glass half full? *Nature Reviews Drug Discovery*, 14(6), pp.424–442.

WIGGANS, D.S., WINITZ, M. & FRUTON, J.S., 1954. Action of cathepsin C on dipeptide esters. *The Yale journal of biology and medicine*, 27(1), pp.11–19.

World Health Organization, 2017. World Malaria Report 2016. pp.1–186.

Yang, P.-Y. et al., 2012. Proteomic profiling and potential cellular target identification of K11777, a clinical cysteine protease inhibitor, in Trypanosoma brucei. *Chemical communications (Cambridge, England)*, 48(6), pp.835–837.

Young, J.A. et al., 2005. The Plasmodium falciparum sexual development transcriptome: a microarray analysis using ontology-based pattern identification. *Molecular & Biochemical Parasitology*, 143(1), pp.67–79.

Yuan, F. et al., 2006. A selective activity-based probe for the papain family cysteine protease dipeptidyl peptidase I/cathepsin C. *Journal of the American Chemical Society*, 128(17), pp.5616–5617.