A genomic island of *Streptomyces coelicolor* with the self-contained regulon of an ECF sigma factor

Camilla M. Kao,[a,][#] Nitsara Karoonuthaisiri,[a,][1] David Weaver,[a,][2] Jonathan A. Vroom,[a,][3]

Shuning A. Gai,[a,][4] Mai-Lan Ho,[a,][5] and Kedar G. Patel[a]

[a]Department of Chemical Engineering, Stanford University, Stanford, California, USA

Running Head: GI 6 of *S. coelicolor* with ECF sigma factor regulon

#Address correspondence to Camilla M. Kao, cammy@stanford.edu.

[ ] Present Address: School of Engineering, Stanford University, Stanford, California, USA

[ ]1 Present Address: Biosensing Technology Research Unit, National Center for Engineering and

Biotechnology, Pathum Thani, Thailand

[ ]2 Present Address: ERM: Environmental Resources Management, San Francisco, California,

USA

[ ]3 Present Address: Molecular and Cellular Engineering, Codexis, Redwood City, California, USA

18 ☐4 Present Address: Molecular Biology, Prothena Biosciences, South San Francisco, California,

19 USA

20 ☐5 Present Address: Department of Radiology, Mayo Clinic, Rochester, Minnesota, USA

21 N.K. and D.W. contributed equally to this work.

Abstract

Streptomycetes constitute the largest genus of actinobacteria, living predominantly in soil and

decaying vegetation. The bacteria are widely known for their filamentous morphologies and

their capacity to synthesize antibiotics and other biologically active molecules. More than a

decade ago, we and others identified 22 genomic islands that *Streptomyces coelicolor* M145

possesses and other *Streptomyces* strains lack. One of these genomic islands, Genomic Island

(GI) 6, encodes an extracytoplasmic function (ECF) sigma factor that we were characterizing in

separate work. Here we report that artificial induction of the ECF sigma factor, which is

encoded by SCO3450, causes the transcription of approximately one-fourth of GI 6, or ~26

mostly contiguous genes, to increase. More than half of the regulon encodes putative enzymes

involved in small molecule metabolism. A putative haloacid dehalogenase is present. Genes

encoding two putative anti-sigma factors flank SCO3450, the three genes residing within the

regulon. Our data suggest that the ECF sigma factor and its regulon are a self-contained

transcriptional unit that can be transferred by horizontal gene transfer. To our knowledge, only

one other example has been identified of an ECF sigma factor and its contiguous regulon

appearing to be transferrable by horizontal gene transfer [18,19]. Because the regulon appears

not to be induced by the 44 growth conditions recently examined by Byung-Kwan Cho and

colleagues [20], if it confers fitness to *S. coelicolor*, the regulon likely does so in as-yet unknown

situations. Those situations might range from scavenging to detoxification to even

communication within microbial communities.

IMPORTANCE *Streptomyces* bacteria grow as hyphae that colonize soil and differentiate into

spores when nutrients become scarce. In their terrestrial habitats, the bacteria encounter

diverse conditions. Presumably so that the bacteria can cope with those conditions, the

chromosomes of streptomycetes are highly dynamic, varying greatly in structure not only

between species but also between closely related strains of a single species. The bacteria also

have large numbers of extracytoplasmic function (ECF) sigma factors, which undoubtedly help

the microorganisms respond to the plethora of challenges coming from the environment. This

work illustrates these two threads of *Streptomyces* biology dovetailing: Genetic adaptability

through horizontal gene transfer seems to have enabled *Streptomyces coelicolor* to acquire a

self-contained transcriptional unit that consists of an ECF sigma factor and its regulon. The

suggested facile movement of the regulon between microbial hosts indicates the value of the

metabolism of small molecules possibly mediated by the regulon.

Introduction

Researchers began studying *Streptomyces* genetics in the second half of the 20th century

because of the filamentous morphologies and biosynthetic capacities of the bacteria [1]. Many

of the *Streptomyces* strains that were studied belong to a group of closely related "blue"

streptomycetes that had been given various names, but probably should all have the same

name [2]. During the early 2000's, we characterized the genomes of six of these "blue" strains

by using DNA microarrays: the sequenced strain [3], *S. coelicolor* M145, which derives from *S.*

coelicolor A3(2), which in turn derives from Waksman's strain 3443 [4]; Sermonti's SE1 (John

Innes (JI) strain 1152) [5]; Bradley's S199 (JI 1153) [6]; *S. lividans* 66 (JI 1326) [7]; *S. lividans*

ISP5434 (JI 2896), which contains the plasmid pIJ101 [8]; and *S. violaceoruber* SANK95570 (JI

3034), which contains the plasmid pSV1 that encodes the biosynthetic genes for the antibiotic

methylenomycin [9]. That study was described in the doctoral dissertation of co-author David

Weaver [10]. Similar findings were reported by Jayapal et al., who examined *S. coelicolor* M145

and *S. lividans* TK21 [11].

By comparing the genome of *S. coelicolor* M145 to the five wild type genomes by using DNA

microarrays, we identified 22 sets of contiguous genes in *S. coelicolor* M145 that are absent in

the wild type strains [10]. We designated these genes Genomic Islands (GIs) 1 to 22. Fig. S1, Fig.

S1, and Fig. S3 show the GIs. The sizes of the GIs range from 3 kb to 150 kb and the number of

ORFs within them from 3 to 148. *S. coelicolor* M145 likely acquired the 22 GIs by horizontal

gene transfer, because they have the following characteristics: Direct repeats flank most of the

GIs; the GIs contain regions with low G + C content; the GIs encode transposable elements such

as invertases, recombinases, and transposons, as well as plasmid-related proteins; some GIs

appear to have inserted into tRNA sequences; and the GIs have slightly lower gene expression

during exponential growth [10,11]. We found that different wild type strains lack different GIs

and have different boundaries at the sites corresponding to the GIs in *S. coelicolor* M145. We

also identified DNA that the wild type strains possess and that *S. coelicolor* M145 lacks. Some of

that DNA (corresponding to GIs 8, 14, 17, 18, and 19 in *S. coelicolor* M145) might be large

segments, because we were unable to amplify the DNA by PCR [10]. The genomic islands have a

relatively uniform distribution across the chromosome of *S. coelicolor* M145, with a slight

abundance in the right chromosome arm [10,11].

A sigma factor is a specialized unit of RNA polymerase that permits the multisubunit enzyme to

initiate transcription selectively. Extracytoplasmic function (ECF) sigma factors constitute the

most abundant, smallest, and most divergent group of sigma factors [12]. Their name derives

from many of their members having roles in sensing and responding to signals generated

outside of the cell or in the cell membrane [13]. The genome of *S. coelicolor* M145 encodes 63

sigma factors, of which 49 belong to the ECF group [3]. During the early 2000's, we sought to

identify regulons of *S. coelicolor* sigma factors by overexpressing each one individually and

analyzing the response of the transcriptome by using DNA microarrays. The ECF sigma factor

encoded by SCO3450 was evaluated as part of that work. While we were unaware of this fact at

that time, SCO3450 is located in GI 6 [10].

The ECF sigma factor encoded by SCO3450 belongs to the subgroup ECF01 as classified by

Thorsten Mascher and colleagues in 2009 [14]. This subgroup includes RpoE-like sigma factors,

of which the best studied is σ^W of *Bacillus subtilis*. Members of the ECF01 subgroup are widely

distributed throughout bacterial phyla, but are absent in most actinobacteria, the phylum

which includes the genus *Streptomyces* [15]. ECF01 sigma factors have been characterized

experimentally to be involved in responses to envelope stress and the production and

detoxification of antimicrobial compounds [14].

Results and Discussion

Genomic Island 6 encodes an ECF sigma factor and the regulon of the sigma factor

Table 1 lists the five wild type *Streptomyces* strains that we studied that lack GI 6 [10].

Fig. 1 shows the boundaries of GI 6 (the left-hand columns) and the levels of transcripts, as

measured by microarrays, after artificial induction of SCO3450 in the strain *S. coelicolor* M600

growing exponentially in a liquid culture (the right-hand columns) [16]. Fifteen minutes after we

induced the sigma factor, transcripts of approximately one-fourth of the genes in GI 6 increased

in abundance. The genes numbered ~26: SCO3437 and SCO3442-3465; SCO3478 is an induced

gene located 14 kb away from the other genes. Cross-hybridization does not explain the result

for SCO3478 because the gene, which encodes a dehydrogenase, lacks sequence similarity to

other *S. coelicolor* genes. SCO3450 is located among the induced genes (Fig. 1, arrow; Fig. S4).

Together these data suggest that the ECF sigma factor and its regulon are a self-contained

transcriptional unit that can be transferred by horizontal gene transfer.

To our knowledge, the only other example of an ECF sigma factor and its regulon appearing to

be transferrable by horizontal gene transfer is the system for cobalt and nickel resistance in the

bacterium *Cupriavidus metallidurans* CH34 [18,19]. There, the plasmid pMOL28 contains six *cnr*

genes that are organized in two adjacent operons of three genes each. The gene *cnrH* encodes

the ECF sigma factor CnrH. Deletion and complementation of *cnrH* indicated that CnrH is

essential for the regulation of the *cnr* system. When CnrH was present but the periplasmic

sensor CnrX and the anti-sigma factor CnrY were absent, high-level constitutive expression was

observed for the *cnr* genes [18].

The regulon of the ECF sigma factor contains conserved putative -35 and -10 promoter regions

Byung-Kwan Cho and colleagues reported in 2016 transcription start sites of *S. coelicolor* genes

induced by 44 growth conditions [20]. Those conditions included growth in rich media and

many kinds of minimal media, growth in the liquid phase and on solid media, rapid growth to

stationary phase, and several kinds of shocks. That study did not identify transcription start

sites for the regulon of the ECF sigma factor encoded by SCO3450. This observation indicates

that the regulon is not induced by the growth conditions examined by Jeong et al. [20]. Perhaps

S. coelicolor uses the regulon for circumstances that are not replicated in the laboratory, such

as interactions in the natural environment among different microbial species.

We used the bioinformatics tool PromoterHunter [21], combined with visual inspection of

nucleotide sequences, to identify possible -35 and -10 promoter regions in the regulon of the

ECF sigma factor. The conserved sequences for the -35 and -10 promoter regions might be

AACGG and CG, respectively (Fig. 2). The promoters of 14 genes contain these sequences

exactly. The data suggest that operons might comprise SCO3445 to SCO3444, SCO3448 to

SCO3446, SCO3455 to SCO3452, and SCO3463 to SCO3465 (see Fig. S4). Alternatively, genes not

listed in Fig. 2 might have promoters with less conserved -35 and -10 regions. Note that

SCO3460 and SCO3478 do not possess the highly conserved promoter regions (see Fig. S4). In

addition, that SCO3438 has the conserved promoter regions supports the notion that this gene

belongs to the regulon of the ECF sigma factor. In Fig. 1, the lack of data for SCO3438 from all of

the microarray hybridizations indicates that its spot on the microarrays was functioning poorly.

The regulon of the ECF sigma factor encodes putative enzymes for small molecule metabolism

Recently we updated the annotations of the regulon of the ECF sigma factor by using

EnsemblBacteria [22] and UniProt BLASTS [23]. Of the approximately 26 genes in the regulon,

15 encode putative enzymes for small molecule metabolism. 11 genes encode putative

membrane and transport proteins. Four genes encode putative regulatory proteins. Four of the

26 genes fall into two categories. Only one gene lacks a putative function. Table S1 lists the

annotations of each gene in the regulon.

Enzymes encoded by the regulon of the ECF sigma factor

Table 2 lists the types of enzymes encoded by the regulon of the ECF sigma factor. The table

excludes SCO3461 and SCO3462, because they lack gene expression data (Fig. 1, right-hand

columns).

The putative enzymes fall into three general groups. One group possibly catalyzes the cleavage

of small molecules. The group includes a glycoside hydrolase, a nucleoside phosphorylase, and

a haloacid dehalogenase. A second group of enzymes might catalyze oxidation and reduction

reactions. The group includes a glutaredoxin-like protein, two oxidoreductases, and a

dehydrogenase. The third group of enzymes might catalyze transfer reactions. The group

includes two methyltransferases, perhaps a phosphatidyltransferase, and two nucleotide-

diphospho sugar transferases. A final gene, SCO3445, encodes a small membrane protein that

has 30-50% identity to small regions of seven larger proteins, a set which includes several

enzymes.

The composition of the enzymes in the regulon of the ECF sigma factor suggests that the

regulon might help to metabolize small molecules. A mixture of compounds might be being

degraded: sugars, nucleosides and nucleotides, and lipids for the purposes of energy and

biosynthesis; halogenated molecules for detoxification. Because the regulon encodes a set of

putative enzymes with diverse substrates, the small molecules might be coming from the

environment, for example, from neighboring cellular compartments or hyphae that are lysing

due to hostile conditions.

Haloacid dehalogenases (HADs) belong to a large superfamily of hydrolases with diverse

substrate specificity [24]. Type II HADs catalyze the hydrolytic dehalogenation of small L-2-

haloalkanoic acids to yield the corresponding D-2-hydroxyalkanoic acids [25]. Because many

Streptomyces bacteria produce halogenated antibiotics [26,27], the dehalogenase encoded by

SCO3446 might serve to defend *S. coelicolor* against competitors in the environment by helping

to catabolize antibiotics.

The following pairs of genes have no significant similarity, such that no cross-hybridization on

the microarrays should have occurred: SCO3452 and SCO3449, which encode the

methyltransferases; SCO3443 and SCO3460, which encode the oxidoreductases; and SCO3464

and SCO3465, which encode the nucleotide-diphospho-sugar transferases (Table 2).

Anti-sigma factors within the regulon of the ECF sigma factor

Two genes that possibly encode anti-sigma factors flank SCO3450, the gene encoding the ECF

sigma factor. The gene product of SCO3451 encodes a 103-amino-acid protein. A UniProt BLAST

shows that the protein has approximately 40% identity with varying coverages to putative anti-

sigma factors of other bacterial species, including putative transmembrane anti-sigma factors.

The Constrained Consensus TOPology (CCTOP) prediction server [28] predicts with a reliability

of 95.9069 that SCO3451 encodes a protein with one transmembrane segment (Fig. S5). The

gene product of SCO3451 also contains a putative zinc-finger found in some anti-sigma factor

proteins (Fig. S6). This zinc finger domain overlaps with the predicted transmembrane domain.

SCO3449 encodes a 106-amino-acid protein. A UniProt BLAST shows that the protein has

approximately 40% identity with coverages around 50% to putative anti-sigma factors in other

bacterial species, including putative transmembrane anti-sigma factors. However, according to

CCTOP, the protein does not contain a transmembrane domain. Like the putative anti-sigma

factor encoded by SCO3451, the putative anti-sigma factor encoded by SCO3449 contains a

possible zinc-finger found in some anti-sigma factor proteins (Fig. S7).

Microarray data are lacking for the expression of SCO3449 (Fig. 1, right-hand columns), such

that inferences about the role of this gene in comparison to those of the induced genes should

be made with caution. However, because microarray data are available for this gene in the

hybridizations that identified the genomic islands (Fig. 1, left-hand columns), the spot on the

microarrays that represented this gene was likely intact. It is possible that a low abundance of

transcripts of SCO3449 in both the reference and experimental samples of mRNA produced

poor signals from the microarray spot of the gene.

Membrane and transport proteins encoded by the regulon of the ECF sigma factor

Four contiguous genes in the regulon of the ECF sigma factor, SCO3453 to SCO3456, encode

proteins that likely constitute an ABC transporter system. UniProt BLASTS show that all of the

proteins have 45-50% identity with coverages greater than 95% to homologs with putative

functions in spermidine and putrescine transport in other bacterial species. In particular, the

species include *Geodermatophilus* and *Wenxinia*. Spermidine is a polyamine involved in cellular

metabolism that can be used to stimulate RNA polymerase. Putrescine attacks S-adenosyl

methionine and converts it to spermidine [29].

The putative transporters Sco3454 and Sco3455 have 34% identity with each other, the

coverage being 44% between the C-terminal portions of the proteins.

Genomic Island 6 consists of four segments with transposases at their boundaries

In addition to the regulon of the ECF sigma factor, we updated the annotations of the other

genes of Genomic Island 6 by using EnsemblBacteria and UniProt BLASTS (Table S2). In the

table, the location of the regulon is denoted by the words "REGULON HERE."

The length of GI 6 is 108 kb. The genomic island consists of four segments of DNA bounded by

transposases (Table S2, pink color). Three of the four segments have coherent putative

functions: the oxidation and reduction of copper (Table S2, blue color); a characterized agarase

encoded by *dagA* [30,31]; and the utilization of sugars (Table S2, yellow color). Included in the

latter segment are a putative *lacI*-family transcriptional regulator and a putative β-

galactosidase. A large fourth segment encodes putative enzymes and many hypothetical

proteins.

The coherent functions of at least three of the four segments of GI 6 indicate that significant

portions of the island might be "active." If so, the regulon of the ECF sigma factor is likely active

as well. While a nucleotide blast of the regulon of the ECF sigma factor yielded no similar

segments of DNA in other sequenced organisms, the value to *S. coelicolor* of the regulon for as-

yet unknown reasons is indicated by the presence of the regulon on a genomic island. It would

be interesting to determine how common the regulon is among natural populations of

microbes and whether the regulon moves easily between hosts.

Materials and Methods

Construction of a S. coelicolor *strain that overexpresses SCO3450*

SCO3450, the gene which encodes the ECF sigma factor harbored by GI 6, was amplified by PCR

and cloned into the conjugative plasmid pIJ6902 [32] by using the restriction sites *Nde*I and

*Bgl*II. These restriction sites placed the gene immediately downstream of the thiostrepton-

inducible promoter *tipA*p. The resulting plasmid was transformed into the methylation-

deficient strain ET12567 with a non-transmissable helper plasmid, pUZ8002, and conjugated

into *S. coelicolor* M600 as described by Kieser et al. [33]. The exconjugants were selected by an

overlay of 50 μg/mL of apramycin.

Time course of the overexpression strain

Supplemented minimal medium (SMM) was inoculated with approximately 5×10^7 spores/mL

of a spore stock. The culture was grown at 30°C to early exponential phase, which

corresponded to a cell density of $OD_{450} \sim 0.5$. A reference sample, designated "0 min," was

harvested. To induce transcription of SCO3450, thiostrepton was added to the culture to a final

concentration of 30 μg/mL. Samples of the culture were harvested 15, 30, 45, 60, and 90

minutes after induction of SCO3450.

Extraction of total RNA from the overexpression strain

Samples of cells from liquid cultures were recovered by filtration on Whatman filter paper (15-

20 mm diameter; catalog #1002 055). RNA was isolated using the modified Kirby mix protocol

as described previously [34] with the following modifications: Harvested volumes ranged

between 5 mL and 20 mL, depending on the cell densities. RNA samples were treated only once

with DNase I (50-70 units; RNase-free, Invitrogen) for 15 minutes at room temperature.

DNA microarray experiments for the overexpression strain

Samples of cDNA were synthesized from total RNA as described previously [34]. cDNA of the

reference sample was labeled with Cy3-CTP. cDNAs of the samples isolated at subsequent time

points were labeled with Cy5-CTP. The cDNAs were hybridized to microarrays as described

previously [34].

Identification of -35 and -10 conserved promoter regions in the regulon of the ECF sigma factor

The tool PromoterHunter [21] was used to examine DNA upstream of the genes in the regulon

of the ECF sigma factor. Weight matrices corresponding to AAC for the -35 promoter region and

CG for the -10 promoter region were used, because they reflect the conserved promoter

regions of the subgroup ECF01 of ECF sigma factors [14]. A global G + C content of 72% was

used. The space between the -35 and -10 regions was specified to be between 17 to 21 base

pairs. Initially DNA segments of 300 base pairs were examined for isolated genes and genes

located at the beginning of likely operons. Visual inspection of sequences returned by

PromoterHunter identified AACGG and CG as possible consensus sequences for the -35 and -10

promoter regions, respectively. PromoterHunter was used to search for these conserved

sequences within DNA segments of 1000 base pairs upstream of all of the genes in the regulon,

in order to ensure that all instances of the sequences were identified.

Acknowledgements

We thank Thomas M. Privalsky for informing us about bioinformatic tools and Mark J. Buttner

and David A. Hopwood for discussions about the data.

Author Contributions

C.M.K. conceived the experiments. D.W. designed the experiments that compared the genomes

of the six *Streptomyces* strains and performed those experiments with J.A.V., M.L.H., and K.G.P.

N.K. designed the experiment overexpressing the ECF sigma factor and performed the

experiment with S.A.G. C.M.K. conducted the recent bioinformatic analyses and wrote the

manuscript.

References

1. Hopwood DA. 2007. Streptomyces in nature and medicine: the antibiotic makers. Oxford

318   University Press, New York, NY.

2.   Kutzner HJ, Waksman SA. 1959. Streptomyces coelicolor Mueller and Streptomyces

violaceoruber Waksman and Curtis, two distinctly different organisms. J Bacteriol

78:528—538.

3.   Bentley SD, Chater KF, Cerdeño-Tárraga AM, Challis GL, Thomson NR, James KD, Harris

DE, Quail MA, Kieser H, Harper D, Bateman A, Brown S, Chandra G, Chen CW, Collins M,

Cronin A, Fraser A, Goble A, Hidalgo J, Hornsby T, Howarth S, Huang CH, Kieser T, Larke

L, Murphy L, Oliver K, O'Neil S, Rabbinowitsch E, Rajandream MA, Rutherford K, Rutter

S, Seeger K, Saunders D, Sharp S, Squares R, Squares S, Taylor K, Warren T, Wietzorrek A,

Woodward J, Barrell BG, Parkhill J, Hopwood DA. 2002. Complete genome sequence of

the model actinomycete *Streptomyces coelicolor* A3(2). Nature 417:141—147.

4.   Hopwood DA. 1999. Forty years of genetics with *Streptomyces*: from *in vivo* through *in

vitro* to *in silico*. Microbiology 145:2183—2202.

5.   Sermonti G, Spada-Sermonti I. 1955. Genetic recombination in *Streptomyces*. Nature

176:121.

6.   Bradley SG. 1960. Reciprocal crosses in Streptomyces coelicolor. Genetics 45:613—619.

7.   Lomovskaya ND, Mkrtumian NM, Gostimskaya NL, Danilenko VN. 1972. Characterization

of temperate actinophage phi C31 isolated from Streptomyces coelicolor A3(2). J Virol

9:258—262.

8.   Kieser T, Hopwood DA, Wright HM, Thompson CJ. 1982. pIJ101, a multi-copy broad host-

range Streptomyces plasmid: functional analysis and development of DNA cloning

vectors. Mol Gen Genet 185:223—228.

9. Aguilar A, Hopwood DA. 1982. Determination of methylenomycin A synthesis by the

pSV1 plasmid from *Streptomyces violaceus-ruber* SANK 95570. J Gen Microbiol

128:1893—901.

10. Weaver D. 2005. PhD thesis. Stanford University, Stanford, CA. Genome plasticity in

Streptomyces coelicolor.

11. Jayapal KP, Lian W, Glod F, Sherman DH, Hu WS. 2007. Comparative genomic

hybridizations reveal absence of large *Streptomyces coelicolor* genomic islands in

Streptomyces lividans. BMC Genomics 8:229.

12. Feklístov A, Sharon BD, Darst SA, Gross CA. 2014. Bacterial sigma factors: a historical,

structural, and genomic perspective. Annu Rev Microbiol 68:357—376.

13. Lonetto MA, Brown KL, Rudd KE, Buttner MJ. 1994. Analysis of the Streptomyces

coelicolor sigE gene reveals the existence of a subfamily of eubacterial RNA polymerase

sigma factors involved in the regulation of extracytoplasmic functions. Proc Natl Acad

Sci U S A 91:7573—7577.

14. Staroń A, Sofia HJ, Dietrich S, Ulrich LE, Liesegang H, Mascher T. 2009. The third pillar of

bacterial signal transduction: classification of the extracytoplasmic function (ECF) σ

factor protein family. Mol Microbiol 74:557—581.

15. ECFfinder. http://ecf.g2l.bio.uni-goettingen.de:8080/ECFfinder/.

16. Takano E, White J, Thompson CJ, Bibb MJ. 1995. Construction of thiostrepton-inducible,

high-copy-number expression vectors for use in *Streptomyces* spp. Gene 166:133—137.

17. Huang J, Lih CJ, Pan KH, Cohen SN. 2001. Global analysis of growth phase responsive

gene expression and regulation of antibiotic biosynthetic pathways in *Streptomyces*

coelicolor using DNA microarrays. Genes Dev 15:3183—3192.

18. Grass G, Grosse C, Nies DH. 2000. Regulation of the *cnr* cobalt and nickel resistance

determinant from *Ralstonia* sp. strain CH34. J Bacteriol 182:1390—1398.

19. Tibazarwa C, Wuertz S, Mergeay M, Wyns L, van der Lelie D. 2000. Regulation of the *cnr*

cobalt and nickel resistance determinant of *Ralstonia eutropha* (*Alcaligenes eutrophus*)

CH34. J Bacteriol 182:1399—1409.

20. Jeong Y, Kim JN, Kim MW, Bucca G, Cho S, Yoon YJ, Kim BG, Roe JH, Kim SC, Smith CP,

Cho BK. 2016. The dynamic transcriptional and translational landscape of the model

antibiotic producer *Streptomyces coelicolor* A3(2). Nat Commun 7:11605.

21. Klucar L, Stano M, Hajduk M. 2010. phiSITE: database of gene regulation in

bacteriophages. Nucleic Acids Res 38:D366—D370.

22. Kersey PJ, Allen JE, Allot A, Barba M, Boddu S, Bolt BJ, Carvalho-Silva D, Christensen M,

Davis P, Grabmueller C, Kumar N, Liu Z, Maurel T, Moore B, McDowall MD, Maheswari

U, Naamati G, Newman V, Ong CK, Paulini M, Pedro H, Perry E, Russell M, Sparrow H,

Tapanari E, Taylor K, Vullo A, Williams G, Zadissia A, Olson A, Stein J, Wei S, Tello-Ruiz M,

Ware D, Luciani A, Potter S, Finn RD, Urban M, Hammond-Kosack KE, Bolser DM, De

Silva N, Howe KL, Langridge N, Maslen G, Staines DM, Yates A. 2018. Ensembl Genomes

2018: an integrated omics infrastructure for non-vertebrate species. Nucleic Acids Res

46:D802—D808.

23. The UniProt Consortium. 2017. UniProt: the universal protein knowledgebase. Nucleic

Acids Res 45:D158—D169.

24. Koonin EV, Tatusov RL. 1994. Computer analysis of bacterial haloacid dehalogenases

defines a large superfamily of hydrolases with diverse specificity: application of an

iterative approach to database search. J Mol Biol 244:125—132.

25. Janssen DB, Oppentocht JE, Poelarends GJ. 2001. Microbial dehalogenation. Curr Opin

Biotechnol 12:254—258.

26. van Pée KH. 1996. Biosynthesis of halogenated metabolites by bacteria. Annu Rev

Microbiol 50:375—399.

27. Rezanka T, Spizek J. 2003. Halogen-containing antibiotics from Streptomycetes, p 309—

353. *In* Atta-ur-Rahman (ed), Studies in natural products chemistry, 1st ed, vol 29.

Elsevier Science.

28. Dobson L, Reményi R, Tusnády GE. 2015. CCTOP: a Consensus Constrained TOPology

prediction web server. Nucleic Acids Res 43:W408—W412.

29. Finn RD, Attwood TK, Babbitt PC, Bateman A, Bork P, Bridge AJ, Chang HY, Dosztányi Z,

El-Gebali S, Fraser M, Gough J, Haft D, Holliday GL, Huang H, Huang X, Letunic I, Lopez R,

Lu S, Marchler-Bauer A, Mi H, Mistry J, Natale DA, Necci M, Nuka G, Orengo CA, Park Y,

Pesseat S, Piovesan D, Potter SC, Rawlings ND, Redaschi N, Richardson L, Rivoire C,

Sangrador-Vegas A, Sigrist C, Sillitoe I, Smithers B, Squizzato S, Sutton G, Thanki N,

Thomas PD, Tosatto SC, Wu CH, Xenarios I, Yeh LS, Young SY, Mitchell AL. 2017. InterPro

in 2017--beyond protein family and domain annotations. Nucleic Acids Res 45:D190—

D199.

30. Temuujin U, Chi WJ, Lee SY, Chang YK, Hong SK. 2011. Overexpression and biochemical

characterization of DagA from *Streptomyces coelicolor* A3(2): an endo-type β-agarase

producing neoagarotetraose and neoagarohexaose. Appl Microbiol Biotechnol 92:749—

759.

31. Bibb MJ, Jones GH, Joseph R, Buttner MJ, Ward JM. 1987. The agarase gene (*dagA*) of

Streptomyces coelicolor A3(2): affinity purification and characterization of the cloned

gene product. J Gen Microbiol 133:2089—2096.

32. Huang J, Shi J, Molle V, Sohlberg B, Weaver D, Bibb MJ, Karoonuthaisiri N, Lih CJ, Kao

CM, Buttner MJ, Cohen SN. 2005. Cross-regulation among disparate antibiotic

biosynthetic pathways of *Streptomyces coelicolor*. Mol Microbiol 58:1276—1287.

33. Kieser T, Bibb MJ, Buttner, MJ, Chater KF, Hopwood DA. 2000. Practical *Streptomyces*

genetics. John Innes Foundation.

34. Weaver D, Karoonuthaisiri N, Tsai HH, Huang CH, Ho ML, Gai S, Patel KG, Huang J, Cohen

SN, Hopwood DA, Chen CW, Kao CM. 2004. Genome plasticity in *Streptomyces*:

identification of 1 Mb TIRs in the *S. coelicolor* A3(2) chromosome. Mol Microbiol

51:1535—1550.

FIG 1 Genomic Island 6 and the regulon of the ECF sigma factor encoded by SCO3450. GI 6 is

108 kb in length. The arrow indicates the location of SCO3450, which encodes a putative ECF

sigma factor. See Table 1 for a list of the strains used to identify GI 6. To obtain the data in the

figure, we used DNA microarrays that contained the 7825 predicted genes in the chromosome

of *S. coelicolor* M145 [17]. For the left-hand columns, the maximum intensity of green

425 corresponds to a 3-fold difference in signal on the microarrays for DNA in *S. coelicolor* M145

426 relative to DNA in the wild type strains. SCO3450 was overexpressed in an exponentially

427 growing liquid culture (SMM) of a *S. coelicolor* M600 derivative, through the use of a

428 thiostrepton-inducible promoter [16]. RNA was harvested from the culture immediately prior to

429 the addition of thiostrepton and every 15 to 30 minutes afterwards for 90 minutes. cDNA of the

430 initial RNA sample (OD_{450} ~ 0.5) was labeled with the green fluorescent dye Cy3. cDNA of RNA

431 isolated from subsequent time points was labeled with the red fluorescent dye Cy5. Yellow

432 microarray spots for a particular time point represented genes with transcript levels equal to

433 the levels of the initial time point. Red and green microarray spots represented genes induced

434 and repressed, respectively, after the addition of thiostrepton. The yellow color is shown here

435 as black for clarity. For the right-hand columns, the maximum intensity of red corresponds to 8-

436 fold induction of a given gene at a particular time relative to the beginning of the time course. A

437 control experiment with a M600 derivative that lacks SCO3450 in the induction plasmid

438 identified genes induced by thiostrepton. Labels on the right indicate ORF numbers and genes

439 with names.

441 **FIG 2** Alignment of putative promoters in the regulon of the ECF sigma factor encoded by

442 SCO3450. Red letters indicate the putative -35 and -10 promoter regions. On the right-hand

443 side, the numbers in parentheses indicate the distance between each -10 promoter region and

444 the start codon of the respective gene.

447   **Table 1. Strains that revealed the genomic islands of _S. coelicolor_ M145.**

Strain	Comment	Ref.
M145	_S. coelicolor_ sequenced "reference" strain, SCP1- SCP2-	[3]
1152	_S. coelicolor_ Sermonti's SE1	[5]
1153	_S. coelicolor_ Bradley's S199 strain	[6]
1326	_S. lividans_ 66	[7]
2896	_S. lividans_ ISP5434-, the strain from which pIJ101 was isolated	[8]
3034	_S. violaceroruber_ SANK95570, harbors pSV1, which encodes methylenomycin biosynthetic genes	[9]

450   **Table 2. Types of enzymes encoded by the regulon of the ECF sigma factor.**

Type	Gene	AA	Putative Function	Other feature
CLEAVAGE				
Glycoside hydrolase	SCO3444	617	glycoside hydrolase family 15/phosphorylase b kinase regulatory chain family; six-hairpin glycosidase	
Nucleoside phosphorylase	SCO3463	262	nucleoside phosphorylase domain	
Dehalogenase	SCO3446	225	haloacid dehydrogenase (HAD)-like domain	
REDOX				
Glutaredoxin	SCO3442	114	glutaredoxin (DNA synthesis?)	
Oxidoreductase	SCO3443	454	pyridine nucleotide-disulphide oxidoreductase; dihydrolipoyl dehydrogenase?	
	SCO3460	505	pyridine nucleotide-disulphide oxidoreductase; dihydrolipoamide dehydrogenase?	
Dehydrogenase	SCO3478	344	D-isomer specific 2-hydroxyacid dehydrogenase	
TRANSFER				
Methyltransferase	SCO3452	359	S-adenosyl-L-methionine-dependent methyltransferase	
	SCO3459	287	S-adenosyl-L-methionine-dependent methyltransferase	
Phosphatidyltransferase?	SCO3457	205	CDP-alcohol phosphatidyltransferase; YnjF?	transmembrane
Nucleotide-diphospho-sugar transferase	SCO3464	210	nucleotide-diphospho-sugar transferases; transferase 1, rSAM/selenodomain-associated	
	SCO3465	236	nucleotide-diphospho-sugar transferase; transferase 2, rSAM/selenodomain-associated	
MISC				
Enzyme?	SCO3445	55	low to moderate homology to small regions of seven larger proteins, which include enzymes	membrane

AA = amino acids

Legends for Supplemental Material

FIG S1 Genomic Islands 1 to 10 of *S. coelicolor* M145. Only the genomic islands are shown and

not the entire chromosome. Table 1 lists the strains used. To obtain these data, we used DNA

microarrays that contained the 7825 predicted genes in the chromosome of *S. coelicolor* M145

[17]. Genomic DNA from *S. coelicolor* M145 was used as the reference sample and labeled with

the green fluorescent dye Cy3. Genomic DNA samples from the five wild type strains each were

labeled with the red fluorescent dye Cy5. The labeled DNA of each wild type strain was mixed

with the labeled DNA of *S. coelicolor* M145 and hybridized to a microarray. Yellow spots on the

microarrays represented genes at equal copy numbers between *S. coelicolor* M145 and the

other strains. Green spots represented genes present in *S. coelicolor* M145 but absent in the

other strains. Red spots represented genes at higher copy numbers in the strains relative to *S.*

coelicolor M145. The yellow color is shown here as black for clarity. The maximum intensity of

green corresponds to a 3-fold difference in signal on the microarrays for DNA in *S. coelicolor*

M145 relative to DNA in the wild type strains. The green bars to the left of the GIs designate

horizontally transferred genes in the genome sequence of *S. coelicolor* M145 that were

predicted by Bentley et al. [3]. Regions of GIs not predicted by Bentley et al. lack a bar. Labels

on the right indicate ORF numbers and genes with names.

FIG S2 Genomic Islands 11 to 18 of *S. coelicolor* M145. See the text of Fig. S1.

FIG S3 Genomic Islands 19 to 22 of *S. coelicolor* M145. See the text of Fig. S1.

FIG S4 The region of the *S. coelicolor* M145 chromosome that contains the regulon of the ECF

sigma factor. The figure was obtained from EnsemblBacteria [22]. SCO3478 is not shown.

FIG S5 Result from CCTOP for the putative anti-sigma factor encoded by SCO3451. The putative

anti-sigma factor is predicted to contain one transmembrane domain. The data were obtained

from the Constrained Consensus TOPology (CCTOP) prediction server [28].

FIG S6 Predicted zinc finger in the putative anti-sigma factor encoded by SCO3451. The data

were obtained from UniProt [23].

FIG S7 Predicted zinc finger in the putative anti-sigma factor encoded by SCO3449. The data

were obtained from UniProt [23].

Table S1. Annotations of the regulon of the ECF Sigma Factor encoded by SCO3450.

Annotations were obtained from EnsemblBacteria [22] and UniProt [23]. For the UniProt

BLASTs, parentheses denote genera and percent identities of similar proteins. Red text denotes

putative enzymes. Blue text denotes proteins with homology to small regions of larger proteins.

Table S2. Annotations of Genomic Island 6, excluding the regulon of the ECF sigma factor

encoded by SCO3450. Annotations were obtained from EnsemblBacteria [22] and UniProt [23].

For the UniProt BLASTs, parentheses denote genera and percent identities of similar proteins.

Red text denotes putative enzymes. Blue text denotes proteins with homology to small regions

of larger proteins. See the text for additional details.

FIG 1 Genomic Island 6 and the regulon of the ECF sigma factor encoded by SCO3450.

```
SCO3437-1  CGTTGCCGACAACGGGATGATGATCCTCGG--CGACCTCAACCA  (694 bp)
SCO3437-2  CGTCGCCGACAACGGGATGATGATCCTCGG--CGACCTCAACCA  (283 bp)
SCO3438    CAGGTCCCGCAACGGCGCACCCTCGTGGGC--CGGTGCCCGCGG  (178 bp)
SCO3442    GTGGTGGTGGAACGGCACCTGGCTGCCGCAGTCCGTGCCGCCCG  (407 bp)
SCO3443    CGACCAGGCCAACGGCGCGCTACCTCGCCACGACGCCGACCCGG  (158 bp)
SCO3445    ACGGCCTGGAAACGGCGTGCTCCTGAGCCGC-CGTGCGTCAATG  (  7 bp)
SCO3448    TCGATGGCGAAACGGACGAGGTCACCGCACGCCGGGTCGCGGCC  (295 bp)
SCO3451    TTCCCGTCGAAACGGTCGATGGCCCGGTAGGCGCGCAGCAAGGT  (312 bp)
SCO3455-1  GGTGTGGGTCAACGGAGAGAACTTCCGCAC--CGGAAAGCAGGC  (883 bp)
SCO3455-2  CTGGACCAGGAACGGCTGCCCGAGCAGTGG--CGAGAGAAGTTC  (150 bp)
SCO3456    GTTTCCGCGCAACGGATGTTCGCGGGACCGTCGCGCGGGAAACA  ( 50 bp)
SCO3457    GTTGCGCGGAAACGGAACACGGCCGGGTGCT-CGCAGGCTTCCC  ( 58 bp)
SCO3458    GCGTGTCCGAAACGGAATGCGTGACCAGCC--CGGCAGGGTTCC  (844 bp)
SCO3459    GTCGCTGGTCAACGGCCCGCTCTCGACCAGGCCGGGGATCGGCA  (611 bp)
SCO3462    CCGAGGCCGTAACGGCCCTCTGGTTTCTCGCCGGCCCCGCTCC   (276 bp)
SCO3463    CCACCCCGAGAACGGAACCCTCGTCCCCGCCTGCGTACAGCACG  (145 bp)
                     -35                    -10
```

503 **FIG 2** Alignment of putative promoters in the regulon of the ECF sigma factor encoded by

504 SCO3450.