Planned missing data design: stronger inferences, increased research efficiency and improved animal welfare in ecology and evolution

Daniel W.A. Noble[1,*] and Shinichi Nakagawa[1, 2,*]

[1] Ecology and Evolution Research Centre, School of Biological, Earth and Environmental Sciences, The University of New South Wales, Kensington NSW 2052, Sydney
[2] Diabetes and Metabolism Division, Garvan Institute of Medical Research, 384 Victoria Street, Darlinghurst, Sydney, NSW 2010, Australia

* To whom correspondence should be addressed:
Daniel Noble: daniel.wa.noble@gmail.com
Shinichi Nakagawa: s.nakagawa@unsw.edu.au

Abstract:

1. Ecological and evolutionary research questions are increasingly requiring the integration of research fields along with larger datasets to address fundamental local and global scale problems. Unfortunately, these agendas are often in conflict with limited funding and a need to balance animal welfare concerns.

2. Planned missing data design (PMDD), where data are randomly and deliberately missed during data collection, is a simple and effective strategy to working under greater research constraints while ensuring experiments have sufficient power to address fundamental research questions. Here, we review how PMDD can be incorporated into existing experimental designs by discussing alternative design approaches and evaluating how data imputation procedures work under PMDD situations.

3. Using realistic examples and simulations of multilevel data we show how a variety of research questions and data types, common in ecology and evolution, can be aided by using a PMDD with data imputation procedures. More specifically, we show how PMDD can improve statistical power in detecting effects of interest even with high levels (50%) of missing data and moderate sample sizes. We also provide examples of how PMDD can facilitate improved animal welfare and potentially alleviate research costs and constraints that would make endeavours for integrative research challenging.

4. Planned missing data designs are still in their infancy and we discuss some of the difficulties in their implementation and provide tentative solutions. Nonetheless, data imputation procedures are becoming more sophisticated and more easily implemented and it is likely that PMDD will be an effective and powerful tool for a wide range of experimental designs, data types and problems in ecology and evolution.

Keywords: data augmentation, multiple imputation, personality, quantitative genetics, mixed effects models, hierarchical models, multilevel modelling, refinement, reduction, multiple working hypotheses.

Introduction

Missing data is a widespread problem in ecological and evolutionary research (Nakagawa & Freckleton 2010; Nakagawa & Freckleton 2011; Ellington *et al.* 2015; Nakagawa 2017), often resulting in the exclusion of a substantial amount of data. This contributes to a major reduction in statistical power and, if the nature of 'missingness' is not considered carefully, leads to biased parameter estimates (Enders 2001b; Graham 2009; Nakagawa & Freckleton 2010; Little *et al.* 2013). Theoretical frameworks for dealing with missing data, however, have received substantial attention and missing data theory is now a well-developed field of research grounded in solid statistical theory (Graham, Hofer & MacKinnon 1996; Enders 2001a; Little & Rubin 2002; Graham 2003; Graham 2009; van Buuren 2012; Little *et al.* 2013). Nonetheless, while social scientists have been at the forefront of applied missing data techniques, ecologists and evolutionary biologists have lagged behind (Nakagawa & Freckleton 2010).

Missing data is traditionally viewed by ecologists and evolutionary biologists with a sense of disdain and annoyance. But, what if including missing data in analyses could be advantageous? Indeed, social scientists have taken a rather different stance to missing data – instead embracing its power to help address fundamental research questions (Graham *et al.* 2006). Planned missing data design (PMDD) is an approach that involves deliberately planning to 'miss' data as an integral part of an experiment. In other words, deliberately not collecting data on certain variables or experimental subjects. While this seems like an odd thing to do, if missing data in the variables of interest is completely random, or can be made random, existing statistical frameworks are known to do an excellent job at estimating parameters and standard errors (Schafer & Graham 2002; van Buuren 2012). The PMDD approach comes with a substantial number of benefits that have been largely ignored by ecologists and evolutionary biologists.

Here, we argue that PMDD has the potential to expand the scope, reduce research costs and improve animal welfare, facilitating higher impact research with greater power. We begin our discussion by briefly introducing missing data theory and then describe a few core statistical tools that can be used to impute / augment (i.e., 'fill in') missing data. Using simulations, we show that, when missing data is 'completely' random, existing data imputation techniques can improve the estimation of parameters and their standard errors – even with hierarchically structured data that is common in ecological and evolutionary research (Enders, Mistler & Keller 2016; Quartagno & Carpenter 2016; Resche-Rigon &

79   White 2016). We then describe PMDD, overviewing some of the different experimental

80   approaches that can be implemented, what they involve and important design considerations.

81   Following from this discussion, we overview the important benefits of utilizing a PMDD and

82   end with a discussion on some of the challenges and limitations to their use – providing

83   suggestions for how these can be rectified.

A brief introduction to missing data theory

Missing data patterns can generally be classified as falling into one of three different types –

based on the different mechanisms generating missing data – missing completely at random

(MCAR), missing at random (MAR) and missing not at random (MNAR) (Rubin 1976; Little

& Rubin 2002; Graham 2009; Nakagawa & Freckleton 2010; van Buuren 2012; Nakagawa

2017). The distinction between these three missing data mechanisms is important to

understand when implementing a PMDD, which we will introduce below. Missing data

(either in response or predictor variables) are considered to be MCAR when missingness is

random with respect to both observed and unobserved (i.e., not collected in the study)

variables (Enders 2001b; Nakagawa 2017). In other words, the observed data is simply a

random sub-sample of complete data (Enders 2001b). In contrast, missing data are considered

MAR when the missing values in a dataset depend on observed values of other variables in

the dataset (Enders 2001b; Graham 2009). For example, if we were interested in

understanding the correlation between survival to 1 year and mass at 6 months, we would

find that individuals that die before 6 months are missing data on mass, but missing data on

mass is correlated with their lifespan, which is known. Missing not at random (MNAR),

however, occurs when missing values depend on unobserved variables that have not been

quantified in the study, or on the variable itself after accounting for the observed data. For

example, we may be missing behavioural data on small sized animals within a population

because they tend to be 'shy' and difficult to capture (e.g, Biro & Dingemanse 2009), in

which case we would be missing both behavioural and morphological data on a non-random

sample of the population. Under these situations, dealing with missing data is difficult

(possibly even impossible) because statistical techniques for estimating parameters when data

are MNAR are difficult to implement given the need to explicitly model the process of

missingness (Schafer & Graham 2002).

Missing data mechanisms have different consequences on statistical results when

missing data is excluded prior to analysis, as is often the case (i.e., referred to as 'complete

case', 'pairwise deletion' or 'listwise deletion'). While MCAR results in a loss of power

when missing data is excluded from an analysis, it does not bias parameter estimates (Enders
2001b; Schafer & Graham 2002; Graham 2009; Nakagawa & Freckleton 2010). In contrast,
when missing data is MAR or MNAR, excluding missing data will result in both a loss of
power and biased parameter estimates (sometimes severly so; Enders 2001b; Schafer &
Graham 2002; Graham 2009; Nakagawa & Freckleton 2010). To better appreciate the impact
missing data can have on sample size (and thus statistical power), assume that we have 10
variables, each containing 5% missing data, and a total complete dataset of $n = 1000$. If we
used all variables in a statistical model we may need to exclude as much as 500 observations,
resulting in a substantial decrease in power and severely compromising our ability to detect
significant effects (see '*Improving power of experimental designs*' for more on this issue).
Statistical techniques for dealing with missing data rely on the assumption of missing data
being MCAR or MAR, and if this assumption is met, then both power and bias in parameter
estimates can be recovered (Enders 2001b; Schafer & Graham 2002; Nakagawa & Freckleton
2010; van Buuren 2012; Ellington *et al.* 2015; Nakagawa 2017).

Statistical procedures for dealing with missing data

Planned missing data design hinges on the ability of researchers to make use of statistical
procedures for handling missing data (Little & Rubin 2002; Graham *et al.* 2006; Enders
2010). It is therefore pertinent that we briefly review existing missing data procedures and
provide some guidance on their implementation when data has been collected using a PMDD.
We do not discuss these topics in great depth as there are a number of important, accessible
reviews and books on these subjects already, which we direct the reader to for more details
(Schafer 1997; Enders 2001b; Little & Rubin 2002; McKnight *et al.* 2007; Allison 2012; van
Buuren 2012; Nakagawa 2017).
135   As mentioned above, imputation methods fall under two broad categories and we
follow the general categorization of McKnight et al. (2007) in classifying them in to those
implementing model-based (MB) techniques and those using multiple imputation (MI) with
the help of Rubin's rules (Rubin 1987; Enders 2010). While we acknowledge that these two
broad categories have some overlap, they generally capture the major differences in the types
of imputation procedures that can be applied. Model-based procedures incorporate both
observed and missing data into a single joint modelling approach that is constrained by the
underlying probability distribution and proceeds through the following steps: 1) the
parameters of a model are estimated using observed data; 2) parameters estimated in step 1
are then used to augment missing data and 3) model parameters are re-assessed conditional

on both the observed and imputed data (Figure 1A; Nakagawa 2017). These steps are re-
iterated until the model converges (i.e., maximum likelihood or stable posterior distribution if
using Bayesian methods) (Figure 1A). Model-based approaches are advantageous in that they
are fast, easily implemented (under the assumption of multivariate normality) and result in
robust parameter estimates and standard errors (McKnight *et al.* 2007).

In contrast, multiple imputation proceeds by generating a set of m complete datasets
where missing data is imputed using variables of interest. Usually $m = 40$–50 imputed
datasets perform well under a variety of situations (Nakagawa & de Villemereuil 2015;
Nakagawa 2017). These m datasets can then be analysed normally (i.e., as if a complete
dataset existed) and the results (i.e., parameter estimates and standard errors) pooled across
the m datasets (Figure 1B; Schafer 1997; Schafer & Olsen 1998; Little & Rubin 2002; van
Buuren & Groothuis-Oudshoorn 2011; van Buuren 2012; Nakagawa 2017). Multiple
imputation provides a number of advantages over MB approaches. First, it is extremely
flexible, easily accommodating different distributions (i.e., Bernoulli, Poisson etc.), variables
and model types if needed. Second, since MI creates m complete datasets it allows one to
separate out the imputation step from the analysis step. In other words, we can have a set of
auxiliary variables (see '*Auxiliary variables to aid in imputation*' below) that are used to
impute missing values and then subsequently use only the variables of biological interest to
run the analysis on m imputed datasets. This is particularly advantageous because including
unnecessary variables in MB procedures can complicate the interpretation of model results
(McKnight *et al.* 2007; Enders 2010). Lastly, MI procedures account better for imputation
uncertainty as variation in parameter estimates across data sets can be explicitly incorporated
in pooled estimates, protecting against type I errors (McKnight *et al.* 2007). Additionally, the
effect of missing data on analysis results (i.e. efficiency) can be explicitly quantified and
presented by deriving statistics summarising the variability in parameter estimates across
imputed datasets (McKnight *et al.* 2007). Given the flexibility, ease of implementation and
their general tendency to produce robust parameter estimates, it is unsurprising that Rubin
(1996) recommends MI over MB procedures.

Auxiliary variables to aid imputation
Auxiliary variables are variables that are not necessarily of interest with respect to the
biological question at hand, but that are correlated with other variables, or missing data itself,
within the dataset (Collins, Schafer & Kam 2001; Graham 2003). Including auxiliary
variables, especially ones that are expected to be predictive of missing data, have been shown

to improve the accuracy and stability of estimates and to reduce their standard error (Enders
2010; Allison 2012; von Hippel & Lynch 2013). The best auxiliary variables are those that
are easy and cheap to collect and that are strongly correlated with a number of other variables
within the data set (Collins, Schafer & Kam 2001; Graham 2003; von Hippel & Lynch 2013).
Collins et al. (2001) have shown that auxiliary variables can be particularly useful when the
missing data is in the response variable, when they change the missing data mechanism from
MNAR to MAR and when the correlation between auxiliary variables and the response is
high ($r = 0.9$). Adding even just 2–3 auxiliary variables can improve imputation procedures
and for the most part, an inclusive analysis strategy where a large number of auxiliary
variables are included in the analysis is recommended (Enders 2010 p.g. 128). However, this
procedure can be slightly more complex than this in practice. Hardt *et al.* (2012) show that
the inclusion of too many (> 10) can start to lead to a downward bias in regression
coefficients and a decrease in precision. In addition, auxiliary variables will have little impact
when the correlations between variables in the dataset are low ($r = 0.10$) (Hardt, Herke &
Leonhart 2012). While the number of auxiliary variables will depend on the specific study
questions being asked, we recommend including auxiliary variables with moderate to high
correlations (0.4–0.8). These will be particularly important when using imputation procedures
where unplanned missing data exits (see section "*Unplanned missing data*" below) to change
missingness from MNAR to MAR.

Experiments in ecology and evolution often collect variables that are not necessarily
of interest, but can be used as auxiliary variables. These variables can include body
dimensions, sex, age or even spatial data. These types of auxiliary variables can be included
in imputation procedures (e.g., MI) with unplanned missing data to ensure that the MAR
assumption is met, but then discarded when testing the biological questions and hypotheses
of interest (Graham 2003). Considering these variables more carefully with respect to their
potential correlations with other variables, and with missing data itself, is an important aspect
of imputation because it can change missing data from MNAR to MAR satisfying
assumptions of imputation procedures. As an illustrative example, consider a field study on
birds, where the spatial position (i.e., latitude and longitude) of nest boxes is known and is
stable through time. Here, the spatial position may not be of interest to researchers but it may
be the case that it is correlated with behaviour (e.g., shyness) and / or body mass because sub-
ordinate animals get pushed to the fringes of habitat by dominant individuals and tend to be
harder to recapture on repeated samples increasing the amount of missing data for these
animals (Holtmann, Santos, Lara & Nakagawa, 2017). One way to use these spatial

coordinates might be to generate a spatial covariance matrix between observations (using, for

example, the SpatialTools package in R – French 2016) and extracting from it principle

components (PCs). Multiple imputation could then make use of the PCs to impute missing

data (e.g. using *mice* or *mi* – Table 3) for individuals that were not measured on a given

sampling occasion. Similar approaches have been developed that make use of phylogenetic

covariance matrices (Nakagawa & de Villemereuil 2015) as well as the relatedness matrices

(Hadfield 2008).

Planned missing data designs and their application in ecology and evolution

Planned missing data designs involve collecting incomplete data from subjects or

observations of subjects on purpose by randomly assigning them to have missing

measurements or measurement occasions (Graham *et al.* 2006; Rhemtulla & Little 2012;

Little & Rhemtulla 2013). Researchers can then utilize model-based or multiple imputation

techniques to fill in missing data such that the data contains complete information for all

variables and experimental units within the dataset. Importantly, PMDD should always

conform to the MCAR assumption because missing data is random by virtue of the

experimental design making it ideal for use with imputation methods (Little & Rhemtulla

2013).

Subset Measurement Design

Planned missing data design was first developed for research utilising questionnaires

or surveys to help deal with participant fatigue, and is particularly useful when there are also

logistical and financial constraints to asking many different questions (Graham, Hofer &

Piccinin 1994; Graham *et al.* 2006). For example, a common type of PMDD called the *multi-*

form design (MFD) involves creating alternative questionnaires that each contain overlapping

questions and a sample of new questions (Graham *et al.* 2006; Little & Rhemtulla 2013).

Combining data on participants across the questionnaires, and then treating the questions

participants were not given as missing information, allows missing data to be imputed based

on the covariance between known answers (Graham *et al.* 2006).

In ecology and evolutionary biology, we often do not use questionnaires to collect

data (aside from the field of ethnobiology; see Albuquerque *et al.* 2014), therefore, an

analogous design is what we refer to as a *subset measurement design* (SMD) (Figure 2A).

Similar to the MFD, a SMD involves quantifying a common set of variables across all

individuals (e.g., body size) and then randomly allocating subjects to be quantified on a
subset of other variables (e.g., hormone concentrations, metabolism etc.) (Figure 2A).
Common variables can be those that are easily or cheaply quantified, such as body size
indices (e.g., mass, body / wing length). In contrast, variables that are expensive or
logistically challenging to quantify (e.g., gene expression, hormone concentrations) can be
randomly sampled on a subset of subjects during the experiment. When using a SMD one
should also consider, *a priori*, any potential interactions (Figure 2A) of interest and whether
the planned missingness provides sufficient power to test these interactions (Enders 2010).
254

Two-Method Design

SMD can also be applied to situations where researchers have a choice between two
variables that might be related to some response variable of interest, but where one is more
easily and cheaply quantified but has larger measurement error (i.e., less correlated with the
response variable) and the second is more logistically challenging to measure but is
considered the 'gold standard' (i.e., lower measurement error / more informative to the
question). The latter design is referred to as a *two-method design* (TMD) in the social
sciences (Little & Rhemtulla 2013). The two method design is often used in the social
sciences when one variable is known to be systematically biased, which might be the case for
self-reporting measures in surveys (Graham et al. 2006). Nonetheless, such approaches can
still be used to help understand the relationship between a response variable and a more
direct, but expensive measure, through the collection of a related, but cheaper and noisier
variable. For example, we may be interested in measuring concentrations of thyroxine (T4) to
understand how it relates to oxidative stress (concentrations of reactive oxygen species –
ROS). Thyroxine is known to impact cell metabolism by being a involved in major hormone
signalling pathways that affect ATP turnover in cells, and so, is expected to be related to
ROS. However, T4 is costly to measure and we may be able to reduce costs by measuring a
similar, more general 'metabolic' variable such as whole-organism resting metabolic rate,
which is cheap and easy to measure but may not be as strong a predictor of ROS itself (e.g.,
Figure 2A – where M is measured more than T4). As such, depending on our question, we
may actually measure more animals on whole-organism metabolic rate and fewer on T4, but
still be able to understand the impact of T4 on oxidative stress itself.
277

Wave Missingness Design

Longitudinal research questions, where repeated measurements on a set of

independent individuals is of interest, can utilize a PMDD called *wave missingness* (Figure

2B). Here, a random group of experimental subjects are not measured at all at a given time

point or measurement occasion (Little & Rhemtulla 2013; Rhemtulla *et al.* 2014). Waves can

also be blocked such that all subjects are measured at the beginning and end measurement

wave (e.g., month 1 and 6 – Figure 2B) with subsamples of animals not measured at all in

middle waves (i.e., month 2–5 – Figure 2B) (i.e., pseudo-randomised missingness; Rhemtulla

& Little 2012; Rhemtulla *et al.* 2014). Wave missingness designs have the potential to

drastically decrease data collection costs because entire sets of individuals lack measurements

at certain occasions. For example, if a study measured a set of traits on 20 animals across 20

populations, all populations might be sampled at months 1 and 6, but only some populations

measured on months 2–5. Alternatively, we may be interested in understanding seasonal

changes in profiles of two hormones across a sample individuals (e.g., testosterone and

corticosterone), sampling the same set of individuals in a population at monthly intervals

over the active season (6 months – Figure 2B). Here, we may take blood from all animals in

months 1 and 6, but avoid sampling blood from a sub-sample of animals between months 2–

5. Sampling animals across all sampling occasions means that we would need to sample

blood 120 times and run 240 different assays. In contrast, with the missingness pattern in

Figure 2B, we can reduce the total number of blood samples by ~20%, and avoid running 66

hormone assays (28% less). This can lead to reduced costs and stress to animals. The specific

wave missingness design utilized will largely depend on the research question and the

constraints faced in executing the study. However, careful attention needs to be paid when

using wave missingness designs as they have been shown to perform poorly in certain

situations with small sample sizes (Rhemtulla & Hancock, 2016).

Considerations for and General Performance of Missing Data Designs

We have overviewed three of the more common designs that can be applied to

experimental systems, however, it is important to note that PMDD's can be diverse and are

often not necessarily mutually exclusive of one another (Enders 2010; Rhemtulla & Little

2012; Little & Rhemtulla 2013; Rhemtulla *et al.* 2014). Combinations of the designs

described above are probably necessary in many real research situations (e.g. Figure 2C).

Regardless of which PMDD is used, researchers should choose the variables that are most

pertinent to the specific hypothesis being tested, or those that are likely to have small effect

sizes (and lower power), as those being measured with as little missing data as possible (i.e.,

having complete measurements on these variables). This ensures that the most pertinent questions can be tested rigorously (Graham *et al.* 2006). In addition, researchers should also consider the hypothesized correlation between variables. More tightly correlated variables ($r > 0.50$) may allow for one to plan for a greater level of missing data then two variables that are weakly correlated.

While PMDD's can be powerful tools, it is still unclear what designs work best at estimating parameters and standard errors across various situations. Graham et al. (2006) and Enders (2010) (pg. 23–36) provide an excellent overview of the power of various PMDD's. Graham et al. (2006) showed that with moderate sample sizes ($n = 200$–300), subset measurement type designs can have >0.80 power in estimating even small effects in many situations (i.e., Hedges' $d > 0.20$). Rhemtulla et al. (2014) also show that with reasonably large sample sizes ($n = 300$ for multi-form design and $n = 500$ for wave missing and hybrid designs) that parameter estimates and standard errors in latent growth models show little bias. In many situations, any loss in power resulting from missing data seems to be rather small relative to the gains in the number of questions that can be tested and the logistical and cost improvements for a given experiment (Enders 2010). Despite simulations showing the benefits of PMDD, missing data itself can affect the information content in the data that is available for estimating parameters and standard errors (Rhemtulla & Hancock, 2016). It is important to recognize that missing information is not a simple function of the amount of missing data, it also depends on which variables are missing and the pattern of missingness (Rhemtulla & Hancock, 2016). Hence, a PMDD that is useful for, say, a linear latent growth curve model, may not be useful if the growth curves are non-linear. Indeed, efficiency – or the ability the to effectively estimate uncertainty in parameters – can be reduced when using PMDD relative to a complete case scenario (Rhemtulla et al. 2016).

Benefits of planned missing data design

The PMDD's outlined above provide a number of important advantages for ecologists and evolutionary biologists. Below we discuss these potential benefits more thoroughly using realistic simulations and examples to support our arguments. Despite the benefits, it is important to recognize, however, that PMDD's do have limitations (see "*Challenges in implementing planned missing data designs*" section below), and despite their potential power in addressing a number of logistical constraints in research, it is unlikely they will be able to optimize all aspects of an experiment. When implementing PMDD it will be

important to identify, *a priori*, what models will be fit to any data collected to make sure that

the estimation of certain parameters can be done with high efficiency under the specific

PMDD (Rhemtulla & Hancock, 2016).

(i) Improving power of experimental designs

We have already indicated above that missing data procedures can improve the power of a

given study by increasing the effective sample size. In the presence of missing data standard

errors are estimated with less efficiency and thus the power to test the significance of an

effect will decline (Little & Rhemtulla 2013). Studies in ecology and evolutionary biology

are known to be under-powered in many cases (Møller & Jennions 2002; Jennions & Møller

2003), and so, this has important consequences for the inferences drawn in a given study.

This is particularly true for multi-level data that often require large sample sizes to achieve

sufficiently high power (van de Pol 2012) or even in genotype–phenotyping mapping studies,

such as GWAS (here using imputation can also improve power; e.g., Marchini & Howie

2010). Integrating a PMDD into an experiment can recover the power lost after excluding

missing data, facilitating the detection of small to moderate effects. To demonstrate how

PMDD can improve inferences, even with multi-level hierarchical data, we conducted a

couple simulations. In the first simulation, assume we are interested in estimating the

between-individual level correlation between two traits (Y_1 = "boldness" and Y_2 = "time to

emerge after predatory attack") using a multi-response model. We simulated three variables

(Y_1, Y_2 and Y_3) that follow a multi-variate normal distribution (MVN) with between (B) and

within (W)-individual covariance matrices (assuming a standard deviation (SD) = 1, which

means the correlation and covariance matrices are the same) as follows:

$$Y_{ijk} = \alpha_k + b_{ik} + w_{ijk} \qquad \text{eqn. 1}$$

$$\begin{bmatrix} Y_{ij1} \\ Y_{ij2} \\ Y_{ij3} \end{bmatrix} \sim MVN\left(\begin{bmatrix} 2 \\ 4 \\ 10 \end{bmatrix}, B + W \right)$$

$$B = \begin{pmatrix} 1 & 0.40 & 0 \\ 0.40 & 1 & 0 \\ -0.05 & -0.40 & 1 \end{pmatrix}, \quad W = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Here, Y_{ijk} is an *j* x 3 matrix containing all observations, *j,* on each individual, *i,* for the three

traits, *k* (Y_1, Y_2, and Y_3); α_k is a 1 x 3 scalar vector containing the intercept for each trait

$[2\ 4\ 10]^T$ (means for $Y_1 - Y_3$); b_{ik} is a *i* x 3 matrix containing the individual specific

random effects or deviations for individual, i, for each trait, k, with $\boldsymbol{b_{ik}} \sim MVN\left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \boldsymbol{B}\right)$.

$\boldsymbol{w_{ijk}}$ is the j observation level deviation for each individual i for the three traits k with

$\boldsymbol{w_{ijk}} \sim MVN\left(\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \boldsymbol{W}\right)$. Note here that Y_3 is not of interest with respect to the question, but

because it is correlated with Y_2 it can be used as an auxiliary variable to improve the

imputation process. For this example, we simulated 5000 datasets under varying sample

sizes. More specifically, the total number of observations ranged between 100 – 1000, with

the number of individuals varying between 10–100, each being measured 10 times. The

levels of missing data varied between 5–50%. We estimated the between– and within–

individual covariance matrices under the above simulation scenarios using MB imputation

(maximum likelihood approaches) in *ASReml–R* (vers. 3.4.1). Missing data was assumed to

be MCAR throughout the data sets, as would be the case in a PMDD, and we evaluated how

well imputation and complete case analyses performed in estimating the covariance between

Y_2 and Y_1 under these varying situations (Figure 3). As expected, given our MCAR

assumption, we did not see much impact on the point estimate (~0.40; Figure 3A & B).

Imputation procedures actually tended to do slightly better at estimating the point estimate as

there was slight estimation error for the complete case analysis with small sample sizes and

high levels of missing data (Figure 3B). This was likely the result of model convergence

problems when one has small sample sizes and high levels of missingness, with 15–26% of

the simulations failing to converge under these conditions (i.e., $n = 100$ & 150 with 50%

missingness). Despite this, we observed improvements in the estimation of standard errors

when imputing data in comparison to the complete case analysis (Figure 2C & D), suggesting

that imputation procedures, even with hierarchical data such as this, can lead to fairly

substantial improvements in power. This is particularly important as many research areas,

such as quantitative genetics and behavioural ecology, are interested variance partitioning

methods to understand trait covariance (Dingemanse & Dochtermann 2013; Brommer &

Class 2017; Careau & Wilson 2017).

Ecological and evolutionary questions are often more complex than simply estimating

variance components. Experiments will often combine experimental manipulations of

independent individuals and repeatedly measure these individuals across their life. To

understand the benefits of PMDD at improving statistical inferences when both fixed and

random effects might be of interest we conducted a second simulation. Assume we have

manipulated the early thermal environment of a sample of lizard eggs – a common approach

in lizard research (e.g., Noble, Stenhouse & Schwanz 2017). We might be interested in

understanding how these thermal environments affect the growth curves of animals within

each treatment (Figure 4). We simulated data assuming that individuals follow a linear

growth trajectory (random regression model), at least over the period in which we measured

their weight, according to the following model:

$$m_{ij} = (\beta_0 + \alpha_i) + \beta_1 T_{ij} + (\beta_2 + s_j)A_{ij} + \beta_3(A_{ij} \cdot T_{ij}) + \varepsilon_{ij} \qquad \text{eqn. 2}$$

where m_{ij} is the mass of individual i for observation j, T_{ij} is a dummy variable ('0' or '1')

indicating whether observation j for individual i belongs to the control group (23ºC) or the

treatment group (26ºC), A_{ij} is the age of individual i at observation j, β_1 is the contrast

between the control group mean at age 0 (β_0) and the treatment mean at age 0, β_2 is the

effect of age on mass, and β_3 is the difference in mass change across age between treatments

(i.e., interaction effect). α_i and s_j are individual level random effects for the random

intercept and slope, respectively, which are assumed to follow an

$\sim MVN\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0.5 & 0.25 \\ 0.25 & 0.4 \end{bmatrix}\right)$. ε_{ij} is the observation-level random effect assumed to follow an

$\sim N(0, 1)$. An example of the simulated data is shown in Figure 4. If we were limited in the

total amount of sampling, say 100 observations, then we can design an experiment where 10

animals (5 / treatment) were each measured 10 times (*Scenario 1*) or we could use a PMDD

whereby we increase the total number of individuals to $n = 20$ (10 / treatment), but instead

measure each individual randomly only five times, instead of 10 (*Scenario 2*) (Figure 4).

Under these two scenarios we simulated 1500 datasets and used *ASReml–R* to estimate the

parameters and their standard errors in eqn. 2. Overall, scenario 2 had a number of benefits

both with respect to estimating fixed and random effects performing nearly as well as

complete data (i.e., 20 individuals measured 10 times – *Scenario 3*). For all fixed effect

estimates, there was between ~10–27% reduction in the standard errors with standard errors

for the slopes and treatment interactions receiving a substantial boost. Interestingly, we also

increase our precision in estimating random slopes (Table 1). Benefits of using a PMDD are

even greater with larger sample sizes (e.g., 500 individuals measured 100 times *versus* 1000

individuals measured 50 times – see supplementary material Table S1), with a 23–29%

reduction across both fixed and random effects. Estimation of random effects and their

precision were improved quite a bit more with larger sample sizes (Table S1). Overall, our

simulations suggest that PMDDs can provide power benefits under realistic experimental

situations that are common in ecology and evolution.

(ii) Improved animal welfare: considering the three 'R's'

Central to research in ecology and evolutionary biology, particularly with respect to research

on vertebrate animals, are issues surrounding animal welfare (Stamp Dawkins 2006; Barnard

2007; Cuthill 2007; Stamp Dawkins 2008; McMahon *et al.* 2012). Biological research on

vertebrates, and indeed any study species, should strive to minimize pain, suffering and

distress caused by experimentation, and we look to the three R's (Refinement, Replacement

and Reduction) to achieve these goals (Cuthill 2007). Planned missing data design can be an

important tool during the experimental planning stages that directly targets two of the three

R's (Refinement and Reduction) to improve animal welfare. Often, PMDD can concurrently

target both Refinement and Reduction at the same time by using less invasive procedures and

allowing data to be collected on fewer experimental subjects (or less often on a given

subject). For example, we could use a SMD to randomly assign the measurement of different

physiological traits (e.g., metabolism, hormones) to a subset of subjects, reducing the number

of subjects quantified on a given physiological measure. Additionally, fewer repeated

physiological measurements can be done on a given subject by randomly assigning a

different temporal sequence of measurements to individuals, refining the experimental design

to reduce stress inflicted by repeated handling. Refinement can further occur by adopting a

TMD where different physiological assays measuring the same construct, say 'innate

immunity', can be done in such a way that no one individual has both measurements, but

rather is randomly assigned to the cheaper less invasive measure. In summary, PMDD can

improve animal welfare without compromising our ability to effectively answer a given

question by designing an experiment that has too little power.

(iii) Reduced research costs

One critical benefit of implementing PMDD is the ability to get a 'bigger bang for your buck'

in terms of the research cost to outcome ratio. The cost savings when using a PMDD can be

substantial, particularly for experiments involving expensive biochemical, proteomic,

metabolomic and genomic work. However, they may also reduce costs for laborious

experiments that require the breeding or measurement of a large number of individuals, such

as is the case for diallel crosses in quantitative genetics (Lynch & Walsh, 1998). For

example, assume we are interested in measuring hormone profiles for 60 individuals across a

six-month activity period, meaning we would have a total of $n = 360$ blood samples (i.e.,

each animal sampled once a month). We might be able to reduce our sample size by 33%

using a PMDD similar to that described in Figure 2C, bringing the total number of blood

samples to $n = 240$. If we assume that the cost of individual reactions to run assays, in

addition to labour costs, was $8 per sample, then we would save $900 for this experiment.

These additional savings could be used to run a follow up experiment or even go towards

assaying a sub-sample of individuals on a second hormone. Alternatively, we may also be

interested in quantifying telomere length to understand cellular senescence using either flow

cytometry (estimated cost / sample = $68) or qPCR (estimated cost / sample = $13) methods

(Nussey *et al.* 2014) to understand patterns of telomere attrition over the season in relation to

hormones. In this example, using a PMDD would save $8160 for flow cytometry methods or

$1560 using qPCR methods (assuming we really had money to do all animals). It also

demonstrates how a TMD can also save costs. Flow cytometry has been identified as a

promising method for accurate, high throughput quantification of telomere length (Nussey *et*

al. 2014). However, it is quite expensive compared to qPCR methods. A PMDD where a

sample of animals are quantified both on qPCR and flow cytometry, and those missing flow

cytometry data are then imputed would lead to substantial cost savings and allow one to

verify the utility of both methods. We therefore view PMDD as a promising approach to

improving cost efficiency.

(iv) Stronger inferences and testing predictions from multiple working hypotheses

Strong inference involves devising alternative hypotheses and then running an experiment or

set of experiments to test alternative predictions generated from these hypotheses (Platt 1964;

Chamberlain 1965). Identifying and testing among alternative hypotheses is the hallmark of

rapid scientific progress, indeed Platt (1964) suggested that this was one of the primary

reasons for the rapid advancement of molecular biology through the 1960s. Nonetheless, it is

clear that ecological and evolutionary studies rarely test predictions from multiple working

hypotheses (Betini, Avgar & Fryxell 2017). Betini et al. (2017) suggest a number of

intellectual and practical barriers impeding the use of multiple working hypotheses, but

particularly relevant for our argument are the barriers limiting one's ability to "execute" such

investigations. Designing fully factorial experiments to disentangle predictions from

alternative hypotheses is a major hurdle, referred to as the "*fallacy of the factorial design*"

(Betini, Avgar & Fryxell 2017), whereby the addition of every new working hypothesis

requires a new treatment or set of measurement variables leading to a geometric increase in

number of required replicates. Including variables or experimental manipulations that test

predictions from alternative hypotheses into a PMDD, can potentially off-set the need for

prohibitively large sample sizes and time-consuming and costly measurements. These

approaches may maximize the information gained from an experiment to facilitate more rapid

scientific progress compared to testing single working hypotheses. Additionally, predictions

from hypotheses can be tested using the same sample of experimental subjects, reducing the

spatial and temporal differences between experiments.

(v) Greater integration between research fields and disciplines

Understanding ecological and evolutionary processes require an integrative,

multidisciplinary approach to tackling research questions (Wake 2009) – including the use of

observational, experimental and theoretical modelling (Wake 2009). Integrative research

questions cut across traditional boundaries, often making use of novel, and sometimes

expensive techniques to help facilitate an understanding of a system or process. Despite its

benefits, it can often be challenging to implement in practice given the different norms in

various research fields, along with the costs and logistical constraints in doing integrative

work. We argue that PMDD may help facilitate more multidisciplinary research efforts as it

alleviates these constraints.

To demonstrate how PMDD can facilitate integrative research, assume that we are

interested in testing Pace-of-life syndrome (POLS) theory (Reale *et al.* 2010). POLS theory

explicitly predicts covariation between physiological, behavioural and life-history traits (Biro

& Stamps 2008; Careau *et al.* 2010; Reale *et al.* 2010). Integration of suites of behavioural

traits can lead to consistent individual differences in behaviour (i.e. personality - Reale *et al.*

2010; Stamps & Groothuis 2010) that can form behavioural syndromes (Sih & Bell 2008;

Stamps & Groothuis 2010; Sih & Del Guidice 2012). Importantly, physiological mechanisms

are thought to be the primary mechanisms (e.g. hormones, metabolism; Biro & Stamps 2008;

Biro & Stamps 2010; Careau *et al.* 2010; Reale *et al.* 2010) underpinning both behavioural

and life-history variation in populations. Pace-of-life theory is integrative, requiring

concurrent measurements of physiological, behavioural and life-history traits to understand

their covariance. It often requires laborious, time intensive, and sometimes costly

measurements of the same individuals over time, posing major hurdles in obtaining the

necessary data required. Additionally, physiological measures (e.g. metabolism, hormones,

ROS, immunity) can be costly to obtain limiting the number of samples that can be collected.

Despite this, it is important to take many physiological measurements (Adamo 2004). A

PMDD may alleviate some of these constraints by reducing the need to measure all variables

across subjects over time.

Challenges in implementing planned missing data designs

As with any new research method, there will be challenges, particularly in establishing the

most suitable approaches that work across a wide diversity of different research questions and

experimental designs in ecology and evolution. Given that PMDD is still new, stimulating

interest in them will be the first step to identifying, solving and implementing solutions to

some of the challenges that crop up. Below we discuss some of the hurdles we see to

implementing a PMDD and suggest some tentative solutions.

Unplanned missing data

Of course, as with any experiment, unplanned missing data will creep into PMDD designs.

Often missing data may be random, such as when a piece of equipment malfunctions during a

set of measurements or when recording errors are identified. Random instances of missing

data, even if unplanned, will not affect the imputation process or the utility of PMDD unless

missing data levels begin to get quite high. However, our simulations show, as well as

others', that imputation procedures perform quite well even with large amounts of missing

data (~50% – see Figure 3). Nonetheless, there are real situations where unplanned missing

data can be MNAR and this will affect any experiment regardless of whether a PMDD is

implemented or not. We have outlined above how data can be made MAR though the use of

auxiliary variables, and these unplanned missing data, can then be imputed normally along

with any planned missing data using the same statistical methods. We therefore advise

colleagues to collect possible auxiliary variables where possible to counter unplanned

missing data.

Imputation with generalised linear mixed effect models

Multiple and model-based (MB) imputation both work well with normally distributed

data, however, in reality variables often are non-normally distributed. While MB procedures

are limited to situations that assume multivariate normality, MI procedures can also work

with non-normal data fit using generalised linear mixed effect models (e.g., Poisson

GLMMs) (Schafer 1997). However, implementation in the context of GLMMs is still under

active development, and in many cases, is restricted to simple random effect structures (van

Buuren & Groothuis-Oudshoorn 2011; Enders, Mistler & Keller 2016; Quartagno &
Carpenter 2016; Audigier & Resche-Rigon 2017). Nonetheless, two-level random regression
models can be run in a number of existing packages (e.g., *mice*) and we believe that the
capacity to run more sophisticated models will grow in the near future.

579   To re-assure readers that imputation can and does work with non-normal
distributions, we provide a simulated example along with sample R code to demonstrate MI
procedures with GLMMs in Box 1. For our hypothetical example, assume we are interested
in provisioning rates (the number of feeding visits by a parent) in a bird species, the fictitious
Missing Capped Warbler (*Sylvia absenscapilla*). We would like to understand the costs of
female provisioning by experimentally manipulating brood sizes ($n = 6$ chicks) in a random
sample of birds compared to a control group, which has normal brood sizes ($n = 3$ chicks)
(Liebl, Browning & Russell 2016). The Missing Capped Warbler is notoriously difficult to
observe as it is found in thick scrub, and so, we placed cameras at random nests during the
first two weeks of the breeding season to observe provisioning rates in the two treatments
over a 5-hour period. Provisioning rates are known to change as chicks develop (Khwaja *et*
al. 2017), and so, the cameras were on each nest for a total of 20 days to understand how the
demands of chicks change, and whether females can keep up with these demands.
Unfortunately, it is a laborious process to observe all the resulting video for 40 birds
measured over 20 days (a total of 4000 hours of video!). We therefore decided to implement
a PMDD, where we randomly sampled a set of videos ($n = 536$ out of 800 videos; ~ 33%
missing data) where provisioning rates can be quantified (cutting the total hours of video
watching to 2680 hours). Planned missing data can be imputed, for example, using the *mice*
package (see Box 1). After a long and laborious field season, we were able to collect data that
shows a tendency for experimentally elevated clutch sizes to have higher rates of
provisioning that increase over the care period (Figure 5). Overall, we see that the imputed
data matches well with the complete dataset, with nearly identical results in this simulation
(see Table in Box 1).

Overcoming psychological barriers to missing data
One of the biggest challenges to implementing PMDDs probably involves the need for
researchers to over-come the 'psychological taboos' around missing data, and the suspicion
of techniques for handling missing data (Enders 2010). We can re-assure readers that missing
data practices are now very well established (Graham *et al.* 2006; Nakagawa 2017), and are
rather painlessly implemented in many commonly used statistical software such as *R*, *SAS*,

SPSS, and *MPlus* (See Table 2 for an overview). In fact, many techniques are implemented

by default when missing data is included as response variables in models for a number of

mixed modelling packages (e.g. model-based procedures in '*MCMCglmm*' and '*ASReml*').

While statistical algorithms vary across these platforms, fairly sophisticated and versatile

ones are now implemented in packages for some of the most widely used platforms (e.g.

'*mice*', '*mi*', '*multimput*' and '*Amelia*' in the R environment – Table 2) that implement MI

algorithms known to perform well under a wide variety of situations (Schafer & Graham

2002; van Buuren 2012; Enders, Mistler & Keller 2016; Quartagno & Carpenter 2016;

Resche-Rigon & White 2016; Audigier *et al.* 2017). These techniques are under active

development (e.g., the *mice* package in R), and so, we envisage the breadth of problems these

tools can tackle to increase and be even easier to apply in the future. Nonetheless, caution is

still needed in their implementation as it is unclear whether imputation procedures perform

well under all circumstances (Nakagawa 2017). Statistical procedures for missing data are

still rarely taught in undergraduate and graduate level courses, so part of the solution will be

to begin educating students and practitioners about how to perform imputation procedures,

explicitly highlighting some of the challenges and caveats that need to be considered.

Nonetheless, there are now excellent resources that provide an in depth look at imputation

procedures (Gelman & Hill 2002; Schafer & Graham 2002; Graham 2009; Enders 2010; Su

et al. 2011; van Buuren & Groothuis-Oudshoorn 2011; Nakagawa 2017).

Uncertainties surrounding the best PMDD

One challenge in implementing PMDD is the uncertainty around what the most appropriate

missing data design is for a given experiment. This is particularly true in ecology and

evolutionary biology because different questions, experimental systems, data structure and

measurement variables may require creative combinations of different PMDD's that we

discuss in our paper – or possibly even new ones! While we argue that the benefits of PMDD

can be substantial it may still be important to test the robustness and power of any given

PMDD through simulations (Rhemtulla & Handcock, 2016). With some very simple

simulated data based on effect sizes and experimental designs relevant to the question at

hand, the power of different PMDD's can be thoroughly tested during the design stage of an

experiment (Enders 2010). This may require researchers thinking carefully about the model

they wish to test and simulating data under this model early on (Rhemtulla & Handcock,

2016). Enders (2010 p.g. 30) provide a nice introduction on how to conduct power analysis

with PMDD's using simulations, and we provide all our R code which we hope can act as a

skeleton for readers to familiarize themselves with simulations to help them sort out the best
PMDD for their particular situation. Additionally, new multi-level simulation packages, such
as SQuID, allow for researchers to simulate hierarchical data easily (Allegue *et al.* 2017).
Data can be downloaded and missing data introduced to evaluate the power of different
PMDD's. While definitive guidelines will depend on the experimental design, question and
covariance between traits, we believe that up to ~30% missing data overall across a wide
variety of different designs will likely not compromise performance of imputation
procedures.

Conclusions and future directions

Our goal was to put planned missing data design on the radar of ecologists and evolutionary
biologists given the substantial number of ethical, logistical and cost saving benefits it may
afford. We have provided some guidance on possible PMDD's that can be implemented in
research programs and shown, with simulations, that even hierarchical / multilevel data
imputation procedures can perform quite well. While it is still unclear whether imputation
procedures in a multilevel framework will work under all circumstances there is increasing
awareness of the need to develop such techniques with hierarchical data and in many cases
existing methods will likely perform well (Drechsler 2015; Quartagno & Carpenter 2016;
Audigier & Resche-Rigon 2017; Audigier *et al.* 2017). We encourage colleagues to begin
thinking about PMDD's and their utility in their research both to improve research quality
and to promote integrative, cost effective research projects in ecology and evolutionary
biology.

Acknowledgements
We would like to thank the I-DEEL lab (Joel Pick, Rose O'Dea, Malgorzata Lagisz, and
Fonti Kar), Diego Barneche and Corey Callaghan for feedback on previous versions of our
manuscript. We would also like to thank the editors, Mijke Rhemtulla and an anonymous
reviewer for very constructive comments that greatly improved our manuscript. DWAN was
supported by an ARC Discovery Early Career Research Award (DE150101774) and SN an
ARC Future Fellowship (FT130100268). The authors declare no conflicts of interest.

Author's contributions

SN and DN conceived the research. DN and SN conducted the simulations. DN wrote the manuscript, with input from SN. SN contributed critically and constructively to the revisions of the manuscript.

Data Accessibility

All R code and simulation results used in the manuscript were provided upon submission.

References

Adamo, S.A. (2004) How should behavioural ecologists interpret measurements of immunity? *Animal Behaviour,* **68,** 1443–1449.

Albuquerque, U.P., Ramos, M.A., Paiva de Lucena, R.F. & Alencar, N.L. (2014) *Methods and techniques in ethnobiology and ethnoecology*. Springer, New York.

Allegue, H., Araya-Ajoy, Y.G., Dingemanse, N.J., Dochtermann, N.A., Garamszegi, L.Z., Nakagawa, S., Réale, D., Schielzeth, H., Westneat, D.F. & Hadfield, J. (2017) Statistical Quantification of Individual Differences (SQuID): an educational and statistical tool for understanding multilevel phenotypic data in linear mixed models. *Methods in Ecology and Evolution,* **8,** 257-267.

Allison, P.D. (2012) Handling missing data by maximum liklihood. *SAS Global Forum,* **Statistics and Data Analysis, Statistical Horizons, Haverford, PA, USA**.

Audigier, V. & Resche-Rigon, M. (2017) micemd: Multiple imputation by chained equations with multilevel data. *R package version 1.1.0.* https://cran.r-project.org/package=micemd.

Audigier, V., White, I.R., Jolani, S., Debray, T.P.A., Quartagno, M., Carpenter, J., van Buuren, S. & Resche-Rigon, M. (2017) Multiple imputation for multilevel data with continuous and binary variables. *arXiv,* 1–38.

Barnard, C. (2007) Ethical regulation and animal science: why animal behaviour is special. *Animal Behaviour,* **74,** 5–13.

Betini, G.S., Avgar, T. & Fryxell, J.M. (2017) Why are we not evaluating multiple competing hypotheses in ecology and evolution? *Royal Society Open Science,* **4,** 160756.

Biro, P.A. & Dingemanse, N.J. (2009) Sampling bias resulting from animal personality. *Trends in Ecology & Evolution,* **24,** 66-67.

Biro, P.A. & Stamps, J.A. (2008) Are animal personality traits linked to life-history productivity? *Trends in Ecology & Evolution,* **23,** 361-368.

Biro, P.A. & Stamps, J.A. (2010) Do consistent individual differences in metabolic rate promoste consistent individual differences in behavior? *Trends in Ecology and Evolution,* **25,** 653-659.

Brommer, J.E. & Class, B. (2017) Phenotypic correlations capture between-individual correlations underlying behavioural syndromes. *Behavioral Ecology and Sociobiology,* **71,** 50.

Butler, D. (2009) asreml: asreml() fits the linear mixed model. *R package version 3.0.* http://www.vsni.co.uk/.

Careau, V., Reale, D., Humphries, M.M. & Thomas, D.W. (2010) The pace of life under artificial selection: Personality, energy expenditure, and longevity are correlated in domestic dogs. *The American Naturalist,* **175,** 753-758.

Careau, V. & Wilson, R.S. (2017) Of Uberfleas and Krakens: Detecting Trade-offs Using Mixed Models. *Integrative and Comparative Biology*.

Chamberlain, T.C. (1965) The method of multiple working hypotheses. *Science,* **148,** 754–759.

Collins, L.M., Schafer, J.L. & Kam, C.-M. (2001) A comparison of inclusive and restrictive strategies in modern missing data procedures. *Psychological Methods,* **6,** 330-351.

Cuthill, I.C. (2007) Ethical regulation and animal science: why animal behaviour is not so special. *Animal Behaviour,* **74,** 15–22.

Dingemanse, N.J. & Dochtermann, N.A. (2013) Quantifying individual variation in behaviour: mixed-effect modelling approaches. *J Anim Ecol,* **82,** 39-54.

Drechsler, J. (2015) Multiple imputation of multilevel missing data - rigor versus simplicity. *Journal of Educational and Behavioral Statistics,* **40,** 69-95.

Ellington, E.H., Bastille-Rousseau, G., Austin, C., Landolt, K.N., Pond, B.A., Rees, E.E., Robar, N. & Murray, D.L. (2015) Using multiple imputation to estimate missing data in meta-regression. *Methods in Ecology and Evolution,* **6,** 153-163.

Enders, C.K. (2001a) The impact of nonnormality on full information maximum-likelihood estimation for structural equation models with missing data. *Psychological Methods,* **6,** 352-370.

Enders, C.K. (2001b) A primer on maximum likelihood algorithms available for use with missing data. *Structural Equation Modeling,* **8,** 128-141.

Enders, C.K. (2010) *Applied missing data analysis*. The Guilford Press, New York, USA.

Enders, C.K., Mistler, S.A. & Keller, B.T. (2016) Multi-level multiple imputation: A review and evoluation of joint modeling and chained equations imputation. *Psychological Methods,* **21,** 222-240.

French, J. (2016) SpatialTools: Tools for spatial data analysis. *R package version 1.0.2.* https://cran.r-project.org/package=SpatialTools.

Gelman, A. & Hill, J. (2002) *Data analysis using regression and multilevel/hierachical models*. Cambridge University Press.

Graham, J.W. (2003) Adding missing-data-relevant variables to FIML-based structural equation models. *Structural Equation Modeling,* **10,** 80-100.

Graham, J.W. (2009) Missing data analysis: Making it work in the real world. *Annual Review of Psycology,* **60,** 549-576.

Graham, J.W., Hofer, S.M. & MacKinnon, D.P. (1996) Maximizing the usefulness of data obtained with planned missing value patterns: An application of maximum likelihood procedures. *Multivariate Behavioral Research,* **31,** 197-218.

Graham, J.W., Hofer, S.M. & Piccinin, A.M. (1994) Analysis with missing data in drug prevention research. *Advances in data analysis for prevention intervention research* (eds L.M. Collins & L. Seitz), pp. 13–63. National Institute on Drug Abuse Research monograph series, Washington, DC, USA.

Graham, J.W., Taylor, B.J., Olchowski, A.E. & Cumsille, P.E. (2006) Planned missing data designs in psychological research. *Psychological Methods,* **11,** 323-343.

Hadfield, J.D. (2008) Estimating evolutionary parameters when viability selection is operating. *Proc Biol Sci,* **275,** 723-734.

Hadfield, J.D. (2010) MCMC methods for multi-response generalised linear mixed models: The MCMCglmm R package. *Journal of Statistical Software,* **33,** 1–22.

Hardt, J., Herke, M. & Leonhart, R. (2012) Auxiliary variables in multiple imputation in regression with missing X: a warning against including too many in small sample research. *BMC Biomedical Research Methodology,* **12,** 1–13.

Holtmann, B., Santos E. S. A., Lara C. E. & Nakagawa S. (2017). Personality-matching
habitat choice, rather than behavioural plasticity, is a likely driver of a phenotype-
environment covariance. Proceedings of Royal Society of London B. 284: 20170943
Honaker, J., King, G. & Blackwell, M. (2011) Amelia II: A program for missing data.
Journal of Statistical Software, **45,** 1–47.
Jennions, M.D. & Møller, A.P. (2003) A survey of the statistical power of research in
behavioral ecology and animal behavior. *Behavioral Ecology,* **14,** 438–445.
Khwaja, N., Preston, S.A.J., Hatchwell, B.J., Briskie, J.V., Winney, I.S. & Savage, J.L.
(2017) Flexibility but no coordination of visits in provisioning riflemen. *Animal*
Behaviour, **125,** 25-31.
Kleinke, K. & Reinecke, J. (2013) *countimp*: Multiple imputation of incomplete count data. *R*
package version 1.0.
Liebl, A.L., Browning, L.E. & Russell, A.F. (2016) Manipulating carer number versus brood
size: complementary but not equivalent ways of quantifying carer effects on
offspring. *Behav Ecol,* **27,** 1247-1254.
Little, T.D., Jorgensen, T.D., Lang, K.M. & Moore, E.W. (2013) On joys of missing data.
Journal of Pediatric Psychology, **doi:10.1093/jpepsy/jst048,** 1-12.
Little, T.D. & Rhemtulla, M. (2013) Planned missing data designs for developmental
researchers. *Child Development Perspectives,* **7,** 199-204.
Little, T.D. & Rubin, D.B. (2002) *Statistical analysis with missing data*. John Wiley & Sons,,
New York, 2nd edition.
Lynch, M. & Walsh, B. (1998). *Genetics and analysis of quantitative traits*. Sinauer
Associates, Sunderland, Massachusetts, USA.
Marchini, J. & Howie, B. (2010) Genotype imputation for genome-wide association studies.
Nat Rev Genet, **11,** 499-511.
McKnight, P.E., Mcknight, K.M., Sidani, S. & Figueredo, A.J. (2007) *Missing data: A gentle*
introduction. The Guilford Press., New York, USA.
McMahon, C.R., Harcourt, R., Bateson, P. & Hindell, M.A. (2012) Animal welfare and
decision making in wildlife research. *Biological Conservation,* **153,** 254–256.
Møller, A.P. & Jennions, M.D. (2002) How much variance can be explained by ecologists
and evolutionary biologists? *Oecologia,* **132,** 492-500.
Nakagawa, S. (2017) Missing data: mechanisms, methods, and messages. *Ecological*
Statistics: Contemporary Theory and Application (eds G.A. Fox, S. Negrete-
Yankelevich & V.J. Sosa), pp. 81–105. Oxford University Press., Great Clarendon
Street, Oxford, UK.
Nakagawa, S. & de Villemereuil, P. (2015) A simple and general method for accounting for
phylogenetic uncertainty via Rubin's rules in comparative analysis. *Peer J. PrePrints,*
3:e1216v1.
Nakagawa, S. & Freckleton, R.P. (2010) Missing in action: the dangers of ignoring missing
data. *Trends in Ecology & Evolution,* **23,** 592-596.
Nakagawa, S. & Freckleton, R.P. (2011) Model averaging, missing data and multiple
imputation: A case study for behavioural ecology. *Behavioral Ecology and*
Sociobiology, **65,** 103-116.
Noble, D.W.A., Stenhouse, V. & Schwanz, L.E. (2017) Developmental temperatures and
phenotypic plasticity in reptiles: a systematic review and meta-analysis. *Biol Rev*
Camb Philos Soc.
Nussey, D.H., Baird, D., Barrett, E., Boner, W., Fairlie, J., Gemmell, N., Hartmann, N., Horn,
T., Haussmann, M., Olsson, M., Turbill, C., Verhulst, S., Zahn, S. & Monaghan, P.
(2014) Measuring telomere length and telomere dynamics in evolutionary biology and
ecology. *Methods in Ecology and Evolution,* **5,** 299-310.

Platt, J.R. (1964) Strong inference *Science,* **146,** 347–353.

Quartagno, M. & Carpenter, J. (2016) *jomo*: A package for multilevel joint modelling
multiple imputation. *R package version 2.2-0*.

Reale, D., Garant, D., Humphris, M.M., Bergeron, P., Careau, V. & Montiglio, P.-O. (2010)
Personality and the emergence of the pace-of-life syndrome concept at the population
level. *Philisophical Transactions of the Royal Society B: Biological Sciences,* **365,**
4051-4063.

Resche-Rigon, M. & White, I.R. (2016) Multiple imputation by chained equations for
systematically and sporadically missing multilevel data. *Stat Methods Med Res*.

Rhemtulla, M., Jia, F., Wu, W. & Little, T.D. (2014) Planned missing designs to optimize the
efficiency of latent growth parameter estimates. *International Journal of Behavioral
Development,* **38,** 423-434.

Rhemtulla, M. & Hancock, G.R. (2016). Planned missing data designs in educational
psycology research. *Educational Psycologist*, **51**, 305–316.

Rhemtulla, M. & Little, T.D. (2012) Planned missing data designs for research in cognitive
development. *Journal of Cognition and Development,* **13,** 425-438.

Rhemtulla, M., Savalei, V. & Little, T.D. (2016). On the asymptotic relative efficiency of
planned missingness designs. *Psychometrika*, **81**, 60–90.

Rubin, D.B. (1976) Inference and missing data. *Biometrika,* **63,** 581-592.

Rubin, D.B. (1987) *Multiple imputation for non-response in surveys*. Wiley, New Yok.

Rubin, D.B. (1996) Multiple imputation after 18+ years. *Journal of the American Statistical
Association,* **91,** 473-489.

Schafer, J.L. (1997) *Analysis of incomplete multivariate data*. Chapman & Hall / CRC Press.,
Boca Raton, USA.

Schafer, J.L. & Graham, J.W. (2002) Missing data: our view of the state of the art.
Psycological Methods, **7,** 147-177.

Schafer, J.L. & Olsen, M.K. (1998) Multiple imputation for multivariate missing-data
probelms: A data analyst's perspective. *Multivariate Behavioural Research,* **33,** 545-
571.

Sih, A. & Bell, A.M. (2008) Insights for behavioural ecology from behavioural syndromes.
Advances in the Study of Animal Behaviour, **38,** 227-281.

Sih, A. & Del Guidice, M. (2012) Linking behavioural syndromes and cognition: a
behavioural ecology perspective. *Philosophical Transactions of the Royal Society of
London B Biological Sciences,* **367,** 2762-2772.

Stamp Dawkins, M. (2006) A user's guide to animal welfare science. *Trends in Ecology &
Evolution,* **21,** 77–82.

Stamp Dawkins, M. (2008) The science of animal suffering. *Ethology,* **114,** 937–945.

Stamps, J.A. & Groothuis, G.G. (2010) The development of animal personality: relevance,
concepts and perspectives. *Biological Reviews (Cambridge),* **85,** 301-325.

Su, Y.-S., Gelman, A., Hill, J. & Yajima, M. (2011) Multiple imputation with diagnostics
(mi) in R: Opening windows into the black box. *Journal of Statistical Software,* **45,**
1–31.

van Buuren, S. (2012) *Flexible imputation of missing data. In: The handbook of advanced
multilevel analysis*. CRC Press, Taylor and Francis Group, Boca Raton, FL, USA, pp
173-196.

van Buuren, S. & Groothuis-Oudshoorn, K. (2011) mice: Multivariate imputation by chained
equations in R. *Journal of Statistical Software,* **45,** 1-64.

van de Pol, M. (2012) Quantifying individual variation in reaction norms: how study design
affects the accuracy, precision and power of random regression models. *Methods in
Ecology and Evolution,* **3,** 268–280.

von Hippel, P. & Lynch, J. (2013) Efficiency gains from using auxiliary variables in
864   imputation. *Cornell University Library,* **arXiv:1311.5249**.
865   Wake, M.H. (2009) Integrative biology: Science for the 21st century. *BioScienec,* **58,** 349–
866   353.
867

Tables and Figures

Table 1 – Effect of two scenarios on the ability to estimate parameters of a mixed model.
Average parameter estimates and their standard errors for both fixed and random effects over
1500 simulated datasets. Example data is shown in figure 4. Abbreviations and symbols as
follows: ρ: correlation between intercept (α) and slope (s); σ_α^2 = variance estimate for
random intercept; σ_s^2 = variance estimate for random slope; trt = treatment; age = change in
mass across age. % SE Decrease = the percent decrease in standard error from Scenario 1
compared to Scenario 2. "True" are the true parameters that the simulation was based on.
Scenario 1 involves a hypothetical situation where 10 lizards are measured 10 times with 0%
missing data. Scenario 2 is a situation where 20 lizards were measures 5 times (i.e., 50%
missing data for each animal. In Scenario 3, 20 lizards are measured 10 times (i.e., a full
complete case analysis for comparison). Note that simulations with larger sample sizes lead
to even better improvements of SE when PMDD are implemented (Table S1).
884
885
886

	Scenario 1		Scenario 2		Scenario 3		% SE Decrease	True
Fixed Effects								
	Est.	*SE*	*Est.*	*SE*	*Est.*	*SE*		
intercept	1.21	0.45	1.22	0.40	1.19	0.31	9.83	1.20
trt	2.01	0.63	1.98	0.57	2.00	0.44	9.82	2
age	1.79	0.28	1.80	0.21	1.79	0.20	27.06	1.8
trt*age	1.61	0.40	1.60	0.29	1.60	0.28	27.06	1.6
Random Effects								
	Est.	*SE*	*Est.*	*SE*	*Est.*	*SE*		
σ_α^2	0.57	0.55	0.62	0.59	0.51	0.34	-6.96	0.50
$\rho(s,\alpha)$	0.24	0.25	0.23	0.21	0.24	0.16	16	0.25
σ_s^2	0.41	0.21	0.41	0.15	0.40	0.14	31.26	0.40

893
894 **Table 2** – Examples of common packages and statistical programs that can be used to deal
895 with missing data. Abbreviations are as follows: SA = stand-alone program; MI = multiple
896 imputation; MB = model-based imputation; B = both, R = response, P = predictors, Y = yes,
897 ? = unknown.
898

Package	Prog.	Algorithms	Response/ Predictor	Multi-level	Reference / link
mi	R	MI	B	Y	Su et al. (2011)
mice	R	MI	B	Y	Van Buuren & Groothuis-Oudshoorn (2011)
micemd	R	MI	B	Y	Audigier & Resche-Rigon (2017)
jomo	R	MI	B	Y	Quartagno & Carpenter (2016)
Amelia	R	MI	B	Y	Honaker et al. (2011)
multimp	R	MI	B	Y	https://github.com/inbo/multimput
MCMCglmm	R	MB	R	Y	Hadfield (2010)
ASReml	R/SA	MB	B	Y	Butler (2009)
SAS	SA	MB/MI	B	Y	https://stats.idre.ucla.edu/sas/seminars/multiple-imputation-in-sas/mi_new_1/
SPSS	SA	MB/MI	B	?	https://www.ibm.com/ms-en/marketplace/spss-missing-values
MPlus	SA	MB/MI	B	Y	https://www.statmodel.com/index.shtml

Figure 1 – Two major types of imputation procedures: **A)** model-based procedures (e.g., full information maximum likelihood, expectation maximization, MCMC) and **B)** multiple imputation procedures. Each large square represents a dataset containing four variables (Y, X1, X2, X3) and n = 10 observations. Small red squares represent missing data and black squares complete data. Model-based procedures **(A)** take both observed and missing data in the analysis under a pre-specified model [E(Y) = **XB**)], augment missing data, estimate parameter estimates (B1, B2, B3) and then re-iterate this process with updated parameters [different coloured B1, B2, B3 and E(Y) = XB] until the model converges on a set of unbiased parameter estimates. Multiple imputation **(B)** uses complete data and often (but not always) uses other variables within the data as predictors of a specific variable. It then imputes using regression equations plausible values of missing data to generate *m* complete datasets. To prevent biased estimates residual error is added to each of the imputed data points (checkered small squares in step 2). These *m* datasets are then analysed with a given model, which can be different from the ones used to impute and, using Rubin's rules (Rubin 1987), pool the parameter estimates across datasets (in this case B1). Abbreviations are as follows: E = expectation of variable, or mean estimate of variable; **X** = design matrix; B = vector of parameter estimates (e.g., B1, B2, B3); ε = residual effect or random error.

Figure 2 – Three examples of planned missing data designs relevant for ecological and evolutionary research. In all cases, 20 individuals are shown along the rows (ID) and variables (e.g., BM) or traits (e.g., T1) measured are shown along columns. Abbreviations: BM = body mass, S = Sex, M = metabolism; TH = thyroxine (T4), T1 = Trait 1; T2 = Trait 2. 'Gray' filled boxes indicate that traits on individuals are measured and 'white' filled boxes are missed data. **A)** Subset measurement designs randomize a set of variables to be measured on a sample of individuals. Body mass (BM) is strongly correlated with all three other variables, and so, is measured on all individuals in the study. Molecular determination of sex is needed with our species, and along with thyroxine, can both be costly to quantify so these traits are measured on a sample of individuals. Metabolism can also be time consuming to measure, and so, it is only quantified on a sub-sample of animals. While we can estimate all main effects (i.e., single variables of interest) with this design, if interactions are of interest then one should have a PMDD that ensures there is enough data to effectively estimate the interaction parameters. **B)** Wave missing design can be applied to longitudinal data. Here, only two traits are quantified once a month on 20 individuals. In each measurement wave (i.e., a month), a random sample of individuals are completely lacking in data on both traits. **C)** Combined design. Longitudinal designs can miss entire sub-samples of individuals at lower rates, but also impose trait-level missingness for other individuals.

Figure 3 – Simulation results of the estimation of a between–individual correlation for two traits [$(Y_1, Y_2) = 0.40$] under varying levels of missing data (5–50%) and sample sizes (total number of observations; $n = 100$–1000, with number of individuals ranging from 10–100 each measured 10 times). Plots show average point estimates (A & B) and their corresponding standard errors (C & D) from 5000 randomly generated datasets when imputing missing data (A & C) or running a complete case analysis (excluding missing data – B & D). Note that convergence problems are more prevalent with small samples and high levels of missing data (see text). Parameter estimates and their precisions are therefore summarised on simulations in which models did converge.

Figure 4 – Example of experimental data on growth rates across age for two simulated scenarios. **A)** Mass of $n = 10$ lizards for a control group incubated at 23ºC and experimental group incubated at 26ºC. In this scenario, each animal was measured 10 times across the first 10 months of age. **B)** Mass of $n = 20$ lizards for a control group 23ºC and experimental group incubated at 26ºC measured five times with 50% of the mass data on each of the 20 animals considered missing ('red' points). In both scenarios, there were main effects of treatment and an interaction between growth across ages and treatment. Individual lizards varied in both their intercept and slope (see text for more details). Data was simulated according to eqn. 2 with the following parameters: $\beta_0 = 1.2$; $\beta_1 = 2$; $\beta_2 = 1.8$; $\beta_3 = 1.6$. In scenario B, missing data was imputed using likelihood based approaches in ASReml-R.

Figure 5– Example data showing the provisioning rates (number of feeding visits within a 5-hour period) for control (3 eggs) and experimentally elevated (6 eggs) brood sizes in the fictitious Missing Capped Warbler across the first 20 days of chick age. Provisions were simulated assuming a Poisson error distribution using the following model: $n_{ij} = (\beta_0 + \alpha_i) + \beta_1 T_{ij} + (\beta_2 + s_j)A_{ij} + \varepsilon_{ij}$ where Provisions = $\log(n_{ij})$. $\beta_0 = 2$; $\beta_1 = 1$; $\beta_2 = 0.01$ with a random intercept and slope variance and covariance matrix as: $\sim MVN\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0.01 & 0 \\ 0 & 0.0025 \end{bmatrix}\right)$.

BOX 1

Multiple imputation (MI) for a few different generalised family types can be implemented in the *mice* package (van Buuren & Groothuis-Oudshoorn 2011). Given that our data (number of visits in 5 hours) is Poisson distributed (or nearly so), and hierarchical in nature, we will need to impute using a new add-on developed in the *countimp* package (Kleinke & Reinecke 2013), which imputes Poisson random regressions. It can be installed, along with other needed packages in R as follows:

```
> link <- "http://www.uni-bielefeld.de/soz/kds/software/countimp_1.0.tar.gz"
> install.packages(link, repos=NULL, type="source")
> library(countimp)
> library(mice)
> library(glmmADMB)
```

Our data, including the missing data is set up as follows:

```
> head(data)
  ind age trt    trt_name provision
1   1   1   1 Experimental        NA
2   1   2   1 Experimental        22
3   1   3   1 Experimental        29
4   1   4   1 Experimental        38
5   1   5   1 Experimental        33
6   1   6   1 Experimental        37
```

In total, we have approximately 33% missing data at the individual level. To impute these missing data, we need to first set up the predictor matrix to define what variables are class (random effect groups) and what are to be used as random and fixed effects:

```
> data$ind <- as.integer(data$ind) # Need to keep class variable as integer
> imp <- mice(data, maxint = 0, printFlag = FALSE) # Do quick run of mice to set up pred matrix
> pred <- imp$pred # Extract the pred matrix
```

Now that we have the predictor matrix, to run a multi-level imputation we need to change the predictor matrix row for provision to define the class variable (i.e., random effect group – set as '–2' – only one level can be included currently), the variable included as both a fixed and random effect (i.e., age – set as '2' because we have a random regression model) and we will set our 'trt' as a fixed effect only (i.e., set as '1') as follows:

```
> pred["provision", ] <- c(-2,2,1,0,0)
```

The '0' is used to tell mice not to include these variables as predictors in the imputation. Now that this is set up we can run multiple imputation telling mice that the 'provision' variable is a 2-level Poisson variable:

```
> imp <- mice(data, m = 20, meth = c("","", "","","2l.poisson"), pred = pred)
```

This will impute missing information in the 'provision' variable creating $m = 20$ 'filled in' datasets for which estimates and standard errors can be pooled as follows:

```
> fit <- do.mira(imp = imp, DV = "provision", fixedeff = "trt+age", randeff = "1 + age",
grp = "ind", id = "ind", fam = "poisson")
> summary(fit)
```

We can compare the overall pooled estimates with estimates from our 'real complete dataset' (assuming we watched all videos):

	Full Dataset		Imputed Dataset	
Fixed Effects			*Pooled Fixed Effects*	
	Estimate	**SE**	**Estimate**	**SE**
Intercept	1.93	0.05	1.93	0.06
trt	1.13	0.06	1.13	0.07
age	0.01	0.01	0.01	0.01
Random Effects			*Pooled Random Effects*	
	Estimate		**Estimate**	
Intercept	0.011		0.015	
Age	0.0025		0.0023	