How the Central American Seaway and an ancient northern passage affected Flatfish diversification

LISA BYRNE[1], FRANÇOIS CHAPLEAU[1], AND STÉPHANE ARIS-BROSOU[1,2,*]

[1]*Department of Biology, University of Ottawa, Ottawa, ON, CANADA;* [2]*Department of Mathematics & Statistics, University of Ottawa, Ottawa, ON, CANADA*

[*]*Correspondence to be sent to: Department of Biology, University of Ottawa, 30 Marie Curie Pvt., Ottawa, ON, CANADA; Email: sarisbro@uottawa.ca*

Abstract

While the natural history of flatfish has been contentious for decades, the mode of diversification of this biologically and economically important group has never been elucidated. To address this question, we assembled the largest molecular data set to date, covering > 300 species (out of *ca*. 800 extant), from 13 of the 14 known families over nine genes, and employed relaxed molecular clocks to uncover their patterns of diversification. As the fossil record of flatfish is contentious, we used sister species distributed on both sides of the American continent to calibrate clock models based on the closure of the Central American Seaway (CAS), and on their current species range. We show that flatfish diversified in two bouts, as species that are today distributed around the Equator diverged during the closure of CAS, while those with a northern range diverged after this, hereby suggesting the existence of a post-CAS closure dispersal for these northern species, most likely along a northern route, a hypothesis fully compatible with paleogeographic reconstructions.

Keywords: Pleuronectiformes, molecular clock, Bayesian dating, vicariance, Central American Seaway, Isthmus of Panama

The Pleuronectiformes, or flatfishes, are a speciose group of ray-finned fish, containing

14 families and over 800 known species (Munroe 2015). Flatfish begin life in the pelagic

zone, but undergo a larval metamorphosis in which one eye, either left or right,

depending on the species, migrates to the other side of the cranium. The adult fish then

adopts a benthic lifestyle. Flatfish have asymmetric, laterally-compressed bodies, and

have lost their swim bladders during transformation. With eyes facing upwards, flatfish

are also capable of protruding them. This singular morphology long puzzled taxonomists

(Norman 1934), and the phylogeny of this group remains poorly resolved.

At the highest taxonomic level, flatfishes are generally considered to be monophyletic,

based on both morphological (Chapleau 1993), and molecular evidence (Berendzen et al.

2002; Pardo et al. 2005; Azevedo et al. 2008; Betancur-R et al. 2013; Harrington et al.

2016). All these studies also support the monophyletic status of most families within the

order, to the exception of the Paralichthyidae. As all molecular studies to date have

essentially focused on the monophyletic status of the order, they were based on as many

representative species of each order. As a result, intra-ordinal relationships, among

genera and even families, are still debated. It can therefore be expected that taking

advantage of both species- and gene-rich evidence, while incorporating paleontological

and/or geological data in the framework of molecular clocks, should help clarify not only

the phylogenetic status of this family (dos Reis et al. 2015), but also – and more critically

here – their evolutionary dynamics.

However, very few flatfish fossils are known, and placed with confidence on the

flatfish evolutionary tree (Parham et al. 2012). As a result, calibrating a molecular clock

becomes challenging. On the other hand, a dense species sampling may enable us to take

advantage of a singular feature of flatfishes: some extant species are found both in the

Pacific and Atlantic oceans. Furthermore, the existence of geminate species pairs of

flatfishes, where sister taxa have one member in each ocean, suggests a speciation event

pre-dating the closure of the Isthmus of Panama, which occurred approximately 12 to 3

million years ago [MYA] (Haug and Tiedemann 1998; ODea et al. 2016). Our driving

hypothesis is then that this information can be used to calibrate molecular clocks, and

allow us to unravel the timing of flatfish evolution, as how rapidly they diversified

remains an unsolved question. We show here that the diversification of flatfish in the sea

surrounding the Americas followed a complex pattern, related to not only the closure of

the Isthmus of Panama, but also to a warming event that opened up a northern route.

Results

To test how the evolutionary dynamics of flatfish were affected by a major geological

event, the closure of the CAS, we reconstructed dated Bayesian phylogenetic trees from a

large data matrix under four relaxed molecular clock models, each one of them being

based on a different calibration scheme (Fig. 1). Under the first model, no calibration

priors were placed on internal nodes. The initial data survey tree, with the rogue

sequences removed (data on GitHub) was used to identify pairs of sister taxa that are split

between the two oceans, with one species in the Atlantic and the other in the Pacific. This

led us to single out twelve pairs of such species. These sister species happened not to be

evenly distributed on the estimated phylogeny (Fig. 2, top), but they are the only

geminate species included in GenBank (as of August 2016). On each pair, we placed

calibration date priors corresponding to the closure of the CAS. An examination of the

posterior distributions of their divergence times suggests that some species had a very

narrow speciation window, where all the mass of the posterior distribution is between 5-3

MYA, while others have a wider distribution (Fig. 3A). Closer inspection of these

distributions further reveals that most of the species with narrow posterior distributions

have a northern range (Fig. 2, and 3A, in blue), while those with the wider posterior

distributions have a "southern" distribution, closer to the Isthmus of Panama (Fig. 2, and

3A, in red). To further assess this observation, we first went back to the original clock

model, with no priors on sister taxa, and were able to validate that even in this case,

northern and southern species showed, to one exception each (*Hippoglossus hippoglossus*

and *H. stenolepis* in the north, and *Poecilopsetta natalensis* and *P. hawaiiensis* in the

south; Fig. 2), shifted posterior distributions (Fig. 3B). The former pair was actually not

estimated as being sister species in any of the four clock models (Fig. 1), while *P.*

natalensis and *P. hawaiiensis,* although inhabiting the Western Atlantic and Eastern

Pacific oceans, occupy ranges that do not extend to the coasts of the Americas as with the

other identified sister taxa. Models with priors placed only on northern (Fig. 3C) or

southern (Fig. 3D) species also showed a similar temporal shift. This shift suggested that

southern species diverged early, before the complete closing of the CAS, while northern

species diverged later, at or possibly after the isthmus was completed. Averaging these

posterior distributions for the northern and southern species, to the exception of the two

outliers noted above, showed these results more clearly (Fig. 3E-H).

In an attempt to tease out these models and their predictions about the exact timing of

divergence between northern and southern species, we assessed model fit by means of

AICM. Even if model ranking based on this measure is known to be unstable (Baele et al.

2012), it is clear that the models with priors only on the northern or on the southern

species perform significantly (> 200 AIC units) more poorly than the two other models,

which may be difficult to tease apart (Table 1). Their predictions suggest that *H.*

hippoglossus and *H. stenolepis*, both northern species, consistently diverged before the

complete closure of the seaway, in tandem with the average southern species, and that the

average northern species diverged in tandem with *P. natalensis* and *P. hawaiiensis* (Fig.

3E-F).

105   Discussion

Our results show that flatfish underwent a first speciation at the time when the CAS

closed, which led to the species that, today, have an equatorial range (Fig. 2), as the

formation of the Isthmus of Panama resulted in a barrier to gene flow leading to their

speciation. Our results also show that species that today have a northern range (Fig. 2)

either emerged at the closure of the seaway, or after its closure. These timing estimates

imply that this second bout of speciation was not caused by gene flow impeded by the

closure of the seaway, but demand an interpretation involving gene flow through a

different route, a northern route, before being interrupted, most likely this time by a

climatic event. Strikingly, the geological evidence is directly in line with our date

estimates. The fossil record suggests that the first aquatic connection between the Pacific

and Arctic (and Atlantic) oceans through the Bering Strait occurred approximately 5.5-

5.4 MYA (Gladenkov et al. 2002) due to a rise in sea levels, linked to tectonic activity

(Marincovich and Gladenkov 2001). This would have permitted the migration of

populations ancestral to today's northern species from one ocean to the other through this

ancient "northern passage." This global warming event, between the late Miocene to

early Pleiocene, was then followed by a significant period of cooling during the Pleiocene

into the Pleistocene (Zachos 2001), leading to periods of repeated glaciations and a

subsequent ice age. These cold events would have resulted in the closure of this ancient

"northern passage," hereby stopping gene flow between the two oceans, and leading to

the more recent speciation of the northern taxa.

For approximately a million years after the first opening of the Bering Strait, water

flowed in a southern direction, until the formation of the Isthmus of Panama occurred

close to the equator (Berta 2012). With the formation of the Isthmus, and the closing of

the CAS, the ocean currents reversed due to a change in global ocean circulation (Haug

and Tiedemann 1998; De Schepper et al. 2015), and have since flowed from the Pacific

to the Arctic (Marincovich 2000). Dispersal or migration from the Pacific into the

Atlantic in a northern direction through this strait is known as the trans-Arctic

interchange (Vermeij 1991). Fossil data also show that the Bering land bridge has been

exposed and submerged on multiple occasions since the Pleistocene (Gladenkov and

Gladenkov 2004). These openings and closings of the Bering Strait could have provided a

mechanism for divergence and the evolution of sister taxa (Taylor and Dodson 1994;

Väinölä 2003).

Our results also have implication at the family level of flatfish. Further significant

global cooling during the Pleistocene resulted in major glaciation events (Zachos et al.

2008) that could be responsible for creating barriers that isolated populations. All of the

remaining sister taxa in our analysis, who have divergence estimates of less than 2 MYA

in our study belong to the family Pleuronectidae (*sensu* Cooper and Chapleau 1998). The

Pleuronectidae are the predominant flatfish family found in cold and temperate seas of

the northern hemisphere (Norman 1934; Cooper and Chapleau 1998). There are far more

Pleuronectidae species in the Pacific Ocean, most of them endemic to the north Pacific

Ocean off of north America and Asia in the region extending from the Bering Strait to the

gulf of California (Norman 1934). None of the arctic species are restricted solely to arctic

waters (Munroe 2015). Munroe also noted that Cooper (in an unpublished manuscript)

identified areas of endemism among the current distribution of the Pleuronectidae. It has

been shown that during the trans-Arctic interchange, there was a far higher number of

species (up to eight times higher) migrating to the Atlantic than to the Pacific (Vermeij

1991). Fossil and phylogenetic evidence suggest the Pacific Ocean as the origin for

diversification of the Pleuronectidae (Munroe 2015) and our phylogenetic results are

highly congruent with this hypothesis.

It is possible that the outliers, the Atlantic and Pacific halibut (*H. hippoglossus* and *H.*

stenolepis, respectively), diverged during one of the first openings of the Bering Strait.

The estimated dates from the molecular clock analysis are approximately 5.5 MYA, in

accordance with the hypothesized dates for the first aquatic connection (Marincovich and

Gladenkov 2001). The remaining sister taxa have a younger age estimate of ~1-2 MYA,

corresponding with global cooling during the Pleistocene and repeated glaciations

(Zachos 2001) which likely formed more barriers to genetic flow. In the second pair of

outliers, *P. hawaiiensis* inhabits waters of the Eastern Central Pacific to the Hawaiian

Islands, while *P. natalensis* inhabits coastal waters of Eastern Africa (Fig. 2). Due to the

far reaching range of *P. natalensis*, and the relatively younger age estimates for

divergence (1-2 MYA), these results beg for future research into other vicariant

hypotheses, or dispersal routes, as these two members of *Poecilopsetta* diverged long

after the Isthmus had closed.

Based on the most extensive multigene sequence alignment available to date across

all flatfish species, we have showed here that the evolutionary dynamics of sister species

that are distributed across the two oceans strongly supported the existence of two bouts of

speciation: one triggered by the closure of the CAS 12-3 MYA, and a second one due to

the closure of an ancient northern passage 5-0.01 MYA. The most intriguing implication

of our results is that other sea critters than the flatfish should have been affected by the

two same geological processes, the closure of the CAS (Bacon et al. 2015) and, most

intriguingly, of the ancient northern passage, so that future studies should be able to

extend our findings to other living forms inhabiting these two oceans.

Methods and Materials

Methods and Materials

Data retrieval and alignment

Prior to retrieving sequence data, GenBank was surveyed to identify all the genes

belonging to species of Pleuronectiform families (as of August 2016), based on

GenBank's taxonomy browser. DNA sequences for a total of 332 flatfish species (out of

over 800 species in the order) were identified and downloaded for five nuclear genes

(KIAA1239, MYH6, RIPK4, RAG1, SH3PX3), and four mitochondrial genes (12S, 16S,

COX1 and CYTB). These represented all the taxa having at least one of these six gene

sequences in GenBank; see Table S1 for the corresponding accession numbers. Diversity

was richly sampled as species from 13 of the 14 families in the order Pleuronectiformes

were included in our catchment. In line with current consensus in flatfish systematics, the

family Psettodidae was chosen as the outgroup to all other taxa (Chapleau 1993).

These sequences were aligned using MUSCLE ver. 3.8.31 (Edgar 2004) on a gene-

by-gene basis. Each alignment was visually inspected with AliView ver. 1.18 (Larsson

2014), and was manually edited where necessary. In particular, large indels (> 10 bp)

were removed prior to all phylogenetic analyses. The 5' and 3' ends of sequences were

also trimmed. The aligned sequences were then concatenated into a single alignment

using a custom R script.

Data pre-processing

To gauge the phylogenetic content of our data set, we performed a first series of

molecular clock analyses on the concatenated data matrix, with all the nine gene

sequences obtained above (12S, 16S, COI, Cyt-b, KIAA1239, MYH6, RIPK4, SH3PX3

and RAG1), and all the 332 taxa representing all families of flatfish. To account for rate

heterogeneity along this alignment, partitions were first identified with PartitionFinder 2

(Lanfear 2012), selecting the best-fit model of nucleotide substitution based on Akaike's

Information Criterion (AIC). This optimal partition scheme was then employed in a first

Bayesian phylogenetic analysis, conducted with BEAST ver. 1.8.3 (Drummond and

Rambaut 2007). This program implements a Markov chain Monte Carlo (MCMC)

sampler, which co-estimates both tree topology and divergence times. As determined by

PartionFinder, a GTR+I+Γ model was applied to each data partition. These models of

evolution across partitions were assumed to be independent (*unlinked*, in BEAST

parlance), while both clock model and tree model partitions were shared (*linked*) by all

partitions. An uncorrelated relaxed clock was employed with a lognormal distribution

prior on rates, and a Yule speciation prior (Drummond et al. 2006). Due to the paucity of

reliably placed fossils on the flatfish tree, a unique calibration point was placed on the

most recent common ancestor (MRCA) of the ingroup, as a mean-one exponential prior,

with an offset of 40 million years reflecting the age of 47.8-42.1 MYA (Chanet 1997). To

stabilize the estimate, a lognormal ln(0, 1.5) prior with a 40 MYA offset was also placed

on the root of the tree (root height). Two separate MCMC samplers were run, each for

10,000,000 generations. Trees were sampled every 5,000 generations, and convergence

was checked visually using Tracer ver. 1.6. (Rambaut et al. 2014). Tree log files from

each run were combined in LogCombiner, after conservatively removing 10% of each

run as burn-in. The resulting maximum a posteriori (MAP) tree was generated with

TreeAnnotator (Drummond and Rambaut 2007).

As the topology of this resulting MAP tree was unconventional, we suspected the

presence of rogue taxa, *i.e.* species evolving either much faster or much slower than the

majority, which can contribute negatively to consensus tree support (Aberer et al. 2013).

Rogue taxa were identified using the RogueNaRok (Aberer et al. 2013) webserver

(http://rnr.h-its.org). The consensus trees from the preliminary analysis using 10,000

iterations were used as the tree set for rogue taxon analysis. A total of 22 rogues were

identified and pruned from the analysis, leaving 310 species.

Molecular Dating

To assess the impact of the closure of the Central American Seaway (CAS; forming the

Isthmus of Panama) on flatfish evolutionary dynamics, a second set of partitioned relaxed

molecular clock analyses was performed (without the rogue sequences). The timing of

the closure of the CAS is estimated to have occurred between 12 and 3 MYA (Duque-

Caro 1990; Coates et al. 1992; Haug and Tiedemann 1998; ODea et al. 2016), and we

used this time window as a prior to inform the relaxed molecular clock-based

phylogenetic reconstructions. The analyses were performed on the same concatenated

data set, with the same single fossil calibration, but we also placed a lognormal prior

ln(3,1.5), that has most of its mass on the 12-3 MYA window, on the MRCA of each pair

of sister taxa. From the initial BEAST analyses, twelve pairs of taxa were selected based

on two criteria: (i) being sister species on that initial tree, (ii) with one species distributed

in the Pacific and one in the Atlantic ocean (Fig. 2). Again, two independent MCMC

samplers were run each for 100 million iterations, with samples taken every 5000 step.

Because these pairs of sister species show a contrasted geographic distribution,

having either a southern (equatorial) or a northern range (Fig. 2), two additional sets of

analyses were performed. In a first set, calibration priors (ln(3,1.5), as *per* above) were

placed only sister taxa that had a geographic range in the southern hemisphere, while in a

second set, identical calibration priors were placed only on sister species with a northern

range. Finally, a set of analyses was performed using no sister taxa calibrations at all. For

each analysis, results from the two MCMC runs were combined using LogCombiner after

removing an even more conservative burn-in of 50%. The final MAP tree was generated

with TreeAnnotator.

In an attempt to rank these different models (priors on all sister taxa; only on southern

taxa; only on southern taxa; no "CAS" priors), a modified Akaike's Information Criterion

(AICM) that accounts for uncertainly in the MCMC sampling (Raftery et al. 2007) was

computed for each model (Baele et al. 2013). These computations were performed in

Tracer for each of the four different calibration models, based on 100 replicates.

Acknowledgments

This work was supported by the University of Ottawa (LB), and the Natural Sciences Research Council of Canada (FC, SAB). We are grateful to Jonathan Dench for discussions and comments, as well as to Compute Canada and Ontario's Center for Advanced Computing for providing us access to their resource.

References

Aberer A.J., Krompass D., Stamatakis A. 2013. Pruning rogue taxa improves phylogenetic accuracy: An efficient algorithm and webservice. Syst. Biol. 62:162–166.

Azevedo M.F.C., Oliveira C., Pardo B.G., Martínez P., Foresti F. 2008. Phylogenetic analysis of the order Pleuronectiformes (Teleostei) based on sequences of 12S and 16S mitochondrial genes. Genet. Mol. Biol. 31:284–292.

Bacon C.D., Silvestro D., Jaramillo C., Smith B.T., Chakrabarty P., Antonelli A. 2015. Biological evidence supports an early and complex emergence of the Isthmus of Panama. Proc Natl Acad Sci. 19112:6110–6115.

Baele G., Li W.L.S., Drummond A.J., Suchard M.A., Lemey P. 2013. Accurate model selection of relaxed molecular clocks in Bayesian phylogenetics. Mol. Biol. Evol. 30:239–243.

Berendzen P.B., Dimmick W.W., McEachran J.D. 2002. Phylogenetic relationships of Pleuronectiformes based on molecular evidence. Copeia. 2002:642–652.

Berta A. 2012. Return to the Sea: The life and evolutionary times of marine mammals.

281   University of California Press.

282 Betancur-R R., Li C., Munroe T.A., Ballesteros J.A., Ortí G. 2013. Addressing gene tree

283   discordance and non-stationarity to resolve a multi-locus phylogeny of the flatfishes

284   (Teleostei: Pleuronectiformes). Syst. Biol. 62:763–785.

285 Chanet B. 1997. A cladistic reappraisal of the fossil flatfishes record: consequences on

286   the phylogeny of the Pleuronectiformes (Osteichthyes: Teleostei). Ann. des Sci. Nat.

287   Zool. Paris. 18:105–117.

288 Chapleau F. 1993. Pleuronectiform relationships : a cladistic reassessment. Bull. Mar. Sci.

289   52:516–540.

290 Coates A.G., Jackson J.B.C., Collins L.S., Cronin T.M., Dowsett J., Bybell L.M., Jung P.,

291   Obando J. a. 1992. Closure of the Isthmus of Panama : The near-shore marine record

292   of Costa Rica and western Panama. Geol. Soc. Am.:814–828.

293 Cooper J.A., Chapleau F. 1998. Monophyly and intrarelationships of the family

294   Pleuronectidae (Pleuronectiformes), with a revised classification. Fish. Bull.

295   96:686–726.

296 Drummond A.J., Ho S.Y.W., Phillips M.J., Rambaut A. 2006. Relaxed phylogenetics and

297   dating with confidence. PLoS Biol. 4:699–710.

298 Drummond A.J., Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by sampling

299   trees. BMC Evol. Biol. 7:214.

300 Duque-Caro H. 1990. Neogene stratigraphy, paleoceanography and paleobiology in

301   northwest South America and the evolu- tion of the Panama Seaway. Palaeogeogr.

302   Palaeoclimatol. Palaeoecol. 77:203–234.

303 Edgar R. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

304   throughput. Nucleic Acids Res. Res.32(5):1792–1797.

305 Gladenkov A.Y., Oleinik A.E., Marincovich L., Barinov K.B. 2002. A refined age for the

306   earliest opening of Bering Strait. Palaeogeogr. Palaeoclimatol. Palaeoecol. 183:321–

307   328.

308 Gladenkov, Gladenkov A.Y. 2004. Onset of connections between the Pacific and Arctic

309   Oceans through the Bering Strait in the Neogene. Stratigr. Geol. Correl. 12:175–187.

310 Harrington R.C., Faircloth B.C., Eytan R.I., Smith W.L., Near T.J., Alfaro M.E.,

311   Friedman M. 2016. Phylogenomic analysis of carangimorph fishes reveals flatfish

asymmetry arose in a blink of the evolutionary eye. BMC Evol. Biol. 16:224.

Haug G.H., Tiedemann R. 1998. Effect of the formation of the Isthmus of Panama on

Atlantic Ocean thermohaline circulation. Nature. 393:673–676.

Larsson A. 2014. AliView: A fast and lightweight alignment viewer and editor for large

datasets. Bioinformatics. 30:3276–3278.

Marincovich L., Gladenkov A.Y. 2001. New evidence for the age of Bering Strait. Quat.

Sci. Rev. 20:329–335.

Marincovich L.J. 2000. Central American paleogeography controlled Pliocene Arctic

Ocean molluscan migrations. Geology. 28:551–554.

Munroe T.A. 2015. Systematic Diversity of the Flatfishes. In: Robin Gibson, Richard

Nash, Audrey Geffen H.V. der V., editor. Flatfishes: biology and exploitation. Wiley.

p. 13–44.

Norman J.R. 1934. A systematic monograph of the flatfishes (Heterosomata). By J.R.

Norman, London.

ODea A., Lessios H.A., Coates A.G., Eytan R.I., Restrepo-Moreno S.A., Cione A.L.,

Collins L.S., de Queiroz A., Farris D.W., Norris R.D., Stallard R.F., Woodburne

M.O., Aguilera O., Aubry M.-P., Berggren W.A., Budd A.F., Cozzuol M.A.,

Coppard S.E., Duque-Caro H., Finnegan S., Gasparini G.M., Grossman E.L.,

Johnson K.G., Keigwin L.D., Knowlton N., Leigh E.G., Leonard-Pingel J.S., Marko

P.B., Pyenson N.D., Rachello-Dolmen P.G., Soibelzon E., Soibelzon L., Todd J.A.,

Vermeij G.J., Jackson J.B.C. 2016. Formation of the Isthmus of Panama. Sci. Adv.

2:e1600883–e1600883.

Pardo B.G., Machordom A., Foresti F., Porto-Foresti F., Azevedo M.F.C., Bañon R.,

Sánchez L., Martínez P. 2005. Phylogenetic analysis of flatfish (Order

Pleuronectiformes) based on mitochondrial 16S rDNA sequences. Sci. Mar. 69:531–

543.

Parham J.F., Donoghue P.C.J., Bell C.J., Calway T.D., Head J.J., Holroyd P.A., Inoue

J.G., Irmis R.B., Joyce W.G., Ksepka D.T., Patané J.S.L., Smith N.D., Tarver J.E.,

Van Tuinen M., Yang Z., Angielczyk K.D., Greenwood J.M., Hipsley C.A., Jacobs

L., Makovicky P.J., Müller J., Smith K.T., Theodor J.M., Warnock R.C.M., Benton

M.J. 2012. Best practices for justifying fossil calibrations. Syst. Biol. 61:346–359.

Raftery A.E., Newton M.A., Satagopan J.M., Krivitsky P.N. 2007. Estimating the
integrated likelihood via posterior simulation using the Harmonic Mean Identity.
Bayesian Stat.:1–45.

Rambaut A., Suchard M.A., Xie D., Drummond A.J. 2014. Tracer v1.6.

dos Reis M., Donoghue P.C.J., Yang Z. 2015. Bayesian molecular clock dating of species
divergences in the genomics era. Nat. Rev. Genet. 17:71–80.

De Schepper S., Schreck M., Beck K.M., Matthiessen J., Fahl K., Mangerud G. 2015.
Early Pliocene onset of modern Nordic Seas circulation related to ocean gateway
changes. Nat. Commun. 6:8659.

Taylor E.B., Dodson J.J. 1994. A molecular analysis biogeography within (genus
Osmevus) of relationships and a species complex of Holarctic fish. Mol. Ecol.
3:235–248.

Väinölä R. 2003. Repeated trans-Arctic invasions in littoral bivalves: Molecular
zoogeography of the *Macoma balthica* complex. Mar. Biol. 143:935–946.

Vermeij G.J. 1991. Anatomy of an invasion: the trans-Arctic interchange. Paleobiology.
17:281–307.

Zachos J. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present.
Science (80-.). 292:686–693.

Zachos J.C., Dickens G.R., Zeebe R.E. 2008. An early Cenozoic perspective on
greenhouse warming and carbon-cycle dynamics. Nature. 451:279–283.

Table 1. AICM values for the phylogenetic analyses using four different calibration

schemes. SE: standard error.

Calibration Model	AICM value	SE
All	374,082.199	+/-1.579
None	374,110.488	+/-3.789
South	374,367.749	+/-0.696
North	374,746.628	+/-3.789

Figure captions

FIG. 1. Phylogenetic trees reconstructed based on relaxed clock models. Four models were employed, representing different specifications of prior distributions set on sister taxa. (**A**) No priors were set on sister taxa. (**B)** Priors were set on all pairs of taxa. (**C**) Priors were set only on sister taxa with a current northern range. (**D)** Priors were set only on sister taxa with a current southern range. Horizontal scale is in million years ago (MYA). The closure of the CAS (12-3 MYA) is shown between vertical gray broken lines.

FIG. 2. Distribution of the geminate species used in this study. Geographic distribution of the twelve pairs of sister species of flatfish used to calibrate the relaxed molecular clock models. Original data come from GBIF (www.gbif.org; accessed Nov 9, 2017). The top panel shows the phylogenetic distribution of these species based on the clock model including prior ages on all twelve pairs of species; scale bar: 5 MYA.

FIG. 3. Posterior densities of divergence times for sister taxa used as calibration points in the relaxed molecular clock analyses, under four different calibration schemes. NONE: no priors were placed on sister taxa; ALL: priors were placed on all pairs of sister taxa; NORTH: priors were placed on only pairs of sister taxa with a northern distribution; SOUTH: priors were placed on only pairs of sister taxa with a southern distribution. Top row (**A-D**) shows all 17 distributions, while bottom row (**E-H**) shows range-averaged distributions (solid) to the exception of outlier pairs (dashed lines). Densities are color-coded for species with northern (blue) and southern (red) range. Dashed vertical gray lines indicate the closure of the CAS (21-3 MYA).

9:L. aspera   8:L.proboscidea

11:H. stenolepis

7:P. quadrituberculatus

1:H. stomata

2:P. hawaiiensis

5:S. maculiferum

3:C. platophrys

6:C. querna

4:C. darwini

9:L. limanda

10:H. platessoides

12:G. cynoglossus

11:H. hippoglossus

7:P. platessa

1:H. oblonga
8:L. ferruginea
3:C. arctifrons
5:S. micrurum
6:C. chittendeni
4:C. minutus

2:P. natalensis

12:Glyptocephalus stelleri/Glyptocephalus cynoglossus
11:Hippoglossus stenolepis/Hippoglossus hippoglossus
10:Hippoglossoides dubius/Hippoglossoides platessoides
9:Limanda aspera/Limanda limanda
8:Limanda ferruginea/Limanda proboscidea
7:Pleuronectes platessa/Pleuronectes quadrituberculatus
6:Cyclopsetta chittendeni/Cyclopsetta querna
5:Syacium maculiferum/Syacium micrurum
4:Citharichthys darwini/Citharichthys minutus
3:Citharichthys arctifrons/Citharichthys platophrys
2:Poecilopsetta hawaiiensis/Poecilopsetta natalensis
1:Hippoglossina oblonga/Hipoglossina stomata

Northern

Southern

6 5 4   3   1   11 12   9 10   8 7   2